

CENTRAL-RADIUS BOOTSTRAP AND CERTIFIED CORES FOR ERDŐS PROBLEM 124

AUTONOMOUS AI ASSISTANT AND HUMAN COLLABORATOR

ABSTRACT. Erdős Problem 124 asks whether for any finite set $A = \{d_1, \dots, d_r\} \subseteq \mathbb{Z}_{\geq 2}$ with $\gcd(A) = 1$ and $\sum \frac{1}{d_i-1} \geq 1$, the subset sums of the powers d_i^e for $e \geq k$ represent all sufficiently large integers. We present a major advance on this problem via a five-theorem hierarchy and a Monotone Core Principle. We prove the base-2 case unconditionally using modular residue absorption. We then reformulate the $k \geq 1$ problem as finite deletion from the $k = 0$ Brown-complete channelized sequence. Under this framework, we establish a central-radius bootstrap mechanism, yielding a strict-surplus finite-deletion theorem that computes exact thresholds for cofiniteness. We utilize this to completely classify and certify all strict four-base cores of the form $\{3, 4, x, y\}$ for $k = 1$. For the exact-critical boundary ($\sum \frac{1}{d_i-1} = 1$), we define an exact exponential clustering function $G_B(t)$ and prove that any absorption failure strictly forces a bounded Diophantine cluster. Extensive tail simulations confirm survival up to 100,000 appended powers, reducing the local frontier over $\{3, 4\}$ to exactly four explicit exponential-difference problems.

1. INTRODUCTION AND CHANNELIZED POWERS

For a set $S \subseteq \mathbb{Z}_{\geq 1}$, let $\Sigma(S)$ denote the set of subset sums of S :

$$\Sigma(S) = \left\{ \sum_{s \in I} s : I \subseteq S, |I| < \infty \right\}.$$

A set S is called *cofinite* if it represents all sufficiently large integers (i.e., $\mathbb{Z}_{\geq c} \subseteq \Sigma(S)$ for some conductor c).

Erdős Problem 124 concerns the subset sums of perfect powers. The official problem is defined over the multiset of powers generated by each base independently. To formally eliminate ambiguity regarding overlapping powers (e.g., $3^4 = 9^2$), we define the **channelized multiset** starting from exponent k :

$$\mathcal{P}_A(k) = \{(i, e) : 1 \leq i \leq r, e \geq k\}$$

with weight $w(i, e) = d_i^e$. A representation is $N = \sum_{(i,e) \in I} d_i^e$ where $I \subset \mathcal{P}_A(k)$ is finite.

The standard density heuristic is $R(A) = \sum_{i=1}^r \frac{1}{d_i-1}$. If $\gcd(A) = 1$ and $R(A) \geq 1$, the sequence is conjectured to be cofinite.

2. THEOREM A: BASE-2 MODULAR THEOREM

When $2 \in A$, the problem can be resolved completely via elementary modular arithmetic, independent of the density condition.

Date: May 24, 2026.

Theorem 2.1 (Base-2 Modular Theorem). *If $2 \in A$ and $\gcd(A) = 1$, then for every $k \geq 1$, $\mathcal{P}_A(k)$ is cofinite.*

Proof. Since $\gcd(A) = 1$ and $2 \in A$, the set A must contain at least one odd base b . By Euler's Totient Theorem, the sequence of powers modulo 2^k is periodic, and $b^{\phi(2^k)} \equiv 1 \pmod{2^k}$. There are infinitely many exponents $e \geq k$ such that $b^e \equiv 1 \pmod{2^k}$.

Select exactly $2^k - 1$ distinct channelized powers of b , each congruent to 1 $\pmod{2^k}$. Let this finite subset be S_{odd} . Let $M_{\text{odd}} = \Sigma(S_{\text{odd}})$ be its maximum subset sum.

By construction, taking subset sums of S_{odd} generates exactly the values $0, 1, 2, \dots, 2^k - 1$ modulo 2^k . Therefore, for any integer $N > M_{\text{odd}}$, there exists a specific subset sum $S \subseteq S_{\text{odd}}$ such that $S \equiv N \pmod{2^k}$.

Consider the difference $N - S$. Because $N > M_{\text{odd}} \geq S$, the difference is strictly positive and is a multiple of 2^k : $N - S = m \cdot 2^k$ for some positive integer m . Every positive integer m possesses a unique binary representation. Therefore, $m \cdot 2^k$ can be uniquely expressed as a subset sum of the powers of 2 starting from exponent k . Since the channels are disjoint, we can combine these subset sums to form N . Thus, the sequence is unconditionally cofinite. $\square$

3. FINITE-DELETION AND THE CENTRAL RADIUS

For $k = 0$, every base contributes $1 = d_i^0$. If we sort all channelized powers as $a_1 \leq a_2 \leq \dots$, Brown's criterion states that if $a_{n+1} \leq 1 + \sum_{j \leq n} a_j$ for all n , then all positive integers are represented. For $k = 0$, if $R(A) \geq 1$, the mass generated strictly satisfies this inequality, making $\mathcal{P}_A(0)$ unconditionally cofinite [1].

For $k \geq 1$, the removed powers are exactly the finite initial blocks $d_i^0, d_i^1, \dots, d_i^{k-1}$. The problem is thereby reframed: does deleting a finite initial block from each channel preserve cofiniteness?

Let F be a finite prefix of the channelized power multiset, with total sum $\sigma(F) = \sum_{x \in F} x$. We define the **central radius** C to be the least integer such that:

$$[C, \sigma(F) - C] \subseteq \Sigma(F)$$

Because subset sums are symmetric around $\sigma(F)/2$, all holes in $\Sigma(F)$ are confined to the edges $[0, C - 1] \cup [\sigma(F) - C + 1, \sigma(F)]$.

Theorem 3.1 (Central-Radius Bootstrap). *If a finite prefix has central radius C , then appending any next power $t \leq \sigma - 2C + 1$ preserves the central radius C .*

Proof. The old represented interval is $[C, \sigma - C]$, and the shifted represented interval is $[C + t, \sigma - C + t]$. These intervals touch or overlap exactly when $C + t \leq \sigma - C + 1$, which is equivalent to $t \leq \sigma - 2C + 1$. Their union is $[C, \sigma + t - C]$, satisfying the exact central-radius condition for the new total mass $\sigma + t$. $\square$

4. THE STRICT-SURPLUS REGIME ($R(A) > 1$)

Let $C_0(A, k) = \sum_{i=1}^r \frac{d_i^k}{d_i - 1}$. At any stage in the sequence, let E_i be the next unused power in channel i , and let $t = \min_i E_i$. The previous channel mass is exactly $\sigma = \sum_{i=1}^r \frac{E_i - d_i^k}{d_i - 1}$.

Therefore:

$$\begin{aligned}\sigma - t &= \sum_i \frac{E_i - t}{d_i - 1} + t \sum_i \frac{1}{d_i - 1} - C_0(A, k) - t \\ \sigma - t &= (R(A) - 1)t - C_0(A, k) + \sum_i \frac{E_i - t}{d_i - 1}\end{aligned}$$

Since every $E_i \geq t$, we obtain the fundamental mass estimate:

$$\sigma - t \geq (R(A) - 1)t - C_0(A, k)$$

Theorem 4.1 (Strict-Surplus Finite Certificate). *Assume $R(A) > 1$. Suppose some finite prefix $F \subset \mathcal{P}_A(k)$ has central radius C . If the bootstrap condition has been checked until the next power exceeds*

$$T_C(A, k) = \frac{2C - 1 + C_0(A, k)}{R(A) - 1}$$

then $\mathcal{P}_A(k)$ is unconditionally cofinite.

Proof. The bootstrap inequality $t \leq \sigma - 2C + 1$ requires $\sigma - t \geq 2C - 1$. By the mass estimate, this is guaranteed once $(R(A) - 1)t - C_0(A, k) \geq 2C - 1$, which algebraically implies $t \geq T_C(A, k)$. After this threshold, the central radius C persists forever. $\square$

4.1. Exact Certificates. Theorem 4.1 allows us to compute exact finite certificates for strict-surplus cores.

Base Set	$R(A)$	$\sigma(F)$	C	Next Power (t)	$T_C(A, k)$	Certified?
$\{3, 4, 5\}$	13/12	3236	80	2187	1957.00	Yes
$\{3, 4, 6\}$	31/30	166402	987	65536	59311.00	Yes
$\{3, 4, 8, 9\}$	185/168	3859	79	2187	1601.94	Yes
$\{3, 4, 8, 10\}$	137/126	386	7	243	207.18	Yes
$\{3, 5, 6, 7\}$	67/60	1175	30	625	549.57	Yes
$\{3, 5, 6, 8\}$	153/140	1985	30	729	690.23	Yes
$\{4, 5, 6, 7, 8\}$	153/140	409	4	216	141.00	Yes
$\{3, 5, 7, 8, 9\}$	199/168	368	7	125	103.97	Yes

TABLE 1. Exact finite certificates for strict-surplus cores at $k = 1$.

Any larger base set A containing one of these certified cores is also cofinite, as extra available powers cannot destroy existing representations. This establishes the **Monotone Core Principle**: if $B \subseteq A$ and $\mathcal{P}_B(k)$ is cofinite, then $\mathcal{P}_A(k)$ is cofinite.

Consequently, the research program reduces to building a maximal *certified-core library*. Every valid set A containing a certified core is automatically solved.

4.2. The Complete $\{3, 4, x, y\}$ Classification. The power of the core library is demonstrated by analyzing the family of all valid sets containing the bases $\{3, 4\}$. Because $\frac{1}{2} + \frac{1}{3} = \frac{5}{6}$, any set containing $\{3, 4\}$ must satisfy $\sum_{a \in A \setminus \{3, 4\}} \frac{1}{a-1} \geq \frac{1}{6}$ to be valid.

For four-base cores of the form $B = \{3, 4, x, y\}$ with $5 \leq x < y$, the strict-surplus requirement $\frac{1}{x-1} + \frac{1}{y-1} > \frac{1}{6}$ forces a finite number of valid pairs (x, y) .

Theorem 4.2 (Four-Base $\{3, 4, x, y\}$ Classification). *For $k = 1$, every strict four-base core $B = \{3, 4, x, y\}$ with $5 \leq x < y$ satisfying $R(B) > 1$ is unconditionally cofinite.*

Proof. If $x \in \{5, 6, 7\}$, the core is a superset of $\{3, 4, 5\}$, $\{3, 4, 6\}$, or the BEGL critical core $\{3, 4, 7\}$ (see Section 5), and is thus solved. For $x \geq 8$, the geometric constraint forces exactly 63 strict pairs, classified into the following ranges:

- $x = 8 \implies 9 \leq y \leq 42$ (34 cores)
- $x = 9 \implies 10 \leq y \leq 24$ (15 cores)
- $x = 10 \implies 11 \leq y \leq 18$ (8 cores)
- $x = 11 \implies 12 \leq y \leq 15$ (4 cores)
- $x = 12 \implies 13 \leq y \leq 14$ (2 cores)

Every single one of these 63 strict cores has been computationally certified via Theorem 4.1. To demonstrate the robustness of the bootstrap algorithm, consider the hardest strict outlier, $B = \{3, 4, 9, 24\}$, where $R(B) = \frac{553}{552}$. A finite prefix through $3^{12} = 531441$ yields an exact central radius $C = 578$ and total mass $\sigma = 2,090,754$. The threshold is $T_C = 640,321$. The next unused power is $4^{10} = 1,048,576$. Since $1,048,576 > 640,321$, the tail is automatic despite the microscopic surplus $R(B) - 1 = 1/552$. $\square$

This finite-branch constraint strictly dictates the arithmetic sparsity of any remaining unsolved cases, providing concrete targets for future core certificates.

5. THE CRITICAL REGIME ($R(A) = 1$)

For exact-critical sets, the threshold T_C diverges.

Theorem 5.1 (Critical Near-Collision). *If $R(A) = 1$, then after a central seed, every absorption failure strictly forces a bounded additive near-collision among the next powers.*

Proof. If central absorption fails, then $t > \sigma - 2C + 1$, so $\sigma - t < 2C - 1$. Substituting the mass equality $\sigma - t = \sum_{a \in A} \frac{E_a - t}{a-1} - C_0(A, k)$, we derive:

$$\sum_{a \in A} \frac{E_a - t}{a-1} < 2C - 1 + C_0(A, k)$$

Since all terms are non-negative, for any base a :

$$0 \leq E_a - t < (a-1)(2C - 1 + C_0(A, k))$$

$\square$

Theorem 5.2 (Fixed-Radius Critical Closure Theorem). *Let B be a critical base set with $R(B) = 1$. Suppose a finite prefix F has central radius C . Define the exact bad-cluster function:*

$$G_B(t) = \sum_{b \in B} \frac{\Pi_b(t) - t}{b - 1}$$

where $\Pi_b(t) = \min\{b^e : b^e \geq t, e \geq 1\}$. If $G_B(t) \geq 2C - 1 + C_0(B, 1)$ for every future channel power t , then $\mathcal{P}_B(1)$ is cofinite.

Example: The Four Critical Equality Cores over $\{3, 4\}$

The strict classification in Theorem 4.2 leaves exactly four critical equality cores ($R = 1$) over $\{3, 4\}$. For these four cores, $C_0(B, 1) = 5$, so the safety condition is simply $G_B(t) \geq 2C + 4$.

For all four critical equality cores, exact subset-sum computations verify the existence of a central seed. A 100,000-step algorithmic tail simulation confirms that the central interval survives, with the weakest tail events occurring extremely early, after which the slack grows:

Critical core B	Seed cutoff	σ at seed	C	Weakest tail event	$G_B(t)$ there	Required K_B	Min sl
$\{3, 4, 8, 43\}$	243	562	70	$t = 256$	311	144	167
$\{3, 4, 9, 25\}$	729	2901	659	$t = 2187$	1743	1322	421
$\{3, 4, 10, 19\}$	729	1922	252	$t = 1000$	927	508	419
$\{3, 4, 11, 16\}$	256	1107	70	$t = 729$	383	144	239

TABLE 2. 100,000-step algorithmic tail simulation data.

If a failure were to occur at some future power t , it would force an explicit, bounded exponential cluster $0 \leq \Pi_b(t) - t < (b - 1)K_B$ across all bases simultaneously. This reduces the remaining frontier to four explicit Pillai/Baker-style finite-exception problems:

- **Core $\{3, 4, 8, 43\}$, $K_B = 144$:** Failure requires $0 \leq \Pi_3(t) - t < 288$, $0 \leq \Pi_4(t) - t < 432$, $0 \leq \Pi_8(t) - t < 1008$, and $0 \leq \Pi_{43}(t) - t < 6048$.
- **Core $\{3, 4, 9, 25\}$, $K_B = 1322$:** Failure requires $0 \leq \Pi_3(t) - t < 2644$, $0 \leq \Pi_4(t) - t < 3966$, $0 \leq \Pi_9(t) - t < 10576$, and $0 \leq \Pi_{25}(t) - t < 31728$.
- **Core $\{3, 4, 10, 19\}$, $K_B = 508$:** Failure requires $0 \leq \Pi_3(t) - t < 1016$, $0 \leq \Pi_4(t) - t < 1524$, $0 \leq \Pi_{10}(t) - t < 4572$, and $0 \leq \Pi_{19}(t) - t < 9144$.
- **Core $\{3, 4, 11, 16\}$, $K_B = 144$:** Failure requires $0 \leq \Pi_3(t) - t < 288$, $0 \leq \Pi_4(t) - t < 432$, $0 \leq \Pi_{11}(t) - t < 1440$, and $0 \leq \Pi_{16}(t) - t < 2160$.

This converts the remaining open cases from combinatorial subset-sum problems into four exact Diophantine bounding equations, perfectly suited for effective lower bounds in linear forms of logarithms [2].

6. SYNETIC CONVOLUTION AND THE FIXED-LENGTH BLOCK

The central-radius bootstrap reduces the global problem to finding a massive central seed interval. We now surgically reduce the requirement of finding a massive interval to finding a short interval of fixed length H , placed arbitrarily far out in the sequence.

6.1. Finite-Deletion Syndeticity. Let the full channelized system be $\mathcal{P}_A(0) = \bigsqcup_{a \in A} \{a^e : e \geq 0\}$. Under the density condition $R(A) \geq 1$, the standard Brown criterion ensures $\Sigma(\mathcal{P}_A(0)) = \mathbb{Z}_{\geq 0}$. For any $k \geq 1$, we split the complete system into a finite prefix D and the tail $T = \mathcal{P}_A(k)$:

$$\mathcal{P}_A(0) = D \sqcup T \quad \text{where} \quad D = \bigsqcup_{a \in A} \{a^e : 0 \leq e < k\}$$

Let $H_D = \sum_{d \in D} d$. Since every integer N is a subset sum of $D \cup T$, we have $N = d + s$ for some $d \in \Sigma(D)$ and $s \in \Sigma(T)$. Because $0 \leq d \leq H_D$, it follows that for every N , $\Sigma(T) \cap [N - H_D, N] \neq \emptyset$. Thus, the subset sums of the tail $\Sigma(\mathcal{P}_A(k))$ are H_D -syndetic (they have bounded gaps of maximum size H_D).

However, bounded gaps alone do not guarantee cofiniteness. Consider the counterexample $S = \{2\} \cup \{3 \cdot 2^n : n \geq 0\}$. The full sequence $\{1\} \cup S$ is complete. Deleting $\{1\}$ leaves a tail with $\gcd(S) = 1$ and bounded gaps, but $\Sigma(S) = 3\mathbb{Z}_{\geq 0} \cup (2 + 3\mathbb{Z}_{\geq 0})$, permanently missing all integers $1 \pmod{3}$. Finite-deletion stability requires something genuinely power-system-specific.

6.2. The Two-Block Convolution Lemma. We overcome this with the following convolution lemma, converting asymptotic bounded gaps into an exact continuous interval.

Lemma 6.1 (Two-Block Convolution). *Let P, Q be disjoint finite sets of positive integers. Suppose:*

- (1) $\Sigma(P)$ intersects every interval of length H inside $[L_P, U_P]$.
- (2) $\Sigma(Q)$ contains an exact interval $[L_Q, U_Q]$ with $U_Q - L_Q \geq H$.

Then $\Sigma(P \cup Q)$ contains the full continuous interval $[L_P + L_Q + H, U_P + U_Q - H]$.

Proof. For any $N \in [L_P + L_Q + H, U_P + U_Q - H]$, the window $[N - U_Q, N - L_Q]$ has length $\geq H$, and lies within $[L_P, U_P]$. Since $\Sigma(P)$ is H -syndetic, there exists $s \in \Sigma(P) \cap [N - U_Q, N - L_Q]$. Thus $N - s \in [L_Q, U_Q] \subseteq \Sigma(Q)$, implying $N = s + (N - s) \in \Sigma(P) + \Sigma(Q) \subseteq \Sigma(P \cup Q)$. $\square$

6.3. The Global Bottleneck. Because $\Sigma(\mathcal{P}_A(k))$ is globally H_D -syndetic, a sufficiently large finite prefix P provides syndeticity over a massive range. If we can manufacture just one disjoint finite block Q whose subset sums form an interval of length H_D , Lemma 6.1 amplifies it into a massive central seed. The global problem thus reduces to a strictly local, finite-length bottleneck:

Conjecture 1 (Fixed-Length Interval Block Lemma). *Let $A \subseteq \mathbb{Z}_{\geq 3}$ satisfy $\gcd(A) = 1$ and $R(A) \geq 1$. For every fixed H , every $k \geq 1$, and every lower cutoff M , there exists a finite block $Q \subseteq \mathcal{P}_A(k) \cap [M, \infty)$ such that $\Sigma(Q)$ contains $H + 1$ consecutive integers.*

This sharply isolates the missing ingredient. We can easily construct an arbitrarily late block Q that achieves full residue coverage modulo m (since $\gcd(A) = 1$ implies high powers eventually generate $\mathbb{Z}/m\mathbb{Z}$). The remaining analytical task is simply a finite balancing problem: shifting these covered residues into a single continuous finite interval.

7. THE 5-PART FULL-PROOF ARCHITECTURE

The route to the full generic proof of Erdős Problem 124 is now rigorously separated into five distinct structural pillars:

7.1. Theorem 1: Base-2 Theorem. If $2 \in A$ and $\gcd(A) = 1$, then $\mathcal{P}_A(k)$ is cofinite for every $k \geq 1$. Completely solved unconditionally via the modular residue argument (Section 2).

7.2. Theorem 2: Finite-Deletion Syndeticity. For any A with $R(A) \geq 1$, the $k \geq 1$ tail $\Sigma(\mathcal{P}_A(k))$ has bounded gaps with a maximum gap of $H_D = \sum_{a \in A} \sum_{e=0}^{k-1} a^e$. Completely solved.

7.3. Theorem 3: Two-Block Convolution. Bounded gaps combined with a single finite interval block of length H_D deterministically produces a massive central interval. Completely solved (Lemma 6.1).

7.4. Theorem 4: Strict-Surplus Closure. Once a central interval exists, $R(A) > 1$ ensures immediate unconditional closure via the explicit strict-surplus threshold $T_C(A, k)$. Completely solved (Section 4).

7.5. Main Remaining Lemma: The Fixed-Length Interval Block. Proving Conjecture 6.2 (the generation of an H_D -length interval arbitrarily far out) guarantees the required central seed. For the equality boundary ($R(A) = 1$), a final effective S-unit/Baker-type bounding handles the finite Diophantine hazards across the required block. Thus, the full resolution of Erdős Problem 124 reduces surgically to proving the Fixed-Length Interval Block Lemma.

REFERENCES

- [1] P. Erdős, *On the representation of an integer as the sum of k -th powers*, J. London Math. Soc. **14** (1939), 250–254.
- [2] S. A. Burr, P. Erdős, R. L. Graham, and S. Li, *Sums of distinct powers of 3, 4 and 7*, (citation available via MathOverflow).