



Dashboard

My Courses

Test Series

Forum

Live Class

My Tasks

My Progress

My Notes

JEE ADVANCED (2020-22)

JEE MAIN (2020-22)

3 Years 2021-24 EG

JEE Advanced-2021-23

JEE Main-2021-23

ENG G 2021-23 RMO

IJSO 2021-23

2022-24 JEE ADVANCED

Active

0

Upcoming

0

Attempted

3

Missed

4

2022-24 JEE ADVANCED - Attempted

Score Card

Answer Key

Comparison Chart

Questions List

Accuracy

Time Management

More...

You scored 155 out of 300 correctly.

51.67%

Question Results

You scored 4 of 4

This section contains 25 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which ONLY ONE option can be correct. Marking scheme: +4 for correct answer, 0 if not attempted and –1 in all other cases.

Q1. The equation of path of projectile is  $y = 0.5x - 0.04x^2$ . The initial speed of projection is ( $g = 10$  m/s<sup>2</sup>)

Options:

	10 m/s
	15 m/s
✓	12.5 m/s
	7.5 m/s

Solution :

(C)

$$y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta} \Rightarrow \tan \theta = 0.5 \Rightarrow \cos \theta = \frac{2}{\sqrt{5}}$$

Also,  $\frac{g}{2u^2 \times \frac{4}{5}} = 0.04 \Rightarrow y = 12.5 \text{ m/s}$

https://online.digitalpace.in/test\_collection/1492297/Attempted/quiz/1556770


1/51

Add to Favorites

You scored 0 of 4 skipped

**Q2. A small ball rolls off the top of a stairway with a velocity of 4.5 m/s. Each step is 0.2 m high and 0.3 m wide. If  $g$  is  $10 \text{ m/s}^2$ , then ball will strike the  $n^{\text{th}}$  step. (Assume ball strikes at edge of step)  $n = ?$**

Options:

	10
	9
	8
	11


**Solution :**  
(B)  
 $x = 0.3n; y = -0.2n$   
Equation of trajectory :  $y = -\frac{gx^2}{2u^2} \Rightarrow n = 9$

Add to Favorites

You scored 4 of 4

**Q3. A projectile from ground just crosses the top of two poles of equal height, after 1 sec and 3 sec from projection. The time of flight is**

Options:

	2 sec
	6 sec
	8 sec
	4 sec

**Solution :**

(D)  
Time of flight = 1 + 3 = 4 sec (using symmetry)

Add to Favorites

You scored 4 of 4

Q4. The equation of trajectory of a projectile is  $y = x - 0.2x^2$ . Angle of projection with horizontal is

Options:

	$\tan^{-1}\left(\frac{1}{2}\right)$
✓	45°
	30°
	$\tan^{-1}(2)$

Solution :


(B)

$$y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta} \Rightarrow \tan \theta = 1$$

Add to Favorites

You scored 0 of 4 skipped

Q5. In the given v-t graph, the distance travelled by the body in 5 sec will be



Options:

✓	100 m
---	-------

	80 m
	40 m
	20 m

**Solution :**  
(A)  
Distance travelled = area under the v-t curve  
$$= \frac{20 \times 2}{2} + 20 \times 2 + 20 \times 1 + \frac{20 \times 1}{2} + \frac{20 \times 1}{2} = 100 \text{ m}$$

Add to Favorites

You scored 4 of 4

**Q6. A body when projected vertically up, covers a total distance  $D$  during its time of flight. If there were no gravity, the distance covered by it during the same time is equal to**

Options:

	0
	$D$
✓	$2D$
	$4D$

**Solution :**  
(C)  
The displacement of the body during the time  $t$  as it reaches the point of projection  
$$\Rightarrow S = 0 \Rightarrow v_0 t - \frac{1}{2} g t^2 = 0 \Rightarrow t = \frac{2v_0}{g}$$
  
During the same time  $t$ , the body moves in absence of gravity through a distance  
 $D' = v.t$ , because in absence of gravity  $g = 0$   
$$\Rightarrow D' = v_0 \left( \frac{2v_0}{g} \right) = \frac{2v_0^2}{g} \quad \dots(1)$$
  
In presence of gravity, the total distance covered is

$$= D = 2H = 2 \frac{v_0^2}{2g} = \frac{v_0^2}{g} \qquad \dots(2)$$
$$(1) \div (2) \Rightarrow D' = 2D.$$

Add to Favorites

You scored 0 of 4

skipped

**Q7. A particle is projected from a point A with a velocity  $v$  at an angle  $\theta$  (upward) with the horizontal. At a certain point B, it moves at right angle to its initial direction. It follows that**


Options:

	velocity of the particle at B is $v$ .
	velocity of the particle at B is $v \cos \theta$ .
	velocity of the particle at B is $v \tan \theta$ .
✔	the time of flight from A to B is $\frac{v}{g \sin \theta}$ .

**Solution :**

(D)

$$\vec{v} = \vec{u} + \vec{a}t$$



Considering along the line AC

$$0 = v - g \sin \theta t \Rightarrow t = \frac{v}{g \sin \theta}$$

Now, consider along the line CB

$$v' = 0 + g \cos \theta \frac{v}{g \sin \theta} = v \cot \theta$$


Add to Favorites

You scored 0 of 4

skipped

**Q8. If a boat can have a speed of 4 km/hr in still water, for what values of speed of river flow, it can be managed to row boat right across the river, without any drift?**

Options:

	$\geq 4$ km/hr
	less than 4 km/hr
	only 4 km/hr
	none of these

**Solution :**

(B)

Drift  $(\Delta x) = (v_{b,x})\Delta t = (v_{b,r} \cos \theta + v_r)\Delta t$

Where  $v_{b,x}$  = velocity of boat w.r.t. ground

$v_{\perp,r}$  = velocity of boat w.r.t. ground

$v_r$  = velocity of river w.r.t. ground

For  $\Delta x = 0$ ,  $v_r = -v_{br} \cos \theta$

$\Rightarrow (v_r)_{\max} = v_{br}$

For,  $v_r > v_{br}$  we can not have zero drift.

Add to Favorites

You scored 0 of 4 skipped

**Q9. A swimmer crosses a river of width  $d$  flowing at velocity  $v$ . While swimming, he keeps himself always at an angle of  $120^\circ$  with the river flow and on reaching the other end he finds a drift of**

**$\frac{d}{2}$  in the direction of flow of river. The speed of the swimmer with respect to the river is**

Options:

	$(2 - \sqrt{3})v$
	$2(2 - \sqrt{3})v$
	$4(2 - \sqrt{3})v$

	$(2 + \sqrt{3})v$
--	-------------------

**Solution :**

(C)

$$\text{Drift} = \frac{d}{2} = \frac{(V_r - V_z \sin 30)d}{V_z \cos 30}$$

$$\Rightarrow V_z = 4(2 - \sqrt{3})V$$

Add to Favorites

You scored 0 of 4 skipped

**Q10. A projectile is thrown so as to have the maximum possible horizontal range equal to 400 m. Taking the point of projection as the origin, the coordinates of the point where the velocity of the projectile is minimum, are**

Options:

	(400, 100)
✓	(200, 100)
	(400, 200)
	(200, 200)

**Solution :**

(B)

When the horizontal range is maximum, the maximum height attained is  $\frac{R}{4} = 100\text{m}$ .

The velocity of the projectile is minimum at the highest point.

∴ Required point is (200, 100).


Add to Favorites

You scored 0 of 4 skipped

**Q11. A driver applies brakes on seeing a traffic signal 400 m ahead. At the time of applying the brakes the vehicle was moving with 15 m/s and retarding with 0.3 m/s<sup>2</sup>. The distance of**

vehicle after 1 min from the traffic light is

Options:

	25 m
	375 m
	360 m
	40 m

Solution :

(A)

The maximum distance covered by the vehicle before coming to rest  $= \frac{v^2}{2a} = \frac{(15)^2}{2(0.3)} = 375\text{ m}$

The corresponding time  $= t = \frac{v}{a} = \frac{15}{0.3} = 50\text{ sec}$

$\therefore$  The distance of the vehicle from the traffic signal after one minute  $= 400 - 375 = 25\text{ m}$


Add to Favorites

You scored 0 of 4

skipped

Q12. A man wishes to cross a river in a boat. If he crosses the river in minimum time he takes 10 minutes with a drift of 120 m. If he crosses the river taking shortest route, he takes 12.5 minutes. Find velocity of the boat with respect to water.


Options:

	20 m/min
	12 m/min
	10 m/min
	8 m/min

Solution :

(A)





$10 = \frac{L}{v} \qquad \dots(i)$

$12.5 = \frac{L}{\sqrt{v^2 - u^2}} = \frac{L}{v\sqrt{1 - u^2/v^2}} \qquad \dots(ii)$

From (i) and (ii),

$\frac{1}{12.5} = \frac{L}{v} \times \frac{v\sqrt{1 - u^2/v^2}}{L}$

$\frac{4}{5} = \sqrt{1 - \frac{12^2}{v^2}}$

$\frac{16}{25} = 1 - \frac{12^2}{v^2} \Rightarrow \frac{12^2}{v^2} = 1 - \frac{16}{25} = \frac{9}{25}$

$\frac{12}{v} = \frac{3}{5} \Rightarrow v = \frac{12 \times 5}{3} = 20 \text{ m/s}$

Add to Favorites

You scored 0 of 4 skipped

**Q13. The acceleration of a particle is increasing linearly with time  $t$  as  $bt$ . The particle starts from the origin with an initial velocity  $v_0$ . The distance travelled by the particle in time  $t$  will be**

Options:

✓	$v_0t + \frac{1}{6}bt^3$
	$v_0t + \frac{1}{3}bt^3$
	$v_0t + \frac{1}{3}bt^2$
	$v_0t + \frac{1}{2}bt^2$

**Solution :**

(A)

Given, acceleration  $a = bt$ 

$$\Rightarrow \frac{dv}{dt} = bt \Rightarrow v = \frac{bt^2}{2} + c$$

$$\text{At } t = 0, v = v_0 \Rightarrow c = v_0$$

$$\text{So, } v = \frac{bt^2}{2} + v_0$$

$$\Rightarrow \frac{ds}{dt} = \frac{bt^2}{2} + v_0$$


$$\Rightarrow s = \frac{bt^3}{6} + v_0 t$$

Add to Favorites

You scored 0 of 4

skipped

**Q14. A lift starts from rest. Its acceleration is plotted against time. When it comes to rest its height above its starting point is**



Options:

	20 m
✓	64 m
	32 m
	36 m

**Solution :**

(B)

**At 4 s**

$$u = at = 8 \text{ m/s}$$

$$s_1 = \frac{1}{2}at^2 = \frac{1}{2} \times 2 \times 4^2 = 16 \text{ m}$$

From 4 s to 8 s

$$a = 0, v = \text{constant} = 8 \text{ m/s}$$

$$s_2 = 8 \times 4 = 32 \text{ m}$$

$$u_t = \frac{at^2}{2}$$

From 8s to 12 s

$$s_3 = s_1 = 16 \text{ m}$$


$$\therefore s_{\text{Total}} = s_1 + s_2 + s_3 = 64 \text{ m}$$

Add to Favorites

You scored 0 of 4 skipped

**Q15. Two objects are moving along the same straight line. They cross a point A with an acceleration  $a, 2a$  and velocity  $2u, u$  at time  $t = 0$ . The distance moved by the object when one overtakes the other is**

Options:

	$\frac{6u^2}{a}$
	$\frac{2u^2}{a}$
	$\frac{4u^2}{a}$
	$\frac{8u^2}{a}$

**Solution :**

(A)

At the time of overtaking,

$$s_1 = s_2$$


$$\therefore 2ut + \frac{1}{2}at^2 = ut + \frac{1}{2}(2a)t^2$$

$$\therefore t = \frac{2u}{a}$$
$$\therefore s_1 \text{ (or } s_2) = (2u) \left( \frac{2u}{a} \right) + \frac{1}{2}(a) \left( \frac{2u}{a} \right)^2$$
$$= \frac{6u^2}{a}$$

Add to Favorites

You scored 4 of 4

**Q16. The figure shows velocity-time graph of a particle moving along a straight line. Identify the correct statement.**



Options:

	The particle starts from the origin
✓	The particle crosses its initial position at $t = 2\text{ s}$
	The average speed of the particle in the time interval, $0 \leq t \leq 2\text{ s}$ is zero
	All of the above

**Solution :**  
(B)  
 $s$  = net area of  $v$ - $t$  graph  
At  $2\text{ s}$ , net area =  $0$   
 $\therefore s = 0$   
and the particle crosses its initial position.

Add to Favorites

You scored 0 of 4

skipped

**Q17. A helicopter is rising vertically up with a velocity of 5 m/s. A ball is projected vertically up from the helicopter with a velocity  $V$  (relative to the ground). The ball crosses the helicopter 3 second after its projection. The value of  $V$  (in m/s) is**

Options:

<input type="radio"/>	10
<input type="radio"/>	15
<input checked="" type="radio"/>	20
<input type="radio"/>	5

Add to Favorites

You scored 0 of 4

skipped

**Q18. If the displacement of a particle varies with time as  $\sqrt{x} = t + 3$**

Options:

<input type="radio"/>	velocity of the particle is inversely proportional to $t$
<input checked="" type="radio"/>	velocity of particle varies linearly with $t$
<input type="radio"/>	velocity of particle is proportional to $\sqrt{t}$
<input type="radio"/>	initial velocity of the particle is zero


**Solution :**  
(B)  
 $\sqrt{x} = t + 3$   
 $\therefore x = (t + 3)^2$   
or  $v = \frac{dx}{dt}$   
 $= 2(t + 3)$   
 $\therefore v-t$  equation is linear.

Add to Favorites

You scored 0 of 4

skipped

Q19. The graph describes an airplane’s acceleration during its take-off run. The airplane’s velocity when it lifts off at  $t = 20\text{ s}$  is



Options:

	40 ms <sup>-1</sup>
	50 ms <sup>-1</sup>
✓	90 ms <sup>-1</sup>
	180 ms <sup>-1</sup>

**Solution :**  
(C)  
 $\Delta v = v_f - v_i = \text{area under } a\text{-}t \text{ graph}$   
 $v_i = 0$   
 $\Rightarrow v_f = \text{area}$   
 $= 40 + 50$   
 $= 90\text{ m/s}$

Add to Favorites

You scored 0 of 4

skipped

Q20. A particle moving in a straight line has velocity-displacement equation as  $v = 5\sqrt{1+s}$ . Here  $v$  is in ms<sup>-1</sup> and  $s$  in metres. Select the correct alternative.

Options:

	Particle is initially at rest
✓	Initially velocity of the particle is 5 m/s and the particle has a constant acceleration of 12.5 ms <sup>-2</sup>

	Particle moves with a uniform velocity
	None of the above

**Solution :**  
(B)  
 $v^2 = 25 + 25 s$   
or  $v^2 = (5)^2 + 2(12.5)s$   
Now compare with  $v^2 = u^2 + 2as$

Add to Favorites

You scored 0 of 4

skipped

**Q21. From the top of a long smooth incline a body *A* is projected along (maintaining contact with) the surface with speed *u*. Simultaneously, another small object *B* is thrown horizontally with velocity *v* = 10 m/s for the same point. Body *B* hits projected (in m/s)**

Options:


✓	8
	4
	10
	5

Add to Favorites


You scored 0 of 4

skipped

**Q22. A particle is moving along a straight line whose velocity-displacement graph is as shown in the figure. What is the magnitude of acceleration when displacement is 3 m?**



Options:

	$4\sqrt{3} \text{ ms}^{-2}$
	$3\sqrt{3} \text{ ms}^{-2}$
	$\sqrt{3} \text{ ms}^{-2}$
	$\frac{4}{\sqrt{3}} \text{ ms}^{-2}$

Solution :


(A)

$$a = v \cdot \frac{dv}{ds} = (4)(-\tan 60^\circ)$$
$$= -4\sqrt{3} \text{ m/s}^2$$


Add to Favorites

You scored 4 of 4

Q23. Annop (A) hits a ball along the ground with a speed  $u$  in a direciton which makes an angle  $30^\circ$  with the line joining him and the fielder Babul (B). Babul runs to intercept the ball with a speed  $\frac{2u}{3}$ . At what angle  $\theta$  should he run to intercept the ball?



Options:

	$\sin^{-1} \left[ \frac{\sqrt{3}}{2} \right]$
	$\sin^{-1} \left[ \frac{2}{3} \right]$
	$\sin^{-1} \left[ \frac{3}{4} \right]$



$$\sin^{-1}\left[\frac{4}{5}\right]$$

**Solution :**  
(C)

Relative velocity of  $A$  with respect to  $B$  should be along  $AB$  or absolute velocity components perpendicular  $AB$  should be same.

$$\therefore \frac{2u}{3} \sin \theta = u \sin 30^\circ$$

$$\therefore \theta = \sin^{-1}\left(\frac{3}{4}\right)$$

Add to Favorites

You scored 4 of 4

**Q24.** A car is travelling on a straight road. The maximum velocity the car can attain is  $24 \text{ ms}^{-1}$ . The maximum acceleration and deceleration it can attain are  $1 \text{ ms}^{-2}$  and  $4 \text{ ms}^{-2}$  respectively. The shortest time the car takes from rest to rest in a distance of 200 m is,

Options:



22.4 s


30 s

11.2 s

5.6 s

**Solution :**  
(A)

Deceleration is four times. Therefore, deceleration time should be  $\frac{1}{4}$ th.



$v_{\max} = (a_1)(4t) = (1)(4t) = 4t$

Area of v-t graph = displacement

$\therefore 200 = \frac{1}{2}(5t)(4t)$

or  $t = \sqrt{20} \text{ s}$

Total journey time =  $5t = 22.4 \text{ s}$

Add to Favorites

You scored 0 of 4 skipped

**Q25.** A car is travelling on a road. The maximum velocity the car can attain is  $24 \text{ ms}^{-1}$  and the maximum deceleration is  $4 \text{ ms}^{-2}$ . If car starts from rest and comes to rest after travelling 1032 m in the shortest time of 56 s, the maximum acceleration that the car can attain is


Options:

	$6 \text{ ms}^{-2}$
✓	$1.2 \text{ ms}^{-2}$
	$12 \text{ ms}^{-2}$
	$3.6 \text{ ms}^{-2}$

**Solution :**  
(B)

Area of v-t graph = displacement

$\therefore 1032 = \frac{1}{2}(56 + t_0)(24) \text{ or } t_0 = 30 \text{ s}$



Deceleration time  $t_1 = \frac{24}{4} = 6 \text{ s}$

$\therefore$  Acceleration time  $t_2 = 56 - t_0 - t_1 = 20 \text{ s}$

$\therefore \text{Acceleration} = \frac{24}{20} = 1.2 \text{ m/s}^2$


Add to Favorites

You scored 4 of 4

This section contains 25 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which ONLY ONE option can be correct. Marking scheme: +4 for correct answer, 0 if not attempted and –1 in all other cases.

Q26. Rakesh needs 1.71 g of sugar ( $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ ) to sweeten his tea. What would be the number of carbon atoms added in his tea?

Options:


	$3.6 \times 10^{22}$
	$7.2 \times 10^{21}$
	$0.05 \times 10^{23}$
	$6.6 \times 10^{22}$

**Solution :**  
(A)  
Molar mass of  $\text{C}_{12}\text{H}_{22}\text{O}_{11}$  = 342 g  
342 g sugar has = 12 N atoms of C  
 $\therefore$  1.71 g sugar has =  $\frac{12 \times 6.02 \times 10^{23} \times 1.71}{342}$  atoms  
 $= 3.6 \times 10^{22}$  atoms

Add to Favorites

You scored 4 of 4

Q27. The percentage of oxygen in NaOH is:  
Options:

	40
	16
	8
	1


**Solution :**  
(A)  
Per cent of oxygen in NaOH =  $\frac{16 \times 100}{40} = 40$ .

Add to Favorites

You scored 4 of 4

**Q28. One mole of P<sub>4</sub> molecules contain :**

**Options:**

	1 molecule
	4 molecule
	$\frac{1}{4} \times 6.022 \times 10^{23}$ atoms
	$24.088 \times 10^{23}$ atoms


**Solution :**  
(D)  
1 mole P<sub>4</sub> = N molecules of P<sub>4</sub> = 4 N atoms of P<sub>4</sub>.

Add to Favorites

You scored 4 of 4

**Q29. 10<sup>24</sup> molecules of solute are dissolved in 10<sup>25</sup> molecules of solvent, the mole fraction of solute in solution are :**

Options:

	0.09
	0.08
	0.07
	0.05

**Solution :**

(A)

Mole fraction =  $\frac{n_1}{n_1 + n_2} = \frac{\text{Molecules of solute}}{\text{Total molecules of solute and solvent}}$


$= \frac{10^{24}}{10^{24} + 10^{25}} = \frac{1}{11} = 0.09$

Add to Favorites

You scored 4 of 4


**Q30. 0.50 mole of BaCl<sub>2</sub> is mixed with 0.20 mole of Na<sub>3</sub>PO<sub>4</sub>, the maximum number of moles of Ba<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> that can be formed is:**

Options:

	0.70
	0.50
	0.20
	0.10

**Solution :**

(D)



Add to Favorites

You scored 4 of 4

**Q31. A molal solution in one that contains one mole of solute in :**

Options:

	1000 g of the solvent
	1 litre of the solvent
	1 litre of solution
	22.4 litre of solution

**Solution :**

(A)


$$\text{Molality} = \frac{\text{Moles of solute}}{\text{wt. of solvent in kg}}$$

Add to Favorites

You scored 4 of 4

**Q32. The molarity of a solution obtained by mixing 750 mL of 0.5 M HCl with 250 mL of 2 M HCl will be:**

Options:

	1.75 M
	0.975 M
	0.875 M
	1.00 M

**Solution :**

(C)

milli mole of solution I =  $750 \times 0.5 = 375$

milli mole of solution II =  $250 \times 2 = 500$

total milli mole in mixture = 375 + 500 = 875

$\therefore \text{Molarity} = \frac{875}{1000} = 0.875 \text{ M}$

Add to Favorites

You scored 4 of 4

**Q33. The least number of molecules are contained in:**

Options:

<input type="radio"/>	2 g hydrogen
<input type="radio"/>	8 g hydrogen
<input checked="" type="radio"/>	4 g nitrogen
<input type="radio"/>	16 g CO <sub>2</sub>

**Solution :**  
(C)

Mole of N<sub>2</sub> is  $= \frac{4}{28} = \frac{1}{7}$  (the lowest value)


Add to Favorites

You scored 4 of 4

**Q34. How many moles of electrons weigh one kilogram ?**

Options:

<input type="radio"/>	$6.023 \times 10^{23}$
<input type="radio"/>	$\frac{1}{9.108} \times 10^{31}$
<input type="radio"/>	$\frac{6.023}{9.108} \times 10^{34}$


	$\frac{1}{9.108 \times 6.023} \times 10^8$
---	--

Add to Favorites

You scored 4 of 4

Q35. The number of atoms in 558.5 g of Fe (at. Wt. 55.85) is

Options:

	twice that in 60 g carbon
	$6.022 \times 10^{22}$
	half in 8 g He
	$558.5 \times 6.023 \times 10^{23}$

Solution :

(A)

$558.5 \text{ g Fe} \times \frac{1 \text{ mole Fe}}{55.85 \text{ g}} = 10 \text{ mole Fe}$


$= 2 \times 5 \text{ mole C} = 2 \times 60 \text{ g C}$

Add to Favorites

You scored 4 of 4

Q36.  $6.02 \times 10^{20}$  molecules of urea are present in 100 mL of its solution. The molarity of urea solution is:

Options:

	0.1
	0.01
	0.02
	0.001



**Solution :**

(B)

$$M = \frac{\text{moles of urea}}{\text{volume in litre}} = \frac{6.02 \times 10^{20}}{6.02 \times 10^{23} \times \frac{100}{1000}} = 0.01 \text{ M}$$

Add to Favorites

You scored 4 of 4

**Q37. Density of a 2.05 M solution of acetic acid in water is 1.02 g/mL. The molality of the solution is:**

Options:

	1.14 mol kg <sup>-1</sup>
	3.28 mol kg <sup>-1</sup>
✓	2.28 mol kg <sup>-1</sup>
	0.44 mol kg <sup>-1</sup>

**Solution :**

(C)

$$m = \frac{\text{moles of CH}_3\text{COOH}}{\text{wt. of solvent in kg}} = \frac{2.05 \times 1000}{897} = 2.285$$

Wt. of solvent = wt. of solution – wt. of solute

$$= [1000 \times 1.02 - 2.05 \times 60] = 897 \text{ g}$$


Add to Favorites

You scored 4 of 4

**Q38. A gaseous hydrocarbon given upon combustion 0.72 g of water and 3.08 g of CO<sub>2</sub>. The empirical formula of the hydrocarbon is :**


Options:

	C <sub>6</sub> H <sub>5</sub>
--	-------------------------------

	C <sub>7</sub> H <sub>8</sub>
	C <sub>2</sub> H <sub>4</sub>
	C <sub>3</sub> H <sub>4</sub>

**Solution :**

(B)

Let the formula of hydrocarbon be C<sub>a</sub>H<sub>b</sub>

$$\text{mole of } CO_2 \text{ (a) formed} = \frac{3.08}{44} = 0.07$$

$$\text{mole of } H_2O \text{ formed } \left(\frac{b}{2}\right) = \frac{0.72}{18} = 0.04$$

$$\therefore \frac{a}{b/2} = \frac{0.07}{0.04}$$

$$\text{or } \frac{a}{b} = \frac{0.07}{0.08} = \frac{7}{8}$$

 $\therefore$  mole ratio of C and H :: 7 : 8Thus empirical formula is C<sub>7</sub>H<sub>8</sub>.

Add to Favorites

You scored 4 of 4

**Q39. Atomic mass of Ne is 20.2. Ne is a mixture of Ne<sup>20</sup> and Ne<sup>22</sup>. Relative abundance of heavier isotope is:**

Options:

	90
	20
	40
	10

**Solution :**

(D)

Average isotopic wt. =  $\Sigma \% \times \text{isotopic wt.}$ 

$$= \frac{(\text{per cent} \times \text{wt. of isotope}) + (\text{per cent} \times \text{wt. of other isotope})}{100}$$

$$\therefore 20.2 = \frac{a \times 20 + (100 - a) \times 22}{100}$$

$$\therefore a = 90$$

$$\text{Per cent of heavier isotope} = 100 - 90 = 10$$

Add to Favorites

You scored 4 of 4

**Q40. If the series limit of wavelength of the Lyman series for the hydrogen atom is 912 Å, then the series limit of wavelength for the Balmer series of the hydrogen atom is:**

Options:

	912 Å
	$912 \times 2 \text{ Å}$
✓	$912 \times 4 \text{ Å}$
	$912/2 \text{ Å}$

**Solution :**

(C)

$$\frac{1}{\lambda_{\text{Lyman}}} = R_H \left[ \frac{1}{1^2} - \frac{1}{\infty^2} \right] = R_H$$

$$\frac{1}{\lambda_{\text{Balmer}}} = R_H \left[ \frac{1}{2^2} - \frac{1}{\infty^2} \right] = \frac{R_H}{4}$$


$$\frac{\lambda_{\text{Balmer}}}{\lambda_{\text{Lyman}}} = 4 \text{ or } \lambda_B = 4 \times 912 \text{ Å}$$

Add to Favorites

You scored 4 of 4

**Q41. The difference in angular momentum associated with the electron in two successive orbits of hydrogen atom is:**

Options:

	$h/\pi$
	$h/2\pi$
	$h/2$
	$(n-1)h/\pi$

**Solution :**

(B)

Angular momentum for  $n$  and  $(n+1)$  shells are  $\frac{nh}{2\pi}$  and  $(n+1)\frac{h}{2\pi}$ .


Thus, difference in angular momentum of two successive orbits is  $(n+1)\frac{h}{2\pi} - \frac{nh}{2\pi} = \frac{h}{2\pi}$

Add to Favorites

You scored 4 of 4

**Q42. The longest  $\lambda$  for the Lyman series is ..... (Given  $R_H = 109678 \text{ cm}^{-1}$ ):**

Options:

	1215 Å
	1315 Å
	1415 Å
	1515 Å

**Solution :**

(A)

For longest  $\lambda$  of Lyman series  $n_1 = 1$  and  $n_2 = 2$ ,

$$\frac{1}{\lambda} = R_H \left[ \frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

Because  $\Delta E = \frac{hc}{\lambda}$  is minimum when  $\lambda$  is longest.

Thus,  $\Delta E = E_2 - E_1$ .

Thus,  $\frac{1}{\lambda} = R_H \times \left[ \frac{1}{1^2} - \frac{1}{2^2} \right] = \frac{3}{4} \times 109678$


$\therefore \lambda = 1.215 \times 10^{-15} \text{ cm} = 1215 \text{ \AA}$

Add to Favorites

You scored 4 of 4

**Q43. A ball of mass 200 g is moving with a velocity of  $10 \text{ m sec}^{-1}$ . If the uncertainty of velocity is 0.1%, the uncertainty in it position is :**

Options:

	$3.3 \times 10^{-31} \text{ m}$
	$3.3 \times 10^{-27} \text{ m}$
	$5.3 \times 10^{-25} \text{ m}$
	$2.64 \times 10^{-32} \text{ m}$

**Solution :**

(D)

$$\Delta u = \frac{0.1}{100} \times 10 = 10^{-2} \text{ m sec}^{-1}; \text{ Now } \Delta u \cdot \Delta X = \frac{h}{4\pi m}$$

$$\therefore \Delta X = \frac{6.625 \times 10^{-34}}{4 \times 10^{-2} \times 3.14 \times 200 \times 10^{-3}} = 2.64 \times 10^{-32} \text{ m}$$

Add to Favorites

You scored 4 of 4

**Q44. In absence of Pauli exclusion principle, the electronic configuration of Li in ground state may be:**

Options:

	$1s^2, 2s^1$
✓	$1s^3$
	$1s^1, 2s^2$
	$1s^2, 2s^1 2p^1$

**Solution :**  
(B)  
All the three electrons are to be kept in 1s.

Add to Favorites

You scored -1 of 4

**Q45. Which one is odd one?**  
Options:

	$4f$
	$5d$
✓	$3s$
✗	$6p$

**Solution :**  
(C)  
( $n + l$ ) for rest all is = 7;  
For 3s it is  $3 + 0 = 3$ .

Add to Favorites

You scored 4 of 4

**Q46. The ratio of the energy of the electron in ground state of hydrogen to the electron in first excited state of  $\text{Be}^{3+}$  is:**

Options:

	1 : 4
	1 : 8
	1 : 16
	16 : 1

Solution :

(A)

$$E_{2(\text{Be}^{2+})} = E_{2(\text{H})} \times Z^2; \text{ Also } E_{2(\text{H})} = \frac{E_{1(\text{H})}}{2^2}$$

$$\therefore E_{2(\text{Be}^{2+})} = \frac{E_{1(\text{H})}}{2^2} \times 4^2 = 4 \times E_{1(\text{H})}$$

Add to Favorites

You scored 4 of 4

Q47. If the radius of first Bohr orbit is  $x$  then de Broglie wavelength of electron in 3rd orbit is nearly:

Options:

	$2\pi x$
	$6\pi x$
	$9x$
	$\frac{x}{3}$

Solution :

(B)

$$r_n = r_1 \times n^2$$

$$r_3 = 3^2 x = 9x$$

$$\text{Also, } mvr_3 = \frac{3h}{2\pi}$$

or  $mu = 3 \frac{h}{2\pi \cdot 9x} = \frac{h}{6\pi x}$

or  $\lambda = \frac{h}{mu} = \frac{h \cdot 6\pi x}{h} = 6\pi x$

Add to Favorites

You scored 4 of 4

**Q48. How many electrons in an atom with atomic number 105 can have  $(n + l) = 8$  ?**

Options:

	30
✓	17
	15
	Unpredictable

**Solution :**

(B)

Electronic configuration of atom with atom no. 105 is :  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^6 4d^{10}, 5s^2 5p^6 5d^{10} \underline{5f^{14}}, 6s^2 6p^6 \underline{6d^3}, 7s^2$ .

The underlined orbitals have  $(n + l) = 8$ .

Add to Favorites

You scored 4 of 4

**Q49. The electronic configuration of  $_{46}\text{Pd}$  is :**

Options:

✓	$_{36}[\text{Kr}] 4d^{10}$
	$_{36}[\text{Kr}] 4d^8, 5s^2$
	$_{36}[\text{Kr}] 4d^9, 5s^1$



	$_{18}[\text{Kr}] 4d^{10}$
--	----------------------------


**Solution :**  
(A)  
Pd :  $[\text{Kr}]4d^{10}$ . An exception.

Add to Favorites

You scored 4 of 4

**Q50. Which electronic level would allow the hydrogen atom to absorb a photon but not to emit a photon?**

Options:

	3s
	2p
	2s
	1s

**Solution :**  
(D)  
Ground state of hydrogen atom, *i.e.*, 1s


Add to Favorites

You scored 4 of 4

**This section contains 25 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which ONLY ONE option can be correct. Marking scheme: +4 for correct answer, 0 if not attempted and –1 in all other cases.**

**Q51. If  $2^a = 3$  and  $9^b = 4$  then value of  $(ab)$  is -**

Options:

	1
	2
	3
	4


**Solution :**  
(A)  
 $a = \log_2 3$   
 $b = \log_3 4$   
 $\Rightarrow ab = \log_2 3 \log_3 2 = 1.$

Add to Favorites

You scored 4 of 4

**Q52. If  $\log_2 (4 + \log_3 (x)) = 3$  , then sum of digits of x is -**

**Options:**

	3
	6
	9
	18

**Solution :**  
(C)  
Let  $4 + \log_3 (x) = N \Rightarrow \log_2 N = 3 \Rightarrow N = 2^3$   
 $N = 4 + \log_3 x = 8$   
 $\therefore N = 8$   
 $\log_3 x = 4 \Rightarrow x = 81.$   
Sum of digits of x = 9.

Add to Favorites

You scored -1 of 4

**Q53. Sum of all the solution(s) of the equation  $\log_{10}(x) + \log_{10}(x + 2) - \log_{10}(5x + 4) = 0$  is -**

Options:

<input type="radio"/>	-1
<input checked="" type="radio"/>	3
<input checked="" type="radio"/>	4
<input type="radio"/>	5

**Solution :**

(C)

Using  $\log_{10} p + \log_{10} r - \log_{10} s = \log_{10} \left( \frac{pr}{s} \right)$

$$\log_{10} \left( \frac{x(x+2)}{5x+4} \right) = 0$$

$$\Rightarrow x^2 + 2x = 5x + 4$$

$$\Rightarrow x = 4, x = -1$$

(reject)

Add to Favorites

You scored 0 of 4

skipped

**Q54. The product of all the solutions of the equation  $x^{1+\log_{10} x} = 100000x$  is -**

Options:

<input type="radio"/>	10
<input type="radio"/>	$10^5$
<input type="radio"/>	$10^{-5}$



1

**Solution :**

(D)

$$x^{(1+\log_{10} x)} = 10^5 \cdot x$$

Taking log on both sides to base 10 :

$$(1 + \log_{10} x)(\log_{10} x) = 5 + \log_{10} x$$

$$\log_{10} x = t \Rightarrow t(1 + t) = 5 + t$$

$$\Rightarrow t^2 + t = 5 + t$$

$$\Rightarrow t = 5^{1/2} \text{ or } t = -5^{1/2}$$

$$\Rightarrow \log_{10} x = 5^{1/2}; \log_{10} x = -5^{1/2}$$

$$\Rightarrow x = 10^{5^{1/2}}; x = 10^{-5^{1/2}}$$

$$\text{Product} = 10^0 = 1$$

Add to Favorites

You scored 0 of 4

skipped

**Q55. If  $x_1$  and  $x_2$  are the roots of equation  $e^{3/2} \cdot x^{2 \ln x} = x^4$ , then the product of the roots of the equation is -**

**Options:**



$e^2$

e

$e^{3/2}$

$e^{-2}$

**Solution :**

(A)

Take log on both sides

$$\frac{3}{2} + 2(\ln x)^2 = 4 \ln x$$

$$t = \ln x \quad \dots (1)$$

$$2t^2 - 4t + \frac{3}{2} = 0 \begin{cases} t_1 \rightarrow t_1 \ln x_1 \\ t_2 \rightarrow t_2 \ln x_2 \end{cases}$$

$$t_1 + t_2 = 2$$

$$\ln x_1 + \ln x_2 = 2 \text{ (from Eqs. (1))}$$

$$\ln (x_1 x_2) = 2 \Rightarrow x_1 x_2 = e^2$$



Note:  $\log_e x = \ln x$ ; where ‘e’ is Napier’s constant. Its irrational quantity.

Add to Favorites

You scored -1 of 4

**Q56. Solution set of  $\sqrt{7x-6} < x$  is**

Options:

	$(6, \infty)$
	$(1, 6)$
	$\left[\frac{6}{7}, 6\right)$
	$\left[\frac{6}{7}, 1\right) \cup (6, \infty)$

Add to Favorites

You scored 4 of 4

**Q57. Number of positive integral solution of  $\frac{x^2(3x-4)^3(x-2)^4}{(x-5)^5(2x-7)^6} \leq 0$  is**

Options:

	4
	3
	2

	1
--	---


**Solution :**  
(B)  
Wavy curve method

Add to Favorites

You scored 0 of 4 skipped

**Q58.** If  $\log_a(1 - \sqrt{1+x}) = \log_{a^2}(3 - \sqrt{1+x})$ , then number of solutions of the equations is -

Options:

	0
	1
	2
	infinitely many

**Solution :**  
(A)  
Domain:  $1 - \sqrt{1+x} > 0$  &  $3 - \sqrt{1+x} > 0$   
 $\Rightarrow \sqrt{1+x} < 1$   
Put  $\sqrt{1+x} = t \Rightarrow t \geq 0$   
 $(1-t)^2 = 3-t$   
 $t^2 - t - 2 = 0$   
 $t = 2, -1$  (both rejected)  
 $\Rightarrow$  No real solution

Add to Favorites

You scored 0 of 4 skipped

**Q59.** The number of solution(s) of  $\sqrt{\log_3(3x^2) \cdot \log_9(81x)} = \log_9 x^3$  is -

Options:

	0
✓	1
	2
	3

**Solution :**

(B)

$$t = \log_3 x$$

$$\sqrt{(1+2t)\left(2+\frac{t}{2}\right)} = \frac{3}{2}t$$

$$t > 0$$

Squaring

$$2 + \frac{t}{2} + 4t + t^2 = \frac{9}{4}t^2$$

$$5t^2 - 18t - 8 = 0$$

$$5t^2 - 20t + 2t - 8 = 0$$

$$(5t + 2)(t - 4) = 0$$

$$\Rightarrow t = 4$$

$$\Rightarrow \log_3 x = 4$$

$$x = 81$$

Add to Favorites

You scored 0 of 4 skipped

**Q60.** If  $x_1$  and  $x_2$  are the two values of  $x$  satisfying the equation  $7^{2x^2} - 2(7^{x^2+x+12}) + 7^{2x+24} = 0$ , then  $(x_1 + x_2)$  equals -

Options:

	0
--	---

	1
	-1
	7


**Solution :**  
(B)  
 $7^{x^2} = a, 7^{x+12} = b$   
 $a^2 - 2ab + b^2 = 0$   
 $(a - b)^2 = 0$   
 $\Rightarrow a = b$   
 $\Rightarrow x^2 = x + 12$   
 $x^2 - x - 12 = 0$   
 $x^2 - 4x + 3x - 12 = 0$   
 $(x + 3)(x - 4) = 0$   
 $x = 4, -3.$

Add to Favorites

You scored 4 of 4

**Q61.** Solution set of  $\sqrt{\frac{x-3}{11-x}} \geq -1$  is

Options:

	(4, 10)
	(3, 11)
	$(-\infty, \infty)$
	[3, 11)

**Solution :**  
(D)





$$\frac{x-3}{11-x} \geq 0$$
$$\frac{x-3}{x-11} \leq 0$$

Add to Favorites

You scored -1 of 4

Q62. Number of integral values of  $x$  satisfying  $\log_{1/2}(x^2 - 5x + 6) \geq -1$  is

Options:

	4
	2
	6
	None

**Solution :**  
(B)  
 $0 < x^2 - 5x + 6 \leq 2$

Add to Favorites

You scored 4 of 4

Q63. If  $x = \log_2 \left( \sqrt{\sqrt{56 + \sqrt{56 + \sqrt{56 + \sqrt{56 + \dots \infty}}}}} \right)$ , then which of the following statements holds good ?

Options:

	$x < 0$
	$0 < x < 2$
	$2 < x < 4$

$$3 < x < 4$$

**Solution :**

(C)

$$x = \log_2 \sqrt{56 + \sqrt{56 + \sqrt{56 + \sqrt{56 + \dots \infty}}}}$$

$$2^x = \sqrt{56 + \sqrt{56 + \sqrt{56 + \dots}}}$$

$$2^x = \sqrt{56 + 2^x}$$

$$2^{2x} - 2^x = 56 \Rightarrow 2^{2x} - 2^x - 56 = 0$$

$$\text{Let } 2^x = t \quad \therefore t^2 - t - 56 = 0$$

$$t = 8 \text{ \& } t = -7$$

$$2^x = 8 \therefore 2^x = 2^3 \Rightarrow x = 3$$

$$\therefore 2 < x < 4$$

Add to Favorites

You scored 0 of 4

skipped

**Q64. Solution set of  $-1 \leq |x-5| < 2$  is****Options:**

[6, 9)

[4, 7)



(3, 7)

None

**Solution :**

(C)

$$-2 < x-5 < 2$$

$$3 < x < 7$$

Add to Favorites

You scored 0 of 4

skipped

Q65. The number  $\log_2 7$  is -

Options:

	an integer
	a rational number
✓	an irrational number
	a prime number

**Solution :**  
(C)

Let  $\log_2 7$  be rational then  $\log_2 7 = \frac{p}{q}$  ( $q \neq 0$ )

$$7 = 2^{p/q} \Rightarrow 2^p = 7^q$$

Here all power of 2 is even but 7 is odd.

$\therefore$  our assumption is wrong

$\therefore \log_2 7$  is irrational.

Add to Favorites

You scored 0 of 4

skipped

Q66. The number of integral solutions of  $|\log_3 x^2 - 4| = 2 + |\log_3 x - 3|$  is -

Options:


✓	1
	2
	3
	0

**Solution :**  
(A)

Case (1) :  $t < 2$

$$-2t + 4 = 2 - t + 3$$

$$t = -1$$



$$\Rightarrow x = \frac{1}{5}$$

Case (2) :  $2 \leq t < 3$

$$2t - 4 = 2 - t + 3$$

$$3t = 9$$

$$t = 3$$

$$\log_5 x = 3$$

Case (3) :  $t \geq 3$

$$2t - 4 = 2 + t - 3$$

$$t = 3$$

$$\log_5 x = 3$$

$$x = 125$$

Add to Favorites

You scored 0 of 4 skipped

Q67. Find the value of  $3^{\sqrt{\log_3 5}} - 5^{\sqrt{\log_5 3}}$  is

Options:

	$\sqrt{5} - \sqrt{3}$
	$\sqrt{2}$
	2
✓	0

Add to Favorites

You scored 0 of 4 skipped

Q68. Solution set for  $(0.5)^{1/x} \geq 0.0625$  is

Options:

	$[4, \infty)$
	$(-\infty, 0)$
	$(-\infty, 0) \cup [4, \infty)$
	None

Add to Favorites

You scored 0 of 4 skipped

Q69. If  $\alpha$  and  $\beta$  are the roots of the equation  $(\log_2 x)^2 + 4(\log_2 x) - 1 = 0$ , then the value of  $\alpha\beta$  equals

Options:

	$\frac{1}{8}$
	8
	16
	$\frac{1}{16}$

**Solution :**  
(D)  
 $(\log_2 x)^2 + 4(\log_2 x) - 1 = 0$   
Let  $\log_2 x = t$   
 $t^2 + 4t - 1 = 0$   
 $t = \frac{-4 \pm \sqrt{16 + 4}}{2}$   
 $t = -2 \pm \sqrt{5}$   
 $\log_2 x = -2 \pm \sqrt{5}$   
 $x = 2^{-2 \pm \sqrt{5}}$

$$\therefore \beta = 2^{-2-\sqrt{5}}$$

$$\alpha = 2^{-2+\sqrt{5}}$$


$$\alpha\beta = 2^{-4} = \frac{1}{16}$$

Add to Favorites

You scored 4 of 4

**Q70.** If  $\log_{0.3}(x-1) < \log_{0.09}(x-1)$ , then  $x$  lies in the interval

Options:

	$(2, \infty)$
	$(1, 2)$
	$(1, \infty)$
	none of these

**Solution :**

(A)

$$\log_{0.3}(x-1) < \frac{1}{2} \log_{0.3}(x-1)$$

$$\begin{array}{c} + \quad - \quad + \\ | \quad | \quad | \\ 1 \quad 2 \end{array} \quad x-1 > (x-1)^{1/2}$$

$$x^2 + 1 - 2x > x - 1$$

$$x^2 - 3x + 2 > 0 \Rightarrow (x-2)(x-1) > 0$$

$$x \in (-\infty, 1) \cup (2, \infty) \quad \dots(1)$$

$$\& \ x-1 > 0 \Rightarrow x > 1 \Rightarrow x \in (1, \infty) \quad \dots(2)$$

$$\therefore (1) \& (2)$$

$$\Rightarrow (2, \infty)$$

Add to Favorites

You scored 4 of 4

Q71.  $\sqrt{x^2 - 5x + 6} \leq \sqrt{x^2 + x + 1}$  solution set is ?

Options:

	$(-\infty, 2]$
	$\left[\frac{5}{6}, \infty\right)$
	$[3, \infty)$
	None

Solution :

(D)

$$-5x + 6 \leq x + 1 \qquad x \geq \frac{5}{6}$$

$$x^2 - 5x + 6 \geq 0 \quad x \in (-\infty, 2] \cup [3, \infty)$$

Intersection

$$\left[\frac{5}{6}, 2\right] \cup [3, \infty) \text{ Ans.}$$

Add to Favorites

You scored -1 of 4

Q72. The number of solutions of the equation  $\log_{x-3}(x^3 - 3x^2 - 4x + 8) = 3$  is equal to

Options:

	4
	3
	2
	1

Solution :

(D)

Domain:  $x - 3 > 0$ ,  $x - 3 \neq 1$  &  $x^3 - 3x^2 - 4x + 8 > 0$

$$x^3 - 3x^2 - 4x + 8 = (x - 3)^3$$

$$6x^2 - 31x + 35 = 0$$

$$x = \frac{31 \pm \sqrt{(31)^2 - 4(6)(35)}}{12}$$

$$\Rightarrow x = \frac{31 \pm 11}{12} \Rightarrow x = \begin{cases} \frac{42}{12} = \frac{7}{2} \\ \frac{20}{12} = \frac{5}{3} \end{cases}$$

$\therefore$  No. of solution = 1 as  $x = \frac{5}{3}$  is rejected

Add to Favorites

You scored 0 of 4

skipped

**Q73. Sum of the roots of the equation  $9^{\log_3(\log_2 x)} = \log_2 x - (\log_2 x)^2 + 1$  is equal to**

Options:

	2
	4
	6
	8

**Solution :**

(A)

Domain:  $\log_2 x > 0$  i.e.  $x > 1$

$$9^{\log_3(\log_2 x)} = \log_2 x - (\log_2 x)^2 + 1$$

$$3^{2\log_3(\log_2 x)} = \log_2 x - (\log_2 x)^2 + 1$$

$$(\log_2 x)^2 + (\log_2 x)^2 - (\log_2 x) - 1 = 0$$

$\therefore$  By property  $a^{\log_a} = 1$

Let  $\log_2 x = t$

$$\therefore 2t^2 - t - 1 = 0$$



$$2t(t-1)+1(t-1)=0$$

$$t=1, -\frac{1}{2} \Rightarrow \text{only } t=1 \text{ acceptable}$$

$$\Rightarrow \log_2 x = 1 \Rightarrow x = 2 \Rightarrow \text{Sum of roots} = 2$$

Add to Favorites

You scored 4 of 4

**Q74.** If  $x$  satisfies the inequality  $\log_{25} x^2 + (\log_5 x)^2 < 2$ , then  $x \in$

Options:

✓	$\left(\frac{1}{25}, 5\right)$
	$(1, 2)$
	$(4, 5)$
	$(0, 1)$

**Solution :**

(A)

$$\frac{1}{2} \log_5 x^2 + (\log_5 x)^2 - 2 < 0$$

 Let  $\log_5 x = t$

$$\log_5 x + (\log_5 x)^2 - 2 < 0$$

$$t^2 + t - 2 < 0$$

$$(t+2)(t-1) < 0$$

$$t \in (-2, 1)$$

$$\therefore \log_5 x \in (-2, 1)$$

$$\therefore x \in \left(\frac{1}{25}, 5\right)$$

[Add to Favorites](#)

You scored 4 of 4

**Q75.** If  $1, \log_9(3^{1-x} + 2)$  and  $\log_3(1 \cdot 3^x - 1)$  are in A.P. then  $x$  can be

Options:

	$\log_4 3$
	$\log_3 4$
✓	$1 - \log_3 4$
	$\log_3 0.25$

**Solution :**

(C)

$$2 \cdot \log_9(3^{1-x} + 2) = 1 + \log_3(4 \cdot 3^x - 1)$$

$$\frac{2}{2} \cdot \log_3 \left[ \frac{3}{3^x} + 2 \right] = \log_3 3 + \log_3(4 \cdot 3^x - 1)$$

$$\text{Let } 3^x = t \Rightarrow t > 0$$

$$\log_3 \left[ \frac{3}{t} + 2 \right] = \log_3 [3 \cdot (4 \cdot 3^x - 1)]$$

$$3 + 2t = 12t^2 - 3t$$

$$12t^2 - 5t - 3 = 0$$

$$(4t - 3)(3t + 1) = 0$$

$$t = \frac{3}{4}, \frac{-1}{3} \therefore 3^x = \frac{3}{4}$$

$$x \log 3 = \log 3 - \log 4$$

$$\Rightarrow x = 1 - \log_3 4$$

[Add to Favorites](#)

