



KEY, 55 Cancri e: single transit




Transit depth 331 ppm (0.033%), duration 1.65 h. Host star 55 Cancri A (G8V), radius 0.943 $R_\odot$.

KEY, 55 Cancri e: light curve over several orbits




Orbital period $P = 0.7365 \text{ d}$. Semi-major axis $a = 0.01544 \text{ AU}$ (follows from P and the host-star mass $0.905 M_{\odot}$ by Kepler's third law).

KEY, WASP-76 b: single transit




Transit depth 1.17%, duration 3.84 h. Host star WASP-76 (F7V), radius 1.756 $R_\odot$.

KEY, WASP-76 b: light curve over several orbits




Orbital period $P = 1.81 \text{ d}$. Semi-major axis $a = 0.033 \text{ AU}$ (follows from P and the host-star mass $1.458 M_{\odot}$ by Kepler's third law).

KEY, TRAPPIST-1 e: single transit




Transit depth 0.50%, duration 0.95 h. Host star TRAPPIST-1 (M8V), radius 0.119 $R_\odot$.

KEY, TRAPPIST-1 e: light curve over several orbits




Orbital period $P = 6.101 \text{ d}$. Semi-major axis $a = 0.02925 \text{ AU}$ (follows from P and the host-star mass $0.090 M_{\odot}$ by Kepler's third law).

KEY, K2-18 b: single transit




Transit depth 0.28%, duration 3.55 h. Host star K2-18 (M2.5V), radius 0.411 $R_{\odot}$.

KEY, K2-18 b: light curve over several orbits




Orbital period $P = 32.94$ d. Semi-major axis $a = 0.1429$ AU (follows from P and the host-star mass $0.359 M_{\odot}$ by Kepler's third law).

KEY, TOI-700 d: single transit




Transit depth 544 ppm (0.054%), duration 3.51 h. Host star TOI-700 (M2.5V), radius 0.421 $R_{\odot}$.

KEY, TOI-700 d: light curve over several orbits




Orbital period $P = 37.42$ d. Semi-major axis $a = 0.1633$ AU (follows from P and the host-star mass $0.415 M_{\odot}$ by Kepler's third law).

KEY, GJ 1214 b: single transit




Transit depth 1.34%, duration 0.90 h. Host star GJ 1214 (M4V), radius 0.216 $R_\odot$.

KEY, GJ 1214 b: light curve over several orbits




Orbital period $P = 1.58$ d. Semi-major axis $a = 0.01505$ AU (follows from P and the host-star mass $0.182 M_{\odot}$ by Kepler's third law).

KEY, LHS 1140 b: single transit




Transit depth 0.54%, duration 2.15 h. Host star LHS 1140 (M4.5V), radius 0.216 $R_{\odot}$.

KEY, LHS 1140 b: light curve over several orbits




Orbital period $P = 24.74$ d. Semi-major axis $a = 0.0946$ AU (follows from P and the host-star mass $0.184 M_{\odot}$ by Kepler's third law).

KEY, TRAPPIST-1 g: single transit



Transit depth 0.75%, duration 1.21 h. Host star TRAPPIST-1 (M8V), radius 0.119 $R_\odot$.

KEY, TRAPPIST-1 g: light curve over several orbits



Orbital period $P = 12.35 \text{ d}$. Semi-major axis $a = 0.04683 \text{ AU}$ (follows from P and the host-star mass $0.090 M_{\odot}$ by Kepler's third law).