

Interval Vertex Deletion Admits Polynomial Kernel

Akanksha Agrawal, Pranabendu Misra, Saket Saurabh, and Meirav Zehavi

TAMREF (Changki Yun)

Sep 10, 2020

Seoul Nat'l Univ

Table of contents

1. Introduction
2. Computing k -redundant set
3. Designing a kernel
4. Main task
5. Epilogue

Intro

Graph Modification Problem

The **Graph Modification Problem** is an intriguing type of problem. Given a class \mathcal{F} of graphs, The \mathcal{F} -modification problem inquires

Given a graph G , Is it possible to add/delete at most k vertices/edges in order to make modified graph belong to \mathcal{F} ?

Each of them are called vertex/edges completion/deletion problems, respectively. There are a variety of special named questions about the **vertex deletion problem**.

- Feedback Vertex Set – \mathcal{F} is consisted of acyclic graphs.
- Maximum Clique Problem – \mathcal{F} is consisted of complete graphs.
- Minimum Vertex Cover – \mathcal{F} is consisted of edgeless graphs.

...most of them looks quite hard.

Graph Modification Problem

Class \mathcal{F} is called **hereditary**, if $G \in \mathcal{F}$ implies that any induced subgraph of G belongs to \mathcal{F} as well.

Theorem (Lewis & Yannakakis, 1980)

- For any nontrivial hereditary graph class \mathcal{F} , \mathcal{F} -modification problem is **NP-hard**.

Known that the problem is *MaxSNP-hard*, which is out of our interest. After all, it is very hard to solve.

Graph Modification Problem

One of the canonical approach to challenge NP-hard problem is examining *FPT* (*Fixed-Parameter Tractibility*).

A problem with *parameter* k is Fixed-Parameter Tractable if it is solvable in $\mathcal{O}(f(k) \cdot n^{\mathcal{O}(1)})$ for some computable function $f(k)$.

- f can be arbitrarily large such as $f(k) = k^{R(k,k^k)}$ as long as it's finite.
- Shoving all the 'hardness' of problem into k .

For example, vertex cover problem with n vertices and m edges may run in $\mathcal{O}(2^k \cdot \min(m, nk))$ time, with a pruning technique – vertex cover is FPT.

Elucidating that \mathcal{F} -modification problem is FPT, received attention for decades — bipartite graph, forests, chordal graphs... and finally interval graph. **(Cao & Marx, 2015)**

Current best complexity is $\mathcal{O}(6^k(n + m))$ by Cao (2016).

Kernelization – Vertex Cover Example

Look at the ‘pruning scheme’ for a given vertex cover instance (G, k) .

1. Ignore all the isolated vertices v and solve for $(G - v, k)$.
2. If there is a vertex v with degree $\geq k$, include v to answer and solve for $(G - v, k - 1)$.
3. If G has more than k^2 edges and neither two rules above is not available, return No.

After the pruning scheme, G has at most $2k^2$ vertices, and applying the previous search tree algorithm gives the overall complexity is about $\mathcal{O}(k^2 2^k + n + m)$. Note that now we may cope with the case $k = 15$ and $n = 10^6$.

Kernelization means a *reduction step* from a parametrized problem instance (x, k) to an equivalent instance (x', k') , where $|x'|$ is bounded by some function $f(k')$.

- If f is $k^{\mathcal{O}(1)}$, we say the problem admits a *polynomial kernel*.

The *reduction step* need not to be polynomial, but it is common for a polynomial kernel to accept a polynomial reduction step.

In fact, kernelizability and FPT is equivalent in some sense. But FPT does not guarantee the existence of a polynomial kernel.


Let the story begin

Now, it's clear to say what is today's goal.

- Interval Graph Vertex Deletion (IVD) is NP-hard. (Lewis & Yannakakis, 1980)
- IVD is FPT. (Cao & Marx, SODA '14)
- IVD admits a **Polynomial Kernel. (Agrawal et al, SODA '19)**


But, we didn't mention a single word about the class *Interval Graph*. We share a few slide to analyze Interval Graphs.

Interval Graph




- Interval Graphs are **Chordal**. i.e. It does not have induced subgraph C_n for $n \geq 4$.
- Surprisingly, interval graphs are distinguishable to add a few type of forbidden induced subgraphs including the chordal condition, called *obstructions*.

Obstructions



- (t, t_ℓ, t_r) forms a Asteroidal Triple (AT)s, two of them are connected by a path avoiding neighbor of the rest.
- t_ℓ, t_r are called *non-shallow terminal*, t is called a *shallow terminal*.
- c_i are called 'center's, and b_i 's are called 'base vertices'.
- 'Base vertices' form a chordless path.

Path Decomposition



- Another way to characterize interval graph is using the path decomposition of graph.
- A path decomposition of G is a path equipped with some 'bags' of $V(G)$, assigning some common bags for each adjacent pair of vertices.
- Rigorous definition goes to the next slide.

Path Decomposition

A path decomposition (P, β) of G is consisted of a path P and ‘bags’ $\beta : V(P) \rightarrow 2^{V(G)}$ such that,

- $\bigcup_{p \in V(P)} \beta(p) = V(G)$ — bags cover the whole vertex set.
- If $uv \in E(G)$, $\exists q \in V(P)$ s.t. $u, v \in \beta(q)$. — adjacent vertices have some common bags.
- $P_v := \{p \in V(P) \mid v \in \beta(p)\}$ is a subpath of P — Each vertex is contained in some consecutive bags.

If $\beta(p)$ are distinct maximal cliques, (P, β) is called a *clique path*.

Theorem.

- Graph G is interval if and only if G has a clique path.

1. Computing Redundant Sets – (Chap 3)

Terminology Setup

Given an IVD instance (G, k) .

- $X \subset V(G)$ is called *solution* if $G - X$ is an interval graph.
- Solution X is called a t -solution if $|X| \leq t$.
- For $\mathcal{W} \subseteq 2^{V(G)}$, $S \subset V(G)$ is a *hitting set* of \mathcal{W} if $W \cap S \neq \emptyset \forall W \in \mathcal{W}$.
- \mathcal{W} is called t -necessary if all t -solutions hits \mathcal{W} .
- Obstruction \mathbb{O} is *covered* by \mathcal{W} , if there is $W \in \mathcal{W}$ s.t. $W \subset V(\mathbb{O})$.
- $M \subset V(G)$ is called t -redundant w.r.t \mathcal{W} , if for every obstruction \mathbb{O} either
 - \mathbb{O} is covered by \mathcal{W} .
 - $|M \cap V(\mathbb{O})| > t$

Terminology Setup

All the other words are generic. What do you mean by *necessary* and *redundant*?

- We will eventually use the $(k+2)$ -necessary family \mathcal{W} and a 9-redundant solution M w.r.t. \mathcal{W} such that $\mathcal{W} \subseteq 2^M$.
- Often we drop a carefully chosen vertex $v \notin M$, and claim that X , the solution of $(G-v, k)$ is a solution for (G, k) .
- If it's *false*, there is an obstruction \mathbb{O} such that $V(\mathbb{O}) \subset G-X$, and definitely $v \in V(\mathbb{O})$. Now we prove that $|V(\mathbb{O}) \cap M| > 9$.
- Otherwise, \mathbb{O} is covered by \mathcal{W} . Thus there is a $Z \in \mathcal{W}$ such that $Z \subseteq V(\mathbb{O})$. Then $X \cup \{v\}$, being a $(k+1)$ -solution, intersects with Z hence $X \cap Z \neq \emptyset$. This contradicts to the choice of \mathbb{O} .

Hence, we only need to consider the *large* obstructions *highly intersecting* with M , which leads to a structural benefit.

Lemma 3.1 (Agrawal)

- Given an IVD instance (G, l) . For a fixed $r \in \mathbb{N}$, it is poly-time attainable to either verify that (G, l) is No-instance, or compute (\mathcal{W}, M) such that
 - $\mathcal{W} \subseteq 2^M$.
 - \mathcal{W} is l -necessary.
 - M is r -redundant w.r.t. \mathcal{W} .
 - M is $(r+1)(6l)^{r+1}$ -solution.

To prove Lemma 3.1, we use the following algorithm as a black-box.

Propo 3.1 (Cao)

- IVD admits a poly-time 6-approximation algorithm. We call this $\text{ApproxIVD}(G)$.

Analyzing tool — copy of a graph

Define $G' := G.\text{copy}(U, t)$ for $U \subset V(G)$ and $t \in \mathbb{N}$, such that

- $V(G') = \{v^0 \mid v \in V(G)\} \cup \{v^i \mid (v, i) \in U \times [t]\}$
- $E(G') = \{u^i v^j \mid uv \in E(G), 0 \leq i, j \leq t\} \cup \{v^i v^j \mid v \in V(G), 0 \leq i < j \leq t\}$

So we make t replicas for vertices in G , and made a clique with clones. We sometimes identify G as a subgraph of G' .

Note that G' is also interval graph for an interval graph G .

In this problem, we specify $G' := G.\text{copy}(U, 6l + 1)$, and analyze $\text{ApproxIVD}(G')$.

Let $A := \text{ApproxIVD}(G')$. The following statement holds.

- If $|A| > 6l$, then $\{U\}$ is l -necessary.
- If $|A| \leq 6l$, Every obstruction \mathcal{O} of G intersects with $(A \cap V(G)) \setminus U$.

Thus, we may obtain a necessary set or augment the redundancy.

Proof.

- If $|A| > 6l$, (G', l) is a No-instance. If $\{U\}$ is not necessary, there is an l -solution L s.t. $L \cap U = \emptyset$. Thus $G - L$ is an interval graph containing U . Thus $(G - L).\text{copy}(U, 6l + 1)$ equals $G' - L$ and it's an interval graph. Contradiction, since L becomes an l -solution for (G', l) .
- If $|A| \leq 6l$, every $b \in V(\mathcal{O}) \cap U$ has a 'twin' $b^{i(b)} \in G' \setminus A$ since b has $6l + 1$ duplicates. Exchanging b and $b^{i(b)}$ gives another obstruction, hence there should be a vertex $c \in (A \cap V(G)) \setminus U$.

The algorithm RedundantIVD

$M_0 := \text{ApproxIVD}(G)$, $\mathcal{W}_0 := \emptyset$, $\mathcal{T}_0 := \{(v) \mid v \in V(M_0)\}$.

If $|V(M_0)| > 6l$ return No. For $i = 1, \dots, r$,

1. $M_i := M_{i-1}$, $\mathcal{W}_i := \mathcal{W}_{i-1}$, $\mathcal{T}_i := \emptyset$.
2. For $\mathbf{v} = (v_0, \dots, v_{i-1}) \in \mathcal{T}_{i-1}$
 - 2.1 $A := \text{ApproxIVD}(G.\text{copy}(\mathbf{v}, 6l + 1))$.
 - 2.2 If $|A| > 6l$, $\mathcal{W}_i.\text{append}(\mathbf{v})$
 - 2.3 Else for each $u \in (A \cap V(G)) \setminus \mathbf{v}$, $\mathcal{T}_i.\text{append}(\mathbf{v} :: u)$, $M_i.\text{append}(u)$

return (M_r, \mathcal{W}_r) .

- It is clear that \mathcal{W}_i is l -necessary and M_i is i -redundant solution w.r.t \mathcal{W}_i , and $\mathcal{W}_i \subseteq 2^{M_i}$.
- Now we analyze the size of M_r, \mathcal{W}_r .
 - $|\mathcal{T}_i| \leq (6l)^{i+1}$.
 - $|M_i| \leq |M_{i-1}| + 6l |\mathcal{T}_i| \leq \sum_{j=0}^i (6l)^{j+1} \leq (i+1)(6l)^{i+1}$.
 - $|\mathcal{W}_i| \leq |\mathcal{T}_{i-1}| \leq (6l)^{i+1}$.

Intermission

Now, we have a 9-redundant $\mathcal{O}(k^{10})$ -solution M w.r.t $k+2$ -necessary family \mathcal{W} . Additionally, we may erase all the singletons in \mathcal{W} and include them in the answer.

However, we still have to bound the kernel in $G - M$. To resolve this, we introduce the concept of **graph modules**.

$C \subseteq V(G)$ is a **module** if all the vertices in C has same neighbor set outside C . (i.e. for all $v \notin C$, $N_G(v) \cap C \in \{\emptyset, C\}$)

We classify the connected components of $G - M$ into module (in G) components and non-module (in G) components. Then we need to:

- Bound the number of distinct components in kernel
- Bound the size of each components in kernel

- Bounding the number of non-module components is easy.
- Bounding the size of module components is easy.
- Bounding the number of module components is involved.
- Bounding the size of non-module components is *dreadful*.

Number of non-module components

Claim.

- It is possible to leave at most $(k+2)|M| = \mathcal{O}(k^{11})$ non-module components.

If not, there is a vertex $w \in M$ such that for at least $k+3$ non-module components E_1, \dots, E_{k+3} , each E_i contains $u_i v_i \in E(G)$ such that $u_i w \in E(G)$ but $v_i w \notin E(G)$. Then for every k -subset S excluding w of G , $G - S$ contains a **Long Claw** centred at w . Thus every k -solution must contain w . So we simply pick w and remove from G until the claim hold.

Size of module components

We introduce a simple proposition beforehand.

- Given an obstruction \mathbb{O} with size > 4 and a module component D , either $V(\mathbb{O}) \subseteq V(D)$ or $|V(\mathbb{O}) \cap V(D)| \leq 1$.

Claim

- For a minimal obstruction \mathbb{O} not covered by \mathcal{W} and a module component of D , $|V(\mathbb{O}) \cap V(D)| \leq 1$ holds.

\mathbb{O} is not covered by \mathcal{W} means $|M \cap V(\mathbb{O})| > 9$, thus $|V(\mathbb{O})| > 4$ holds. Also D is interval graph, hence $V(\mathbb{O}) \not\subseteq V(D)$. Thus $|V(\mathbb{O}) \cap V(D)| \leq 1$.

Size of module components

If there is a module component D with $|V(D)| > k + 1$, we just erase an arbitrary vertex $v \in V(D)$ and solve for $(G - v, k)$.

If S is a k -solution for $(G - v, k)$, we claim that S is a k -solution for G as well. If not, take a minimal obstruction \mathbb{O} of $G - S$, and v must be included.

Since $S + v$ is $(k + 1)$ -solution, S hits \mathcal{W} by its $(k + 2)$ -necessity. Also, $\mathcal{W} \subset 2^M$ guarantees that S hits \mathcal{W} . S and \mathbb{O} are disjoint, hence \mathbb{O} is not covered by \mathcal{W} .

By the previous lemma, $\{v\} = V(\mathbb{O}) \cap V(D)$. Since D is a module component, $\mathbb{O} - v + u$ is also obstruction for all $u \in V(D)$. As $|V(D)| > k + 1$, $|V(D) \setminus S \setminus v| \geq 1$ means there is an obstruction $\mathbb{O}' = \mathbb{O} - v + u'$ for some $u' \in V(D) \setminus S \setminus v$, which contradicts to that S is a k -solution for $G \setminus v$.

Main

- 9-redundant solution M w.r.t. $(k+2)$ -necessary family \mathcal{W} : $\mathcal{O}(k^{10})$.
- Number of non-module components: $\mathcal{O}(k^{11})$
- Size of each module component: $k+1$
- Number of module components: $\mathcal{O}(k^{102})$.
- Size of each non-module component: ???

Main Lemma (Lemma 4.3)

Variables

- (G, k) : IVD instance
- \hat{M} : some arbitrary vertex set. $M' := M \cup \hat{M}$, which is still 9-redundant.
- $\hat{\mathcal{C}}$: set of module compo's in $G - M'$.

In polynomial time,

We can do either

- Find an equivalent instance (G', k) such that G' is a strictly induced subgraph of G
- Find a set $B \subseteq V(\hat{\mathcal{C}})$ with $|B| \leq 8(k+1)^2 |M'|^{10}$ such that for all k -subset $S \subset V(G)$ which has a corresponding obstruction \mathcal{O}_S such that $S \cap \mathcal{O}_S = \emptyset$ and not covered by \mathcal{W} , there is an obstruction \mathcal{O}' such that $\mathcal{O}' \cap S = \emptyset$ and $\mathcal{O}' \cap (V(\hat{\mathcal{C}}) \setminus B) = \emptyset$.

Power of Lemma 4.3

To reduce the module component size, just put $\hat{M} = \emptyset$.

- If Lemma 4.3 gives (G', k) , just solve for it.
- If Lemma 4.3 gives the set B , delete arbitrary $v \in V(\hat{\mathcal{C}}) \setminus B$ and solve $(G - v, k)$.

We prove that S , a k -solution for $(G - v, k)$ is also a k -solution for (G, k) . Similarly assume the contrary, and then S hits \mathcal{W} and the minimal obstruction \mathcal{O}_S in $G - S$ is not covered by \mathcal{W} . Thus applying the Lemma 4.3, there is an obstruction \mathcal{O}' with $\mathcal{O}' \cap S = \emptyset$ and $\mathcal{O}' \cap (V(\hat{\mathcal{C}}) \setminus B) = \emptyset$. Where $v \in V(\hat{\mathcal{C}}) \setminus B$, the existence of \mathcal{O}' contradicts that $S + v$ is a $k + 1$ -solution.

Thus, the reduction rule guarantees that

$$|V(\hat{\mathcal{C}})| \leq 8(k+1)^2 |M'|^{10} = \mathcal{O}(k^{102}).$$

So our final goal is to construct such B to prove the Lemma 4.3. we switch to the general case with $\hat{M} \neq \emptyset$ again.

A handy lemma (Lemma 4.6)

Lemma 4.6 (Agrawal)

- For $u \neq v \in V(G)$ with $uv \notin E(G)$, then either $\{u, v\} \in \mathcal{W}$ or $G[(N_G(u) \cap N_G(v)) - M']$ is a clique.

Assume $\{u, v\} \notin \mathcal{W}$ and there exists a pair of vertices $x, y \in (N_G(u) \cap N_G(v)) - M'$ such that $xy \notin E(G)$. Then $\mathcal{O} := G[\{u, v, x, y\}]$ is a hole C_4 . \mathcal{O} is not covered by \mathcal{W} since $x, y \notin M'$, $\{u, v\} \notin \mathcal{W}$ and there are no singletons in \mathcal{W} . Thus $|M' \cap V(\mathcal{O})| > 9$, but $|V(\mathcal{O})| = 4$ leads to a contradiction.

Consequence of Lemma 4.6

Henceforth, for each non-adjacent u, v with $\{u, v\} \notin \mathcal{W}$, there is at most one module component $C_{uv} \in \hat{\mathcal{C}}$ such that $N_G(u) \cap N_G(v) \cap V(C_{uv}) \neq \emptyset$. Denote

$$\mathcal{C}^* := \{C_{uv} \mid uv \notin E(G), \{u, v\} \notin \mathcal{W}, N_G(u) \cap N_G(v) \cap V(C_{uv}) \neq \emptyset\}$$

Then $|V(\mathcal{C}^*)| \leq (k+1)|M'|^2 = \mathcal{O}(k^{21})$, which is small enough. so we just care for module components in $\mathcal{C}' := \hat{\mathcal{C}} \setminus \mathcal{C}^*$.


- Given a vertex $v \in V(\mathcal{C}')$ and $u \neq w \in N_G(v) \cap M'$, $\{u, w\} \in \mathcal{W}$ or $uw \in E(G)$.

Classification Lemma (Lemma 4.11)

Variables

- $C \in \mathcal{C}'$: a module component.
- \mathbb{O} : a minimal obstruction not covered by \mathcal{W} , intersecting with C .

Then $\{t\} = V(\mathbb{O}) \cap V(C)$ and \mathbb{O} is an **AW**, having t as a terminal.



Since \mathbb{O} is not covered, $|V(\mathbb{O})| > 9$ and thus \mathbb{O} is either \dagger -AW, \ddagger -AW or a hole. Since $N_G(t) \subset V(C) \cup M'$ and Lemma 4.6, for any $u, w \in N_G(t) \cap V(\mathbb{O})$ is adjacent, since $\{u, w\} \notin \mathcal{W}$. The figure implies that t must be a terminal.

Handling non-shallow terminals

- Remark that t_l, t_r are non-shallow terminals, while t is a shallow terminal.

Lemma 4.13 (Agrawal)

- There is a set $A' \subseteq V(\mathcal{C}')$ such that $|A'| \leq (k+1) |M'|^4$, all non-shallow terminals in $V(\mathcal{C}') \setminus A'$ has $k+1$ alternatives.
- Rigorously, if an obstruction \mathbb{O} is not covered by \mathcal{W} and \mathbb{O} is an AW having a non-shallow terminal $v \in V(\mathcal{C}') \setminus A'$, there are $k+1$ elements $a \in A'$ such that $\mathbb{O} - v + a$ is still an obstruction.
- So, if S the k -solution of $G \setminus v$ is not a k -solution for (G, k) ...

We won't look at the proof and the construction. It's just a case-work.

Lemma 4.17 (Agrawal)

- There is a set $A'' \subseteq V(\mathcal{C}')$ such that $|A''| \leq 4(k+1)^2 |M'|^{10}$, all shallow terminals in $V(\mathcal{C}') \setminus (A' \cup A'')$ has $k+1$ alternatives.
- Rigorously, if an obstruction \mathbb{O} is not covered by \mathcal{W} and \mathbb{O} is an AW having a shallow terminal $v \in V(\mathcal{C}') \setminus (A' \cup A'')$, there are $k+1$ elements $a \in A''$ such that $\mathbb{O} - v + a$ is still an obstruction.
- So, if S the k -solution of $G \setminus v$ is not a k -solution for (G, k) ...

... Let's peek at this! (optional)

Observation 4.6

Let P be a chordless path. We say that P is covered by \mathcal{W} if there is $W \in \mathcal{W}$ with $W \subset V(P)$, it's same as the obstructions.

- Let P be an chordless path in $G[V(G) - V(C)]$ for some $C \in \hat{\mathcal{C}}'$, not covered by \mathcal{W} Then for all $v \in V(C)$ $|N_G(v) \cap V(P)| \leq 2$, and they are adjacent.
- If $u, v \in N_G(v) \cap V(P)$, since $\{u, v\} \notin \mathcal{W}$ $(u, v) \in E(G)$. Thus, u, v must be adjacent and $N_G(v) \cap V(P)$ cannot contain more than three elements to maintain chord-free property.

Handling shallow terminals

Define $M'' := M' \cup V(\mathcal{C}^*) \cup A'$.

For all 2-subset S of M' , denote $A_S := \{v \in V(\mathcal{C}') \setminus A' \mid S \subseteq N_G(v)\}$. It intuitively means the candidate of vertices can be a shallow terminal of AW, having S as center.

Denote $V(M'') = \{v_1, \dots, v_p\}$ and $E(G[M'']) = \{e_1, \dots, e_q\}$. We define a incidence vector $\text{inc}(u) \in \mathbb{F}_2^{p+q+1}$ for all $u \in V(\mathcal{C}') \setminus A'$ as:

- $\text{inc}(u)_i$ is 1 iff $v_i \in N_G(u)$. ($1 \leq i \leq p$)
- $\text{inc}(u)_{p+i}$ is 1 iff both endpoints of e_i is in $N_G(u)$. ($1 \leq i \leq q$)
- $\text{inc}(u)_{p+q+1} = 1$.

inc_i completely describes the possible incidence relation of u .

Basis Marking Strategy

For every 2-subset $S \subset M'$, define the **multiset** of vectors $\mathbf{V}_S := \{\text{inc}(u) \mid u \in A_S\}$. Then we gradually build up the bases:

1. $\mathbf{U}_S^0 := \emptyset$.
2. For $i = 1, \dots, k+1$, build the basis \mathbf{B}_S^i for the vector subspace $\mathbf{V}_S \setminus \mathbf{U}_S^{i-1}$, and assign $\mathbf{U}_S^i = \mathbf{U}_S^{i-1} \cup \mathbf{B}_S^i$.
3. For every $\mathbf{v} \in \mathbf{U}_S^{k+1}$, choose an arbitrary vertex $u_{\mathbf{v}}$ such that $\text{inc}(u) = \mathbf{v}$.
4. Denote $A_S'' := \{u_{\mathbf{v}} \mid \mathbf{v} \in \mathbf{U}_S^{k+1}\}$.

Then merge all A_S'' into a single set $A'' := \bigcup_S A_S''$. Note that $|A_S''| \leq (k+1)(p+q) \leq (k+1)p^2$ since A_S'' is $k+1$ -union of vector spaces with $\dim(p+q)$. Thus $|A''| \leq (k+1)|M'|^2 |M''|^2 \leq 4(k+1)^2 |M'|^{10}$.

Define a vector $\mathbf{n}^{X,Y}$, such that $\mathbf{n}_i^{X,Y} = 1$ if and only if $i \in X$ or $i - p \in Y$ or $i = p + q + 1$. i.e. \mathbf{n}_i extracts the integer from vertex set X and the edge set Y .

For a path P , $\mathbf{n}^P := \mathbf{n}^{V(P) \cap M'', E(P) \cap E(G[M''])}$.

Lemma 4.16 (Agrawal)

- Let P be an induced path in $G[V(G) \setminus V(C)]$ for some $C \in \mathcal{C}'$, such that P is not covered by \mathcal{W} . For all $u \in V(C)$, $\mathbf{n}^P \cdot \text{inc}(u) = 1 \pmod{2}$ iff $N_G(u) \cap V(P) = \emptyset$.

Since the only possible case is $|N_G(u) \cap V(P)| = 0, 1, 2$, examining each case gives the proof.

Lemma 4.17 (Restated)

- Let $w \in V(\hat{\mathcal{C}}') \setminus (A' \cup A'')$, and \mathbb{O} be an AW with the shallow terminal w . Let S be the center of \mathbb{O} , then for all $i \in [k+1]$ there is a $v \in \mathbf{B}_S^i$ such that $\mathbb{O} - w + u_v$ is still an obstruction.

Since $w \in A_S$ but $w \notin A''$, $\text{inc}(w) \in \mathbf{V}_S$ is not spent by \mathbf{U}_S^i . Thus $\text{inc}(w) \in \mathbf{V}_S \setminus \mathbf{U}_S^{i-1}$ for all $i \in [k+1]$.

Thus

$$\text{inc}(w) = \text{inc}(b_{i1}) + \cdots + \text{inc}(b_{it})$$

for some $\text{inc}(b_{ij}) \in \mathbf{B}_S^i$. Taking a dot product with the chordless path P of \mathbb{O} ,

$$1 = \mathbf{n}^P \cdot \text{inc}(b_{i1}) + \cdots + \mathbf{n}^P \cdot \text{inc}(b_{it})$$

indicates that there is some j such that $\mathbf{n}^P \cdot \text{inc}(b_{ij}) = 1$. Then $\mathbb{O} - w + b_{ij}$ is still an obstruction, and we may choose j for all $i \in [k+1]$.

Hence, all the vertices in $V(\hat{\mathcal{C}}) \setminus (V(\mathcal{C}^*) \cup A' \cup A'')$ is a terminal of some AW, and there are $k + 1$ 'replica obstructions' in $(V(\mathcal{C}^*) \cup A' \cup A'')$.

Since $|V(\mathcal{C}^*) \cup A' \cup A''| = \mathcal{O}(k^{102})$, our main Lemma 4.3 is proved.

Epilogue

- 9-redundant solution M w.r.t. $(k+2)$ -necessary family \mathcal{W} : $\mathcal{O}(k^{10})$.
- Number of non-module components: $\mathcal{O}(k^{11})$
- Size of each module component: $k+1$
- Number of module components: $\mathcal{O}(k^{102})$.
- Size of each non-module component: ???

In current state, we may kernelize the subset with additional parameter –
the vertex cover size.

- 9-redundant solution M w.r.t. $(k+2)$ -necessary family \mathcal{W} : $\mathcal{O}(k^{10})$.
- Number of non-module components: $\mathcal{O}(k^{11})$
- Size of each module component: $k+1$
- Number of module components: $\mathcal{O}(k^{102})$.
- Size of each non-module component: 0

In current state, we may kernelize the subset with additional parameter – the vertex cover size.

Introducing M as a vertex cover, all the components in $G - M$ are independent set.

But we know that, the vertex cover can grow arbitrarily large. Thus for completion, we have to bound the size of each non-module component.

How to?

First, we adopt the clique-path decomposition of each non-module component. Then we may deal with each independent question:

- 9-redundant solution M w.r.t. $(k+2)$ -necessary family \mathcal{W} : $\mathcal{O}(k^{10})$.
- Number of non-module components: $\mathcal{O}(k^{11})$
- Size of each module component: $k+1$
- Number of module components: $\mathcal{O}(k^{102})$.
- Size of each non-module component: $?_1 \times ?_2$
 - Maximum size of bags in clique-path: $?_1$ (section 5, pp.21-40)
 - Maximum clique-path length in each non-module component: $?_2$ (section 6, pp.40-62)

How to?

First, we adopt the clique-path decomposition of each non-module component. Then we may deal with each independent question:

- 9-redundant solution M w.r.t. $(k+2)$ -necessary family \mathcal{W} : $\mathcal{O}(k^{10})$.
- Number of non-module components: $\mathcal{O}(k^{11})$
- Size of each module component: $k+1$
- Number of module components: $\mathcal{O}(k^{102})$.
- Size of each non-module component: $?_1 \times ?_2$
 - Maximum size of bags in clique-path: $?_1$ (section 5, pp.21-40)
 - Maximum clique-path length in each non-module component: $?_2$ (section 6, pp.40-62)

$?_1 = \mathcal{O}(k^{101})$, $?_2 = \mathcal{O}(k^{1629})$. Several marking strategies are used for each cases.

Hence, we obtain size $\mathcal{O}(k^{1741})$ kernel for IVD.

- One simpler problem is Chordal Vertex Deletion, which has $\mathcal{O}(k^{13})$ kernel found by Agrawal et al.
- The author believes there is about $\mathcal{O}(k^{10})$ kernel and leaves it to an open problem.
- The accessible paper is quite faulty... There are some calculation errors and typos.