



US008552000B2

(12) **United States Patent**
Ohno et al.(10) **Patent No.:** **US 8,552,000 B2**(45) **Date of Patent:** **Oct. 8, 2013**(54) **THERAPEUTIC AGENT FOR SENILE
DEMENTIA**(75) Inventors: **Yukihiro Ohno**, Osaka (JP); **Takeo
Ishiyama**, Suita (JP)(73) Assignee: **Dainippon Sumitomo Pharma Co.,
Ltd.**, Osaka-shi (JP)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.(21) Appl. No.: **13/409,075**(22) Filed: **Feb. 29, 2012**(65) **Prior Publication Data**

US 2012/0165341 A1 Jun. 28, 2012

Related U.S. Application Data(62) Division of application No. 12/140,927, filed on Jun.
17, 2008, now Pat. No. 8,148,379, which is a division
of application No. 10/562,039, filed as application No.
PCT/JP2004/009095 on Jun. 22, 2004, now
abandoned.(30) **Foreign Application Priority Data**

Jun. 23, 2003 (JP) 2003-178386

(51) **Int. Cl.**
A61K 31/497 (2006.01)(52) **U.S. Cl.**
USPC **514/254.02**(58) **Field of Classification Search**
None
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**

5,444,073 A	8/1995	Perregaard et al.
5,532,372 A	7/1996	Saji et al.
5,780,632 A	7/1998	Saji et al.
6,964,962 B2	11/2005	Wong et al.
7,067,507 B2	6/2006	Pulley et al.
2003/0050307 A1	3/2003	Ruhland et al.
2004/0014972 A1	1/2004	Gottschlich et al.
2006/0025422 A1	2/2006	Nakamura et al.
2006/0111429 A1	5/2006	Fish et al.
2009/0176800 A1	7/2009	Ishiyama
2010/0105692 A1	4/2010	Moheno et al.

FOREIGN PATENT DOCUMENTS

EP	0 464 846 A1	1/1992
JP	05-17440 A	1/1993
JP	6-504787 A	6/1994
JP	08-333368	12/1996
JP	2000-281576	10/2000
JP	2003-135074 A	5/2003
JP	2003-160583	6/2003
JP	2003-519226 A	6/2003
WO	WO 93/16073	8/1993
WO	WO 95/34306	12/1995

WO	WO 96/14297	5/1996
WO	WO 99/52519	10/1999
WO	WO 02/20491 A1	3/2002
WO	WO 02/22581 A1	3/2002
WO	WO 02/24166 A1	3/2002
WO	WO 03/066039 A1	8/2003
WO	WO 2004/017973	3/2004
WO	WO 2004/113333	12/2004
WO	WO 2007/124757 A2	11/2007
WO	WO 2008/124030	10/2008


OTHER PUBLICATIONS

Dwork. American Journal of Psychiatry, 1998, 155:11, 1536-1543.*
Togo. Journal of Neuroimmunology, 2002, 124, 83-92.*
"Delirium, Dementia, Amnesia, Cognitive Disorders," http://www.nlm.nih.gov/cgi/mesh/2009/MB_cgi?mode=&term m,+Dementia,+Amnestic,+Cognitive+Disorders&field=entry, accessed Jul. 1, 2009.
Alphas, Larry, "An industry perspective on the NIMH Consensus Statement on negative symptoms," Schizophrenia Bulletin, vol. 32, No. 2, pps. 225-230, (2006).
Approval Labeling Text, NDA 21-487, for NAMENDA™ (memantine hydrochloride) (2003).
Barber, Teresa A., et al., "Memantine ameliorates scopolamine-induced amnesia in chicks trained on taste-avoidance learning," Neurobiology of Learning and Memory, vol. 93, pp. 540-545, (2010).
Bejar, Corina, et al., "Effect of rivastigmine on scopolamine-induced memory impairment in rats," European Journal of Pharmacology, vol. 383, pp. 231-240, (1999).
Botero, Hector M. et al., "Structure—Activity Relationships in a Series of Bisquaternary Bisphthalimidine Derivatives Modulating the Muscarinic M2-Receptor Allosterically," J. Med. Chem., vol. 43, pp. 2155-2164, (2000).
Bowen, D.M., "Traditional pharmacotherapy may succeed in alzheimer's disease," Trends in Neurosciences, vol. 15, No. 3, pp. 84-85, (1992).

(Continued)


Primary Examiner — Noble Jarrell(74) *Attorney, Agent, or Firm* — Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P.(57) **ABSTRACT**

A therapeutic/preventive agent for cognitive dysfunctions, which comprises as an active ingredient an imide derivative of the following formula [1]:



[1]

wherein Z is a group of the formula [2]:



[2]

D is a group of $-(\text{CH}_2)_p-\text{A}-(\text{CH}_2)_q-$; G is $=\text{N}-$, $-\text{CH}-$, etc.; and Ar is an aromatic heterocyclic group, etc.**11 Claims, 2 Drawing Sheets**

(56)

References Cited

OTHER PUBLICATIONS

- Clinton et al., Altered transcript expression of NMDA receptor associated postsynaptic proteins in the thalamus of subject with schizophrenia, *Am. J. Psychiatry*, vol. 160, No. 6, pp. 1100-1109, (Jun. 2003).
- Clinton et al., "Thalamic expression of NMDA receptor-associated postsynaptic density proteins in schizophrenia," *Society for Neuroscience*, Program No. 754.4, (2003), (online) (abstract only).
- Cloninger, "The discovery of susceptibility genes for mental disorders," *Proc. Natl. Acad. Sci.*, vol. 99, No. 21, pp. 13365-13367, (Oct. 15, 2002).
- Corbett, "Clozapine but not haloperidol antagonizes an MK-801 discriminative stimulus cue," *Pharmacol. Biochem. Behav.*, vol. 51(2-3), pp. 561-564, (1995).
- Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV™) pp. 273-278, 285, and 286 (1994), published by the American Psychiatric Association, Washington D.C.
- Didriksen, et al., "Antipsychotic potential of the Gly T-1 inhibitor NFPS," *Society Neuroscience Abstract*, vol. 2002, abstract No. 893. 1, (2002).
- Doggrell, Sheila A. et al., "Treatment of dementia with neurotransmission modulation," *Expert Opinion on Investigational Drugs*, vol. 12, No. 10, pp. 1633-1654, (2003).
- Duka, Theodora, "Scopolamine-induced amnesia in humans: Lack of effects of the benzodiazepine receptor antagonist β -carboline ZK 93426," *Journal of Psychopharmacology*, vol. 6, No. 3, pp. 382-388, Abstract, (1992).
- Ebihara, Mitsuru et al., "Togo Scicchoshono Dobutsu Model," *Igaku no Ayumi*, vol. 208, No. 3, pp. 138-142, (2004).
- Emre, Murat, M.D. et al., "Rivastigmine for dementia associated with parkinson's disease," *The New England Journal of Medicine*, vol. 351, No. 24, pp. 2509-2518, (2004).
- English translation of Office Action from the Chinese Patent Office in Appln. No. 200480017534.X dated Jan. 29, 2010.
- English translation of Office Action from the Japanese Patent Office in Appln. No. 2005-507314 dated Jun. 29, 2010.
- English translation of Second Office Action from the Chinese Patent Office in Appln. No. 200480017534.X dated Jan. 29, 2010.
- Enomoto et al., "Development of antipsychotics by using animal model," *Brain Science*, vol. 25, No. 5, pp. 437-444 (2003).
- EP Official Action for Corresponding EP Application No. 04 746 564.6-2117 dated Aug. 27, 2010.
- EP Official Action for Corresponding EP Application No. 04 746 564.6-2117 dated Nov. 20, 2009.
- EP Search Report for European Patent Application No. 04746564.6 dated Mar. 2, 2009.
- Fabre, Serge et al., "Protein Kinase C Inhibitors; Structure—Activity Relationships in K252c-Related Compounds," *Bioorg. Med. Chem.*, vol. 1, No. 3, pp. 193-196, (1993).
- Fernandez, Hubert H. et al., "Pharmacotherapy of dementia with Lewy bodies," *Expert Opinion on Pharmacotherapy*, vol. 4, No. 11, pp. 2027-2037, (2003).
- Final Office Action in U.S. Appl. No. 10/525,021 mailed Mar. 5, 2010.
- Final Office Action in U.S. Appl. No. 10/525,021 mailed Sep. 17, 2008.
- Final Office Action in U.S. Appl. No. 12/140,927 dated Jul. 10, 2009.
- Final Office Action in U.S. Appl. No. 12/140,927 dated Jul. 18, 2011.
- Final Office Action in U.S. Appl. No. 12/401,958 mailed Apr. 5, 2010.
- Friedman, Joseph I., "Cholinergic targets for cognitive enhancement in schizophrenia: focus on cholinesterase inhibitors and muscarinic agonists," *Psychopharmacology*, 174, pp. 45-53, (2004).
- Geyer et al., "Animal behavior models of the mechanisms underlying antipsychotic atypicality," *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, vol. 27, pp. 1071-1079, (2003).
- Goff et al., "The emerging role of glutamate in the pathophysiology and treatment of schizophrenia," *Am. J. Psychiatry*, vol. 158, No. 9, pp. 1367-1377, (Sep. 2001).
- Harrod et al., "MK-801 induced retrieval, but not acquisition, deficits for passive avoidance conditioning," *Pharmacology, Biochemistry, and Behavior*, vol. 69, pp. 585-593, (2001).
- Harvey et al., "Cognition in schizophrenia: from basic science to clinical treatment," *Psychopharmacology*, vol. 169, pp. 213-214, (2003).
- Harvey et al., "Cognitive functioning in schizophrenia: a consensus statement on its role in the definition and evaluation of effective treatments for the illness," *J. Clin. Psychiatry*, vol. 65, pp. 361-372, (2004).
- Hyman et al., "What are the right targets for psychopharmacology?" *Science*, vol. 299, pp. 350-351, (Jan. 17, 2003).
- Ibrahim et al., "Ionotropic glutamate receptor binding and subunit mRNA expression in thalamic nuclei in schizophrenia," *Am. J. Psychiatry*, vol. 159, No. 11, pp. 1811-1823, (Nov. 2000).
- Ishizumi, Kikuo, et al., "Succinimide Derivatives. II. Synthesis and Antipsychotic Activity of N-[4-[4-(1,2-Benzisothiazol-3-yl)-1-piperazinyl]butyl]-1,2-cis-cyclohexanedicarboximide (SM-9018) and Related Compounds," *Chem. Pharm. Bull.*, vol. 43, No. 12, pp. 2139-2151, (1995).
- Jellinger, Kurt A., "The Pathology of Ischemic-Vascular Dementia: An Update," *Journal of the Neurological Sciences* 203-204, pp. 153-157, (2002).
- Kahle, Philipp J. et al., "The Emerging Utility of Animal Models of Chronic Neurodegenerative Diseases," *Emerging Therapeutic Targets*, vol. 5, No. 1, 125-132, (2001).
- Kasper et al., "Cognitive effects and antipsychotic treatment," *Psychoneuroendocrinology*, vol. 28, pp. 27-38, (2003).
- Kay, Stanley R. et al., "The Positive and Negative Syndrome Scale (PANSS) for Schizophrenia," *Schizophrenia Bulletin*, vol. 13, No. 2, pp. 261-276, (1987).
- Krystal et al., "NMDA receptor antagonist effects, cortical glutamatergic function, and schizophrenia: toward a paradigm shift in medication development," *Psychopharmacology*, vol. 169, pp. 215-233, (2003).
- Lindenmayer et al., "A new five factor model of schizophrenia," *Psychiatric Quarterly*, vol. 65, No. 4, pp. 299-322, (1994).
- Malenka et al., "Long term potentiation—A decade of progress?" *Science*, vol. 285, pp. 1870-1874, (Sep. 17, 1999).
- Meltzer et al., "Cognition, schizophrenia, and the atypical antipsychotic drugs," *Proc. Natl. Acad. Sci.*, vol. 96, No. 24, pp. 13591-13593, (Nov. 23, 1999).
- Metthey Y, et al., "Synthesis of 11-Aminodibenzol[b,f][1,4]thiazepines and Fluoro Derivatives," *Journal of Heterocyclic Chemistry*, vols. 03-04 No. 34, pp. 465-467, (1997).
- Misane et al., "Selective 5-HT_{1A} Antagonists WAY 10065 and NAD-299 Attenuate the Impairment of Passive Avoidance Caused by Scopolamine in the Rat," *Neuropsychopharmacology* 28, pp. 253-264, (2003).
- Miyachi, Hiroyuki et al., "Novel Biological Response Modifiers: Phthalimides with Tumor Necrosis Factor- α Production-Regulating Activity," *J. Med. Chem.*, vol. 40, pp. 2858-2865, (1997).
- Miyamoto et al., "Hyperfunction of dopaminergic and serotonergic neuronal systems in mice lacking the NMDA receptor E1 subunit," *Journal of Neuroscience*, vol. 21, No. 2, pp. 750-757, (Jan. 15, 2001).
- Moghaddam, "Bringing order to the glutamate chaos in schizophrenia," *Neuron*, vol. 40, pp. 881-884, (Dec. 4, 2003).
- Mohn et al., "Mice with reduced NMDA receptor expression display behaviors related to schizophrenia," *Cell*, vol. 98, pp. 427-436, (1999).
- Myhrer, "Neurotransmitter systems involved in learning and memory in the rat: a meta-analysis based on studies of four behavioral tasks," *Brain Research Reviews*, vol. 41, pp. 268-287, (2003).
- Nakagawa et al., "Ethanol-induced state-dependent learning is mediated by 5-hydroxytryptamine₃ receptors but not by N-methyl-D-aspartate receptor complex," *Brain Research*, vol. 706, pp. 227-232, (1996).
- Noda et al., "Clozapine, but not haloperidol, reverses working memory impairment induced by chronic PCP administration in rats: a new model for cognitive dysfunction in schizophrenia," *Abstracts Society Neuroscience*, vol. 26, Nos. 1-2, pp. 6533, (2000).

(56)

References Cited

OTHER PUBLICATIONS

- Norman, Mark H. et al., "Effect of Linking Bridge Modifications on the Antipsychotic Profile of Some Phthalimide and Isoindolinone Derivatives," *Journal of Medical Chemistry*, vol. 39, No. 1, pp. 149-57, (1996).
- Office Action in Japanese Application No. 2005-507314 issued on Jun. 29, 2010 (4 pages).
- Office Action in U.S. Appl. No. 10/525,021 mailed Dec. 17, 2007.
- Office Action in U.S. Appl. No. 10/525,021 mailed Jun. 12, 2009.
- Office Action in U.S. Appl. No. 10/562,039 mailed Mar. 18, 2008.
- Office Action in U.S. Appl. No. 10/589,804 mailed Dec. 11, 2008.
- Office Action in U.S. Appl. No. 12/140,927 (continuation of U.S. Appl. No. 10/562,039) mailed Oct. 3, 2008.
- Office Action in U.S. Appl. No. 12/140,927 dated Nov. 10, 2010.
- Office Action in U.S. Appl. No. 12/401,958 (continuation of U.S. Appl. No. 10/589,804) mailed Oct. 1, 2009.
- Parnetti, et al., "Cholinergic precursors in the treatment of cognitive impairment of vascular origin: Ineffective approaches or need for re-evaluation?," *Journal of the Neurological Sciences*, vol. 257, pp. 264-269, (2007).
- Perricone v. Medicis Pharm. Corp.*, 432 F.3d 1368 (Fed. Cir. 2005).
- Perry, Elaine et al., "Acetylcholine in Mind: a Neurotransmitter Correlate of Consciousness?," *TINS*, vol. 22, No. 6, pp. 273-280 (1999).
- Prescribing information for "Exelon® (rivastigmine tartrate) Capsules and Oral Solution," (31 pages), (2006).
- Prescribing Information for Aricept® (donepezil hydrochloride) (14 pages) (2010).
- Puttrese, et al., "Localized deletion of the NR1 gene in mouse prefrontal cortex impairs spatial memory," *Society Neuroscience Abstract* vol. 2003, abstract No. 964.19, (2003).
- Reingold, Jennifer L. et al., "Rivastigmine for the Treatment of Dementia Associated with Parkinson's Disease," *Neuropsychiatric Disease and Treatment* vol. 3, pp. 775-783, (2007).
- Roman, Gustavo C. et al., "Donepezil in Vascular Dementia: Combined Analysis of Two Large-Scale Clinical Trials," *Dementia and Geriatr. Cogn. Disord.*, vol. 20 pp. 337-344, (2005).
- Romero, Arthur G. et al., "Synthesis of Metabolically Stable Arylpiperazine 5-HT_{1A} Receptor Agonists," *Bioorganic & Medicinal Chemistry Letters*, vol. 2, No. 12, pp. 1703-1706, (1992).
- Sharma et al., "Cognitive function in schizophrenia deficits, functional consequences, and future treatment," *Psychiatr. Clin. N. Am.*, vol. 26, pp. 25-40, (2003).
- Shinkei Kairomu Keisei to Kofunsei Synapse Kasosei ni Kansura Kodogakuteki Kenkyu, pp. 13-2, with partial English language translation, (2003).
- Small, David H., "Acetylcholinesterase Inhibitors for the Treatment of Dementia in Alzheimer's Disease: Do We Need New Inhibitors?," *Expert Opinion on Emerging Drugs*, vol. 10, No. 4, pp. 817-823, (2005).
- The Lancet, "The Treatment of Senile Insanity," *Lancet Limited*, London, GB LNKD-DOI:10.1016/S0140-6736 (01) 05083-8, vol. 208, No. 5381, pp. 820-882, (Oct. 16, 1926).
- Thomas, Elizabeth et al., "Specific Impairments in Visuospatial Working and Short-Term Memory Following Low-Dose Scopolamine Challenge in Healthy Older Adults," *Neuropsychologia*, vol. 46, pp. 2476-2484, (2008).
- Tokita, Kenichi et al., "Combination of a Novel Antidementia Drug FK960 with Donepezil Synergistically Improves Memory Deficits in Rats," *Pharmacology, Biochemistry and Behavior*, vol. 73, pp. 511-519, (2002).
- Tokuda, et al., "Effects of SM-13496, an atypical antipsychotic agent, on MK-801-induced learning deficit in rats," *J. Pharmacol Sciences*, vol. 94, supplement 1, p. 163P, (2004).
- Turetsky et al., "Memory-Delineated Subtypes of Schizophrenia: Relationship to Clinical, Neuroanatomical, and Neurophysiological Measures," *Neuropsychology*, vol. No. 4, pp. 481-490 (2002).
- U.S. Appl. No. 10/525,021, filed Feb. 18, 2005.
- U.S. Appl. No. 10/562,039, filed Dec. 22, 2005.
- U.S. Appl. No. 10/589,804, filed Aug. 17, 2006.
- Weiss et al., "The effects of second-generation antipsychotics on cognitive functioning and psychosocial outcome in schizophrenia," *Psychopharmacology*, vol. 162, pp. 11-17 (2002).
- Xu Taixiang et al., "Status quo and Development of Alzheimer's Disease," *Acta Academiae Medicinae Qingdao Universitatis*, vol. 37, No. 4, pp. 355-357, (2001).
- Biederman, Joseph, et al., "Risperidone treatments for ADHD in children and adolescents with bipolar disorder," *Neuropsychiatric Diseases and Treatment*, vol. 4, No. 1, pp. 203-207 (2008).
- Bruno, et al., "The α_2c -adrenergic receptor mediates hyperactivity of colobomo mice, a model of attention deficit hyperactivity disorder," *Neurobiology of Disease*, vol. 23, pp. 679-688, (2006).
- Center for Drug Evaluation and Research, *Pharmacology Reviews at FDA*, pp. 1-260, (Oct. 2010).
- EP Search Report for European Patent Application No. 11160001.1. 2123 dated Jul. 19, 2011.
- Erhart, Stephen M., et al., "Treatment of schizophrenia negative symptoms: future prospects," *Schizophrenia Bulletin*, vol. 32, No. 2, pp. 234-237, 2006.
- European Neuropsychopharmacology, "P.3.155 Efficacy of lurasidone (SM-13496) in the treatment of schizophrenia: results of two, phase 2, placebo-controlled studies," vol. 15, pp. S522-S523, (2005).
- International Search Report for International Application No. PCT/JP2004/009095 dated Aug. 24, 2004.
- International Search Report for International Application No. PCT/JP2011/062314 dated Jun. 28, 2011.
- Ishiyama, T., et al., "Effects on sm-13496, a novel serotonin-dopamine antagonist, and other antipsychotics on cognitive performance in rat passive avoidance test," abstract, vol. 23, (2003).
- Japanese Office Action in corresponding Japanese Application No. 2006-510283 dated May 31, 2011.
- Kane, John, "Commentary: Consensus statement on negative symptoms," *Schizophrenia Bulletin*, vol. 32, No. 2, pp. 223-224, (2006).
- Kirkpatrick, Brian, et al., "The NIMH-MATRICS consensus statement on negative symptoms," vol. 32, No. 2, pp. 214-219, (2006).
- Laughren, Thomas, et al., "Food and Drug Administration perspective on negative symptoms in schizophrenia as a target for a drug treatment claim," *Schizophrenia Bulletin*, vol. 32, No. 2, pp. 220-222, (2006).
- Masi, Gabriele, et al., "Aripiprazole monotherapy in children and young adolescents with pervasive developmental disorders," *CNS Drugs*, vol. 23, No. 6, pp. 511-521, (2009).
- Meyer, Jonathan, M. et al., "Lurasidone: a new drug in development for schizophrenia," *Expert Opinion on Investigational Drugs*, vol. 18, No. 11, pp. 1715-1726, (2006).
- Nippon-Shinkei-Seishin-Yakurigaku Zasshi (JPn. J. Neuropsychopharmacol.) 23: 296 (2003).
- Office Action in U.S. Appl. No. 10/525,021 mailed Aug. 29, 2007.
- Office Action in U.S. Appl. No. 13/113,703, mailed Nov. 22, 2011.
- Ogasa et al., "SM-13496 in patients with acute exacerbation of schizophrenia: A two-dose double-blind phase II comparison with placebo," *Schizophrenia Research*, vol. 60, No. 1, pp. 297, (2003).
- Poster exhibited at the 18th European College of Neuropsychopharmacology Congress, Oct. 23-26, 2005.
- Powell, Susan, B., et al., "RO-10-5824 is a selective dopamine D4 receptor agonist that increases novel object exploration in C57 mice," *Neuropharmacology*, vol. 44, pp. 473-481, (2003).
- Protais, P. et al., "Climbing behavior induced by apomorphine in mice: a simple test for the study of dopamine receptors in striatum," *Psychopharmacology*, vol. 50, pp. 1-6, (1976).
- Russell, Vivienne, A., et al., "Animal models of attention-deficit hyperactivity disorder," *Behavioral and Brain Functions*, vol. 1, No. 9, pp. 1-17, (2005).
- Rutten, K., et al., "Selective PDE inhibitors rolipram and sildenafil improve object retrieval performance in adult cynomolgus macaques," *Psychopharmacology*, vol. 196, pp. 643-648, (2008).
- Snyder, Peter J. et al., "Reversal of scopolamine-induced deficits with a single dose of donepezil, an acetylcholinesterase inhibitor," *Alzheimer's & Dementia*, vol. 1, pp. 126-135, (2005).
- Takahashi, Satoshi et al., "Anti-Dementia Drugs and Vascular Dementia," *Rinsho-Seishinigaku, (Clinical Psychiatry)*, vol. 31, No. 10, pp. 1189-1193, (2002).

(56)

References Cited

OTHER PUBLICATIONS

Wang, D., et al. "Synergistic effect of galantamine with risperidone on impairment of social interaction in phencyclidine-treated mice as a schizophrenic animal model," *Neuropharmacology*, vol. 52, pp. 1179-1187 (2007).

Wise, L.E., et al., "Reversal learning in the 8—arm radial maze in rats is impaired by subchronic administration of the non-competitive

NMDA antagonist ketamine", Society for Neuroscience, abstract, vol. 2002, (2002).

Woolley et al., "Selective dopamine D4 receptor agonist (A-412997) improves cognitive performance and stimulates motor activity without influencing reward-related behaviour in rat," *Behavioural Pharmacology*, vol. 19, Iss. 8, pp. 765-776, (Dec. 2008).

EP Communication for Corresponding EP Application No. 04 746 564.6-2117 dated Mar. 23, 2012.

* cited by examiner

FIG. 1





FIG. 2



ethylidene may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom},

or an acid addition salt thereof.

(2) The therapeutic/preventive agent for cognitive dysfunctions according to the above (1), which is a therapeutic agent for senile dementia.


(3) A therapeutic/preventive agent for cognitive dysfunctions, which comprises as an active ingredient an imide derivative of the above formula [1], wherein —Ar is an aromatic heterobicyclic group, naphthyl, benzoyl, phenoxy or phenylthio and G is N, CH or COH, or —Ar is biphenyl-methylidene and G is a carbon atom (said aromatic heterobicyclic group, naphthyl, benzoyl, phenoxy, phenylthio and biphenylmethylidene may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom), or an acid addition salt thereof.

(4) The therapeutic/preventive agent for cognitive dysfunctions according to the above (3), which is a therapeutic agent for senile dementia.

(5) A therapeutic/preventive agent for cognitive dysfunctions, which comprises as an active ingredient an imide derivative of the above formula [1], wherein —Ar is an aromatic heterocyclic group condensed with a benzene ring, or naphthyl, benzoyl, phenoxy or phenylthio (said aromatic heterocyclic group condensed with a benzene ring, naphthyl, benzoyl, phenoxy, and phenylthio may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom), and G is N, CH or COH, or an acid addition salt thereof.


(6) The therapeutic/preventive agent for cognitive dysfunctions according to the above (5), which is a therapeutic agent for senile dementia.

(7) A therapeutic/preventive agent for cognitive dysfunctions, which comprises as an active ingredient an imide derivative of the above formula [1], wherein Z is a group of the formula [4]:




in which —L— is a single bond or a double bond, E is a lower alkylene optionally substituted by a lower alkyl, or an oxygen atom, R⁵ is a hydrogen atom or a lower alkyl, and B is the same as defined in the above (1);

a group of the formula [5]:




in which —L—, E, R⁵ and B are as defined above;

a group of the formula [6]:




in which R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵ are independently a hydrogen atom or a lower alkyl, or the adjacent two groups of R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵ may combine each other to form a double bond, and B is as defined above;

a group of the formula [7]:



in which R¹⁶ and R¹⁷ are independently a hydrogen atom or a lower alkyl, or R¹⁶ and R¹⁷ may combine each other to form a saturated hydrocarbon ring, and R⁵ and B are as defined above; or

a group of the formula [8]:




in which B is as defined above,

or an acid addition salt thereof.

(8) The therapeutic/preventive agent for cognitive dysfunctions according to the above (7), which is a therapeutic agent for senile dementia.

(9) A therapeutic/preventive agent for cognitive dysfunctions, which comprises as an active ingredient an imide derivative of the formula [9]:



or an acid addition salt thereof.

(10) The therapeutic/preventive agent for cognitive dysfunctions according to the above (9), which is a therapeutic agent for senile dementia.

5

(11) The therapeutic/preventive agent for cognitive dysfunctions according to the above (2), (4), (6), (8) or (10), which is a therapeutic agent for the Alzheimer type dementia.

(12) The therapeutic/preventive agent for cognitive dysfunctions according to the above (2), (4), (6), (8) or (10), which is a therapeutic agent for the cerebrovascular dementia.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 shows the effects of the imide derivative on rats in one step-through passive avoidance test where the acetylcholine receptor blocker scopolamine was used for inducing amnesia, and indicates the step-through latency during the training (*: $P < 0.05$ vs the group treated with 0.5% MC+scopolamine (Steel's test)).

FIG. 2 shows the effects of the imide derivatives on rats in one step-through passive avoidance test where the acetylcholine receptor blocker scopolamine was used for inducing amnesia, and indicates the step-through latency during the test (*: $P < 0.05$ vs the group treated with 0.5% MC+scopolamine (Steel's test), #: $P < 0.01$ vs the group treated with 0.5% MC+saline solution (Mann-Whitney test)).

BEST MODE FOR CARRYING OUT THE INVENTION

Each group of the imide derivative of the formula [1] of the present invention are explained in detail.

The lower alkylene for Z and A includes, for example, ones having not more than 3 carbon atoms such as methylene, ethylene, trimethylene, etc.

The hydrocarbon ring for Z and A includes, for example, a cycloalkane or cycloalkene having not more than 7 carbon atoms. The cycloalkane having not more than 7 carbon atoms includes, for example, cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, etc. The cycloalkene having not more than 7 carbon atoms includes, for example, cyclopentene, cyclohexene, cycloheptene, etc.

The hydrocarbon ring being cross-linked with a lower alkylene or an oxygen atom for Z and A includes, for example, rings having not more than 10 carbon atoms such as bicyclo[1.1.1]pentane, bicyclo[2.1.1]hexane, bicyclo[2.1.1]hex-2-ene, bicyclo[2.2.1]heptane, bicyclo[2.2.1]hept-2-ene, bicyclo[2.2.2]octane, bicyclo[2.2.2]oct-2-ene, bicyclo[4.1.1]octane, bicyclo[4.1.1]oct-2-ene, bicyclo[4.1.1]oct-3-ene, bicyclo[3.2.1]octane, bicyclo[3.2.1]oct-2-ene, bicyclo[3.2.1]oct-3-ene, bicyclo[3.2.1]oct-6-ene, bicyclo[3.2.2]nonane, bicyclo[3.2.2]non-2-ene, bicyclo[3.2.2]non-3-ene, bicyclo[3.2.2]non-6-ene, 2-oxabicyclo[1.1.1]butane, 2-oxabicyclo[2.1.1]pentane, 2-oxabicyclo[2.1.1]pent-4-ene, 7-oxabicyclo[2.2.1]hexane, 7-oxabicyclo[2.2.1]hex-2-ene, 7-oxabicyclo[4.1.1]heptane, 7-oxabicyclo[4.1.1]hept-2-ene, 7-oxabicyclo[4.1.1]hept-3-ene, 8-oxabicyclo[3.2.1]heptane, 8-oxabicyclo[3.2.1]hept-2-ene, 8-oxabicyclo[3.2.1]hept-3-ene, 8-oxabicyclo[3.2.1]hept-6-ene, etc.

The aromatic hydrocarbon ring for Z includes, for example, ones having not more than 10 carbon atoms such as phenyl ring, naphthyl ring, etc.

The binding position of the hydrocarbon ring for A includes, for example, -1,1-, -1,2-, -1,3-, -1,4-, etc.

The aromatic hydrocarbon group for —Ar includes, for example, ones having not more than 10 carbon atoms such as phenyl, naphthyl, etc. The aromatic heterocyclic group for —Ar includes, for example, an aromatic heteromonocyclic group and an aromatic heterobicyclic group.

The aromatic heteromonocyclic group includes, for example, ones having not more than 6 carbon atoms, and

6

further having the same or different 1 to 4 heteroatoms selected from a nitrogen atom, an oxygen atom and a sulfur atom, such as pyridyl, pyrimidinyl, thiazolyl, oxazolyl, isoxazolyl, isothiazolyl, furyl, imidazolyl, etc.

The aromatic heterobicyclic group includes, for example, ones having not more than 10 carbon atoms, and further having the same or different 1 to 5 heteroatoms selected from a nitrogen atom, an oxygen atom and a sulfur atom, such as benzolog-fused rings (e.g., benziso-thiazolyl, benzisoxazolyl, benzofuranyl, quinolyl, isoquinolyl, indolyl, indazolyl, benzimidazolyl, benzoxazolyl, etc.), naphthyridinyl, puteridyl, thienofuranyl, imidazothiophen-yl, imidazofuranyl, etc.

The alkyl includes, for example, ones having not more than 6 carbon atoms, and preferably lower alkyl groups having not more than 4 carbon atoms, such as methyl, ethyl, propyl, 2-propyl, butyl, etc. The lower alkyl includes, for example, ones having not more than 4 carbon atoms, such as methyl, ethyl, propyl, 2-propyl, butyl, etc.

The lower alkoxy includes, for example, ones having not more than 4 carbon atoms, such as methoxy, ethoxy, propoxy, 2-propoxy, butoxy, etc.

The halogen atom is fluorine, chlorine, bromine, iodine.


The present compound [1] may have stereoisomers and/or an optical isomer. The present invention also includes a mixture of these isomers or each isolated isomer.

The preferable group for —Ar is an aromatic heterobicyclic group, or naphthyl, benzoyl, phenoxy or phenylthio (in these cases, G is N, CH, or COH), or biphenylmethylidene (in this case, G is a carbon atom), where said aromatic heterobicyclic group, naphthyl, benzoyl, phenoxy, phenylthio and biphenylmethylidene may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom.

The more preferable group for —Ar is a benzolog-fused ring, naphthyl, benzoyl, phenoxy, or phenyl (said benzolog-fused ring, naphthyl, benzoyl, phenoxy, and phenylthio may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom), and in this case, G is N, CH or COH.

The further preferable group for —Ar is benzisothiazolyl, benzisoxazolyl, isoquinolyl, benzofuranyl, indazolyl or indolyl (said benzisothiazolyl, benzisoxazolyl, isoquinolyl, benzofuranyl, indazolyl and indolyl may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom), and in this case, G is N, CH or COH.


The preferable group for Z is, for example, a group of the formula [4]:




(in which —L— is a single bond or a double bond, E is a lower alkylene optionally substituted by a lower alkyl, or an oxygen atom, R^5 is a hydrogen atom or a lower alkyl, and B is a carbonyl or a sulfonyl);

7

a group of the formula [5]:




(in which -L-, E, R⁵ and B are as defined above);
a group of the formula [6]:




(in which R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵ are independently a hydrogen atom or a lower alkyl, or the adjacent two groups of R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵ may combine each other to form a double bond, and B is as defined above);

a group of the formula [7]:



(in which R¹⁶, R¹⁷ are independently a hydrogen atom or a lower alkyl, or R¹⁶ and R¹⁷ may combine each other to form a saturated hydrocarbon ring, and R⁵ and B are as defined above); or

a group of the formula [8]:



(in which B is as defined above), etc.


Then, the saturated hydrocarbon ring formed by combining R¹⁶ and R¹⁷ includes, for example, a cycloalkane having not more than 7 carbon atoms, such as cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, etc.

8

The preferable group for Z is, for example, a group of the formula [10]:

[5]
5

[10]



10


(in which -L'- is a single bond, E is a lower alkylene optionally substituted by a lower alkyl, or an oxygen atom, R⁵ is a hydrogen atom or a lower alkyl, and B is a carbonyl or a sulfonyl);

a group of the formula [11]:

[6]

20

[11]




25

(in which -L'-, E, R⁵ and B are as defined above);
a group of the formula [12]:

30

[12]



[7]


40

(in which R^{6'}, R^{7'}, R^{8'}, R^{9'}, R^{10'}, R^{11'}, R^{12'}, R^{13'}, R^{14'}, R^{15'} are independently a hydrogen atom or a lower alkyl, and B is as defined above);

a group of the formula [7]:

45

[7]




50

[8]

(in which R¹⁶, R¹⁷, R⁵ and B are as defined above); or
a group of the formula [8]:

60

[8]



(B is as defined above).

The imide derivative of the present invention or an acid addition salt thereof may be prepared, for example, by the method disclosed in JP Patent No. 2800953 1 as mentioned above.

The imide derivative of the present invention may be used in the form of a pharmaceutically acceptable acid addition salt thereof. Inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, etc. or organic acids such as fumaric acid, citric acid, tartaric acid, succinic acid, etc. may be exemplified as an acid for forming addition salts.

The imide derivative or a pharmaceutically acceptable acid addition salt thereof, which is the active compound of the present invention, may be administered at a dose suitable for necessity of each case in a conventional dosage form. For example, it can be administered orally in the form of tablets, capsules, syrups, suspension, etc. or parenterally in the form of injection preparation such as liquid preparations (e.g., solutions, emulsions, suspension, patches, etc.).

In addition, the above-mentioned suitable dosage forms may be prepared by mixing an active compound with a conventional pharmaceutically acceptable carriers, excipients, binders, stabilizers, etc. When used in the form of injection, it may additionally contain buffering agents, solubilizers, isotonic agents, etc.

The dose and the frequency of the administration of the present therapeutic agent may vary according to the dosage forms, or the severity of the diseases to be treated. For example, the imide derivative is orally administered at a dose of 1 to 200 mg per day in an adult, which is administered once a day or divided into several dosage units.

The diseases to which the therapeutic agent of the present invention is effective are the Alzheimer type and the cerebrovascular dementia, more particularly, various senile dementias (e.g., dementia with Lewy bodies, dementia from Pick's disease, dementia from Creutzfeldt-Jakob disease, dementia from Huntington's chorea, dementia from Parkinson's disease, etc.) including multiple infarct dementia, dementia caused by cerebral infarction, Binswanger disease, dementia caused by stroke, amyloid angiopathy, ischemic dementia. Further, the therapeutic agent of the present invention shows the improving activity of cognitive dysfunctions accompanied by acetylcholine neuronal dysfunctions, and hence, it may be used in the treatment of traumatic cognitive dysfunctions, dementia in Down syndrome, schizophrenial cognitive dysfunctions, or cognitive dysfunctions accompanied by acetylcholine neuronal dysfunctions from any cause, in addition to the treatment of senile dementia.

EXAMPLES

The present invention is illustrated in more detail by Examples, but the present invention should not be construed to be limited thereto.

Example 1

Method

Male Wistar rats (7 weeks old) were used. As a test medicament, (1R,2S,3R,4S)—N-[(1R,2R)-2-[4-(1,2-benzisothiazol-3-yl)-1-piperazinyl-methyl]-1-cyclohexylmethyl]-2,3-bicycl[2.2.1]heptanedicarboxylimide (Compound A) was suspended in 0.5% methyl cellulose (MC) solution. As an agent for inducing amnesia, scopolamine (manufactured by Wako Pure Chemical Industries, Ltd., product No. 198-07901), which is an acetylcholine receptor blocker, was dissolved in saline solution (manufactured by TERUMO COR-

PORATION). Compound A at a dose of 3 mg/kg or 30 mg/kg, or 0.5% MC as a control was orally administered to the animals one hour prior to the training step in the one step-through passive avoidance test, and then, scopolamine at a dose of 0.5 mg/kg or a saline solution as a control was subcutaneously administered to the animals 30 minutes prior to the training step. The volume of each solution to be administered was 5 ml/kg each.

The one step-through passive avoidance test in rats was carried out in the following manners with using an apparatus consisting of a light-dark box and an electric stimulator (manufactured by O'hara & Co., Ltd., product no. PA-2030A) as an experimental apparatus. Namely, on Day 1, after the medicament and the agent for inducing amnesia were administered, the rats were put into the light box of the experimental apparatus where the back of each rat was directed to the dark box. Then, 10 seconds later, a guillotine door set at the border between the dark box and the light box was opened. Due to the habits of the rats, once the rats entered into the dark box, the guillotine door was quickly closed. At three seconds after the entering into the dark box, an electroconvulsive shock (0.5 mA, for 3 seconds) was given to the rats. Again, the guillotine door was opened, and after the rats spontaneously returned to the light box, the animals were transferred into the home cage. The period between the time just after the guillotine door was opened and the time at which the rats entered into the dark box was measured as a step-through latency. As to the animals which did not enter into the dark room even after 300 seconds, the training was terminated, and those animals were dropped in the following experiment for the reasons of training failure.

On Day 2 of the experiment, a test was carried out about 24 hours after the training. The procedures of the test were carried out in the same manner to the training step except that an electroconvulsive shock was not given. The step-through latency in the test was measured up to 300 seconds, and the step-through latency over 300 seconds was regarded as 300 seconds. FIG. 1 and FIG. 2 show the effects of Compound A on the scopolamine-induced cognitive/memory dysfunction models in the one step-through passive avoidance test when it was orally administered at a dosage of 3 mg/kg or 30 mg/kg. FIG. 1 shows the step-through latency during the training step, and FIG. 2 shows the step-through latency during the test. The number of the animals was 15 per group, and the data was expressed in mean±SEM.

(Results)

The agent for inducing amnesia, scopolamine did not affect on the step-through latency during the training. Compound A slightly shortened the step-through latency during the training step at a dose of 3 mg/kg. During the test, the animals treated with scopolamine showed a significantly shorter step-through latency as compared to the animals treated with saline solution (cognitive/memory dysfunction inducing effect). In the group treated with both of Compound A at a dose of 30 mg/kg and scopolamine, the step-through latency was significantly extended. That is, it was observed that Compound A exhibited an improving effect of scopolamine-induced cognitive/memory dysfunctions. Thus, it was found that the imide derivatives may exhibit an improving activity of scopolamine-induced cognitive/memory dysfunctions, and as a result, it may become apparent that the present invention may provide a therapeutic method for senile dementia and a therapeutic agent for said method.

INDUSTRIAL APPLICABILITY


According to the present invention, it was found that the imide derivatives may exhibit an improving activity of sco-

11


polamine-induced cognitive/memory dysfunctions, and as a result, it has become apparent that the present invention may provide a therapeutic method for senile dementia and a therapeutic agent for said method.

The invention claimed is:

1. A method for treatment of senile dementia, which comprises administering an effective amount of an imide derivative of the formula [1]:




wherein Z is a group of the formula [2]:



in which B is a carbonyl or a sulfonyl; R^1 , R^2 , R^3 and R^4 are independently a hydrogen atom or a lower alkyl, provided that R^1 and R^2 , or R^1 and R^3 may combine each other to form a hydrocarbon ring, or R^1 and R^3 may combine each other to form an aromatic hydrocarbon ring; said hydrocarbon ring may optionally be cross-linked with a lower alkylene or an oxygen atom; said lower alkylene and hydrocarbon ring may optionally be substituted by at least one alkyl; and n is 0 or 1,

D is a group of the formula [3]:



in which A is a hydrocarbon ring optionally be cross-linked with a lower alkylene or an oxygen atom; said lower alkylene or said hydrocarbon ring may optionally be substituted by at least one alkyl; and p and q are independently 0, 1 or 2,

G is N, CH or COH, and —Ar is an aromatic heterocyclic group, an aromatic hydrocarbon group, benzoyl, phenoxy, or phenylthio, or

G is a carbon atom, and —Ar is biphenylmethylidene, where said aromatic heterocyclic group aromatic hydrocarbon group, benzoyl, phenoxy, phenylthio, and biphenylmethylidene may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom,

or an acid addition salt thereof.


2. The method according to claim 1, wherein —Ar is an aromatic heterobicyclic group, naphthyl, benzoyl, phenoxy or phenylthio, and G is N, CH or COH, or —Ar is biphenylmethylidene, and G is a carbon atom, wherein said aromatic heterobicyclic group, naphthyl, benzoyl, phenoxy, phenylthio and biphenylmethylidene may optionally be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom.

3. The method according to claim 1, wherein —Ar is an aromatic heterocyclic group condensed with a benzene ring, or naphthyl, benzoyl, phenoxy or phenylthio, wherein said aromatic heterocyclic group condensed with a benzene ring, naphthyl, benzoyl, phenoxy, and phenylthio may optionally

12


be substituted by at least one group selected from a lower alkyl, a lower alkoxy and a halogen atom, and G is N, CH or COH.

4. The method according to claim 1, wherein Z is a group of the formula [4];




in which -L- is a single bond or a double bond, E is a lower alkylene optionally substituted by a lower alkyl, or an oxygen atom, R^5 is a hydrogen atom or a lower alkyl, and B is a carbonyl or a sulfonyl;

a group of the formula [5]:




in which -L- is a single bond or a double bond, E is a lower alkylene optionally substituted by a lower alkyl, or an oxygen atom, R^5 is a hydrogen atom or a lower alkyl, and B is a carbonyl or a sulfonyl;

a group of the formula [6]:



in which R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} are independently a hydrogen atom or a lower alkyl, or the adjacent two groups of R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} may combine each other to form a double bond, and B is a carbonyl or a sulfonyl;


a group of the formula [7]:



in which R^{16} and R^{17} are independently a hydrogen atom or a lower alkyl, or R^{16} and R^{17} may combine each other to form a saturated hydrocarbon ring, R^5 is a hydrogen atom or a lower alkyl, and B is a carbonyl or a sulfonyl; or

13

a group of the formula [8]:



in which B is a carbonyl or a sulfonyl.


5. The method according to claim 1, wherein the senile dementia is selected from the group consisting of Alzheimer type dementia, cerebrovascular dementia, dementia with Lewy bodies, dementia from Pick's disease, dementia from Creutzfeldt-Jakob disease, dementia from Huntington's chorea and dementia from Parkinson's disease.

6. The method according to claim 1, wherein the senile dementia is Alzheimer type dementia.

7. The method according to claim 1, wherein the senile dementia is selected from the group consisting of cerebrovascular dementia and dementia with Lewy bodies.

14

8. A method for treatment of senile dementia, which comprises administering an effective amount of an imide compound of the formula [9]:



or an acid addition salt thereof, and

wherein the senile dementia is selected from the group consisting of cerebrovascular dementia, dementia with Lewy bodies, dementia from Pick's disease, dementia from Creutzfeldt-Jakob disease and dementia from Huntington's chorea.

9. The method according to claim 8, wherein the senile dementia is selected from the group consisting of cerebrovascular dementia and dementia with Lewy bodies.

10. The method according to claim 8, wherein the senile dementia is cerebrovascular dementia.

11. The method according to claim 8, wherein the senile dementia is dementia with Lewy bodies.

* * * * *