



US 20080287364A1

(19) **United States**(12) **Patent Application Publication****Bufe et al.**(10) **Pub. No.: US 2008/0287364 A1**(43) **Pub. Date: Nov. 20, 2008**(54) **BITTER TASTE RECEPTORS****Publication Classification**

(76) Inventors: **Bernd Bufo**, Ferch (DE); **Thomas Hofmann**, Munster-Roxel (DE); **Dietmar Krautwurst**, Bergholz-Rehbrücke (DE); **Christina Kuhn**, Bergholz-Rehbrücke (DE); **Wolfgang Meyerhof**, Norderstedt (DE)

Correspondence Address:

SALIWANCHIK LLOYD & SALIWANCHIK
A PROFESSIONAL ASSOCIATION
PO BOX 142950
GAINESVILLE, FL 32614-2950 (US)

(51) **Int. Cl.**

A61K 38/00 (2006.01)
C12N 15/11 (2006.01)
C12N 15/00 (2006.01)
C12N 5/06 (2006.01)
A01K 67/027 (2006.01)
C12P 21/04 (2006.01)
C12N 15/87 (2006.01)
A23L 1/00 (2006.01)
A61P 43/00 (2006.01)
C07K 14/00 (2006.01)
C07K 16/18 (2006.01)
C07H 21/04 (2006.01)
G01N 33/53 (2006.01)
A61K 31/70 (2006.01)

(21) Appl. No.: **12/168,551**

(52) **U.S. Cl. 514/12**; 536/23.5; 435/320.1; 435/325;
800/13; 435/69.1; 435/455; 530/350; 530/387.9;
536/24.5; 435/7.2; 514/44; 426/531

(22) Filed: **Jul. 7, 2008****Related U.S. Application Data**

(62) Division of application No. 10/528,630, filed on Jun. 29, 2005, now Pat. No. 7,413,867, filed as application No. PCT/EP03/10691 on Sep. 25, 2003.

(60) Provisional application No. 60/413,298, filed on Sep. 25, 2002.

(57)

ABSTRACT

The present invention relates to bitter-taste receptors and their role in bitter taste transduction. The invention also relates to assays for screening molecules that modulate, e.g. suppress or block, bitter taste transduction, or enhance bitter taste response.

BITTER TASTE RECEPTORS

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application is a divisional application of co-pending application Ser. No. 10/528,630, filed Mar. 22, 2005; which is a National Stage Application of international Application Number PCT/EP2003/010691, filed Sep. 25, 2003; which claims the benefit of U.S. Provisional Application Ser. No. 60/413,298, filed Sep. 25, 2002, all of which are incorporated herein in their entirety.

BACKGROUND OF THE INVENTION

[0002] Investigators have recently turned their attention to understanding the biological mechanisms of taste, and in particular bitter taste. For a review of the literature see, for example, *Science* 291, 1557-1560. (2001); *Cell* 100, 607-610 (2000); *Neuron* 25, 507-510 (2000); *Nature* 413, 219-225. (2001); and *J Biol. Chem.* 277, 1-4 (2001).

[0003] Bitter taste is aversive, and as such provides humans with a mechanism of protection against poisonous substances, which are generally bitter-tasting compounds. More subtly, bitter-tastants also affect the palatability of food, beverages, thereby influencing human nutritional habits as is more fully discussed by Drewnowski in "The Science and Complexity of Bitter Taste", *Nutr. Rev.* 59, 163-169 (2001). They also affect the palatability of other ingestibles such as orally administered pharmaceuticals and nutraceuticals. Understanding the mechanism of bitter taste transduction has implications for the food and pharmaceutical industries. If the bitter taste transduction pathway can be manipulated, it may be possible to suppress or eliminate bitter taste to render foods more palatable and increase patient compliance in oral pharmaceuticals.

[0004] Taste transduction involves the interaction of molecules, i.e., tastants with taste receptor-expressing cells which reside in the taste buds located in the papillae of the tongue. Taste buds relay information to the brain on the nutrient content of food and the presence of poisons. Recent advances in biochemical and physiological studies have enabled researchers to conclude that bitter taste transduction is mediated by so-called G-protein coupled receptors (GPCRs). GPCRs are 7 transmembrane domain cell surface proteins that amplify signals generated at a cell surface when the receptor interacts with a ligand (a tastant) whereupon they activate heterotrimeric G-proteins. The G-proteins are protein complexes that are composed of alpha and beta-gamma subunits. They are usually referred to by their alpha subunits and classified generally into 4 groups: $G_{\alpha s}$, $G_{\alpha i}$, $G_{\alpha q}$ and $G_{\alpha 12}$. The $G_{\alpha q}$ type couple with GPCRs to activate phospholipase C which leads to the increase in cellular Ca^{2+} . There are many G_q -type G-proteins that are promiscuous and can couple to GPCRs, including taste receptors, and these so-called "promiscuous" G-proteins are well known to the man skilled in the art. These G-proteins dissociate into alpha and beta-gamma subunits upon activation, resulting in a complex cascade of cellular events that results in the cell producing cell messengers, such as calcium ions, that enable the cells to send a signal to the brain indicating a bitter response.

[0005] There is also anatomical evidence that GPCRs mediate bitter taste transduction: clusters of these receptors are found in mammalian taste cells containing gustducin. Gustducin is a G-protein subunit that is implicated in the

perception of bitter taste in mammals, see for example Chandrashekar, J. et al., *Cell* 100, 703-711 (2000); Matsunami H. et al., *Nature* 404, 601-604 (2000); or Adler E. et al., *Cell* 100, 693-702 (2000). cDNAs encoding such GPCRs have been identified, isolated, and used as templates to compare with DNA libraries using in-silico data-mining techniques to identify other related receptors. In this manner it has been possible to identify a family of related receptors, the so-called T2R family of receptors, that have been putatively assigned as bitter receptors.

[0006] To-date, however, it is not clear as to whether all the bitter taste receptors have been discovered. Further, of those that have been discovered, many have not been matched, or paired, with ligands, and applicant is aware of very few published studies wherein rigorous matching has been undertaken. Chandrashekar, J. et al. in *Cell* 100, 703-711 (2000), has expressed a human T2R receptor, the so-called hT2R4 receptor, in heterologous systems and looked at the in vitro response of this receptor. They found that it provided a response to the bitter compounds denatonium and 6-n-propyl-2-thiouracil. However, the concentrations of bitter tastants needed to activate the hT2R4 receptor were two orders of magnitude higher than the thresholds reported in human taste studies, and so it is not clear that the protein encoded by the hT2R4 gene is a functional bitter receptor. The authors of the Chandrashekar et al. article also looked at a number of mouse T2R receptors with a range of stock bitter-tasting chemicals of disparate chemical structure. However, no study has looked at receptor responses to bitter ligands that are problematic in the food and pharmaceutical industries, and means of suppressing the bitter response to these ligands.

[0007] The universe of compounds that provoke a bitter response in humans is structurally very diverse. Therefore, if research into bitter receptors is to be of any practical significance to the food and pharmaceutical industries, all bitter receptors will need to be identified, and once identified, there has to be a rigorous understanding of how specific receptors are matched to particular structural classes of bitter compounds. Unfortunately, although much basic research has been conducted in the area of bitter taste receptors, there are potentially many more bitter receptors to be discovered, and little is still known as to whether the known members of the human T2R family of bitter receptors actually respond to bitter tastants, and if so what, if any, specificity they show to ligand substructures.

BRIEF SUMMARY OF THE INVENTION

[0008] The present invention relates to bitter-taste receptors and their role in bitter taste transduction. The invention also relates to assays for screening molecules that modulate, e.g. suppress or block bitter taste transduction, or enhance bitter taste response.

DETAILED DESCRIPTION OF THE INVENTION

[0009] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. In case of conflict, the present document, including definitions, will control. Preferred methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used

in the practice or testing of the present invention. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety. The materials, methods, and examples disclosed herein are illustrative only and not intended to be limiting.

[0010] Other features and advantages of the invention, e.g., screening for compounds that inhibit bitter taste, will be apparent from the following description, from the drawings and from the claims.

[0011] Surprisingly, applicant has now found a new group of putative bitter taste receptors, and in respect of certain known bitter receptors, applicant has found that they respond with specificity towards classes of bitter compounds that are important in the food and pharmaceutical industries.

[0012] In a first aspect of the invention there is provided a new group of putative bitter receptors. The genetics of bitter tasting has been extensively studied in mice and rats. Therefore, applicant compared the nucleotide sequences encoding polypeptides previously proposed to be bitter receptors with publicly available human nucleic acid sequences in the NCBI database using the BLAST® search methodology (Parameters: Expect=0.01, Filter=default). Surprisingly, the search identified 24 DNA sequences (from human chromosomes 5, 7, and 12) that, because of their homology to a mouse nucleic acid sequence that encodes a polypeptide (T2R5) previously designated as a bitter receptor, we designated as bitter receptor-encoding. Bitter taste receptors were originally assigned identifiers starting with the three characters "T2R" (identifying the receptor family) followed by a number (e.g., 1, 2, 3, etc.) that identifies a particular receptor, e.g., T2R5. More recently a different system has been used in which the identifiers start with five characters "TAS2R" (identifying the receptor family) followed, as previously, by a number (e.g., 1, 2, 3, etc.) that identifies a particular receptor, e.g., TAS2R5. A lower case letter in front of the identifier indicates the species of the receptor (e.g., "h" for human, "r" for rat, and "m" for mouse). Thus, for example, mTAS2R5 is a mouse bitter receptor and hTAS2R2 is a human bitter receptor. For consistency the new TAS2R identifier system is used throughout the rest of this application.

[0013] Of the 24 coding sequences identified by the search, 12 are believed to be novel; the polypeptides encoded by these novel sequences are designated hTAS2R38-41, and 43-50. The DNA sequences encoding the polypeptides are assigned SEQ ID NOs: 2 (hTAS2R38), 4 (hTAS2R39), 6 (hTAS2R40), 8 (hTAS2R41), 10 (hTAS2R43), 12 (hTAS2R44), 14 (hTAS2R45), 16 (hTAS2R46), 18 (hTAS2R47), 20 (hTAS2R48), 22 (hTAS2R49), and 24 (hTAS2R50), respectively, and the amino acid sequences of the polypeptides are assigned SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, and 23, respectively.

[0014] The 12 additional (possibly novel) sequences identified by the search encode polypeptides, which are designated hTAS2R1, 4, 5, 7-10, 13, 14, 16, 3, 42 and 60. The DNA sequences encoding the polypeptides are assigned SEQ ID NOs: 26 (hTAS2R1), 28 (hTAS2R4), 30 (hTAS2R5), 32 (hTAS2R7), 34 (hTAS2R8), 36 (hTAS2R9), 38 (hTAS2R10), 40 (hTAS2R13), 42 (hTAS2R14), 44 (hTAS2R16), 46 (hTAS2R3), 48 (hTAS2R42), and 50 (hTAS2R60), respectively, and the amino acid sequences of the polypeptides are assigned SEQ ID NOs: 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47 and 49, respectively.

[0015] Thus, one aspect of the present invention is a polynucleotide selected from the group consisting of

[0016] (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NOs 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, and 23;

[0017] (b) polynucleotides having the coding sequence, as shown in SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, and 24 encoding at least the mature form of the polypeptide;

[0018] (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to said polypeptide, and said fragment or derivative has bitter substance binding activity;

[0019] (d) polynucleotides which are at least 50% identical to a polynucleotide as defined in any one of (a) to (c) and which code for a polypeptide having bitter substance binding activity; and

[0020] (e) polynucleotides the complementary strand of which hybridizes, preferably under stringent conditions to a polynucleotide as defined in any one of (a) to (d) and which code for a polypeptide having bitter substance binding activity;

or the complementary strand of such a polynucleotide.

[0021] A polypeptide that exhibits bitter substance binding activity is a polypeptide that has at least 20% (e.g., at least: 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100% or even more) of the ability of the respective full-length TAS2R to bind to a given bitter substance. Binding assays and bitter substances are described herein below.

[0022] In a preferred embodiment the polynucleotide of the present invention encodes a polypeptide that still exhibits essentially the same activity as the respective mature bitter taste receptor, i.e. has "bitter taste receptor activity". Preferably the polypeptide has at least 20% (e.g., at least: 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100% or even more) of the ability of the respective full-length TAS2R to release intracellular calcium in a heterologous cell expression system like, for example, HEK293/15-cells, which stably express the alpha-subunit of promiscuous G-proteins, e.g. the mouse G₁₅ subunit, in response to bitter tastants, which is dependent on the expression of polypeptides encoded by the polynucleotides of the present invention. The amount of intracellular calcium release can be monitored by, for example, the in vitro FLIPR assay described herein below.

[0023] The TAS2R nucleic acid molecules of the invention can be DNA, cDNA, genomic DNA, synthetic DNA, or RNA, and can be double-stranded or single-stranded, the sense and/or an antisense strand. Segments of these molecules are also considered within the scope of the invention, and can be produced by, for example, the polymerase chain reaction (PCR) or generated by treatment with one or more restriction endonucleases. A ribonucleic acid (RNA) molecule can be produced by in vitro transcription.

[0024] The polynucleotide molecules of the invention can contain naturally occurring sequences, or sequences that differ from those that occur naturally, but, due to the degeneracy of the genetic code, encode the same polypeptide (for example, the polypeptides with SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, and 23). In addition, these nucleic acid

molecules are not limited to coding sequences, e.g., they can include some or all of the non-coding sequences that lie upstream or downstream from a coding sequence.

[0025] The polynucleotide molecules of the invention can be synthesized *in vitro* (for example, by phosphoramidite-based synthesis) or obtained from a cell, such as the cell of a bacteria mammal. The nucleic acids can be those of a human but also derived from a non-human primate, mouse, rat, guinea pig, cow, sheep, horse, pig, rabbit, dog, or cat as long as they fulfill the criteria set out above. Combinations or modifications of the nucleotides within these types of nucleic acids are also encompassed.

[0026] In addition, the isolated nucleic acid molecules of the invention encompass segments that are not found as such in the natural state. Thus, the invention encompasses recombinant nucleic acid molecules incorporated into a vector (for example, a plasmid or viral vector) or into the genome of a heterologous cell (or the genome of a homologous cell, at a position other than the natural chromosomal location). Recombinant nucleic acid molecules and uses therefore are discussed further below.

[0027] A polynucleotide belonging to a family of any of the TAS2R disclosed herein or a protein can be identified based on its similarity to the relevant TAS2R gene or protein, respectively. For example, the identification can be based on sequence identity. In certain preferred embodiments the invention features isolated nucleic acid molecules which are at least 50% (or 55%, 65%, 75%, 85%, 95%, or 98%) identical to: (a) a nucleic acid molecule that encodes the polypeptide of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, or 23; (b) the nucleotide sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, or 24; and (c) a nucleic acid molecule which includes a segment of at least 30 (e.g., at least 30, 40, 50, 60, 80, 100, 125, 150, 175, 200, 250, 300, 400, 500, 600, 700, 800, 850, 900, 950, 1000, or 1010) nucleotides of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, or 24.

[0028] The determination of percent identity between two sequences is accomplished using the mathematical algorithm of Karlin and Altschul, *Proc. Natl. Acad. Sci. USA* 90, 5873-5877, 1993. Such an algorithm is incorporated into the BLASTN and BLASTP programs of Altschul et al. (1990) *J. Mol. Biol.* 215, 403-410. BLAST nucleotide searches are performed with the BLASTN program, score=100, wordlength=12, to obtain nucleotide sequences homologous to HIN-1-encoding nucleic acids. BLAST protein searches are performed with the BLASTP program, score=50, wordlength=3, to obtain amino acid sequences homologous to the TAS2R polypeptide. To obtain gapped alignments for comparative purposes, Gapped BLAST is utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25, 3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs.

[0029] Hybridization can also be used as a measure of homology between two nucleic acid sequences. A nucleic acid sequence encoding any of the TAS2R disclosed herein, or a portion thereof, can be used as a hybridization probe according to standard hybridization techniques. The hybridization of a TAS2R probe to DNA or RNA from a test source (e.g., a mammalian cell) is an indication of the presence of the relevant TAS2R DNA or RNA in the test source. Hybridization conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y., 6.3.1-6.3.6, 1991. Moderate hybridization conditions are defined as equivalent to hybridization in 2×

sodium chloride/sodium citrate (SSC) at 30° C., followed by a wash in 1 X SSC, 0.1% SDS at 50° C. Highly stringent conditions are defined as equivalent to hybridization in 6×

[0030] An "isolated DNA" is either (1) a DNA that contains sequence not identical to that of any naturally occurring sequence, or (2), in the context of a DNA with a naturally-occurring sequence (e.g., a cDNA or genomic DNA), a DNA free of at least one of the genes that flank the gene containing the DNA of interest in the genome of the organism in which the gene containing the DNA of interest naturally occurs. The term therefore includes a recombinant DNA incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote. The term also includes a separate molecule such as a cDNA where the corresponding genomic DNA has introns and therefore a different sequence; a genomic fragment that lacks at least one of the flanking genes; a fragment of cDNA or genomic DNA produced by polymerase chain reaction (PCR) and that lacks at least one of the flanking genes; a restriction fragment that lacks at least one of the flanking genes; a DNA encoding a non-naturally occurring protein such as a fusion protein, mutein, or fragment of a given protein; and a nucleic acid which is a degenerate variant of a cDNA or a naturally occurring nucleic acid. In addition, it includes a recombinant nucleotide sequence that is part of a hybrid gene, i.e., a gene encoding a non-naturally occurring fusion protein. It will be apparent from the foregoing that isolated DNA does not mean a DNA present among hundreds to millions of other DNA molecules within, for example, cDNA or genomic DNA libraries or genomic DNA restriction digests in, for example, a restriction digest reaction mixture or an electrophoretic gel slice.

[0031] A further aspect of the present invention is a vector containing the polynucleotide(s) of the present invention or a protein encoded by a polynucleotide of the present invention. The term "vector" refers to a protein or a polynucleotide or a mixture thereof which is capable of being introduced or of introducing the proteins and/or nucleic acid comprised into a cell. It is preferred that the proteins encoded by the introduced polynucleotide are expressed within the cell upon introduction of the vector.

[0032] In a preferred embodiment the vector of the present invention comprises plasmids, phagemids, phages, cosmids, artificial mammalian chromosomal constructs, or knock-in constructs, viruses, in particular adenoviruses, vaccinia viruses, attenuated vaccinia viruses, canary pox viruses, lentivirus (Chang, L. J. and Gay, E. E. (2000) *Curr. Gene Therap.* 1:237-251), herpes viruses, in particular Herpes simplex virus (HSV-1, Carlezon, W. A. et al. (2000) *Crit. Rev. Neurobiol.*), baculovirus, retrovirus, adeno-associated-virus (AAV, Carter, P. J. and Samulski, R. J. (2000) *J. Mol. Med.* 6:17-27), rhinovirus, human immune deficiency virus (HIV), filovirus and engineered versions thereof (see, for example, Cobinger G. P. et al (2001) *Nat. Biotechnol.* 19:225-30), virosomes, "naked" DNA liposomes, and nucleic acid coated particles, in particular gold spheres. Particularly preferred are viral vectors like adenoviral vectors or retroviral vectors (Lindemann et al. (1997) *Mol. Med.* 3:466-76 and Springer et al. (1998) *Mol. Cell.* 2:549-58). Liposomes are usually small unilamellar or multilamellar vesicles made of cationic, neutral and/or anionic lipids, for example, by ultrasound treatment of liposomal suspensions. The DNA can, for example,

be ionically bound to the surface of the liposomes or internally enclosed in the liposome. Suitable lipid mixtures are known in the art and comprise, for example, DOTMA (1,2-Dioleoyloxypropyl-3-trimethylammoniumbromide) and DPOE (Dioleoylphosphatidyl-ethanolamine) which both have been used on a variety of cell lines.

[0033] Nucleic acid coated particles are another means for the introduction of nucleic acids into cells using so called "gene guns", which allow the mechanical introduction of particles into the cells. Preferably the particles itself are inert, and therefore, are in a preferred embodiment made out of gold spheres.

[0034] In a further aspect the polynucleotide of the present invention is operatively linked to expression control sequences allowing expression in prokaryotic and/or eukaryotic host cells. The transcriptional/translational regulatory elements referred to above include but are not limited to inducible and non-inducible, constitutive, cell cycle regulated, metabolically regulated promoters, enhancers, operators, silencers, repressors and other elements that are known to those skilled in the art and that drive or otherwise regulate gene expression. Such regulatory elements include but are not limited to regulatory elements directing constitutive expression like, for example, promoters transcribed by RNA polymerase III like, e.g., promoters for the snRNA U6 or scRNA 7SK gene, the cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40 adenovirus, viral promoter and activator sequences derived from, e.g., NBV, HCV, HSV, HPV, EBV, HTLV, MMTV or HIV; which allow inducible expression like, for example, CUP-1 promoter, the tet-repressor as employed, for example, in the tet-on or tet-off systems, the lac system, the trp system; regulatory elements directing tissue specific expression, preferably taste bud specific expression, e.g., PLC β 2 promoter or gust-ducin promoter, regulatory elements directing cell cycle specific expression like, for example, cdc2, cdc25C or cyclin A; or the TAC system, the TRC system, the major operator and promoter regions of phage λ , the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast a- or a-mating factors.

[0035] As used herein, "operatively linked" means incorporated into a genetic construct so that expression control sequences effectively control expression of a coding sequence of interest.

[0036] Similarly, the polynucleotides of the present invention can form part of a hybrid gene encoding additional polypeptide sequences, for example, a sequence that functions as a marker or reporter. Examples of marker and reporter genes include β -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neo^r, G418^r), dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding β -galactosidase), and xanthine guanine phosphoribosyltransferase (XGPRT). As with many of the standard procedures associated with the practice of the invention, skilled artisans will be aware of additional useful reagents, for example, additional sequences that can serve the function of a marker or reporter. Generally, the hybrid polypeptide will include a first portion and a second portion; the first portion being a TAS2R polypeptide and the second portion being, for example, the reporter described above or an Ig constant region or part of an Ig constant region, e.g., the CH2 and CH3 domains of IgG2a heavy chain. Other

hybrids could include an antigenic tag or His tag to facilitate purification and/or detection. Recombinant nucleic acid molecules can also contain a polynucleotide sequence encoding a TAS2R polypeptide operatively linked to a heterologous signal sequence. Such signal sequences can direct the protein to different compartments within the cell and are well known to someone of skill in the art. A preferred signal sequence is a sequence that facilitates secretion of the resulting protein.

[0037] Another aspect of the present invention is a host cell genetically engineered with the polynucleotide or the vector as outlined above. The host cells that may be used for purposes of the invention include but are not limited to prokaryotic cells such as bacteria (for example, *E. coli* and *B. subtilis*), which can be transformed with, for example, recombinant bacteriophage DNA, plasmid DNA, or cosmid DNA expression vectors containing the polynucleotide molecules of the invention; simple eukaryotic cells like yeast (for example, *Saccharomyces* and *Pichia*), which can be transformed with, for example, recombinant yeast expression vectors containing the polynucleotide molecule of the invention; insect cell systems like, for example, Sf9 of Hi5 cells, which can be infected with, for example, recombinant virus expression vectors (for example, baculovirus) containing the polynucleotide molecules of the invention; *Xenopus* oocytes, which can be injected with, for example, plasmids; plant cell systems, which can be infected with, for example, recombinant virus expression vectors (for example, cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV)) or transformed with recombinant plasmid expression vectors (for example, Ti plasmid) containing a TAS2R nucleotide sequence; or mammalian cell systems (for example, COS, CHO, BHK, HEK293, VERO, HeLa, MDCK, Wi38, and NIH 3T3 cells), which can be transformed with recombinant expression constructs containing, for example, promoters derived, for example, from the genome of mammalian cells (for example, the metallothionein promoter) from mammalian viruses (for example, the adenovirus late promoter and the vaccinia virus 7.5K promoter) or from bacterial cells (for example, the tet-repressor binding site employed in the tet-on and tet-off systems). Also useful as host cells are primary or secondary cells obtained directly from a mammal and transfected with a plasmid vector or infected with a viral vector. Depending on the host cell and the respective vector used to introduce the polynucleotide of the invention the polynucleotide can integrate, for example, into the chromosome or the mitochondrial DNA or can be maintained extrachromosomally like, for example, episomally or can be only transiently comprised in the cells.

[0038] In a preferred embodiment, the TAS2R encoded by the polynucleotides of the present invention and which are expressed by such cells are functional, i.e., upon binding to one or more bitter molecules they trigger an activation pathway in the cell. The cells are preferably mammalian (e.g., human, non-human primate, horse, bovine, sheep, goat, pig, dog, cat, goat, rabbit mouse, rat, guinea pig, hamster, or gerbil) cells, insect cells, bacterial cells, or fungal (including yeast) cells.

[0039] A further aspect of the present invention is a transgenic non-human animal containing a polynucleotide, a vector and/or a host cell as described above. The animal can be a mosaic animal, which means that only part of the cells making up the body comprise polynucleotides, vectors, and/or cells of the present invention or the animal can be a transgenic animal which means that all cells of the animal comprise the

polynucleotides and/or vectors of the present invention or are derived from a cell of the present invention. Mosaic or transgenic animals can be either homo- or heterozygous with respect to the polynucleotides of the pre-sent invention contained in the cell. In a preferred embodiment the transgenic animals are either homo- or heterozygous knock-out or knock-in animals with respect to the genes which code for the proteins of the present invention. The animals can in principal be any animal, preferably, however, it is a mammal, selected from the group of non-human pimate horse, bovine, sheep, goat, pig, dog, cat, rabbit, mouse, rat, guinea pig, hamster, or gerbil. Another aspect of the present invention is a process for producing a polypeptide encoded by a polynucleotide of the present invention comprising: culturing the host cell described above and recovering the polypeptide encoded by said polynucleotide. Preferred combinations of host cells and vectors are outlined above and further combination will be readily apparent to someone of skill in the art. Depending on the intended later use of the recovered peptide a suitable cell type can be chosen. Eukaryotic cells are preferably chosen, if it is desired that the proteins produced by the cells exhibit an essentially natural pattern of glycosylation and prokaryotic cells are chosen, if, for example, glycosylation or other modifications, which are normally introduced into proteins only in eukaryotic cells, are not desired or not needed.

[0040] A further aspect of the invention is a process for producing cells capable of expressing at least one of the bitter taste receptor polypeptides comprising genetically engineering cells in vitro with at least one of the vectors described above, wherein said bitter taste receptor polypeptide(s) is(are) encoded by a polynucleotide of the present invention. Another aspect of the invention is a polypeptide having the amino acid sequence encoded by a polynucleotide of the invention or obtainable by the process mentioned above. The polypeptides of the invention include all those disclosed herein and functional fragments of these polypeptides. "Polypeptide" and "protein" are used interchangeably and mean any peptidelinked chain of amino acids, regardless of length or posttranslational modification. As used herein, a functional fragment of a TAS2R is a fragment of the TAS2R that is shorter than the full-length TAS2R but that has at least 20% (e.g., at least: 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100% or even more) of the ability of the full-length TAS2R to bind to a bitter substance to which the full-length TAS2R binds. Binding assays and bitter substances are described herein. Further bitter substances can be identified by the binding assays and bitter taste receptor activity assays described herein. The polypeptides embraced by the invention also include fusion proteins that contain either a full-length TAS2R polypeptide or a functional fragment of it fused to an unrelated amino acid sequence. The unrelated sequences can be additional functional domains or signal peptides. Signal peptides are described in greater detail and exemplified below.

[0041] The polypeptides can be any of those described above but with not more than 50 (e.g., not more than: 50, 45, 40, 35, 30, 25, 20, 15, 14, 13, 12, 11, 10, nine, eight, seven, six, five, four, three, two, or one) conservative substitutions. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine, and leucine; aspartic acid and glutamic acid; asparagine, glutamine, serine and threonine; lysine, histidine and arginine; and phenylalanine and tyrosine. All that is required of a polypeptide having one or more conservative substitutions is

that it has at least 20% (e.g., at least: 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100% or even more) of the ability of the wild-type, full-length TAS2R to bind to a bitter substance, preferably the ability to release intracellular calcium, when expressed in a cellular system.

[0042] The polypeptides can be purified from natural sources (e.g., blood, serum, plasma, tissues or cells such as normal tongue cells or any cell that naturally produces the relevant TAS2R polypeptides). Smaller peptides (less than 50 amino acids long) can also be conveniently synthesized by standard chemical means. In addition, both polypeptides and peptides can be produced by standard in vitro recombinant DNA techniques and in vivo transgenesis, using nucleotide sequences encoding the appropriate polypeptides or peptides. Methods well-known to those skilled in the art can be used to construct expression vectors containing relevant coding sequences and appropriate transcriptional/translational control signals. See, for example, the techniques described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2nd Ed.) [Cold Spring Harbor Laboratory, N.Y., 1989], and Ausubel et al., *Current Protocols in Molecular Biology* [Green Publishing Associates and Wiley Interscience, N.Y., 1989].

[0043] Polypeptides and fragments of the invention also include those described above, but modified for in vivo use by the addition, at the amino- and/or carboxyl-terminal ends, of blocking agents to facilitate survival of the relevant polypeptide in vivo. This can be useful in those situations in which the peptide termini tend to be degraded by proteases prior to cellular uptake. Such blocking agents can include, without limitation, additional related or unrelated peptide sequences that can be attached to the amino and/or carboxyl terminal residues of the peptide to be administered. This can be done either chemically during the synthesis of the peptide or by recombinant DNA technology by methods familiar to artisans of average skill.

[0044] Alternatively, blocking agents such as pyroglutamic acid or other molecules known in the art can be attached to the amino and/or carboxyl terminal residues, or the amino group at the amino terminus or carboxyl group at the carboxyl terminus can be replaced with a different moiety. Likewise, the peptides can be covalently or noncovalently coupled to pharmaceutically acceptable "carrier" proteins prior to administration.

[0045] Also of interest are peptidomimetic compounds that are designed based upon the amino acid sequences of the functional peptides or peptide fragments. Peptidomimetic compounds are synthetic compounds having a three-dimensional conformation (i.e., a "peptide motif") that is substantially the same as the three-dimensional conformation of a selected peptide. The peptide motif provides the peptidomimetic compound with the ability to bind to a bitter compound in a manner qualitatively identical to that of the TAS2R functional fragment from which the peptidomimetic was derived. Peptidomimetic compounds can have additional characteristics that enhance their therapeutic utility, such as increased cell permeability and prolonged biological half-life.

[0046] The peptidomimetics typically have a backbone that is partially or completely nonpeptide, but with side groups that are identical to the side groups of the amino acid residues that occur in the peptide on which the peptidomimetic is based. Several types of chemical bonds, e.g., ester, thioester, thioamide, retroamide, reduced carbonyl, dimethylene and ketomethylene bonds, are known in the art to be generally

useful substitutes for peptide bonds in the construction of protease-resistant peptidomimetics.

[0047] The term “isolated” polypeptide or peptide fragment as used herein refers to a polypeptide or a peptide fragment which either has no naturally-occurring counterpart or has been separated or purified from components which naturally accompany it, e.g., in tissues such as tongue, pancreas, liver, spleen, ovary, testis, muscle, joint tissue, neural tissue, gastrointestinal tissue or tumor tissue, or body fluids such as blood, serum, or urine. Typically, the polypeptide or peptide fragment is considered “isolated” when it is at least 70%, by dry weight, free from the proteins and other naturally-occurring organic molecules with which it is naturally associated. Preferably, a preparation of a polypeptide (or peptide fragment thereof) of the invention is at least 80%, more preferably at least 90%, and most preferably at least 99%, by dry weight, the polypeptide (or the peptide fragment thereof), respectively, of the invention. Thus, for example, a preparation of polypeptide x is at least 80%, more preferably at least 90%, and most preferably at least 99%, by dry weight, polypeptide x. Since a polypeptide that is chemically synthesized is, by its nature, separated from the components that naturally accompany it, the synthetic polypeptide is “isolated.”

[0048] An isolated polypeptide (or peptide fragment) of the invention can be obtained, for example, by extraction from a natural source (e.g., from tissues or bodily fluids); by expression of a recombinant nucleic acid encoding the polypeptide; or by chemical synthesis. A polypeptide that is produced in a cellular system different from the source from which it naturally originates is “isolated,” because it will necessarily be free of components which naturally accompany it. The degree of isolation or purity can be measured by any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

[0049] A further aspect of the invention is an antibody, which specifically binds to the polypeptide encoded by a polynucleotide of the invention or obtainable by the process mentioned above. The term “antibody” comprises monoclonal and polyclonal antibodies and binding fragments thereof, in particular Fc-fragments as well as so called “single-chain-antibodies” (Bird R. E. et al (1988) Science 242:423-6), chimeric, humanized, in particular CDR-grafted antibodies, and dia or tetrabodies (Holliger P. et al (1993) Proc. Natl. Acad. Sci. U.S.A. 90:6444-8). Also comprised are immunoglobulin like proteins that are selected through techniques including, for example, phage display to specifically bind to the polypeptides of the present invention. Preferred antibodies bind to the extracellular domain of bitter receptors and in particular to those domains responsible for binding to bitter tastants.

[0050] In yet another embodiment there is provided a molecule, or collections of molecules containing a molecule, that act to antagonise aforementioned receptors in particular the bitter taste response, and methods for screening for such molecules.

[0051] Therefore, a further aspect of the invention is a nucleic acid molecule which specifically hybridizes to a polynucleotide of the present invention. In particular this nucleic acid molecule is an inhibiting RNA. Preferred inhibiting RNAs are antisense constructs hybridizing to a polynucleotide of the present invention, RNAi, siRNA or a ribozyme. The design of such inhibiting RNAs would be readily apparent to someone of skill in the art.

[0052] Another type of antagonist/inhibitor against the polypeptides of the present invention is an antibody, which is preferably directed against the extracellular domain of the

respective bitter taste receptor and even more preferably binds to the site(s) of the receptor that interact(s) with the bitter substance(s) essentially without triggering the release of intracellular calcium. Further antagonists to the bitter taste response of a receptor are fragments of the receptor which have the capability to bind to the bitter substances as defined above. Such fragments can bind to the bitter substance and, thus, competitively antagonize the activity of the respective TAS2R. If such antagonists are, for example, employed within foodstuff to suppress the bitter taste of a specific bitter substance they might be exposed to a proteolytic environment and in this case the modifications of the polypeptides outlined above could be used to stabilize the competitive bitter receptor antagonist. However, various additional modifications, which stabilize such fragments will be readily apparent to the skilled person.

[0053] Antagonists and agonists of the bitter taste receptors described herein are of great importance for specific stimulation of a given bitter taste receptor or to antagonize it. The bitter taste response of the receptor is elicited by the specific binding of the respective bitter substance. Therefore, the present invention is also directed at a process for isolating a compound that binds to a polypeptide encoded by a polynucleotide selected from the group consisting of:

[0054] (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NOs 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47 and 49;

[0055] (b) polynucleotides having the coding sequence, as shown in SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48 and 50 encoding at least the mature form of the polypeptide;

[0056] (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to said polypeptide, and said fragment or derivative has bitter substance binding activity;

[0057] (d) polynucleotides which are at least 50% identical to a polynucleotide as defined in any one of (a) to (c) and which code for a polypeptide having bitter substance binding activity; and

[0058] (e) polynucleotides the complementary strand of which hybridizes, preferably under stringent conditions to a polynucleotide as defined in any one of (a) to (d) and which code for a polypeptide having bitter substance binding activity;

comprising:

[0059] (1) contacting said polypeptide or a host cell genetically engineered with said polynucleotide or with a vector containing said polynucleotide with a compound;

[0060] (2) detecting the presence of the compound which binds to said polypeptide; and

[0061] (3) determining whether the compound binds said polypeptide.

[0062] A polynucleotide employed in this process is in preferred embodiments of the invention at least 50% (or 55%, 65%, 75%, 85%, 95%, or 98%) identical to: (a) a nucleic acid molecule that encodes the polypeptide of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47 or 49; (b) the nucleotide sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48 or 50; and has a length of at least

30 (e.g., at least 30, 40, 50, 60, 80, 100, 125, 150, 175, 200, 250, 300, 400, 500, 600, 700, 800, 850, 900, 950, 1000, or 1010) of the nucleotides of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, or 50.

[0063] Furthermore for all of the above described hTAS2Rs, which can be employed in a process for isolating

binding compounds, with the exception of hTAS2R40, single nucleotide polymorphisms are known. 79 of these are listed in Table I below. 61 of those result in an amino acid change. Polynucleotides or polypeptides that differ from the respectively in SEQ ID 1-50 indicated sequences by the nucleotide and amino acid change as indicated in Table I can similarly be employed for the process of the present invention.

TABLE I

Gen +		Substitution		Position		
Accession No.	Name of SNP	Base	Amino acid	Base pair	Amino acid	Allelic frequency
hTAS2R1	rs2234231	C/T	P/L	128	43	unknown
NM019599	rs41469	G/A	R/H	332	111	A 0.46/G 0.54
	rs223432	G/A	C/Y	422	141	unknown
	rs2234233	C/T	R/W	616	206	C 0.87/T 0.13
	rs2234234	C/T	S/S	675	225	unknown
	rs2234235	T/C	L/L	850	284	unknown
hTAS2R3	rs227009	C/T	G/G	807	369	unknown
NM016943						
hTAS2R4	ss3181498	G/A	R/Q	8	3	unknown
NM016944	rs2233996	G/C	R/R	9	3	unknown
	rs2233997	A/C	Y/C	17	6	unknown
	rs2233998	T/C	F/S	20	7	unknown
	rs2233999	T/A	F/L	186	62	unknown
	rs2234000	C/T	T/M	221	74	C 0.94/T 0.56
	rs2234001	G/C	V/L	286	96	C 0.78/G 0.22
	rs2234002	G/A	S/N	512	171	A 0.78/G 0.22
	rs2234003	A/G	I/V	571	191	unknown
	rs2234013	G/A	G/S	58	20	unknown
	rs2227264	G/T	S/I	77	26	unknown
hTAS2R5	rs2234014	C/T	P/L	338	113	unknown
NM018980	rs2234015	G/A	R/Q	638	213	unknown
	rs2234016	G/T	R/L	294	881	unknown
	rs3759251	A/T	T/S	787	263	A 0.97/T 0.03
	rs3759252	C/A	I/I	828	276	unknown
	rs619381	G/A	M/I	912	304	unknown
hTAS2R8	ss2391467	G/A	L/L	549	183	unknown
NM023918	rs2537817	A/G	M/V	922	308	unknown
hTAS2R9	rs3741845	T/C	V/A	560	187	C 0.73/T 0.27
NM23917	rs3944035	C/T	L/F	910	304	unknown
	rs2159903	C/T	P/L	926	309	unknown
	rs597468	C/T	T/M	467	156	unknown
hTAS2R10						
NM23921						
hTAS2R13	ss1478988	A/G	N/S	776	259	C 0.73/T 0.27
NM23920						
hTAS2R14	rs3741843	G/A	R/R	375	125	A 0.97/G 0.03
NM23922						
hTAS2R16	rs2233988	C/T	T/T	300	100	unknown
NM016945	rs2692396	G/C	V/V	303	101	unknown
	rs2233989	T/C	L/L	460	154	unknown
	rs846664	T/G	N/K	516	172	A 0.71/C 0.29
	rs860170	G/A	R/H	665	222	A 0.55/G 0.45
	PTC Paper	G/A	V/I	886	296	G 0.38/A 0.62
hTAS2R38	rs1726866	T/C	V/A	785	262	G 0.38/T 0.62
AF494321	rs713598	C/T	A/P	49	145	C 0.36/G 0.64
	hTAS2R38 SNP1	A/T	N/I	557	186	C 0.60/G 0.40
	hTAS2R39 SNP1	A/AA	frameshift	967	323	unknown
hTAS2R39						
AF494230						
hTAS2R41	rs1404635	A/G	T/T	189	64	unknown
AF494232	hTAS2R41 SNP1	T/C	L/P	380	127	unknown
	hTAS2R41 SNP2	A/G	S/S	885	295	unknown
hTAS2R42	rs1650017	G/C	A/P	931	311	unknown
AX097739	rs1669411	T/C	N/N	930	310	unknown
	rs1669412	G/A	R/Q	875	292	unknown
	rs1451772	A/G	Y/C	794	265	unknown
	rs1669413	G/T	G/W	763	255	unknown
	rs1650019	A/G	L/L	561	187	unknown
	rs3759246	G/C	R/T	893	298	unknown
	hTAS2R43 SNP1	C/G	S/W	104	35	unknown
hTAS2R43	hTAS2R43 SNP2	G/A	R/H	635	212	unknown
AF494237	hTAS2R43 SNP3	G/C	T/T	663	221	unknown

TABLE I-continued

Gen +		Substitution		Position		Allelic frequency
Accession No.	Name of SNP	Base	Amino acid	Base pair	Amino acid	
hTAS2R44	rs3759247	G/A	W/stop	900	300	unknown
AF494228	rs3759246	G/C	R/T	893	298	unknown
	hTAS2R44 SNP1	A/T	M/L	162	484	unknown
	hTAS2R44 SNP2	T/A	F/Y	869	290	unknown
	hTAS2R44 SNP3	G/A	V/M	899	297	unknown
hTAS2R45	rs3759247	A/G	G/stop	900	300	unknown
AF494226	rs3759246	G/C	R/T	893	298	unknown
	rs3759245	C/T	R/C	712	238	unknown
	rs3759244	T/C	F/L	703	235	unknown
hTAS2R46	rs2708381	G/A	W/stop	749	250	unknown
AF494227	rs2708380	T/A	L/M	682	228	unknown
	rs2598002	T/G	F/V	106	36	unknown
	hTAS2R46 SNP1	A/T	Q/H	888	296	unknown
	hTAS2R46 SNP2	A/G	M/V	889	297	unknown
	hTAS2R46 SNP3	T/C	F/F	108	36	unknown
hTAS2R47	rs2597924	G/A	R/H	920	307	unknown
AF494233	rs1669405	T/G	L/W	842	281	unknown
	rs2599404	T/G	F/L	756	252	unknown
	rs2600355	T/G	V/V	54	18	unknown
hTAS2R48	rs1868769	T/C	L/L	418	140	unknown
AF494234						
hTAS2R49	hTAS2R49	A/G	K/R	164	55	unknown
AF494236	SNP1					
hTAS2R50	rs1376521	A/G	Y/C	608	203	G 0.66/A0.34
AF494235	hTAS2R50 SNP1	A/G	P/P	777	259	unknown

[0064] A polypeptide that exhibits bitter substance binding activity is a polypeptide that has at least 20% (e.g., at least: 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100% or even more) of the ability of the respective full-length TAS2R to bind to a given bitter substance. Binding assays and bitter substances are described herein.

[0065] The term “contacting” in the context of the present invention means any interaction between the compound with the polypeptide of the invention, whereby any of the at least two components can be independently of each other in a liquid phase, for example in solution, or in suspension or can be bound to a solid phase, for example, in the form of an essentially planar surface or in the form of particles, pearls or the like. In a preferred embodiment a multitude of different compounds are immobilized on a solid surface like, for example, on a compound library chip and the protein of the present invention is subsequently contacted with such a chip. In another preferred embodiment the cells genetically engineered with the polynucleotide of the invention or with a vector containing such a polynucleotide express the bitter taste receptor at the cell surface and are contacted separately in small containers, e.g., microtitre plates, with various compounds.

[0066] Detecting the presence and the binding of the compound to the polypeptide can be carried out, for example, by measuring a marker that can be attached either to the protein or to the compound. Suitable markers are known in the art and comprise, for example, fluorescence, enzymatic or radioactive markers. The binding of the two components can, however, also be measured by the change of an electrochemical parameter of the binding compound or of the protein, e.g. a change of the redox properties of either the protein or the binding compound, upon binding. Suitable methods of detecting such changes comprise, for example, potentiomet-

ric methods. Further methods for detecting and/or measuring the binding of the two components to each other are known in the art and can without limitation also be used to measure the binding of the compound to the polypeptide. The effect of the binding of the compound on the activity of the polypeptide can also be measured by assessing changes in the cells that express the polypeptides, for example, by assaying the intracellular release of calcium upon binding of the compound.

[0067] As a further step after measuring the binding of a compound and after having measured the binding strength of at least two different compounds at least one compound can be selected, for example, on grounds of a higher binding strength or on grounds of the detected intracellular release of calcium.

[0068] The thus selected compound is then in a preferred embodiment modified in a further step. Modification can be effected by a variety of methods known in the art, which include without limitation the introduction of one or more, preferably two, three or four novel side chains or residues or the exchange of one or more functional groups like, for example, introduction or exchange of halogens, in particular F, Cl or Br; the introduction or exchange of lower alkyl residues, preferably having one to five carbon atoms like, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or iso-pentyl residues; lower alkenyl residues, preferably having two, three, four or five carbon atoms; lower alkynyl residues, preferably having two, three, four or five carbon atoms, which can in a preferred embodiment be further substituted with F, Cl, Br, NH₂, NO₂, OH, SH, NH, CN, aryl, heteroaryl, COH or COOH group; or the introduction of, for example, one or more residue(s) selected from the group consisting of NH₂, NO₂, OH, SH, NH, CN, aryl, alkylaryl, heteroaryl, alkylheteroaryl, COH or COOH group.

[0069] The thus modified binding substances are then individually tested with the method of the pre-sent invention, i.e.

they are contacted with the polypeptide as such or with the polypeptide expressed in a cell, and subsequently binding of the modified compounds is measured. In this step both the binding per se can be measured and/or the effect of the function of the protein like, e.g. the intracellular calcium release. If needed the steps of selecting the compound, modifying the compound, contacting the compound with a polypeptide of the invention and measuring the binding of the modified compound to the polypeptide can be repeated a third or any given number of times as required. The above described method is also termed "directed evolution" of the compound since it involves a multitude of steps including modification and selection, whereby binding compounds are selected in an "evolutionary" process optimizing their capabilities with respect to a particular property, e.g. its binding activity, its ability to activate, inhibit or modulate the activity, in particular inhibit the intracellular release of calcium mediated by the polypeptides of the present invention.

[0070] Of particular interest are compounds that antagonize the bitter taste receptor activity of the TAS2Rs disclosed and described herein. The specification thereby enables the skilled person to design intelligent compound libraries to screen for antagonists to the bitter response of these receptors, which in turn enables the development of compounds and compositions to suppress or eliminate bitter tasting components of foods, in particular animal foods, nutrients and dietary supplements and pharmaceutical or homeopathic preparations containing such phyto-chemicals. Similarly, the invention also enables the skilled person to screen for additional bitter ligands, or even to screen for compounds that enhance a bitter response, such as might be useful in the food industry. Therefore, another aspect of the invention is a process for isolating an antagonist of the bitter taste receptor activity of the polypeptide encoded by a polynucleotide selected from the group consisting of:

[0071] (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47 and 49;

[0072] (b) polynucleotides having the coding sequence, as shown in SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48 and 50 encoding at least the mature form of the polypeptide;

[0073] (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to said polypeptide, and said fragment or derivative has bitter taste receptor activity;

[0074] (d) polynucleotides which are at least 50% identical to a polynucleotide as defined in any one of (a) to (c) and which code for a polypeptide having bitter taste receptor activity; and

[0075] (e) polynucleotides the complementary strand of which hybridizes, preferably under stringent conditions to a polynucleotide as defined in any one of (a) to (d) and which code for a polypeptide having bitter taste receptor activity;

comprising:

[0076] (1) contacting said polypeptide or a host cell genetically engineered with said polynucleotide or with a vector containing said polynucleotide with a potential antagonist;

[0077] (2) determining whether the potential antagonists antagonizes the bitter taste receptor activity of said polypeptide.

[0078] The polynucleotide employed in this process encodes a polypeptide that still exhibits essentially the same activity as the respective mature bitter taste receptor, i.e. has "bitter taste receptor activity". Preferably the polypeptide has at least 20% (e.g., at least: 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100% or even more) of the activity of the respective full-length TAS2R. One preferred way of measuring TAS2R activity is the ability to release intracellular calcium in a heterologous cell expression system like, for example, (HEK293/15) that stably expresses the alpha-subunit of promiscuous G-proteins, e.g. the mouse G₁₅ subunit or chimeric, in response to bitter tastants, which is dependent on the expression of polypeptides encoded by the polynucleotides of the present invention. The amount of intracellular calcium released can be monitored by, for example, the in vitro FLIPR assay described herein but also by the measurement of one of a variety of other parameters including, for example, IP₃ or cAMP. Additional ways of measuring G-protein coupled receptor activity are known in the art and comprise without limitation electrophysiological methods, transcription assays, which measure, e.g. activation or repression of reporter genes which are coupled to regulatory sequences regulated via the respective G-protein coupled signaling pathway, such reporter proteins comprise, e.g., CAT or LUC; assays measuring internalization of the receptor; or assays in frog melanophore systems, in which pigment movement in melanophores is used as a read out for the activity of adenylate cyclase or phospholipase C (PLC), which in turn are coupled via G-proteins to exogenously expressed receptors (see, for example, McClintock T. S. et al. (1993) *Anal. Biochem.* 209: 298-305; McClintock T. S. and Lerner M. R. (1997) *Brain Res. Brain, Res. Protoc.* 2: 59-68, Potenza M N (1992) *Pigment Cell Res.* 5: 372-328, and Potenza M. N. (1992) *Anal. Biochem.* 206: 315-322)

[0079] As described above with the exception of hTAS2R40, single nucleotide polymorphisms are known for all of the above hTAS2R5, which can be employed in a process for isolating an antagonist of the bitter taste receptor activity. Polynucleotides or polypeptides that differ from the respectively in SEQ ID 1-50 indicated sequences by the nucleotide and amino acid change as indicated in Table I can similarly be employed for the process of the present invention.

[0080] The term "contacting" has the meaning as outlined above. A potential antagonist is a substance which lowers the respective bitter taste receptor activity determined in the absence of the antagonist by at least 10% (e.g., at least: 1%; 15%; 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100%) once contacted with the bitter taste receptor.

[0081] In a preferred embodiment the process further comprises the contacting of the polypeptide with an agonist of the respective bitter taste receptor activity. The contacting of the bitter taste receptor with the agonist can be carried out prior, concomitantly or after contacting the polypeptide with the potential antagonist.

[0082] It has been demonstrated by the inventors that the bitter receptors hTAS2R10, hTAS2R14, hTAS2R16, hTAS2R38, hTAS2R43, hTAS2R44, hTAS2R45, hTAS2R46 and hTAS2R48 respond with specificity to (a) defined class(es) of ligand(s) that include a class of useful phyto-chemicals

in a functional expression assay. Therefore, in an even more preferred embodiment the polypeptides and agonist employed together in above process are selected from the group consisting of:

[0083] (a) the polypeptide encoded by the polynucleotide outlined above as determined by SEQ ID NO: 1 and SEQ ID NO: 2 and the agonist selected from the group consisting of acetyl-thiourea, N,N-dimethylthioformamide, N,N'-diphenylthiourea, N-ethylthiourea, 2-imidazolidinethione, 4(6)-methyl-2-thiouracil, N-methylthiourea, phenylthiocarbamid, 6-phenyl-2-thiouracil, 6-propyl-2-thiouracil, tetramethylthiourea, thioacetamide, thioacetanilide, 2-thiobarbituric acid, and 2-thiouracil and functional derivatives thereof;

[0084] (b) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 9 and SEQ ID NO: 10 and the agonist selected from the group consisting of saccharin and functional derivatives thereof;

[0085] (c) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 11 and SEQ ID NO: 12 and the agonist selected from the group consisting of saccharin and acesulfame K and functional derivatives thereof;

[0086] (d) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 13 and SEQ ID NO: 14 and the agonist selected from the group consisting of absinthine and functional derivatives thereof;

[0087] (e) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 15 and SEQ ID NO: 16 and the agonist selected from the group consisting of absinthine and functional derivatives thereof;

[0088] (f) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 19 and SEQ ID NO: 20 and the agonist selected from the group consisting of absinthine and functional derivatives thereof;

[0089] (g) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 37 and SEQ ID NO: 38 and the agonist selected from the group consisting of strychnine, brucine, denatonium benzoate, and absinthine and functional derivatives thereof;

[0090] (h) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 41 and SEQ ID NO: 42 and the agonist selected from the group consisting of tyrosine, preferably L-tyrosine, and other bitter tasting amino acids including, e.g., leucine, histidine, phenylalanine and tryptophan, and functional derivatives thereof; and

[0091] (i) the polypeptide encoded by the polynucleotide of claim 1 or 2 as determined by SEQ ID NO: 43 and SEQ ID NO: 44 and the agonist selected from the group consisting of naphthyl- β -D-glucoside, phenyl- β -D-glucoside, salicin, helicin, arbutin, 2-nitrophenyl- β -D-glucoside, 4-nitrophenyl- β -D-glucoside, methyl- β -D-glucoside, esculin, 4-nitrophenyl- β -D-thioglucoside, 4-nitrophenyl- β -D-mannoside, and amygdalin and functional derivatives thereof.


[0092] The term "functional derivatives thereof" refers to substances, which are derived from the respectively indicated bitter substance by chemical modification and which elicit at least 20% (e.g., at least: 20%; 30%; 40%; 50%; 60%; 70%; 80%; 90%; 95%; 98%; 99%; 99.5%; or 100% or even more) of the bitter taste receptor activity, if compared to the respective unmodified bitter substance. Chemical modification includes without limitation the introduction of one or more, preferably two, three or four novel side chains or residues or

the exchange of one or more functional groups like, for example, introduction or exchange of H; linear or branched alkyl, in particular lower alkyl (C₁, C₂, C₃, C₄, and C₅, e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or iso-pentyl); substituted linear or branched alkyl, in particular lower substituted alkyl; linear or branched alkenyl, in particular lower alkenyl (C₂, C₃, C₄ and C₅, e.g. ethenyl, 1-propenyl, 2-propenyl, iso-propenyl, 1-butenyl, 2-butenyl, 3-butenyl; substituted linear or branched alkenyl, in particular lower substituted alkenyl; linear or branched alkynyl, in particular lower alkynyl (C₂, C₃, C₄ and C₅); substituted linear or branched alkynyl, in particular lower substituted alkynyl; linear or branched alkanol, in particular lower alkanol (C₁, C₂, C₃, C₄, and C₅); linear or branched alkanal, in particular lower alkanal (C₁, C₂, C₃, C₄, and C₅, e.g. COH, CH₂COH, CH₂CH₂COH; aryl, in particular phenyl; substituted aryl, in particular substituted aryl; heteroaryl; substituted heteroaryl; alkylaryl, in particular benzyl; substituted alkylaryl; in particular substituted benzyl; alkylheteroaryl; substituted alkylheteroaryl; aminoalkyl, C₁, C₂, C₃, C₄ and C₅, e.g. —NHCH₃, —NHCH₂CH₃, —N(CH₃)₂; substituted aminoalkyl; aminoketone, in particular —NHCOCH₃; substituted aminoketone; aminoaryl, in particular —NH-Ph; substituted aminoaryl, in particular substituted —NH-Ph; CN; NH₂; Halogen, in particular F, Cl, and Br; NO₂; OH; SH; NH; CN; or COOH group. If the residues mentioned above are substituted they are preferably mono, di, or tri substituted with a substituent selected from the group of halogen, in particular F, Cl, and Br, NH₂, NO₂, OH, SH, NH, CN, aryl, alkylaryl, heteroaryl, alkylheteroaryl, COH or COOH.

[0093] In particular the hTAS2R16 receptor has been shown to respond specifically to a narrow class of interesting phyto-chemicals selected from the group consisting of bitter beta-glucopyranosides and mannopyranosides.

[0094] The beta-glucopyranosides and beta-mannosepyranosides are a group of bitter compounds consisting of a hydrophobic residue attached to glucose and mannose, respectively, by a beta-glycosidic bond.

[0095] Preferred compounds that bind to the hTAS2R16 taste receptor are chosen from beta glucopyranosides and beta-mannopyranosides defined by the formula:



[0096] These compounds were studied in vitro (see Table I and IV below) and also by human panelists (see Table I below) as is described in greater detail below.

[0097] From these studies certain inferences can be drawn regarding the affinity of the compounds towards activation of the hTAS2R16 receptor. Thus, for the promotion of activation the steric position at C2 can be either alpha or beta and the beta-configuration of the glycosidic bond and the alpha steric position of the hydroxyl group at C4 of the pyranose ring are preferred. Whereas R can be hydrogen, it is preferred that R is a substituent selected from C₁-C₈ alkyl which may be branched, linear or cyclic as appropriate; lower alkenyl resi-

dues, preferably having two, three, four or five carbon atoms; lower alkynyl residues, preferably having two, three, four or five carbon atoms, which can in a preferred embodiment be further substituted with F, Cl, Br, NH_2 , NO_2 , OH, SH, NH, CN, aryl, heteroaryl, COH or COOH group; heteroaryl, e.g. benzofuran and coumarin; aryl, e.g. phenyl, naphthyl; or the same of other sugar residue, e.g. glucopyranoside, which itself can carry a substituent R with the meaning as outlined above. Bulkier groups at C1 may increase the activation of the receptor. The aryl or heteroaryl may be further substituted with one or more substituents. Preferred substituents of the aryl or heteroaryl group are F, Br, Cl, NO_2 , lower alkyl with one, two, three, four, five, six, seven or eight carbon atoms and CH_2OH . The phenyl group is preferably mono, di, or trisubstituted in ortho, para and/or meta position(s). The substituent at C6 is shown as an hydroxyl group above. However, the compounds activity as agonists are little effected by further or alternative substitution at this position, and there is design freedom at this part of the compound. Furthermore, without intending to be bound by theory, it is thought that the substituent "R" is not responsible for bitterness in these compounds. Rather, bitterness is thought to derive from a hydrogen acceptor and donor site provided by two hydroxyl groups on the ring. In another embodiment the β -glycosidic bond of the compounds outlined above can be a S-glycosidic bond, as exemplified by the bitter substance 4-nitrophenyl- β -D-thioglycoside.

[0098] Most preferred compounds are selected from the group consisting of naphthyl- β -D-glucoside, phenyl- β -D-glucoside, salicin, helicin, arbutin, 2-nitrophenyl- β -D-glucoside, 4-nitrophenyl- β -D-glucoside, methyl- β -D-glucoside, esculin, 4-nitrophenyl- β -D-thioglycoside, 4-nitrophenyl- β -D-mannoside, and amygdalin.

[0099] The beta-glucopyranosides are phytonutrients that represent an important class of compounds found in plant-derived foods that may be useful as dietary supplements, or in functional foods or medicaments for the prevention of disease states. However, due to their bitter after-taste they are aversive to consumers and so they are routinely removed from foods during production and processing as is further described in Drewnowski, A. & Gomez-Carneros, C. Bitter taste, phytonutrients, and the consumer: a review. *Am. J. Clin. Nutr.* 72, 1424-1435 (2000). Removal is laborious and therefore expensive. The alternative is to mask the off-flavor using encapsulation technologies or organoleptic compounds as masking agents. However, encapsulation technology may not be appropriate in pharmaceuticals as this may affect the absorption characteristics of the active compound, whereas the use of masking agents may impart their own characteristic flavor which may unbalance the flavor of food or beverages.

[0100] Without wishing to be bound by any particular theory as to their mechanism of action, applicant believes that the bitter receptors activate a G-protein and thereby initiate the aforementioned cellular activation cascade as a result of conformational changes in the receptor after binding by a ligand. Potential antagonists of the bitter response will contain functionality (i.e., will compete for binding at the receptor, and/or act at another binding site through an allosteric mechanism, and/or stabilize the receptor in the inactive conformation, and/or bind reversibly or irreversibly, and/or weaken receptor G protein interaction, and/or interfere with G protein activation).


[0101] Similarly, in another embodiment of the invention, it has been found that the so-called hTAS2R10 receptor is acti-

vated by strychnine, and strychnine analogues such as brucine as well as by denatonium benzoate, absinthine and other alkaloids with (a) ring system(s). Strychnine and its analogues are also useful phytochemicals that find use in medicines and homeopathic treatments.

[0102] In another embodiment of the invention, it has been found that the so-called hTAS2R14 receptor is activated by tyrosine, in particular L-tyrosine, and other bitter tasting amino acids including leucine, histidine, phenylalanine and tryptophan.


[0103] In another embodiment of the invention, it has been found that the so-called hTAS2R38 receptor is activated by acetylthiourea, N,N-dimethylthioformamide, N,N'-diphenylthiourea, N-ethylthiourea, 2-imidazolidinethione, 4(6)-methyl-2-thiouracil, N-methylthiourea, phenylthio-carbamid, 6-phenyl-2-thiouracil, 6-propyl-2-thiouracil, tetramethylthiourea, thioacetamide, thioacetanilide, 2-thiobarbituric acid, and 2-thiouracil.

[0104] From these studies certain inferences can be drawn regarding the affinity of the compounds, which activate the hTAS2R38 receptor. Thus, for the promotion of activation derivatives of 2-thiouracil according to following formula are preferred compounds.



[0105] Whereas R in this formula can be hydrogen, it is preferred that R is a substituent selected from C_1 - C_{10} alkyl, which may be branched, linear or cyclic as appropriate, particularly preferred alkyls are methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or iso-pentyl residues; lower alkenyl residues, preferably having two, three, four or five carbon atoms; lower alkynyl residues, preferably having two, three, four or five carbon atoms, which can in a preferred embodiment be further substituted with F, Cl, Br, NH_2 , NO_2 , OH, SH, NH, CN, aryl, heteroaryl, COH or COOH group; heteroaryl, e.g. benzofuran and coumarin; aryl, e.g. phenyl, naphthyl; F, Cl, Br, NH_2 , NO_2 , OH, SH, NH, CN, aryl, alkylaryl, heteroaryl, alkylheteroaryl, COH or COOH group. In a further embodiment the carbon atom at the 4 position can substituted with $-\text{O}-\text{R}_1$, in which R_1 can have the same meaning as outlined above for R.

[0106] Another general structure of compounds having affinity for hTAS2R38 and which are thus suitable for activation of hTAS2R38 is depicted by the following formula:



[0107] In this formula R_2 , R_3 , and R_4 can each independently of each other have the meaning H; alkyl, in particular lower alkyl (C_1 - C_5 , e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or iso-pentyl); substituted alkyl; alkenyl, in particular lower alkenyl (C_2 - C_5); sub-

stituted alkenyl; alkynyl, in particular lower alkynyl (C_2 - C_5); substituted alkynyl; alkanal, in particular lower alkanal (e.g. $-COCH_3$, $-COCH_2CH_3$, $-COCH_2CH_2CH_3$); aryl, in particular phenyl; substituted aryl; heteroaryl; substituted heteroaryl; alkylaryl, in particular benzyl; substituted alkylaryl; alkylheteroaryl; substituted alkylheteroaryl aminoalkyl, in particular $-NHCH_3$, $-NHCH_2CH_3$, $-N(CH_3)_2$; substituted aminoalkyl; aminoketone, in particular $-NHCOCH_3$; substituted aminoketone; aminoaryl, in particular $-NH-Ph$; substituted aminoaryl; CN; NH_2 ; Halogen, in particular F, Cl, and Br; NO_2 . In a preferred embodiment R_2 or R_3 and R_4 can form a ring, preferably a four, five, six, seven or eight membered hetero cycle, which in a preferred embodiment is an aromatic hetero cycle. The residue of R_2 or R_3 , which is not involved in the formation of the ring structure can have any of the meanings as outlined above. In a further preferred embodiment at least one of R_2 or R_3 has the meaning alkanal, preferably lower alkanal as outlined above. In case that only one of R_2 or R_3 has the meaning alkanal, than the other substituent preferably has the meaning H.

[0108] In a preferred embodiment R_2 is selected from the group consisting of H, CH_3 and Ph, R_3 is selected from the group of H, CH_3 and Ph and R_4 is selected from the group consisting of H, CH_3 , $NH-Ph$, $-NHCH_2CH_3$, $-NHCH_2CH_3$, $-NHCH_3$, and $-N(CH_3)_2$.

[0109] In another embodiment of the invention, it has been found that the so-called hTAS2R43 receptor is activated by saccharin, derivatives thereof and other sulfonamides.

[0110] In another embodiment of the invention, it has been found that the so-called hTAS2R44 receptor is activated by saccharin and acesulfame K, derivatives thereof and other sulfonamides.

[0111] In another embodiment of the invention, it has been found that the so-called hTAS2R45, hTAS2R46 and hTAS2R48 receptor is activated by absinthine derivatives thereof and other sulfonamides.

[0112] The skilled person will appreciate that having regard to the structure-function information provided by the present invention, it is possible to compile libraries of molecules to find inhibitors of the bitter response of the disclosed hTAS2R in particular of the hTAS2R10, 14, 16, 38, 43, 44, 45, 46, and 48, which are triggered by the above outlined specific bitter substance(s). Such inhibitors, and libraries comprising same, form other aspects of the present invention. A still farther aspect of the invention relates to the use of such inhibitors in food or pharmaceutical compositions containing bitter tastants such as referred to herein above, for the elimination or suppression of bitter taste perception.

[0113] In practicing the various aspects and embodiments of the present invention in relation to cloning receptors, elucidating ligand-receptor pairs, and finding modulators of the bitter response of receptors, recourse is made to conventional techniques in molecular biology, microbiology and recombinant technology. Accordingly, the skilled person is fully apprised of such techniques and as such they are hereafter treated only summarily in order to more fully describe the context of the present invention.

[0114] In order to express cDNAs encoding the receptors, one typically subclones receptor cDNA into an expression vector that contains a strong promoter to direct transcription, a transcription/translation terminator, and a ribosome-binding site for translational initiation. Suitable bacterial promoters are well known in the art, e.g., *E. Coli*, *Bacillus* sp., and *Salmonella*, and kits for such expression systems are com-

mercially available. Similarly eukaryotic expression systems for mammalian cells, yeast, and insect cells are well known in the art and are also commercially available. The eukaryotic expression vector may be, for example an adenoviral vector, an adeno-associated vector, or a retroviral vector.

[0115] In addition to the promoter, the expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the receptor-encoding nucleic acid in host cells. A typical expression cassette thus contains a promoter operatively linked to the nucleic acid sequence encoding the receptor and signals required for efficient polyadenylation of the transcript, ribosome binding sites, and translation termination. The nucleic acid sequence encoding the receptor may typically be linked to a membrane-targeting signal such as the N-terminal 45 amino acids of the rat Somatostatin-3 receptor sequence to promote efficient cell-surface expression of the recombinant receptor. Additional elements of the cassette may include, for example enhancers.

[0116] An expression cassette should also contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes.

[0117] The particular expression vector used to transport the genetic information into the cell is not particularly critical. Any of the conventional vectors used for expression in eukaryotic or prokaryotic cells may be used. Standard bacterial expression vectors include plasmids such as pBR322 based plasmids, pSKF, pET23D, and fusion expression systems such as GST and LacZ, but there are many more known in the art to the skilled person that can be usefully employed.

[0118] Expression vectors containing regulatory elements from eukaryotic viruses are typically used in eukaryotic expression vectors, e.g., SV40 vectors, papilloma virus vectors, and vectors derived from Epstein-Barr virus. Other exemplary eukaryotic vectors include pMSG, pAV009/A.sup.+, pMTO10/A.sup.+, pMAMneo-5, baculovirus pDSVE, pcDNA3.1, pIRES and any other vector allowing expression of proteins under the direction of the SV40 early promoter, SV40 late promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

[0119] Some expression systems have markers that provide gene amplification such as thymidine kinase, hygromycin B phosphotransferase, and dihydrofolate reductase. Alternatively, high yield expression systems not involving gene amplification are also suitable.

[0120] The elements that are typically included in expression vectors also include a replicon that functions in *E. Coli*, a gene encoding drug resistance to permit selection of bacteria that harbor recombinant plasmids, and unique restriction sites in nonessential regions of the plasmid to allow insertion of eukaryotic sequences. The particular drug resistance gene chosen is not critical, any of the many drug resistance genes known in the art are suitable. The prokaryotic sequences are optionally chosen such that they do not interfere with the replication of the DNA in eukaryotic cells, if necessary.

[0121] Standard transfection methods can be used to produce bacterial, mammalian, yeast or insect cell lines that express large quantities of the receptor, which are then purified using standard techniques.

[0122] Any of the well known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell. It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one gene into the host cell capable of expressing the receptor.

[0123] After the expression vector is introduced into the cells, the transfected cells may be cultured under conditions favoring expression of the receptor, which is recovered from the culture using standard techniques. For example the cells may be burst open either mechanically or by osmotic shock before being subject to precipitation and chromatography steps, the nature and sequence of which will depend on the particular recombinant material to be recovered. Alternatively, the recombinant protein may be recovered from the culture medium in which the recombinant cells had been cultured.

[0124] The activity of any of the receptors described herein can be assessed using a variety of in vitro and in vivo assays to determine functional, chemical, and physical effects, e.g., measuring ligand binding, secondary messengers (e.g., cAMP, cGMP, IP₃, DAG, or Ca²⁺) ion flux, phosphorylation levels, transcription levels, neurotransmitter levels, and the like. Furthermore, such assays can be used to test for inhibitors of the receptors as is well known in the art.

[0125] Samples or assays that are treated with a potential receptor inhibitor may be compared to control samples without the test compound, to examine the extent of modulation. Control samples (untreated with inhibitors) are assigned a relative receptor activity value of 100. Inhibition of receptor activity is achieved when the receptor activity value relative to the control is lower, and conversely receptor activity is enhanced when activity relative to the control is higher.

[0126] The effects of the test compounds upon the function of the receptors can be measured by examining any of the parameters described above. Any suitable physiological change that affects receptor activity can be used to assess the influence of a test compound on the receptors of this invention. When the functional consequences are determined using intact cells or animals, one can measure a variety of effects such as changes in intracellular secondary messengers such as Ca²⁺, IP₃ or cAMP.

[0127] Preferred assays for G-protein coupled receptors include cells that are loaded with ion sensitive dyes to report receptor activity. In assays for identifying modulatory compounds, changes in the level of ions in the cytoplasm or membrane voltage will be monitored using an ion sensitive or membrane voltage fluorescent indicator, respectively. For G-protein coupled receptors, promiscuous G-proteins such as G.alpha. 15 and G.alpha. 16 and chimeric G-proteins can be used in the assay of choice (see, for example, Wilkie et al., *Proc. Nat. Acad. Sci. USA* 88, 10049-10053 (1991)). Such promiscuous G-proteins allow coupling of a wide range of receptors.

[0128] Receptor activation typically initiates subsequent intracellular events, e.g., increases in second messengers such as IP₃, which releases intracellular stores of calcium ions. Activation of some G-protein coupled receptors stimulates the formation of inositol triphosphate (1133) through phos-

pholipase C-mediated hydrolysis of phosphatidylinositol (Berridge & Irvine, *Nature* 312, 315-21 (1984)). IP₃ in turn stimulates the release of intracellular calcium ion stores. Thus, a change in cytoplasmic calcium ion levels, or a change in second messenger levels such as IP₃ can be used to assess G-protein coupled receptor function. Cells expressing such G-protein coupled receptors may exhibit increased cytoplasmic calcium levels as a result of contribution from both intracellular stores and via activation of ion channels, in which case it may be desirable, although not necessary, to conduct such assays in calcium-free buffer, optionally supplemented with a chelating agent such as EGTA, to distinguish fluorescence response resulting from calcium release from internal stores.

[0129] In a preferred embodiment, receptor activity is measured by expressing the receptor in a heterologous cell with a promiscuous G-protein, such as G.alpha. 15, 16, or a chimeric G-protein that links the receptor to a phospholipase C signal transduction pathway. Optionally the cell line is HEK-293, although other mammalian cells are also preferred such as CHO and COS cells. Modulation of taste transduction is assayed by measuring changes in intracellular Ca²⁺ levels, which change in response to modulation of the receptor signal transduction pathway via administration of a molecule that associates with the receptor. Changes in Ca²⁺ levels are optionally measured using fluorescent Ca²⁺ indicator dyes and fluorometric imaging.

[0130] The type of assay described above with respect to G-protein coupled bitter taste receptors can, however, also be employed for the identification of binding compounds, in particular agonists or antagonists of any G-protein coupled signalling molecule, in particular G-protein coupled receptor. Therefore, another aspect of the present invention relates to a process for the identification of agonists or antagonists of G-protein coupled signalling molecules comprising the steps of:

[0131] (1) contacting a cell comprising a promiscuous G-protein like, for example, G.alpha.15, 16, or a chimeric G-protein, and a G-protein coupled signalling molecule, in particular receptor, with a the potential agonist or antagonists of the signalling molecule;

[0132] (2) determining whether the potential agonist or antagonists agonizes or antagonizes the activity of the signalling molecule.

[0133] The activity of the signalling molecule and the increase or decrease of that activity in response to the potential agonist or antagonist can be determined as outlined above with respect to the identification of bitter receptor taste activity. The respectively indicated percent increases or decreases of the activity, which are required to qualify as antagonist or agonist do apply mutatis mutandis. Additionally the term "contacting" has the meaning as outlined above. Preferably the signalling molecule and/or the promiscuous G-protein has been introduced into the cell. The type of cell, which are preferred are those indicated above.

[0134] In yet another embodiment, the ligand-binding domains of the receptors can be employed in vitro in soluble or solid-state reactions to assay for ligand binding. Ligand binding in a receptor, or a domain of a receptor, can be tested in solution, in a bilayer membrane attached to a solid phase in a lipid monolayer or vesicles. Thereby, the binding of a modulator to the receptor, or domain, can be observed using changes in spectroscopic characteristics, e.g. fluorescence,

absorbance or refractive index; or hydrodynamic (e.g. shape), chromatographic, or solubility properties, as is generally known in the art.

[0135] The compounds tested as modulators of the receptors can be any small chemical compound, or a biological entity, such as a protein, sugar, nucleic acid or lipid. Typically, test compounds will be small chemical molecules. Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although knowledge of the ligand specificity of an individual receptor would enable the skilled person to make an intelligent selection of interesting compounds. The assays may be designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays). The skilled person will understand that there are many suppliers of libraries of chemical compounds.

[0136] Assays may be run in high throughput screening methods that involve providing a combinatorial chemical or peptide library containing a large number of potential therapeutic, or tastant compounds (that are potential ligand compounds). Such libraries are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as lead compounds to further develop modulators for final products, or can themselves be used as actual modulators.

[0137] A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

[0138] Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art and no more needs to be stated here.

[0139] In the high throughput assays of the invention, it is possible to screen up to several thousand different modulators or ligands in a single day. In particular, each well of a microtiter plate can be used to run a separate assay against a selected potential modulator, or, if concentration or incubation time effects are to be observed, every 5-10 wells can test a single modulator. Thus, a single standard microtiter plate can assay about 100 (e.g., 96) modulators. If 1536 well plates are used, then a single plate can easily assay from about 100 to about 1500 different compounds. It is possible to assay several different plates per day; assay screens for up to about 6,000-20,000 different compounds is possible using the integrated systems of the invention.

[0140] Lead compounds found by assay technology herein above described, or development compounds formed from such leads can be administered directly to a human subject to modulate bitter taste. Alternatively, such compounds can be formulated with other ingredients of preparations to be taken orally, for example, foods, including animal food, and beverages, pharmaceutical or nutraceutical or homeopathic preparations.

[0141] Therefore, another aspect of the invention is a process for the production of foodstuffs or any precursor material or additive employed in the production of foodstuffs comprising the steps of the above described processes for the identification of a compound binding to hTAS2R or an antagonist of hTAS2R and the subsequent step of admixing the identified compound or antagonist with foodstuffs or any precursor material or additive employed in the production of foodstuffs.

[0142] Bitter taste is a particular problem when orally administering pharmaceuticals, which often have an unpleasant bitter taste. In particular in elderly persons, children and chronically ill patients this taste can lead to a lack of compliance with a treatment regimen. In addition in veterinary applications the oral administration of bitter tasting pharmaceuticals can be problematic. Therefore, a further aspect of the invention is a process for the production of a nutraceutical or pharmaceutical composition comprising the steps of the processes of a compound binding to hTAS2R or an antagonist of hTAS2R and the subsequent step of formulating the compound or antagonist with an active agent in a pharmaceutically acceptable form.

[0143] Consequently, a further aspect of the invention is a foodstuff, in particular animal food, or any precursor material or additive employed in the production of foodstuffs comprising an antagonist/inhibitor described above, preferably an antibody directed against one of the hTAS2Rs described herein, the extracellular domain of one of the hTAS2Rs described herein or an inhibiting RNA.

[0144] Also comprised is a nutraceutical or pharmaceutical composition comprising an antagonist/inhibitor as described above, preferably an antibody directed against one of the hTAS2Rs described herein, the extracellular domain of one of the hTAS2Rs described herein or an inhibiting RNA and an active agent, which preferably inhibits a bitter taste, and optionally a pharmaceutically acceptable carrier.

[0145] The amount of compound to be taken orally must be sufficient to effect a beneficial response in the human subject, and will be determined by the efficacy of the particular taste modulators and the existence, nature, and extent of any adverse side-effects that accompany the administration of a particular compound. There now follows a series of examples that serve to illustrate the invention, not to limit.

[0146] A further aspect of the present invention is the use of a polynucleotide as described above, a vector as described above, an antibody as described above or an antagonist/inhibitor of as described above, preferably an antibody directed against one of the hTAS2Rs described herein, the extracellular domain of one of the hTAS2Rs described herein or an inhibiting RNA for the manufacture of a medicament for the treatment of an abnormally increased or decreased sensitivity towards a bitter substance.

[0147] Techniques associated with detection or regulation of genes are well known to skilled artisans. Such techniques can be used, for example, for basic research on bitter receptors and to diagnose and/or treat disorders associated with aberrant bitter receptor expression.

[0148] The following examples are merely illustrative of the present invention and should not be construed to limit the scope of the invention as indicated by the appended claims in any way. The contents of the U.S. provisional application Ser. No. 60/413,298 the priority of which is claimed is hereby incorporated by reference in its entirety.

EXAMPLE 1

Cloning of the hTAS2R Genes

[0149] Human genomic DNA was isolated from HEK293 cells using the E.Z.N.A. Blood DNA Kit II (Qiagen) and the

various hTAS2Rs were amplified by PCR using gene-specific primers that span the complete coding region of the individual hTAS2R genes. Reaction parameters were: 4 cycles; 1 min, 94° C.; 1 min, 64° C.; 1.5 min 68° C. using Advantage 2 polymerase (Clontech). 5% of the reaction served then as template for further amplification with Pfu DNA polymerase (Promega): 30 cycles; 1 min, 94 C; 1 min, 64° C.; 3 min, 72° C. The hTAS2R amplicons were then sub-cloned into a cassette based on pcDNA5-FRT (Invitrogen). The cloning cassette contains the first 45 amino acids of the rat somatostatin type 3 receptor (as is further described by Meyerhof et al., *Proc. Nat. Acad. Sci. USA*, 89, 10267-10271 (1992)) as a cell surface-targeting signal at the N-terminus. The C-terminus contained the herpes simplex virus (HSV) glycoprotein D epitope which does not interfere with signaling of heptahelical receptors and can be used for immunocytochemistry using an antibody that binds specifically to the HSV glycoprotein D epitope (see Roosterman et al, *J. Neuroendocrinol.*, 9, 741-751 (1997)). Comparison of the DNA sequences of at least four clones identified mutations generated during PCR and this avoided picking mutated clones. We compared the amino acid sequences using the AlignX program of the Vector NTT™ Suite (InforMax).

[0150] Using the above-described method, DNA sequences encoding all 24 bitter receptors identified by applicant were cloned. As indicated above, they were derived by a PCR-based method using genomic DNA as the template. Since all of the 24 genomic sequences lack introns, the DNA clones obtained had the same sequences as corresponding cDNA clones derived by reverse transcription-PCR (RT-PCR) of mRNA from cells expressing the relevant polypeptides would have.

EXAMPLE 2

Immunocytochemistry

[0151] Batches of HE 93 cells were separately transiently transfected with expression vectors (pCDN5/FRT; Invitrogen) containing each of the 24 above described coding sequences using lipofectamine 2000 (Invitrogen) and aliquots of the resulting cell populations were separately seeded on polylysine-coated coverslips. At 24 h post transfection they were washed with phosphate buffered saline (PBS), cooled on ice and added 20 microgram/ml biotin-labeled concanavalin A (Sigma) for 1 h, which binds to cell surface glycoproteins. Thereafter, the cells were fixed for 5 min in methanol/acetone (1:1) and then permeabilized for 4 min with 0.25% Triton X-100. In order to reduce nonspecific binding the coverslips were incubated in 2% goat serum. Thereafter, anti-HSV glycoprotein D antiserum (Novagen, 1:10,000) was added to detect the chimeric receptors that, as described above, would have a HSV glycoprotein epitope fused to their C-termini, and Texas Red-Avidin D (Vector, 1:200) has added to stain the cell surface and incubation continued overnight at 4° C. Such C-termini are intracellular and for this reason it is necessary to permeabilize the cells to permit entry of the HSV glycoprotein D epitope-specific antibody molecules into them. After washing (5x in PBS, RT) Alexa488-conjugated goat anti-mouse antiserum (Molecular Probes, 1:1000) was added and incubation continued at room temperature for 1 h. Finally, the cells were embedded in Fluorescent Mounting Medium (Dako) and analyzed using a Leica TCS SP2 Laser Scan Inverted microscope. The preparations were scanned sequentially with an argon/krypton

laser (488 nm) to excite the Alexa488 dye and with a green-helium-neon laser (543 nm) to excite the Texas Red dye. The spectral detector recorded light emission at 510-560 nm and 580-660 nm, respectively. Images of 1024×1024 pixels were processed with Corel PHOTO-PAINT 10.0 (Corel Corporation) and printed on a Tektronix color laser printer. The immunocytochemical data permitted calculation of the proportion of cells expressing recombinant receptors (green fluorescent cells divided by total cell number in a microscopic field) and the proportion of cells that display expression of TAS2Rs at the plasma membrane level (number of cells with colocalization of green and red fluorescence divided by the number of green fluorescent cells). Of the 24 transfectant lines tested, all were found to express the encoded polypeptides. The proportion of receptor-expressing cells in the various transfectant lines ranged from about 10% to about 35%.

EXAMPLE 3

Heterologous Expression of hTAS2R Receptors

[0152] A fluorescence imaging plate reader (FLIPR, Molecular Devices) was used to functionally screen cell populations transiently transfected with expression vectors encoding the above-described 24 bitter receptors and to establish concentration-response curves for hTAS2R16 and hTASR10. The single-cell calcium imaging technique was also employed to demonstrate receptor selectivity and cross-desensitization. For the FLIPR experiments the HEK293/15 cells were grown to 50% confluence. The cells were then seeded at a density of 3×10^3 cells per well into 96-well black-wall, clear-bottom microtiter plates (Greiner). After 48 h the cells in each well were transfected using Lipofectamine 2000 and 24-30 h later were loaded with Fluo4AM (Molecular Probes). Thereafter they were stimulated with bitter compounds (SigmaAldrich, further purified by reversed-phase HPLC to >99% purity). Calcium signals were recorded simultaneously from each well at 1 Hz at 510 nm after excitation at 488 nm and the recordings were corrected for cell density. The responses of five wells containing cells expressing the same receptor and that received the same stimulus (i.e., the same compound at the same concentration) were averaged. Calcium traces were subtracted that were determined in triplicate of mock-transfected cells stimulated with the same concentration of tastant. The calculations rest on at least four independent transfection experiments. Plots of the amplitudes versus concentrations fitted by nonlinear regression to the function $f(x)=100/(1+(EC_{50}/x)^n)$, with x =agonist concentration and nH =Hill coefficient permitted calculation of EC_{50} values and threshold values of activation.

[0153] EC_{50} and threshold values obtained with hTAS2R16-expressing transfectants are shown in Table 1 below and the results are described in Example 4.

[0154] In separate experiments, hTAS2R10-expressing transfectants were found to have a threshold of activation of approximately 0.1 μ M and a EC_{50} of 5-20 μ M using strychnine as the test compound. Similar results were obtained with brucine.

[0155] Single-cell Ca^{2+} imaging was performed with the hTAS2R16-transfected HEK293/15 cells as described in *Cell* 95, 917-926 (1998), but with the following modifications: The Till Photonics imaging system (Munich, Germany) was used in which a monochromator is connected by a quartz fiber lightguide and an epifluorescence condenser to an inverted Olympus IX50 microscope equipped with a UApo/340 40×1.

35 oil-immersion lens. 30 h post-transfection, FURA-2AM-loaded cells were sequentially illuminated in 5 s intervals for 3-10 ms, first at 340 nm, then at 380 nm, online ratioed light emissions at 510 nm (340/380) and monitored the images via an intensified, cooled CCD camera. The 5 s interval camera pictures of all cells in the microscope field of vision permanently were stored and analyzed offline. 10-15% of all cells in the camera field responded to agonists in transient transfection experiments. The proportion of responders was about half of that found by immunocytochemistry, probably reflecting a sub-optimal signal transduction. Responses were not observed in mocktransfected cells. Isoproterenol (10 micro-Molar) was used at the end of all experiments to stimulate endogenous betaadrenergic receptors, proving a functional $G_{\alpha 13}$ dependent signal transduction cascade.

[0156] For RT-PCR and in-situ hybridization work, human RNA (Clontech) was purchased or it was isolated from surgical tongue specimens with peqGOLD RNAPure (Peqlab) and the preparations digested with DNase I (Invitrogen). Following cDNA synthesis (Smart cDNA synthesis Kit, Clontech) hTAS2R16 cDNA was PCR-amplified (39 cycles, 1 min 94° C., 1 min 64° C., 1 min 72° C.) using specific forward and reverse primers with overhangs containing EcoRI or NotI sites SEQ ID Nos 51 and 52 and the amplicons analyzed on agarose gels. Subcloning and sequencing demonstrated the identity of the amplified bands. Approximately 15 micrometer cryo-sections of human tongue specimens containing vallate papilla at 65° C. were processed and hybridized with a hTAS2R16 riboprobe spanning the complete coding region and generated from hTAS2R16 cDNA. The in-situ hybridization method used was essentially the same as that described in *Nature*, 413, 631-635 (2001) except that the riboprobe was conjugated with biotin and an alkaline phosphatase-avidin conjugate was used for detection. This experiment indicated that TAS2R16 mRNA is expressed in vallate papilla which are known to perceive bitter taste.

EXAMPLE 4

Human Taste Experiments

[0157] 15 experienced panelists in a sensory panel room at 22-25° C. determined bitter thresholds on three different sessions using a triangle test with tap water as solvent, according to methodology set out in *J. Agric. Food Chem.*, 49, 231-238 (2001), or Mailgaard M et al, "Sensory Evaluation Techniques" (CRC Press LLC, New York 1999). For dose-response relations, bitter tastant concentration series were presented to 10 trained panelists in random order. The panelists ranked the samples in increasing order of intensity and, for each concentration, evaluated bitterness intensity on a scale from 0 to 5 (ref. 24). The dose-response curves of three different sessions were averaged. The intensity values between individuals and separate sessions differed by not more than 0.5 units.

[0158] To investigate adaptation, the 8 panelists first maintained aqueous solutions (5 ml) of phenyl-β-D-glucopyranoside (8 mM), phenyl-α-D-glucopyranoside (180 mM), salicin (8 mM), or helicin (8 mM) for 15 s in their oral cavities and evaluated the bitter intensity as described above. After 30 min, they kept a denatonium benzoate solution (5 ml, 0.0003 mM) for 15 s in their mouth and evaluated its bitterness. The panelists spat off the denatonium benzoate solution, took up the phenyl-β-D-glucopyranoside or the phenyl-α-D-glucopyranoside solutions orally for 120 s or 180 s and judged

their bitterness intensity after 15, 30, 60, 120 and 180s. Thereafter, the panelists spat off these solutions and then sequentially took up salicin, helicin (5 ml, 8 mM) and denatonium benzoate (5 ml, 0.0003 mM) and evaluated bitterness intensities of these solutions after 15 s. After an additional 30 min, the first experiment was repeated. The data of three different sessions for each panelist were averaged. Intensity values between individuals and separate sessions differed by not more than ±0.5 units.

[0159] Results of in vitro assays (FLIPR) and human taste experiments are shown in Table 1 below.

TABLE II

Compound	Threshold Value (mM)		EC ₅₀ (mM)	
	FLIPR	Human	FLIPR	Human
1	0.07 +/- 0.02	0.1 +/- 0.05	1.1 +/- 0.1	0.7 +/- 0.2
2	0.07 +/- 0.02	0.2 +/- 0.1	1.4 +/- 0.2	1.1 +/- 0.3
3	0.3 +/- 0.1	0.4 +/- 0.1	2.3 +/- 0.4	2.2 +/- 0.7
4	0.5 +/- 0.2	0.9 +/- 0.3	5.8 +/- 0.9	5.4 +/- 1.8
5	1.5 +/- 0.5	n.d.	n.d.	n.d.
6	0.4 +/- 0.1	0.2 +/- 0.1	1.0 +/- 0.1	1.4 +/- 0.4
7	15 +/- 6	32 +/- 11	n.d.	320 +/- 108
8	2.3 +/- 0.9	n.d.	20 +/- 3.4	n.d.
9	4 +/- 2	4 +/- 1	n.d.	n.d.
10	n.r.	40 +/- 13	n.r.	n.d.
11	n.r.	9 +/- 3	n.r.	50 +/- 17

1 = phenyl-beta-D-glucopyranoside; 2 = salicin; 3 = helicin; 4 = arbutin; 5 = 2-nitro-phenyl-beta-D-glucopyranoside; 6 = naphthyl-beta-D-pyranoside; 7 = methyl-beta-D-lucopyranoside; 8 = amygdalin; 9 = esculin; 10 = phenyl-beta-D-galactopyranoside; 11 = phenyl-alpha-D-glucopyranoside.
n.d. = not data due to solubility problems or toxicity or artifacts in vitro.
n.r. = No response up to 100 mM.

[0160] The FLIPR results provide the threshold concentration of the compounds (nM) at which point the receptor detects the compounds. The EC₅₀ results express the concentration of the compound wherein the receptor signal is at 50%, and is a representation of the affinity of a receptor for a compound.

[0161] The results show that the in vitro FLIPR measurements for salicin closely resemble the human taste study results. This bitter-tasting compound has known anti-pyretic and analgesic action, and the results suggest that in vitro assays using hTAS2R16 may represent a useful tool to find compounds that suppress or eliminate the bitter response to this compound. Also, for all the other tested beta-glucopyranosides, the close correspondence of Threshold Concentration and EC₅₀ results suggest that hTAS2R16 is a cognate human receptor for these class of bitter compounds. In contrast, the related structures (see compounds 10 and 11) show 90- to 400-fold higher Threshold Concentrations, which indicates that this receptor is rather selective, and that these bitter compounds activate different receptors.

[0162] Adaptation frequently occurs in sensory systems and means that stimuli elicit reduced responses upon prolonged or repeated stimulus presentations. Repeated stimulation of hTAS2R16-expressing cells with phenyl-beta-D-glucopyranoside resulted in largely diminished responses to salicin as well. This cross-desensitization occurred among the other tested beta-pyranosides and was fully reversible. It resembles homologous desensitization of agonist-occupied heptahelical receptors mediated by GRKs, i.e. specific kinases, and arresting. We also observed adaptation in the human test panel that initially scored phenyl-beta-D-glucopyranoside, salicin and helicin as equally intensely bitter.

The bitterness of phenyl-beta-D glucopyranoside declined during prolonged stimulation and the test panel perceived salicin and helicin also as less bitter, but not the unrelated bitter substance denatonium benzoate, which cannot activate TAS2R16. Adaptation was fully reversible. On the opposite, the phenyl-alpha-D-glucopyranoside failed to cross-adapt with all tested beta-D-glucopyranosides, although its own bitter response desensitized strongly. This indicates that beta-glucopyranosides signal through a common mechanism most likely involving hTAS2R16 as a bitter taste receptor while the alpha-isomer activates a separate receptor. A recent human psychophysical study also revealed cross-adaptation amongst two bitter amino acids but not between the two bitter amino acids and urea, suggesting the existence of distinct receptors for the bitter amino acids and urea. Although most, if not all, bitter receptors are present in the same subset of taste receptor cells, adaptation to specific bitter stimuli can be explained if bitter receptors were subject to homologous desensitization.

EXAMPLE 5

Heterologous Expression of hTAS2R

[0163] Transient transfection of TAS2Rs into HEK-293T- α 16gustducin44 cells. We cloned the DNAs of all human putative bitter responsive receptors into pcDNA5/FRT (Invitrogen) by PCR-methods and transiently transfected the plasmids with lipofectamine 2000 (Invitrogen) into HEK-293T- α 16gustducin44 cells grown to 50% confluence. These cells stably express a chimeric G protein constructed from human α 16 and rat gustducin. Finally, we seeded the transfected cells at a density of 3×10^3 cells per well into 96-well black-wall, clear-bottom microtiter plates (Greiner).

[0164] Co-transfection of TAS2Rs with gustducin and phospholipase-C β 2 into HEK-293 cells. Alternatively, we transfected simultaneously plasmid DNAs encoding one of the TAS2Rs, phospholipase-C β 2 and α -gustducin into HEK-293 cells using the lipofectamine method. Additional cotrans-

fection of G-protein β and γ -subunits may improve the bitter tastant-induced responses. Thereafter, the transfected cells were seeded at a density of 3×10^3 cells per well into 96-well black-wall, clear-bottom microtiter plates (Greiner).

[0165] Fluorometric Imaging Plate Reader (FLIPR) assay 24-30 h later, the cells were loaded with 4 μ M FLUO-4/AM (Molecular Probes) and 0.04% Pluronic F-127 (Molecular Probes) in HEPES-buffered saline (HBS), 140 mM NaCl, 5 mM KCl, 2.5 mM CaCl_2 , 10 mM HEPES, 10 mM glucose and 2.5 mM probenecide, pH 7.4, for 1 hour at 37° C. Thereafter, cells were gently washed in HBS by an automated plate washer (Denley Cellwash, Labsystem) and transferred to the FLIPR (Molecular Devices). The FLIPR integrates an argon laser excitation source, a 96-well pipettor, and a detection system utilizing a Charged Coupled Device imaging camera. Fluorescence emissions from the 96 wells were monitored at an emission wavelength of 510 nm, after excitation with 488 nm (F488). Fluorescence data were collected 1 min before and 10 min after stimulation. Data were collected every 6 s before and every 1 s after agonist stimulation. 50 μ l of 3 \times concentrated agonists were delivered within 2 by the integrated 96-well pipettor to the wells containing 100 μ l HBS. Agonist responses were quantified using the amplitudes of the fluorescence peaks. We averaged the responses of five wells containing cells expressing the same receptor and that received the same stimulus. Calcium traces were determined in triplicate of mock-transfected cells stimulated with the same concentration of tastant. EC_{50} values and plots of the amplitudes versus concentrations were derived from fitting the data by nonlinear regression to the function $f(x) = 100/[1 + (\text{EC}_{50}/x)^{nH}]$, where x is the agonist concentration and nH is the Hill coefficient. The results for hTAS2R10 (Table II), hTAS2R14 (Table III), hTAS2R16 (Table IV), hTAS2R38 (Table V), hTAS2R43 (Table VI), hTAS2R44 (Table VII), hTAS2R45 (Table VIII), hTAS2R46 (Table IX) and hTAS2R (Table X) are shown below.

TABLE III

Identified agonists of hTAS2R10



Substance	Structure	Approx. thresh- old [mM]	EC_{50} [mM]
Strychnine*		0.003	0.04
Bucine		0.01	0.06

TABLE III-continued

<u>Identified agonists of hTAS2R10</u>			
Substance	Structure	Approx. thresh- old [mM]	EC ₅₀ [mM]
Denatonium benzoate		0.003	0.07
Absinthine		0.01	

TABLE IV

<u>Identified agonists of hTAS2R14</u>		
Substance	Structure	Reacts at
L-Tyrosine		1 mM

TABLE V

<u>Identified agonists of hTAS2R16</u>			
Substance	Structure	Threshold [mM]	EC ₅₀ [mM]
Naphthyl-β-D-Glucoside		0.4 ± 0.1	1.0 ± 0.1
Phenyl-β-D-Glucoside		0.07 ± 0.02	1.1 ± 0.1

TABLE V-continued








Identified agonists of hTAS2R16			
Substance	Structure	Threshold [mM]	EC ₅₀ [mM]
Salicin		0.07 ± 0.02	1.4 ± 0.2
Helicin		0.3 ± 0.1	2.3 ± 0.4
Arbutin		0.5 ± 0.2	5.8 ± 0.9
2-Nitrophenyl-β-D-Glucoside		0.3 – 1	Not determined
4-Nitrophenyl-β-D-Glucoside		1 – 3	Not determined
Methyl-β-D-Glucoside*		15 ± 6	32 ± 11
Esculin		4 ± 2	Not determined

TABLE V-continued




Identified agonists of hTAS2R16			
Substance	Structure	Threshold [mM]	EC ₅₀ [mM]
4-Nitrophenyl-β-D-Thioglucoside		1 – 5	Not determined
4-Nitrophenyl-β-D-Mannoside		1 – 3	Not determined
Amygdalin		2.3 ± 0.9	20 ± 3.4

TABLE VI






Identified agonists of hTAS2R38			
Substance	Structure	Approx. Threshold [μM]	EC ₅₀ [μM]
Acetylthiourea		2	15
N,N-Dimethyl-thioformamide		10	55
N,N'-Diphenylthiourea		0.3	2.3
N-Ethylthiourea		30	260
2-Imidazolidinethione (=N,N'-Ethylenethiourea)		10	not determined

TABLE VI-continued










Identified agonists of hTAS2R38			
Substance	Structure	Approx. Threshold [μ M]	EC ₅₀ [μ M]
4(6)-Methyl-2-thiouracil		20	180
N-Methylthiourea		100	estimated 600-800
Phenylthiocarbamid (PTC)		0.3	2
6-Phenyl-2-thiouracil		0.15	0.5
6-Propyl-2-thiouracil (PROP)		0.3	2
Tetramethylthiourea		10-30	100
Thioacetamide		100	not determined
Thioacetanilide		3	18
2-Thiobarbituric acid		reacts at 10 mM	

TABLE VI-continued


<u>Identified agonists of hTAS2R38</u>			
Substance	Structure	Approx. Threshold [μ M]	EC ₅₀ [μ M]
2-Thiouracil		300	estimated 2000

TABLE VII



<u>Identified agonists of hTAS2R43</u>			
Substance	Structure	Approx. Threshold [mM]	EC ₅₀ [mM]
Saccharin		0.2	1.1
Acesulfame K		No response up to 10 mM	

TABLE VIII



<u>Identified agonists of hTAS2R44</u>			
Substance	Structure	Approx. Threshold [mM]	EC ₅₀ [mM]
Saccharin		0.2	estimated 2-5
Acesulfame K		0.5	3

TABLE IX


<u>Identified agonists of hTAS2R45</u>			
Substance	Structure	Approx. Threshold [mM]	EC ₅₀ [mM]
Absinthine		0.003	Not determined

TABLE X



Identified agonists of hTAS2R46			
Substance	Structure	Approx. Thresh- old [mM]	EC ₅₀ [mM]
Absinthine		0.001	Not determined

TABLE XI

Identified agonists of hTAS2R48			
Substance	Structure	Approx. Thresh- old [mM]	EC ₅₀ [mM]
Absinthine		0.03	Not determined

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 52

<210> SEQ ID NO 1

<211> LENGTH: 333

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Met Leu Thr Leu Thr Arg Ile Arg Thr Val Ser Tyr Glu Val Arg Ser
1 5 10 15

Thr Phe Leu Phe Ile Ser Val Leu Glu Phe Ala Val Gly Phe Leu Thr
20 25 30

Asn Ala Phe Val Phe Leu Val Asn Phe Trp Asp Val Val Lys Arg Gln
35 40 45

Ala Leu Ser Asn Ser Asp Cys Val Leu Leu Cys Leu Ser Ile Ser Arg
50 55 60

Leu Phe Leu His Gly Leu Leu Phe Leu Ser Ala Ile Gln Leu Thr His
65 70 75 80

Phe Gln Lys Leu Ser Glu Pro Leu Asn His Ser Tyr Gln Ala Ile Ile

-continued

85					90					95					
Met	Leu	Trp	Met	Ile	Ala	Asn	Gln	Ala	Asn	Leu	Trp	Leu	Ala	Ala	Cys
			100					105					110		
Leu	Ser	Leu	Leu	Tyr	Cys	Ser	Lys	Leu	Ile	Arg	Phe	Ser	His	Thr	Phe
			115				120					125			
Leu	Ile	Cys	Leu	Ala	Ser	Trp	Val	Ser	Arg	Lys	Ile	Ser	Gln	Met	Leu
			130				135					140			
Leu	Gly	Ile	Ile	Leu	Cys	Ser	Cys	Ile	Cys	Thr	Val	Leu	Cys	Val	Trp
	145					150					155				160
Cys	Phe	Phe	Ser	Arg	Pro	His	Phe	Thr	Val	Thr	Thr	Val	Leu	Phe	Met
			165					170					175		
Asn	Asn	Asn	Thr	Arg	Leu	Asn	Trp	Gln	Asn	Lys	Asp	Leu	Asn	Leu	Phe
			180					185					190		
Tyr	Ser	Phe	Leu	Phe	Cys	Tyr	Leu	Trp	Ser	Val	Pro	Pro	Phe	Leu	Leu
			195				200					205			
Phe	Leu	Val	Ser	Ser	Gly	Met	Leu	Thr	Val	Ser	Leu	Gly	Arg	His	Met
			210				215					220			
Arg	Thr	Met	Lys	Val	Tyr	Thr	Arg	Asn	Ser	Arg	Asp	Pro	Ser	Leu	Glu
			225				230					235			240
Ala	His	Ile	Lys	Ala	Leu	Lys	Ser	Leu	Val	Ser	Phe	Phe	Cys	Phe	Phe
			245					250					255		
Val	Ile	Ser	Ser	Cys	Val	Ala	Phe	Ile	Ser	Val	Pro	Leu	Leu	Ile	Leu
			260					265					270		
Trp	Arg	Asp	Lys	Ile	Gly	Val	Met	Val	Cys	Val	Gly	Ile	Met	Ala	Ala
			275				280					285			
Cys	Pro	Ser	Gly	His	Ala	Ala	Ile	Leu	Ile	Ser	Gly	Asn	Ala	Lys	Leu
			290				295					300			
Arg	Arg	Ala	Val	Met	Thr	Ile	Leu	Leu	Trp	Ala	Gln	Ser	Ser	Leu	Lys
			305				310					315			320
Val	Arg	Ala	Asp	His	Lys	Ala	Asp	Ser	Arg	Thr	Leu	Cys			
			325					330							

<210> SEQ ID NO 2

<211> LENGTH: 999

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

atgttgactc taactcgcat ccgcaactgtg tcctatgaag tcaggagtac atttctgttc	60
atttcagtcc tggagtttgc agtgggggtt ctgaccaatg ccttcgtttt ctgggtgaat	120
ttttgggatg tagtgaagag gcaggcactg agcaacagtg attgtgtgct gctgtgtctc	180
agcatcagcc ggcttttcct gcattggactg ctgttcctga gtgctatcca gcttaccac	240
ttccagaagt tgagtgaacc actgaaccac agctaccaag ccatcatcat gctatggatg	300
attgcaaacc aagccaacct ctggcttgct gectgcctca gectgcttta ctgctccaag	360
ctcatccgtt tctctcacac ctctctgacg tgcctggcaa gctgggtctc caggaagatc	420
tcccagatgc tcttgggtat tattctttgc tctgcatct gcaactgtcct ctgtgtttgg	480
tgctttttta gcagacctca cttcacagtc acaactgtgc tattcatgaa taacaataca	540
aggctcaact ggcagaataa agatctcaat ttattttatt cctttctctt ctgctatctg	600
tggtctgtgc ctcctttctt attgtttctg gtttcttctg ggatgctgac tgtctccctg	660

-continued

```

ggaaggcaca tgaggacaat gaaggtctat accagaaact ctcgtgaccc cagcctggag    720
gccacacatta aagccctcaa gtctcttgtc tcctttttct gcttctttgt gatatacatcc    780
tgtgttgccet tcattctctgt gccctactgt attctgtggc gcgacaaaat aggggtgatg    840
gtttgtgttg ggataatggc agcttggtccc tctgggcatg cagccatcct gatctcaggc    900
aatgccaaagt tgaggagagc tgtgatgacc attctgctct gggctcagag cagcctgaag    960
gtaagagccg accacaaggc agattcccg acactgtgc                                999

```

<210> SEQ ID NO 3

<211> LENGTH: 337

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```

Met Leu Gly Arg Cys Phe Pro Pro Asp Thr Lys Glu Lys Gln Gln Leu
1      5      10      15
Arg Met Thr Lys Leu Cys Asp Pro Ala Glu Ser Glu Leu Ser Pro Phe
20     25     30
Leu Ile Thr Leu Ile Leu Ala Val Leu Leu Ala Glu Tyr Leu Ile Gly
35     40     45
Ile Ile Ala Asn Gly Phe Ile Met Ala Ile His Ala Ala Glu Trp Val
50     55     60
Gln Asn Lys Ala Val Ser Thr Ser Gly Arg Ile Leu Val Phe Leu Ser
65     70     75     80
Val Ser Arg Ile Ala Leu Gln Ser Leu Met Met Leu Glu Ile Thr Ile
85     90     95
Ser Ser Thr Ser Leu Ser Phe Tyr Ser Glu Asp Ala Val Tyr Tyr Ala
100    105    110
Phe Lys Ile Ser Phe Ile Phe Leu Asn Phe Cys Ser Leu Trp Phe Ala
115    120    125
Ala Trp Leu Ser Phe Phe Tyr Phe Val Lys Ile Ala Asn Phe Ser Tyr
130    135    140
Pro Leu Phe Leu Lys Leu Arg Trp Arg Ile Thr Gly Leu Ile Pro Trp
145    150    155    160
Leu Leu Trp Leu Ser Val Phe Ile Ser Phe Ser His Ser Met Phe Cys
165    170    175
Ile Asn Ile Cys Thr Val Tyr Cys Asn Asn Ser Phe Pro Ile His Ser
180    185    190
Ser Asn Ser Thr Lys Lys Thr Tyr Leu Ser Glu Ile Asn Val Val Gly
195    200    205
Leu Ala Phe Phe Phe Asn Leu Gly Ile Val Thr Pro Leu Ile Met Phe
210    215    220
Ile Leu Thr Ala Thr Leu Leu Ile Leu Ser Leu Lys Arg His Thr Leu
225    230    235    240
His Met Gly Ser Asn Ala Thr Gly Ser Asn Asp Pro Ser Met Glu Ala
245    250    255
His Met Gly Ala Ile Lys Ala Ile Ser Tyr Phe Leu Ile Leu Tyr Ile
260    265    270
Phe Asn Ala Val Ala Leu Phe Ile Tyr Leu Ser Asn Met Phe Asp Ile
275    280    285
Asn Ser Leu Trp Asn Asn Leu Cys Gln Ile Ile Met Ala Ala Tyr Pro

```

-continued

290	295	300	
Ala Ser His Ser Ile Leu Leu Ile Gln Asp Asn Pro Gly Leu Arg Arg			
305	310	315	320
Ala Trp Ser Gly Phe Ser Phe Asp Phe Ile Phe Thr Gln Lys Ser Gly			
	325	330	335

Leu

<210> SEQ ID NO 4
 <211> LENGTH: 1013
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

atgctagga gatgttttcc tccagacacc aaagagaagc aacagctcag aatgactaaa	60
ctctgcgac ctgcagaaag tgaattgtcg ccattttctca tcaccttaat ttagcagtt	120
ttacttgctg aatacctcat tggatcatt gcaaattggt tcatcatggc tatacatgca	180
gctgaatggg ttcaaaaataa ggcaagtttcc acaagtggca ggatcctggt ttctctgagt	240
gtatccagaa tagctctcca aagcctcatg atgttagaaa ttaccatcag ctcaacctcc	300
ctaagttttt attctgaaga cgtgtatat tatgcattca aaataagttt tatattctta	360
aatttttgta gcctgtggtt tgcctgctgg ctgagtttct tctactttgt gaagattgcc	420
aattttcctt accccctttt cctcaaaactg aggtggagaa ttaactggatt gataccctgg	480
cttctgtggc tgtccgtgtt tatttccttc agtcacagca tgttctgcat caacatctgc	540
actgtgtatt gtaacaattc ttccctatc cactcctcca actccactaa gaaaacatac	600
ttgtctgaga tcaatgtggt cggctctggc tttttcttta acctggggat tgtgactcct	660
ctgatcatgt tcatcctgac agccaccctg ctgatcctct ctctcaagag acacacccta	720
cacatgggaa gcaatgccac aggggtccaac gacccagca tggaggctca catgggggcc	780
atcaaagcta tcagctactt tctcattctc tacattttca atgcagttgc tctgtttatc	840
tacctgtcca acatgtttga catcaacagt ctgtggaata atttgtgcca gatcatcatg	900
gctgcctacc ctgccagcca ctcaattcta ctgattcaag ataaccctgg gctgagaaga	960
gcctggagcg gcttcagctt cgacttcato tttacccaaa agagtggact ctg	1013

<210> SEQ ID NO 5
 <211> LENGTH: 323
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Met Ala Thr Val Asn Thr Asp Ala Thr Asp Lys Asp Ile Ser Lys Phe	
1	15
Lys Val Thr Phe Thr Leu Val Val Ser Gly Ile Glu Cys Ile Thr Gly	
20	30
Ile Leu Gly Ser Gly Phe Ile Thr Ala Ile Tyr Gly Ala Glu Trp Ala	
35	45
Arg Gly Lys Thr Leu Pro Thr Gly Asp Arg Ile Met Leu Met Leu Ser	
50	60
Phe Ser Arg Leu Leu Leu Gln Ile Trp Met Met Leu Glu Asn Ile Phe	
65	80
Ser Leu Leu Phe Arg Ile Val Tyr Asn Gln Asn Ser Val Tyr Ile Leu	

-continued

85					90					95						
Phe	Lys	Val	Ile	Thr	Val	Phe	Leu	Asn	His	Ser	Asn	Leu	Trp	Phe	Ala	
100					105					110						
Ala	Trp	Leu	Lys	Val	Phe	Tyr	Cys	Leu	Arg	Ile	Ala	Asn	Phe	Asn	His	
115					120					125						
Pro	Leu	Phe	Phe	Leu	Met	Lys	Arg	Lys	Ile	Ile	Val	Leu	Met	Pro	Trp	
130					135					140						
Leu	Leu	Arg	Leu	Ser	Val	Leu	Val	Ser	Leu	Ser	Phe	Ser	Phe	Pro	Leu	
145					150					155					160	
Ser	Arg	Asp	Val	Phe	Asn	Val	Tyr	Val	Asn	Ser	Ser	Ile	Pro	Ile	Pro	
165					170					175						
Ser	Ser	Asn	Ser	Thr	Glu	Lys	Lys	Tyr	Phe	Ser	Glu	Thr	Asn	Met	Val	
180					185					190						
Asn	Leu	Val	Phe	Phe	Tyr	Asn	Met	Gly	Ile	Phe	Val	Pro	Leu	Ile	Met	
195					200					205						
Phe	Ile	Leu	Ala	Ala	Thr	Leu	Leu	Ile	Leu	Ser	Leu	Lys	Arg	His	Thr	
210					215					220						
Leu	His	Met	Gly	Ser	Asn	Ala	Thr	Gly	Ser	Arg	Asp	Pro	Ser	Met	Lys	
225					230					235					240	
Ala	His	Ile	Gly	Ala	Ile	Lys	Ala	Thr	Ser	Tyr	Phe	Leu	Ile	Leu	Tyr	
245					250					255						
Ile	Phe	Asn	Ala	Ile	Ala	Leu	Phe	Leu	Ser	Thr	Ser	Asn	Ile	Phe	Asp	
260					265					270						
Thr	Tyr	Ser	Ser	Trp	Asn	Ile	Leu	Cys	Lys	Ile	Ile	Met	Ala	Ala	Tyr	
275					280					285						
Pro	Ala	Gly	His	Ser	Val	Gln	Leu	Ile	Leu	Gly	Asn	Pro	Gly	Leu	Arg	
290					295					300						
Arg	Ala	Trp	Lys	Arg	Phe	Gln	His	Gln	Val	Pro	Leu	Tyr	Leu	Lys	Gly	
305					310					315					320	

Gln Thr Leu

<210> SEQ ID NO 6

<211> LENGTH: 969

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

atggcaacgg tgaacacaga tgccacagat aaagacatat ccaagttcaa ggtcaccttc	60
actttggtgg tctccggaat agagtgcac accggcatcc ttgggagtgg cttcatcacg	120
gccatctatg gggctgagtg ggccaggggc aaaacactcc ccaactggtga ccgcattatg	180
ttgatgctga gcttttccag gctcttgcta cagatttgga tgatgctgga gaacattttc	240
agtcgtctat tccgaattgt ttataaccaa aactcagtgt atatcctctt caaagtcac	300
actgtctttc tgaaccattc caatctctgg tttgctgcct ggctcaaagt cttctattgt	360
cttagaattg caaacttcaa tcattccttg ttcttctgta tgaagaggaa aatcatagtg	420
ctgatgcctt ggcttctcag gctgtcagtg ttggtttcct taagcttcag ctttctcttc	480
tcgagagatg tcttcaatgt gtatgtgaat agctccattc ctatccccct ctccaactcc	540
acggagaaga agtactcttc tgagaccaat atggtcaacc tgggtatttt ctataacatg	600
gggatcttcg ttcctctgat catgttcac ctggcagcca ccctgctgat cctctctctc	660

-continued

```

aagagacaca ccctacacat gggaagcaat gccacaggt ccagggaccc cagcatgaag 720
gctcacatag gggccatcaa agccaccagc tactttctca tcctctacat tttcaatgca 780
attgctctat ttctttccac gtccaacatc tttgacactt acagtctctg gaatatatttg 840
tgcaagatca tcatggctgc ctaccctgcc ggccactcag tacaactgat cttgggcaac 900
cctgggctga gaagagcctg gaagcgggtt cagcaccaag ttctctctta cctaaaaggg 960
cagactctg 969

```

<210> SEQ ID NO 7

<211> LENGTH: 307

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```

Met  Gln  Ala  Ala  Leu  Thr  Ala  Phe  Phe  Val  Leu  Leu  Phe  Ser  Leu  Leu
 1          5          10          15

Ser  Leu  Leu  Gly  Ile  Ala  Ala  Asn  Gly  Phe  Ile  Val  Leu  Val  Leu  Gly
          20          25          30

Arg  Glu  Trp  Leu  Arg  Tyr  Gly  Arg  Leu  Leu  Pro  Leu  Asp  Met  Ile  Leu
          35          40          45

Ile  Ser  Leu  Gly  Ala  Ser  Arg  Phe  Cys  Leu  Gln  Leu  Val  Gly  Thr  Val
          50          55          60

His  Asn  Phe  Tyr  Tyr  Ser  Ala  Gln  Lys  Val  Glu  Tyr  Ser  Gly  Gly  Leu
65          70          75          80

Gly  Arg  Gln  Phe  Phe  His  Leu  His  Trp  His  Phe  Leu  Asn  Ser  Ala  Thr
          85          90          95

Phe  Trp  Phe  Cys  Ser  Trp  Leu  Ser  Val  Leu  Phe  Cys  Val  Lys  Ile  Ala
          100          105          110

Asn  Ile  Thr  His  Ser  Thr  Phe  Leu  Trp  Leu  Lys  Trp  Arg  Phe  Leu  Gly
          115          120          125

Trp  Val  Pro  Trp  Leu  Leu  Leu  Gly  Ser  Val  Leu  Ile  Ser  Phe  Ile  Ile
          130          135          140

Thr  Leu  Leu  Phe  Phe  Trp  Val  Asn  Tyr  Pro  Val  Tyr  Gln  Glu  Phe  Leu
145          150          155          160

Ile  Arg  Lys  Phe  Ser  Gly  Asn  Met  Thr  Tyr  Lys  Trp  Asn  Thr  Arg  Ile
          165          170          175

Glu  Thr  Tyr  Tyr  Phe  Pro  Ser  Leu  Lys  Leu  Val  Ile  Trp  Ser  Ile  Pro
          180          185          190

Phe  Ser  Val  Phe  Leu  Val  Ser  Ile  Met  Leu  Leu  Ile  Asn  Ser  Leu  Arg
          195          200          205

Arg  His  Thr  Gln  Arg  Met  Gln  His  Asn  Gly  His  Ser  Leu  Gln  Asp  Pro
          210          215          220

Ser  Thr  Gln  Ala  His  Thr  Arg  Ala  Leu  Lys  Ser  Leu  Ile  Ser  Phe  Leu
225          230          235          240

Ile  Leu  Tyr  Ala  Leu  Ser  Phe  Leu  Ser  Leu  Ile  Ile  Asp  Ala  Ala  Lys
          245          250          255

Phe  Ile  Ser  Met  Gln  Asn  Asp  Phe  Tyr  Trp  Pro  Trp  Gln  Ile  Ala  Val
          260          265          270

Tyr  Leu  Cys  Ile  Ser  Val  His  Pro  Phe  Ile  Leu  Ile  Phe  Ser  Asn  Leu
          275          280          285

Lys  Leu  Arg  Ser  Val  Phe  Ser  Gln  Leu  Leu  Leu  Leu  Ala  Arg  Gly  Phe
          290          295          300

```

-continued

Trp Val Ala
305<210> SEQ ID NO 8
<211> LENGTH: 921
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```
atgcaagcag cactgacggc cttcttcgtg ttgctcttta gcctgctgag tcttctgggg    60
attgcagcga atgggttcat tgtgctggtg ctgggcaggg agtggctgcg atatggcagg    120
ttgctgcctt tggatatgat cctcattagc ttgggtgcct cccgcttctg cctgcagttg    180
gttgggacag tgcacaactt ctactactct gcccagaagg tcgagtactc tgggggtctc    240
ggccgacagt tcttccatct acactggcac ttcttgaact cagccacctt ctggttttgc    300
agctggctca gtgtcctggt ctgtgtgaag attgctaaca tcacacactc caccttcctg    360
tggtgaagt  ggaggttctt aggggtgggtg ccttggtctc tgttgggtct tgcctgac    420
tccttcacat taacctgctt gtttttttgg gtgaactacc ctgtatatca agaattttta    480
attagaaaat tttctgggaa catgacctac aagtgaata caaggataga aacatactat    540
ttcccatccc tgaaactggt catctggtca attccttttt ctgtttttct ggtctcaatt    600
atgctgttaa ttaattctct gaggaggcat actcagagaa tgcagcacia cgggcacagc    660
ctgcaggacc ccagcaccca ggctcacacc agagctctga agtccctcat ctcttctctc    720
attcttttat ctctgtcctt tctgtccctg atcattgatg ccgcaaaatt tatctccatg    780
cagaacgact tttactggcc atggcaaatt gcagtctacc tgtgcatatc tgtccatccc    840
ttcatcctca tcttcagcaa cctcaagctt cgaagcgtgt tctcacagct cctgttgttg    900
gcaaggggct tctgggtggc c                                         921
```

<210> SEQ ID NO 9
<211> LENGTH: 309
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```
Met Ile Thr Phe Leu Pro Ile Ile Phe Ser Ser Leu Val Val Val Thr
 1             5             10            15
Phe Val Ile Gly Asn Phe Ala Asn Gly Phe Ile Ala Leu Val Asn Ser
      20             25             30
Ile Glu Ser Phe Lys Arg Gln Lys Ile Ser Phe Ala Asp Gln Ile Leu
      35             40             45
Thr Ala Leu Ala Val Ser Arg Val Gly Leu Leu Trp Val Leu Leu Leu
      50             55             60
Asn Trp Tyr Ser Thr Val Leu Asn Pro Ala Phe Asn Ser Val Glu Val
      65             70             75             80
Arg Thr Thr Ala Tyr Asn Ile Trp Ala Val Ile Asn His Phe Ser Asn
      85             90             95
Trp Leu Ala Thr Thr Leu Ser Ile Phe Tyr Leu Leu Lys Ile Ala Asn
      100            105            110
Phe Ser Asn Phe Ile Phe Leu His Leu Lys Arg Arg Val Lys Ser Val
      115            120            125
```

-continued

Ile	Leu	Val	Met	Leu	Leu	Gly	Pro	Leu	Leu	Phe	Leu	Ala	Cys	His	Leu
130						135					140				
Phe	Val	Ile	Asn	Met	Asn	Glu	Ile	Val	Arg	Thr	Lys	Glu	Phe	Glu	Gly
145					150					155				160	
Asn	Met	Thr	Trp	Lys	Ile	Lys	Leu	Lys	Ser	Ala	Met	Tyr	Phe	Ser	Asn
				165					170					175	
Met	Thr	Val	Thr	Met	Val	Ala	Asn	Leu	Val	Pro	Phe	Thr	Leu	Thr	Leu
				180				185					190		
Leu	Ser	Phe	Met	Leu	Leu	Ile	Cys	Ser	Leu	Cys	Lys	His	Leu	Lys	Lys
	195						200					205			
Met	Gln	Leu	Arg	Gly	Lys	Gly	Ser	Gln	Asp	Pro	Ser	Thr	Lys	Val	His
210						215					220				
Ile	Lys	Ala	Leu	Gln	Thr	Val	Ile	Ser	Phe	Leu	Leu	Cys	Ala	Ile	
225				230						235				240	
Tyr	Phe	Leu	Ser	Ile	Met	Ile	Ser	Val	Trp	Ser	Phe	Gly	Ser	Leu	Glu
				245					250					255	
Asn	Lys	Pro	Val	Phe	Met	Phe	Cys	Lys	Ala	Ile	Arg	Phe	Ser	Tyr	Pro
			260					265					270		
Ser	Ile	His	Pro	Phe	Ile	Leu	Ile	Trp	Gly	Asn	Lys	Lys	Leu	Lys	Gln
		275					280					285			
Thr	Phe	Leu	Ser	Val	Phe	Trp	Gln	Met	Arg	Tyr	Trp	Val	Lys	Gly	Glu
	290					295					300				
Lys	Thr	Ser	Ser	Pro											
305															

<210> SEQ ID NO 10

<211> LENGTH: 927

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

atgataactt ttctacccat cattttttcc agtctggtag tggttacatt tgttattgga	60
aattttgcta atgggttcac agcactggta aattccattg agtcgttcaa gagacaaaag	120
atctcctttg ctgaccaaata tctcactgct ctggcggctc ccagagttgg tttgctctgg	180
gtattattat taaactggta ttcaactgtg ttgaatccag cttttaatag tgtagaagta	240
agaactactg cttataatat ctgggcagtg atcaaccatt tcagcaactg gcttgctact	300
acctcagca tattttattt gctcaagatt gccaatctct ccaactttat tttctttcac	360
ttaaagagga gagttaagag tgtcattctg gtgatgttgt tggggccttt gctatttttg	420
gcttgctcgc tttttgtgat aaacatgaat gagattgtgc ggacaaaaga atttgaagga	480
aacatgactt ggaagatcaa attgaagagt gcaatgtact tttcaaatat gactgtaacc	540
atggtagcaa acttagtacc cttcactctg acctactat cttttatgct gttaactctgt	600
tctttgtgta aacatctcaa gaagatgcag ctccgtggta aaggatctca agatcccagc	660
acgaaggtcc acataaaagc ttgcaaaact gtgatctcct tcctcttggt atgtgccatt	720
tactttctgt ccataatgat atcagtttgg agttttggaa gtctggaaaa caaacctgtc	780
ttcatgttct gcaaagctat tagattcagc tatccttcaa tccaccatt catcctgatt	840
tggggaaaca agaagctaaa gcagactttt ctttcagttt tttggcaaat gaggtactgg	900
gtgaaaggag agaagacttc atctcca	927

-continued

<210> SEQ ID NO 11
 <211> LENGTH: 309
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

```

Met Thr Thr Phe Ile Pro Ile Ile Phe Ser Ser Val Val Val Val Leu
1      5      10      15

Phe Val Ile Gly Asn Phe Ala Asn Gly Phe Ile Ala Leu Val Asn Ser
20      25      30

Ile Glu Arg Val Lys Arg Gln Lys Ile Ser Phe Ala Asp Gln Ile Leu
35      40      45

Thr Ala Leu Ala Val Ser Arg Val Gly Leu Leu Trp Val Leu Leu Leu
50      55      60

Asn Trp Tyr Ser Thr Val Phe Asn Pro Ala Phe Tyr Ser Val Glu Val
65      70      75      80

Arg Thr Thr Ala Tyr Asn Val Trp Ala Val Thr Gly His Phe Ser Asn
85      90      95

Trp Leu Ala Thr Ser Leu Ser Ile Phe Tyr Leu Leu Lys Ile Ala Asn
100     105     110

Phe Ser Asn Leu Ile Phe Leu His Leu Lys Arg Arg Val Lys Ser Val
115     120     125

Ile Leu Val Met Leu Leu Gly Pro Leu Leu Phe Leu Ala Cys Gln Leu
130     135     140

Phe Val Ile Asn Met Lys Glu Ile Val Arg Thr Lys Glu Tyr Glu Gly
145     150     155     160

Asn Met Thr Trp Lys Ile Lys Leu Arg Ser Ala Val Tyr Leu Ser Asp
165     170     175

Ala Thr Val Thr Thr Leu Gly Asn Leu Val Pro Phe Thr Leu Thr Leu
180     185     190

Leu Cys Phe Leu Leu Leu Ile Cys Ser Leu Cys Lys His Leu Lys Lys
195     200     205

Met Gln Leu His Gly Lys Gly Ser Gln Asp Pro Ser Thr Lys Val His
210     215     220

Ile Lys Ala Leu Gln Thr Val Ile Phe Phe Leu Leu Leu Cys Ala Val
225     230     235     240

Tyr Phe Leu Ser Ile Met Ile Ser Val Trp Ser Phe Gly Ser Leu Glu
245     250     255

Asn Lys Pro Val Phe Met Phe Cys Lys Ala Ile Arg Phe Ser Tyr Pro
260     265     270

Ser Ile His Pro Phe Ile Leu Ile Trp Gly Asn Lys Lys Leu Lys Gln
275     280     285

Thr Phe Leu Ser Val Leu Arg Gln Val Arg Tyr Trp Val Lys Gly Glu
290     295     300

Lys Pro Ser Ser Pro
305

```

<210> SEQ ID NO 12
 <211> LENGTH: 927
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

-continued

```

atgacaactt ttatacccat cttttttcc agtgtgtag tgggtctatt tgtattgga      60
aattttgcta atgggttcat agcattggta aattccattg agcgggtcaa gagacaaaag    120
atctcttttg ctgaccagat tctcactgct ctggcggctc ccagagttag tttgctctgg    180
gtattattat taaattggta ttcaactgtg tttaatccag ctttttatag tgtagaagta    240
agaactactg cttataatgt ctgggcagta accggccatt tcagcaactg gcttgctact    300
agcctcagca tattttatatt gctcaagatt gccaatctct ccaaccttat ttttcttcac   360
ttaagagga gagttaagag tgcattctg gtgatgctgt tggggccttt actatttttg    420
gcttgctaac tttttgtgat aaacatgaaa gagattgtac ggacaaaaga atatgaagga    480
aacatgactt ggaagatcaa attgaggagt gcagtgtacc tttcagatgc gactgtaacc    540
acgctaggaa acttagtgcc cttcactctg accctgctat gttttttgct gttaatctgt    600
tctctgtgta aacatctcaa gaagatgcag ctccatgcta aaggatctca agatcccagc   660
accaaggtcc acataaaaagc ttgcaaact gtgatctttt tcctcttggt atgtgccgtt    720
tactttctgt ccataatgat atcagtttgg agttttggga gtctggaaaa caaacctgtc    780
ttcatgttct gcaaagctat tagattcagc tacccttcaa tccaccatt catcctgatt    840
tggggaaaca agaagctaaa gcagactttt ctttcagttt tgccggcaagt gaggtactgg    900
gtgaaaggag agaagccttc atctcca                                     927

```

<210> SEQ ID NO 13

<211> LENGTH: 299

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

```

Met Ile Thr Phe Leu Pro Ile Ile Phe Ser Ile Leu Val Val Val Thr
1           5           10          15

Phe Val Ile Gly Asn Phe Ala Asn Gly Phe Ile Ala Leu Val Asn Ser
20          25          30

Thr Glu Trp Val Lys Arg Gln Lys Ile Ser Phe Ala Asp Gln Ile Val
35          40          45

Thr Ala Leu Ala Val Ser Arg Val Gly Leu Leu Trp Val Leu Leu Leu
50          55          60

Asn Trp Tyr Ser Thr Val Leu Asn Pro Ala Phe Cys Ser Val Glu Leu
65          70          75          80

Arg Thr Thr Ala Tyr Asn Ile Trp Ala Val Thr Gly His Phe Ser Asn
85          90          95

Trp Pro Ala Thr Ser Leu Ser Ile Phe Tyr Leu Leu Lys Ile Ala Asn
100         105         110

Phe Ser Asn Leu Ile Phe Leu Arg Leu Lys Arg Arg Val Lys Ser Val
115         120         125

Ile Leu Val Val Leu Leu Gly Pro Leu Leu Phe Leu Ala Cys His Leu
130         135         140

Phe Val Val Asn Met Asn Gln Ile Val Trp Thr Lys Glu Tyr Glu Gly
145         150         155         160

Asn Met Thr Trp Lys Ile Lys Leu Arg Arg Ala Met Tyr Leu Ser Asp
165         170         175

Thr Thr Val Thr Met Leu Ala Asn Leu Val Pro Phe Thr Val Thr Leu
180         185         190

```

-continued

Ile	Ser	Phe	Leu	Leu	Leu	Val	Cys	Ser	Leu	Cys	Lys	His	Leu	Lys	Lys
		195					200					205			
Met	Gln	Leu	His	Gly	Lys	Gly	Ser	Gln	Asp	Pro	Ser	Thr	Lys	Val	His
	210					215					220				
Ile	Lys	Val	Leu	Gln	Thr	Val	Ile	Ser	Phe	Phe	Leu	Leu	Arg	Ala	Ile
225					230					235					240
Tyr	Phe	Val	Ser	Val	Ile	Ile	Ser	Val	Trp	Ser	Phe	Lys	Asn	Leu	Glu
				245					250					255	
Asn	Lys	Pro	Val	Phe	Met	Phe	Cys	Gln	Ala	Ile	Gly	Phe	Ser	Cys	Ser
			260					265					270		
Ser	Ala	His	Pro	Phe	Ile	Leu	Ile	Trp	Gly	Asn	Lys	Lys	Leu	Lys	Gln
		275					280					285			
Thr	Tyr	Leu	Ser	Val	Leu	Trp	Gln	Met	Arg	Tyr					
	290					295									

<210> SEQ ID NO 14

<211> LENGTH: 897

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

```

atgataactt ttctgcccat catattttcc attctagtag tggttacatt tgttattgga      60
aattttgcta atggcttcat agcgttggtta aattccaccg agtgggtgaa gagacaaaag      120
atctcctttg ctgaccaaat tgtcactgct ctggcgggtct ccagagttgg tttgctctgg      180
gtgttattat taaattggta ttcaactgtg ttgaatccag cttttttag tagtagaatta      240
agaactactg cttataatat ctgggcagta accggccatt tcagcaactg gcctgctact      300
agcctcagca tatttttatt gctcaagatt gccaatctct ccaaccttat ttttcttcgc      360
ttaaagagga gagttaagag tgtcattctg gtggtgctgt tggggccttt gctatttttg      420
gcttgctcgc tttttgtggt aaacatgaat cagattgtat ggacaaaaga atatgaagga      480
aacatgactt ggaagatcaa attgaggcgt gcaatgtacc tttcagatac gactgtaacc      540
atgctagcaa acttagtacc ctttactgta accctgatat cttttctgct gtagtctgt      600
tctctgtgta aacatctcaa gaagatgcag ctccatggca aaggatctca agatccagat      660
accaaggtcc acataaaaagt tttgcaaact gtgatctcct tcttcttggt acgtgccatt      720
tactttgtgt ctgtaataat atcagtttgg agttttaaga atctggaaaa caaacctgtc      780
ttcatgttct gccaaactat tggattcagc tgttcttcag cccacccggt catcctgatt      840
tggggaaaca agaagctaaa gcagacttat ctttcagttt tgtggcaaat gaggtac      897

```

<210> SEQ ID NO 15

<211> LENGTH: 299

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

Met	Ile	Thr	Phe	Leu	Pro	Ile	Ile	Phe	Ser	Ile	Leu	Ile	Val	Val	Thr
1			5					10					15		
Phe	Val	Ile	Gly	Asn	Phe	Ala	Asn	Gly	Phe	Ile	Ala	Leu	Val	Asn	Ser
		20					25					30			
Ile	Glu	Trp	Phe	Lys	Arg	Gln	Lys	Ile	Ser	Phe	Ala	Asp	Gln	Ile	Leu
	35						40					45			

-continued

Thr	Ala	Leu	Ala	Val	Ser	Arg	Val	Gly	Leu	Leu	Trp	Val	Leu	Val	Leu
50						55					60				
Asn	Trp	Tyr	Ala	Thr	Glu	Leu	Asn	Pro	Ala	Phe	Asn	Ser	Ile	Glu	Val
65					70					75				80	
Arg	Ile	Thr	Ala	Tyr	Asn	Val	Trp	Ala	Val	Ile	Asn	His	Phe	Ser	Asn
				85					90					95	
Trp	Leu	Ala	Thr	Ser	Leu	Ser	Ile	Phe	Tyr	Leu	Leu	Lys	Ile	Ala	Asn
			100					105					110		
Phe	Ser	Asn	Leu	Ile	Phe	Leu	His	Leu	Lys	Arg	Arg	Val	Lys	Ser	Val
		115					120					125			
Val	Leu	Val	Ile	Leu	Leu	Gly	Pro	Leu	Leu	Phe	Leu	Val	Cys	His	Leu
	130					135					140				
Phe	Val	Ile	Asn	Met	Asn	Gln	Ile	Ile	Trp	Thr	Lys	Glu	Tyr	Glu	Gly
145					150					155					160
Asn	Met	Thr	Trp	Lys	Ile	Lys	Leu	Arg	Ser	Ala	Met	Tyr	Leu	Ser	Asn
			165						170					175	
Thr	Thr	Val	Thr	Ile	Leu	Ala	Asn	Leu	Val	Pro	Phe	Thr	Leu	Thr	Leu
			180					185					190		
Ile	Ser	Phe	Leu	Leu	Leu	Ile	Cys	Ser	Leu	Cys	Lys	His	Leu	Lys	Lys
		195					200					205			
Met	Gln	Leu	His	Gly	Lys	Gly	Ser	Gln	Asp	Pro	Ser	Met	Lys	Val	His
	210					215					220				
Ile	Lys	Ala	Leu	Gln	Thr	Val	Thr	Ser	Phe	Leu	Leu	Leu	Cys	Ala	Ile
225					230					235					240
Tyr	Phe	Leu	Ser	Ile	Ile	Met	Ser	Val	Trp	Ser	Phe	Glu	Ser	Leu	Glu
			245						250					255	
Asn	Lys	Pro	Val	Phe	Met	Phe	Cys	Glu	Ala	Ile	Ala	Phe	Ser	Tyr	Pro
			260					265					270		
Ser	Thr	His	Pro	Phe	Ile	Leu	Ile	Trp	Gly	Asn	Lys	Lys	Leu	Lys	Gln
		275						280				285			
Thr	Phe	Leu	Ser	Val	Leu	Trp	Gln	Met	Arg	Tyr					
		290				295									

<210> SEQ ID NO 16

<211> LENGTH: 897

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

atgataactt ttctgcccat cattttttcc attctaatag tggttacatt tgtgattgga	60
aattttgcta atggcttcat agcattggta aattccattg agtggtttaa gagacaaaag	120
atctcttttg ctgaccaaatt tctcactgct ctggcagtct ccagagtggg tttactctgg	180
gtattagtat taaattggta tgcaactgag ttgaatccag cttttaacag tatagaagta	240
agaattactg cttacaatgt ctgggcagta atcaaccatt tcagcaactg gcttgctact	300
agcctcagca tattttattt gctcaagatt gccaatctct ccaaccttat tttcttcac	360
ttaaagagga gagttaagag tgtgtttctg gtgatactat tggggccttt gctatttttg	420
gtttgtcatc tttttgtgat aaacatgaat cagattatat ggacaaaaga atatgaagga	480
aacatgactt ggaagatcaa actgaggagt gcaatgtacc tttcaaatac aacggttaac	540
atcctagcaa acttagttcc cttcactctg accctgatat cttttctgct gttaactctgt	600

```
<210> SEQ ID NO 17
<211> LENGTH: 308
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

[illegible]

-continued

305

<210> SEQ ID NO 18

<211> LENGTH: 926

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

```
atgataactt ttctgcccat cttttttcc attctaata taggtatatt tgttattgga      60
aattttgcta atggcttcat agcattggta aattccattg agtgggtcaa gagacaaaag    120
atctcctttg ttgaccaaatt tctcactgct ctggcggctct ccagagttgg tttgctctgg   180
gtgttattac tacattggta tgcaactcag ttgaatccag ctttttatag tgtagaagta    240
agaattactg cttataatgt ctgggcagta accaaccatt tcagcagctg gcttgctact    300
agcctcagca tgttttatatt gctcaggatt gccaatctct ccaaccttat ttttcttcgc   360
ataaagagga gagttaagag tgttgttctg gtgatactgt tggggccttt gctatttttg    420
gtttgtcatc tttttgtgat aaacatggat gagactgtat ggacaaaaga atatgaagga    480
aacgtgactt ggaagatcaa attgaggagt gcaatgtacc attcaaatat gactctaacc    540
atgctagcaa actttgtacc cctcactctg accctgatat cttttctgct gttaactctgt   600
tctctgtgta aacatctcaa gaagatgcag ctccatggca aaggatctca agatcccagc   660
accaaggtcc acataaaaagc tttgaaaact gtgacctcct ttcttctggt atgtgccatt    720
tactttctgt ccatgatcat atcagtttgt aattttggga ggctggaaaa gcaacctgtc   780
ttcatgttct gccaagetat tatattcagc tatecttcaa cccaccatt catcctgatt    840
ttgggaaaca agaagctaaa gcagattttt ctttcagttt tgccggcatgt gaggtactgg   900
gtgaaagaca gaagccttcg tctcca                                     926
```

<210> SEQ ID NO 19

<211> LENGTH: 298

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

```
Met Met Cys Phe Leu Leu Ile Ile Ser Ser Ile Leu Val Val Phe Ala
1           5           10          15

Phe Val Leu Gly Asn Val Ala Asn Gly Phe Ile Ala Leu Val Asn Val
20          25          30

Ile Asp Trp Val Asn Thr Arg Lys Ile Ser Ser Ala Glu Gln Ile Leu
35          40          45

Thr Ala Leu Val Val Ser Arg Ile Gly Leu Leu Trp Val Met Leu Phe
50          55          60

Leu Trp Tyr Ala Thr Val Phe Asn Ser Ala Leu Tyr Gly Leu Glu Val
65          70          75          80

Arg Ile Val Ala Ser Asn Ala Trp Ala Val Thr Asn His Phe Ser Met
85          90          95

Trp Leu Ala Ala Ser Leu Ser Ile Phe Cys Leu Leu Lys Ile Ala Asn
100         105         110

Phe Ser Asn Leu Ile Ser Leu His Leu Lys Lys Arg Ile Lys Ser Val
115         120         125

Val Leu Val Ile Leu Leu Gly Pro Leu Val Phe Leu Ile Cys Asn Leu
130         135         140
```

-continued

Ala Val Ile Thr Met Asp Glu Arg Val Trp Thr Lys Glu Tyr Glu Gly
 145 150 155 160

Asn Val Thr Trp Lys Ile Lys Leu Arg Asn Ala Ile His Leu Ser Ser
 165 170 175

Leu Thr Val Thr Thr Leu Ala Asn Leu Ile Pro Phe Thr Leu Ser Leu
 180 185 190

Ile Cys Phe Leu Leu Leu Ile Cys Ser Leu Cys Lys His Leu Lys Lys
 195 200 205

Met Arg Leu His Ser Lys Gly Ser Gln Asp Pro Ser Thr Lys Val His
 210 215 220

Ile Lys Ala Leu Gln Thr Val Thr Ser Phe Leu Met Leu Phe Ala Ile
 225 230 235 240

Tyr Phe Leu Cys Ile Ile Thr Ser Thr Trp Asn Leu Arg Thr Gln Gln
 245 250 255

Ser Lys Leu Val Leu Leu Leu Cys Gln Thr Val Ala Ile Met Tyr Pro
 260 265 270

Ser Phe His Ser Phe Ile Leu Ile Met Gly Ser Arg Lys Leu Lys Gln
 275 280 285

Thr Phe Leu Ser Val Leu Trp Gln Met Thr
 290 295

<210> SEQ ID NO 20

<211> LENGTH: 897

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

```

atgatgtgtt ttctgctcat catttcatca attctggtag tgtttgcaatt tgttcttgga      60
aatgttgcca atggettcac agccctagta aatgtcattg actgggttaa cacacgaaag      120
atctcctcag ctgagcaaat tctcactgct ctggtggtct ccagaattgg tttactctgg      180
gtcatgttat tccttttgga tgcaactgtg ttttaattctg ctttatatgg tttagaagta      240
agaattgttg cttctaatagc ctgggctgta acgaaccatt tcagcatgtg gcttgcgtct      300
agcctcagca tattttgttt gctcaagatt gccaatctct ccaaccttat ttctctccac      360
ctaaagaaga gaattaagag tgttgttctg gtgatactgt tggggccctt ggtatctttg      420
atttgtaatc ttgctgtgat aaccatggat gagagagtgt ggacaaaaga atatgaagga      480
aatgtgactt ggaagatcaa atgaggaat gcaatacacc tttcaagctt gactgtaact      540
actctagcaa acctcatacc ctttactctg agcctaatat gttttctgct gttaatctgt      600
tctctttgta aacatctcaa gaagatgcgg ctccatagca aaggatctca agatcccagc      660
accaaggtcc atataaaagc tttgcaaact gtgacctcct tctcatgtt atttgccatt      720
tactttctgt gtataatcac atcaacttgg aatcttagga cacagcagag caaacttgta      780
ctctgctttt gccaaactgt tgcaatcatg tatccttcat tccactcatt cactctgatt      840
atgggaagta ggaagctaaa acagaccttt ctttcagttt tgtggcagat gacacgc      897

```

<210> SEQ ID NO 21

<211> LENGTH: 309

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

-continued

```

Met Met Ser Phe Leu His Ile Val Phe Ser Ile Leu Val Val Val Ala
1           5           10           15

Phe Ile Leu Gly Asn Phe Ala Asn Gly Phe Ile Ala Leu Ile Asn Phe
20           25           30

Ile Ala Trp Val Lys Arg Gln Lys Ile Ser Ser Ala Asp Gln Ile Ile
35           40           45

Ala Ala Leu Ala Val Ser Lys Val Gly Leu Leu Trp Val Ile Leu Leu
50           55           60

His Trp Tyr Ser Thr Val Leu Asn Pro Thr Ser Ser Asn Leu Lys Val
65           70           75           80

Ile Ile Phe Ile Ser Asn Ala Trp Ala Val Thr Asn His Phe Ser Ile
85           90           95

Trp Leu Ala Thr Ser Leu Ser Ile Phe Tyr Leu Leu Lys Ile Val Asn
100          105          110

Phe Ser Arg Leu Ile Phe His His Leu Lys Arg Lys Ala Lys Ser Val
115          120          125

Val Leu Val Ile Val Leu Gly Ser Leu Phe Phe Leu Val Cys His Leu
130          135          140

Val Met Lys His Thr Tyr Ile Asn Val Trp Thr Glu Glu Cys Glu Gly
145          150          155          160

Asn Val Thr Trp Lys Ile Lys Leu Arg Asn Ala Met His Leu Ser Asn
165          170          175

Leu Thr Val Ala Met Leu Ala Asn Leu Ile Pro Phe Thr Leu Thr Leu
180          185          190

Ile Ser Phe Leu Leu Leu Ile Tyr Ser Leu Cys Lys His Leu Lys Lys
195          200          205

Met Gln Leu His Gly Lys Gly Ser Gln Asp Pro Ser Thr Lys Ile His
210          215          220

Ile Lys Ala Leu Gln Thr Val Thr Ser Phe Leu Ile Leu Leu Ala Ile
225          230          235          240

Tyr Phe Leu Cys Leu Ile Ile Ser Phe Trp Asn Phe Lys Met Arg Pro
245          250          255

Lys Glu Ile Val Leu Met Leu Cys Gln Ala Phe Gly Ile Ile Tyr Pro
260          265          270

Ser Phe His Ser Phe Ile Leu Ile Trp Gly Asn Lys Thr Leu Lys Gln
275          280          285

Thr Phe Leu Ser Val Leu Trp Gln Val Thr Cys Trp Ala Lys Gly Gln
290          295          300

Asn Gln Ser Thr Pro
305

```

<210> SEQ ID NO 22

<211> LENGTH: 927

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

```

atgatgagtt ttctacacat tgttttttcc attctagtag tggttgcatt tattcttgga      60
aattttgcc aatggctttat agcactgata aatttcattg cctgggtcaa gagacaaaag      120
atctcctcag ctgatcaaat tattgtctgt ctggcagtct ccaaagtggg tttgctctgg      180
gtaatattat tacattggta ttcaactgtg ttgaatccaa cttcatctaa tttaaaagta      240

```

-continued

```

ataattttta tttctaatagc ctgggcagta accaatcatt tcagcatctg gcttgctact    300
agcctcagca tattttattt gctcaagatc gtcaatttct ccagacttat ttttcatcac    360
ttaaaaagga aggctaagag tgtagtcttg gtgatagtgt tggggctctt gttctttttg    420
gtttgtcacc ttgtgatgaa acacacgtat ataaatgtgt ggacagaaga atgtgaagga    480
aacgtaactt ggaagatcaa actgaggaat gcaatgcacc tttccaactt gactgtagcc    540
atgctagcaa acttgatacc attcactctg accctgatat cttttctgct gttaacttac    600
tctctgtgta aacatctgaa gaagatgcag ctccatggca aaggatctca agatcccagc    660
accaagatcc acataaaaagc tctgaaaact gtgacctcct tctcatattt acttgccatt    720
tactttctgt gtctaatacat atcgtttttg aattttaaga tgccgacaaa agaaattgtc    780
ttaatgcttt gccaaagcttt tggaatcata tatccatcat tccactcatt cattctgatt    840
tgggggaaca agacgctaaa gcagaccttt ctttcagttt tgtggcaggt gacttgctgg    900
gcaaaaggac agaaccagtc aactcca                                     927

```

<210> SEQ ID NO 23

<211> LENGTH: 299

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

```

Met Ile Thr Phe Leu Tyr Ile Phe Phe Ser Ile Leu Ile Met Val Leu
 1             5             10            15

Phe Val Leu Gly Asn Phe Ala Asn Gly Phe Ile Ala Leu Val Asn Phe
 20            25            30

Ile Asp Trp Val Lys Arg Lys Lys Ile Ser Ser Ala Asp Gln Ile Leu
 35            40            45

Thr Ala Leu Ala Val Ser Arg Ile Gly Leu Leu Trp Ala Leu Leu Leu
 50            55            60

Asn Trp Tyr Leu Thr Val Leu Asn Pro Ala Phe Tyr Ser Val Glu Leu
 65            70            75            80

Arg Ile Thr Ser Tyr Asn Ala Trp Val Val Thr Asn His Phe Ser Met
 85            90            95

Trp Leu Ala Ala Asn Leu Ser Ile Phe Tyr Leu Leu Lys Ile Ala Asn
100            105            110

Phe Ser Asn Leu Leu Phe Leu His Leu Lys Arg Arg Val Arg Ser Val
115            120            125

Ile Leu Val Ile Leu Leu Gly Thr Leu Ile Phe Leu Val Cys His Leu
130            135            140

Leu Val Ala Asn Met Asp Glu Ser Met Trp Ala Glu Glu Tyr Glu Gly
145            150            155            160

Asn Met Thr Gly Lys Met Lys Leu Arg Asn Thr Val His Leu Ser Tyr
165            170            175

Leu Thr Val Thr Thr Leu Trp Ser Phe Ile Pro Phe Thr Leu Ser Leu
180            185            190

Ile Ser Phe Leu Met Leu Ile Cys Ser Leu Tyr Lys His Leu Lys Lys
195            200            205

Met Gln Leu His Gly Glu Gly Ser Gln Asp Leu Ser Thr Lys Val His
210            215            220

Ile Lys Ala Leu Gln Thr Leu Ile Ser Phe Leu Leu Leu Cys Ala Ile

```


-continued

225	230	235	240
Phe Phe Leu Phe Leu Ile Val Ser Val Trp Ser Pro Arg Arg Leu Arg			
	245	250	255
Asn Asp Pro Val Val Met Val Ser Lys Ala Val Gly Asn Ile Tyr Leu			
	260	265	270
Ala Phe Asp Ser Phe Ile Leu Ile Trp Arg Thr Lys Lys Leu Lys His			
	275	280	285
Thr Phe Leu Leu Ile Leu Cys Gln Ile Arg Cys			
	290	295	

<210> SEQ ID NO 24
 <211> LENGTH: 897
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

```

atgataactt ttctatacat ttttttttca attctaataa tggttttatt tgttctcgga      60
aactttgcc aatggcttcat agcactggta aatttcattg actgggtgaa gagaaaaaag      120
atctcctcag ctgaccaaat tctcactgct ctggcgggtct ccagaattgg tttgctctgg      180
gcattattat taaattggta tttaactgtg ttgaatccag ctttttatag tgtagaatta      240
agaattactt cttataatgc ctgggttgta accaaccatt tcagcatgtg gcttgctgct      300
aacctcagca tattttattt gctcaagatt gccaatctct ccaaccttct ttttcttcat      360
ttaaagagga gagttaggag tgtcattctg gtgatactgt tggggacttt gatatttttg      420
gtttgtcatc ttcttgtggc aaacatggat gagagtatgt gggcagaaga atatgaagga      480
aacatgactg ggaagatgaa attgaggaat acagtacatc tttcatatth gactgtaact      540
accctatgga gcttcatacc ctttactctg tccctgatat cttttctgat gctaactctg      600
tctctgtata aacatctcaa gaagatgcag ctccatggag aaggatcgca agatctcagc      660
accaaggtcc acataaaagc ttgcaaaact ctgatctcct tcctcttggt atgtgccatt      720
ttctttctat tcctaactgt ttccggtttg agtcctagga ggctgcggaa tgacccagtt      780
gtcatgggta gcaaggctgt tggaaacata tatcttgcat tcgactcatt catcctaatt      840
tggaagaacca agaagctaaa acacaccttt cttttgattt tgtgtcagat taggtgc      897
  
```

<210> SEQ ID NO 25
 <211> LENGTH: 299
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Met Leu Glu Ser His Leu Ile Ile Tyr Phe Leu Leu Ala Val Ile Gln			
1	5	10	15
Phe Leu Leu Gly Ile Phe Thr Asn Gly Ile Ile Val Val Val Asn Gly			
	20	25	30
Ile Asp Leu Ile Lys His Arg Lys Met Ala Pro Leu Asp Leu Leu Leu			
	35	40	45
Ser Cys Leu Ala Val Ser Arg Ile Phe Leu Gln Leu Phe Ile Phe Tyr			
	50	55	60
Val Asn Val Ile Val Ile Phe Phe Ile Glu Phe Ile Met Cys Ser Ala			
	65	70	75
Asn Cys Ala Ile Leu Leu Phe Ile Asn Glu Leu Glu Leu Trp Leu Ala			

-continued

85								90					95				
Thr	Trp	Leu	Gly	Val	Phe	Tyr	Cys	Ala	Lys	Val	Ala	Ser	Val	Arg	His		
			100						105					110			
Pro	Leu	Phe	Ile	Trp	Leu	Lys	Met	Arg	Ile	Ser	Lys	Leu	Val	Pro	Trp		
		115					120					125					
Met	Ile	Leu	Gly	Ser	Leu	Leu	Tyr	Val	Ser	Met	Ile	Cys	Val	Phe	His		
		130					135					140					
Ser	Lys	Tyr	Ala	Gly	Phe	Met	Val	Pro	Tyr	Phe	Leu	Arg	Lys	Phe	Phe		
145						150					155				160		
Ser	Gln	Asn	Ala	Thr	Ile	Gln	Lys	Glu	Asp	Thr	Leu	Ala	Ile	Gln	Ile		
				165						170				175			
Phe	Ser	Phe	Val	Ala	Glu	Phe	Ser	Val	Pro	Leu	Leu	Ile	Phe	Leu	Phe		
			180					185					190				
Ala	Val	Leu	Leu	Leu	Ile	Phe	Ser	Leu	Gly	Arg	His	Thr	Arg	Gln	Met		
		195						200					205				
Arg	Asn	Thr	Val	Ala	Gly	Ser	Arg	Val	Pro	Gly	Arg	Gly	Ala	Pro	Ile		
	210						215				220						
Ser	Ala	Leu	Leu	Ser	Ile	Leu	Ser	Phe	Leu	Ile	Leu	Tyr	Phe	Ser	His		
225						230					235				240		
Cys	Met	Ile	Lys	Val	Phe	Leu	Ser	Ser	Leu	Lys	Phe	His	Ile	Arg	Arg		
			245							250				255			
Phe	Ile	Phe	Leu	Phe	Phe	Ile	Leu	Val	Ile	Gly	Ile	Tyr	Pro	Ser	Gly		
		260						265					270				
His	Ser	Leu	Ile	Leu	Ile	Leu	Gly	Asn	Pro	Lys	Leu	Lys	Gln	Asn	Ala		
		275					280						285				
Lys	Lys	Phe	Leu	Leu	His	Ser	Lys	Cys	Cys	Gln							
	290						295										

<210> SEQ ID NO 26

<211> LENGTH: 897

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

```

atgctagagt ctcacctcat tatctatctt cttcttgcag tgatacaatt tcttcttggg    60
atcttcacaa atggcatcat tgtggtggtg aatggcattg acttgatcaa gcacagaaaa    120
atggctccgc tggatctcct tctttcttgt ctggcagttt ctagaatttt tctgcagttg    180
ttcatcttct acgttaatgt gattgttato ttcttcatag aattcatcat gtgttctgcg    240
aattgtgcaa ttctcttatt tataaatgaa ttggaacttt ggcttgccac atggctcggc    300
gttttctatt gtgccaaggt tgcacagctc cgtcacccac tcttcatctg gttgaagatg    360
aggatatcca agctgggtccc atggatgata ctgggggtctc tgctatatgt atctatgatt    420
tgtgttttcc atagcaaata tgcagggttt atggtcccat acttctaag gaaatttttc    480
tcccaaaatg ccacaattca aaaagaagat aactggcta tacagatttt ctcttttgtt    540
gctgagttct cagtgcattt gcttatcttc ctttttgctg ttttgctctt gattttctct    600
ctggggaggc acaccgggca aatgagaaac acagtggccg gcagcagggt tcttggcagg    660
ggcgacccca tcagcgcgtt gctgtctatc ctgtccttcc tgatcctcta cttctcccac    720
tgcatgataa aagtttttct ctcttctcta aagtttcaca tcagaaggtt catctttctg    780
ttcttcatcc ttgtgattgg tatataccct tctggacact ctctcatctt aatttttaga    840

```

-continued

aatcctaaat tgaaacaaaa tgcaaaaaag ttctctctcc acagtaagtg ctgtcag 897

<210> SEQ ID NO 27

<211> LENGTH: 299

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

Met Leu Arg Leu Phe Tyr Phe Ser Ala Ile Ile Ala Ser Val Ile Leu
1 5 10 15Asn Phe Val Gly Ile Ile Met Asn Leu Phe Ile Thr Val Val Asn Cys
20 25 30Lys Thr Trp Val Lys Ser His Arg Ile Ser Ser Ser Asp Arg Ile Leu
35 40 45Phe Ser Leu Gly Ile Thr Arg Phe Leu Met Leu Gly Leu Phe Leu Val
50 55 60Asn Thr Ile Tyr Phe Val Ser Ser Asn Thr Glu Arg Ser Val Tyr Leu
65 70 75 80Ser Ala Phe Phe Val Leu Cys Phe Met Phe Leu Asp Ser Ser Ser Val
85 90 95Trp Phe Val Thr Leu Leu Asn Ile Leu Tyr Cys Val Lys Ile Thr Asn
100 105 110Phe Gln His Ser Val Phe Leu Leu Leu Lys Arg Asn Ile Ser Pro Lys
115 120 125Ile Pro Arg Leu Leu Leu Ala Cys Val Leu Ile Ser Ala Phe Thr Thr
130 135 140Cys Leu Tyr Ile Thr Leu Ser Gln Ala Ser Pro Phe Pro Glu Leu Val
145 150 155 160Thr Thr Arg Asn Asn Thr Ser Phe Asn Ile Ser Glu Gly Ile Leu Ser
165 170 175Leu Val Val Ser Leu Val Leu Ser Ser Ser Leu Gln Phe Ile Ile Asn
180 185 190Val Thr Ser Ala Ser Leu Leu Ile His Ser Leu Arg Arg His Ile Gln
195 200 205Lys Met Gln Lys Asn Ala Thr Gly Phe Trp Asn Pro Gln Thr Glu Ala
210 215 220His Val Gly Ala Met Lys Leu Met Val Tyr Phe Leu Ile Leu Tyr Ile
225 230 235 240Pro Tyr Ser Val Ala Thr Leu Val Gln Tyr Leu Pro Phe Tyr Ala Gly
245 250 255Met Asp Met Gly Thr Lys Ser Ile Cys Leu Ile Phe Ala Thr Leu Tyr
260 265 270Ser Pro Gly His Ser Val Leu Ile Ile Ile Thr His Pro Lys Leu Lys
275 280 285Thr Thr Ala Lys Lys Ile Leu Cys Phe Lys Lys
290 295

<210> SEQ ID NO 28

<211> LENGTH: 897

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

-continued

```

atgcttcggt tattctatatt ctctgctatt attgcctcag ttattttaaa tttttagga      60
atcattatga atctgtttat tacagtggtc aattgcaaaa ctgggtcaa aagccataga      120
atctcctctt ctgataggat tctgttcagc ctgggcatca ccaggtttct tatgctggga      180
ctattttctgg tgaacaccat ctacttcgtc tcttcaaata cggaaaggtc agtctacctg      240
tctgcttttt ttgtgttttg tttcatgttt ttggactcga gcagtgtctg gtttgtgacc      300
ttgtcgaata tcttgtactg tgtgaagatt actaacttcc aacctcagt gtttctcctg      360
ctgaagcgga atatctcccc aaagatcccc aggctgctgc tggcctgtgt gctgatttct      420
gctttcacca ctgacctgta catcacgctt agccaggcat caccctttcc tgaacttgtg      480
actacgagaa ataacacatc atttaatatc agtgagggca tcttgtcttt agtggtttct      540
ttggtcttga gctcatctct ccagttcatc attaatgtga cttctgcttc ctgctaata      600
cactccttga ggagacatat acagaagatg cagaaaaatg ccactggttt ctggaatccc      660
cagacggaag ctcatgtagg tgctatgaag ctgatggtct atttctcat cctctacatt      720
ccatattcag ttgtaccct ggtccagtat ctcccctttt atgcagggat ggatatgggg      780
accaaatcca ttgtctgat ttttgccacc ctttactctc caggacattc tgttctcatt      840
attatcacac atcctaaact gaaaacaaca gcaaagaaga ttctttgttt caaaaaa      897

```

<210> SEQ ID NO 29

<211> LENGTH: 299

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

```

Met Leu Ser Ala Gly Leu Gly Leu Leu Met Leu Val Ala Val Val Glu
 1             5             10            15

Phe Leu Ile Gly Leu Ile Gly Asn Gly Ser Leu Val Val Trp Ser Phe
 20            25            30

Arg Glu Trp Ile Arg Lys Phe Asn Trp Ser Ser Tyr Asn Leu Ile Ile
 35            40            45

Leu Gly Leu Ala Gly Cys Arg Phe Leu Leu Gln Trp Leu Ile Ile Leu
 50            55            60

Asp Leu Ser Leu Phe Pro Leu Phe Gln Ser Ser Arg Trp Leu Arg Tyr
 65            70            75            80

Leu Ser Ile Phe Trp Val Leu Val Ser Gln Ala Ser Leu Trp Phe Ala
 85            90            95

Thr Phe Leu Ser Val Phe Tyr Cys Lys Lys Ile Thr Thr Phe Asp Arg
100           105           110

Pro Ala Tyr Leu Trp Leu Lys Gln Arg Ala Tyr Asn Leu Ser Leu Trp
115           120           125

Cys Leu Leu Gly Tyr Phe Ile Ile Asn Leu Leu Leu Thr Val Gln Ile
130           135           140

Gly Leu Thr Phe Tyr His Pro Pro Gln Gly Asn Ser Ser Ile Arg Tyr
145           150           155           160

Pro Phe Glu Ser Trp Gln Tyr Leu Tyr Ala Phe Gln Leu Asn Ser Gly
165           170           175

Ser Tyr Leu Pro Leu Val Val Phe Leu Val Ser Ser Gly Met Leu Ile
180           185           190

Val Ser Leu Tyr Thr His His Lys Lys Met Lys Val His Ser Ala Gly
195           200           205

```

-continued

Arg Arg Asp Val Arg Ala Lys Ala His Ile Thr Ala Leu Lys Ser Leu
 210 215 220

Gly Cys Phe Leu Leu Leu His Leu Val Tyr Ile Met Ala Ser Pro Phe
 225 230 235 240

Ser Ile Thr Ser Lys Thr Tyr Pro Pro Asp Leu Thr Ser Val Phe Ile
 245 250 255

Trp Glu Thr Leu Met Ala Ala Tyr Pro Ser Leu His Ser Leu Ile Leu
 260 265 270

Ile Met Gly Ile Pro Arg Val Lys Gln Thr Cys Gln Lys Ile Leu Trp
 275 280 285

Lys Thr Val Cys Ala Arg Arg Cys Trp Gly Pro
 290 295

<210> SEQ ID NO 30
 <211> LENGTH: 897
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

```

atgctgagcg ctggcctagg actgctgatg ctggtggcag tggttgaatt tctcatcggt    60
ttaattggaa atggaagcct ggtggtctgg agttttagag aatggatcag aaaattcaac    120
tggtcctcat ataacctcat taccctgggc ctggtggct gccgatttct cctgcagtgg    180
ctgatcattt tggacttaag cttgtttcca cttttccaga gcagccgttg gcttcgctat    240
cttagtatct tctgggtcct ggtaagccag gccagcttat ggtttgccac cttcctcagt    300
gtcttctatt gcaagaagat cagcaccttc gatcgcccg cctacttggt gctgaagcag    360
agggcctata acctgagtct ctggtgcctt ctgggtact ttataatcaa ttgttactt    420
acagtccaaa ttggettaac attctatcat cctccccaag gaaacagcag cattcggtat    480
ccctttgaaa gctggcagta cctgtatgca tttcagtcga attcaggaag ttatttgcct    540
ttagtggtgt ttcttgtttc ctctgggatg ctgattgtct ctttgatac acaccacaag    600
aagatgaagg tcatttcagc tggtaggagg gatgtccggg ccaaggctca catcactgcg    660
ctgaagtctt tgggtgctt cctcttactt cacctggttt atatcatggc cagcccttc    720
tccatcacct ccaagactta tcctcctgat ctcaccagtg tcttcactct ggagacactc    780
atggcagcct atccttctct tcattctctc atattgatca tggggattcc tagggtgaag    840
cagacttgtc agaagatcct gtggaagacg gtgtgtgctc ggagatgctg gggccca    897

```

<210> SEQ ID NO 31
 <211> LENGTH: 318
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

Met Ala Asp Lys Val Gln Thr Thr Leu Leu Phe Leu Ala Val Gly Glu
 1 5 10 15

Phe Ser Val Gly Ile Leu Gly Asn Ala Phe Ile Gly Leu Val Asn Cys
 20 25 30

Met Asp Trp Val Lys Lys Arg Lys Ile Ala Ser Ile Asp Leu Ile Leu
 35 40 45

Thr Ser Leu Ala Ile Ser Arg Ile Cys Leu Leu Cys Val Ile Leu Leu
 50 55 60

-continued

Asp Cys Phe Ile Leu Val Leu Tyr Pro Asp Val Tyr Ala Thr Gly Lys
 65 70 75 80
 Glu Met Arg Ile Ile Asp Phe Phe Trp Thr Leu Thr Asn His Leu Ser
 85 90 95
 Ile Trp Phe Ala Thr Cys Leu Ser Ile Tyr Tyr Phe Phe Lys Ile Gly
 100 105 110
 Asn Phe Phe His Pro Leu Phe Leu Trp Met Lys Trp Arg Ile Asp Arg
 115 120 125
 Val Ile Ser Trp Ile Leu Leu Gly Cys Val Val Leu Ser Val Phe Ile
 130 135 140
 Ser Leu Pro Ala Thr Glu Asn Leu Asn Ala Asp Phe Arg Phe Cys Val
 145 150 155 160
 Lys Ala Lys Arg Lys Thr Asn Leu Thr Trp Ser Cys Arg Val Asn Lys
 165 170 175
 Thr Gln His Ala Ser Thr Lys Leu Phe Leu Asn Leu Ala Thr Leu Leu
 180 185 190
 Pro Phe Cys Val Cys Leu Met Ser Phe Phe Leu Leu Ile Leu Ser Leu
 195 200 205
 Arg Arg His Ile Arg Arg Met Gln Leu Ser Ala Thr Gly Cys Arg Asp
 210 215 220
 Pro Ser Thr Glu Ala His Val Arg Ala Leu Lys Ala Val Ile Ser Phe
 225 230 235 240
 Leu Leu Leu Phe Ile Ala Tyr Tyr Leu Ser Phe Leu Ile Ala Thr Ser
 245 250 255
 Ser Tyr Phe Met Pro Glu Thr Glu Leu Ala Val Ile Phe Gly Glu Ser
 260 265 270
 Ile Ala Leu Ile Tyr Pro Ser Ser His Ser Phe Ile Leu Ile Leu Gly
 275 280 285
 Asn Asn Lys Leu Arg His Ala Ser Leu Lys Val Ile Trp Lys Val Met
 290 295 300
 Ser Ile Leu Lys Gly Arg Lys Phe Gln Gln His Lys Gln Ile
 305 310 315

<210> SEQ ID NO 32

<211> LENGTH: 954

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

atggcagata aagtcagac tactttattg ttcttagcag ttggagagtt ttcagtggg 60
 atcttaggga atgcattcat tggattggta aactgcatgg actgggtcaa gaagaggaaa 120
 attgcctcca ttgatttaat cctcacaagt ctggccatat ccagaatttg tctattgtgc 180
 gtaatactat tagattgttt tatattgttg ctatatccag atgtctatgc cactggtaaa 240
 gaaatgagaa tcattgactt cttctggaca ctaaccaatc atttaagtat ctgggttgca 300
 acctgctca gcatttacta tttcttcaag ataggtaatt tctttcacc acttttctc 360
 tggatgaagt ggagaattga cagggtgatt tcttggtatc tactgggttg cgtggttctc 420
 tctgtgttta ttagccttcc agccactgag aatttgaacg ctgatttcag gttttgtgtg 480
 aaggcaaaga ggaaacaaa cttaacttgg agttgcagag taaataaac tcaacatgct 540
 tctaccaagt tatttctcaa cctggcaacg ctgtccctc tttgtgtgtg cctaagtcc 600

-continued

```

ttttctctct tgatcctctc cctgcccaga catatcaggc gaatgcagct cagtgccaca    660
gggtgcagag accccagcac agaagcccat gtgagagccc tgaaagctgt catttccttc    720
cttctctctc ttattgecta ctatttgctc ttctcattg ccacctccag ctactttatg    780
ccagagacgg aattagctgt gatttttggt gattccatag ctctaactta cccctcaagt    840
cattcattta tcctaatact ggggaacaat aaattaagac atgcattctc aaaggtgatt    900
tggaagtaa  tgtctattct aaaaggaaga aaattccaac aacataaaca aatc        954

```

<210> SEQ ID NO 33

<211> LENGTH: 309

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

```

Met Phe Ser Pro Ala Asp Asn Ile Phe Ile Ile Leu Ile Thr Gly Glu
1           5           10          15

Phe Ile Leu Gly Ile Leu Gly Asn Gly Tyr Ile Ala Leu Val Asn Trp
20          25          30

Ile Asp Trp Ile Lys Lys Lys Lys Ile Ser Thr Val Asp Tyr Ile Leu
35          40          45

Thr Asn Leu Val Ile Ala Arg Ile Cys Leu Ile Ser Val Met Val Val
50          55          60

Asn Gly Ile Val Ile Val Leu Asn Pro Asp Val Tyr Thr Lys Asn Lys
65          70          75          80

Gln Gln Ile Val Ile Phe Thr Phe Trp Thr Phe Ala Asn Tyr Leu Asn
85          90          95

Met Trp Ile Thr Thr Cys Leu Asn Val Phe Tyr Phe Leu Lys Ile Ala
100         105         110

Ser Ser Ser His Pro Leu Phe Leu Trp Leu Lys Trp Lys Ile Asp Met
115         120         125

Val Val His Trp Ile Leu Leu Gly Cys Phe Ala Ile Ser Leu Leu Val
130         135         140

Ser Leu Ile Ala Ala Ile Val Leu Ser Cys Asp Tyr Arg Phe His Ala
145         150         155         160

Ile Ala Lys His Lys Arg Asn Ile Thr Glu Met Phe His Val Ser Lys
165         170         175

Ile Pro Tyr Phe Glu Pro Leu Thr Leu Phe Asn Leu Phe Ala Ile Val
180         185         190

Pro Phe Ile Val Ser Leu Ile Ser Phe Phe Leu Leu Val Arg Ser Leu
195         200         205

Trp Arg His Thr Lys Gln Ile Lys Leu Tyr Ala Thr Gly Ser Arg Asp
210         215         220

Pro Ser Thr Glu Val His Val Arg Ala Ile Lys Thr Met Thr Ser Phe
225         230         235         240

Ile Phe Phe Phe Phe Leu Tyr Tyr Ile Ser Ser Ile Leu Met Thr Phe
245         250         255

Ser Tyr Leu Met Thr Lys Tyr Lys Leu Ala Val Glu Phe Gly Glu Ile
260         265         270

Ala Ala Ile Leu Tyr Pro Leu Gly His Ser Leu Ile Leu Ile Val Leu
275         280         285

Asn Asn Lys Leu Arg Gln Thr Phe Val Arg Met Leu Thr Cys Arg Lys

```

-continued

290	295	300	
Ile Ala Cys Met Ile			
305			
<210> SEQ ID NO 34			
<211> LENGTH: 927			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 34			
atgttcagtc ctgcagataa catctttata atcctaataa ctggagaatt catactagga			60
atattgggga atggatacat tgcactagtc aactggattg actggattaa gaagaaaaag			120
atttccacag ttgactacat ccttaccaat ttagttatcg ccagaatttg tttgatcagt			180
gtaatggttg taaatggcat tgtaatagta ctgaaccag atgtttatac aaaaaataaa			240
caacagatag tcattttttac cttctggaca ttgccaact acttaaatat gtggattacc			300
acctgcctta atgtcttcta tttctgaag atagccagtt cctctcatcc actttttctc			360
tggctgaagt ggaaaattga tatggtggtg cactggatcc tgctgggatg ctttgccatt			420
tccttggttg tcagccttat agcagcaata gtactgagtt gtgattatag gtttcatgca			480
attgccaaac ataaaagaaa cattactgaa atgttccatg tgagtaaaat accatacttt			540
gaacccttaa ctctctttaa cctgtttgca attgtccat ttattgtgtc actgatatca			600
tttttccttt tagtaagatc tttatggaga cataccaagc aaataaaaact ctatgctacc			660
ggcagtagag accccagcac agaagttcat gtgagagcca ttaaaactat gacttcattt			720
atcttctttt ttttctata ctatatttct tctattttga tgacctttag ctatcttatg			780
acaaaataca agttagctgt ggagtttga gagattgcag caattctcta cccctgggt			840
cactcactta ttttaattgt tttaaataat aaactgaggc agacatttgt cagaatgctg			900
acatgtagaa aaattgcctg catgata			927

<210> SEQ ID NO 35
 <211> LENGTH: 312
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

Met Pro Ser Ala Ile Glu Ala Ile Tyr Ile Ile Leu Ile Ala Gly Glu			
1 5 10 15			
Leu Thr Ile Gly Ile Trp Gly Asn Gly Phe Ile Val Leu Val Asn Cys			
20 25 30			
Ile Asp Trp Leu Lys Arg Arg Asp Ile Ser Leu Ile Asp Ile Ile Leu			
35 40 45			
Ile Ser Leu Ala Ile Ser Arg Ile Cys Leu Leu Cys Val Ile Ser Leu			
50 55 60			
Asp Gly Phe Phe Met Leu Leu Phe Pro Gly Thr Tyr Gly Asn Ser Val			
65 70 75 80			
Leu Val Ser Ile Val Asn Val Val Trp Thr Phe Ala Asn Asn Ser Ser			
85 90 95			
Leu Trp Phe Thr Ser Cys Leu Ser Ile Phe Tyr Leu Leu Lys Ile Ala			
100 105 110			
Asn Ile Ser His Pro Phe Phe Phe Trp Leu Lys Leu Lys Ile Asn Lys			
115 120 125			

-continued

Val Met Leu Ala Ile Leu Leu Gly Ser Phe Leu Ile Ser Leu Ile Ile
130 135 140

Ser Val Pro Lys Asn Asp Asp Met Trp Tyr His Leu Phe Lys Val Ser
145 150 155 160

His Glu Glu Asn Ile Thr Trp Lys Phe Lys Val Ser Lys Ile Pro Gly
165 170 175

Thr Phe Lys Gln Leu Thr Leu Asn Leu Gly Val Met Val Pro Phe Ile
180 185 190

Leu Cys Leu Ile Ser Phe Phe Leu Leu Leu Phe Ser Leu Val Arg His
195 200 205

Thr Lys Gln Ile Arg Leu His Ala Thr Gly Phe Arg Asp Pro Ser Thr
210 215 220

Glu Ala His Met Arg Ala Ile Lys Ala Val Ile Ile Phe Leu Leu Leu
225 230 235 240

Leu Ile Val Tyr Tyr Pro Val Phe Leu Val Met Thr Ser Ser Ala Leu
245 250 255

Ile Pro Gln Gly Lys Leu Val Leu Met Ile Gly Asp Ile Val Thr Val
260 265 270

Ile Phe Pro Ser Ser His Ser Phe Ile Leu Ile Met Gly Asn Ser Lys
275 280 285

Leu Arg Glu Ala Phe Leu Lys Met Leu Arg Phe Val Lys Cys Phe Leu
290 295 300

Arg Arg Arg Lys Pro Phe Val Pro
305 310

<210> SEQ ID NO 36

<211> LENGTH: 936

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

```

atgccaaagt caatagaggc aatatatatt attttaattg ctggtgaatt gaccataggg      60
atttggggaa atggattcat tgtactagtt aactgcattg actggctcaa aagaagagat      120
atttccttga ttgacatcat cctgatcagc ttggccatct ccagaatctg tctgctgtgt      180
gtaatatcat tagatggctt ctttatgctg ctctttccag gtacatatgg caatagcgtg      240
ctagtaagca ttgtgaatgt tgtctggaca ttgccaata attcaagtct ctggtttact      300
tcttgctca gtatctttcta ttactcaag atagccaata tatcgacccc atttttcttc      360
tggtgaagc taaagatcaa caaggtcattg cttgcgattc ttctggggtc ctttcttatt      420
tctttaatta ttagtggttc aaagaatgat gatatgtggt atcacctttt caaagtcagt      480
catgaagaaa acattacttg gaaattcaaa gtgagtaaaa ttccaggtac tttcaaacag      540
ttaacctga acctgggggt gatggttccc tttatccttt gctgatctc atttttcttg      600
ttacttttct ccctagttag acacaccaag cagattcgac tgcattgctac agggttcaga      660
gaccccagta cagaggccca catgagggcc ataaaggcag tgatcatctt tctgctcttc      720
ctcatcgtgt actaccagct ctttcttggt atgaccteta gcgctctgat tcttcaggga      780
aaattagtgt tgatgattgg tgacatagta actgtcattt tcccatcaag ccattcattc      840
attctaatta tgggaatatg caagttgagg gaagcttttc tgaagatgtt aagatttgtg      900
aagtgtttcc ttagaagaag aaagcctttt gttcca                                936

```

-continued

<210> SEQ ID NO 37
<211> LENGTH: 307
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 37
Met Leu Arg Val Val Glu Gly Ile Phe Ile Phe Val Val Val Ser Glu
1 5 10 15
Ser Val Phe Gly Val Leu Gly Asn Gly Phe Ile Gly Leu Val Asn Cys
20 25 30
Ile Asp Cys Ala Lys Asn Lys Leu Ser Thr Ile Gly Phe Ile Leu Thr
35 40 45
Gly Leu Ala Ile Ser Arg Ile Phe Leu Ile Trp Ile Ile Ile Thr Asp
50 55 60
Gly Phe Ile Gln Ile Phe Ser Pro Asn Ile Tyr Ala Ser Gly Asn Leu
65 70 75 80
Ile Glu Tyr Ile Ser Tyr Phe Trp Val Ile Gly Asn Gln Ser Ser Met
85 90 95
Trp Phe Ala Thr Ser Leu Ser Ile Phe Tyr Phe Leu Lys Ile Ala Asn
100 105 110
Phe Ser Asn Tyr Ile Phe Leu Trp Leu Lys Ser Arg Thr Asn Met Val
115 120 125
Leu Pro Phe Met Ile Val Phe Leu Leu Ile Ser Ser Leu Leu Asn Phe
130 135 140
Ala Tyr Ile Ala Lys Ile Leu Asn Asp Tyr Lys Met Lys Asn Asp Thr
145 150 155 160
Val Trp Asp Leu Asn Met Tyr Lys Ser Glu Tyr Phe Ile Lys Gln Ile
165 170 175
Leu Leu Asn Leu Gly Val Ile Phe Phe Phe Thr Leu Ser Leu Ile Thr
180 185 190
Cys Ile Phe Leu Ile Ile Ser Leu Trp Arg His Asn Arg Gln Met Gln
195 200 205
Ser Asn Val Thr Gly Leu Arg Asp Ser Asn Thr Glu Ala His Val Lys
210 215 220
Ala Met Lys Val Leu Ile Ser Phe Ile Ile Leu Phe Ile Leu Tyr Phe
225 230 235 240
Ile Gly Met Ala Ile Glu Ile Ser Cys Phe Thr Val Arg Glu Asn Lys
245 250 255
Leu Leu Leu Met Phe Gly Met Thr Thr Thr Ala Ile Tyr Pro Trp Gly
260 265 270
His Ser Phe Ile Leu Ile Leu Gly Asn Ser Lys Leu Lys Gln Ala Ser
275 280 285
Leu Arg Val Leu Gln Gln Leu Lys Cys Cys Glu Lys Arg Lys Asn Leu
290 295 300
Arg Val Thr
305

<210> SEQ ID NO 38
<211> LENGTH: 921
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

-continued

```

atgctacgtg tagtgaagg catcttcatt tttgtttag ttagtgagtc agtgtttggg    60
gttttgggga atggatttat tggacttgta aactgcattg actgtgcaa gaataagtta    120
tctacgattg gctttattct caccggctta gctatttcaa gaatttttct gatatggata    180
ataattacag atggatttat acagatatct tctccaaata tatatgcctc cggtaaccta    240
attgaatata ttagttactt ttgggtaatt ggtaatcaat caagtatgtg gtttgccacc    300
agcctcagca tcttctattt cctgaagata gcaaattttt ccaactacat atttctctgg    360
ttgaagagca gaacaaatat ggttcttccc ttcgatagat tattcttact tatttcacg    420
ttacttaatt ttgcatacat tgogaagatt cttaatgatt ataaaatgaa gaatgacaca    480
gtctgggagc tcaacatgta taaaagttaa tactttatta aacagatttt gctaaatctg    540
ggagtcattt tcttctttac actatcccta attacatgta tttttttaat catttccctt    600
tggagacaca acaggcagat gcaatcgaat gtgacaggat tgagagactc caacacagaa    660
gctcatgtga aggcaatgaa agttttgata tctttcatca tctcttttat cttgtatttt    720
ataggcatgg ccatagaaat atcatgtttt actgtgagag aaaacaaact gctgcttatg    780
tttggaaatg caaccacagc catctatccc tgggggcact catttatctt aattctagga    840
aacagcaagc taaagcaagc ctctttgagg gtactgcagc aattgaagtg ctgtgagaaa    900
aggaaaaatc tcagagtcac a                                           921

```

<210> SEQ ID NO 39

<211> LENGTH: 303

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

```

Met Glu Ser Ala Leu Pro Ser Ile Phe Thr Leu Val Ile Ile Ala Glu
1           5           10          15

Phe Ile Ile Gly Asn Leu Ser Asn Gly Phe Ile Val Leu Ile Asn Cys
20        25        30

Ile Asp Trp Val Ser Lys Arg Glu Leu Ser Ser Val Asp Lys Leu Leu
35        40        45

Ile Ile Leu Ala Ile Ser Arg Ile Gly Leu Ile Trp Glu Ile Leu Val
50        55        60

Ser Trp Phe Leu Ala Leu His Tyr Leu Ala Ile Phe Val Ser Gly Thr
65        70        75        80

Gly Leu Arg Ile Met Ile Phe Ser Trp Ile Val Ser Asn His Phe Asn
85        90        95

Leu Trp Leu Ala Thr Ile Phe Ser Ile Phe Tyr Leu Leu Lys Ile Ala
100       105       110

Ser Phe Ser Ser Pro Ala Phe Leu Tyr Leu Lys Trp Arg Val Asn Lys
115       120       125

Val Ile Leu Met Ile Leu Leu Gly Thr Leu Val Phe Leu Phe Leu Asn
130       135       140

Leu Ile Gln Ile Asn Met His Ile Lys Asp Trp Leu Asp Arg Tyr Glu
145       150       155       160

Arg Asn Thr Thr Trp Asn Phe Ser Met Ser Asp Phe Glu Thr Phe Ser
165       170       175

Val Ser Val Lys Phe Thr Met Thr Met Phe Ser Leu Thr Pro Phe Thr
180       185       190

```

-continued

Val Ala Phe Ile Ser Phe Leu Leu Leu Ile Phe Ser Leu Gln Lys His
 195 200 205

Leu Gln Lys Met Gln Leu Asn Tyr Lys Gly His Arg Asp Pro Arg Thr
 210 215 220

Lys Val His Thr Asn Ala Leu Lys Ile Val Ile Ser Phe Leu Leu Phe
 225 230 235 240

Tyr Ala Ser Phe Phe Leu Cys Val Leu Ile Ser Trp Ile Ser Glu Leu
 245 250 255

Tyr Gln Ser Thr Val Ile Tyr Met Leu Cys Glu Thr Ile Gly Val Phe
 260 265 270

Ser Pro Ser Ser His Ser Phe Leu Leu Ile Leu Gly Asn Ala Lys Leu
 275 280 285

Arg Gln Ala Phe Leu Leu Val Ala Ala Lys Val Trp Ala Lys Arg
 290 295 300

<210> SEQ ID NO 40
 <211> LENGTH: 909
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

```

atggaaagtg cctgccgag tatcttcact cttgtaataa ttgcagaatt cataattggg    60
aatttgagca atggatttat agtactgatc aactgcattg actgggtcag taaaagagag    120
ctgtcctcag tcgataaaact cctcattatc ttggcaatct ccagaattgg gctgatctgg    180
gaaatattag taagtgtggt ttagctctg cattatctag ccatatttgt gtctggaaca    240
ggattaagaa ttatgatttt tagctggata gtttctaata acttcaatct ctggcttgct    300
acaatcttca gcactcttcta ttgctcaaaa atagcgagtt tctctagccc tgcttttctc    360
tatttgaagt ggagagtaaa caaagtgatt ctgatgatac tgctaggaac cttggtcttc    420
ttatttttaa atctgataca aataaacatg catataaaag actggctgga ccgatatgaa    480
agaaacacaa cttggaattt cagtatgagt gactttgaaa cattttcagt gtcggtcaaa    540
ttcactatga ctatgttcag tctaacacca tttactgtgg ccttcatctc ttttctcctg    600
ttaattttct cctgcagaaa acatctccag aaaatgcaac tcaattacaa aggacacaga    660
gaccccgagg ccaaggtcca tacaaatgcc ttgaaaattg tgatctcatt ccttttattc    720
tatgctagtt tctttctatg tgttctcata tcatggattt ctgagctgta tcagagcaca    780
gtgatctaca tgctttgtga gacgattgga gtcttctctc cttcaagcca ctcttttctt    840
ctgattctag gaaacgctaa gttaagacag gcctttcttt tgggtggcagc taaggtatgg    900
gctaaacga                                     909
  
```

<210> SEQ ID NO 41
 <211> LENGTH: 317
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

Met Gly Gly Val Ile Lys Ser Ile Phe Thr Phe Val Leu Ile Val Glu
 1 5 10 15

Phe Ile Ile Gly Asn Leu Gly Asn Ser Phe Ile Ala Leu Val Asn Cys
 20 25 30

-continued

Ile	Asp	Trp	Val	Lys	Gly	Arg	Lys	Ile	Ser	Ser	Val	Asp	Arg	Ile	Leu
	35						40					45			
Thr	Ala	Leu	Ala	Ile	Ser	Arg	Ile	Ser	Leu	Val	Trp	Leu	Ile	Phe	Gly
	50					55					60				
Ser	Trp	Cys	Val	Ser	Val	Phe	Phe	Pro	Ala	Leu	Phe	Ala	Thr	Glu	Lys
65					70					75				80	
Met	Phe	Arg	Met	Leu	Thr	Asn	Ile	Trp	Thr	Val	Ile	Asn	His	Phe	Ser
				85					90					95	
Val	Trp	Leu	Ala	Thr	Gly	Leu	Gly	Thr	Phe	Tyr	Phe	Leu	Lys	Ile	Ala
			100					105					110		
Asn	Phe	Ser	Asn	Ser	Ile	Phe	Leu	Tyr	Leu	Lys	Trp	Arg	Val	Lys	Lys
		115					120					125			
Val	Val	Leu	Val	Leu	Leu	Leu	Val	Thr	Ser	Val	Phe	Leu	Phe	Leu	Asn
	130					135					140				
Ile	Ala	Leu	Ile	Asn	Ile	His	Ile	Asn	Ala	Ser	Ile	Asn	Gly	Tyr	Arg
145					150					155					160
Arg	Asn	Lys	Thr	Cys	Ser	Ser	Asp	Ser	Ser	Asn	Phe	Thr	Arg	Phe	Ser
				165					170					175	
Ser	Leu	Ile	Val	Leu	Thr	Ser	Thr	Val	Phe	Ile	Phe	Ile	Pro	Phe	Thr
			180					185					190		
Leu	Ser	Leu	Ala	Met	Phe	Leu	Leu	Leu	Ile	Phe	Ser	Met	Trp	Lys	His
		195					200					205			
Arg	Lys	Lys	Met	Gln	His	Thr	Val	Lys	Ile	Ser	Gly	Asp	Ala	Ser	Thr
	210					215						220			
Lys	Ala	His	Arg	Gly	Val	Lys	Ser	Val	Ile	Thr	Phe	Phe	Leu	Leu	Tyr
225					230					235					240
Ala	Ile	Phe	Ser	Leu	Ser	Phe	Phe	Ile	Ser	Val	Trp	Thr	Ser	Glu	Arg
				245						250				255	
Leu	Glu	Glu	Asn	Leu	Ile	Ile	Leu	Ser	Gln	Val	Met	Gly	Met	Ala	Tyr
			260					265					270		
Pro	Ser	Cys	His	Ser	Cys	Val	Leu	Ile	Leu	Gly	Asn	Lys	Lys	Leu	Arg
		275					280					285			
Gln	Ala	Ser	Leu	Ser	Val	Leu	Leu	Trp	Leu	Arg	Tyr	Met	Phe	Lys	Asp
	290					295					300				
Gly	Glu	Pro	Ser	Gly	His	Lys	Glu	Phe	Arg	Glu	Ser	Ser			
305					310					315					

<210> SEQ ID NO 42

<211> LENGTH: 951

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

atgggtggtg tcataaagag catatttaca ttctgttttaa ttgtggaatt tataattgga	60
aatttaggaa atagtttcat agcactggtg aactgtattg actgggtcaa ggaagaaaag	120
atctcttcgg ttgatcggat cctcactgct ttggcaatct ctgaattag cctggtttgg	180
ttaatattcg gaagctggtg tgtgtctgtg tttttcccag ctttatttgc cactgaaaaa	240
atgttcagaa tgcttactaa tatctggaca gtgatcaatc attttagtgt ctggttagct	300
acaggcctcg gtacttttta tttttcaag atagccaatt tttctaactc tatttttctc	360
tacctaaggt ggagaggtaa aaaggtggtt ttggtgctgc ttcttgtgac ttcggtcttc	420

-continued

```

ttgtttttaa atattgcact gataaacatc catataaatg ccagtatcaa tggatacaga    480
agaacaaga cttgcagttc tgattcaagt aactttacac gattttccag tcttattgta    540
ttaaccagca ctgtgttcat tttcataccc tttactttgt cctgggaat gtttcttctc    600
ctcatcttct ccatgtggaa acatcgcaag aagatgcagc acactgtcaa aatatccgga    660
gacgccagca ccaaagccca cagaggagtt aaaagtgtga tcactttctt cctactctat    720
gccattttct ctctgtcttt tttcatatca gtttgacct ctgaaagggt ggaggaaaat    780
ctaattattc tttcccaggt gatgggaatg gcttatcctt catgtcactc atgtgttctg    840
attcttggaa acaagaagct gagacaggcc tctctgtcag tgctactgtg gctgaggtac    900
atgttcaaag atggggagcc ctcaggtcac aaagaattta gagaatcatc t          951

```

<210> SEQ ID NO 43

<211> LENGTH: 291

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

```

Met Ile Pro Ile Gln Leu Thr Val Phe Phe Met Ile Ile Tyr Val Leu
 1             5             10             15

Glu Ser Leu Thr Ile Ile Val Gln Ser Ser Leu Ile Val Ala Val Leu
      20             25             30

Gly Arg Glu Trp Leu Gln Val Arg Arg Leu Met Pro Val Asp Met Ile
      35             40             45

Leu Ile Ser Leu Gly Ile Ser Arg Phe Cys Leu Gln Trp Ala Ser Met
 50             55             60

Leu Asn Asn Phe Cys Ser Tyr Phe Asn Leu Asn Tyr Val Leu Cys Asn
65             70             75             80

Leu Thr Ile Thr Trp Glu Phe Phe Asn Ile Leu Thr Phe Trp Leu Asn
      85             90             95

Ser Leu Leu Thr Val Phe Tyr Cys Ile Lys Val Ser Ser Phe Thr His
100            105            110

His Ile Phe Leu Trp Leu Arg Trp Arg Ile Leu Arg Leu Phe Pro Trp
115            120            125

Ile Leu Leu Gly Ser Leu Met Ile Thr Cys Val Thr Ile Ile Pro Ser
130            135            140

Ala Ile Gly Asn Tyr Ile Gln Ile Gln Leu Leu Thr Met Glu His Leu
145            150            155            160

Pro Arg Asn Ser Thr Val Thr Asp Lys Leu Glu Asn Phe His Gln Tyr
165            170            175

Gln Phe Gln Ala His Thr Val Ala Leu Val Ile Pro Phe Ile Leu Phe
180            185            190

Leu Ala Ser Thr Ile Phe Leu Met Ala Ser Leu Thr Lys Gln Ile Gln
195            200            205

His His Ser Thr Gly His Cys Asn Pro Ser Met Lys Ala His Phe Thr
210            215            220

Ala Leu Arg Ser Leu Ala Val Leu Phe Ile Val Phe Thr Ser Tyr Phe
225            230            235            240

Leu Thr Ile Leu Ile Thr Ile Ile Gly Thr Leu Phe Asp Lys Arg Cys
245            250            255

Trp Leu Trp Val Trp Glu Ala Phe Val Tyr Ala Phe Ile Leu Met His
260            265            270

```

-continued

Ser Thr Ser Leu Met Leu Ser Ser Pro Thr Leu Lys Arg Ile Leu Lys
 275 280 285

Gly Lys Cys
 290

<210> SEQ ID NO 44
 <211> LENGTH: 873
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 44

```

atgataccca tccaactcac tgtcttcttc atgatcatct atgtgcttga gtccttgaca    60
attattgtgc agagcagcct aattgttgca gtgctgggca gagaatggct gcaagtcaga    120
aggctgatgc ctgtggacat gattctcatc agcctgggca tctctcgctt ctgtctacag    180
tgggcatcaa tgctgaacaa tttttgctcc tattttaatt tgaattatgt actttgcaac    240
ttaacaatca cctgggaatt tttaatatc cttacattct ggtaaacag cttgcttacc    300
gtgttctact gcacaaaggc ctcttcttcc acccatcaca tctttctctg gctgaggtgg    360
agaattttga gggtgtttcc ctggatatta ctgggttctc tgatgattac ttgtgtaaca    420
atcatccctt cagctattgg gaattacatt caaattcagt tactcaccat ggagcatcta    480
ccaagaaaca gcaactgaac tgacaaaact gaaaatttcc atcagtatca gttccaggct    540
catacagttg cattgggtat tcctttcatc ctgttctctg cctccaccat ctttctcatg    600
gcatcactga ccaagcagat acaacatcat agcactgggc actgcaatcc aagcatgaaa    660
gcgcacttca ctgccttgag gtcccttgcc gtcttattta ttgtgtttac ctcttacttt    720
ctaaccatac tcatcaccat tataggtact ctatttgata agagatgttg gttatgggtc    780
tggaagcctt ttgtctatgc tttcatctta atgcattcca cttcactgat gctgagcagc    840
cctacgttga aaaggattct aaagggaag tgc                                873

```

<210> SEQ ID NO 45
 <211> LENGTH: 316
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

```

Met Met Gly Leu Thr Glu Gly Val Phe Leu Ile Leu Ser Gly Thr Gln
1          5          10          15

Phe Thr Leu Gly Ile Leu Val Asn Cys Phe Ile Glu Leu Val Asn Gly
20        25        30

Ser Ser Trp Phe Lys Thr Lys Arg Met Ser Leu Ser Asp Phe Ile Ile
35        40        45

Thr Thr Leu Ala Leu Leu Arg Ile Ile Leu Leu Cys Ile Ile Leu Thr
50        55        60

Asp Ser Phe Leu Ile Glu Phe Ser Pro Asn Thr His Asp Ser Gly Ile
65        70        75        80

Ile Met Gln Ile Ile Asp Val Ser Trp Thr Phe Thr Asn His Leu Ser
85        90        95

Ile Trp Leu Ala Thr Cys Leu Gly Val Leu Tyr Cys Leu Lys Ile Ala
100       105       110

Ser Phe Ser His Pro Thr Phe Leu Trp Leu Lys Trp Arg Val Ser Arg
115       120       125

```

-continued

Val Met Val Trp Met Leu Leu Gly Ala Leu Leu Leu Ser Cys Gly Ser
130 135 140

Thr Ala Ser Leu Ile Asn Glu Phe Lys Leu Tyr Ser Val Phe Arg Gly
145 150 155 160

Ile Glu Ala Thr Arg Asn Val Thr Glu His Phe Arg Lys Lys Arg Ser
165 170 175

Glu Tyr Tyr Leu Ile His Val Leu Gly Thr Leu Trp Tyr Leu Pro Pro
180 185 190

Leu Ile Val Ser Leu Ala Ser Tyr Ser Leu Leu Ile Phe Ser Leu Gly
195 200 205

Arg His Thr Arg Gln Met Leu Gln Asn Gly Thr Ser Ser Arg Asp Pro
210 215 220

Thr Thr Glu Ala His Lys Arg Ala Ile Arg Ile Ile Leu Ser Phe Phe
225 230 235 240

Phe Leu Phe Leu Leu Tyr Phe Leu Ala Phe Leu Ile Ala Ser Phe Gly
245 250 255

Asn Phe Leu Pro Lys Thr Lys Met Ala Lys Met Ile Gly Glu Val Met
260 265 270

Thr Met Phe Tyr Pro Ala Gly His Ser Phe Ile Leu Ile Leu Gly Asn
275 280 285

Ser Lys Leu Lys Gln Thr Phe Val Val Met Leu Arg Cys Glu Ser Gly
290 295 300

His Leu Lys Pro Gly Ser Lys Gly Pro Ile Phe Ser
305 310 315

<210> SEQ ID NO 46

<211> LENGTH: 948

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

```

atgatgggac tcaccgaggg ggtgttcctg attctgtctg gcaactcagtt cacactggga      60
attctggtca attgtttcat tgagttggtc aatggtagca gctggttcaa gaccaagaga      120
atgtctttgt ctgacttcat catcaccacc ctggcactct tgaggatcat tctgctgtgt      180
attatcttga ctgatagttt ttaataagaa ttctctccca acacacatga ttcagggata      240
ataatgcaaa ttattgatgt ttctctggaca tttacaaacc atctgagcat ttggcttgcc      300
acctgtcttg gtgtcctcta ctgcctgaaa atcgccagtt tctctcacc caccattcctc      360
tggctcaagt ggagagtttc tagggtgatg gtatggatgc tgttgggtgc actgctctta      420
tctgtgggta gtaccgcac tctgatcaat gagtttaagc tctattctgt ctttagggga      480
attgaggcca ccaggaatgt gactgaacac ttcagaaaga agaggagtga gtattatctg      540
atccatgttc ttgggactct gtggtacctg cctcccttaa ttgtgtccct ggccctctac      600
tctttgtctc tcttctccct ggggaggcac acacggcaga tgctgcaaaa tgggacaagc      660
tccagagatc caaccactga ggcccacaag agggccatca gaatcatcct ttcttctctc      720
tttctcttct tactttactt tcttgcttct ttaattgcat catttggtta tttctacca      780
aaaaccaaga tggctaagat gattggcgaa gtaatgacaa tgttttatcc tgctggccac      840
tcattttatt tcattctggg gaacagtaag ctgaagcaga catttgtagt gatgctccgg      900
tgtgagtcct gtcactgaa gccctgatcc aagggaacca ttttctct      948

```

-continued

<210> SEQ ID NO 47
<211> LENGTH: 314
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47
Met Ala Thr Glu Leu Asp Lys Ile Phe Leu Ile Leu Ala Ile Ala Glu
1 5 10 15
Phe Ile Ile Ser Met Leu Gly Asn Val Phe Ile Gly Leu Val Asn Cys
20 25 30
Ser Glu Gly Ile Lys Asn Gln Lys Val Phe Ser Ala Asp Phe Ile Leu
35 40 45
Thr Cys Leu Ala Ile Ser Thr Ile Gly Gln Leu Leu Val Ile Leu Phe
50 55 60
Asp Ser Phe Leu Val Gly Leu Ala Ser His Leu Tyr Thr Thr Tyr Arg
65 70 75 80
Leu Gly Lys Thr Val Ile Met Leu Trp His Met Thr Asn His Leu Thr
85 90 95
Thr Trp Leu Ala Thr Cys Leu Ser Ile Phe Tyr Phe Phe Lys Ile Ala
100 105 110
His Phe Pro His Ser Leu Phe Leu Trp Leu Arg Trp Arg Met Asn Gly
115 120 125
Met Ile Val Met Leu Leu Ile Leu Ser Leu Phe Leu Leu Ile Phe Asp
130 135 140
Ser Leu Val Leu Glu Ile Phe Ile Asp Ile Ser Leu Asn Ile Ile Asp
145 150 155 160
Lys Ser Asn Leu Thr Leu Tyr Leu Asp Glu Ser Lys Thr Leu Phe Asp
165 170 175
Lys Leu Ser Ile Leu Lys Thr Leu Leu Ser Leu Thr Ser Phe Ile Pro
180 185 190
Phe Ser Leu Ser Leu Thr Ser Leu Leu Phe Leu Phe Leu Ser Leu Val
195 200 205
Arg His Thr Arg Asn Leu Lys Leu Ser Ser Leu Gly Ser Arg Asp Ser
210 215 220
Ser Thr Glu Ala His Arg Arg Ala Met Lys Met Val Met Ser Phe Leu
225 230 235 240
Phe Leu Phe Ile Val His Phe Phe Ser Leu Gln Val Ala Asn Trp Ile
245 250 255
Phe Phe Met Leu Trp Asn Asn Lys Tyr Ile Lys Phe Val Met Leu Ala
260 265 270
Leu Asn Ala Phe Pro Ser Cys His Ser Phe Ile Leu Ile Leu Gly Asn
275 280 285
Ser Lys Leu Arg Gln Thr Ala Val Arg Leu Leu Trp His Leu Arg Asn
290 295 300
Tyr Thr Lys Thr Pro Asn Ala Leu Pro Leu
305 310

<210> SEQ ID NO 48
<211> LENGTH: 942
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

-continued

```

atggccaccg aattggacaa aatctttctg attctggcaa tagcagaatt catcatcagc    60
atgctgggga atgtgttcat tggactggta aactgctctg aagggatcaa gaacccaaaag    120
gtctttctcag ctgacttcat cctcacctgc ttggctatct ccacaattgg acaactgttg    180
gtgatactgt ttgattcatt tctagtggga cttgcttcac atttatatac cacatataga    240
ctaggaaaaa ctgttattat gctttggcac atgactaatc acttgacaac ctggcttgcc    300
acctgcctaa gcattttcta tttctttaag atagcccact tccccactc ccttttcctc    360
tggctgaggt ggaggatgaa cggaatgatt gttatgcttc ttatattgtc ttgttctta    420
ctgatttttg acagtttagt gctagaaata tttattgata tctcactcaa tataatagat    480
aaaagtaatc tgactttata tttagatgaa agtaaaactc tctttgataa actctctatt    540
ttaaaaactc ttctcagctt gaccagtgtt atcccccttt ctctgtccct gacctccttg    600
ctttttttat ttctgtcctt ggtgagacat actagaaatt tgaagctcag ttccctgggc    660
tctagagact ccagcacaga ggcccatagg agggccatga aaatggtgat gtctttcctt    720
ttcctcttca tagttcattt tttttcctta caagtggcca attggatatt ttttatgttg    780
tggaacaaca agtacataaa gtttgtcatg ttagccttaa atgcctttcc ctctgtccac    840
tcatttattc tcattctggg aaacagcaag ctgcgacaga cagctgtgag gctactgtgg    900
catcttagga actatacaaa aacaccaaat gctttacctt tg                               942

```

<210> SEQ ID NO 49

<211> LENGTH: 318

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49

```

Met Asn Gly Asp His Met Val Leu Gly Ser Ser Val Thr Asp Lys Lys
1           5           10          15

Ala Ile Ile Leu Val Thr Ile Leu Leu Leu Arg Leu Val Ala Ile
          20          25          30

Ala Gly Asn Gly Phe Ile Thr Ala Ala Leu Gly Val Glu Trp Val Leu
          35          40          45

Arg Arg Met Leu Leu Pro Cys Asp Lys Leu Leu Val Ser Leu Gly Ala
          50          55          60

Ser Arg Phe Cys Leu Gln Ser Val Val Met Gly Lys Thr Ile Tyr Val
65          70          75          80

Phe Leu His Pro Met Ala Phe Pro Tyr Asn Pro Val Leu Gln Phe Leu
          85          90          95

Ala Phe Gln Trp Asp Phe Leu Asn Ala Ala Thr Leu Trp Ser Ser Thr
          100         105         110

Trp Leu Ser Val Phe Tyr Cys Val Lys Ile Ala Thr Phe Thr His Pro
          115         120         125

Val Phe Phe Trp Leu Lys His Lys Leu Ser Gly Trp Leu Pro Trp Met
          130         135         140

Leu Phe Ser Ser Val Gly Leu Ser Ser Phe Thr Thr Ile Leu Phe Phe
145         150         155         160

Ile Gly Asn His Arg Met Tyr Gln Asn Tyr Leu Arg Asn His Leu Gln
          165         170         175

Pro Trp Asn Val Thr Gly Asp Ser Ile Arg Ser Tyr Cys Glu Lys Phe
          180         185         190

```

-continued

Tyr Leu Phe Pro Leu Lys Met Ile Thr Trp Thr Met Pro Thr Ala Val
 195 200 205
 Phe Phe Ile Cys Met Ile Leu Leu Ile Thr Ser Leu Gly Arg His Arg
 210 215 220
 Lys Lys Ala Leu Leu Thr Thr Ser Gly Phe Arg Glu Pro Ser Val Gln
 225 230 235 240
 Ala His Ile Lys Ala Leu Leu Ala Leu Leu Ser Phe Ala Met Leu Phe
 245 250 255
 Ile Ser Tyr Phe Leu Ser Leu Val Phe Ser Ala Ala Gly Ile Phe Pro
 260 265 270
 Pro Leu Asp Phe Lys Phe Trp Val Trp Glu Ser Val Ile Tyr Leu Cys
 275 280 285
 Ala Ala Val His Pro Ile Ile Leu Leu Phe Ser Asn Cys Arg Leu Arg
 290 295 300
 Ala Val Leu Lys Ser Arg Arg Ser Ser Arg Cys Gly Thr Pro
 305 310 315

<210> SEQ ID NO 50
 <211> LENGTH: 957
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

```

atgaatggag accacatggt tctaggatct tcggtgactg acaagaaggc catcatcttg      60
gttaccattt tactcctttt acgcctggta gcaatagcag gcaatggctt catcactgct      120
gctctgggag tggagtgggt gctacggaga atgttgttgc cttgtgataa gttatttggt      180
agcctagggg cctctcgctt ctgtctgcag tcagtggtaa tgggtaagac catttatggt      240
ttcttgcatc cgatggcctt cccatacaac cctgtactgc agtttctagc ttccagtggt      300
gacttcttga atgctgccac cttatgggtc tctacctggc tcagtgtctt ctatttgtgtg      360
aaaattgcta ccttcaccca cctgtcttc ttctggctaa agcacaagtt gtctgggtgg      420
ctaccatgga tgctcttcag ctctgtaggg ctctccagct tcaccacat tctatttttc      480
ataggcaacc acagaatgta tcagaactat ttaaggaacc atctacaacc ttggaatgtc      540
actggcgata gcatacggag ctactgtgag aaattctatc tcttccctct aaaaatgatt      600
acttggaaca tgcccactgc tgtctttttc atttgcata ttttgcctat cacatctctg      660
ggaagacaca ggaagaaggc tctccttaca acctcaggat tccgagagcc cagtgtgcag      720
gcacacataa aggcctctgt ggcctctctc tcttttgcca tgctcttcat ctcatatttc      780
ctgtcactgg tggtcactgc tgcaggtatt tttccacctc tggactttaa attctgggtg      840
tgaggagtcg tgatttatct gtgtgcagca gttcacccca tcattctgct cttcagcaac      900
tgcaggctga gagctgtgct gaagagtctg cgttcctcaa ggtgtgggac accttga      957
  
```

<210> SEQ ID NO 51
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer comprising EcoRI restriction site
 for PCR amplification of hTAS2R16

<400> SEQUENCE: 51

- continued

cctgggaatt ttttaatatc cttacattct ggt

33

<210> SEQ ID NO 52
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer comprising NotI restriction site
 for PCR amplification of hTAS2R16

<400> SEQUENCE: 52

gaagcgcgct ttcattgctt

19

We claim:

1. A polynucleotide selected from the group consisting of:
 - (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NO: 12;
 - (b) polynucleotides having the coding sequence, as shown in SEQ ID NO: 11 encoding at least the mature form of the polypeptide;
 - (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to said polypeptide, and said fragment or derivative has bitter substance binding activity;
 - (d) polynucleotides which are at least 50% identical to a polynucleotide as defined in any one of (a) to (c) and which code for a polypeptide having bitter substance binding activity; and
 - (e) polynucleotides the complementary strand of which hybridizes under stringent conditions to a polynucleotide as defined in any one of (a) to (d) and which code for a polypeptide having bitter substance binding activity;
 or the complementary strand of such a polynucleotide.
2. The polynucleotide of claim 1 which is DNA, genomic DNA or RNA.
3. A vector containing the polynucleotide of claim 1.
4. The vector of claim 3 in which the polynucleotide is operatively linked to one or more expression control sequences allowing expression in prokaryotic and/or eukaryotic host cells.
5. A host cell genetically engineered to comprise a polynucleotide of claim 1.
6. A transgenic non-human animal comprising a polynucleotide of claim 1.
7. A process for producing a polypeptide encoded by a polynucleotide of claim 1, comprising: culturing a host cell genetically engineered to comprises a polynucleotide of claim 1 and recovering the polypeptide encoded by said polynucleotide.
8. A process for producing cells capable of expressing at least one bitter taste receptor polypeptide comprising genetically engineering cells in vitro with a vector containing a polynucleotide of claim 1, wherein said bitter taste receptor polypeptide(s) is(are) encoded by a polynucleotide of claim 1.
9. A polypeptide having the amino acid sequence encoded by a polynucleotide selected from the group consisting of:
 - (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NO: 12;
 - (b) polynucleotides having the coding sequence, as shown in SEQ ID NO: 11 encoding at least the mature form of the polypeptide;
 - (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to
10. An antibody that specifically binds to the polypeptide of claim 9.
11. A nucleic acid molecule which specifically hybridizes to a polynucleotide of claim 1.
12. An antagonist/inhibitor against the polypeptide of claim 9 which is an antibody, the extracellular domain of the polypeptide of claim 9 or a fragment thereof, or an inhibiting RNA.
13. The antagonist/inhibitor of claim 12, wherein said inhibiting RNA is an antisense construct, RNAi, siRNA or a ribozyme.
14. A process for isolating a compound that binds to a polypeptide encoded by a polynucleotide selected from the group consisting of:
 - (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NO: 12;
 - (b) polynucleotides having the coding sequence, as shown in SEQ ID NO: 11 encoding at least the mature form of the polypeptide;
 - (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to

- said polypeptide, and said fragment or derivative has bitter substance binding activity;
- (d) polynucleotides which are at least 50% identical to a polynucleotide as defined in any one of (a) to (c) and which code for a polypeptide having bitter substance binding activity; and
 - (e) polynucleotides the complementary strand of which hybridizes under stringent conditions to a polynucleotide as defined in any one of (a) to (d) and which code for a polypeptide having bitter substance binding activity;

comprising:

- (1) contacting said polypeptide or a host cell genetically engineered with said polynucleotide or with a vector containing said polynucleotide with a compound;
- (2) detecting the presence of the compound which binds to said polypeptide; and
- (3) determining whether the compound binds said polypeptide.

15. A process for isolating an antagonist of the bitter taste receptor activity of the polypeptide encoded by a polynucleotide selected from the group consisting of:

- (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NO: 12;
- (b) polynucleotides having the coding sequence, as shown in SEQ ID NO: 11 encoding at least the mature form of the polypeptide;
- (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to said polypeptide, and said fragment or derivative has bitter taste receptor activity;
- (d) polynucleotides which are at least 50% identical to a polynucleotide as defined in any one of (a) to (c) and which code for a polypeptide having bitter taste receptor activity; and
- (e) polynucleotides the complementary strand of which hybridizes under stringent conditions to a polynucleotide as defined in any one of (a) to (d) and which code for a polypeptide having bitter taste receptor activity;

comprising:

- (1) contacting said polypeptide or a host cell genetically engineered with said polynucleotide or with a vector containing said polynucleotide with a potential antagonist; and
- (2) determining whether the potential antagonist antagonizes the bitter taste receptor activity of said polypeptide.

16. The process of claim **15** further comprising the contacting of the polypeptide with an agonist of the respective bitter taste receptor activity.

17. The process of claim **16** in which said contacting with an agonist is carried out prior to, concomitantly with, or after step (1).

18. The process of claim **16** in which said agonist is selected from the group consisting of saccharin and acesulfame K and functional derivatives thereof.

19. A process for the production of a food or any precursor material or additive employed in the production of foodstuffs comprising the steps of the process of claim **14** and the subsequent step of admixing the identified compound or antagonist

with foodstuffs or any precursor material or additive employed in the production of foodstuffs.

20. A process for the production of a nutraceutical or pharmaceutical composition comprising the steps of the process of claim **14** and the subsequent step of formulating the compound or antagonist with an active agent in a pharmaceutically acceptable form.

21. A food stuff or any precursor material or additive employed in the production of foodstuffs comprising an antagonist/inhibitor of claim **12**.

22. A nutraceutical or pharmaceutical composition comprising an antagonist/inhibitor of claim **12** and an active agent and optionally a pharmaceutically acceptable carrier.

23. A method for treating an abnormally increased or decreased sensitivity towards a bitter substance wherein said method comprises administering, to a subject in need of such treatment, one or more agents selected from the group consisting of:

- 1) A polynucleotide selected from the group consisting of:
 - (a) polynucleotides encoding at least the mature form of the polypeptide having the deduced amino acid sequence as shown in SEQ ID NO: 12;
 - (b) polynucleotides having the coding sequence, as shown in SEQ ID NO: 11 encoding at least the mature form of the polypeptide;
 - (c) polynucleotides encoding a fragment or derivative of a polypeptide encoded by a polynucleotide of any one of (a) to (b), wherein in said derivative one or more amino acid residues are conservatively substituted compared to said polypeptide, and said fragment or derivative has bitter substance binding activity;
 - (d) polynucleotides which are at least 50% identical to a polynucleotide as defined in any one of (a) to (c) and which code for a polypeptide having bitter substance binding activity; and
 - (e) polynucleotides the complementary strand of which hybridizes under stringent conditions to a polynucleotide as defined in any one of (a) to (d) and which code for a polypeptide having bitter substance binding activity;

or the complementary strand of such a polynucleotide;

- 2) a vector comprising a polynucleotide of part 1) above;
- 3) an antibody that specifically binds to a polypeptide that is encoded by a polynucleotide of part 1) above and which is obtainable by culturing a host cell genetically engineered to comprise a polynucleotide of part 1) above, and recovering the polypeptide encoded by said polynucleotide; and
- 4) an antagonist/inhibitor against the polypeptide of part 3) above, which is an antibody, the extracellular domain of the polypeptide of part 3) above, or a fragment thereof, or an inhibiting RNA.

24. A process for the production of a food or any precursor material or additive employed in the production of foodstuffs comprising the steps of the process of claim **15** and the subsequent step of admixing the identified compound or antagonist with foodstuffs or any precursor material or additive employed in the production of foodstuffs.

25. A process for the production of a nutraceutical or pharmaceutical composition comprising the steps of the process of claim **15** and the subsequent step of formulating the compound or antagonist with an active agent in a pharmaceutically acceptable form.

* * * * *