

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 March 2009 (05.03.2009)

PCT

(10) International Publication Number  
**WO 2009/027473 A1**

(51) International Patent Classification:

A61K 31/4164 (2006.01) A61P 31/10 (2006.01)  
A61K 31/4412 (2006.01) A61P 31/12 (2006.01)  
A61K 45/06 (2006.01) A01N 43/54 (2006.01)  
A61P 31/04 (2006.01) A01N 43/40 (2006.01)

(21) International Application Number:

PCT/EP2008/061312

(22) International Filing Date: 28 August 2008 (28.08.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

07115421.5 31 August 2007 (31.08.2007) EP

(71) Applicant (for all designated States except US):  
JANSSEN PHARMACEUTICA NV [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BOSSELAERS, Jan Pieter Hendrik** [BE/BE]; Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **BYLEMANS, Dany Leopold Jozefien** [BE/BE]; Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **KEMPEN, Tony Mathilde Jozef** [BE/BE]; Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **LEITHOFF, Hans Bernhard** [DE/DE]; Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **THYS, Amber Paula Marcella** [BE/BE]; Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). **VAN DER**

**FLAAS, Mark Arthur Josepha** [BE/BE]; Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE).

(74) Common Representative: **JANSSEN PHARMACEUTICA NV**; Turnhoutseweg 30, B-2340 Beerse (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

**Published:**

- with international search report

(54) Title: COMBINATIONS OF IMAZALIL AND HYDROXYPYRIDONES

(57) Abstract: The present invention relates to combinations of imazalil, or a salt thereof, and hydroxypyridones which provide an improved biocidal effect. More particularly, the present invention relates to compositions comprising a combination of imazalil, or a salt thereof, together with one or more hydroxypyridones selected from 1-hydroxy-2-pyridinone, ciclopirox, ciclopirox olamine, piroctone, piroctone olamine and rilopirox in respective proportions to provide a synergistic biocidal effect. Compositions comprising these combinations are useful for the protection of any living or non-living material, such as crops, plants, fruits, seeds, objects made of wood, thatch or the like, engineering material, biodegradable material and textiles against deterioration due to the action of microorganisms such as bacteria, fungi, yeasts, algae, viruses, and the like.

WO 2009/027473 A1

## COMBINATIONS OF IMAZALIL AND HYDROXYPYRIDONES

---

**[0001]** The present invention relates to combinations of imazalil, or a salt thereof,  
5 and hydroxypyridones which provide an improved biocidal effect. More particularly,  
the present invention relates to compositions comprising a combination of imazalil, or  
a salt thereof, together with one or more hydroxypyridones selected from 1-hydroxy-2-  
pyridinone, ciclopirox, ciclopirox olamine, piroctone, piroctone olamine and rilopirox in  
10 respective proportions to provide a synergistic biocidal effect. Compositions  
comprising these combinations are useful for the protection of any living or non-living  
material, such as crops, plants, fruits, seeds, objects made of wood, thatch or the like,  
engineering material, biodegradable material and textiles against deterioration due to  
the action of microorganisms such as bacteria, fungi, yeasts, algae, virusses, and the  
like.

15  
**[0002]** Microorganisms are extremely useful, and even indispensable, in processes  
such as, e.g. alcoholic fermentation, ripening of cheese, baking of bread, production of  
penicillin, purification of waste water, production of biogas, and the like. However,  
microorganisms can also be harmful or highly dangerous; by causing infectious  
20 diseases, by forming poisonous or carcinogenic metabolites and by attacking valuable  
materials, disturbing production processes, or impairment of the quality of products.

**[0003]** Biocides or microbiocides are a broad and diverse group of compounds which  
are able to control microorganisms : *i.e.* to eliminate, kill, or inhibit microorganisms, or  
25 to reduce the growth or proliferation of microorganisms such as bacteria, fungi, yeasts  
and algae. An important group of the biocides are the bactericides and fungicides.  
Since bacteria and fungi occur everywhere, their destructive activity (biodeterioration)  
is basically unavoidable. Nevertheless objects can be protected with the aid of  
compounds that prevent the multiplication of bacteria or fungi at the relevant sites,  
30 either by killing them or inhibiting their development.

**[0004]** Fungicidal combinations comprising the antifungal agents imazalil,  
pyrimethanil or thiabendazole have been disclosed in, e.g. EP-0,336,489 which  
describes combinations of imazalil and propiconazole, WO-99/12422 which describes  
35 combinations of imazalil and epoxiconazole and WO-03/011030 which describes  
fungicidal compositions comprising pyrimethanil and imazalil.

**[0005]** WO-2004/105662 discloses microbicidal compositions comprising ciclopirox olamine as a microbicidal agent, and *Mycoeses*, 40, 243 - 247 (1997) discloses that hydroxypyridones chemically form a class of antimycotics which include ciclopirox and rilopirox.

5


**[0006]** It has now been found that the combination of imazalil (hereinafter referred to as component I) and one or more hydroxypyridones selected from 1-hydroxy-2-pyridinone, ciclopirox, ciclopirox olamine, piroctone, piroctone olamine and rilopirox (hereinafter referred to as a component II), has a synergistic effect on the control of microorganisms.

10

**[0007]** Imazalil, component (I), is a systemic fungicide with protective and curative action and is used to control a wide range of fungi on fruit, vegetables, and ornamentals, including powdery mildew on cucumber and black spot on roses.

15





Imazalil is also used as a seed dressing and for post harvest treatment of citrus, banana, and other fruit to control storage decay. It is the generic name of the compound 1-[2-(2,4-dichlorophenyl)-2-(2-propenyloxy)ethyl]-1*H*-imidazole, which compound may be represented by the formula



20

**[0008]** The hydroxypyridones (II) have the following structure :

1-hydroxy-2-pyridinone (II-a) (CAS 822-89-9)	<p style="text-align: right;">(II-a)</p>
ciclopirox (II-b) (CAS 29342-05-0)	<p style="text-align: right;">(II-b)</p>

ciclopirox olamine (II-c) (CAS 41621-49-2)	 <div style="text-align: right;">(II-c)</div>
piroctone (II-d) (CAS 50650-76-5)	 <div style="text-align: right;">(II-d)</div>
piroctone olamine (II-e) (CAS 68890-66-4)	 <div style="text-align: right;">(II-e)</div>
rilopirox (II-f) (CAS 104153-37-9)	 <div style="text-align: right;">(II-f)</div>

**[0009]** 1-Hydroxy-2-pyridinone, component (II-a), is an antimycotic agent that can be used in antidandruff shampoos.

- 5 **[0010]** The antifungal agent imazalil (I) may be present in its free base form or in the form of an acid addition salt, the latter being obtained by reaction of the base form with an appropriate acid. Appropriate acids comprise, for example, inorganic acids, such as the hydrohalic acids, *i.e.* hydrofluoric, hydrochloric, hydrobromic and hydroiodic, sulfuric acid, nitric acid, phosphoric acid, phosphinic acid and the like; or organic acids,
- 10 such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxy-butanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethane-sulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxy-benzoic and the like acids.
- 15

**[0011]** Particular salt forms of imazalil (I) are the sulfate, phosphate, acetate, nitrate or phosphite salts.

- [0012]** Imazalil (I) has one asymmetric carbon atom and can therefore be used in the embodied compositions in the form of a mixture of both enantiomers, in particular a racemic mixture, or in the form of a substantially pure (R)- or (S)-enantiomer. The term "substantially pure" as used hereinbefore means a purity (either chemical or optical), as determined by methods conventional in the art such as high performance liquid chromatography or optical methods, of at least about 96%, preferably at least 98% and more preferably at least 99%.
- [0013]** The compositions of the present invention have biocidal activity against a broad range of microorganisms such as bacteria, fungi, yeasts and viruses. Bacteria include Gram-positive and Gram-negative bacteria. Fungi include e.g. wood-discoloring fungi, wood-destroying fungi, and phytopathogenic fungi. Viruses include HIV, SARS and bird flu.
- [0014]** The biocidal compositions of the present invention are useful in the preservation of wood, wood products, leather, natural or synthetic textile, fibers, non-wovens, technical textile, plasticized materials and non-plasticized thermoplastics as polypropylene, polyvinylchloride, etc..., paper, wall paper, insulation material, laminates, amino moulding compounds, paints and coatings, fabrics, floor coverings, synthetic fibres like plasticized polymers, hessian, rope and cordage and biodegradable materials and protect said materials against attack and destruction by bacteria or fungi. As wood or wood products which can be preserved with the compositions according to the present invention is considered, for example, wood products such as timber, lumber, railway sleepers, telephone poles, fences, wood coverings, wicker-work, windows and doors, plywood, particle board, waferboards, chipboard, joinery, timber used above ground in exposed environments such as decking and timber used in ground contact or fresh water or salt water environments, bridges or wood products which are generally used in housebuilding, construction and carpentry. As biodegradable materials besides wood which can benefit from treatment with the compositions of the invention include cellulosic material such as cotton.
- [0015]** The biocidal compositions of the present invention are useful in the prevention of microbial contamination or biofilm formation in several industrial processes like gaskets, pipes and tubings in contact with fluids or involved in fluid transport, conveyor belts, surfaces and plastic components used in food transport, processing or production, and medical activities like medical equipment and devices like catheters, pacemakers, implants, surgery equipment and sterile textile.

**[0016]** The biocidal compositions of the present invention are useful in the prevention of hygienic concerns like unwanted bacterial, fungal or algal growth on surfaces, safety problems like the presence of Legionella in closed water systems, Nosocomial infections in hospitals, the presence of Multi-Resistant Staphylococcus aureus (MRSA), odor problems like in fabrics like socks, towels, protective uniforms, shoe linings or in filters or floor coverings. The invention is as well possible to protect areas or items coated with an ultra-hygienic polymer like for the manufacture of electrical devices such as light switches and switch plates; sanitary ware such as toilet seats; and door handles, handrails, baby-changing tables, telephones, and other end-use applications where the highest levels of sanitary protection are needed.

**[0017]** The biocidal compositions of the present invention are useful in the prevention of bacterial, fungal or algal growth on surfaces and herewith causing aesthetical problems for the materials considered

**[0018]** In an embodiment, the present invention relates to a method of controlling microbial growth on wood, wood products and biodegradable materials, which comprises applying an antimicrobially effective amount of a composition comprising a combination of component (I) and one or more components (II) in respective proportions to provide a synergistic biocidal effect, to the wood, wood products, leather, natural or synthetic textile, fibers, non-wovens, technical textile, plasticized materials and non-plasticized thermoplastics as polypropylene, polyvinylchloride, etc..., paper, wall paper, insulation material, laminates, amino moulding compounds, paints and coatings, fabrics, floor coverings, synthetic fibres like plasticized polymers, hessian, rope and cordage.

**[0019]** The biocidal compositions of the present invention are also useful to protect engineering materials against microorganisms. Engineering materials which are intended to be protected can be glues, sizes, paints and plastic articles, cooling lubricants, aqueous hydraulic fluids and other non-living materials which can be infested with, or decomposed by, microorganisms.

**[0020]** In an embodiment, the present invention relates to a method of controlling microbial growth on engineering materials, which comprises applying an antimicrobially effective amount of a composition comprising a combination of component (I) and one or more components (II) in respective proportions to provide a synergistic biocidal effect, to the engineering materials to be treated.

**[0021]** The biocidal compositions according to the present invention can also be used to protect plants, or parts of plants, e.g. fruit, blossoms, flowers, foliage, stems, roots, cuttings, tubers of plants, fruit and seeds.

5 **[0022]** As examples of the wide variety of culture plants in which the combinations of components (I) and (II) according to the present invention can be used, there may be named for example cereals, e.g. wheat, barley, rye, oats, rice, sorghum and the like; beets, e.g. sugar beet and fodder beet; pome and stone fruit and berries, e.g. apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries;  
10 leguminous plants, e.g. beans, lentils, peas, soy beans; oleaginous plants, e.g. rape, mustard, poppy, olive, sunflower, coconut, castor-oil plant, cocoa, ground-nuts; cucurbitaceae, e.g. pumpkins, gherkins, melons, cucumbers, squashes; fibrous plants, e.g. cotton, flax, hemp, jute; citrus fruit, e.g. orange, lemon, grapefruit, mandarin; vegetables, e.g. spinach, lettuce, asparagus, brassicaceae such as cabbages and  
15 turnips, carrots, onions, tomatoes, potatoes, hot and sweet peppers; laurel-like plants, e.g. avocado, cinnamon, camphor tree; or plants such as maize, tobacco, nuts, coffee, sugar-cane, tea, vines, hops, bananas, rubber plants, as well as ornamental plants, e.g. flowers, shrubs, deciduous trees and evergreen trees such as conifers. This enumeration of culture plants is given with the purpose of illustrating the invention and  
20 not to delimiting it thereto.

**[0023]** In an embodiment, the present invention relates to a method of controlling microbial growth on plants, parts of plants, fruit and seeds, which comprises applying an antimicrobially effective amount of a composition comprising a combination of  
25 component (I) and one or more components (II) in respective proportions to provide a synergistic biocidal effect, to the plants, parts of plants, fruit and seeds to be treated.

**[0024]** The relative proportions of component (I) and a component (II) in compositions comprising a combination of component (I) and one of the components  
30 (II) are those proportions which result in a synergistic biocidal effect, when compared to a composition including, as an active ingredient, either component (I) alone or one of the components (II) alone. As will be understood by those skilled in the art, the said synergistic effect may be obtained within various proportions of components (I) and (II) in the composition, depending on the kind of microorganism towards which effect is  
35 measured and the substrate to be treated. Based on the teachings of the present application, determination of the synergistic effect of such combinations can be performed according to the procedures of the Poison Plate Assay as described in

Experiment 1. As a general rule, however, it may be said that for most micro-organisms the suitable proportions by weight of the amount of component (I) to component (II) in the active composition should lie in the range from 10:1 to 1:10. Particularly, this range is from 4:1 to 1:4, more particularly from 3:1 to 1:3 or 2:1 to 1:2.

5 Another particular ratio of component (I) to component (II) in the compositions of the present invention is a 1:1 ratio between component (I) and component (II).

**[0025]** The quantity of each of the active ingredients in the compositions according to the present invention will be so that a synergistic biocidal effect is obtained. In particular it is contemplated that the ready to use compositions of the present invention

10 comprise component (I) in a range from 10 to 50.000 mg/l. The component (II) is present in an amount ranging from 10 to 50.000 mg/l or mg/kg depending upon the specific activity of the selected component (II).

**[0026]** The concentration of component (I) and one of the components (II) in the ready to use compositions is also dependent upon the specific conditions wherein these compositions are used. For instance, in foliar treatment a solution is sprayed directly onto the leaves wherein the concentration of component (I) ranges from 100 mg to 250 mg/l. Potatoes are treated with a composition comprising component (I) in an amount of about 7500 mg/l in such a manner that a solution of 2 litres is used for

20 treating 1.000 kg. In the treatment of seed the compositions used comprise component (I) in an amount of about 50 g/l in such a manner that 100 kg of seed is treated with a solution of 100 ml to 200 ml. In the post-harvest treatment of fruit compositions are used comprising component (I) in an amount ranging from 250 to

25 500 mg/l in dip treatment, from 500 to 1.000 mg/l in spray treatment, and from 1.000 to 2.000 mg/l in wax treatment.

**[0027]** The compositions according to the present invention comprise as a combination of component (I) and one or more components (II) in respective proportions to provide a synergistic biocidal effect, and optionally one or more

30 acceptable carriers.

**[0028]** These carriers are any material or substance with which the composition of components (I) and (II) is formulated in order to facilitate its application/dissemination to the locus to be treated, for instance by dissolving, dispersing, or diffusing the said composition, and/or to facilitate its storage, transport or handling without impairing its antifungal effectiveness. Said acceptable carriers may be a solid or a liquid or a gas which has been compressed to form a liquid, *i.e.* the compositions of this invention can

35



suitably be used as concentrates, emulsions, emulsifiable concentrates, oil miscible suspension concentrates, oil-miscible liquid, soluble concentrates, solutions, granulates, dusts, sprays, aerosols, pellets, or powders.

5     **[0029]** In many instances the biocidal compositions to be used directly can be obtained from concentrates, such as e.g. emulsifiable concentrates, suspension concentrates, or soluble concentrates, upon dilution with aqueous or organic media, such concentrates being intended to be covered by the term composition as used in the definitions of the present invention. Such concentrates can be diluted to a ready to  
10 use mixture in a spray tank shortly before use. Preferably the compositions of the invention should contain from about 0.01 to 95% by weight of the combination of components (I) and (II). More preferably this range is from 0.1 to 90% by weight. Most preferably this range is from 1 to 80% by weight, depending on the type of formulation to be selected for specific application purposes, as further explained in  
15 details hereinafter.

**[0030]** An emulsifiable concentrate is a liquid, homogeneous formulation of the components (I) and (II) to be applied as an emulsion after dilution in water. A suspension concentrate is a stable suspension of the active ingredients in a fluid  
20 intended for dilution with water before use. A soluble concentrate is a liquid, homogeneous formulation to be applied as a true solution of the active ingredients after dilution in water.

**[0031]** The fungicidal compositions of the present invention can also be formulated  
25 as waxes for use as a cover or coating of e.g. fruit, in particular citrus fruit.

**[0032]** Biocidal compositions according to the present invention can be of use in post-harvest treatment of fruit, especially citrus fruit. In the latter instance, the fruit will be sprayed with or dipped or drenched into a liquid formulation or the fruit may be  
30 coated with a waxy composition. The latter waxy composition conveniently is prepared by thoroughly mixing a suspension concentrate with a suitable wax. The formulations for spray, dip or drench applications may be prepared upon dilution of a concentrate such as, e.g. an emulsifiable concentrate, a suspension concentrate or a soluble liquid, with an aqueous medium. Such concentrate in most instances consists  
35 of the active ingredients, a dispersing or suspending agent (surfactant), a thickening agent, a small amount of organic solvent, a wetting agent, optionally some anti-freeze agent, and water.

[0033] The biocidal compositions of the present invention can also be used for protecting seed against fungi. To that effect the present fungicidal compositions can be coated on seed, in which case the seed grains are drenched consecutively with a liquid composition of the active ingredients or if they are coated with a previously combined composition. The compositions can also be sprayed or atomised onto the seed using e.g. a spinning disc atomiser.

[0034] The combination of components (I) and (II) is preferably applied in the form of compositions wherein both said components are intimately admixed in order to ensure simultaneous administration to the materials to be protected. Administration or application of both components (I) and (II) can also be a "sequential-combined" administration or application, *i.e.* component (I) and one or more components (II) are administered or applied alternatively or sequentially in the same place in such a way that they will necessarily become admixed together at the locus to be treated. This will be achieved namely if sequential administration or application takes place within a short period of time e.g. within less than 24 hours, preferably less than 12 hours. This alternative method can be carried out for instance by using a suitable single package comprising at least one container filled with a formulation comprising the active component (I) and at least one container filled with a formulation comprising an active component (II). Therefore the present invention also encompasses a product containing :

- (a) a composition comprising component (I) (*i.e.* imazalil) and
- (b) a composition comprising as a component (II), (a hydroxypyridone selected from 1-hydroxy-2-pyridinone, ciclopirox, ciclopirox olamine, piroctone, piroctone olamine and rilopirox); as a combination for simultaneous or sequential use, wherein said compositions (a) and (b) are in respective proportions to provide a synergistic biocidal effect. Such products may consist of a suitable package comprising separate containers wherein each container comprises component (I) or component (II), preferably in formulated form. Such formulated forms in general have the same composition as described for the formulations containing both active ingredients.

[0035] Appropriate carriers and adjuvants for use in the compositions of the present invention may be solid or liquid and correspond to suitable substances known in the art of formulation, such as, for example natural or regenerated mineral substances, solvents, dispersants, surfactants, wetting agents, adhesives, thickeners, binders, fertilizers or anti-freeze agents.

**[0036]** Apart from both the aforementioned components (I) and (II), the compositions according to the present invention may further comprise other active ingredients, e.g. other microbiocides, in particular fungicides, and also insecticides, acaricides, nematocides, herbicides, plant growth regulators and fertilizers.

5

**[0037]** The components (I) and (II) are used in unmodified form or, preferably, together with the adjuvants conventionally employed in the art of formulation. They are therefore formulated following art-known procedures to emulsifiable concentrates, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations in e.g. polymer substances. As with the nature of the compositions, the methods of application, such as spraying, atomizing, dusting, scattering or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances.

10

**[0038]** The formulations, *i.e.* the compositions, preparations or mixtures comprising the active ingredients and, where appropriate, a solid or liquid adjuvant, are prepared in known manner, e.g. by homogeneously mixing and/or grinding the active ingredients with extenders, e.g. solvents, solid carriers and, where appropriate, surface-active compounds (surfactants).

15

20

**[0039]** Suitable solvents are aromatic hydrocarbons, preferably the fractions containing 8 to 12 carbon atoms, e.g. dimethylbenzene mixtures or substituted naphthalenes, phthalates such as dibutyl phthalate or dioctyl phthalate, aliphatic or alicyclic hydrocarbons such as cyclohexane or paraffins, alcohols and glycols and their ethers and esters, such as ethanol, ethylene glycol, ethylene glycol monomethyl or monoethyl ether, ketones such as cyclohexanone, strongly polar solvents such as *N*-methyl-2-pyrrolidone, dimethylsulfoxide or dimethylformamide, as well as vegetable oils or epoxidised vegetable oils such as epoxidised coconut oil or soybean oil; or water.

25

30

**[0040]** The solid carriers used e.g. for dusts and dispersible powders are normally natural mineral fillers such as calcite, talcum, kaolin, montmorillonite or attapulgite. In order to improve the physical properties it is also possible to add highly dispersed silicic acid or highly dispersed absorbent polymers. Suitable granulated absorbent carriers are of the porous type, for example pumice, broken brick, sepiolite or bentonite; and suitable nonsorbent carriers are materials such as calcite or sand. In addition, a great number of pregranulated materials of inorganic or organic nature can be used, e.g. especially dolomite or pulverised plant residues.

35

5     **[0041]** Suitable surface-active compounds to be used in the compositions of the present invention are non-ionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. The term "surfactants" will also be understood as comprising mixtures of surfactants.

10     **[0042]** Appropriate carriers and adjuvants for use in the compositions of the present invention may be solid or liquid and correspond to suitable substances known in the art for preparing formulations for treating plants or their loci, or for treating plant products, in particular for treating wood, such as, for example, natural or regenerated mineral substances, solvents, dispersants, surfactants, wetting agents, adhesives, thickeners, binders, fertilizers, anti-freeze agents, repellents, colour additives, corrosion inhibitors, water-repelling agents, siccatives, UV-stabilizers and other active ingredients.

15     **[0043]** Suitable anionic surfactants can be both water-soluble soaps and water-soluble synthetic surface-active compounds.

20     **[0044]** Suitable soaps are the alkali metal salts, earth alkaline metal salts or unsubstituted or substituted ammonium salts of higher fatty acids (C<sub>10</sub>-C<sub>22</sub>), e.g. the sodium or potassium salts of oleic or stearic acid, or of natural fatty acid mixtures which can be obtained e.g. from coconut oil or tallow oil. In addition, there may also be mentioned fatty acid methyltaurin salts.

25     **[0045]** More frequently, however, so-called synthetic surfactants are used, especially fatty sulfonates, fatty sulfates, sulfonated benzimidazole derivatives or alkylarylsulfonates. The fatty sulfonates or sulfates are usually in the form of alkali metal salts, earth alkaline metal salts or unsubstituted or substituted ammonium salts and contain an alkyl radical having from 8 to 22 carbon atoms said alkyl also comprising radicals derived from acyl radicals, e.g. the sodium or calcium salt of lignosulfonic acid, of dodecylsulfate or of a mixture of fatty alcohol sulfates obtained from natural fatty acids. These compounds also comprise the salts of sulfuric acid esters and sulfonic acids of fatty alcohol/ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain 2 sulfonic acid groups and one fatty acid radical containing 8 to 22 carbon atoms. Examples of alkylarylsulfonates are the sodium, calcium or triethanolamine salts of dodecylbenzene sulfonic acid, dibutylnaphthalene-sulfonic acid, or of a naphthalene-sulfonic acid/formaldehyde condensation product. Also suitable are corresponding phosphates, e.g. salts of the

35

phosphoric acid ester of an adduct of p-nonylphenol with 4 to 14 moles of ethylene oxide, or phospholipids.

5 **[0046]** Non-ionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, or saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 10 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols.

10 **[0047]** Further suitable non-ionic surfactants are the water-soluble adducts of polyethylene oxide with polypropylene glycol, ethylenediaminopolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which adducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit.

15 **[0048]** Representative examples of non-ionic surfactants are nonylpheno-lpolyethoxy ethanols, castor oil polyglycol ethers, polypropylene/polyethylene oxide adducts, tributylphenoxypolyethoxyethanol, polyethylene glycol and octylphenoxypolyethoxyethanol. Fatty acid esters of polyethylene sorbitan, such as polyoxyethylene sorbitan trioleate, are also suitable non-ionic surfactants.

**[0049]** Particularly advantageous additives useful to improve the application and reduce the dose of the active ingredients, are the natural (animal or plant) or synthetic phospholipids of the cephalin or lecithin type such as, for example, phosphatidyl-ethanolamine, phosphatidylserine, phosphatidylglycerine, lysolecithin, or cardiolipin. Such phospholipids may be obtained from animal or plant cells, in particular from brain-, heart- or liver tissue, egg yolks or soy beans. Appropriate such phospholipids are for instance, phosphatidylcholine mixtures. Synthetic phospholipids are for instance, dioctanoylphosphatidylcholine and dipalmitoylphosphatidylcholine.

30

#### Experimental part

##### Experiment 1 : poison plate assay

Name of component (I) : - imazalil (I)

35 Name of component (II) :  
 - 1-hydroxy-2-pyridinone (II-a)  
 - ciclopirox (II-b)  
 - ciclopirox olamine (II-c)  
 - piroctone (II-d)  
 - piroctone olamine (II-e)  
 40 - rilopirox (II-f)

	Test models :	Fungi: to each well of a 24-well microtiter plate, 1000 µl of Potato Dextrose Agar (PDA: 4g potato infusion, 20g bacto dextrose and 15g bacto agar in 1 liter deionised water) medium containing the appropriate combination of the test compounds in one of the concentrations of a dose series is added. The nutrient medium is inoculated with the test fungi by adding a spore/mycelium suspension (10 µl) or a small piece of agar from the margin of an actively growing colony and incubated under dark at 27°C with 70 % relative humidity. The growth of the fungi is evaluated after two weeks.		
5				
10				
15		Bacteria: to each well of a 24-well microtiter plate, 1000 µL Tryptose Agar (TA, 20g bacto tryptose agar, 5g sodium chloride, 1g bacto dextrose and 15g agar in 1 liter deionised water), containing the test compounds in one of the concentrations of a dose series is added. The nutrient medium is inoculated by adding 10 µl of an actively growing liquid culture of the test bacteria in tryptose broth (without agar), and incubated under dark at 27°C with 70 % relative humidity. The growth is evaluated after five days.		
20				
25	Concentrations:	For the combination experiments, two overlapping dose series were used, in order to maximise the chance of the MIC values falling within the test concentration range for all combinations:		
30		Series 1 : 50.00 - 37.50 - 28.13 - 21.09 - 15.82 - 11.87 - 8.90 - 6.67 - 5.01 - 3.75 - 2.82 - 2.11 ppm		
35		Series 2 : 25.00 - 18.75 - 14.06 - 10.55 - 7.91 - 5.93 - 4.45 - 3.34 - 2.50 - 1.88 - 1.41 - 1.06 ppm		
40	Test combinations :	% product A	+	% product B
		100	+	0
		80	+	20
		66	+	33
		50	+	50
		33	+	66
		20	+	80
		0	+	100

Test species:

Bacteria:

*Staphylococcus aureus* (g+) LMG 8064

Fungi:

5	<i>Botrytis cinerea</i>	BC03019
	<i>Chaetomium globosum</i>	ATCC6205
	<i>Coriolus versicolor</i>	CTB863A
	<i>Humicola grisea</i>	MG28
	<i>Ulocladium atrum</i>	IMI214699a

10

MIC values (minimum inhibitory concentration in ppm total active ingredient) were noted and synergy was calculated using the Synergy Index method described by Kull et al. (Kull, F.C., P.C. Eismann, H.D. Sylvestrowicz, and R.L. Mayer (1961) "Mixtures of quaternary ammonium compounds and long-chain fatty acids as antifungal agents" *Applied Microbiology* 9: 538-541; also see Zwart Voorspuij, A.J., and C.A.G. Nass (1957) "Some aspects of the notions additivity, synergism and antagonism in the simultaneous activity of two antibacterial agents in vitro" *Arch. intern.*

15

*Pharmacodynamie* 109: 211-228; Steinberg, D.C. (2000) "Measuring synergy" *cosmetics & Toiletries* 115(11): 59-62; and Lada, A., A.N. Petrocci, H.A. Green, and J.J. Merianos (1977) "Antimicrobial composition" US Patent 4061750, 3pp.):

20

$$\text{Synergy Index (SI)} = \frac{Q_a}{Q_A} + \frac{Q_b}{Q_B}$$

wherein:

- $Q_A$  is the concentration of compound A in ppm, acting alone, which produced an end point (e.g. MIC),
- $Q_a$  is the concentration of compound A in ppm, in the mixture, which produced an end point (e.g. MIC),
- $Q_B$  is the concentration of compound B in ppm, acting alone, which produced an end point (e.g. MIC),
- $Q_b$  is the concentration of compound B in ppm, in the mixture, which produced an end point (e.g. MIC).

25

30

When the Synergy Index is greater than 1.0, antagonism is indicated. When the SI is equal to 1.0, additivity is indicated. When the SI is less than 1.0, synergism is demonstrated.

35

Table 1: MIC-values (minimum inhibitory concentration in ppm) and synergy index of combinations of imazalil (component (I)) with 1-hydroxy-2-pyridinone (component (II-a))

	% (I) + % (II-a)	MIC-values in ppm	Synergy Index
<i>Staphylococcus aureus</i>	100 + 0	25.00	-
	80 + 20	25.00	0.83
	66 + 33	25.00	0.72
	50 + 50	18.75	0.43
	33 + 66	14.06	0.25
	20 + 80	18.75	0.24
	0 + 100	160.00	-



Claims

1. A composition comprising a combination of imazalil or a salt thereof, as component (I), and as a component (II) a hydroxypyridone selected from  
5 1-hydroxy-2-pyridinone (II-a), ciclopirox (II-b), ciclopirox olamine (II-c), piroctone (II-d), piroctone olamine (II-e) and rilopirox (II-f);  
whereby component (I) and one of the components (II) are in respective proportions to provide a synergistic biocidal effect.
- 10 2. A composition as claimed in claim 1 wherein the hydroxypyridone is 1-hydroxy-2-pyridinone (II-a).
3. A composition as claimed in claim 1 or 2 wherein the ratio by weight of component (I) to component (II) ranges from 10:1 to 1:10.
- 15 4. A composition as claimed in claim 3 wherein the ratio by weight of component (I) to component (II) ranges from 4:1 to 1:4.
5. A composition as claimed in claim 4 wherein the ratio by weight of component (I) to component (II) ranges from 2:1 to 1:2.
- 20 6. A composition according to any of claims 1 to 5 wherein the amount of component (I) is present in a range from 10 to 50.000 mg/l and the amount of component (II) is present in a range from 10 to 50.000 mg/l.
- 25 7. Method of controlling microbial growth on wood, wood products and biodegradable materials, which comprises applying an antimicrobially effective amount of a composition as claimed in any of claims 1 to 6, to the wood, wood products and biodegradable materials to be treated.
- 30 8. Method of controlling microbial growth on engineering materials, which comprises applying an antimicrobially effective amount of a composition as claimed in any of claims 1 to 6, to the engineering materials to be treated.
- 35 9. A process for preparing a synergistic composition as claimed in any of claims 1 to 6, characterized in that the component (I) and one or more components (II) are intimately mixed with another.

10. A product containing
- (a) a composition comprising component (I), imazalil; and
  - (b) a composition comprising as a component (II), a hydroxypyridone selected from 1-hydroxy-2-pyridinone (II-a), ciclopirox (II-b), ciclopirox olamine (II-c),
- 5      piroctone (II-d), piroctone olamine (II-e) and rilopirox (II-f), as a combination for simultaneous or sequential use, wherein said compositions (a) and (b) are in respective proportions to provide a synergistic biocidal effect.

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2008/061312

## A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/4164 A61K31/4412 A61K45/06 A61P31/04 A61P31/10  
A61P31/12 A01N43/54 A01N43/40

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K A61P A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99/12422 A (JANSSEN PHARMACEUTICA NV [BE]; BOSSELAERS JAN PIETER HENDRIK [BE]; GAR) 18 March 1999 (1999-03-18) cited in the application the whole document	1-10
Y	WO 03/011030 A (BAYER CROPS SCIENCE SA [FR]; GARNIER ALAIN JOSEPH JEAN FLOR [BE]; VERBRU) 13 February 2003 (2003-02-13) cited in the application the whole document	1-10
Y	WO 2004/105662 A (PROGRAM FOR APPROPRIATE TECHNO [US]; CAMUS-BABLON FLORENCE [US]; ZANEV) 9 December 2004 (2004-12-09) abstract page 7, paragraphs 19,20	1-10
	-/--	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 November 2008

Date of mailing of the international search report

12/11/2008

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040,  
Fax: (+31-70) 340-3016

Authorized officer

Ansaldo, M

# INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2008/061312

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>KORTING H C ET AL: "The hydroxypyridones: A class of antimycotics of its own" MYCOSES, vol. 40, no. 7-8, November 1997 (1997-11), pages 243-247, XP009094077 ISSN: 0933-7407 the whole document</p>	1-10

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2008/061312

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9912422	A	18-03-1999	AT 241905 T	15-06-2003
			AU 741544 B2	06-12-2001
			AU 1145599 A	29-03-1999
			BG 64109 B1	30-01-2004
			BG 104203 A	31-08-2001
			BR 9812630 A	22-08-2000
			CA 2302428 A1	18-03-1999
			DE 69815363 D1	10-07-2003
			EE 200000099 A	15-12-2000
			HU 0003478 A2	28-02-2001
			NO 20001170 A	02-05-2000
			NZ 503118 A	25-05-2001
			PL 339175 A1	04-12-2000
			SK 2802000 A3	14-08-2000
			TR 200000623 T2	21-08-2000
			UA 62978 C2	16-10-2000
			US 6174911 B1	16-01-2001
			ZA 9808167 A	22-03-2000
WO 03011030	A	13-02-2003	AT 290786 T	15-04-2005
			BR 0211787 A	03-08-2004
			CA 2451865 A1	13-02-2003
			CN 1535112 A	06-10-2004
			DE 60203299 D1	21-04-2005
			DE 60203299 T2	19-01-2006
			EP 1411766 A1	28-04-2004
			ES 2236560 T3	16-07-2005
			FR 2828065 A1	07-02-2003
			HK 1067494 A1	06-10-2006
			HU 0401252 A2	28-09-2004
			MA 26140 A1	01-04-2004
			MX PA04000874 A	03-06-2004
			NZ 530348 A	28-10-2005
			PL 365478 A1	10-01-2005
			PT 1411766 T	30-06-2005
			US 2004167144 A1	26-08-2004
			ZA 200400700 A	15-10-2004
WO 2004105662	A	09-12-2004	NONE	