

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 June 2009 (11.06.2009)

PCT

(10) International Publication Number
WO 2009/073203 A1

(51) International Patent Classification:

C07D 217/04 (2006.01) *C07D 405/06* (2006.01)
C07D 217/06 (2006.01) *C07D 405/12* (2006.01)
C07D 217/22 (2006.01) *C07D 413/06* (2006.01)
C07D 401/04 (2006.01) *C07D 417/06* (2006.01)
C07D 401/06 (2006.01) *A61K 31/472* (2006.01)
C07D 401/10 (2006.01) *A61P 29/00* (2006.01)
C07D 401/12 (2006.01)

(US). **MONENSCHEN, Holger** [DE/US]; 711 Hunt Circle, Camarillo, California 93012 (US). **NGUYEN, Thomas** [US/US]; 468 Calle Veracruz, Thousand Oaks, California 91320 (US). **NISHIMURA, Nobuko** [JP/US]; 24050 Armita Street, West Hills, California 91304 (US). **TAMAYO, Nuria** [ES/US]; 4394 Camino de la Rosa, Newbury Park, California 91320 (US).

(21) International Application Number:

PCT/US2008/013375

(74) Agent: **PERSON, Richard V.**; Amgen Inc., M/s 28-2-c, One Amgen Center Drive, Thousand Oaks, California 91320-1799 (US).

(22) International Filing Date:

4 December 2008 (04.12.2008)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/005,457 4 December 2007 (04.12.2007) US
61/008,892 21 December 2007 (21.12.2007) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): **AMGEN INC.** [US/US]; One Amgen Center Drive, Thousand Oaks, California 91320-1799 (US).


Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report

(54) Title: TRP-M8 RECEPTOR LIGANDS AND THEIR USE IN TREATMENTS



(57) Abstract: Tetrahydroisoquinoline compounds of formula (I), and compositions containing them, for the treatment of acute, inflammatory and neuropathic pain, dental pain, general headache, migraine, cluster headache, mixed-vascular and non-vascular syndromes, tension headache, general inflammation, arthritis, rheumatic diseases, osteoarthritis, inflammatory bowel disorders, inflammatory eye disorders, inflammatory or unstable bladder disorders, psoriasis, skin complaints with inflammatory components, chronic inflammatory conditions, inflammatory pain and associated hyperalgesia and allodynia, neuropathic pain and associated hyperalgesia and allodynia, diabetic neuropathy pain,

causalgia, sympathetically maintained pain, deafferentation syndromes, asthma, epithelial tissue damage or dysfunction, herpes simplex, disturbances of visceral motility at respiratory, genitourinary, gastrointestinal or vascular regions, wounds, burns, allergic skin reactions, pruritus, vitiligo, general gastrointestinal disorders, gastric ulceration, duodenal ulcers, diarrhea, gastric lesions induced by necrotising agents, hair growth, vasomotor or allergic rhinitis, bronchial disorders or bladder disorders.

TRP-M8 RECEPTOR LIGANDS AND THEIR USE IN TREATMENTS

This application claims the benefit of two U.S. Provisional Application No.'s 61/005,457, filed December 4, 2007 and 61/008,892 filed December 21, 5 2007, which are hereby incorporated by reference.

Background

Cold sensation is derived from activation of the somatosensory system by a cold stimulus. Calcium imaging and patch clamp experiments in dissociated trigeminal and dorsal root ganglia neurons have revealed that cold stimuli induced 10 calcium influx, suggesting the direct opening of a calcium-permeable ion channels by cold (Thut et al., 2003; Reid, 2005). A recently cloned non-selective cation channel, TRPM8 (transient receptor potential melastatin 8) or trp-p8 (identified as a prostate-specific gene, up-regulated in prostate cancer and other malignancies, (Tsavaler et al., 2001)) is activated by cold stimulus of 10 to 24° C temperature 15 (McKemy et al., 2002; Peier et al., 2002). In addition, TRPM8 is also activated by compounds that elicit cool sensation such as menthol, icilin (AG-3-5) (McKemy et al., 2002), and the endogenous lipid PIP₂ (Rohacs et al., 2005). Correlating with the cold sensitivity of both A delta and C-fibers, TRPM8 is 20 highly expressed in sensory neurons of the trigeminal and dorsal root ganglia (McKemy et al., 2002; Peier et al., 2002; Thut et al., 2003). TRPM8 is also expressed in nerve fibers innervating urinary bladder in guinea pigs (Tsukimi et al., 2005) and humans (Mukerji et al., 2006) and believed to contribute to the bladder hypersensitivity.

Activation mechanism of TRPM8 by menthol and icilin appears to differ. 25 Icilin requires calcium for robust activation of TRPM8, whereas menthol and cold do not (Chuang et al., 2004). Typically, activation by all these agonists follows a period of calcium-dependent desensitization. The domain swap analysis of chicken and rat TRPM8 and further mutational studies revealed that determinants of icilin sensitivity map to a region of TRPM8 that corresponds to the capsaicin 30 binding site in TRPV1 transmembrane domain 3 to 4 region (Chuang et al., 2004).

Cold allodynia and mechanical hyperalgesia are associated with neuropathic pain in humans and in rodent models of neuropathic and

chemotherapy-induced pain. TRPM8 is shown to mediate the analgesia by agonists such as menthol and icilin (by desensitization of the receptor) during experimental neuropathic pain in rodents (Proudfoot et al., 2006). Further, attenuation of cold sensation and cold allodynia after chronic constriction injury 5 model of neuropathic pain in TRPM8 knockout mice (Colburn et al., 2007; Dhaka et al., 2007) suggests that antagonists of TRPM8 may be considered as pain therapeutics for chemotherapy-induced pain, neuropathic pain and bladder disorders.

Mint oil that contains menthol, an agonist of TRPM8 has been reported to 10 alleviate pain in post-herpetic neuralgia (Davies et al., 2002), a neuropathic pain condition. Furthermore, oral or intracerebroventricular injection of menthol decreased nociceptive responses to hot-plate test and acetic acid-induced writhing in mice (Galeotti et al., 2002). These responses are believed to be mediated by the activation and desensitization of the TRPM8. These observations and the 15 knockout mice studies indicate that TRPM8 modulation by antagonists might be beneficial for patients experiencing neuropathic pain.


Summary

The present invention comprises a new class of compounds useful in the 20 treatment of diseases, such as TRPM8-mediated diseases and other maladies, such as inflammatory or neuropathic pain and diseases involving sensory nerve function such as asthma, rheumatoid arthritis, osteoarthritis, inflammatory bowel disorders, urinary incontinence, migraine and psoriasis. In particular, the compounds of the invention are useful for the treatment of acute, inflammatory 25 and neuropathic pain, dental pain, general headache, migraine, cluster headache, mixed-vascular and non-vascular syndromes, tension headache, general inflammation, arthritis, rheumatic diseases, osteoarthritis, inflammatory bowel disorders, anxiety, depression, inflammatory eye disorders, inflammatory or unstable bladder disorders, psoriasis, skin complaints with inflammatory 30 components, chronic inflammatory conditions, inflammatory pain and associated hyperalgesia and allodynia, neuropathic pain and associated hyperalgesia and allodynia, diabetic neuropathy pain, causalgia, sympathetically maintained pain,

deafferentation syndromes, asthma, epithelial tissue damage or dysfunction, herpes simplex, disturbances of visceral motility at respiratory, genitourinary, gastrointestinal or vascular regions, wounds, burns, allergic skin reactions, pruritus, vitiligo, general gastrointestinal disorders, gastric ulceration, duodenal
5 ulcers, diarrhea, gastric lesions induced by necrotising agents, hair growth, vasomotor or allergic rhinitis, bronchial disorders or bladder disorders.

Accordingly, the invention also comprises pharmaceutical compositions comprising the compounds, methods for the treatment of TRP-M8-receptor-mediated diseases, such as inflammatory or neuropathic pain, asthma, rheumatoid
10 arthritis, osteoarthritis, inflammatory bowel disorders, urinary incontinence, migraine and psoriasis diseases, using the compounds and compositions of the invention, and intermediates and processes useful for the preparation of the compounds of the invention.

The compounds of the invention are represented by the following general
15 structure:




or a pharmaceutically acceptable salt thereof, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^f , J and X are defined below.

20 The foregoing merely summarizes certain aspects of the invention and is not intended, nor should it be construed, as limiting the invention in any way. All patents, patent applications and other publications recited herein are hereby incorporated by reference in their entirety.

Detailed Description

One aspect of the current invention relates to compounds having the general structure:



5 or any pharmaceutically-acceptable salt thereof, wherein:

J is $-N(R^a)(CR^cR^c)_n-$, $-O(CR^cR^c)_n-$, $-S(CR^cR^c)_n-$ or $-(CR^cR^c)_n-$;

m is 0, 1 or 2;

n is independently in each instance 0, 1, 2 or 3;

o is 0 or 1

10 X is O, NR^a or S;

R¹ is selected from H, halo, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$, $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$,

15 $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$ and $-NR^aC_{2-6}alkOR^a$; or R¹ is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,

C₁₋₄haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$,


20 $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$, $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$ and $-NR^aC_{2-6}alkOR^a$;

25 R² is selected from H, halo, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$, $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$ and $-NR^aC_{2-6}alkOR^a$; or

R^2 is C_{1-6} alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C_{1-6} alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C_{1-8} alk, C_{1-4} haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ;

5 R^3 is

10



; or

R^3 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, substituted with 0, 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F , Cl , Br , CN , OH and NH_2 ; or R^3 is benzyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F , Cl , Br , CN , OH and NH_2 ; or R^3 is napthyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F , Cl , Br , CN , OH and NH_2 ; or R^3 is benzyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F , Cl , Br , CN , OH and NH_2 ; or R^3 is napthyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F , Cl , Br , CN , OH and NH_2 ; or R^3 is benzyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F , Cl , Br , CN , OH and NH_2 ; or R^3 is napthyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F , Cl , Br , CN , OH and NH_2 ; or R^3 is C_{1-11} alkyl, C_{6-11} cycloalkyl or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C_{1-6} alkyl and ring are substituted by 0, 1 or 2 oxo groups and the C_{1-11} alkyl,


C₆₋₁₁cycloalkyl and ring are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a,

5 -NRR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or if J is -N(R^a)(CR^cR^c)_n-, -O(CR^cR^c)_n- or -S(CR^cR^c)_n-, then R⁵ may also be C₃₋₅cycloalkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or if J is -N(R^a)(CR^cR^c)_n-, then R⁵ may also be H;

10 15 R⁶ is independently in each instance, selected from F, Cl, Br, I, C₂₋₆alk, cyano, -OR^a, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁶ is C₁₋₆alk substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁶ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

20 25 30 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

- R^7 is selected from C_{1-6} alk, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)R^b$, $-N(R^a)S(=O)_2R^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ; or R^7 is C_{1-6} alk substituted by 1, 2 or 3 substituents selected from C_{1-4} haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)R^b$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ; or R^7 is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C_{1-8} alk, C_{1-4} haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)R^b$, $-N(R^a)S(=O)_2R^a$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ;
- R^a is independently, at each instance, H or R^b ;
- R^b is independently, at each instance, phenyl, benzyl or C_{1-6} alk, the phenyl, benzyl and C_{1-6} alk being substituted by 0, 1, 2 or 3 substituents selected from halo, C_{1-4} alk, C_{1-3} haloalk, $-OC_{1-4}$ alk, $-NH_2$, $-NHC_{1-4}$ alk, and $-N(C_{1-4}$ alk) C_{1-4} alk;
- R^c is independently, at each instance, H, halo, C_{1-4} alk, C_{1-4} haloalk, $-OC_{1-4}$ alk, $-OC_{1-4}$ haloalk, $-NH_2$, $-NHC_{1-4}$ alk or $-N(C_{1-4}$ alk) C_{1-4} alk; and
- R^f is independently in each instance H or F.
- Another aspect of the current invention relates to compounds having the general structure:



or any pharmaceutically-acceptable salt thereof, wherein:

J is $-N(R^a)(CR^cR^c)_n-$, $-O(CR^cR^c)_n-$, $-S(CR^cR^c)_n-$ or $-(CR^cR^c)_n-$;

m is 0, 1 or 2;

n is independently in each instance 0, 1, 2 or 3;

o is 0 or 1

5 X is O, NR^a or S;

R¹ is selected from H, halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,

-C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

-OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,

-N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,

10 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R¹ is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,

C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,

15 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

-OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,

-N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,

-N(R^a)S(=O)₂R^b, -N(R^a)S(=O)NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

R² is selected from H, halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,

20 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

-OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,

-N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,

-N(R^a)S(=O)₂R^b, -N(R^a)S(=O)NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R² is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,

25 C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,


-C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

-OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,

30 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,

-N(R^a)S(=O)₂R^b, -N(R^a)S(=O)NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

R³ is



; or

- R³ is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, substituted with 0, 1, 2 or 3 groups selected from
- 5 C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂; or R³ is benzyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂; or R³ is naphthyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk,
- 10 -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN and NH₂; or R³ is benzyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂;
- 15 R⁴ is selected from H, C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂;
- 15 R⁵ is C₁₋₁₁alkyl, C₆₋₁₁cycloalkyl or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alkyl and ring are substituted by 0, 1 or 2 oxo groups and the C₁₋₁₁alkyl,
- 20 C₆₋₁₁cycloalkyl and ring are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a,
- 25 -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or if J is -N(R^a)(CR^cR^c)_n-, then R⁵ may also be H;
- R⁶ is independently in each instance, selected from F, Cl, Br, I, C₂₋₆alk, cyano, -OR^a, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -S(=O)R^b,

$\text{-S(=O)}_2\text{R}^b$, $\text{-S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{R}^a$, $\text{-N(R}^a\text{)C(=O)R}^b$, $\text{-N(R}^a\text{)C(=O)OR}^b$,
 $\text{-N(R}^a\text{)C(=O)NR}^a\text{R}^a$, $\text{-N(R}^a\text{)C(=NR}^a\text{)NR}^a\text{R}^a$, $\text{-N(R}^a\text{)S(=O)}_2\text{R}^b$,
 $\text{-N(R}^a\text{)S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{C}_{2-6}\text{alkNR}^a\text{R}^a$ and $\text{-NR}^a\text{C}_{2-6}\text{alkOR}^a$; or R^6 is C_{1-6}alk
 substituted by 1, 2 or 3 substituents selected from $\text{C}_{1-4}\text{haloalk}$, halo, cyano, nitro,
5 -C(=O)R^b , -C(=O)OR^b , $\text{-C(=O)NR}^a\text{R}^a$, $\text{-C(=NR}^a\text{)NR}^a\text{R}^a$, -OR^a , -OC(=O)R^b ,
 $\text{-OC(=O)NR}^a\text{R}^a$, $\text{-OC}_{2-6}\text{alkNR}^a\text{R}^a$, $\text{-OC}_{2-6}\text{alkOR}^a$, -SR^a , -S(=O)R^b , $\text{-S(=O)}_2\text{R}^b$,
 $\text{-S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{R}^a$, $\text{-N(R}^a\text{)C(=O)R}^b$, $\text{-N(R}^a\text{)C(=O)OR}^b$, $\text{-N(R}^a\text{)C(=O)NR}^a\text{R}^a$,
 $\text{-N(R}^a\text{)C(=NR}^a\text{)NR}^a\text{R}^a$, $\text{-N(R}^a\text{)S(=O)}_2\text{R}^b$, $\text{-N(R}^a\text{)S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{C}_{2-6}\text{alkNR}^a\text{R}^a$
 and $\text{-NR}^a\text{C}_{2-6}\text{alkOR}^a$; or R^6 is a saturated, partially saturated or unsaturated 5-, 6-
10 or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S,
 wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C_{1-8}alk ,
 $\text{C}_{1-4}\text{haloalk}$, halo, cyano, nitro, -C(=O)R^b , -C(=O)OR^b , $\text{-C(=O)NR}^a\text{R}^a$,
 $\text{-C(=NR}^a\text{)NR}^a\text{R}^a$, -OR^a , -OC(=O)R^b , $\text{-OC(=O)NR}^a\text{R}^a$, $\text{-OC}_{2-6}\text{alkNR}^a\text{R}^a$,
 $\text{-OC}_{2-6}\text{alkOR}^a$, -SR^a , -S(=O)R^b , $\text{-S(=O)}_2\text{R}^b$, $\text{-S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{R}^a$,
15 $\text{-N(R}^a\text{)C(=O)R}^b$, $\text{-N(R}^a\text{)C(=O)OR}^b$, $\text{-N(R}^a\text{)C(=O)NR}^a\text{R}^a$, $\text{-N(R}^a\text{)C(=NR}^a\text{)NR}^a\text{R}^a$,
 $\text{-N(R}^a\text{)S(=O)}_2\text{R}^b$, $\text{-N(R}^a\text{)S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{C}_{2-6}\text{alkNR}^a\text{R}^a$ and $\text{-NR}^a\text{C}_{2-6}\text{alkOR}^a$;
 R^7 is selected from C_{1-6}alk , cyano, -C(=O)R^b , -C(=O)OR^b , $\text{-C(=O)NR}^a\text{R}^a$,
 $\text{-C(=NR}^a\text{)NR}^a\text{R}^a$, -S(=O)R^b , $\text{-S(=O)}_2\text{R}^b$, $\text{-S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{R}^a$, $\text{-N(R}^a\text{)C(=O)R}^b$,
 $\text{-N(R}^a\text{)C(=O)OR}^b$, $\text{-N(R}^a\text{)C(=O)NR}^a\text{R}^a$, $\text{-N(R}^a\text{)C(=NR}^a\text{)NR}^a\text{R}^a$, $\text{-N(R}^a\text{)S(=O)}_2\text{R}^b$,
20 $\text{-N(R}^a\text{)S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{C}_{2-6}\text{alkNR}^a\text{R}^a$ and $\text{-NR}^a\text{C}_{2-6}\text{alkOR}^a$; or R^7 is C_{1-6}alk
 substituted by 1, 2 or 3 substituents selected from $\text{C}_{1-4}\text{haloalk}$, halo, cyano, nitro,
 -C(=O)R^b , -C(=O)OR^b , $\text{-C(=O)NR}^a\text{R}^a$, $\text{-C(=NR}^a\text{)NR}^a\text{R}^a$, -OR^a , -OC(=O)R^b ,
 $\text{-OC(=O)NR}^a\text{R}^a$, $\text{-OC}_{2-6}\text{alkNR}^a\text{R}^a$, $\text{-OC}_{2-6}\text{alkOR}^a$, -SR^a , -S(=O)R^b , $\text{-S(=O)}_2\text{R}^b$,
 $\text{-S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{R}^a$, $\text{-N(R}^a\text{)C(=O)R}^b$, $\text{-N(R}^a\text{)C(=O)OR}^b$, $\text{-N(R}^a\text{)C(=O)NR}^a\text{R}^a$,
25 $\text{-N(R}^a\text{)C(=NR}^a\text{)NR}^a\text{R}^a$, $\text{-N(R}^a\text{)S(=O)}_2\text{R}^b$, $\text{-N(R}^a\text{)S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{C}_{2-6}\text{alkNR}^a\text{R}^a$
 and $\text{-NR}^a\text{C}_{2-6}\text{alkOR}^a$; or R^7 is a saturated, partially saturated or unsaturated 5-, 6-
 or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S,
 wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C_{1-8}alk ,
 $\text{C}_{1-4}\text{haloalk}$, halo, cyano, nitro, -C(=O)R^b , -C(=O)OR^b , $\text{-C(=O)NR}^a\text{R}^a$,
30 $\text{-C(=NR}^a\text{)NR}^a\text{R}^a$, -OR^a , -OC(=O)R^b , $\text{-OC(=O)NR}^a\text{R}^a$, $\text{-OC}_{2-6}\text{alkNR}^a\text{R}^a$,
 $\text{-OC}_{2-6}\text{alkOR}^a$, -SR^a , -S(=O)R^b , $\text{-S(=O)}_2\text{R}^b$, $\text{-S(=O)}_2\text{NR}^a\text{R}^a$, $\text{-NR}^a\text{R}^a$,

-N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;


R^a is independently, at each instance, H or R^b;

5 R^b is independently, at each instance, phenyl, benzyl or C₁₋₆alk, the phenyl, benzyl and C₁₋₆alk being substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄alk, C₁₋₃haloalk, -OC₁₋₄alk, -NH₂, -NHC₁₋₄alk, and -N(C₁₋₄alk)C₁₋₄alk;

R^c is independently, at each instance, H, halo, C₁₋₄alk, C₁₋₄haloalk, -OC₁₋₄alk, -OC₁₋₄haloalk, -NH₂, -NHC₁₋₄alk or -N(C₁₋₄alk)C₁₋₄alk; and

R^f is independently in each instance H or F.

10 Another aspect of the current invention relates to compounds having the general structure:



or any pharmaceutically-acceptable salt thereof, wherein:

J is -N(R^a)(CR^cR^c)_n-, -O(CR^cR^c)_n-, -S(CR^cR^c)_n or -(CR^cR^c)_n;

15 m is 0, 1 or 2;

n is independently in each instance 0, 1, 2 or 3;


o is 0 or 1

X is O, NR^a or S;

20 R¹ is selected from H, halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R¹ is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

- OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,
 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;
 R² is selected from H, halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
 5 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,
 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,
 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or
 R² is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered
 10 ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk
 and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,
 C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,
 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,
 15 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

R³ is



; or

- R³ is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered
 20 monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 1, 2, 3 or 4
 atoms selected from N, O and S, substituted with 0, 1, 2 or 3 groups selected from
 C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk,
 -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂; or R³ is benzyl substituted with 1,
 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk,
 25 -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂; or
 R³ is napthyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk,
 -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F,
 Cl, Br, CN and NH₂; or R³ is benzyl substituted with 1, 2 or 3 groups selected

from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂;

R⁴ is selected from H, C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂;

R⁵ is C₁₋₁₁alkyl, C₆₋₁₁cycloalkyl or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alkyl and ring are substituted by 0, 1 or 2 oxo groups and the C₁₋₁₁alkyl, C₆₋₁₁cycloalkyl and ring are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or if J is -N(R^a)(CR^cR^c)_n-, then R⁵ may also be H;

R⁶ is independently in each instance, selected from F, Cl, Br, I, C₂₋₆alk, cyano, -OR^a, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁶ is C₁₋₆alk substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁶ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,

- N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;
- R⁷ is selected from C₁₋₆alk, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
 -C(=NR^a)NR^aR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b,
- 5 -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b,
 -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁷ is C₁₋₆alk
 substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro,
 -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b,
 -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b,

10 -S(=O)NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a,
 -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a
 and -NR^aC₂₋₆alkOR^a; or R⁷ is a saturated, partially saturated or unsaturated 5-, 6-
 or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S,
 wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,

15 C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,
 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)NR^aR^a, -NR^aR^a,
 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

20 R^a is independently, at each instance, H or R^b;


R^b is independently, at each instance, phenyl, benzyl or C₁₋₆alk, the phenyl,
 benzyl and C₁₋₆alk being substituted by 0, 1, 2 or 3 substituents selected from
 halo, C₁₋₄alk, C₁₋₃haloalk, -OC₁₋₄alk, -NH₂, -NHC₁₋₄alk, and -N(C₁₋₄alk)C₁₋₄alk;

R^c is independently, at each instance, H, halo, C₁₋₄alk, C₁₋₄haloalk,

25 -OC₁₋₄alk, -OC₁₋₄haloalk, -NH₂, -NHC₁₋₄alk or -N(C₁₋₄alk)C₁₋₄alk; and

R^f is independently in each instance H or F.

Another aspect of the current invention relates to compounds having the general structure:




or any pharmaceutically-acceptable salt thereof, wherein:

- 5 J is $-N(R^a)(CR^cR^c)_n-$, $-O(CR^cR^c)_n-$, $-S(CR^cR^c)_n-$ or $-(CR^cR^c)_n-$;
 m is 0, 1 or 2;
 n is independently in each instance 0, 1, 2 or 3;
 o is 0 or 1
 X is O, NR^a or S;
- 10 R¹ is selected from H, halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,
 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,
 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or
- 15 R¹ is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,
 C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,
- 20 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,
 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;
- 25 R² is selected from H, halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
 -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,
 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,
 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or
 R² is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered

ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

5 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

R³ is



10 R³ is selected from 4-pyrimidinyl, 2-pyridinyl, both of which are substituted with 0, 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂; or R³ is benzyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk,

15 -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂;


R⁴ is selected from H, C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂;

R⁵ is C₁₋₁₁alkyl, C₆₋₁₁cycloalkyl or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered

20 bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alkyl and ring are substituted by 0, 1 or 2 oxo groups and the C₁₋₁₁alkyl, C₆₋₁₁cycloalkyl and ring are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a,

25 -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or if J is -N(R^a)(CR^cR^c)_n-, then R⁵ may also be H;


- R^6 is selected from C_{2-6} alk, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)R^b$, $-N(R^a)S(=O)NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ; or R^6 is C_{1-6} alk substituted by 1, 2 or 3 substituents selected from C_{1-4} haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)R^b$, $-N(R^a)S(=O)_2R^b$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ; or R^6 is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C_{1-6} alk is and C_{1-6} alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C_{1-8} alk, C_{1-4} haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)R^b$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ;
- R^a is independently, at each instance, H or R^b ;
- R^b is independently, at each instance, phenyl, benzyl or C_{1-6} alk, the phenyl, benzyl and C_{1-6} alk being substituted by 0, 1, 2 or 3 substituents selected from halo, C_{1-4} alk, C_{1-3} haloalk, $-OC_{1-4}$ alk, $-NH_2$, $-NHC_{1-4}$ alk, and $-N(C_{1-4}$ alk) C_{1-4} alk;
- R^c is independently, at each instance, H, halo, C_{1-4} alk, C_{1-4} haloalk, $-OC_{1-4}$ alk, $-OC_{1-4}$ haloalk, $-NH_2$, $-NHC_{1-4}$ alk or $-N(C_{1-4}$ alk) C_{1-4} alk; and
- R^f is independently in each instance H or F.
- In one embodiment, in conjunction with any above or below embodiments, J is $-N(R^a)-$, $-O-$, $-S-$ or $-CH_2-$.
- In another embodiment, in conjunction with any above or below embodiments, o is 1.
- In another embodiment, in conjunction with any above or below embodiments, R³ is



In another embodiment, in conjunction with any above or below embodiments, R³ is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, substituted with 0, 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂; or R³ is benzyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂.

In another embodiment, in conjunction with any above or below embodiments, R³ is napthyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN and NH₂.


In another embodiment, in conjunction with any above or below embodiments, R³ is



R⁶ is C₁₋₆alk substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁶ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or

4 atoms selected from N, O and S, wherein the C₁₋₆alk is and C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, 5 -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R³ is



10

R⁶ is C₁₋₆alk substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, 15 -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁶ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk is and C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, 20 cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, m is 0.

In another embodiment, in conjunction with any above or below embodiments, m is 1 or 2; and R⁴ is independently in each instance selected from 5 CH₃, CF₃, F, Cl or -CH₂CH₂- (forming a spiro cyclopropyl).

In another embodiment, in conjunction with any above or below embodiments, R⁵ is C₁₋₁₁alkyl or C₆₋₁₁cycloalkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, 10 -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below 15 embodiments, R⁵ is C₁₋₁₁alkyl or C₅₋₁₁cycloalkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, 20 -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, J is -N(R^a)(CR^cR^c)_n-, -O(CR^cR^c)_n- or -S(CR^cR^c)_n-; and R⁵ is 25 C₃₋₅cycloalkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

30 In another embodiment, in conjunction with any above or below embodiments, J is -N(R^a)(CR^cR^c)_n-; and R⁵ is C₃₋₅cycloalkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b,

-C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R⁵ is cyclopropyl substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R⁵ is C₃₋₆alkyl.

In another embodiment, in conjunction with any above or below embodiments, R⁵ is C₃₋₆alkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R⁵ is C₅₋₁₁cycloalkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R⁵ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the ring is substituted by 0, 1

or 2 oxo groups and the ring is additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a,

5 -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R⁵ is an unsaturated 6-membered monocyclic ring containing 1 or 2 N atoms and the ring is substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,

10 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R⁵ is phenyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,

15 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, J is -N(R^a)(CR^cR^c)_n-.

25 In another embodiment, in conjunction with any above or below embodiments, J is -O(CR^cR^c)_n-.

In another embodiment, in conjunction with any above or below embodiments, J is -NR^a-.

30 In another embodiment, in conjunction with any above or below embodiments, J is -O-.

In another embodiment, in conjunction with any above or below embodiments, J is -CH₂-.

In another embodiment, in conjunction with any above or below embodiments, R¹ is selected from halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, 5 -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R¹ is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected 10 from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a 15 and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R¹ is H or F.

In another embodiment, in conjunction with any above or below embodiments, R¹ is H.

20 In another embodiment, in conjunction with any above or below embodiments, R² is selected from halo, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a 25 and -NR^aC₂₋₆alkOR^a; or R² is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, 30 -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a,

-N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

In another embodiment, in conjunction with any above or below embodiments, R² is H or F.

5 In another embodiment, in conjunction with any above or below embodiments, R² is H.

In another embodiment, in conjunction with any above or below embodiments, R^f is H.

10 In another embodiment, in conjunction with any above or below embodiments, R³ is selected from 4-pyrimidinyl, 2-pyridinyl, both of which are substituted with 0, 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂.

15 In another embodiment, in conjunction with any above or below embodiments, R³ is benzyl substituted with 1, 2 or 3 groups selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂.

20 In another embodiment, in conjunction with any above or below embodiments, m is 1; and R⁴ is selected from C₁₋₆alk, C₁₋₃haloalk, -OC₁₋₆alk, -N(C₁₋₆alk)C₁₋₆alk, -NHC₁₋₆alk, -NC(=O)C₁₋₆alk, F, Cl, Br, CN, OH and NH₂;

25 In another embodiment, in conjunction with any above or below embodiments, R⁶ is selected from C₃₋₆alk, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

30 In another embodiment, in conjunction with any above or below embodiments, R⁶ is C₁₋₆alk substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,

-N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
-N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

- In another embodiment, in conjunction with any above or below
embodiments, R⁶ is a saturated, partially saturated or unsaturated 5-, 6- or
5 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S,
wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,
C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a,
-C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,
-OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a,
10 -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
-N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

Another aspect of the invention relates to a method of treating acute,
inflammatory and neuropathic pain, dental pain, general headache, migraine,
cluster headache, mixed-vascular and non-vascular syndromes, tension headache,
15 general inflammation, arthritis, rheumatic diseases, osteoarthritis, inflammatory
bowel disorders, depression, anxiety, inflammatory eye disorders, inflammatory or
unstable bladder disorders, psoriasis, skin complaints with inflammatory
components, chronic inflammatory conditions, inflammatory pain and associated
hyperalgesia and allodynia, neuropathic pain and associated hyperalgesia and
20 allodynia, diabetic neuropathy pain, causalgia, sympathetically maintained pain,
deafferentation syndromes, asthma, epithelial tissue damage or dysfunction,
herpes simplex, disturbances of visceral motility at respiratory, genitourinary,
gastrointestinal or vascular regions, wounds, burns, allergic skin reactions,
pruritus, vitiligo, general gastrointestinal disorders, gastric ulceration, duodenal
25 ulcers, diarrhea, gastric lesions induced by necrotising agents, hair growth,
vasomotor or allergic rhinitis, bronchial disorders or bladder disorders, comprising
the step of administering a compound according to Claim 1.

Another aspect of the invention relates to a pharmaceutical composition
comprising a compound according to Claim 1 and a pharmaceutically-acceptable
30 diluent or carrier.


Another aspect of the invention relates to the use of a compound according
to any of the above embodiments as a medicament.

Another aspect of the invention relates to the use of a compound according to any of the above embodiments in the manufacture of a medicament for the treatment of acute, inflammatory and neuropathic pain, dental pain, general headache, migraine, cluster headache, mixed-vascular and non-vascular syndromes, tension headache, general inflammation, arthritis, rheumatic diseases, osteoarthritis, inflammatory bowel disorders, anxiety, depression, inflammatory eye disorders, inflammatory or unstable bladder disorders, psoriasis, skin complaints with inflammatory components, chronic inflammatory conditions, inflammatory pain and associated hyperalgesia and allodynia, neuropathic pain and associated hyperalgesia and allodynia, diabetic neuropathy pain, causalgia, sympathetically maintained pain, deafferentation syndromes, asthma, epithelial tissue damage or dysfunction, herpes simplex, disturbances of visceral motility at respiratory, genitourinary, gastrointestinal or vascular regions, wounds, burns, allergic skin reactions, pruritus, vitiligo, general gastrointestinal disorders, gastric ulceration, duodenal ulcers, diarrhea, gastric lesions induced by necrotising agents, hair growth, vasomotor or allergic rhinitis, bronchial disorders or bladder disorders.

The compounds of this invention may have in general several asymmetric centers and are typically depicted in the form of racemic mixtures. This invention is intended to encompass racemic mixtures, partially racemic mixtures and separate enantiomers and diasteromers.

Unless otherwise specified, the following definitions apply to terms found in the specification and claims:

“C_{α-β}alk” means an alkyl group comprising a minimum of α and a maximum of β carbon atoms in a branched, cyclical or linear relationship or any combination of the three, wherein α and β represent integers. The alkyl groups described in this section may also contain one or two double or triple bonds. A designation of C₀alk indicates a direct bond. Examples of C₁₋₆alkyl include, but are not limited to the following:



Where the term “C_{α-β}alkyl” and “C_{α-β}cycloalkyl” are used, they relate to acyclic saturated alkyls and cyclic saturated alkyls, respectively.

- “Benzo group”, alone or in combination, means the divalent radical C₄H₄=, one representation of which is -CH=CH-CH=CH-, that when vicinally attached to 5 another ring forms a benzene-like ring--for example tetrahydronaphthylene, indole and the like.


The terms “oxo” and “thioxo” represent the groups =O (as in carbonyl) and =S (as in thiocarbonyl), respectively.

“Halo” or “halogen” means a halogen atoms selected from F, Cl, Br and I.

- 10 “C_v-whaloalk” means an alk group, as described above, wherein any number--at least one--of the hydrogen atoms attached to the alk chain are replaced by F, Cl, Br or I.


The group N(R^a)R^a and the like include substituents where the two R^a groups together form a ring, optionally including a N, O or S atom, and include groups

- 15 such as:




The group N(C_{α-β}alk)C_{α-β}alk, wherein α and β are as defined above, include substituents where the two C_{α-β}alk groups together form a ring, optionally including a N, O or S atom, and include groups such as:

20



“Heterocycle” means a ring comprising at least one carbon atom and at least one other atom selected from N, O and S. Examples of heterocycles that may be found in the claims include, but are not limited to, the following:



5

and



"Pharmaceutically-acceptable salt" means a salt prepared by conventional means, and are well known by those skilled in the art. The "pharmacologically acceptable salts" include basic salts of inorganic and organic acids, including but not limited to hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, malic acid, acetic acid, oxalic acid, tartaric acid, citric acid, lactic acid, fumaric acid, succinic acid, maleic acid, salicylic acid, benzoic acid, phenylacetic acid, mandelic acid and the like. When

- compounds of the invention include an acidic function such as a carboxy group, then suitable pharmaceutically acceptable cation pairs for the carboxy group are well known to those skilled in the art and include alkaline, alkaline earth, ammonium, quaternary ammonium cations and the like. For additional examples
- 5 of "pharmacologically acceptable salts," *see infra* and Berge et al., J. Pharm. Sci. 66:1 (1977).
- "Saturated, partially-saturated or unsaturated" includes substituents saturated with hydrogens, substituents completely unsaturated with hydrogens and substituents partially saturated with hydrogens.
- 10 "Leaving group" generally refers to groups readily displaceable by a nucleophile, such as an amine, a thiol or an alcohol nucleophile. Such leaving groups are well known in the art. Examples of such leaving groups include, but are not limited to, N-hydroxysuccinimide, N-hydroxybenzotriazole, halides, triflates, tosylates and the like. Preferred leaving groups are indicated herein where appropriate.
- 15 "Protecting group" generally refers to groups well known in the art which are used to prevent selected reactive groups, such as carboxy, amino, hydroxy, mercapto and the like, from undergoing undesired reactions, such as nucleophilic, electrophilic, oxidation, reduction and the like. Preferred protecting groups are indicated herein where appropriate. Examples of amino protecting groups include, but are not
- 20 limited to, aralkyl, substituted aralkyl, cycloalkenylalkyl and substituted cycloalkenyl alkyl, allyl, substituted allyl, acyl, alkoxycarbonyl, aralkoxycarbonyl, silyl and the like. Examples of aralkyl include, but are not limited to, benzyl, ortho-methylbenzyl, trityl and benzhydryl, which can be optionally substituted with halogen, alkyl, alkoxy, hydroxy, nitro, acylamino, acyl and the like, and salts, such
- 25 as phosphonium and ammonium salts. Examples of aryl groups include phenyl, naphthyl, indanyl, anthracenyl, 9-(9-phenylfluorenyl), phenanthrenyl, durenyl and the like. Examples of cycloalkenylalkyl or substituted cycloalkenylalkyl radicals, preferably have 6-10 carbon atoms, include, but are not limited to, cyclohexenyl methyl and the like. Suitable acyl, alkoxycarbonyl and aralkoxycarbonyl groups
- 30 include benzyloxycarbonyl, t-butoxycarbonyl, iso-butoxycarbonyl, benzoyl, substituted benzoyl, butyryl, acetyl, trifluoroacetyl, trichloro acetyl, phthaloyl and the like. A mixture of protecting groups can be used to protect the same amino


group, such as a primary amino group can be protected by both an aralkyl group and an aralkoxycarbonyl group. Amino protecting groups can also form a heterocyclic ring with the nitrogen to which they are attached, for example, 1,2-bis(methylene)benzene, phthalimidyl, succinimidyl, maleimidyl and the like and where these heterocyclic groups can further include adjoining aryl and cycloalkyl rings. In addition, the heterocyclic groups can be mono-, di- or tri-substituted, such as nitrophthalimidyl. Amino groups may also be protected against undesired reactions, such as oxidation, through the formation of an addition salt, such as hydrochloride, toluenesulfonic acid, trifluoroacetic acid and the like. Many of the amino protecting groups are also suitable for protecting carboxy, hydroxy and mercapto groups. For example, aralkyl groups. Alkyl groups are also suitable groups for protecting hydroxy and mercapto groups, such as tert-butyl.

Silyl protecting groups are silicon atoms optionally substituted by one or more alkyl, aryl and aralkyl groups. Suitable silyl protecting groups include, but are not limited to, trimethylsilyl, triethylsilyl, triisopropylsilyl, tert-butyldimethylsilyl, dimethylphenylsilyl, 1,2-bis(dimethylsilyl)benzene, 1,2-bis(dimethylsilyl)ethane and diphenylmethylsilyl. Silylation of an amino groups provide mono- or di-silylamino groups. Silylation of aminoalcohol compounds can lead to a N,N,O-trisilyl derivative. Removal of the silyl function from a silyl ether function is readily accomplished by treatment with, for example, a metal hydroxide or ammonium fluoride reagent, either as a discrete reaction step or in situ during a reaction with the alcohol group. Suitable silylating agents are, for example, trimethylsilyl chloride, tert-butyl-dimethylsilyl chloride, phenyldimethylsilyl chloride, diphenylmethyl silyl chloride or their combination products with imidazole or DMF. Methods for silylation of amines and removal of silyl protecting groups are well known to those skilled in the art. Methods of preparation of these amine derivatives from corresponding amino acids, amino acid amides or amino acid esters are also well known to those skilled in the art of organic chemistry including amino acid/amino acid ester or aminoalcohol chemistry.

Protecting groups are removed under conditions which will not affect the remaining portion of the molecule. These methods are well known in the art and

include acid hydrolysis, hydrogenolysis and the like. A preferred method involves removal of a protecting group, such as removal of a benzyloxycarbonyl group by hydrogenolysis utilizing palladium on carbon in a suitable solvent system such as an alcohol, acetic acid, and the like or mixtures thereof. A t-
 5 butoxycarbonyl protecting group can be removed utilizing an inorganic or organic acid, such as HCl or trifluoroacetic acid, in a suitable solvent system, such as dioxane or methylene chloride. The resulting amino salt can readily be neutralized to yield the free amine. Carboxy protecting group, such as methyl, ethyl, benzyl, tert-butyl, 4-methoxyphenylmethyl and the like, can be removed
 10 under hydrolysis and hydrogenolysis conditions well known to those skilled in the art.

It should be noted that compounds of the invention may contain groups that may exist in tautomeric forms, such as cyclic and acyclic amidine and guanidine groups, heteroatom substituted heteroaryl groups ($Y' = O, S, NR$), and
 15 the like, which are illustrated in the following examples:



and though one form is named, described, displayed and/or claimed herein, all the tautomeric forms are intended to be inherently included in such name, description, display and/or claim.

20 Prodrugs of the compounds of this invention are also contemplated by this invention. A prodrug is an active or inactive compound that is modified

- chemically through in vivo physiological action, such as hydrolysis, metabolism and the like, into a compound of this invention following administration of the prodrug to a patient. The suitability and techniques involved in making and using prodrugs are well known by those skilled in the art. For a general discussion of
- 5 prodrugs involving esters see Svensson and Tunek Drug Metabolism Reviews 165 (1988) and Bundgaard Design of Prodrugs, Elsevier (1985). Examples of a masked carboxylate anion include a variety of esters, such as alkyl (for example, methyl, ethyl), cycloalkyl (for example, cyclohexyl), aralkyl (for example, benzyl, p-methoxybenzyl), and alkylcarbonyloxyalkyl (for example, pivaloyloxymethyl).
- 10 Amines have been masked as arylcarbonyloxymethyl substituted derivatives which are cleaved by esterases in vivo releasing the free drug and formaldehyde (Bungaard J. Med. Chem. 2503 (1989)). Also, drugs containing an acidic NH group, such as imidazole, imide, indole and the like, have been masked with N-acyloxymethyl groups (Bundgaard Design of Prodrugs, Elsevier (1985)).
- 15 Hydroxy groups have been masked as esters and ethers. EP 039,051 (Sloan and Little, 4/11/81) discloses Mannich-base hydroxamic acid prodrugs, their preparation and use.

The specification and claims contain listing of species using the language like “selected from . . . and . . .” and “is . . . or . . .” (sometimes referred to as

20 Markush groups). When this language is used in this application, unless otherwise stated it is meant to include the group as a whole, or any single members thereof, or any subgroups thereof. The use of this language is merely for shorthand purposes and is not meant in any way to limit the removal of individual elements or subgroups as needed.

Experimental


Unless otherwise noted, all materials were obtained from commercial suppliers and used without further purification. All parts are by weight and temperatures are in degrees centigrade unless otherwise indicated. All microwave assisted reactions were conducted with a Smith Synthesizer from Biotage. Mass spectral data was determined by electrospray ionization technique. All examples were purified to >90% purity as determined by high-performance liquid chromatography. Unless otherwise stated, reactions were run at RT.

10 The following abbreviations are used:


DCE	dichloroetane
DCM	dichloromethane
DIAE	diisopropyl ethylamine
DMSO -	dimethyl sulfoxide
15 DMF -	<i>N,N</i> -dimethylformamide
THF -	tetrahydrofuran
Et ₂ O -	diethyl ether
EtOAc -	ethyl acetate
MeOH -	methyl alcohol
20 EtOH -	ethyl alcohol
MeCN -	acetonitrile
MeI -	iodomethane
NMP -	1-methyl-2-pyrrolidinone
TFA -	trifluoroacetic acid
25 SFC -	supercritical fluid chromatography
h -	h
min -	min
mL	milliliters
μL	microliters

g	grams
μg	micrograms
mg	milligrams
μmoL	micromolars


5

Scheme 1

10




Example 1: Ethyl 1-(4-(dimethylamino)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate



15 Step 1. **Ethyl 1-isoquinoline-carboxylate chloride**


Isoquinoline (5.50 g, 42.5 mmol) was dissolved into 50 mL of anhydrous THF and ethyl chloroformate (4.47 mL, 46.8 mmol) was added slowly. The resulting pale yellow mixture was stirred at RT for 30 min. Then, the resulting precipitate was collected via filtration through a cintered glass funnel and dried under vacuum for

20 2 days to yield the title compound as a pale yellow solid.



Step 2. Ethyl 1-(4-(dimethylamino)phenyl)isoquinoline-2(1*H*)-carboxylate


- 5 To a 25-mL round-bottomed flask containing the above ethyl 1-isoquinoline-carboxylate chloride (0.584 g, 2.46 mmol) in anhydrous THF (10 mL) was added with 4-*N,N*-dimethylaminophenylmagnesium bromide (0.5 M solution in THF, 7.50 mL, 3.75 mmol) under a stream of N₂ and the mixture was stirred at RT for 1.5 h. This mixture was quenched with saturated NH₄Cl and extracted with EtOAc. The organic layer was washed with water and brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. The crude product was purified by silica gel chromatography (0-50% EtOAc in hexanes) to give the title compound as pale yellow oil. MS (ESI pos. ion) *m/z*: 323 (M+1).
- 10




15

Step 3. Ethyl 1-(4-(dimethylamino)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

- A solution of ethyl 1-(4-(dimethylamino)phenyl)isoquinoline-2(1*H*)-carboxylate (79 mg, 0.24 mmol) in EtOH (4 mL) was stirred with 10% Pd/C (0.062 g, 0.58 mmol) under hydrogen atmosphere at RT for 14 h. The reaction mixture was filtered through a celite pad and the filtrate was concentrated in vacuo. The crude product was purified by silica gel chromatography (0-30% EtOAc in hexanes) to provide the title compound as a clear oil. MS (ESI pos. ion) *m/z*: 325 (M+1).
- 20




Example 2: Ethyl 1-(4-biphenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate



Step 1. Ethyl 1-(4-biphenyl)isoquinoline-2(1*H*)-carboxylate

- 5 A round-bottomed flask containing isoquinoline (0.545 g, 4.22 mmol) in anhydrous THF (10 mL) was charged with ethyl chloroformate (0.450 mL, 4.71 mmol) under a stream of N₂ and the mixture was stirred at RT for 15 min. 4-Biphenylmagnesium bromide (0.5 M solution in THF, 16.0 mL, 8.0 mmol) was added dropwise and the mixture was stirred at RT for 1.5 h, this mixture was quenched with saturated NH₄Cl and extracted with EtOAc. The organic layer was washed with water and brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. The crude product was purified by silica gel chromatography (0-20% EtOAc in hexanes) to give ethyl 1-(4-biphenyl)isoquinoline-2(1*H*)-carboxylate as a white solid. MS (ESI pos. ion) *m/z*: 356 (M+1).
- 10




15

Step 2. Ethyl 1-(4-biphenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

A solution of ethyl 1-(4-biphenyl)isoquinoline-2(1*H*)-carboxylate (1.05 g, 2.94 mmol) in EtOH (10 mL) and EtOAc (7 mL) was stirred with 10% Pd/C (0.272 g, 2.56 mmol) under hydrogen atmosphere at RT for 14 h. Then the mixture was transferred to a pressure tube and hydrogenated at 46 psi of hydrogen for 4 h. The reaction mixture was filtered through a celite pad and the filtrate was concentrated in vacuo. The crude product was purified by silica gel chromatography (0-20% EtOAc in hexanes) to provide the title compound as a clear oil. MS (ESI pos. ion) *m/z*: 358 (M+1).

10




15

Step 1. Ethyl 1-(4-(trifluoromethyl)phenyl)isoquinoline-2(1*H*)-carboxylate

The 4-trifluoromethylphenyl Grignard reagent was prepared by adding 1-bromo-4-(trifluoromethyl)benzene (1.3 mL, 9.2 mmol) to a suspension of magnesium turnings (240 mg, 10.0 mmol), a catalytic amount of iodine in THF (10 mL) and stirred at RT for 1.5 h. A separate round-bottomed flask containing isoquinoline (1.1 g, 8.3 mmol) in anhydrous THF (10 mL) was charged with ethyl chloroformate (0.88 mL, 9.2 mmol) under a stream of N₂ and the mixture was stirred at RT for 15 min, and then cooled to 0 °C. The previously made Grignard reagent was then cannulated into this solution dropwise and the reaction mixture was stirred for 1 h at 0 °C followed by 1 h at RT. The reaction mixture was quenched with saturated NH₄Cl and extracted with EtOAc. The organic layer was


washed with brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. The crude product was purified by silica gel chromatography (10% EtOAc in hexanes) to give ethyl 1-(4-(trifluoromethyl)phenyl)isoquinoline-2(1*H*)-carboxylate as a white solid. MS (ESI pos. ion) *m/z*: 348 (M+1).



5

Step 2. Ethyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

A solution of ethyl 1-(4-(trifluoromethyl)phenyl)isoquinoline-2(1*H*)-carboxylate (2.0 g, 5.7 mmol) in EtOH (20 mL) was stirred with 10% Pd/C (0.6 g, 5.7 mmol) under a hydrogen atmosphere at RT for 2 h. The reaction mixture was filtered through a celite pad and the filtrate was concentrated in vacuo to provide the title compound as colorless oil. MS (ESI pos. ion) *m/z*: 350 (M+1).




Example 4:

15

(1*S*,3*R*)-*N*-(4-Fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide and (1*R*,3*S*)-*N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]


20



Step1: (4-(Trifluoromethyl)phenyl)magnesium bromide

To a solution of magnesium turnings (238 mg, 9.8 mmol) and iodine (10 mg) in THF (20 mL) under N_2 was added 4-bromobenzotrifluoride (1.24 mL, 8.9 mmol).

After the addition was completed, the mixture was stirred at RT for 2 h to give the title compound, which was used in the next step without further purification.




**Step 2: 2-Benzyl-3-methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydro-
5 isoquinoline**

To a solution of 3-methylisoquinoline (1.0 g, 6.98 mmol) in THF (6 mL) was added benzyl bromide (0.78 mL, 6.63 mmol). After the addition was completed, the mixture was heated at 70 °C under N₂ for 16 h. The mixture was cooled down to RT and THF (10 mL). Then, the mixture was cooled to 0 °C and (4-
10 (trifluoromethyl)phenyl)magnesium bromide was added, and then stirred at RT for 16 h. After quenching with saturated ammonium chloride solution, the mixture was extracted several times with EtOAc. The combined organic extracts were dried over MgSO₄ and concentrated. The residue was mixed with silica gel and the solid mixture was purified by silica gel flash column chromatography (30%-
15 100% EtOAc/hexane, then 20% MeOH in DCM) to give a mixture of 2-benzyl-3-methyl-1-(4-(trifluoromethyl)phenyl)-1,2-dihydroisoquinoline & 2-benzyl-3-methyl-1-(4-(trifluoromethyl)phenyl)-isoquinolinium, which was used without further purification.

This mixture was dissolved in THF (45 mL) and sodium borohydride (2.6 g, 68.9
20 mmol) was added. The resulting mixture was stirred at RT 15 min, and then glacial acetic acid (1 mL) was added. The mixture was stirred at RT for 2 h, and then at 55 °C for an additional 2 h. After cooling down to RT, it was basified with NaOH (10 N) (to pH>12). The organic layer was collected and the aqueous layer was extracted several times with EtOAc. The combined organic extracts were then dried over MgSO₄ and concentrated. The residue was dissolved MeOH (5 mL) and the solution mixture was purified by preparative HPLC (0% MeCN 0.1% TFA/ H₂O 0.1% TFA). All fractions of the product were combined and MeCN was removed in vacuo. The aqueous phase was basified with NaHCO₃ and


extracted with EtOAc. The combined organic extracts were dried over MgSO₄, concentrated, and dried to give the title compound as a yellow oil.



Step 3: **3-Methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline hydrogen chloride**


To a solution of 2-benzyl-3-methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (485 mg, 1.27 mmol) in MeOH (5 mL) was added palladium, 10 wt. % on activated carbon (41 mg, 0.38 mmol). The resulting mixture was then stirred under H₂ for 16 h. Then, 5 N HCl (0.2 mL) was added followed by palladium, 10wt. % on activated carbon (41 mg, 0.38 mmol). The resulting mixture was stirred at RT under H₂ (45 psi) for 30 min. The mixture was filtered through celite and the celite was washed with MeOH (2 x 5 mL). The combined filtrates were concentrated to give the title compound as a light yellow solid (HCl salt), which was used in the next step without further purification.

15



Step 4: **N-(4-Fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**


To a solution of 3-methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline hydrogen chloride (370 mg, 1.27 mmol) and DIEA (1.1 mL, 6.35 mmol) in DCM (5 mL) was added 4-fluorophenyl isocyanate (143 µL, 1.27 mmol). The mixture was then stirred at RT for 0.5 h. Water (10 mL) was added and the mixture was extracted with DCM (2 x 15 mL). The combined organic extracts were dried over MgSO₄ and concentrated to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 429 (M+H).



Step 5: **(1*S*,3*R*)-*N*-(4-Fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide and (1*R*,3*S*)-*N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]**

Isomers of *N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide were separated by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the title compounds as a light yellow solid. MS (ESI, positive ion) m/z: 429 (M+H).

10




10

**Example 5: (1*R*,3*R*)-*N*-(4-Fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide
(1*S*,3*S*)-*N*-(4-Fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]**


Isomers of *N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (example 4) were separated by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the title compounds as a white solid. MS (ESI, positive ion) m/z: 429 (M+H).

20

**Example 6:**



(1*R*,3*S*)-*N*-(4-Fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]

Enantiomers, (1*S*,3*R*)-*N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide & (1*R*,3*S*)-*N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (example 4) were separated using chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a colorless solid. MS (ESI, positive ion) m/z: 429 (M+H).

**Example 7:**

(1*R*,3*R*)-*N*-(4-Fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]


Enantiomers (1*R*,3*R*)-*N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide & (1*S*,3*S*)-*N*-(4-fluorophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (example 5) were separated using chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a colorless solid. MS (ESI, positive ion) m/z: 429 (M+H).

**Example 8:****8-Fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**


5

Step 1: 2-Bromo-6-fluorobenzaldehyde

A solution of diisopropylamine (7.9 mL, 55 mmol) in anhydrous tetrahydrofuran (70 mL) cooled to 0 °C was treated dropwise with butyllithium (1.6 M in hexanes, 10 34.9 mL, 56 mmol). The mixture was stirred at 0 °C for 15 min, then it was cooled to -78 °C, and 1-bromo-3-fluoro-benzene (6.1 mL, 56 mmol) was added over 10 min. The mixture was stirred at -78 °C for 1 h, then anhydrous *N,N*-dimethylformamide (4.9 mL, 64 mmol) was added dropwise over 5 min. The mixture was stirred for another 20 min, then acetic acid (8 mL) and water (180 mL) were added, warmed up to RT, and extracted with ethyl acetate (2 x 175 mL). The combined organic layers were washed with water (2 x 75 mL), brine (100 mL), dried over anhydrous sodium sulfate, and concentrated to yield the crude product which was purified by silica gel chromatography (20-30% EtOAc in hexanes) to give 2-bromo-6-fluorobenzaldehyde as a pale yellow solid. MS (ESI, pos. ion) *m/z*: 204 (M+1).

**Step 2: 2-Fluoro-6-(2-(trimethylsilyl)ethynyl)benzaldehyde**

A solution of 2-bromo-6-fluorobenzaldehyde (3.3 g, 16 mmol) and (trimethylsilyl)acetylene (4.5 mL, 32 mmol) in anhydrous *N,N*-dimethylformamide (60 mL) was treated with triethylamine (3.5 mL, 25 mmol), copper(I) iodide (0.3 g, 1.6 mmol), and Pd(*PPh*₃)₂Cl₂ (0.6 g, 0.80 mmol) under 5 argon atmosphere. The mixture was stirred at RT for 2.5 h. The mixture was poured into water (150 mL) and extracted with ethyl acetate (2 x 150 mL). The combined organic layers were washed with water (2 x 75 mL), brine (100 mL) then dried over anhydrous sodium sulfate and evaporated to yield the crude product which was purified by silica gel chromatography (20-30% EtOAc in 10 hexanes) to give 2-fluoro-6-(2-(trimethylsilyl)ethynyl)benzaldehyde 2-bromo-6-fluorobenzaldehyde as a brown oil. MS (ESI, pos. ion) *m/z*: 221 (M+1).




Step 3: 8-Fluoroisoquinoline

15 2-Fluoro-6-(2-(trimethylsilyl)ethynyl)benzaldehyde (2.3 g, 10.5 mmol) was dissolved in ammonia (2M in methanol, 15 mL, 30 mmol) and the resulting mixture was heated at 80 °C in a sealed tube for 2.5 h. The solvent was removed under high vacuum and the resulting crude product purified by silica gel chromatography (20-30% EtOAc in hexanes) to give 8-fluoroisoquinoline as a 20 pale yellow solid. MS (ESI, pos. ion) *m/z*: 148 (M+1).




Step 4: 2-Benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-isoquinolium bromide

25 A solution of 8-fluoroisoquinoline (0.4 g, 2.7 mmol) in acetonitrile (5 mL) was treated with benzyl bromide (0.37 mL, 3.0 mmol) and the solution was refluxed for 3 h. The reaction mixture was cooled down to RT and concentrated to give 2-benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-isoquinolium bromide as a pale yellow solid. MS (ESI, pos. ion) *m/z*: 238 (M⁺).



Step 5: 2-Benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2-dihydroisoquinoline

- 5 The 4-trifluoromethylphenyl Grignard reagent was prepared by adding 1-bromo-4-(trifluoromethyl)benzene (1.2 mL, 8.3 mmol) to a suspension of magnesium turnings (0.3 g, 11 mmol) and catalytic amount of iodine in THF (10 mL) at RT. A different round-bottomed flask containing 2-benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-isoquinolium bromide (0.87 g, 2.7 mmol) in anhydrous tetrahydrofuran (15 mL) was cooled to 0 °C, and to this flask was added the Grignard reagent, (4-(trifluoromethyl)phenyl)magnesium bromide over a period of 5 min. The reaction mixture was allowed to stir at 0 °C for 1 h after which the cooling bath was removed and allowed to stir the reaction at RT for 16 h. The reaction was quenched with saturated ammonium chloride (35 mL), followed by 10 ethyl acetate (125 mL), stirred the resulting mixture for 10 min, then transferred to a separatory funnel, added ethyl acetate (100 mL), water (50 mL), brine (50 mL). The ethyl acetate layer was separated, dried over anhydrous sodium sulfate and concentrated to yield 2-benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2-dihydroisoquinoline as a brown oil. MS (ESI, pos. ion) *m/z*: 384 (M+1).
- 15




- 20 Step 6: 2-Benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline

A 250-mL round-bottomed flask containing a solution of 2-benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2-dihydroisoquinoline (1.0 g, 2.6 mmol) in anhydrous tetrahydrofuran (20 mL) was treated with sodium borohydride (0.3 g, 8.4 mmol)

25


at RT. After 20 min, acetic acid (3.3 mL) was added and the reaction was stirred at RT for 2.5 h. The reaction mixture was concentrated and the resulting residue was taken up in ethyl acetate (150 mL), water (25 mL), 1N sodium hydroxide, and brine (35 mL). The ethyl acetate layer was separated, dried over anhydrous sodium sulfate, and concentrated to yield the crude product. The crude product was purified by silica gel chromatography (20-30% EtOAc in hexanes) to give 2-benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline as a colorless oil. MS (ESI, pos. ion) m/z : 386 (M+1).



10


Step 7: 8-Fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline


A solution of 2-benzyl-8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.7 g, 2.1 mmol) in EtOH (20 mL) was stirred with 20% Pd(OH)₂/C (0.20 g) under hydrogen atmosphere at 50 psi and at RT for 3 h. The reaction mixture was filtered through a celite pad and the filtrate was concentrated in vacuo. The residue was purified by preparative HPLC [gradient 10-90 % MeCN (0.1 % TFA)/ H₂O (0.1 % TFA)] to give the pure product which was dissolved in methanol (10 mL) and neutralized by passing the solution through a Polymer Lab-HCO₃ Macroporous resin cartridge, and the filtrate was concentrated to give 8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline as an off-white solid. MS (ESI pos. ion) m/z : 296 (M+1).




Step 8: 8-Fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

A solution of 8-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline e (0.4 g, 1.4 mmol) and diisopropylethylamine (0.5 mL, 2.9 mmol) in anhydrous 1,2-dichloroethane (10 mL) was treated with 4-fluorophenyl isocyanate (0.14 mL, 1.2 mmol) and stirred at RT for 16 h. The solvent was removed under 5 vacuum and the residue was purified by preparative HPLC [gradient 10-90 % MeCN (0.1 % TFA)/ H₂O (0.1 % TFA)] to give the pure product which was dissolved in methanol (10 mL) and neutralized by passing the solution through a Polymer Lab-HCO₃ Macroporous resin cartridge, and the filtrate was concentrated to give 8-fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 433 10 (M+1).

Scheme 2

**Example 9:**

***N*-Phenyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**




5

Step 1. *N*-Phenethyl-4-(trifluoromethyl)benzamide


To a solution of phenylethylamine (14.0 g, 14.6 mL, 0.11 mol) and DIEA (20.1 mL, 0.11 mol in DCM (500 mL) at 0 °C was added 4-(trifluoromethyl)benzoyl chloride (17.2 mL, 0.11 mol) dropwise. After the addition was completed, the reaction mixture was stirred at 0 °C and warmed up itself to RT for 16 h. An off-white solid precipitated was observed. The off-white solid was collected by filtration. The solid was then washed with DCM (2 x 50 mL), and dried under vacuum to give the title compound as a white solid. MS (ESI, positive ion) m/z: 294 (M+H).

15

**Step 2: 1-(4-(Trifluoromethyl)phenyl)-3,4-dihydroisoquinoline**


To a 100-mL round-bottomed flask was added *N*-phenethyl-4-(trifluoromethyl)benzamide (5.0 g, 17 mmol), phosphoric pentoxide (0.526 mL, 8.52 mmol), and polyphosphoric acid (60.0 g). The reaction mixture was heated to 165 °C for 2 h. Then, the hot solution was carefully poured into ice/water and a solution of KOH (20%, 40 mL) was added to break up the oil. The mixture was

stirred at RT for 5 min and an additional amount of KOH solution was added until pH 7. Ethyl ether (250 mL) was added and the mixture was stirred at RT for 15 min. The organic layer was collected and the aqueous layer was extracted with EtOAc (2 x 200 mL). The combined organic extracts were dried over MgSO₄ and 5 concentrated in vacuo to give the title compound as orange oil, which was used in the next step without further purification. MS (ESI, positive ion) m/z: 276 (M+H).



Step 3: 1-(4-(Trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline


To a solution of 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (4.69 g, 17 mmol) in MeOH (24 mL) was added sodium borohydride (1.94 g, 51.1 mmol) 10 slowly at 0 °C . Then, the mixture was stirred at 0 °C for 15 min and at RT for 2 h. The solvent was then removed and H₂O (20 mL) was added to the residue. A saturated NaHCO₃ solution (150 mL) was added slowly and the mixture was extracted with EtOAc (2 x 200 mL). The combined organic extracts were dried 15 over MgSO₄ and concentrated. The residue was mixed with silica gel (4:1 ; residue:silica gel) and purified by silica gel chromatography (20%-100% EtOAc/hexane) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 278 (M+H).



20 **Step 4. N-Phenyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (500 mg, 1.80 mmol) and DIEA (0.314 mL, 1.80 mmol) in DCM (9 mL) was added phenyl isocyanate (0.196 mL, 1.80 mmol). The resulting mixture was stirred at


RT for 3 h. Then, MeOH (50 mL) was added and the mixture was concentrated. The residue was purified by silica gel flash column chromatography (0%-50% EtOAc/hexane) to give the title compound as a light yellow solid. The solid was then washed with MeOH (3 x 20 mL) to give 440 mg of the title compound as a white solid. MS (ESI, positive ion) m/z: 397 (M+H).
5



Example 10:

2-Phenyl-1-(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)ethanone

10 To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.36 mmol, example 9 (step 3) and DIEA (62.8 μ L, 0.36 mmol) in DCM (2 mL) was added phenylacetyl chloride (47.7 μ L, 0.36 mmol). The resulting mixture was stirred at RT for 16 h. Then, H₂O (3 mL) was added and the mixture was extracted with DCM (2 x 5 mL). The combined organic extracts were
15 concentrated and the residue was dissolved in MeOH (1 mL). The solution mixture was purified by preparative HPLC (10-100% of MeCN (0.1%TFA)/H₂O (0.1% TFA) to give the title compound as an off-white thick oil. MS (ESI, positive ion) m/z: 396 (M+H).




20

Example 11:

Phenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate


To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.36 mmol, example 9 (step 3) and DIEA (62.8 μ L, 0.36 mmol) in DCM (2 mL) was added phenyl chloroformate (45.2 μ L, 0.36 mmol). The resulting mixture was reacted under the same conditions as described for example 10 to give 112 mg (78% yield) of the title compound as an off-white oil. MS (ESI, positive ion) m/z: 398 ($M+H$).



Example 12:

10 ***N*-Benzyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.36 mmol, example 9 (step 3) and DIEA (62.8 μ L, 0.36 mmol) in DCM (2 mL) was added benzyl isocyanate (44.2 μ L, 0.36 mmol). The resulting mixture was reacted under the same conditions as described for example 10 to give the title compound as a white solid. MS (ESI, positive ion) m/z: 411 ($M+H$).




Example 13:

20 ***N*-Cyclohexyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.36 mmol, example 9 (step 3) and DIEA (63 μ L, 0.36 mmol) in DCM (2 mL) was added isocyanatocyclohexane (46 μ L, 0.36 mmol). The resulting


mixture was reacted under the same conditions as described for example 10 give the title compound as a white solid. MS (ESI, positive ion) m/z: 403 (M+H).



Example 14:


- 5 **N-(4-Fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.36 mmol, example 9 (step 3) and DIEA (62.8 μ L, 0.36 mmol) in DCM (2 mL) was added 4-fluorophenyl isocyanate (40.5 μ L, 0.36 mmol). The resulting mixture was reacted under the same conditions as described for example 10 to give the title compound as a white solid. MS (ESI, positive ion) m/z: 415 (M+H).



Example 15: 2-(4-Fluorophenyl)-1-(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1*H*)-yl)ethanone

15 To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.36 mmol, example 9 (step 3) and DIEA (63 μ L, 0.36 mmol) in DCM (2 mL) was added 4-fluorophenylacetyl chloride (62 μ L, 0.36 mmol). The resulting mixture was reacted under the same conditions as described for example 10 give the title compound as a white oil. MS (ESI, positive ion) m/z: 414 (M+H).




Example 16:


4-Nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (200 mg, 0.721 mmol, example 9 (step 3) and DIEA (126 µL, 0.721 mmol) in DCM (4 mL) was added 4-nitrophenyl chloroformate (145 mg, 0.721 mmol). The resulting mixture was stirred at RT for 16 h. Then, H₂O (10 mL) was added and the mixture was extracted with DCM (2 x 5 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was purified by silica gel flash column chromatography (0%-50% EtOAc/hexane) to give the title compound as a white oil. MS (ESI, positive ion) m/z: 443 (M+H).


15

**Example 17: N-Isopropyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

To a solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (70 mg, 0.252 mmol, example 9 (step 3) and DIEA (44 µL, 0.252 mmol) in DCM (1.6 mL) was added 2-isocyanatopropane (25 µL, 0.252 mmol). The resulting mixture was reacted under the same conditions as described for example 10 to give the title compound as a white solid. MS (ESI, positive ion) m/z: 363 (M+H).


**Example 18:****1-(4-(Trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

- A solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.5 g, 5 mmol) (Example 9, step 3) in dichloromethane (10 mL) was treated with pyridine (1.4 mL, 17 mmol), triethylamine (0.5 mL, 4 mmol), potassium cyanate (0.3 g, 3 mmol), glacial acetic acid (1.00 mL, 17 mmol) and the resulting suspension was stirred at RT for 12 h. The reaction mixture was concentrated to a gummy residue which was taken up in dichloromethane (75 mL), washed with brine (35 mL), dried over anhydrous sodium sulfate, removed dichloromethane, and dried under high vacuum to yield (4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 321 (M+1).

15 **Example 19:****1-(4-(Trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carbothioamide**

- A solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.5 g, 2 mmol) (Example 9, step 3) in dry tetrahydrofuran (10 mL) was treated with 1,1'-thiocarbonyldiimidazole (0.5 g, 3 mmol) and stirred at RT for 4 h. The solvent was removed and the resulting residue was dissolved in ammonia (2.0 M solution in methanol, 15 mL, 30 mmol) and stirred the resulting solution in a pressure tube at RT for 1 h followed by heating at 55 °C for 5 h. The solvent was removed

under vacuum and the residue was purified by preparative HPLC [gradient 10-90 % MeCN (0.1 % TFA)/ H₂O (0.1 % TFA)] to give 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carbothioamide as a white solid. MS (ESI, pos. ion) *m/z*: 337 (M+1).




5

Example 20:

10 **(3*R*)-*tert*-Butyl 3-(1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroiso-
quinoline-2-carboxamido)piperidine-1-carboxylate**


A mixture of 4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroiso-
quinoline-2(1*H*)-carboxylate (350 mg, 0.79 mmol, example 15) and (*R*)-1-Boc-3-
amino-piperidine (1.6 mL, 7.91 mmol) was subjected to a microwave irradiation
15 at 120 °C for 45 min. Then, MeOH (2 mL) was added to the mixture and the
mixture was filtered. The solution mixture was purified by preparative HPLC
(10%-100% of MeCN (0.1%TFA)/H₂O (0.1% TFA)). All fractions of the product
were combined, saturated NaHCO₃ (1 mL) was added and the solvents were
removed. Then, an additional amount of saturated NaHCO₃ (10 mL) was added
20 and the aqueous mixture was extracted with EtOAc (2 x 10 mL). The combined
organic extracts were dried over MgSO₄ and concentrated to give the title
compound as a white solid. MS (ESI, positive ion) *m/z*: 504 (M+H).

**Example 21:**

***tert*-Butyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-**

5 **carboxylate**

To a mixture of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.361 mmol, example 9 (step 3) and DIEA (62.8 μL , 0.361 mmol), and DCM (1 mL) was added di-*tert*-butyldicarbonate (82.9 μL , 0.361 mmol). The resulting mixture was stirred at RT for 16 h. Then, H_2O (1 mL) was added and the mixture was extracted with DCM (2 x 2 mL). The combined organic extracts were dried over MgSO_4 and concentrated. The residue was purified by silica gel chromatography (0%-100% EtOAc/hexane) to give the title compound as an off-white oil. MS (ESI, positive ion) m/z: 322 (M-*tert*-butyl).




15

Example 22:

***N*-Methyl-*N*-phenyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

To a solution of *N*-phenyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (100 mg, 0.252 mmol, example 9 (step 3) in THF (1 mL) was added sodium hydride (12 mg, 0.505 mmol) slowly. The resulting mixture was stirred at RT for 15 min and iodomethane (16 μL , 0.252 mmol) was added. The mixture was stirred at RT for 16 h. Then, H_2O (2 mL) was added and the mixture was extracted with EtOAc (2 x 2 mL). The combined organic extracted were


dried over MgSO₄ and concentrated. The residue was then dissolved in DMSO (1 mL) and the solution mixture was purified by preparative HPLC (10%-100% of MeCN (0.1%TFA)/H₂O (0.1% TFA) to give the title compound as an off-white solid. MS (ESI, positive ion) m/z: 411 (M+H).



5

Example 23:(1*R*,2*S*,5*R*)-2-Isopropyl-5-methylcyclohexyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

A solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.361 mmol, example 9 (step 3) and (1*R*,2*S*,5*R*)-2-isopropyl-5-
10 methylcyclohexyl carbonochloride (76.5 μL, 0.361 mmol) in DCM (1 mL) was stirred at RT for 16 h. Then, H₂O (2 mL) was added and the mixture was extracted with EtOAc (2 x 5 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was purified by silica gel flash column chromatography (0%-100% EtOAc/hexane) to give the title compound as an off-
15 white oil. MS (ESI, positive ion) m/z: 460 (M+H).




Example 24:

N-((R)-Piperidin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

20 A solution of (3*R*)-*tert*-butyl 3-(1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-2-carboxamido)piperidine-1-carboxylate (44 mg, 87 μmol, example 20) in TFA (30% in DCM, 2 mL) was stirred at RT for 30 min. The solvents were removed and EtOAc (5 mL) and saturated NaHCO₃ (3 mL) were added to the residue. The solution mixture was then stirred at RT for 10 min. The organic

layer was collected and the aqueous layer was extracted with EtOAc (2 x 2 mL). The combined organic extracts were dried over MgSO₄, concentrated, and dried under vacuum to give the title compound as an off-white solid. MS (ESI, positive ion) m/z: 404 (M+H).




5


Example 25:

2-Hydroxy-2-methyl-1-(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)propan-1-one


To a solution of (4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (240 mg, 0.87 mmol, example 9 (step 3) and DIEA (151 µL, 0.87 mmol) in DCM (3 mL) was added 1-chlorocarbonyl-1-methylethyl acetate (125 µL, 0.87 mmol) slowly. After addition, the mixture was stirred at RT for 16 h. Then, an additional amount of 1-chlorocarbonyl-1-methylethyl acetate (60 µL) was added and the mixture was stirred at RT for 16 h. Then, H₂O (2 mL) was added and the mixture was stirred at RT for 10 min. The organic layer was collected and the aqueous layer was extracted with EtOAc (2 x 3 mL). The combined organic extracts were then dried over MgSO₄ and concentrated to give a crude intermediate. Then, the crude intermediate was dissolved in MeOH (10 mL), and then potassium carbonate (239 mg, 1.7 mmol) was added. The resulting mixture was then stirred at RT for 16 h. Then, the mixture was filtered and the solid was washed with DCM (2 x 5 mL). The combined filtrates were concentrated and the residue was purified by silica gel flash column chromatography (0%-100% EtOAc/hexane) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 364 (M+H).

**Example 26:*****N*-(Pyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

- 5 A solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (97 mg, 0.350 mmol, example 9 (step 3), DIEA (61 μ L, 0.35 mmol), and 3-isocyanatopyridine (42 mg, 0.35 mmol) in DCM (1 mL) was stirred at RT for 16 h. A white precipitate was observed. The reaction mixture was then filtered and the solid was dried *in vacuo* to give the title compound as a white solid. MS (ESI, positive ion) m/z: 398 ($M+H$).
- 10

**Example 27: (R)-*N*-(Pyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**


- Enantiomers of *N*-(pyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (1.3 g, racemic mixture) were separated using chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 398 ($M+H$).
- 15



- 20 **Example 28:**

(S)-N-(Pyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide


Enantiomers of *N*-(pyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (1.3 g, racemic mixture) were separated by using 5 chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 398 (M+H).



10 **Example 29:**


(R)-N-(4-Fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Enantiomers of *N*-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (100 mg, racemic mixture) were separated using 15 chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 415 (M+H).




20 **Example 30:**

(R)-N-(Pyridin-2-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide



Step 1: (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline


Enantiomers of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (2.120 g, 7.46 mmol, racemic mixture) were separated using chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 278 (M+H).



Step 2: 4-Nitrophenyl pyridin-2-ylcarbamate


10 To a solution of 2-aminopyridine (2.0 mL, 21.25 mmol) in DCM (35 mL) at 0 °C was added a solution of 4-nitrophenyl chloroformate (4.28g, 21.25 mmol) in DCM (35 mL) and pyridine (1.73 mL, 21.25 mmol, anhydrous). After the addition was completed, the reaction mixture was stirred at RT for 16 h under an atmosphere of nitrogen. A white precipitated was observed and it was collected by filtration.

15 The filtrate was washed with DCM (3 x 50 mL) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 260 (M+H).




Step 3: (*R*)-N-(Pyridin-2-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

To a solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.361 mmol) in MeCN (2 mL) was added 4-nitrophenyl pyridin-2-ylcarbamate (467 mg, 1.80 mmol). The resulting mixture was then heated at 70 °C for 16 h, cooled to RT and filtered. The white solid obtained was washed with DCM (2 x 2 mL). The combined filtrates were then concentrated and the residue purified by silica gel flash column chromatography (0%-100% EtOAc/hexane) to give the title compound as a colorless solid. MS (ESI, positive ion) m/z: 398 (M+H).




10 **Example 31:**

(*R*)-*N*-(Pyridin-4-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(*1H*)-carboxamide




Step 1: 4-Nitrophenyl pyridin-4-ylcarbamate

15 To a solution of 4-aminopyridine (1.97 g, 20.9 mmol) in DCM (75 mL) at 0 °C was added 4-nitrophenyl chloroformate (4.21 g, 20.9 mmol) and pyridine (1.7 mL, 20.9 mmol). Then, the mixture was stirred at RT overnight. The white precipitated was filtered off and washed with DCM (2 x 50 mL), and then dried in vacuo to give the title compound as a white solid, which was used in the next step without further purification. MS (ESI, positive ion) m/z: 260 (M+H).



Step 2: (*R*)-*N*-(Pyridin-4-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

To a solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (200 mg, 0.721 mmol) in MeCN (3 mL) was added 4-nitrophenyl pyridin-4-ylcarbamate (561 mg, 2.164 mmol). The resulting mixture was heated to 55 °C overnight. After cooling down to RT the white precipitated obtained was filtered off and washed with DCM (2 x 1 mL). The combined filtrates were then concentrated and the residue purified by silica gel flash column chromatography (0%-100% EtOAc/hexane) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 398 (M+H).




Example 32:


***N*-((*R*)-1-Methylpiperidin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

To a solution of *N*-((*R*)-piperidin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (30 mg, 74 µmol, example 24) in DCE (0.5 mL) was added formaldehyde (28 µL, 372 µmol). The resulting mixture was stirred at RT for 1 h and then sodium triacetoxyborohydride (16 mg, 74 µmol) was added.

The mixture was then stirred at RT for 4 h. Then, saturated NaHCO₃ (0.4 mL) was added and the mixture was extracted with EtOAc (2 x 3 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was purified by silica gel flash column chromatography (0%-10% MeOH (2M NH₃)/EtOAc) to give the title compound as an off-white solid. MS (ESI, positive ion) m/z: 418 (M+H).

**Example 33:**


- (*R*)-2-Methyl-1-oxo-1-(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1*H*)-yl)propan-2-yl acetate
- To a solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (207 mg, 747 µmol, example 30 (step 1) and DIEA (130 µL, 747 µmol) in DCM (2.5 mL) was added 1-chlorocarbonyl-1-methylethyl acetate (119 µL, 821 µmol). The mixture was stirred at RT for 3 h. Then, 1-chlorocarbonyl-1-methylethyl acetate (50µL) was added and the mixture was stirred at RT for 16 h. MeOH (1 mL) was added and the mixture was filtered. The filtrate was purified by preparative HPLC (0%-100% MeCN 0.1% TFA/water 0.1% TFA) to give the title compound in a solution of H₂O 0.1% TFA and MeCN 0.1% TFA. The solvent was removed and saturated NaHCO₃ (5 mL) was added. The mixture was extracted with EtOAc (2 x 30 mL). The combined organic extracts were dried over MgSO₄, concentrated, and dried in vacuo to give the title compound as a white solid. MS (ESI, positive ion) m/z: 406 (M+H).

**Example 34:**

- (*R*)-1-(1-(4-(Trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1*H*)-yl)ethanone


To a solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 361 µmol, example 30 (step 1) in DCM (2 mL) was added DIEA (62.8

μL , 361 μmol) and acetic acid anhydride (34.0 μL , 361 μmol). The resulting mixture was then stirred at RT for 16 h. Then, H_2O (1 mL) was added and the mixture was extracted with DCM (2 x 5 mL). The combined organic extracts were dried over MgSO_4 and concentrated. The residue was purified by silica gel 5 flash column chromatography (0%-100% EtOAc/hexane) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 320 ($\text{M}+\text{H}$).




Example 35:

10 (R) -*N*-(Pyrimidin-5-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroiso-
quinoline-2(1*H*)-carboxamide




Step 1: 4-Nitrophenyl pyrimidin-5-ylcarbamate

To a solution of pyrimidin-5-amine (190 mg, 2.0 mmol) in pyrimidine (3 mL) was added 4-nitrophenyl chloroformate (403 mg, 2.0 mmol). The resulting mixture 15 was stirred at RT overnight, and then heated to 60 °C overnight. After cooling to RT, the solvent was removed under vacuum to give the title compound as a brown solid, which was used in the next step without further purification. MS (ESI, positive ion) m/z: 261 ($\text{M}+\text{H}$).




20 Step 2: (R) -*N*-(Pyrimidin-5-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroiso-
quinoline-2(1*H*)-carboxamide

- A solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (100 mg, 0.30 mmol, example 30 (step 1)) and 4-nitrophenyl pyrimidin-5-ylcarbamate (93.8 mg, 0.36 mmol) in MeCN (3 mL) was heated to 70 °C overnight, and then was subjected to microwave irradiation at 120 °C for 15 min.
- 5 The mixture was filtered and the filtrate was purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the title compound in solution of H₂O (0.1% TFA) and MeCN (0.1% TFA). The solvent was partially removed and saturated NaHCO₃ (10 mL) was added. The mixture was extracted with EtOAc (2 x 5 mL). The combined organic extracts were dried over MgSO₄, concentrated,
- 10 and dried in vacuo to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 399 (M+H).




Example 36:

- 15 *N*-Isopropyl-4,4-dimethyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide




Step 1. 2-Methyl-2-phenylpropanenitrile

- A solution of phenylacetonitrile (5.0 mL, 43 mmol) and sodium hydroxide (7 g, 20 0.17 mol) in dimethylsulfoxide (40 mL):water (7 mL) was cooled to 0 °C and then treated with methyl iodide (11 mL, 0.17 mol). After stirring at RT for 1 h, the reaction mixture was quenched with water (15 mL) and extracted with EtOAc. The organic layer was washed with water, brine, dried over MgSO₄, filtered, and concentrated in vacuo to provide 2-methyl-2-phenylpropanenitrile as a yellow oil.




Step 2. 2-Methyl-2-phenylpropan-1-amine hydrochloride

- To a stirred suspension of lithium aluminum hydride (2.44 g, 64.2 mmol) in dry ether (100 mL) at RT was added under a nitrogen atmosphere 2-methyl-2-phenylpropanenitrile (5.83 g, 40.2 mmol) in dry ether (50 mL) over 30 min. Then, it was heated under reflux for 3 h and allowed to cool to RT. The reaction mixture was then treated sequentially with water (2.3 mL), NaOH (5 N, 2.3 mL) and water (6.9 mL), stirred at RT for 1.5 h and the solids were filtered off. The filtrate was dried over Na_2SO_4 and the solids were removed by filtration. Then the solution was treated with 1N HCl in ether (40 mL) and the white precipitated was separated by filtration. 2-Methyl-2-phenylpropan-1-amine hydrochloride was obtained as a white solid.




Step 3: *N*-(2-Methyl-2-phenylpropyl)-4-(trifluoromethyl)benzamide

- To a suspension of 2-methyl-2-phenylpropan-1-amine hydrochloride (5.32 g, 28.6 mmol) in dry DCM (60 mL) at 0 °C was added triethylamine (8.3 mL, 60.1 mmol) and then slowly 4-(trifluoromethyl)benzoyl chloride (4.48 mL, 30.1 mmol). Then, the mixture was stirred at RT for 45 min, diluted with EtOAc, washed with 10% HCl, sat NaHCO_3 and water. After drying over Na_2SO_4 the solvent was removed under vacuum and the title compound was obtained as a white solid. MS (ESI, positive ion) m/z: 322 ($\text{M}+\text{H}$).



Step 4: 4,4-Dimethyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline


A 100-mL round-bottomed flask was charged with *N*-(2-methyl-2-phenylpropyl)-4-(trifluoromethyl)benzamide (2.69 g, 8.39 mmol), phosphorus pentaoxide (1.88 g, 16.8 mmol), toluene (40 mL) and phosphorous oxychloride (6.3 mL, 67.1 mmol). The reaction mixture was heated at 110 °C for 16 h and allowed to cool 5 down to RT. Then it was poured over ice-water and neutralized with NaOH (10%). The organic phase was taken and the aqueous phase was extracted several times with EtOAc. The combined organic layers were washed with sat NaHCO₃, water, brine and dried over Na₂SO₄. The solvent was removed under vacuum and the product used without further purification.



10


Step 5: 4,4-Dimethyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline

A 50-mL round-bottomed flask was charged with 4,4-dimethyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (2.52 g, 8.3 mmol) and dry 15 MeOH (10 mL). After cooling down to 0 °C sodium borohydride (943 mg, 24.9 mmol) was added and the mixture was stirred at RT for 45 min. Then, the solvent was removed and the residue redissolved in EtOAc, washed with sat NaHCO₃, water and dried over Na₂SO₄. The residue was purified by silica gel chromatography (25-35% EtOAc in hexanes) to give the title compound as a 20 white solid. MS (ESI, positive ion) m/z: 306 (M+H).



Step 6: *N*-Isopropyl-4,4-dimethyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide



A 25-mL round-bottomed flask was charged with 4,4-dimethyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (366 mg, 1.2 mmol), isopropyl isocyanate (141 μ L, 1.4 mmol), DCE (5 mL) and stirred at RT for 1 h. Then the solvent was removed under vacuum and the product was purified by 5 recrystallization from a mixture of ether/hexane. MS (ESI, positive ion) m/z: 391 (M+H).



Example 37:

10 **N-(4-Fluorophenyl)-4,4-dimethyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**


A 25-mL round-bottomed flask was charged with 4,4-dimethyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (118 mg, 0.39 mmol), 4-fluorophenyl isocyanate (43.4 μ L, 0.39 mmol) and DCE (5 mL). The reaction 15 mixture was heated at 80 °C for 2 h and the solvent was then removed under vacuum. Purification by silica gel chromatography (20-30% EtOAc in hexanes) provided the title compound as a white solid. MS (ESI, positive ion) m/z: 443 (M+H).



20 **Example 38:**

(1S,4S)-N-(4-Fluorophenyl)-4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide and (1R,4R)-N-(4-fluorophenyl)-4-


methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide



5


Step 1: 2-Phenylpropan-1-amine hydrochloride

To a stirred suspension of lithium aluminum hydride (2.94 g, 77.4 mmol) in dry ether (100 mL) at RT was added, under nitrogen, alpha-methylbenzyl cyanide (6.4 mL, 48.4 mmol) in dry ether (50 mL) over a 30 min period. Then it was
10 heated under reflux for 3 h, cooled to 0 °C and treated sequentially with water (2.3 mL), NaOH (5N, 2.3 mL) and water (6.9 mL). The mixture was then stirred at RT for 1.5 h, filtered and the filtrate dried over Na₂SO₄ the solids were removed by filtration. The solution was treated with 1N HCl in ether (40 mL) and the white precipitated was separated by filtration to provide the title compound as a white
15 solid.




Step 2: *N*-(2-Phenylpropyl)-4-(trifluoromethyl)benzamide

A 250-mL round-bottomed flask was charged with 2-phenylpropan-1-amine hydrochloride (7.75 g, 45.1 mmol), DCM (75 mL) and cooled down to 0 °C.
20 Triethylamine (13.8 mL, 99.3 mmol) was added followed by slow addition of 4-(trifluoromethyl)benzoyl chloride (7.4 mL, 49.7 mmol). After the addition was completed the mixture was stirred at RT for 1h, diluted with EtOAc, washed with 10% HCl, sat NaHCO₃, sat NaCl and dried over Na₂SO₄. The white residue was
25 recrystallized from ether:hexanes and the title compound was obtained as a white solid. MS (ESI, positive ion) m/z: 308 (M+H).




Step 3: 4-Methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline

A 250-mL round-bottomed flask was charged with *N*-(2-phenylpropyl)-4-(trifluoromethyl)benzamide (5.19 g, 16.9 mmol), phosphorous pentoxide (4.8 g, 5 33.8 mmol) and toluene (75 mL). The reaction mixture was heated at 110 °C overnight and allowed to cool down to RT. Then it was poured over ice-water and neutralized with NaOH (10%). The organic phase was separated and the aqueous phase was extracted several times with EtOAc. The combined organics were washed with satd. NaHCO₃, water, brine and dried over Na₂SO₄. The solvent was 10 removed under vacuum and the product used without further purification.



Step 4: 4-Methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline


A 100-mL round-bottomed flask was charged with 4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (4.89 g, 16.9 mmol) and MeOH (30 mL). After cooling down to 0 °C, sodium borohydride (1.93 g, 50.7 mmol) 15 was added and the mixture was stirred at RT for 45 min. Then, the solvent was removed and the residue redissolved in EtOAc, washed with sat NaHCO₃, water and dried over Na₂SO₄. The mixture was used crude in the next step.



Step 5:

- (1*S*,4*S*)-*N*-(4-Fluorophenyl)-4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide and (1*R*,4*R*)-*N*-(4-fluorophenyl)-4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**
- 5


A 25-mL round-bottomed flask was charged with 4-methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (1.24 g, 4.3 mmol), DCE (10 mL) and 4-fluorophenyl isocyanate (528 μ L, 4.7 mmol) and stirred at RT for 10 30 min. The solvent was then removed under vacuum and the residue purified by flash chromatography (10-20% EtOAc in hexanes). The title compounds were further purified by preparative HPLC (10%-100% of MeCN (0.1%TFA)/H₂O (0.1% TFA) (white solid). MS (ESI, positive ion) m/z: 429 (M+H).



15 **Example 39:**


- (1*S*,4*R*)-*N*-(4-Fluorophenyl)-4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide and (1*R*,4*S*)-*N*-(4-Fluorophenyl)-4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

20 The title compounds were obtained as the second fraction after column chromatography from example 38 (step 5) (white solid). MS (ESI, positive ion) m/z: 429 (M+H).

**Example 40:**

- 5 **(1*R*,4*R*)-*N*-(4-Fluorophenyl)-4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

Enantiomers of example 38 were separated using chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 429 (M+H).




10


Example 41:

- (1*S*,4*S*)-*N*-(4-Fluorophenyl)-4-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**

Enantiomers of example 38 were separated using chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 429 (M+H).

**Example 42:**


5-Fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide



5


Step 1: *N*-(2-Fluorophenethyl)-4-(trifluoromethyl)benzamide

A 250-mL, round-bottomed flask containing a solution of 2-fluorophenethylamine (4.7 mL, 35.9 mmol) and diisopropylethylamine (6.3 mL, 35.9 mmol) in dichloromethane (35 mL) was cooled to 0 °C and treated dropwise with 4-(trifluoromethyl)benzoyl chloride (5.4 mL, 35.9 mmol) over 5 min. The ice-bath was removed after 2 h and the reaction mixture was stirred overnight. The precipitate formed was collected by filtration. The filtrate was diluted with dichloromethane (100 mL) and transferred to a separatory funnel, extracted with water (35 mL), brine (25 mL), and 1N HCl (25 mL). The dichloromethane layer was separated and formation of more precipitate was observed which was collected by filtration. The solids collected were combined and dried under vacuum to yield *N*-(2-fluorophenethyl)-4-(trifluoromethyl)benzamide as an off-white solid. MS (ESI pos. ion) *m/z*: 312 (M+H).



Step 2: 5-Fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline


A suspension of *N*-(2-fluorophenethyl)-4-(trifluoromethyl)benzamide (2.0 g, 6.5 mmol) and phosphoric pentoxide (2.7 g, 19.1 mmol) in anhydrous toluene (40 mL) was heated at 125 °C for 4 h. At this stage more phosphoric pentoxide (2.7 g, 19.1 mmol) was added and continued stirring the reaction at 125 °C for additional 16 h. The reaction was cooled to RT, added ethyl acetate (100 mL), and the resulting slurry was slowly poured into ice-water (100 mL). Then more ethyl acetate (75 mL) was added and the resulting suspension was treated with KOH (3.0 N) until pH ~ 11.0 was achieved. The ethyl acetate layer was separated, dried over anhydrous sodium sulfate, and concentrated under vacuum to provide 5-fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline as a yellow oil. MS (ESI pos. ion) *m/z*: 294 (M+H).



15


Step 3: 5-Fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline

A solution of 5-fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (1.9 g, 6.6 mmol) in methanol (40 mL) from step 2 was cooled to 0 °C and sodium borohydride (0.6 g, 17 mmol) was added in small portions over 5 min and the reaction mixture was stirred at RT for 16 h. The reaction was quenched with ethyl acetate (20 mL) and concentrated to yield a reddish-yellow residue which was taken up in ethyl acetate (100 mL), water (30 mL), and saturated sodium bicarbonate (30 mL). After stirring for 15 min, it was transferred to a separatory funnel, the organic layer was separated, dried over anhydrous sodium sulfate, and concentrated to yield the crude product. The crude product was purified by silica gel chromatography (20-30% EtOAc in hexanes) to give 5-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline as a pale-yellow solid. MS (ESI pos. ion) *m/z*: 296 (M+H).



Step 4: 5-Fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

A solution of 5-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.27 g, 0.897 mmol) and diisopropylethylamine (0.2 mL, 0.919 mmol) in anhydrous dichloromethane (6 mL) was treated with 4-fluorophenyl isocyanate (0.105 mL, 0.934 mmol) and stirred at RT for 16 h. The solvent was removed in vacuo and the residue was purified by preparative HPLC [gradient 10-90 % MeCN (0.1 % TFA)/ H₂O (0.1 % TFA)] to give the pure product which was dissolved in methanol (10 mL) and neutralized by passing the solution through a Polymer Lab-HCO₃ Macroporous resin cartridge, and the filtrate was concentrated to give 5-fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as a colorless oil. MS (ESI pos. ion) *m/z*: 433 (M+H).




15

Example 43:

(R)-5-Fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


Purification of racemic 5-fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide on chiral SFC using the following conditions (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) provided (R)-5-fluoro-N-(4-fluorophenyl)-1-(4-

(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 433 (M+H).




Example 44:

- 5 **6-Fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**




Step 1: N-(3-fluorophenethyl)-4-(trifluoromethyl)benzamide

- A 250-mL, round-bottomed flask containing a solution of 2-(3-fluorophenyl)ethanamine (1.3 mL, 9.6 mmol) and diisopropylethylamine (1.7 mL, 9.6 mmol) in dichloromethane (32 mL) was cooled to 0 °C and treated dropwise with 4-(trifluoromethyl)benzoyl chloride (2.0 mL, 9.6 mmol) over 5 min. The ice-bath was removed after 2 h and the reaction mixture was stirred overnight. The precipitate formed was collected by filtration. The filtrate was diluted with dichloromethane (100 mL) and transferred to a separatory funnel and extracted with water (35 mL). The dichloromethane layer was separated and formation of precipitate was observed which was collected by filtration. The filtrate was concentrated to yield more solid and the combined solids were dried under vacuum to yield *N*-(3-fluorophenethyl)-4-(trifluoromethyl)benzamide as a white solid. MS (ESI pos. ion) *m/z*: 312 (M+H).



Step 2: 6-fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline

A suspension of *N*-(3-fluorophenethyl)-4-(trifluoromethyl)benzamide (2.8 g, 8.9 mmol) and phosphoric pentoxide (0.6 g, 0.45 mmol) in polyphosphoric acid (62.0 g, 8.9 mmol) was heated at 170 °C for 75 min. The hot reaction mixture was poured into ice and KOH (20% W/V, 50 mL) was added followed by diethyl ether (100 mL) and stirred for 5 mins. Additional KOH was added until pH ~ 11.0 was achieved. Then, EtOAc (200 mL) was added and the mixture continued stirring at RT for an additional 5 min. and the ethyl acetate layer was separated. The aqueous layer was back-extracted with ethyl acetate (3 x 200 mL). The combined ethyl acetate layers were dried over anhydrous sodium sulfate, and concentrated under vacuum to provide 6-fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline as a yellow oil. MS (ESI pos. ion) *m/z*: 294 (M+H).




Step 3: 6-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline

A solution of 6-fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (2.6 g, 8.9 mmol) in methanol (50 mL) from step 2 was cooled to 0 °C and sodium borohydride (1.0 g, 2.7 mmol) was added in small portions over 5 min and the reaction mixture was stirred at RT for 16 h. The reaction was quenched with ethyl acetate (20 mL) and concentrated to yield a reddish-yellow residue which was taken up in ethyl acetate (100 mL), water (30 mL), and saturated sodium bicarbonate (30 mL). After stirring for 15 min, it was transferred to a separatory


funnel, the organic layer was separated, dried over anhydrous sodium sulfate, and concentrated to yield the crude product. The crude product was purified by silica gel chromatography (20-30% EtOAc in hexanes) to give 6-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline as a pale-yellow solid.

- 5 MS (ESI pos. ion) m/z : 296 (M+H).




Step 4: 6-fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

A solution of 6-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.3 g, 1.0 mmol) and diisopropylethylamine (0.2 mL, 0.919 mmol) in anhydrous dichloromethane (6 mL) was treated with 4-fluorophenyl isocyanate (0.14 g, 1.0 mmol) and stirred at RT for 16 h. Formation of a white precipitate was observed which was collected by filtration. The solid was dried under high vacuum to give 6-fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide (0.35 g, 80%). MS (ESI pos. ion) m/z : 433 (M+H).



Example 45:


20 **7-Fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**



Step 1: *N*-(4-fluorophenethyl)-4-(trifluoromethyl)benzamide

A 250-mL, round-bottomed flask containing a solution of 4-fluorophenethylamine (4.8 mL, 37 mmol) and diisopropylethylamine (6.4 mL, 37 mmol) in dichloromethane (40 mL) was cooled to 0 °C and treated dropwise with 4-(trifluoromethyl)benzoyl chloride (5.5 mL, 37 mmol) over 5 min. The ice-bath was removed after 2 h and the reaction mixture was stirred overnight. The precipitate formed was collected by filtration. The filtrate was diluted with dichloromethane (100 mL) and transferred to a separatory funnel, extracted with water (35 mL), brine (25 mL), and 1N HCl (25 mL). The dichloromethane layer was separated and formation of more precipitate was observed which was collected by filtration. The solids collected were combined and dried under vacuum to yield *N*-(4-fluorophenethyl)-4-(trifluoromethyl)benzamide as an off-white solid. MS (ESI pos. ion) *m/z*: 312 (M+H).


15



Step 2: 7-Fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline


A suspension of *N*-(4-fluorophenethyl)-4-(trifluoromethyl)benzamide (5.3 g, 17.1 mmol) and phosphoric pentoxide (1.2 g, 8.6 mmol) in polyphosphoric acid (40 g) was heated at 165 °C for 4 h and The reaction was cooled to RT and ice was added followed by ethyl acetate (100 mL), KOH (3N, 25 mL), and water (75 mL). The mixture was stirred for 15 min and the organic layer was separated. The aqueous layer was neutralized with KOH (3N) till pH ~ 7.0 was achieved and extracted with ethyl acetate (250 mL). The ethyl acetate layer was separated,

dried over anhydrous sodium sulfate, and concentrated under vacuum to provide 7-fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline. MS (ESI pos. ion) m/z : 294 (M+H).




- 5 Step 3: **7-Fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline**
A solution of 7-fluoro-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (0.6 g, 2.1 mmol) in methanol (10 mL) from step 2 was cooled to 0 °C and sodium borohydride (0.2 g, 6.2 mmol) was added in small portions over 5 min and the reaction mixture was stirred at RT for 16 h. The reaction was quenched with ethyl acetate (10 mL) and concentrated to yield a light-yellow residue which was dissolved in ethyl acetate (40 mL), water (10 mL), and saturated sodium bicarbonate (20 mL). After stirring for 15 min, it was transferred to a separatory funnel, the organic layer was separated, dried over anhydrous sodium sulfate, and concentrated to yield the crude product. The crude product was purified by silica gel chromatography (20-30% EtOAc in hexanes) to give 7-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline as a pale-yellow solid.
10 MS (ESI pos. ion) m/z : 296 (M+H).
- 15

MS (ESI pos. ion) m/z : 296 (M+H).




- Step 4: **7-fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**
20 A solution of 7-fluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.09 g, 0.3 mmol) and diisopropylethylamine (0.06 mL, 0.32 mmol) in anhydrous dichloromethane (3 mL) was treated with 4-fluorophenyl isocyanate

- (0.04 mL, 0.4 mmol) and stirred at RT for 16 h. The solvent was removed in vacuo and the residue was purified by preparative HPLC [gradient 10-90 % MeCN (0.1 % TFA)/ H₂O (0.1 % TFA)] to give the pure product which was dissolved in methanol (10 mL) and neutralized by passing the solution through a
- 5 Polymer Lab-HCO₃ Macroporous resin cartridge, and the filtrate was concentrated to give 5-fluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as an off-white solid. MS (ESI pos. ion) *m/z*: 433 (M+H).



10 **Example 46:**

1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamidine dihydrochloride




Step 1: Bis-boc-guanidine derivative

- 15 A solution of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.6 g, 2.3 mmol) and 1,3-bis(tert-butoxycarbonyl)-2-methyl-2-thiopseudourea (0.75 g, 2.6 mmol) in dichloromethane (12 mL) was treated with mercuric chloride (0.645 g, 2.4 mmol) followed by triethylamine (0.5 mL, 3.5 mmol) and stirred the reaction at RT for 16 h. Formation of a very fine grayish-white precipitate was seen. The precipitate was removed by filtration through a sintered funnel, washed with 25 mL dichloromethane. The combined filtrates were transferred to a
- 20

separatory funnel, washed with water (50 mL), brine (50 mL), dried over anhydrous sodium sulfate and, concentrated on the rotary evaporator to yield the crude product. The crude product was purified by silica gel chromatography (20-30% EtOAc in hexanes) to give the desired product as an off-white solid. MS (ESI pos. ion) m/z : 296 (M+H).

5



Step 2: 1-(4-(Trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamidine dihydrochloride

A solution of the product from step 1 (0.9 g, 2 mmol) in 1,4-dioxane (10 mL) was treated with hydrogen chloride (4.0M solution in 1,4-dioxane, 20 mL, 80 mmol) was stirred at RT for 16 h. During work up the solvent was removed under high vacuum and the resulting gummy solid was re-dissolved in methanol (50 mL), and concentrated again on the rotary evaporator. The resulting solid was stirred with hexanes (75 mL) for 30 min, after which the solid was collected by filtration. The solid 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamidine dihydrochloride (0.62 g, 87% yield) was collected as an off-white solid. MS (ESI pos. ion) m/z : 320 (M+H).

10

General procedure for examples (47-65)

15

To a solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (70 mg, 0.25 mmol, example 30, step 1) in DCM (0.8 mL) and pyridine (0.2 mL) was added the corresponding acid chloride (0.3 mmol). The reaction mixture was stirred at RT for 16 h. Then, H₂O (0.1 mL) and MeOH (0.5 mL) were added. The solvents were removed under vacuum and the residue was dissolved in DMSO (1 mL). The solution mixture was purified by preparative HPLC to provide the target compound. The product was dissolved in MeOH (1 mL) and washed

20

25


through PL-HCO₃ MP-resin, the resin was further washed with MeOH (2 x 0.4 mL). The combined filtrates were then concentrated and dried in vacuo to give the title compounds.

Example	Structure	Name	MS (ESI, positive ion) $M+H$
47		(<i>R</i>)-Furan-3-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	372
48		(<i>R</i>)-(1-Methyl-1 <i>H</i> -indazol-3-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	436
49		(<i>R</i>)-(4-Methyloxazol-5-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	387
50		(<i>R</i>)-2,2-Dimethyl-1-(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)propan-1-one	362

51		(R)-Furan-2-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)methanone	372
52		(R)-Isoxazol-5-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)methanone	373
53		(R)-2-(4-Methyl-1H-pyrazol-1-yl)-1-(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)ethanone	400
54		(R)-(1-Methyl-1H-pyrazol-4-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)methanone	386
55		(R)-Pyridin-3-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)methanone	383


56		(<i>R</i>)-(1-Methyl-1 <i>H</i> -imidazol-5-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	386
57		(<i>R</i>)-(4-Methyl-1,2,3-thiadiazol-5-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	404
58		(<i>R</i>)-(4-(1 <i>H</i> -Pyrazol-1-yl)phenyl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	448
59		(<i>R</i>)-Pyridin-4-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	383
60		(<i>R</i>)-(4-Fluorophenyl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	400
61		(<i>R</i>)-3,3-Dimethyl-1-(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)butan-1-one	376

62		(<i>R</i>)-Cyclopentyl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	374
63		(<i>R</i>)-Pyridin-2-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	383
64		(<i>R</i>)-(2-Methyl-2 <i>H</i> -indazol-3-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	436
65		(<i>R</i>)-(5-Methylisoxazol-4-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1 <i>H</i>)-yl)methanone	387

**Example 66:**

- 5 (*R*)-*N*-Cyclohexyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Enantiomers, *N*-cyclohexyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide (example 13) were separated using chiral SFC (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 403 (M+H).




5


Example 67:

(*R*)-(1-Methyl-1*H*-pyrazol-3-yl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1*H*)-yl)methanone


A solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (58 mg, 0.21 mmol, example 30, step 1) in DMF (1 mL) was added 1-methyl-1*H*-pyrazole-3-carboxylic acid (Fluorochem, 34 mg, 0.27 mmol), 1-hydroxybenzotriazole hydrate (42 mg, 0.27 mmol), and *N*-(isopropylimino)methylene)propan-2-amine (66 µL, 0.42 mmol). The resulting mixture was then stirred at RT for 16 h. Then, the mixture was filtered and the filtrate was purified by preparative HPLC to give the target compound. The product was dissolved in MeOH and passed through PL-HCO₃ MP resin and the resin was washed with MeOH (2 x 0.3 mL). The combined filtrates were concentrated and dried under vacuum to give the title compound as a light-yellow solid. MS (ESI, positive ion) m/z: 386 (M+H).

**Example 68:**

(*R*)-*N*-(4,4-Difluorocyclohexyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide


5 Step 1: **4-Nitrophenyl 4,4-difluorocyclohexylcarbamate**

To a solution of 4,4-difluorocyclohexanamine hydrochloride (Matrix scientific, 250 mg, 1.45 mmol) in DCM (5 mL) was added 4-nitrophenyl chloroformate (367 mg, 1.82 mmol) and DIEA (507 μ L, 0.29 mmol). The resulting mixture was then 10 stirred at RT for 16 h. Then, MeOH (2 mL) was added and the solvents were removed. The residue was purified by silica gel flash column chromatography (0%-100% EtOAc/hexane) to give the title compound as a yellow solid.




Step 2: **(*R*)-*N*-(4,4-Difluorocyclohexyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**
15 To a solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (80 mg, 0.29 mmol) in MeCN (2 mL) was added 4-nitrophenyl 4,4-difluorocyclohexylcarbamate (217 mg, 0.72 mmol). The resulting mixture was then

subjected to microwave irradiation at 120 °C for 15 min. Then, MeOH (0.5 mL) was added and the mixture was filtered. The filtrate was purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the title compound, which was dissolved in MeOH (1 mL). The solution was then passed through PL-5 HCO₃ and the resin was washed with MeOH (2 x 0.3 mL). The combined filtrates were concentrated and dried in vacuo to give the title compound as a light-yellow solid. MS (ESI, positive ion) m/z: 439 (M+H).




Example 69:

- 10 (*IR*)-2-(tricyclo[3.3.1.1~3,7~]dec-1-ylcarbonyl)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline
A solution of (*R*)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.05 g, 0.2 mmol) in DMF (1 mL) was treated with N-((isopropylimino)methylene)propan-2-amine (0.07 mL, 0.5 mmol), 1*H*-benzo[d][1,2,3]triazol-1-ol hydrate 15 (0.04 g, 0.2 mmol), and 1-adamantanecarboxylic acid (TCI, 0.04 g, 0.2 mmol). The resulting mixture was heated at 65 °C for 12 h. The reaction was cooled to RT and the filtrate was purified by preparative HPLC [gradient 10-90 % MeCN (0.1 % TFA)/ H₂O (0.1 % TFA)] to give the pure product. It was dissolved in methanol (10 mL) and neutralized by passing the solution through a Polymer Lab-20 HCO₃ Macroporous resin cartridge, and the filtrate was concentrated to give (*IR*)-2-(tricyclo[3.3.1.1~3,7~]dec-1-ylcarbonyl)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline as a colorless solid. MS (ESI, positive ion) m/z: 440 (M+H).

Example 70:**1-(3,4-Difluorophenyl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**5 **Step 1. 3,4-Difluoro-N-phenethylbenzamide**

To a stirred solution of phenethylamine (1.01 mL, 7.95 mmol) and DIEA (1.38 mL, 7.95 mmol) in 20 mL of DCM, 3,4-difluorobenzoyl chloride (1.00 mL, 7.95 mmol) was added dropwise. The clear solution was stirred for 1 h. The reaction mixture was concentrated and the crude product was purified by silica gel chromatography (0-30% EtOAc in hexanes) to give the title compound as a white solid. ¹H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.57 (ddd, *J*=10.63, 7.56, 2.12 Hz, 1 H) 7.31 - 7.44 (m, 3 H) 7.15 – 7.31 (m, 4 H) 6.02 (br. s., 1 H) 3.67 - 3.78 (m, 2 H) 2.94 (t, *J*=6.87 Hz, 2 H).

10




15

Step 2. 1-(3,4-Difluorophenyl)-3,4-dihydroisoquinoline

To a 250-mL round-bottomed flask, 3,4-difluoro-N-phenethylbenzamide (1.97 g, 7.54 mmol) and phosphorus pentoxide (3.25 g, 22.9 mmol) were suspended into 40 mL of toluene. Phosphorus oxychloride (3.51 mL, 37.7 mmol) was added and the mixture was refluxed for overnight. The mixture was cooled to RT. The supernatant contained only the starting material and was removed. Ice water (75 mL) was added to the residue in the flask and stirred to break up the solid into powder.


20 EtOAc was added and the pH was adjusted to ~12 with 3N aqueous potassium hydroxide. The layers were separated and the aqueous phase was

extracted with EtOAc. The combined organic phases were washed with water and saturated aqueous sodium chloride. The organic phase was dried over sodium sulfate, filtered and concentrated in vacuo to afford a yellow viscous oil. The crude product was purified by silica gel chromatography (20-70% EtOAc in hexanes) to afford the title compound as a clear oil. 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.17 – 7.52 (m, 7 H) 3.78 - 3.92 (m, 2 H) 2.73 - 2.89 (m, 2 H)



10 Step 3. **1-(3,4-Difluorophenyl)-1,2,3,4-tetrahydroisoquinoline**


To a stirred solution of 1-(3,4-difluorophenyl)-3,4-dihydroisoquinoline (1.34 g, 5.51 mmol) in 20 mL of MeOH, sodium borohydride (0.4366 g, 11.5 mmol) was added. The mixture was stirred for 1 h. MeOH was evaporated and the residue was partitioned between EtOAc and saturated aqueous sodium bicarbonate. The aqueous phase was extracted with EtOAc. The combined organic phases were washed with water and saturated aqueous sodium chloride. The organic phase was dried over sodium sulfate, filtered and concentrated in vacuo to afford a white solid. The crude product was purified by silica gel chromatography (0-5% isopropanol (with 10% ammonium hydroxide) in CHCl₃) to afford the title compound as a white solid. 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 6.99 - 7.22 (m, 6 H) 6.74 (d, *J*=7.60 Hz, 1 H) 5.08 (s, 1 H) 3.18 - 3.32 (m, 1 H) 2.96 - 3.18 (m, 2 H) 2.76 - 2.90 (m, 1 H).



Step 4. 1-(3,4-Difluorophenyl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


To a stirred solution of 1-(3,4-difluorophenyl)-1,2,3,4-tetrahydroisoquinoline

- 5 (0.1019 g, 0.415 mmol) in 3 mL of DCM, 4-fluorophenyl isocyanate (0.0550 mL, 0.484 mmol) was added. The clear solution was stirred for 1 h. The solvent was evaporated and residue was taken into small amount of DCM. The resulting precipitate was collected to yield the title compound as a white solid. The filtrate was concentrated and purified by silica gel chromatography (0-30% EtOAc in hexanes) to afford another batch of the title compound as a white solid. MS (ESI pos. ion) *m/z*: 383 (M+1). 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.16 - 7.37 (m, 6 H) 6.93 - 7.13 (m, 5 H) 6.54 (s, 1 H) 6.40 (s, 1 H) 3.66 (t, *J*=6.28 Hz, 2 H) 2.77 - 3.08 (m, 2 H).




15 **Example 71:**

1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-isopropyl-3,4-dihydroisoquinoline-2(1H)-carboxamide



Step 1. 3-Fluoro-N-phenethyl-4-(trifluoromethyl)benzamide


To a 100-mL round-bottomed flask, 3-fluoro-4-(trifluoromethyl)benzoic acid (1.036 g, 4.98 mmol) and 2-phenylethanamine (0.700 mL, 5.53 mmol) were dissolved into 25 mL of DMF. Triethylamine (0.840 mL, 6.03 mmol) was added followed by 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (0.955 g, 4.98 mmol) and 1-hydroxybenzotriazole (0.915 g, 5.97 mmol). The mixture was stirred for 28 h. The reaction mixture was poured into 80 mL of saturated aqueous sodium bicarbonate. The aqueous phase was extracted with EtOAc. The combined organic phases were washed with saturated aqueous sodium bicarbonate, water and saturated aqueous sodium chloride. The organic phase was dried over sodium sulfate, filtered and concentrated in vacuo to afford a yellowish solid. DCM was added to the crude product and the resulting solid was collected via filtration to afford the title compound as a white solid. The filtrate was concentrated and purified by silica gel chromatography (100% EtOAc) to afford another batch of the title compound as a white solid. ^1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.66 (t, *J*=7.45 Hz, 1 H) 7.46 - 7.59 (m, 2 H) 7.28 - 7.41 (m, 3 H) 7.20 - 7.26 (m, 2 H) 6.14 (br. s., 1 H) 3.69 - 3.81 (m, 2 H) 2.96 (t, *J*=6.87 Hz, 2 H).




Step 2. 1-(3-Fluoro-4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline

To a stirred mixture of 3-fluoro-N-phenethyl-4-(trifluoromethyl)benzamide (1.27 g, 4.08 mmol) in 30 mL of toluene, phosphorus pentoxide (1.751 g, 12.3 mmol) and phosphorus oxychloride (1.90 mL, 20.4 mmol) were added. The mixture was refluxed for overnight. The reaction mixture was cooled to RT. Supernatant was removed and the ice-water was added to the flask containing the residue. The mixture was stirred to break the chunks into powder. EtOAc was added and the pH was adjusted to ~12 with 3N aqueous potassium hydroxide. The layers were separated and the aqueous phase was extracted with EtOAc. The combined

organic phases were washed with water and saturated aqueous sodium chloride. The organic phase was dried over sodium sulfate, filtered and concentrated in vacuo to afford 0.887 g of yellow viscous oil. The crude product was purified by silica gel chromatography (0-30% EtOAc in hexanes) to afford the title compound 5 as a clear oil. ^1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.63 - 7.73 (m, 1 H) 7.39 - 7.54 (m, 3 H) 7.31 (d, *J*=7.60 Hz, 2 H) 7.16 - 7.24 (m, 1 H) 3.84 - 3.96 (m, 2 H) 2.77 - 2.89 (m, 2 H).




Step 3. **1-(3-Fluoro-4-(Trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline**
10 To a stirred solution of 1-(3-fluoro-4-(trifluoromethyl)phenyl)-3,4-dihydroiso-
quinoline (0.750 g, 2.6 mmol) in 10 mL of MeOH, sodium borohydride (0.197 g,
5.2 mmol) was added. The mixture was stirred for 2.5 h. MeOH was evaporated
and the residue was partitioned between EtOAc and saturated aqueous sodium
bicarbonate. The aqueous phase was extracted with EtOAc. The combined organic
15 phases were washed with water and saturated aqueous sodium chloride. The
organic phase was dried over sodium sulfate, filtered and concentrated in vacuo to
afford a white solid. The crude product was purified by silica gel chromatography
(30-70% EtOAc in hexanes to afford the title compound as a white solid. ^1H
NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.56 (t, *J*=7.63 Hz, 1 H) 7.05 - 7.24
20 (m, 5 H) 6.73 (d, *J*=7.63 Hz, 1 H) 5.16 (s, 1 H) 3.19 - 3.29 (m, 1 H) 2.97 - 3.16
(m, 2 H) 2.79 - 2.89 (m, 1 H).



Step 4. 1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-isopropyl-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a stirred solution of 1-(3-fluoro-4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.1020 g, 0.345 mmol) in 3 mL of DCM, isopropyl isocyanate


- 5 (0.0400 mL, 0.407 mmol) was added. The solution was stirred for 1 h. Solvent was evaporated and the residue was purified by silica gel chromatography (0-50% EtOAc in hexanes) then again with 10-40% EtOAc in hexanes to afford the title compound as a white solid. MS (ESI pos. ion) *m/z*: 381.1 (M+1). 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.48 (t, *J*=7.75 Hz, 1 H) 7.19 – 7.32 (m, 2 H) 10 7.01 - 7.14 (m, 2 H) 6.55 (s, 1 H) 4.30 (s, 1 H) 3.98 - 4.13 (m, 1 H) 3.38 - 3.63 (m, 2 H) 2.69 - 2.98 (m, 2 H) 1.13 - 1.25 (m, 6 H).



Example 72:


1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

- To a stirred solution of 1-(3-fluoro-4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.0845 g, 0.286 mmol, Example 71 Step 3) in 3 mL of DCM, 4-fluorophenyl isocyanate (0.0358 mL, 0.315 mmol) was added. The solution was stirred for 1 h. Solvent was evaporated and the residue was purified by silica gel chromatography (0-50% EtOAc in hexanes) to yield the title compound as a white solid. MS (ESI pos. ion) *m/z*: 433.1 (M+1). 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.51 (t, *J*=7.75 Hz, 1 H) 7.22 - 7.39 (m, 6 H) 7.06 - 7.19 (m, 2 H) 6.96 - 7.05 (m, 2 H) 6.63 (s, 1 H) 6.45 (s, 1 H) 3.59 - 3.81 (m, 2 H) 2.76 - 3.08 (m, 2 H).

**Example 73:**


(S)-1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

- 5 After chiral separation (SFC) of the enantiomers of the previous example 72, the title compound was obtained. MS (ESI pos. ion) m/z : 433.1 (M+1).

**Example 74:**

(R)-1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

- 10 After chiral separation (SFC) of the enantiomers of the previous example 72, the title compound was obtained. MS (ESI pos. ion) m/z : 433.1 (M+1).




- 15 **Example 75:**

1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-(pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

1-(3-fluoro-4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.1500 g, 0.508 mmol, Example 71 Step 3) was dissolved in 3 mL CH₂Cl₂ and DIEA (0.0884 mL, 0.508 mmol) and 3-pyridyl isocyanate (0.0732 g, 0.610 mmol) were added at RT. The mixture was stirred over night and hydrolyzed with 2 mL water.

- 5 The reaction was extracted 3 times with EtOAc (3x 20 mL), dried over MgSO₄ and evaporated. The crude product was purified via glass column chromatography providing 1-(3-fluoro-4-(trifluoromethyl)phenyl)-N-(pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as a white solid. MS (ESI, positive ion) m/z: 416.0 (M+H). The majority of the material was submitted for chiral purification (SFC):


10



Example 76:

(S)-1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-(pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


- 15 After chiral separation (SFC) of the enantiomers of the previous example the title compound was obtained. MS (ESI, positive ion) m/z: 416.0 (M+H)




Example 77:

(R)-1-(3-Fluoro-4-(trifluoromethyl)phenyl)-N-(pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

- 20 After chiral separation (SFC) of the enantiomers of the previous example the title compound was obtained. MS (ESI, positive ion) m/z: 416.0 (M+H)

**Example 78:**

4,4-Difluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide




5

Step 1. 2-Benzyl-1-(4-(trifluoromethyl)phenyl)-1,2-dihydroisoquinolin-3(4H)-one

4-(Trifluoromethyl)benzaldehyde (1.87 mL, 14.0 mmol) and benzylamine (1.53 mL, 14.0 mmol) were dissolved into 40 mL of DCM and 4Å molecular sieve was added. The mixture was left under nitrogen for 3 days. Molecular sieve was removed via filtration. 25 mL of DCM was used to wash the filter cake. To the filtrate, phenylacetyl chloride (1.90 mL, 14.4 mmol) was added and the pale yellow solution was stirred for 2 h. Then trifluoromethanesulfonic acid (6.20 mL, 70.1 mmol) was added. The mixture was stirred for 1.5 h then poured into a mixture of ice (~100 mL) and 5N sodium hydroxide (~50 mL). The phases were separated and the aqueous phase was extracted with DCM. The combined organic phases were dried over sodium sulfate, filtered and concentrated in vacuo to afford a yellow oil. The crude product was purified by silica gel chromatography (0-30% EtOAc in hexanes) twice to afford the title compound as a light yellow semi-solid. 1H NMR (300 MHz, DMSO-*d*₆) δ ppm 7.56 - 7.74 (m, 4 H) 7.41 (d,


J=6.87 Hz, 1 H) 7.13 - 7.33 (m, 8 H) 5.80 (s, 1 H) 5.30 (d, *J*=15.20 Hz, 1 H) 3.81 - 4.01 (m, 2 H) 3.66 - 3.78 (m, 1 H).



Step 2. 2-Benzyl-4,4-difluoro-1-(4-(trifluoromethyl)phenyl)-1,2-dihydroiso-


5 *quinolin-3(4H)-one*

To a stirred solution of 2-benzyl-1-(4-(trifluoromethyl)phenyl)-1,2-dihydroisoquinolin-3(4H)-one (1.472 g, 3.86 mmol) in 20 mL of THF, lithium bis(trimethylsilyl)amide (1.0M in THF/ethyl benzene, 8.49 mL, 8.49 mmol) was added slowly at -78 °C. The solution was stirred at -78 °C for 30 min, then n-fluorobenzene-sulfonimide (2.799 g, 8.875 mmol) in a total of 15 mL THF was added slowly. After the solution was stirred at -78 °C for 1 h, the cold bath was removed and the reaction mixture was allowed to warm up to RT. EtOAc was added followed by 100 mL saturated aqueous NH₄Cl. Layers were separated and the aqueous phase was extracted with EtOAc. The organic phase was washed with water and saturated aqueous sodium chloride. The aqueous washes were combined and back-extracted with EtOAc. The combined organic phases were dried over sodium sulfate, filtered and concentrated in vacuo to afford 2.90 g of slurry. DCM and methyl *t*-butyl ether were added to the crude product and the resulting white precipitate was removed via filtration. The filtrate was concentrated to give a light yellow residue. The crude product was purified by silica gel chromatography (0-30% EtOAc in hexanes) to afford the title compound as a pale yellow solid. 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.83 - 7.91 (m, 1 H) 7.65 (d, *J*=8.04 Hz, 2 H) 7.30 - 7.52 (m, 7 H) 7.19 - 7.26 (m, 2 H) 7.07 (s, 1 H) 5.65 (d, *J*=14.91 Hz, 1 H) 5.53 (d, *J*=3.51 Hz, 1 H) 3.64 (dd, *J*=14.91, 1.61 Hz, 1 H).




Step 3. 2-Benzyl-4,4-difluoro-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline

To a stirred solution of borane tetrahydrofuran complex (1M in THF, 0.750 mL, 5 0.750 mmol) in 5 mL of THF, 2-benzyl-4,4-difluoro-1-(4-(trifluoromethyl)- phenyl)-1,2-dihydroisoquinolin-3(4H)-one (0.103 g, 0.246 mmol) in a total of 5 mL THF was added dropwise. The clear solution was refluxed for 5 h. After the reaction mixture was cooled to 0 °C, 5 mL of 5 N hydrochloric acid was added. The solution was stirred at 0 °C to RT over 1 h. Then the most organic solvent was 10 evaporated and the pH was adjusted to ~8 by solid sodium bicarbonate. The aqueous phase was extracted with EtOAc. The combined organic phases were washed with water and saturated aqueous sodium chloride. The organic phase was dried over sodium sulfate, filtered and concentrated in vacuo to afford the title compound as a white solid. 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.76
15 (d, *J*=7.75 Hz, 1 H) 7.62 (d, *J*=8.18 Hz, 2 H) 7.46 (d, *J*=8.04 Hz, 2 H) 7.28 - 7.42 (m, 7 H) 6.77 (d, *J*=7.75 Hz, 1 H) 4.80 (d, *J*=3.65 Hz, 1 H) 3.82 (dd, *J*=13.59, 1.75 Hz, 1 H) 3.37 - 3.54 (m, 2 H) 2.87 - 3.06 (m, 1 H).




Step 4. 4,4-Difluoro-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a 50-mL round-bottomed flask, 2-benzyl-4,4-difluoro-1-(4-(trifluoromethyl)-phenyl)-1,2,3,4-tetrahydroisoquinoline (0.0418 g, 0.10 mmol) and Pd/C (0.0729 g, 0.69 mmol) were mixed into 5 mL of EtOH. The reaction mixture was evacuated under vacuum and refilled with hydrogen (4 times). The mixture was hydrogenated under balloon pressure of hydrogen for 4 h. The catalyst was removed via filtration through a pad of Celite. The filtrate was concentrated to give 0.0291 g of clear film. This product was dissolved in DCM (~ 2 mL) and 1-isocyanato-4-(trifluoromethyl)benzene (0.0200 mL, 0.13 mmol) was added and the mixture was stirred for 30 min. The reaction mixture was directly purified by silica gel chromatography (0-30% EtOAc in hexanes) twice to afford the title compound as a white solid. MS (ESI pos. ion) *m/z*: 451.1 (M+1) and 473.1 (M+23). 1H NMR (300 MHz, CHLOROFORM-*d*) δ ppm 7.79 - 7.88 (m, 1 H) 7.60 (d, *J*=8.18 Hz, 2 H) 7.47 - 7.56 (m, 2 H) 7.41 (d, *J*=8.18 Hz, 2 H) 7.28 - 7.36 (m, 2 H) 7.14 - 7.22 (m, 1 H) 6.96 - 7.07 (m, 2 H) 6.84 (s, 1 H) 6.54 (s, 1 H) 4.07 - 4.23 (m, 1 H) 3.53 - 3.76 (m, 1 H).



Example 79:

1-(4-Chlorophenyl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide




20

Step 1: N-(4-Chlorobenzylidene)-2-phenylethanamine


To a rb-flask containing MgSO₄ (3.06 g, 25.4 mmol) in benzene (10 mL) was added 4-chlorobenzaldehyde (2.02 g, 14.4 mmol) and 4-chlorobenzaldehyde (2.02

g, 14.4 mmol) and the suspension stirred at RT 2 h. The suspension was filtered and the solid washed with benzene (2 x 2 mL). The combined filtrates were concentrated to give the crude imine as a yellow/white solid.



5 **Step 2: 1-(4-Chlorophenyl)-1,2,3,4-tetrahydroisoquinoline**


To a solution of trifluoromethanesulfonic acid (15.3 mL, 173 mmol) at 0 °C was added N-(4-chlorobenzylidene)-2-phenylethanamine (3.38 g) and the reaction was heated to 120 °C and stirred 17 h under N₂. The reaction was poured into ice water (100 mL) and basified with 5N NaOH (45 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 100 mL). The combined organic layers were washed with saturated NaCl (100 mL), dried (MgSO₄) and concentrated to give a yellow/brown oil. The residue was purified by ISCO (120 g SiO₂, 0-50% EtOAc/hexane) to give 1-(4-chlorophenyl)-1,2,3,4-tetrahydroisoquinoline as a tan solid.




15

Step 3: **1-(4-Chlorophenyl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

To a solution of 1-(4-chlorophenyl)-1,2,3,4-tetrahydroisoquinoline (150 mg, 615 µmol) in CH₂Cl₂ (2.0 mL) at RT was added 1-fluoro-4-isocyanatobenzene (85.8 µL, 739 µmol). The reaction was stirred 3 h and then directly purified by reverse phase HPLC to give the title compound as a white solid. MS (ESI pos. ion) *m/z*: 381 (M+1).

**Example 80:**


(1*R*,3*S*)-1-(4-Chlorophenyl)-N-(4-fluorophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]



5

Step 1: **(*S*)-N-(4-Chlorobenzylidene)-1-phenylpropan-2-amine** [stereochemistry is arbitrarily assigned]


To a solution of (*S*)-1-phenylpropan-2-amine (626 mg, 4630 µmol) in benzene (10 mL) was added 4-chlorobenzaldehyde (651 mg, 4630 µmol) and MgSO₄ (1000 mg, 8308 µmol) and the suspension stirred at RT 4 h. The suspension was filtered and the solid washed with benzene (2 x 2 mL). The combined filtrates were concentrated to give (*S*)-N-(4-chlorobenzylidene)-1-phenylpropan-2-amine, which was carried on into the next step without purification.



15 Step 2: **(1*R*,3*S*)-1-(4-Chlorophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline** [stereochemistry is arbitrarily assigned]


To a flask with trifluoromethanesulfonic acid (3.00 mL, 34 mmol) at 0 °C was added dropwise (*S*)-N-(4-chlorobenzylidene)-1-phenylpropan-2-amine (1.20 g,

1.61 mmol) and the reaction was heated to 120 °C and stirred 19 h under N₂. The reaction was cooled, poured into ice water (20 mL), and basified with 5 N NaOH (6 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 10 mL). The combined organic layers were washed with saturated NaCl (20 mL), dried (MgSO₄), and concentrated to give a yellow/brown oil. The residue was purified by ISCO (40 g SiO₂, 0-100% EtOAc/hexane) to give (1*R*,3*S*)-1-(4-chlorophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline as a light yellow oil.




Step 3: (1*R*,3*S*)-1-(4-Chlorophenyl)-N-(4-fluorophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]

To a solution of (1*R*,3*S*)-1-(4-chlorophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline (60 mg, 233 μmol) in CH₂Cl₂ (1.0 mL) at RT was added 1-fluoro-4-isocyanatobenzene (42 μL, 361 μmol). The reaction was stirred 4 h and directly purified by reverse phase HPLC to give (1*R*,3*S*)-1-(4-chlorophenyl)-N-(4-fluorophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide as a light purple solid. MS (ESI pos. ion) *m/z*: 395 (M+1).




Example 81:

(1*R*,3*S*)-1-(4-Bromophenyl)-N-(4-fluorophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]



Step 1: (S)-N-(4-Bromobenzylidene)-1-phenylpropan-2-amine [stereochemistry is arbitrarily assigned]


To a solution of (S)-1-phenylpropan-2-amine (323 mg, 2389 µmol) in benzene (5 mL) was added 4-bromobenzaldehyde (442 mg, 2389 µmol) and MgSO₄ (500 mg, 4154 µmol) and the suspension was stirred at RT 4 h. The suspension was filtered and the solid washed with benzene (2 x 2 mL). The combined filtrates were concentrated to give (S)-N-(4-bromobenzylidene)-1-phenylpropan-2-amine which was used without purification in the next step.



10

Step 2: (1*R*,3*S*)-1-(4-bromophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline [stereochemistry is arbitrarily assigned]


To a flask with trifluoromethanesulfonic acid (3.00 mL, 34 mmol) at 0 °C was added dropwise the (S)-N-(4-bromobenzylidene)-1-phenylpropan-2-amine and the reaction was heated to 120 °C and stirred 19 h under N₂. The reaction was poured into ice water (20 mL) and basified with 5 N NaOH (6 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 10 mL). The combined organic layers were washed with saturated NaCl (20 mL), dried (MgSO₄), and concentrated to give a yellow/brown oil (410 mg). H-NMR shows 92:8 diastereomeric ratio. The residue was purified by ISCO (40 g SiO₂, 0-100% EtOAc/hexane) to give the title compound as a light yellow oil.



Step 3: (1*R*,3*S*)-1-(4-Bromophenyl)-N-(4-fluorophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]

- To a solution of (1*R*,3*S*)-1-(4-bromophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline (48.5 mg, 160 µmol) in CH₂Cl₂ (1.0 mL) at RT was added 1-fluoro-4-isocyanatobenzene (28.9 µL, 249 µmol). The reaction was stirred 4 h and directly purified by reverse phase HPLC to give (1*R*,3*S*)-1-(4-bromophenyl)-N-(4-fluorophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 439, 441 (M+1).


10



Example 82:


((1*R*,3*S*)-1-(4-Chlorophenyl)-N-(4-cyanophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]

- To a solution of (1*R*,3*S*)-1-(4-chlorophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline (50 mg, 194 µmol, example 80, step 2) in CH₂Cl₂ (1.0 mL) at RT was added 4-isocyanatobenzonitrile (42 mg, 291 µmol). The reaction was stirred 5 h and directly purified by reverse phase HPLC to give (1*R*,3*S*)-1-(4-chlorophenyl)-N-(4-cyanophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 402 (M+1).


**Example 83:**

(1*R*,3*S*)-1-(4-Bromophenyl)-N-(4-cyanophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide [stereochemistry is arbitrarily assigned]

- 5 To a solution of (1*R*,3*S*)-1-(4-bromophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline (50 mg, 165 µmol, example 81, step 2) in CH₂Cl₂ (1.0 mL) at RT was added 4-isocyanatobenzonitrile (37 mg, 256 µmol). The reaction was stirred 5 h and directly purified by reverse phase HPLC to give (1*R*,3*S*)-1-(4-bromophenyl)-N-(4-cyanophenyl)-3-methyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 446, 448 (M+1).
- 10

**Example 84:**

4-((1*R*,3*S*)-2-(4-Fluorophenylcarbamoyl)-3-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)benzoic acid [stereochemistry is arbitrarily assigned]




15

Step 1: **(*S*)-N-(4-iodobenzylidene)-1-phenylpropan-2-amine**

To a solution of (*S*)-1-phenylpropan-2-amine (0.500 g, 3698 µmol) in benzene (5 mL) was added 4-iodobenzaldehyde (858 mg, 3698 µmol) and MgSO₄ (1.000 g,

8308 µmol) and the suspension stirred at RT 4 h. The suspension was filtered and the solid washed with benzene (2 x 2 mL). The combined filtrates were concentrated to give (*S*)-N-(4-iodobenzylidene)-1-phenylpropan-2-amine as a yellow oil which was used without purification in the next step.




5

Step 2: (1*R*,3*S*)-1-(4-Iodophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline

[stereochemistry is arbitrarily assigned]

To a flask with the crude N-(4-iodobenzylidene)-1-phenylpropan-2-amine (1.17 g) was added trifluoromethanesulfonic acid (6.00 mL, 67805 µmol) and the reaction 10 was heated to 120 °C and stirred 5 h under N₂. The reaction was cooled to RT, poured into ice water (20 mL), and basified with 5 N NaOH (15 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 10 mL). The combined organic layers were washed with saturated NaCl (20 mL), dried (MgSO₄), and concentrated to give a yellow/brown oil. The residue was purified by ISCO (40 g 15 SiO₂, 0-100% EtOAc/hexane) to give (1*R*,3*S*)-1-(4-iodophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline as a brown oil.




Step 3: 4-((1*R*,3*S*)-3-Methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)benzoic acid

20 [stereochemistry is arbitrarily assigned]


To a vial with (1*R*,3*S*)-1-(4-iodophenyl)-3-methyl-1,2,3,4-tetrahydroisoquinoline (56.2 mg, 161 µmol) under N₂ was added THF (1.0 mL) and the reaction was cooled to -78 °C. *n*-butyllithium (129 µL, 322 µmol) was added and the reaction

stirred 5 min. CO₂ gas was bubbled through the solution and the reaction warmed to RT and stirred 20 min. The reaction mixture was quenched with MeOH (1 mL) and diluted with DMF (1 mL). The reaction mixture was directly purified by HPLC to give 4-((1*R*,3*S*)-3-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)benzoic acid as a white solid.



Step 4: 4-((1*R*,3*S*)-2-(4-Fluorophenylcarbamoyl)-3-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)benzoic acid [stereochemistry is arbitrarily assigned]


- To a solution of 4-((1*R*,3*S*)-3-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)benzoic acid (22 mg, 82 µmol) in CH₂Cl₂ (0.5 mL) at RT was added 1-fluoro-4-isocyanatobenzene (28 mg, 206 µmol). The reaction was stirred 5 h and directly purified by HPLC to give 4-((1*R*,3*S*)-2-((4-fluorophenyl)carbamoyl)-3-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)benzoic acid as a white solid. MS (ESI pos. ion) *m/z*: 405 (M+1).



Example 85:


- N-(4-fluorophenyl)-1-(4-(trifluoromethoxy)phenyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxamide**
- A round-bottomed flask containing a solution of 1-bromo-4-(trifluoromethoxy)benzene (0.093 mL, 626 µmol) and dry THF (2.5 mL) was cooled to -78 °C. The

solution was treated with 2.5M butyllithium in toluene (0.25 mL, 625 μmol) and after 2 min a solution of 3,4-dihydroisoquinoline (76 mg, 579 μmol) in THF (1 mL) was added. After 2 min LC-MS shows a predominant peak (MS (ESI pos. ion) m/z : 294 ($M+1$)). The reaction was quenched with sat'd NH₄Cl (2 mL). The 5 organic layer was separated and the aqueous layer extracted with EtOAc (2 mL). The combined organic layers were dried over MgSO₄. The solution was then treated with 1-fluoro-4-isocyanatobenzene (0.06 mL, 528 μmol). After 16 h, a solid was filtered and the solids washed with EtOAc (2 mL) to give 1,3-bis(4-fluorophenyl)urea as a white solid. The filtrate was concentrated in vacuo and 10 taken up in 1.5 mL of DMF and purified by reverse-phase preparative HPLC (Shimadzu) on a Phenomenex Gemini® column (5 micron, C18, 110 Å, Axia, 100x50 mm) eluting at 90 mL/min with an linear gradient of 10% to 100% MeCN (0.1%TFA) in water (0.1% TFA) over 30 min to give N-(4-fluorophenyl)-1-(4-(trifluoromethoxy)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide (40 mg, 15 16% yield), as colorless film. MS (ESI pos. ion) m/z : 431 ($M+1$).



Example 86:

N-(4-Fluorophenyl)-1-(4-methoxyphenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide




20

Step 1: 1-(4-Methoxyphenyl)-1,2,3,4-tetrahydroisoquinoline

To a round-bottomed flask containing ethyl 1-(4-methoxyphenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (199 mg, 639 μmol) and EtOH (10 mL) was added

1N NaOH (5 mL). The solution was stirred at 100 °C. After 3 h, a pellet of KOH was added and heating continued. After a further 16 h the condenser was removed and 2 pellets of solid KOH were added. After a further 24 h, LC-MS shows ~50% conversion. 10 mL of water was added and the condenser was replaced. After a further 24 h, the reaction was allowed to cool and extracted with CH₂Cl₂ (3 X 15 mL) and the combined organics concentrated in vacuo to give a mixture of starting material and 1-(4-methoxyphenyl)-1,2,3,4-tetrahydroisoquinoline (170 mg) as a golden oil. The crude product was used in the next step. MS (ESI pos. ion) *m/z*: 240 (M+1).


10



Step 2: N-(4-fluorophenyl)-1-(4-methoxyphenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


To a round-bottomed flask containing crude 1-(4-methoxyphenyl)-1,2,3,4-tetrahydroisoquinoline (25 mg, 104 μmol) and CH₂Cl₂ (10 mL) was added 1-fluoro-4-isocyanatobenzene (0.03 mL, 264 μmol). The solution was stirred at RT. After 1 h, the reaction was concentrated in vacuo and taken up in DMF and purified by reverse-phase preparative HPLC (Shimadzu) on a Phenomenex Gemini™ column (5 micron, C18, 110 Å, Axia, 100x50 mm) eluting at 90 mL/min with a linear gradient of 10% to 100% MeCN (0.1%TFA) in water (0.1% TFA) over 10 min to give crude product, that was lyophilized and resubjected to the same purification conditions to give N-(4-fluorophenyl)-1-(4-methoxyphenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide, as a white solid that contained 0.66 eq of TFA. MS (ESI pos. ion) *m/z*: 377 (M+1).

25


**Example 87:**

(R)-Benzo[c][1,2,5]oxadiazol-5-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)methanone


5

**Step1: N-Phenethyl-4-(trifluoromethyl)benzamide**

To a solution of phenylethylamine (14.0 g, 14.6 mL, 115.5 mmol) and diea (20.1 mL, 115.5 mmol) in DCM (500 mL) at 0°C was added 4-(trifluoromethyl)benzoyl chloride (17.2 mL, 115.5 mmol) dropwise. After addition, the reaction mixture was then stirred at 0°C and warmed up itself to RT overnight. An off-white solid precipitation was observed. The off-white solid was collected by filtration. The solid was then washed with DCM (2 x 50 mL), and dried under vacuum to give 31.8 g of the title compound as a white solid. MS (ESI, positive ion) m/z: 294 (M+H).


**Step2: 1-(4-(Trifluoromethyl)phenyl)-3,4-dihydroisoquinoline**

To a round-bottomed flask was added n-phenethyl-4-(trifluoromethyl)benzamide (5.0 g, 17 mmol), phosphoric pentoxide (0.526 mL, 8.52 mmol), and polyphosphoric acid (60.0 g). The reaction mixture was then heated to 165°C for 2 h. A brown thick solution was observed. Then, the hot oil solution was carefully 5 poured into ice in an erlenmeyer flask and a solution of KOH (20%, 40 mL) was added to break the oil. The mixture was stirred at RT for 5 min and an additional KOH solution was added until pH 7. Then, ethyl ether (250 mL) was added and the mixture was stirred at RT for 15 min. The organic layer was collected and the aqueous layer was extracted with EtOAc (2 x 200 mL). The combined organic 10 extracts were dried over MgSO₄ and concentrated to give the title compound as an orange oil, which was used in the next step without purification requirement. MS (ESI, positive ion) m/z: 276 (M+H).




15 Step 3: **1-(4-(Trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline**


To a solution of 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (4.693 g, 17 mmol) (example 1b) in MeOH (24 mL) was added sodium borohydride (1.94 g, 51.1 mmol) slowly at 0°C . After addition, the mixture was stirred at 0 °C for 15 min and at RT for 2 h. The solvent was then removed and H₂O (20 mL) was 20 added to the residue. Then, a saturated NaHCO₃ solution (150 mL) was added slowly and the mixture was extracted with EtOAc (2 x 200 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was mixed with silica gel (4:1 ; residue:silica gel). The solid mixture was purified by silica gel flash column chromatography using ISCO instrument (solid loading, 25 20%-100% EtOAc/hexane) to give the title compound as a white solid. MS (ESI, positive ion) m/z: 278 (M+H).



- Step 4: **(R)-1-(4-(Trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline**
Enantiomers of 1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline
(2.120 g, 7.46 mmol, racemic mixture) were separated by using chiral SFC
5 (Chiralcel AD-H (250 x 21 Mm), 45% methanol/CO₂ (100 bar), 65 mL/min, 220 nm) to give 877 mg of the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 278 (M+H).
- Step 5: **(R)-Benzo[c][1,2,5]oxadiazol-5-yl(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinolin-2(1H)-yl)methanone**
- 10 To a solution of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (50 mg, 180 µmol) and diea (31 µL, 180 µmol) in DCM (1 mL) was added benzo[c][1,2,5]oxadiazole-5-carbonyl chloride (43 mg, 234 µmol). The resulting mixture was then stirred at RT for overnight. Then, the mixture was concentrated and the residue was dissolved in MeOH (1 mL). The solution mixture was then
15 purified by preparative HPLC (0%-100% MeCN 0.1% TFA / H₂O 0.1% TFA) to give a desired product, which was then dissolved in MeOH (1 mL). The solution was then washed through PL-HCO₃ resin and the resin was then washed with MeOH (1 mL). The combined washings were concentrated and dried under vacuum to give the title compound as a white solid. MS (ESI, positive ion) m/z:
20 424 (M+H).


**Example 88:****(R)-4-Nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate**

- 5 To a solution of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (1550 mg, 5590 μ mol) and diea (1071 μ L, 6149 μ mol) in DCM (36 mL) at 0 °C was added 4-nitrophenyl chloroformate (1465 mg, 7267 μ mol, Aldrich). After addition, the reaction mixture was stirred at RT for overnight. Then, H₂O (60 mL) was added and the mixture was stirred at RT for 15 min. The organic layer was collected and the aqueous was extracted with DCM (1 x 50 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was then mixed with silica gel (4:1, residue: silica gel) and the solid mixture was purified by silica gel flash column chromatography using ISCO instrument (solid loading, 0%-100% EtOAc/hexane) to give the compound as an off-white solid. MS (ESI, positive ion) m/z: 443 (M+H).
- 10
- 15

**Example 89:****(R)-N-(4-Cyano-2-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

- 20 A solution of (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (50 mg, 113 μ mol, example 88) and 4-amino-3-fluorobenzonitrile (17 mg, 124 μ mol) in MeCN (0.7 mL) was subjected to a microwave irradiation at 120 °C for 15 min and at 150 °C for 15 min. Then, NaH


(60% dispersion in mineral oil, 20 mg) was added and the mixture was then stirred at RT for overnight. Then, H₂O (1.0 mL) was added and the mixture was extracted with EtOAc (2 x 1.5 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was then dissolved in a solution of 5 DMSO and MeOH (1:1, 0.1 mL). The solution mixture was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give a desired product, which was then dissolved in MeOH (1.0 mL). The solution was then washed through PL-HCO₃ MP-resin and the resin was washed with MeOH (2 x 0.5 mL). The combined washings were concentrated and dried under vacuum to 10 give the title compound as a colorless solid. MS (ESI, positive ion) m/z: 440 (M+H).



Example 90:

(R)-N-(4-Cyano-3-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide
15 To a solution of (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (70 mg, 158 μmol, example 88) in MeCN (0.5 mL) was added 4-amino-2-fluorobenzonitrile (65 mg, 475 μmol). The resulting mixture was then subjected to a microwave irradiation at 180 °C for 15 min.
20 Then, sodium hydride, 60% dispersion in mineral oil (22 μL, 475 μmol) was added and the mixture was stirred at RT for overnight. Then, H₂O (0.5 mL) was added and the solvents were removed. The residue was then dissolved in a solution of MeOH and DMSO (1:1, 1.0 mL). The solution mixture was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to 25 give the desired product, which was then dissolved in MeOH (1.0 mL). The solution was then washed through PL-HCO₃ MP resin (polymerlabs, 200 mg/6 mL tube) and the resin was washed with MeOH (2 x 0.5 mL). The combined

washings were then concentrated and dried under vacuum to give the title compound as a yellow solid. MS (ESI, positive ion) m/z: 440 (M+H).



Example 91:


- 5 **(R)-N-(4-Cyano-2-methylphenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

To a solution of 4-amino-3-methylbenzonitrile (30 mg, 226 µmol) in THF (2 mL) was added sodium hydride, 60% dispersion in mineral oil (15 µL, 339 µmol). The resulting mixture was then stirred at RT for 2 min. Then, (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (100 mg, 226 µmol, example 88) was added and the mixture was stirred at RT for overnight. Then, 26 mg of NaH (60% dispersion in mineral oil) was added and the mixture was stirred at RT for overnight. Then, H₂O (0.2 mL) was added and the solvents were removed. The residue was then dissolved in a solution of DMSO and MeOH (1:1, 1.0 mL). The solution was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the desired product, which was dissolved in MeOH (1.0 mL). The solution was then washed through PL-HCO₃ MP resin (polymerlabs, 200 mg/6 mL tube) and the resin was then washed with MeOH (2 x 0.5 mL). The combined washings were then concentrated and dried under vacuum to give the title compound as a yellow solid.

10 MS (ESI, positive ion) m/z: 436 (M+H).

15


20



Example 92:

(R)-1-(4-(Trifluoromethyl)phenyl)-N-(6-(trifluoromethyl)pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a solution of (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroiso-
5 quinoline-2(1H)-carboxylate (70 mg, 158 µmol, example 88) in MeCN (0.5 mL)
was added 3-amino-6-(trifluoromethyl)pyridine (77 mg, 475 µmol). The resulting
mixture was then subjected to a microwave irradiation at 150 °C for 15 min and at
180 °C for 15 min. Then, sodium hydride, 60% dispersion in mineral oil (18 mg,
475 µmol) was added and the mixture was stirred at RT for overnight. Then, H₂O
10 (0.5 mL) was added and the solvents were removed. The residue was then
dissolved in a solution of MeOH and DMSO (1:1. 1.0 mL). The solution mixture
was purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA)
to give the desired product, which was then dissolved in MeOH (1.0 mL). The
solution was then washed through PL-HCO₃ MP resin (polymerlabs, 200 mg/6
15 mL tube) and the resin was washed with MeOH (2 x 0.5 mL). The combined
washings were then concentrated and dried under vacuum to give the title
compound as a yellow solid. MS (ESI, positive ion) m/z: 466 (M+H).




20 Example 93:

(R)-N-(3-Cyano-4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydro-isoquinoline-2(1H)-carboxamide


A solution of 5-amino-2-fluorobenzonitrile (62 mg, 452 µmol) and sodium hydride, 60% dispersion in mineral oil (26 mg, 678 µmol) in THF (1.0 mL) was stirred at RT for 15 min. Then, (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (100 mg, 226 µmol, example 88) was added and the mixture was stirred at RT for overnight. Then, a solution of H₂O and MeOH (1:1, 0.2 mL) was added and the solvents were removed. The residue

was then dissolved in a solution of DMSO and MeOH (1:1, 1 mL). The solution mixture was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the desired product, which was dissolved in MeOH (1 mL). The solution was then washed through PL-HCO₃ MP resin (polymerlabs, 5 200 mg/6 mL tube) and the resin was washed with MeOH (2 x 0.5 mL). The combined washings were then concentrated and dried under vacuum to give the title compound as an orange solid. MS (ESI, positive ion) m/z: 440 (M+H).




Example 94:

- 10 **(R)-N-(5-cyano-6-(2,2,2-trifluoroethoxy)pyridin-3-yl)-1-(4-(trifluoro-methyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**
To a solution of 5-amino-2-(2,2,2-trifluoroethoxy)nicotinonitrile (49 mg, 226 µmol) in THF (1.0 mL) was added sodium hydride, 60% dispersion in mineral oil (26 mg, 678 µmol). The resulting mixture was then stirred at RT for 15 min.
- 15 Then, (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (100 mg, 226 µmol, example 88) was added and the mixture was stirred at RT for overnight. Then, a solution of H₂O and MeOH (1:1, 0.2 mL) was added and the solvents were removed. The residue was then dissolved in a solution of DMSO and MeOH (1:1, 1 mL). The solution mixture was then 20 purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the desired product, which was dissolved in MeOH (1.0 mL). The solution was then washed through PL-HCO₃ MP resin (polymerlabs, 200 mg/6 mL tube) and the resin was washed with MeOH (2 x 0.5 mL). The combined washings were then concentrated and dried under vacuum to give the title compound as a 25 light yellow solid. MS (ESI, positive ion) m/z: 521 (M+H).

**Example 95:**

(R)-N-(4-(Dimethylcarbamoyl)phenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

- 5 To a solution of 4-amino-N,N-dimethylbenzamide (74 mg, 452 µmol) in THF (0.5 mL) was added sodium hydride, 60% dispersion in mineral oil (19 µL, 452 µmol). The resulting mixture was then subjected to a microwave irradiation at 150 °C for 5 min. Then, (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (100 mg, 226 µmol, example 88) was added and the mixture was stirred at RT for overnight. Then, H₂O (0.5 mL) was added slowly and the mixture was extracted with EtOAc (2 x 1 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was dissolved in DMSO (1.0 mL) and the solution mixture was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/ H₂O 0.1% TFA) to give the desired compound,
- 10 which was dissolved in MeOH (1.0 mL). Then, the solution was washed through PL-HCO₃ MP-Resin (200 mg per 6 mL tube) and the resin was washed with MeOH (2 x 0.3 mL). The combined washings were concentrated and dried under vacuum to give the title compound as light brown solid. MS (ESI, positive ion) m/z: 468 (M+H).
- 15




20

Example 96:

(R)-N-(6-Cyanopyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


To a solution of 5-amino-2-pyridinecarbonitrile (81 mg, 678 µmol) in THF (1.0 mL) was added sodium hydride, 60% dispersion in mineral oil (17 mg, 452 µmol). The resulting mixture was then subjected to a microwave irradiation at 160 °C for 15 min. Then, (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (100 mg, 226 µmol, example 88) was added and the mixture was stirred at RT for overnight. Then, H₂O (0.5 mL) was added and the mixture was extracted with EtOAc (2 x 1 mL). The combined organic extracts were dried over MgSO₄ and concentrated. The residue was dissolved in DMSO (1.0 mL) and the solution was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/ H₂O 0.1% TFA) to give the desired compound, which was dissolved in MeOH (1.0 mL). The solution was then washed through PL-HCO₃ MP-Resin (200 mg per 6 mL tube) and the resin was washed with MeOH (2 x 0.3 mL). The combined washings were concentrated and dried under vacuum to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 423 (M+H).
15 (M+H).



Example 97:

(R)-N-(6-chloropyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

20




Step 1: 4-Nitrophenyl 6-chloropyridin-3-ylcarbamate

To a solution of 5-amino-2-chloropyridine (162 mg, 1262 µmol) and DIEA (275 µL, 1578 µmol) in DCM (6.0 mL) was added 4-nitrophenyl chloroformate (212

mg, 1052 µmol). The resulting mixture was then stirred at RT for overnight. A yellow precipitation was observed. The yellow solid was collected by filtration to give the title compound as a light yellow solid, which was used in the next step without purification requirement.

5 Step 2: **(R)-N-(6-Chloropyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

A solution of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (50 mg, 180 µmol, example 87, Step3) and 4-nitrophenyl 6-chloropyridin-3-yl-carbamate (119 mg, 406 µmol) in toluene (1.0 mL) was subjected to a microwave 10 irradiation at 180 °C for 30 min. Then, DMSO (1.0 mL) was added and the solution mixture was filtered. The filtrate was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/ H₂O 0.1% TFA) to give the title compound, which was dissolved in MeOH (1.0 mL). Then, the solution was washed through PL-HCO₃ MP-Resin (200 mg per 6 mL tube) and the resin was washed with MeOH 15 (2 x 0.3 mL). The combined washings were concentrated and dried under vacuum to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 432 (M+H).




Example 98:

20 **(R)-N-(2-Methoxyethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroiso-quinoline-2(1H)-carboxamide**

To a solution of (R)-4-nitrophenyl 1-(4-(trifluoromethyl)phenyl)-3,4-dihydroiso-quinoline-2(1H)-carboxylate (170 mg, 384 µmol, example 88) in MeCN (0.2 mL) was added 2-methoxyethylamine (36.7 µL, 423 µmol). The resulting mixture was 25 then subjected to a microwave irradiation at 180 °C for 15 min. Then, 2-methoxyethylamine (0.02 mL) was added and the mixture was subjected to a microwave irradiation at 180 °C for an additional 15 min. Then, the solvent was removed and the residue was dissolved in a solution of DMSO and MeOH (1:1, 1.0 mL). The


solution mixture was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the desired product, which was then dissolved in MeOH (2.0 mL). The solution was washed through PL-HCO₃ MP resin (200 mg per 6 mL tube) and the resin was washed with MeOH (2 x 0.5 mL). The combined washings were then concentrated and dried under vacuum to give the title compound as a yellow solid. MS (ESI, positive ion) m/z: 379 (M+H).



Example 99:

(R)-1-(4-(trifluoromethyl)phenyl)-N-((S)-1,1,1-trifluoropropan-2-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a solution of (S)-1,1,1-trifluoropropan-2-amine (90.9 μ l, 884 μ mol) in 4:1 CH₂Cl₂:THF (5 mL) is added di(1H-imidazol-1-yl)methanone (215 mg, 1326 μ mol) and the flask was sealed and stirred at room temperature for 1.5 h. (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (307 mg, 1105 μ mol) was added and the reaction stirred at 2 h. The reaction was directly purified by reverse phase HPLC to give (R)-1-(4-(trifluoromethyl)phenyl)-N-((S)-1,1,1-trifluoropropan-2-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as a white powder. MS (ESI pos. ion) m/z: 417 (M+1).




Example 100:

(R)-N-Butyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


To a solution of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (198 mg, 714 μ mol, Example 87, Step 3) in DCM (2.0 mL) was added n-butyl iso-

cyanate (88 μ L, 785 μ mol). The resulting mixture was then stirred at RT for overnight. Then, the solvent was removed and the residue was mixed with silica gel. The solid mixture was the purified by silica gel flash column chromatography using ISCO instrument (solid loading, 0%-100% EtOAc/hexane) to give the title compound as a colorless solid. MS (ESI, positive ion) m/z: 377 ($M+H$).



Example 101:

(R)-N-(2,2,2-Trifluoroethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide



Step 1: 4-Nitrophenyl 2,2,2-trifluoroethylcarbamate

To a solution of 4-nitrophenyl chloroformate (4578 mg, 22715 μ mol) and DIEA (2638 μ L, 15143 μ mol) in DCM (100 mL) at 0 °C was added 2,2,2-trifluoroethylamine (1500 μ L, 15143 μ mol). The resulting mixture was then stirred at 0 °C for 4 h and at RT for overnight. Then, the mixture was concentrated and a white precipitation was observed. The mixture was then filtered to give the title compound as a white solid, which was used in the next step without purification requirement.


Step 2: (R)-N-(2,2,2-Trifluoroethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a solution of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (150 mg, 541 μ mol, Example 87, Step 3) in MeCN (2 mL) was added 4-

nitrophenyl 2,2,2-trifluoroethylcarbamate (286 mg, 1082 µmol). The resulting mixture was then subjected to a microwave irradiation at 100 °C for 15 min. Then, the solvent was removed and the residue was dissolved in a solution of DMSO and MeOH (1:1, 2 mL). The solution mixture was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give the desired product, which was then dissolved in MeOH (2.0 mL). The solution was then washed through PL-HCO₃ MP-resin and the resin was then washed with MeOH (2 x 2 mL). The combined washings were concentrated and dried under vacuum to give the title compound as a light yellow solid. MS (ESI, positive ion) m/z: 403 (M+H).

General procedure of examples (102-103)

To a solution of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (180 µmol, example 87, Step3) and DIEA (31 µL, 180 µmol) in DCM (1 mL) was added isocyanate (25 µL, 216 µmol). The resulting mixture was then stirred at RT for overnight. Then, the mixture was mixed with silica gel (1:1) and the solid mixture was purified by silica gel flash column chromatography using ISCO instrument (solid loading, 0%-100% EtOAc/hexane) to provide the title compound.

Example	Structure	Name	MS (ESI, positive ion) M+H
102		(R)-N-tert-butyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide	377


103		(R)-N-(4-cyanophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide	422
-----	--	--	-----

General procedure of examples (104-108)


To a solution of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline 5 (252 µmol, example 87, Step 3) in DCM (1.7 mL) was added isocynate (278 µmol). The resulting mixture was then stirred at RT for overnight. Then, the solvent (DCM) was removed and the residue was dissolved in a solution of MeOH and DMSO (1:1, 1.0 mL). The solution mixture was then purified by preparative HPLC (0%-100% MeCN 0.1% TFA/H₂O 0.1% TFA) to give a desired product, 10 which was then dissolved in MeOH (1.0 mL). The solution was then washed through PL-HCO₃ MP resin (polymerlabs, 200 mg/6 mL tube). The resin was then washed with MeOH (2 x 0.5 mL). The combined washings were then concentrated and dried under vacuum to give the title compound.

104		(R)-N-(2-(trifluoromethyl)phenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide	465
105		(R)-N-(3-cyanophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide	422

106		(R)-N-(2-cyanophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide	422
107		(R)-N-(3-(trifluoromethyl)phenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide	465
108		(R)-N-(4-fluoro-3-(trifluoromethyl)phenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide	483

**Example 109:**

(R)-N-(Thiazol-2-yl)-1-(4-trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

**Step 1: 4-Nitrophenyl thiazol-2-yl-ylcarbamate:**

To a stirred mixture of thiazol-2-amine (1.00 g, 9.99 mmol) and pyridine (1.21 mL, 15.0 mmol) in DCM (30 mL) at 0 °C was added a solution of 4-nitrophenyl


chloroformate (2.42 g, 12.0 mmol) in DCM (10 mL). The reaction mixture was continued to stir overnight, concentrated and taken up in H₂O. The tan solid was filtered, air dried, and used in the next step. MS (ESI, positive ion) m/z: 266 (M+1).

5

Step 2: (R)-N-(Thiazol-2-yl)-1-(4-trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

A mixture of 4-nitrophenyl thiazol-2-ylcarbamate (0.0957 g, 0.361 mmol) and (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.100 g, 0.361 mmol) in p-dioxane (2 mL) was heated at 110 °C in 2h. The reaction mixture was cooled, concentrated and purified by ISCO (30% EtOAc/hexanes) to give the title compound. MS (ESI, positive ion) m/z: 404 (M+1).


15




Example 110:

(R)-N-(1-(Trifluoromethyl)cyclopropyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

A mixture of 1-(trifluoromethyl) cyclopropane carboxylic acid (0.36 g, 2.3 mmol), 20 DIEA (0.41 mL, 2.3 mmol), and diphenylphosphoryl azide (0.51 mL, 2.3 mmol) in p-dioxane (8 mL) was stirred at RT for 1h. (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.500 g, 1.8 mmol) was added and the mixture was heated to reflux for 16 h. The mixture was cooled, concentrated and purified by ISCO (0-40% EtOAc/hexanes) to give the title compound. MS (ESI, positive ion) m/z: 429 (M+1).


**Example 111:****(R)-N-Cyclopropyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

- 5 To a stirred solution of triphosgene (0.05 mL, 0.4 mmol) in DCM (1 mL) at 0 °C was added a mixture of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.200 g, 0.7 mmol) and diisopropylethylamine (0.1 mL, 0.8 mmol) in DCM (1 mL). After stirring at 0 °C in 30 min, cyclopropylamine (0.04 mL, 0.7 mmol) was added and the mixture was continued to stir at RT for 24 h. H₂O was 10 added and layers were separated. The organic extracts were dried over MgSO₄, concentrated and purified by ISCO (20% EtOAc/hexanes) to give the white solid title compound. MS (ESI, positive ion) m/z: 361 (M+1).

**Example 112:****(R)-(4,4-Difluorocyclohexyl)(1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-yl) methanone.**

- To a stirred mixture of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.205 g, 0.74 mmol), DIEA (0.14 mL, 0.81 mmol), and 4,4-difluorocyclohexanecarboxylic acid (0.13 g, 0.81 mmol) in DMF (4 mL) was added 20 HATU (0.31 g, 0.81 mmol). The reaction mixture was continued to stir for 2h, quenched with H₂O, and extracted with DCM (3x). The organic extracts were

dried over Na_2SO_4 , concentrated and purified by ISCO (20% EtOAc/hexanes) to give the white solid. MS (ESI, positive ion) m/z: 424 ($M+1$).




Example 113:

- 5 **(R)-N-Isobutyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.**

To a stirred mixture of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.100 g, 0.36 mmol) and DIEA (0.082 mL, 0.47 mmol) in THF (2 mL) was added 4-nitrophenyl chloroformate (0.080 g, 0.40 mmol). The reaction mixture was stirred for 30 min, 2-methylpropan-1-amine (0.036 mL, 0.36 mmol) was added and stirring was continued for 14h, then heated at reflux for 24 h, cooled, taken up in H_2O , extracted with EtOAc (3x), dried over MgSO_4 , concentrated and purified by ISCO (20% EtOAc/hexanes) to give the title compound. MS (ESI, positive ion) m/z: 377 ($M+1$).

10
15




Example 114:

- (R)-N-Cyclopentyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.**

20 A mixture of (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.100 g, 0.361 mmol), N-ethyl-N-isopropylpropan-2-amine (0.0754 mL, 0.433 mmol), and isocyanatocyclopentane (0.0421 g, 0.379 mmol) in DCM (4 mL) was

stirred at RT for 24 h. The mixture was concentrated and purified by ISCO (20% EtOAc/hexanes) to give the white solid. MS (ESI, positive ion) m/z: 389 (M+1).




5 **Example 115:**

(R)-N-(Tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.


This compound was prepared under similar conditions as described for example 111. MS (ESI, positive ion) m/z: (405 (M+1).

10



Example 116:


(R)-N-Cyclopentyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide. This compound was prepared under similar conditions as
15 described for example 111. MS (ESI, positive ion) m/z: 391 (M+1).



Example 117:

(R)-N-(Pyridazin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 111. MS (ESI, positive ion) m/z: 399 (M+1).




5

Example 118:

((R)-Tetrahydrofuran-2-yl)((R)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-yl) methanone.


This compound was prepared under similar conditions as described for example 10 112. MS (ESI, positive ion) m/z: 376 (M+1).



Example 119:


(R)-N-((R)-3,3-Dimethylbutan-2-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 112. MS (ESI, positive ion) m/z: 405 (M+1).

**Example 120:**

(R)-N-((S)-3,3-Dimethylbutan-2-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.


- 5 This compound was prepared under similar conditions as described for example 112. MS (ESI, positive ion) m/z: 405 (M+1).

**Example 121:**

- 10 (R)-N-((S)-3,3-Methylbutan-2-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.


This compound was prepared under similar conditions as described for example 112. MS (ESI, positive ion) m/z: 391 (M+1).

15

**Example 122:**

(R)-N-(1-(Methylcyclopropyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 110. MS (ESI, positive ion) m/z: 375 (M+1).




5 **Example 123:**

(R)-N-(2-Methylbut-3-yn-2yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 111. MS (ESI, positive ion) m/z: 387 (M+1).


10



Example 124:


(R)-N-(1-(Cyanocyclopropyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

15 This compound was prepared under similar conditions as described for example 110. MS (ESI, positive ion) m/z: 386 (M+1).

**Example 125:**


(S)-N-tert-Butyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

- 5 This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 377 (M+1).

**Example 126:**

- 10 **(R)-N-(1,2,3,4-Tetrahydronaphthalen-1-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.**

This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 451 (M+1).




15

Example 127:

(R)-Ethyl 4-(1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-2-carboxamido)benzoate.


This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 469 (M+1).



5

Example 128:

(*R*)-4-(1-(4-(Trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-2-carboxamido)benzoic acid. A mixture of (*R*)-ethyl 4-(1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-2-carboxamido)benzoate (0.210 g, 0.4 mmol) and NaOH (0.4 mL, 2 mmol) in EtOH (5 mL) was stirred at RT in 24 h. The mixture was concentrated, taken up in H₂O and neutralized with 10% HCl. The white solid was collected by filtration and dried. MS (ESI, positive ion) m/z: 441 (M+1).




15

Example 129:


(*S*)-N-(Pyrimidin-5-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 110. MS (ESI, positive ion) m/z: 399 (M+1).

**Example 130:**


(R)-N-((S)-2,3-Dihydro-1H-inden-1-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

- 5 This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 437 (M+1).

**Example 131:**

- 10 (R)-N-((R)-2,3-Dihydro-1H-inden-1-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 437 (M+1).




15

Example 132:

(R)-N-((R)-1-Cyclohexylethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 431 (M+1).




5 **Example 133:**

(R)-N-((S)-1-cyclohexylethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 431 (M+1).


10



Example 134:


15 **(R)-Ethyl 2-(1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-2-carboxamido)propanoate.**

This compound was prepared under similar conditions as described for example 114. MS (ESI, positive ion) m/z: 421 (M+1).

**Example 135**

5 **(R)-2-(1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline-2-carboxamido)propanoic acid.**


This compound was prepared under similar conditions as described for example 128. MS (ESI, positive ion) m/z: 393 (M+1).



10 **Example 136:**


(S)-1-(4-(trifluoromethyl)phenyl)-N-((S)-1,1,1-trifluoropropan-2-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 109. MS (ESI, positive ion) m/z: 417 (M+1).

**Example 137:**


(R)-1-(4-(trifluoromethyl)phenyl)-N-((R)-1,1,1-trifluoropropan-2-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

- 5 This compound was prepared under similar conditions as described for example 109. MS (ESI, positive ion) m/z: 417 (M+1).

**Example 138:**


(R)-N-(Dimethylcyclopropyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 110. MS (ESI, positive ion) m/z: 389 (M+1).

**Example 139:**

(R)-N-(1,1,1-Trifluoro-2-methylpropan-2-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.


- 5 This compound was prepared under similar conditions as described for example 110. MS (ESI, positive ion) m/z: 431 (M+1).

**Example 140:**


(R)-N-(2,2-Difluorocyclopropyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide.

This compound was prepared under similar conditions as described for example 110. MS (ESI, positive ion) m/z: 397 (M+1).

15

**Example 141:**

(R)-Pyrrolidin-2-yl((R)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-yl) methanone. To a stirred mixture of (R)-1-(tert-butoxy-carbonyl)pyrrolidine-2-carboxylic acid (0.078 g, 0.36 mmol), (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.100 g, 0.36 mmol) and DIEA (0.047 g, 0.36 mmol) in DMF (2 mL) was added HATU (0.14 g, 0.36 mmol). The reaction mixture was continued to stir overnight at RT, hydrolyzed with H₂O, extracted with DCM (3x), dried over Na₂SO₄, concentrated and purified by ISCO (20% EtOAc/hexanes) to give the white solid which was dissolved in p-dioxane (2 mL) and added 1 mL of 4M HCl in p-dioxane. The mixture was continued to stir at RT overnight, concentrated, taken up in H₂O, and neutralized by saturated aqueous NaHCO₃, and filtered the light yellow solid which was dried by air. MS (ESI, positive ion) m/z: 375 (M+1).




15


Example 142:

(S)-Pyrrolidin-2-yl((R)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-yl) methanone.

This compound was prepared under similar conditions as described for example 141. MS (ESI, positive ion) m/z: 375 (M+1).

**Example 143:**


3-Methyl-N-(pyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide



5


Step 1: 1-Bromo-2-(2,2-dimethoxypropyl)benzene

2-Bromophenylacetone (4.45000 g, 21 mmol) was dissolved in 10 mL MeOH and trimethyl orthoformate (11 mL, 104 mmol) and hydrogen chloride, 4.0M solution in 1,4-dioxane (5.2 mL, 21 mmol) were added. The mixture was stirred for 1h and neutralized carefully with DIEA (pH 7). The mixture was evaporated and filtered through a plug of silica providing 1-bromo-2-(2,2-dimethoxypropyl)benzene as a colorless liquid.




15 **Step 2: (R)-N-(4-(Trifluoromethyl)benzylidene)-2-methylpropane-2-sulfinamide**

(R)-2-methylpropane-2-sulfinamide (31.6g, 260 mmol), 4-(trifluoromethyl)-benzaldehyde (22.7g, 130 mmol) and cupric sulfate (83.2g, 521 mmol) were added to 300 mL DCM and the mixture was stirred for 36h. The mixture was filtered through a pad of celite and the filtrate was evaporated and purified via 20 silica plug filtration. (R)-N-(4-(trifluoromethyl)benzylidene)-2-methylpropane-2-sulfinamide was obtained as a white solid.



Step 3: (R)-N-((2-(2,2-Dimethoxypropyl)phenyl)(4-(trifluoromethyl)phenyl)methyl)-2-methylpropane-2-sulfinamide


- 5 1-Bromo-2-(2,2-dimethoxypropyl)benzene (4.00 g, 15.43 mmol) was dissolved in 150 mL THF and tert-butyllithium (18.160 mL, 30.871 mmol) was added dropwise at -78 °C. After addition a solution of (R)-N-(4-(trifluoromethyl)-benzylidene)-2-methylpropane-2-sulfinamide (8.5608 g, 30.871 mmol) in THF (10 mL) was added and stirring was continued for 10 min in the cold. The reaction
 10 was hydrolyzed with 50 mL water and the mixture was warmed to RT. TLC showed a 1:1 mixture of the 2 diastereomers. No attempt was undertaken to separate the isomers. Glass column chromatography provided the title compound as a mixture of diastereomers. MS (ESI, positive ion) m/z: 412.0 (M+1; hydrolysis of the dimethyl acetal to the corresponding ketone).



15

- Step 4: 3-Methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline**
 (R)-N-((2-(2,2-Dimethoxypropyl)phenyl)(4-(trifluoromethyl)phenyl)methyl)-2-methylpropane-2-sulfinamide (6.650000 g, 15 mmol) was dissolved in 150 mL of DCE and hydrogen chloride, 4.0m solution in 1,4-dioxane (11 mL, 44 mmol) was added. Stirring was continued for 20 min and trimethyl orthoformate (8.0 mL, 73 mmol) and sodium triacetoxyborohydride (11 g, 51 mmol) were added. TLC-analysis showed a complex product mixture but LCMS showed the formation of


some desired product along with a lot of fully aromatized product. The crude was semi-prurified via glass colm chrom and used w/o further clean up in the next steps. MS (ESI, positive ion) m/z: 292.0 (M+1)



5 **Step 5: 3-Methyl-N-(pyridin-3-yl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

3-Methyl-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (0.100 g, 0.343 mmol, semi-pure from the previous reaction) was dissolved in 3 mL DCM and DIEA (0.119 mL, 0.687 mmol) and 3-isocyanatopyridine (79 mg, 0.659

10 mmol) were added. Stirring was continued for 2h and the mixture was hydrolyzed with 3 drops of water, diluted with 3 mL DMF and injected on the HPLC for purification. The title compound was obtained. No stereochemical assignment was performed.




15 **Example 144:**

N-(4-Cyanophenyl)-3-methyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide



The reaction was performed under similar conditions as described for example

143. The title compound were obtained. No stereochemical assignment was

20 performed.


**Example 145:**

N-tert-Butyl-5-methoxy-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


5 Synthetic Scheme:

Step 1: N-(2-Methoxyphenethyl)-4-(trifluoromethyl)benzamide

- To a 100 mL round-bottomed flask was added 2-(2-methoxyphenyl)ethylamine (1510 mg, 10 mmol, Aldrich), pyridine (10 mL), 4-(trifluoromethyl)benzoyl chloride (2 mL, 10 mmol, Fluka). The reaction mixture was stirred at RT for 1 h.
- 5 The reaction mixture was diluted with water (20 mL) and extracted with EtOAc (2 x 50 mL). The organic extract was washed with saturated NaHCO₃ (20 mL), saturated NaCl (20 mL), dried over Na₂SO₄, filtered and concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 20% EtOAc/CH₂Cl₂ to give N-(2-methoxyphenethyl)-4-(trifluoromethyl)benzamide.
- 10 MS (ESI pos. ion) *m/z*: 324 (M+1).

**Step 2: 5-Methoxy-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline**

- To a 100 mL round-bottomed flask was added N-(2-methoxyphenethyl)-4-(trifluoromethyl)benzamide (1.41 g, 4361 µmol, from step 1), toluene (30 mL), phosphorus pentoxide (1077 µL, 17445 µmol, Aldrich). The reaction mixture was stirred at reflux for overnight (ca 18 h). The mixture was cooled down to RT and poured it to ice (ca 100 g). The mixture was neutralized with KOH until pH 12. The mixture was extracted with EtOAc (2 x 100 mL). The organic extract was
- 15 washed with saturated NaCl (50 mL), dried over Na₂SO₄, filtered and concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 20% EtOAc/hexanes to give 5-methoxy-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline. MS (ESI pos. ion) *m/z*: 306 (M+1).
- 20




Step 3: 5-Methoxy-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline


To a 50 mL round-bottomed flask was added 5-methoxy-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (189 mg, 619 µmol, from step 2), MeOH (3 mL), sodium borohydride (21.8 µL, 619 µmol, Aldrich). The reaction mixture was stirred at RT for 1 h. The reaction mixture was diluted with water (5 mL) and extracted with EtOAc (2 × 20 mL). The organic extract was washed with saturated NaCl (5 mL), dried over Na₂SO₄, filtered and concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 80% EtOAc/hexanes to give 5-methoxy-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline. MS (ESI pos. ion) *m/z*: 308 (M+1).

Step 4: N-tert-Butyl-5-methoxy-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a 50 mL round-bottomed flask was added 5-methoxy-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (44 mg, 143 µmol, from step 3), CH₂Cl₂ (2 mL), t-butylisocyanate (16 µL, 143 µmol, Aldrich). The solution was stirred at RT for 5 h. The solvent was removed in vacuo and the residue was purified by silica gel chromatography, eluting with 20% EtOAc/hexanes to give N-tert-butyl-5-methoxy-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide. MS (ESI pos. ion) *m/z*: 407 (M+1).

**Example 146:**


5-Bromo-N-tert-butyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide



5


Step 1: N-(2-Bromophenethyl)-4-(trifluoromethyl)benzamide

To a 100 mL round-bottomed flask was added 2-bromophenethylamine (1800 mg, 9 mmol, Aldrich), pyridine (5 mL), 4-(trifluoromethyl)benzoyl chloride (2 mL, 9 mmol, Fluka). The reaction mixture was stirred at RT for 1 h. The reaction mixture was diluted with water (30 mL) and extracted with EtOAc (2 x 50 mL). The organic extract was washed with water (30 mL), saturated NaCl (30 mL), dried over Na_2SO_4 , filtered and concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 30% EtOAc/hexanes to give N-(2-bromophenethyl)-4-(trifluoromethyl)benzamide. MS (ESI pos. ion) m/z : 372 (M+1).



Step 2. 5-Bromo-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline


- To a 100 mL round-bottomed flask was added N-(2-bromophenethyl)-4-(trifluoromethyl)benzamide (1.6 g, 4299 µmol, from step 1), toluene (50 mL), phosphorus pentaoxide (1061 µL, 17196 µmol, Aldrich). The reaction mixture was stirred at reflux for 5 h. The mixture was cooled down and then poured into ice (ca 50 g), and extracted with EtOAc (2 x 50 mL). The organic extract was washed with water (30 mL), saturated NaCl (30 mL), dried over Na₂SO₄, filtered and concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 10% EtOAc/hexanes to give 5-bromo-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline. MS (ESI pos. ion) *m/z*: 354 (M+1).



Step 3: 5-Bromo-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline


- To a 100 mL round-bottomed flask was added 5-bromo-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline (1.18 g, 3332 µmol, from step 2), MeOH (5 mL), sodium borohydride (117 µL, 3332 µmol, Aldrich). The reaction mixture was stirred at RT for 1 h. The reaction mixture was diluted with water (30 mL) and extracted with EtOAc (2 x 40 mL). The organic extract was washed with water (20 mL), saturated NaCl (20 mL), dried over Na₂SO₄, filtered and concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with

40% EtOAc/hexanes to give 5-bromo-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline. MS (ESI pos. ion) *m/z*: 356 (M+1).



5 **Step 4. 5-Bromo-N-tert-butyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

To a 50 mL round-bottomed flask was added 5-bromo-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (44 mg, 124 µmol, from step 3), CH₂Cl₂ (2 mL), 2-isocyanato-2-methylpropane (12 mg, 124 µmol, Aldrich). The reaction mixture was stirred at RT for 2 h. The solvent was removed in vacuo and the residue was purified by silica gel chromatography, eluting with 50% EtOAc/hexanes to give 5-bromo-N-tert-butyl-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide. MS (ESI pos. ion) *m/z*: 455 (M+1).




15 **Example 147:**

5-bromo-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide


To a 50 mL round-bottomed flask was added 5-bromo-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (44 mg, 124 µmol, from step 3, example 146), CH₂Cl₂ (2 mL), 4-fluorophenyl isocyanate (17 mg, 124 µmol, Aldrich). The reaction mixture was stirred at RT for 1 h. The solvent was removed in vacuo and the residue was purified by silica gel chromatography, eluting with 50%

EtOAc/hexanes to give 5-bromo-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide. MS (ESI pos. ion) *m/z*: 493 (M+1).




5 **Example 148:**

(R)-N-(4-Fluorophenyl)-N-(2-methoxyethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide



Step 1: 4-Fluoro-N-(2-methoxyethyl)aniline


- 10 To a 50 mL round-bottomed flask was added 4-fluoroaniline (383 µL, 3971 µmol, Aldrich), 2-bromoethyl methyl ether (552 µL, 3971 µmol, Aldrich), sodium carbonate (667 mg, 7943 µmol), DMF (2 mL). The reaction mixture was stirred at 80 °C for overnight (ca 17 h). The mixture was cooled down to RT. The reaction mixture was diluted with water (30 mL) and extracted with EtOAc (2 x 40 mL).
- 15 The organic extract was washed with water (20 mL), saturated NaCl (20 mL), dried over Na₂SO₄, filtered and concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 30% EtOAc/hexanes to give 4-fluoro-N-(2-methoxyethyl)benzenamine. MS (ESI pos. ion) *m/z*: 170 (M+1).



Step 2: N-(4-Fluorophenyl)-N-(2-methoxyethyl)-1H-imidazole-1-carboxamide


To a 50 mL round-bottomed flask was added 4-fluoro-N-(2-methoxyethyl)-benzenamine (512 mg, 3026 µmol, from step 1), 1,1'-carbonyldiimidazole (736 mg, 4539 µmol, Aldrich), THF (2 mL). The reaction mixture was stirred at 80 °C for 28 h. The solvent was removed in vacuo and the residue was purified by silica gel chromatography, eluting with 80% EtOAc/hexanes to give N-(4-fluorophenyl)-N-(2-methoxyethyl)-1H-imidazole-1-carboxamide. MS (ESI pos. ion) *m/z*: 264 (M+1).

10




Step 3: 1-((4-Fluorophenyl)(2-methoxyethyl)carbamoyl)-3-methyl-1H-imidazol-3-ium iodide

To a 50 mL round-bottomed flask was added N-(4-fluorophenyl)-N-(2-methoxyethyl)-1H-imidazole-1-carboxamide (91 mg, 346 µmol, from step 2), acetonitrile (2 mL), methyl iodide (86 µL, 1383 µmol, Aldrich). The solution was stirred at RT for 24 h. The solvent was removed in vacuo to give the product, which was used for the next step reaction without purification.




Step 4: (R)-N-(4-Fluorophenyl)-N-(2-methoxyethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a 50 mL round-bottomed flask was added 1-((4-fluorophenyl)(2-methoxyethyl)carbamoyl)-3-methyl-1H-imidazol-3-ium iodide (70 mg, 173 µmol, from step 3), (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (24 mg, 87 µmol), CH₂Cl₂ (1 mL), triethylamine (12 µL, 87 µmol). The reaction mixture was stirred at 40 °C for 28 h. The solvent was removed in vacuo and the residue was purified by silica gel chromatography, eluting with 30% EtOAc/hexanes to give (R)-N-(4-fluorophenyl)-N-(2-methoxyethyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide. MS (ESI pos. ion) *m/z*: 473 (M+1).




Example 149:

(R)-N-(2-Aminoethyl)-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide




Step 1: 2-(2-(4-Fluorophenylamino)ethyl)isoindoline-1,3-dione

- To a 50 mL round-bottomed flask was added n-(2-bromoethyl)phthalimide, 95% (508 mg, 1999 µmol, Aldrich), 4-fluoroaniline (193 µL, 1999 µmol, Aldrich),
- 5 DMF (2 mL), sodium carbonate (336 mg, 3999 µmol). The reaction mixture was stirred at 80 °C for overnight. The reaction mixture was diluted with water (20 mL) and extracted with EtOAc (2 x 40 mL). The organic extract was washed with water (20 mL), saturated NaCl (20 mL), dried over Na₂SO₄, filtered and concentrated in vacuo and the residue was purified by silica gel chromatography,
- 10 eluting with 30% EtOAc/hexanes to give 2-(2-(4-fluorophenylamino)ethyl)isoindoline-1,3-dione. MS (ESI pos. ion) *m/z*: 285 (M+1).



- Step 2: N-(2-(1,3-Dioxoisoindolin-2-yl)ethyl)-N-(4-fluorophenyl)-1H-imidazole-1-carboxamide**
- To a 50 mL round-bottomed flask was added 2-(2-(4-fluorophenylamino)ethyl)isoindoline-1,3-dione (204 mg, 718 µmol, from step 1), 1,1'-carbonyldiimidazole (175 mg, 1076 µmol, Aldrich), THF (2 mL). The reaction mixture was stirred at 80 °C for 24 h. The solvent was removed in vacuo and the residue was purified by
- 20 silica gel chromatography, eluting with 80% EtOAc/hexanes to give


N-(2-(1,3-dioxoisooindolin-2-yl)ethyl)-N-(4-fluorophenyl)-1H-imidazole-1-carboxamide. MS (ESI pos. ion) *m/z*: 379 (M+1).



5 Step 3: **1-((2-(1,3-Dioxoisooindolin-2-yl)ethyl)(4-fluorophenyl)carbamoyl)-3-methyl-1H-imidazol-3-ium iodide**

To a 50 mL round-bottomed flask was added N-(2-(1,3-dioxoisooindolin-2-yl)ethyl)-N-(4-fluorophenyl)-1H-imidazole-1-carboxamide (110 mg, 291 µmol, from step 2), acetonitrile (2 mL), methyl iodide (72.4 µL, 1163 µmol, Aldrich).

10 The reaction mixture was stirred at RT for 24 h. The solvent was removed in vacuo to give the product (151 mg, 99.8% yield), which was used for the next step reaction without purification.




Step 4: **(R)-N-(2-(1,3-Dioxoisooindolin-2-yl)ethyl)-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

To a 50 mL round-bottomed flask was added (R)-1-(4-(trifluoromethyl)phenyl)-1,2,3,4-tetrahydroisoquinoline (80 mg, 290 µmol), 1-((2-(1,3-dioxoisooindolin-2-yl)ethyl)(4-fluorophenyl)carbamoyl)-3-methyl-1H-imidazol-3-ium iodide (151 mg, 290 µmol, from step 3), CH2Cl2 (2 mL). The reaction mixture was stirred at


40 °C for 48 h. The solvent was removed in vacuo and the residue was purified by silica gel chromatography, eluting with 30% EtOAc/hexanes to give (R)-N-(2-(1,3-dioxoisoindolin-2-yl)ethyl)-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide. MS (ESI pos. ion) *m/z*: 588

5 (M+1).




Step 5: (R)-N-(2-Aminoethyl)-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a 50 mL round-bottomed flask was added (R)-N-(2-(1,3-dioxoisoindolin-2-yl)ethyl)-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide (16 mg, 27 µmol, from step 4), EtOH (1 mL), hydrazine, anhydrous (3.5 µL, 109 µmol, Aldrich). The solution was stirred at RT for 18 h. The solvent was removed in vacuo and the residue was purified by silica gel chromatography, eluting with 10%MeOH/CH₂Cl₂ + 1% ammonia (37% in water) to give (R)-N-(2-aminoethyl)-N-(4-fluorophenyl)-1-(4-(trifluoromethyl)phenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide. MS (ESI pos. ion) *m/z*: 458 (M+1).




Example 150:

20 **1-(6-Chloropyridin-3-yl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**



Step 1: Benzyl 1-(6-chloropyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

To a pear-shaped flask was added 5-bromo-2-chloropyridine (440 mg, 2286 µmol), dry THF (10 mL), and an ice bath. After stirring for 5 min, a 2M solution of isopropylmagnesium chloride (1.15 mL, 2300 µmol) was added. The solution was allowed to stir in the ice bath for 2 h, then transferred to a solution of 3,4-dihydroisoquinoline (165 mg, 1258 µmol), dry THF (5 mL), and benzyl chloroformate (0.20 mL, 1350 µmol) that had been stirring in an ice bath for 15 min. The solution was allowed to warm to RT. After 1 h, LC-MS indicated only partial conversion. Another 2 eq of the Grignard were prepared as described above and added to the reaction. After a further 15 h, LC-MS shows no further progress. The reaction was poured into sat'd Rochelle's salt and extracted with EtOAc (3 X 20 mL). The combined organics were washed with brine and concentrated in vacuo and adsorbed onto a plug of silica gel and chromatographed through a Redi-Sep® pre-packed silica gel column (12 g), eluting with 0% to 20% EtOAc in hexanes, to provide crude benzyl 1-(6-chloropyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxylate as a yellow oil. The crude oil was carried into the next step.




20

Step 2: 1-(6-Chloropyridin-3-yl)-1,2,3,4-tetrahydroisoquinoline

To a round-bottomed flask was added crude benzyl 1-(6-chloropyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (170 mg, 449 µmol), CH_2Cl_2 (10 mL) and TFA (10 mL). The reaction was allowed to stir at RT. After 3 days, LC-MS


shows little progress. The reaction was concentrated in vacuo to remove CH₂Cl₂ and the resulting solution allowed to stir for another 3 days, then cautiously poured into 10% Na₂CO₃ and extracted with CH₂Cl₂ (3 X 20 mL). The combined organic layers were concentrated in vacuo and purified by reverse-phase preparative HPLC (Shimadzu) on a Phenomenex Gemini® column (5 micron, C18, 110 Å, Axia, 100x50 mm) eluting at 90 mL/min with an linear gradient of 10% to 100% MeCN (0.1%TFA) in water (0.1% TFA) over 20 min to give crude 1-(6-chloropyridin-3-yl)-1,2,3,4-tetrahydroisoquinoline as a white solid. The crude solid was carried into the next step. MS (ESI pos. ion) *m/z*: 245 (M+1).

10




Step 3: **1-(6-chloropyridin-3-yl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

To a round-bottomed flask was added crude 1-(6-chloropyridin-3-yl)-1,2,3,4-tetrahydroisoquinoline (7 mg, 29 µmol), CH₂Cl₂ (3 mL), and 1 drop of 1-fluoro-4-isocyanatobenzene. The solution was stirred overnight, then concentrated in vacuo and purified by reverse-phase preparative HPLC (Shimadzu) on a Phenomenex Gemini® column (10 micron, C18, 110 Å, Axia, 100x30 mm) eluting at 45 mL/min with an linear gradient of 10% to 100% MeCN (0.1%TFA) in water (0.1% TFA) over 10 min to give 1-(6-chloropyridin-3-yl)-N-(4-fluorophenyl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as a TFA salt. MS (ESI pos. ion) *m/z*: 382 (M+1).

**Example 151:**


N-(4-Fluorophenyl)-1-(6-(trifluoromethyl)pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide



5

Step 1: Benzyl 1-(6-(trifluoromethyl)pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxylate


To a solution of 5-bromo-2-(trifluoromethyl)pyridine (258 mg, 1144 µmol) in THF (5 mL) at 0 °C was added isopropylmagnesium chloride (572 µL, 1144 µmol) and the red/brown solution was stirred 2 h under N₂. To a solution of 3,4-dihydroisoquinoline (75 mg, 572 µmol) in THF (2 mL) at 0 °C was added benzyl chloroformate (90 µL, 629 µmol) and the reaction was warmed to RT and stirred 15 min. The solution of Grignard reagent was added to the iminium salt and the reaction warmed to RT and stirred 1 h. The reaction was quenched with saturated NH₄Cl (20 mL) and extracted with CH₂Cl₂ (3 x 20 mL). The combined organic layers were dried (MgSO₄) and concentrated to give a yellow oil. Purification by ISCO (12 g SiO₂, 0-50% EtOAc/hexane) gives benzyl 1-(6-(trifluoromethyl)pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxylate as a yellow oil.



Step 2: 1-(6-(Trifluoromethyl)pyridin-3-yl)-1,2,3,4-tetrahydroisoquinoline

To a solution of benzyl 1-(6-(trifluoromethyl)pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (90 mg, 218 µmol) in MeOH (3 mL) at RT under N₂

- 5 was added 10% palladium on carbon (46 mg, 44 µmol) and suspension was sparged with H₂ 15 min. The reaction was stirred under H₂ at RT 3.5 h. The reaction was filtered through a pad of Celite® (2 x 5 mL EtOAc wash) and concentrated to give 1-(6-(trifluoromethyl)pyridin-3-yl)-1,2,3,4-tetrahydroisoquinoline which was carried on to the next step without purification.




10

Step 3: N-(4-fluorophenyl)-1-(6-(trifluoromethyl)pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

- The crude 1-(6-(trifluoromethyl)pyridin-3-yl)-1,2,3,4-tetrahydroisoquinoline was dissolved in CH₂Cl₂ (1.0 mL) and 1-fluoro-4-isocyanatobenzene (32 µL, 273 µmol) was added. The reaction was stirred 17 h under N₂ at RT. The reaction was directly purified by HPLC to give N-(4-fluorophenyl)-1-(6-(trifluoromethyl)pyridin-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 416 (M+1).


20

**Example 152:****N-(4-Fluorophenyl)-1-(1H-indol-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide**

5


Step 1: 1-(1H-Indol-3-yl)-3,4-dihydroisoquinolin-2(1H)-yl)ethanone

To a solution of 3,4-dihydroisoquinoline (4.00 g, 30.5 mmol) and CH₃CN (50 mL) at 0 °C was added acetyl chloride (2.13 mL, 30.5 mmol) over 2 min. The reaction was stirred 30 min and 1H-indole (3.57 g, 30.5 mmol) and triethylamine (4.25 mL, 30.5 mmol) in CH₃CN (5 mL) were added and the reaction warmed to RT. The reaction was stirred under N₂ 3 h, then quenched with 10% HCl (100 mL), and extracted with CH₂Cl₂ (3 x 100 mL). The combined organic layers were washed with NaHCO₃ (100 mL), dried (Na₂SO₄), and concentrated to give the crude product as a white solid (9.2 g). The solid was dissolved in CH₂Cl₂ (50 mL) to give a suspension which was filtered and the solid collected. The white solid (3.55 g) was recrystallized from hot acetone (230 mL) and cooled at -20 °C overnight to give 1-(1H-indol-3-yl)-3,4-dihydroisoquinolin-2(1H)-yl)ethanone as a white crystalline solid.



Step 2: 1-(1H-Indol-3-yl)-1,2,3,4-tetrahydroisoquinoline

To a solution 1-(1-(1H-indol-3-yl)-3,4-dihydroisoquinolin-2(1H)-yl)ethanone (50 mg, 172 µmol) in MeOH (1.0 mL) was added 5 M HCl (1.0 mL, 5000 µmol). The reaction was heated to 100 °C under microwave 30 min. The reaction was concentrated to give 1-(1H-indol-3-yl)-1,2,3,4-tetrahydroisoquinoline, which was used in the next step without purification.



Step 3: N-(4-Fluorophenyl)-1-(1H-indol-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide

To a solution of 1-(1H-indol-3-yl)-1,2,3,4-tetrahydroisoquinoline (16.3 mg, 66 µmol) in CH₂Cl₂ (0.5 mL) at RT was added 1-fluoro-4-isocyanatobenzene (10 µL, 66 µmol). The reaction was stirred 48 h, the suspension dissolved in DMF and purified by HPLC to give N-(4-fluorophenyl)-1-(1H-indol-3-yl)-3,4-dihydroisoquinoline-2(1H)-carboxamide as a white solid. MS (ESI pos. ion) *m/z*: 386 (M+1).

Absolute stereochemistries of the examples were determined using Vibrational Circular Dichroism (VCD, Biotools, Inc.). Comparison of experimental VCD data with ab initio DFT calculations provide for the assigned absolute stereochemistry.

Assays

Luminescence readout assay for measuring intracellular calcium. Stable CHO cell lines expressing human TRPM8 were generated using tetracycline inducible T-

RExTM expression system from Invitrogen, Inc (Carlsbad, CA). In order to enable a luminescence readout based on intracellular increase in calcium (Le Poul et al., 2002), each cell line was also co-transfected with pcDNA3.1 plasmid containing jelly fish aequorin cDNA. Twenty four h before the assay, cells were seeded in 5 96-well plates and TRP channel expression was induced with 0.5 µg/ml tetracycline. On the day of the assay, culture media was removed and cells were incubated with assay buffer (F12 containing 30 mM HEPES for TRPM8 containing 15 µM coelenterazine (P.J.K, Germany) for 2 h. Potential antagonists were added and cells were incubated for 2.5 min prior to adding 10 agonist, 1 µM Icillin, or 1 min prior to addition of cold buffer (<10 °C). The luminescence was measured by a CCD camera based FLASH-luminometer built by Amgen, Inc. Compound activity was calculated using GraphPad Prism 4.01 (GraphPad Software Inc, San Diego, CA).

The following compounds exhibit IC₅₀ values of less than 5µM in the 15 above assay using the cold buffer and human TRPM8: Examples 1-22, 24-65, 70-72, 78, 87-107, 109, 111-124, 145-149.

Icillin Biochemical Challenge Model:

Male Sprague Dawley rats (275–500g, Harlan, n=4-6/treatment) were 20 administered icillin in 2% HPMC / 1% HPbCD at the following doses: 0.1, 0.3, 1.0, 3.0, 10.0 mg/kg, i.p.; 0.32, 1.0, 3.2, 10, 32 mg/kg, p.o. Spontaneous wet dog shakes were counted over 30 min post-icillin. Example compounds were tested (i.v., p.o.) to assess the ability to block the spontaneous wet dog shake phenomena induced by Icillin.

25

CCI model

Surgery – A chronic constriction injury (CCI) was produced as previously described (Bennett & Xie, 1988). Under gaseous anesthesia with a mixture of iso-flurane (3% for induction and 2% for maintenance) in O₂, the sciatic nerve was 30 exposed at the mid-thigh level proximal to the sciatic trifurcation. Four chromic gut ligatures (4-0) were tied loosely around nerve, 1-2 mm apart such that the vascular supply was not compromised.

- Behavioral testing – A behavioral test was performed to estimate cold-induced ongoing pain as previously described (Choi et al., 1994). The rat was placed under a transparent plastic cover on an aluminum plate (IITC PE34, Woodland, CA) which was kept at a cold temperature ($5 \pm 0.5^{\circ}\text{C}$). After 2 min of adaptation, the 5 cumulative duration of time that the rat lifted the foot off the plate for the next 5 min was measured. Foot lifts associated with locomotion or grooming were not counted. Seven to 9 days after the CCI surgery, baseline of the cold-induced ongoing pain was measured. Any rat showing a cold-induced ongoing pain less than 100sec out of 300sec observation period was eliminated from the study.
- 10 Twenty four h after the baseline measurement, test compound, positive control, morphine (2mg/kg, Sigma, St. Louis) or a vehicle (saline or 2%HPMC/1% Tween 80) was administered orally (test compound) or subcutaneously (morphine). Two hrs (test compound) or 30 mins (morphine) after the drug administration, the cold-induced ongoing pain was measured again.
- 15
- Chung model**
- Surgery - Spinal nerve ligation surgery was performed as previously described (Kim & Chung, 1992). Briefly, under gaseous anesthesia with a mixture of isoflurane (3% for induction and 2% for maintenance) in O_2 , the spinal nerve injury 20 was produced by ligating the left L5 and L6 spinal nerves taking special care to avoid any possible damage to the L4 spinal nerve or surrounding area. Additional treatments were performed to increase the development of mechanical allodynia. First, L5 spinal nerve was cut approximately 1 mm distal to the suture as described by Li et al. (2000). Second, immediately after ligation and cut, the L4 25 spinal nerve was lightly manipulated by slightly stretching it with a fine hooked glass rod and gently sliding the hook back and forth 20 times along the nerve as described by Lee et al. (2003). The whole surgery procedure from anesthesia to the clipping of the incised skin took at most 15 min.
- Behavioral testing - Two weeks later, mechanical sensitivity was measured by 30 determining the median 50% foot withdrawal threshold for von Frey filaments using the up-down method (Chaplan et al., 1994). The rats were placed under a plastic cover ($9 \times 9 \times 20 \text{ cm}$) on a metal mesh floor. The area tested was the

- middle glabrous area between the footpads of the plantar surface of the hind paw. The plantar area was touched with a series of 9 von Frey hairs with approximately exponentially incremental bending forces (von Frey values: 3.61, 3.8, 4.0, 4.2, 4.41, 4.6, 4.8, 5.0 and 5.2; equivalent to: 0.41, 0.63, 1.0, 1.58, 2.51, 4.07, 6.31, 10 and 15.8g). The von Frey hair was presented perpendicular to the plantar surface with sufficient force to cause slight bending, and held for approximately 3-4 seconds. Abrupt withdrawal of the foot (paw flinching, shaking or licking for more than 1 sec.) was recorded as a response. Any rat showing a mechanical threshold of more than 3.16g or less than 0.7g after surgery was eliminated from the study. After measuring basal threshold, test compound, positive control gabapentin (Sigma, St. Louis) or a vehicle (saline or 2%HPMC/1% Tween 80) was administered orally (test compound) or intraperitoneally (gabapentin). The measurement of the tactile threshold was reassessed at 1.5 and 2 hrs after drug administration.
- 15 Data - Since the von Frey filament set was calibrated on a logarithmic scale by the vendor (Stoelting) and our selection of 9 filaments for the up-down method was also based on near equal logarithmic intervals (Dixon et al., 1980), data were treated using logarithmic values in every aspect (statistical treatment as well as plotting). However, an equivalent gram value scale is labeled on the Y-axis of the figures for convenience. Data are expressed as mean ± standard error of the mean (S.E.M.).
- 20

For the treatment of TRP-M8-receptor-diseases, such as acute, inflammatory and neuropathic pain, dental pain, general headache, migraine, cluster headache, mixed-vascular and non-vascular syndromes, tension headache, 25 general inflammation, arthritis, rheumatic diseases, osteoarthritis, inflammatory bowel disorders, inflammatory eye disorders, inflammatory or unstable bladder disorders, psoriasis, skin complaints with inflammatory components, chronic inflammatory conditions, inflammatory pain and associated hyperalgesia and allodynia, neuropathic pain and associated hyperalgesia and allodynia, diabetic 30 neuropathy pain, causalgia, sympathetically maintained pain, deafferentation syndromes, asthma, epithelial tissue damage or dysfunction, herpes simplex, disturbances of visceral motility at respiratory, genitourinary, gastrointestinal or

- vascular regions, wounds, burns, allergic skin reactions, pruritus, vitiligo, general gastrointestinal disorders, gastric ulceration, duodenal ulcers, diarrhea, gastric lesions induced by necrotising agents, hair growth, vasomotor or allergic rhinitis, bronchial disorders or bladder disorders, the compounds of the present invention
- 5 may be administered orally, parentally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional pharmaceutically acceptable carriers, adjuvants, and vehicles. The term parenteral as used herein includes, subcutaneous, intravenous, intramuscular, intrasternal, infusion techniques or intraperitoneally.
- 10 Treatment of diseases and disorders herein is intended to also include the prophylactic administration of a compound of the invention, a pharmaceutical salt thereof, or a pharmaceutical composition of either to a subject (*i.e.*, an animal, preferably a mammal, most preferably a human) believed to be in need of preventative treatment, such as, for example, pain, inflammation and the like.
- 15 The dosage regimen for treating TRP-M8-receptor-mediated diseases, cancer, and/or hyperglycemia with the compounds of this invention and/or compositions of this invention is based on a variety of factors, including the type of disease, the age, weight, sex, medical condition of the patient, the severity of the condition, the route of administration, and the particular compound employed.
- 20 Thus, the dosage regimen may vary widely, but can be determined routinely using standard methods. Dosage levels of the order from about 0.01 mg to 30 mg per kilogram of body weight per day, preferably from about 0.1 mg to 10 mg/kg, more preferably from about 0.25 mg to 1 mg/kg are useful for all methods of use disclosed herein.
- 25 The pharmaceutically active compounds of this invention can be processed in accordance with conventional methods of pharmacy to produce medicinal agents for administration to patients, including humans and other mammals.
- For oral administration, the pharmaceutical composition may be in the form of, for example, a capsule, a tablet, a suspension, or liquid. The
- 30 pharmaceutical composition is preferably made in the form of a dosage unit containing a given amount of the active ingredient. For example, these may contain an amount of active ingredient from about 1 to 2000 mg, preferably from

about 1 to 500 mg, more preferably from about 5 to 150 mg. A suitable daily dose for a human or other mammal may vary widely depending on the condition of the patient and other factors, but, once again, can be determined using routine methods.

5 The active ingredient may also be administered by injection as a composition with suitable carriers including saline, dextrose, or water. The daily parenteral dosage regimen will be from about 0.1 to about 30 mg/kg of total body weight, preferably from about 0.1 to about 10 mg/kg, and more preferably from about 0.25 mg to 1 mg/kg.

10 Injectable preparations, such as sterile injectable aqueous or oleaginous suspensions, may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed, including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

15 Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable non-irritating excipient such as cocoa butter and polyethylene glycols that are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

20 A suitable topical dose of active ingredient of a compound of the invention is 0.1 mg to 150 mg administered one to four, preferably one or two times daily. For topical administration, the active ingredient may comprise from 0.001% to 10% w/w, e.g., from 1% to 2% by weight of the formulation, although it may comprise as much as 10% w/w, but preferably not more than 5% w/w, and more preferably from 0.1% to 1% of the formulation.

25 Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin (e.g., liniments,

lotions, ointments, creams, or pastes) and drops suitable for administration to the eye, ear, or nose.

For administration, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate for the indicated route of administration. The compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanoic acids, stearic acid, talc, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, acacia, gelatin, sodium alginate, polyvinyl-pyrrolidine, and/or polyvinyl alcohol, and tableted or encapsulated for conventional administration. Alternatively, the compounds of this invention may be dissolved in saline, water, polyethylene glycol, propylene glycol, ethanol, corn oil, peanut oil, cottonseed oil, sesame oil, tragacanth gum, and/or various buffers. Other adjuvants and modes of administration are well known in the pharmaceutical art. The carrier or diluent may include time delay material, such as glyceryl monostearate or glyceryl distearate alone or with a wax, or other materials well known in the art.

The pharmaceutical compositions may be made up in a solid form (including granules, powders or suppositories) or in a liquid form (*e.g.*, solutions, suspensions, or emulsions). The pharmaceutical compositions may be subjected to conventional pharmaceutical operations such as sterilization and/or may contain conventional adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers, buffers etc.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, *e.g.*, lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert

diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting, sweetening, flavoring, and perfuming agents.

Compounds of the present invention can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or non-racemic mixtures thereof. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, *e.g.*, by formation of diastereoisomeric salts, by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric, and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts. A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting compounds of the invention with an optically pure acid in an activated form or an optically pure isocyanate. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active compounds of the invention can likewise be obtained by using active starting materials. These isomers may be in the form of a free acid, a free base, an ester or a salt.

Likewise, the compounds of this invention may exist as isomers, that is compounds of the same molecular formula but in which the atoms, relative to one another, are arranged differently. In particular, the alkylene substituents of the compounds of this invention, are normally and preferably arranged and inserted into the molecules as indicated in the definitions for each of these groups, being read from left to right. However, in certain cases, one skilled in the art will appreciate that it is possible to prepare compounds of this invention in which these substituents are reversed in orientation relative to the other atoms in the molecule. That is, the substituent to be inserted may be the same as that noted above except that it is inserted into the molecule in the reverse orientation. One skilled in the art will

appreciate that these isomeric forms of the compounds of this invention are to be construed as encompassed within the scope of the present invention.

The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. The salts include, but are not limited to,

5 the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate,

10 methansulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, palmoate, pectinate, persulfate, 2-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, mesylate, and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates

15 like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids that may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulfuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid. Other examples include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic bases.

25 Also encompassed in the scope of the present invention are pharmaceutically acceptable esters of a carboxylic acid or hydroxyl containing group, including a metabolically labile ester or a prodrug form of a compound of this invention. A metabolically labile ester is one which may produce, for example, an increase in blood levels and prolong the efficacy of the corresponding

30 non-esterified form of the compound. A prodrug form is one which is not in an active form of the molecule as administered but which becomes therapeutically active after some in vivo activity or biotransformation, such as metabolism, for

example, enzymatic or hydrolytic cleavage. For a general discussion of prodrugs involving esters see Svensson and Tunek Drug Metabolism Reviews 165 (1988) and Bundgaard Design of Prodrugs, Elsevier (1985). Examples of a masked carboxylate anion include a variety of esters, such as alkyl (for example, methyl, 5 ethyl), cycloalkyl (for example, cyclohexyl), aralkyl (for example, benzyl, p-methoxybenzyl), and alkylcarbonyloxyalkyl (for example, pivaloyloxymethyl). Amines have been masked as arylcarbonyloxymethyl substituted derivatives which are cleaved by esterases *in vivo* releasing the free drug and formaldehyde (Bungaard J. Med. Chem. 2503 (1989)). Also, drugs containing an acidic NH 10 group, such as imidazole, imide, indole and the like, have been masked with N-acyloxymethyl groups (Bundgaard Design of Prodrugs, Elsevier (1985)). Hydroxy groups have been masked as esters and ethers. EP 039,051 (Sloan and Little, 4/11/81) discloses Mannich-base hydroxamic acid prodrugs, their preparation and use. Esters of a compound of this invention, may include, for example, the 15 methyl, ethyl, propyl, and butyl esters, as well as other suitable esters formed between an acidic moiety and a hydroxyl containing moiety. Metabolically labile esters, may include, for example, methoxymethyl, ethoxymethyl, iso-propoxymethyl, α -methoxyethyl, groups such as α -((C₁-C₄)alkyloxy)ethyl, for example, methoxyethyl, ethoxyethyl, propoxyethyl, iso-propoxyethyl, etc.; 2-oxo- 20 1,3-dioxolen-4-ylmethyl groups, such as 5-methyl-2-oxo-1,3,dioxolen-4-ylmethyl, etc.; C₁-C₃ alkylthiomethyl groups, for example, methylthiomethyl, ethylthiomethyl, isopropylthiomethyl, etc.; acyloxymethyl groups, for example, pivaloyloxymethyl, α -acetoxymethyl, etc.; ethoxycarbonyl-1-methyl; or α -acyloxy- α -substituted methyl groups, for example α -acetoxymethyl.

25 Further, the compounds of the invention may exist as crystalline solids which can be crystallized from common solvents such as ethanol, N,N-dimethyl-formamide, water, or the like. Thus, crystalline forms of the compounds of the invention may exist as polymorphs, solvates and/or hydrates of the parent compounds or their pharmaceutically acceptable salts. All of such forms likewise 30 are to be construed as falling within the scope of the invention.

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or


more compounds of the invention or other agents. When administered as a combination, the therapeutic agents can be formulated as separate compositions that are given at the same time or different times, or the therapeutic agents can be given as a single composition.

5 The foregoing is merely illustrative of the invention and is not intended to limit the invention to the disclosed compounds. Variations and changes which are obvious to one skilled in the art are intended to be within the scope and nature of the invention which are defined in the appended claims.

10 From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

We Claim:

1. A compound having the structure:



5 or any pharmaceutically-acceptable salt thereof, wherein:

J is $-N(R^a)(CR^cR^c)_n-$, $-O(CR^cR^c)_n-$, $-S(CR^cR^c)_n-$ or $-(CR^cR^c)_n-$;

m is 0, 1 or 2;

n is independently in each instance 0, 1, 2 or 3;

o is 0 or 1

10 X is O, NR^a or S;

R¹ is selected from H, halo, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$,

$-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$,

$-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$,

$-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$,

15 $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$ and $-NR^aC_{2-6}alkOR^a$; or
R¹ is C₁₋₆alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C₁₋₆alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk,

C₁₋₄haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$,

20 $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$,
 $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$,
 $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$,
 $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$ and $-NR^aC_{2-6}alkOR^a$;

R² is selected from H, halo, cyano, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$,

25 $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}alkNR^aR^a$,
 $-OC_{2-6}alkOR^a$, $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$,
 $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$,
 $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}alkNR^aR^a$ and $-NR^aC_{2-6}alkOR^a$; or

R^2 is C_{1-6} alk or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C_{1-6} alk and ring are substituted by 0, 1, 2 or 3 substituents selected from C_{1-8} alk,

5 C_{1-4} haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$,


$-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a ,

$-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$,

$-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$,

$-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ;

R^3 is



10

; or

R^3 is a saturated, partially-saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 1, 2, 3 or 4 atoms selected from N, O and S, substituted with 0, 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk,

15 $-N(C_{1-6}$ alk) C_{1-6} alk, F, Cl, Br, CN, OH and NH₂; or R^3 is benzyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F, Cl, Br, CN, OH and NH₂; or R^3 is napthyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F,

20 Cl, Br, CN and NH₂; or R^3 is benzyl substituted with 1, 2 or 3 groups selected from C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F, Cl, Br, CN, OH and NH₂;

25 R^4 is selected from H, C_{1-6} alk, C_{1-3} haloalk, $-OC_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, $-NHC_{1-6}$ alk, $-NC(=O)C_{1-6}$ alk, $-N(C_{1-6}$ alk) C_{1-6} alk, F, Cl, Br, CN, OH and NH₂;

R^5 is C_{1-11} alkyl, C_{6-11} cycloalkyl or a saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the C_{1-6} alkyl and ring are substituted by 0, 1 or 2 oxo groups and the C_{1-11} alkyl,

C₆₋₁₁cycloalkyl and ring are additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a,

5 -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or if J is -N(R^a)(CR^cR^c)_n-, -O(CR^cR^c)_n- or -S(CR^cR^c)_n-, then R⁵ may also be C₃₋₅cycloalkyl substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=O)OR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NRR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or if J is -N(R^a)(CR^cR^c)_n-, then R⁵ may also be H;

10 15 R⁶ is independently in each instance, selected from F, Cl, Br, I, C₂₋₆alk, cyano, -OR^a, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁶ is C₁₋₆alk substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a

20 25 and -NR^aC₂₋₆alkOR^a; or R⁶ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

30

R⁷ is selected from C₁₋₆alk, cyano, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁷ is C₁₋₆alk substituted by 1, 2 or 3 substituents selected from C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a; or R⁷ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the ring is substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a,

15 -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a, -N(R^a)S(=O)R^b, -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)₂NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a;

R^a is independently, at each instance, H or R^b;

R^b is independently, at each instance, phenyl, benzyl or C₁₋₆alk, the phenyl,

20 benzyl and C₁₋₆alk being substituted by 0, 1, 2 or 3 substituents selected from halo, C₁₋₄alk, C₁₋₃haloalk, -OC₁₋₄alk, -NH₂, -NHC₁₋₄alk, and -N(C₁₋₄alk)C₁₋₄alk;

R^c is independently, at each instance, H, halo, C₁₋₄alk, C₁₋₄haloalk,

-OC₁₋₄alk, -OC₁₋₄haloalk, -NH₂, -NHC₁₋₄alk or -N(C₁₋₄alk)C₁₋₄alk; and

R^f is independently in each instance H or F.


25

2. A compound according to Claim 1, wherein J is -N(R^a)-, -O-, -S- or -CH₂-.

3. A compound according to Claim 2, wherein o is 1.

30

4. A compound according to Claim 3, wherein
 R^3 is



and

- R^6 is C_{1-6} alk substituted by 1, 2 or 3 substituents selected from C_{1-4} haloalk,
5 halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$,
 $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$,
 $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$,
 $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$,
 $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a ; or R^6 is a saturated,
10 partially saturated or unsaturated 5-, 6- or 7-membered ring containing 0, 1, 2, 3 or
4 atoms selected from N, O and S, wherein the C_{1-6} alk is and C_{1-6} alk and ring are
substituted by 0, 1, 2 or 3 substituents selected from C_{1-8} alk, C_{1-4} haloalk, halo,
cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$, $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$,
 $-OC(=O)R^b$, $-OC(=O)NR^aR^a$, $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$,
15 $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$, $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$,
 $-N(R^a)C(=O)NR^aR^a$, $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$,
 $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a and $-NR^aC_{2-6}$ alk OR^a .

5. A compound according to Claim 2, wherein R^5 is C_{1-11} alkyl or
20 C_{6-11} cycloalkyl both of which are substituted by 0, 1, 2 or 3 substituents selected
from C_{1-8} alk, C_{1-4} haloalk, halo, cyano, nitro, $-C(=O)R^b$, $-C(=O)OR^b$,
 $-C(=O)NR^aR^a$, $-C(=NR^a)NR^aR^a$, $-OR^a$, $-OC(=O)R^b$, $-OC(=O)NR^aR^a$,
 $-OC_{2-6}$ alk NR^aR^a , $-OC_{2-6}$ alk OR^a , $-SR^a$, $-S(=O)R^b$, $-S(=O)_2R^b$, $-S(=O)_2NR^aR^a$,
 $-NR^aR^a$, $-N(R^a)C(=O)R^b$, $-N(R^a)C(=O)OR^b$, $-N(R^a)C(=O)NR^aR^a$,
25 $-N(R^a)C(=NR^a)NR^aR^a$, $-N(R^a)S(=O)_2R^b$, $-N(R^a)S(=O)_2NR^aR^a$, $-NR^aC_{2-6}$ alk NR^aR^a
and $-NR^aC_{2-6}$ alk OR^a .

6. A compound according to Claim 2, wherein R⁵ is a saturated, partially saturated or unsaturated 5-, 6- or 7-membered monocyclic or 8-, 9-, 10- or 11-membered bicyclic ring containing 0, 1, 2, 3 or 4 atoms selected from N, O and S, wherein the ring is substituted by 0, 1 or 2 oxo groups and the ring is
5 additionally substituted by 0, 1, 2 or 3 substituents selected from C₁₋₈alk, C₁₋₄haloalk, halo, cyano, nitro, -C(=O)R^b, -C(=O)OR^b, -C(=O)NR^aR^a, -C(=NR^a)NR^aR^a, -OR^a, -OC(=O)R^b, -OC(=O)NR^aR^a, -OC₂₋₆alkNR^aR^a, -OC₂₋₆alkOR^a, -SR^a, -S(=O)R^b, -S(=O)₂R^b, -S(=O)₂NR^aR^a, -NR^aR^a, -N(R^a)C(=O)R^b, -N(R^a)C(=O)OR^b, -N(R^a)C(=O)NR^aR^a, -N(R^a)C(=NR^a)NR^aR^a,
10 -N(R^a)S(=O)₂R^b, -N(R^a)S(=O)NR^aR^a, -NR^aC₂₋₆alkNR^aR^a and -NR^aC₂₋₆alkOR^a.

7. A method of treating acute, inflammatory and neuropathic pain, dental pain, general headache, migraine, cluster headache, mixed-vascular and non-vascular syndromes, tension headache, general inflammation, arthritis, rheumatic diseases, osteoarthritis, inflammatory bowel disorders, depression, anxiety, inflammatory eye disorders, inflammatory or unstable bladder disorders, psoriasis, skin complaints with inflammatory components, chronic inflammatory conditions, inflammatory pain and associated hyperalgesia and allodynia, neuropathic pain and associated hyperalgesia and allodynia, diabetic neuropathy
15 pain, causalgia, sympathetically maintained pain, deafferentation syndromes, asthma, epithelial tissue damage or dysfunction, herpes simplex, disturbances of visceral motility at respiratory, genitourinary, gastrointestinal or vascular regions, wounds, burns, allergic skin reactions, pruritus, vitiligo, general gastrointestinal disorders, gastric ulceration, duodenal ulcers, diarrhea, gastric lesions induced by
20 necrotising agents, hair growth, vasomotor or allergic rhinitis, bronchial disorders or bladder disorders, comprising the step of administering a compound according to Claim 1.

8. A pharmaceutical composition comprising a compound according
30 to Claim 1 and a pharmaceutically-acceptable diluent or carrier.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2008/013375

A. CLASSIFICATION OF SUBJECT MATTER

INV.	C07D217/04	C07D217/06	C07D217/22	C07D401/04	C07D401/06
	C07D401/10	C07D401/12	C07D405/06	C07D405/12	C07D413/06
	C07D417/06	A61K31/472	A61P29/00		

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 801 067 A (YAMANOUCHI PHARMA CO LTD [JP]) 15 October 1997 (1997-10-15) Claims 1 and 11 and tables -----	1-8
X	DATABASE CA [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ZHOU, ZHISHAN ET AL: "Syntheses of 1,2-disubstituted tetrahydroisoquinoline derivatives and their antiarrhythmic effects" XP002519323 retrieved from STN Database accession no. 1988:94357 examples -/-	1-6,8



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

16 March 2009

Date of mailing of the international search report

25/03/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Sahagún Krause, H

INTERNATIONAL SEARCH REPORT

International application No PCT/US2008/013375

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	& ZHOU, ZHISHAN ET AL: "Syntheses of 1,2-disubstituted tetrahydroisoquinoline derivatives and their antiarrhythmic effects" SHANGHAI YIKE DAXUE XUEBAO , 14(1), 15-20 CODEN: SYDXEE; ISSN: 0257-8131, 1987, -----	
X	WO 2007/098608 A (LI CHAO-JUN [CA]; MCLEOD PATRICIA [CA]; LI ZHIPING [CA]; FENG JIANQING) 7 September 2007 (2007-09-07) compounds 4a, 5a, 4f and 5f, page 25 two first compounds in the last line in page 4 -----	1-6
X	WO 01/29030 A (CENTRE NAT RECH SCIENT [FR]; INNOVATIONSAGENTUR GMBH BU TEC [AT]; UNIV) 26 April 2001 (2001-04-26) pages 20-25, claims 1 and 13 -----	1-8
X	US 6 177 443 B1 (MADSEN PETER [DK] ET AL) 23 January 2001 (2001-01-23) claim 1, examples 80-82, 85, 86, 129-131 -----	1-6,8
X	VENKOV A P ET AL: "New modification of the intramolecular [alpha]-amidoalkylation for the synthesis of 2-acyl-1,2,3,4-tetrahydroisoquinolines" SYNTHESIS 1989 DE, no. 1, 1989, pages 59-61, XP002519321 ISSN: 0039-7881 compound 5f, page 59 -----	1-6
X	VENKOV A P ET AL: "Synthesis of 1-phenyl-2-acyl-tetrahydroisoquinolines by intermolecular [alpha]-amidoalkylation reaction" SYNTHETIC COMMUNICATIONS 1992 US, vol. 22, no. 1, 1992, pages 125-134, XP002519322 ISSN: 0039-7911 compounds 5, 6a, 7a, 8a, 9, 10, 11a, 15-20 -----	1-6
X	NAITO RYO ET AL: "Synthesis and Antimuscarinic Properties of Quinuclidin-3-yl 1,2,3,4-Tetrahydroisoquinoline-2-carboxylate Derivatives as Novel Muscarinic Receptor Antagonists" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON.; US, vol. 48, 20 October 2005 (2005-10-20), pages 6597-6606, XP002435582 ISSN: 0022-2623 compounds 32, 33, 12a-b, 13a-b, 15a-b and introduction -----	1-8

-/-

INTERNATIONAL SEARCH REPORT

International application No PCT/US2008/013375

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	DATABASE CHEMCATS CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 13 June 2008 (2008-06-13), XP002519324 Order numbers: kam-021285 & "Aurora Screening Library" 13 June 2008 (2008-06-13), AURORA FINE CHEMICALS LLC , SAN DIEGO, CA, 92126 USA -----	1-6
P,X	US 2005/054651 A1 (NATARAJAN SATEESH K [US] ET AL) 10 March 2005 (2005-03-10) isoquinolines in pages 64, 66, 68, claim 2, paragraph [0002] -----	1-6
A	US 2005/070570 A1 (GARCIA GABRIEL [DE] ET AL) 31 March 2005 (2005-03-31) the whole document -----	1-8
A		1-8

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2008/013375

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
EP 0801067	A	15-10-1997		AT 233761 T AU 695616 B2 AU 4355396 A CA 2208839 A1 CN 1171109 A DE 69529844 D1 DE 69529844 T2 DE 122004000048 I1 DK 801067 T3 ES 2193208 T3 FI 972775 A HU 77006 A2 WO 9620194 A1 LU 91133 A9 NL 300141 I1 NO 973027 A NO 2005012 I1 NO 2005016 I1 NZ 298144 A PL 321019 A1 PT 801067 T RU 2143432 C1 US 6174896 B1 US 6017927 A	15-03-2003 20-08-1998 19-07-1996 04-07-1996 21-01-1998 10-04-2003 04-03-2004 21-04-2005 30-06-2003 01-11-2003 22-08-1997 02-03-1998 04-07-1996 29-03-2005 01-04-2004 28-08-1997 06-06-2005 29-08-2005 27-04-1998 24-11-1997 31-07-2003 27-12-1999 16-01-2001 25-01-2000
WO 2007098608	A	07-09-2007		CA 2643493 A1	07-09-2007
WO 0129030	A	26-04-2001		AT 245155 T AU 7932000 A CA 2387972 A1 DE 60003941 D1 DE 60003941 T2 EP 1224182 A1 FR 2800072 A1 JP 2003512373 T US 6649626 B1	15-08-2003 30-04-2001 26-04-2001 21-08-2003 06-05-2004 24-07-2002 27-04-2001 02-04-2003 18-11-2003
US 6177443	B1	23-01-2001		NONE	
US 2005054651	A1	10-03-2005		NONE	
US 2005070570	A1	31-03-2005		NONE	