



US 20110250254A1

(19) **United States**(12) **Patent Application Publication****Scheie et al.**(10) **Pub. No.: US 2011/0250254 A1**(43) **Pub. Date: Oct. 13, 2011**(54) **ANTIMICROBIAL COMPOSITIONS AND USES**(76) Inventors: **Anne Aamdal Scheie**, Amdalsverk (NO); **Tore Benneche**, Oslo (NO); **Jessica Lonn-Stensrud**, Lysaker (NO); **Jan Skramstad**, Rykkin (NO)(21) Appl. No.: **13/122,994**(22) PCT Filed: **Oct. 9, 2009**(86) PCT No.: **PCT/EP2009/063210**§ 371 (c)(1),  
(2), (4) Date:**Jun. 17, 2011**(30) **Foreign Application Priority Data**

Oct. 9, 2008 (GB) ..... 0818547.2


**Publication Classification**(51) **Int. Cl.**

**A61K 31/381** (2006.01)  
**A61Q 11/00** (2006.01)  
**A61F 2/00** (2006.01)  
**A01N 25/00** (2006.01)  
**A61F 13/00** (2006.01)  
**A01N 25/34** (2006.01)


**C07D 333/32** (2006.01)  
**A01N 43/10** (2006.01)  
**C07D 409/06** (2006.01)  
**C07D 409/14** (2006.01)  
**A61Q 15/00** (2006.01)  
**A61P 31/00** (2006.01)  
**A01P 1/00** (2006.01)  
**A01P 3/00** (2006.01)  
**C08F 228/06** (2006.01)  
**A61K 8/49** (2006.01)

(52) **U.S. Cl. .... 424/405; 424/65; 424/49; 424/408; 549/62; 514/445; 549/66; 549/64; 549/59; 514/444; 526/256**(57) **ABSTRACT**

An agent comprising the compound according to general formula (I) wherein  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are each independently H or a substituent, and wherein at least one of  $R_1$ ,  $R_3$  and  $R_4$  is halogen, cyano, cyanate, thiocyanate or  $C_1$ - $C_6$  haloalkyl.



**AI-1 Quorum-sensing inhibition by furanone or thiophenone –  
Assessed by inhibition of bioluminescence**






Figure 1






Figure 2


**A**



**B**



**Figure 3**

**Figure 4**

AI-1 Quorum-sensing inhibition by furanone or thiophenone –  
Assessed by inhibition of bioluminescence




Figure 5

# AI-2 Quorum-sensing inhibition by furanone or thiophenone – Assessed by inhibition of bioluminescence




Figure 6

## ANTIMICROBIAL COMPOSITIONS AND USES

[0001] The present invention relates to chemical compounds and polymers incorporating such chemical compounds for use as antimicrobial agents, more particularly for blocking or interfering with quorum-sensing microbial communication and/or preventing or inhibiting biofilm formation.

### BACKGROUND OF THE INVENTION

[0002] Microbes, and in particular bacteria, are known to form biofilms under conditions where there is a combination of bacteria, moisture, nutrients and a suitable surface. Biofilm formation and biofouling create problems and economic losses in domestic, industrial and health fields. Various industrial processes and installations may be affected, such as submarine installations and shipping, various engineering industries, oil processing and manufacturing, the food and beverage industry, the pharmaceutical industry, water systems, cooling towers, heat exchangers, chain lubrication systems and the like. Biofilms also cause problems in relation to medical devices and implants and cause various human and animal infections. It is generally necessary to use a harsh treatment to remove and kill an established biofilm. The reason for this is that bacteria situated in a biofilm structure are protected from established antibacterial treatments. Biofilm formation is believed to involve activation and/or down regulation of a number of genes in response to communication signalling molecules. Gene expression is different in biofilms as compared to free-flowing, planktonic, bacteria.


[0003] One approach to the disruption or inhibition of bacterial biofilms is described in U.S. Pat. No. 6,726,898. Compositions and methods of treating periodontal disease are described which employ furanones or furanone derivatives which inhibit or disrupt the glycocalyx matrix of the bacterial biofilm.

[0004] Outside the field of antimicrobial compositions Halvorsen et al describe in Synthetic Communications, 2007, 37(7), 1167-1177 oligothiophene compounds for use in materials with non-linear optical properties. Synthesis of various thiophenones is also described in Jakobsen et al in Tetrahedron, 1963, 19, 1867-1882. Hornfeldt and Gronowitz describe in Arkiv foer Kemi, 1963, 21(22), 239-57 the synthesis of further thiophenes without any indication of their utility.

[0005] A need exists to find new agents with antimicrobial properties, in particular for use in the prevention or disruption of biofilm formation, which are more effective than those described in the prior art.


### SUMMARY OF THE INVENTION

[0006] Accordingly, in a first aspect, the present invention provides an agent comprising the compound according to general formula (I):



(I)


[0007] wherein  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are each independently H or a substituent, and wherein at least one of  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  is halogen, cyano, cyanate, thiocyanate or  $C_1$ - $C_6$  haloalkyl. In this aspect of the invention, it is preferred that the compound is other than



[0008] For example, when  $R_1$  or  $R_2$  is Br, it is preferred that one of  $R_3$  and  $R_4$  is not Ph when the other is H.

[0009] It has surprisingly been found that compounds according to the present invention act more effectively in inhibition of biofilm formation, as compared with those of the prior art.


[0010] In a further aspect, the present invention provides an agent for use in medicine, which agent comprises the compound of general formula (II):



(II)

[0011] wherein X is O, S, NH or NR', in which R' is an optionally substituted  $C_1$ - $C_6$  alkyl group;  $R_1$ ,  $R_2$ ,  $R_3$ , and  $R_4$  are each independently H or a substituent, and wherein the compound is capable of blocking or interfering with quorum-sensing microbial communication.

[0012] In a further aspect the present invention provides an agent for use in medicine, which agent comprises the compound of general formula (II):



(II)

[0013] wherein X is O, S, NH or NR', in which R' is an optionally substituted  $C_1$ - $C_6$  alkyl group;  $R_1$ ,  $R_2$ ,  $R_3$ , and  $R_4$  are each independently H or a substituent, and wherein the compound is capable of preventing or inhibiting biofilm formation.

[0014] By appropriate selection of substituent groups on the compound of general formula (II), compounds are provided which are capable of blocking or interfering with quorum-sensing microbial communication or preventing or inhibiting biofilm formation. Each of these properties is described in further detail herein, together with tests to demonstrate whether or not the compounds exhibit these proper-




ties. Quorum-sensing microbial communication is thought to be mediated by a signalling pathway that is activated as a response to cell density. Such signalling is found in both gram-positive and gram-negative micro-organisms. Perception by bacteria of a quorum-sensing signal occurs at a concentration threshold and it is thought that the bacterial population then responds to the signal. The signal molecules of quorum-sensing systems are thought to be highly specific. It is thought that quorum-sensing systems play a part in biofilm formation. Accordingly, by using the agents of the present invention, quorum-sensing signalling may be blocked or interfered with, thereby interfering with the behaviour of the bacterial population.

**[0015]** A further advantage of interfering with quorum-sensing signalling is that preferred compounds according to the invention which do this do not exert selective pressure on the bacterial population. The bacteria are not killed; instead, their phenotypes are regulated. Accordingly, antimicrobial resistance development is unlikely to result.

**[0016]** In one arrangement, the compound of general formula (II) has a substituent X which is O. Such compounds are relatively straightforward to synthesize and show inhibitory activity towards biofilm formation and towards quorum-sensing.

**[0017]** Each substituent of the compound of general formula (II) may be independently selected from halogen, cyano, cyanate, thiocyanate, alkyl, alkoxy, haloalkyl, alkyl ester, alkylsilyl, alkenyl, alkynyl, aryl, or arylalkyl, which may be substituted or unsubstituted, optionally interrupted by one or more heteroatoms, straight chain or branched chain. Generally, the substituents may have up to six carbon atoms so that the alkyl group is typically a  $C_1$  to  $C_6$  alkyl substituent, the alkoxy substituent is typically a  $C_1$  to  $C_6$  alkoxy substituent and so on. Generally speaking, larger substituent groups may be more likely not to have the activity of blocking or interfering with quorum-sensing microbial communication or preventing or inhibiting biofilm formation. The aryl substituent is preferably a monocyclic aryl, such as phenyl.

**[0018]** It is preferred that each substituent is independently selected from halogen, haloalkyl, alkoxy, alkyl ester, phenyl or



wherein  $R_5$ ,  $R_6$  and  $R_7$  are each independently H, Br, Cl, OMe, or CHO.

**[0019]** Compounds of high activity have been found where  $R_4$  is a substituent, rather than H. It is further preferred that at least one of  $R_1$ ,  $R_3$  and  $R_4$  is halogen, cyano, cyanate, thiocyanate or  $C_1$  to  $C_6$  haloalkyl and, more preferably, at least one of  $R_1$  and  $R_4$  is halogen or  $C_1$  to  $C_6$  haloalkyl.  $R_1$  may also be  $-\text{CH}_2-\text{O}-\text{CO}-(\text{CH}_2)_2-\text{COOH}$  or thienyl.

**[0020]** In one arrangement it is preferred that  $R_1$  and  $R_2$  are each independently H or halogen.



**[0021]** In one arrangement it is preferred that  $R_3$  is H, halogen,  $C_1$  to  $C_6$  haloalkyl or phenyl. It is also preferred that  $R_4$  is halogen,  $C_1$  to  $C_6$  haloalkyl or phenyl.

**[0022]** Compounds of high activity have been found where the halogen is Br or I, particularly Br. It is particularly preferred that  $R_1$  is Br. It is also particularly preferred that  $R_4$  is Br. It is particularly preferred that  $R_2$  is H. It is also particularly preferred that  $R_3$  is H. In a further arrangement it is


preferred that  $R_1$  and  $R_4$  are each Br and  $R_2$  and  $R_3$  are each H. Alternatively,  $R_3$  and  $R_4$  are each Br and  $R_1$  and  $R_2$  are each H. In a further preferred arrangement,  $R_1$ ,  $R_3$  and  $R_4$  are each Br and  $R_2$  is H. In a further preferred arrangement  $R_1$  is  $-\text{CH}_2-\text{O}-\text{CO}-(\text{CH}_2)_2-\text{COOH}$  and  $R_4$  is Br. In a further preferred arrangement  $R_1$  is thiophenyl and  $R_4$  is Br. In a further preferred arrangement  $R_3$  is methyl and  $R_4$  is Br. In a further preferred arrangement  $R_3$  is H and  $R_4$  is thiocyanate.


**[0023]** In a further aspect the present invention provides a polymer which comprises a compound as defined herein. An advantage of using a polymeric composition comprising the compound is that surfaces may be treated with a polymer or polymer-forming composition so as to inhibit or prevent biofilm formation thereon. The compound may be incorporated into the polymer as a side chain or in the main chain of the polymer, for example copolymerised with another comonomer to form a copolymer. In one arrangement, the polymer may therefore comprise one or more side chain functional groups comprising the compound wherein the backbone of the polymer is typically a known polymeric backbone such as a polyacrylate, polymethacrylate, polycrotonate, polyvinyl alcohol, polyvinyl acetate, polystyrene, acrylonitrile or siloxane.

**[0024]** In one arrangement, the polymer may be obtainable by polymerising the compound according to general formula (III) with the compound according to general formula (IV).



**[0025]** In another arrangement, the compound may be obtainable by polymerising the compound according to general formula (V) with the compound according to general formula (VI):





**[0026]** In a further aspect, the present invention provides a process for manufacturing a polymer comprising the compound as defined herein.

**[0027]** Uses of the Agent or Polymer

**[0028]** The agent or polymer of the invention has a wide variety of uses in different fields.

**[0029]** The agent or polymer may be used in medicine in the form described herein or as a pharmaceutically-acceptable salt, ester or prodrug thereof. Pharmaceutically-acceptable salts and esters are well known to those skilled in the pharmaceutical field and they include suitable acid addition salts, base addition salts and esters which are non-toxic to the recipient. A prodrug form may comprise the agent or polymer as a derivative which becomes active only when metabolised by the recipient. Pharmaceutical compositions may be formulated comprising the agent or polymer optionally incorporating a pharmaceutically-acceptable excipient, diluent or carrier, the exact nature of which may be selected according to the intended route of administration. Other ingredients suitable for pharmaceutical use may also be incorporated, as is well known in the pharmaceutical field, such as solvents, buffers, surfactants, preservative agents, and so on.

**[0030]** The agent or polymer according to the invention may be used as an antimicrobial, for example in the prevention or treatment of microbial infection. Microbial infections include bacterial or fungal infections and, in particular, those which involve quorum-sensing microbial communication or biofilm formation. The agent or polymer may interfere with quorum-sensing microbial communication so as to treat or prevent a condition mediated by microbes which are regulated by quorum-sensing communication.

**[0031]** The agent or polymer may be used in conjunction with one or more further antimicrobial agents such as antibiotics or antifungals. In this way, the invention provides a composition comprising an agent or polymer as described herein and one or more further antimicrobial agents as a combined preparation for simultaneous, separate or sequential use in the prevention or treatment of microbial infection. The two components of the combined preparation may be administered separately from one another either at the same time or at separate times. Sequential administration may involve two or more sequential treatments. Where a simultaneous treatment is required, the composition may comprise the components either mixed together or stored separately. The combined preparation may be provided in kit form for convenient use.

**[0032]** The agent, polymer or composition may be used in the treatment of oral conditions, topical infections, respiratory infections, eye infections, ear infections or localised organ infections.

**[0033]** Each of these conditions typically involves biofilm formation and/or microbial quorum-sensing communication. Oral conditions include periodontitis, gingivitis and dental

caries. At least some of these conditions need not be addressed using a pharmaceutical composition and may instead be addressed using a personal care product. For example, oral conditions may be treated or prevented using a dentifrice or mouthwash. Topical infections may be treated or prevented using shampoo, soap or deodorant or cosmetic composition. Eye infections may be treated or prevented using a contact lens solution.


**[0034]** The invention further provides a personal care product comprising an agent or polymer as defined herein which is a personal hygiene article, shampoo, soap, deodorant, dentifrice, mouthwash, contact lens solution or cosmetic composition. Such personal care products may be made in a conventional way by incorporating into conventional ingredients the agent or polymer as defined herein.

**[0035]** In a further aspect, the present invention provides an antimicrobial surface cleanser which comprises an agent or polymer as defined herein. The antimicrobial surface cleanser may be formulated for use on an inanimate surface or on the surface of the skin of a human or animal. The antimicrobial surface cleanser may be a disinfectant or a cleaning composition.

**[0036]** In the case of the surface of the skin of an animal or human, it is frequently necessary to ensure that the skin is completely free of microorganisms so that their carriage to other humans or animals is prevented or inhibited.

**[0037]** The antimicrobial surface cleanser may be applied to inanimate surfaces of a very wide variety. Such surfaces include worktops, floors, food preparation tools and equipment surfaces and medical equipment surfaces.

**[0038]** In a further aspect, there is provided a coating composition comprising the agent or polymer as defined herein. In one arrangement, the coating composition is capable of binding covalently to a surface. The coating composition may be in the form of any conventional coating composition such as a paint. In one arrangement, the coating composition comprises a polymer or forms a polymer from suitable reactants on the surface to be treated. In one arrangement, the coating composition comprises the agent covalently linked to the group  $\text{Si}(\text{OR}_5)_3$ , wherein each  $\text{R}_5$  is independently substituted or unsubstituted  $\text{C}_1$ - $\text{C}_6$  hydrocarbyl. The agent may be covalently linked to the group  $\text{Si}(\text{OR}_5)_3$  by a linker, which linker may comprise a substituted or unsubstituted alkyl, alkenyl, alkynyl, alkylaryl, arylalkyl or aryl linker optionally interrupted by one or more heteroatoms such as O, N and S. In one arrangement the linker comprises  $-\text{CH}_2-\text{O}-\text{CO}-\text{NH}-(\text{CH}_2)_3-$ . The linker preferably attaches to the agent at the  $\text{R}_1$  position so that the  $\text{R}_1$  substituent is the moiety-linker- $\text{Si}(\text{OR}_5)_3$ .  $\text{R}_5$  is typically ethyl. In one arrangement, the coating composition comprises a compound according to general formula (VII)



[0039] The coating composition may be used as an antibiofouling composition in various applications. Biofouling may occur on marine vessels, submarine installations, pipelines, waterpipes, industrial machines or installations, water systems, cooling towers, heat exchangers, chain lubrication systems, oil or gas platforms, fish farming installations or surfaces, machines, tools or devices used in food production. Biofilm formation is particularly undesirable in these situations. Biocorrosion of the surface may arise over time. By applying a suitable composition to the surface, biofilm formation may be inhibited or prevented. The composition may be painted onto the relevant surface or may be reacted with or polymerised onto the relevant surface. Treatment may be made to the surface in situ or prior to assembly.

[0040] In a further aspect, medical devices or implants may be coated with the coating composition of the invention and so coated medical devices or implants are provided. Such devices or implants include catheters, artificial heart valves, surgical pins, pacemaker capsules, prosthetic joints, stents, shunts, endotracheal or gastrointestinal tubes, surgical or dental instruments, surgical suture, dental implants, electrodes, dialysis devices and bandages.

#### DETAILED DESCRIPTION OF THE INVENTION

[0041] The present invention will now be described in further detail, by way of example only, with reference to the following Examples and accompanying drawings, in which:

[0042] FIG. 1 compares the biofilm inhibitory activity of furanone and thiophenone against *Staphylococcus epidermidis*

[0043] FIG. 2 shows a bar chart comparing biofilm inhibitory activity of polymer coatings according to the invention;

[0044] FIG. 3 shows the effect of thiophenone coatings on bacteria desorbed from steel substrates;

[0045] FIG. 4 shows the effect of thiophenone coatings on biofilm growth on steel substrates;

[0046] FIG. 5 shows inhibition of AI-1 quorum-sensing by furanone and thiophenone according to the invention; and

[0047] FIG. 6 shows inhibition of AI-2 quorum-sensing by furanone and thiophenone according to the invention.

#### EXAMPLES

##### Example 1

[0048] This example relates to the synthesis of the thiophenones. The compound codes (e.g. Thio101) correspond to those set out in Table 1.

[0049] Thio101 and Thio102

[0050] (E)- and (Z)-5-Bromomethylenethiophen-2(5H)-one. Acetyl bromide (0.5 mL) was added dropwise at 0° C. to a solution of 5-formyl-2-methoxythiophene<sup>1</sup> (142 mg) in CDCl<sub>3</sub> (1.0 mL). The mixture was stirred at 0° C. for 1.5 h before it was evaporated. The crude product was purified by flash chromatography using hexane/ethyl acetate 5:1 as eluent. Yield (E)-5-bromomethylenethiophen-2(5H)-one: 9 mg. Yield (Z)-5-bromomethylenethiophen-2(5H)-one: 86 mg. The identity of the compounds were confirmed by mass spectrometry and NMR.

[0051] Thio103

[0052] (Z)-5-Choromethylenethiophen-2(5H)-one. Acetyl chloride (2 mL) was added to 5-formyl-2-methoxythiophene<sup>1</sup> (142 mg). The mixture was stirred at room temperature over night before it was evaporated. The crude product was purified by flash chromatography using hexane/ethyl acetate 5:1

as eluent. Yield: (30 mg). The identity of the compound was confirmed by mass spectrometry and NMR.

[0053] Thio104

[0054] (Z)-5-Acetyloxymethylenethiophen-2(5H)-one.

Acetyl bromide (246 mg) was added dropwise at 0° C. to a solution of 5-formyl-2-methoxythiophene<sup>1</sup> (142 mg) in CDCl<sub>3</sub> (1.0 mL). The mixture was stirred at room temperature over night before it was evaporated. The crude product was purified by flash chromatography using hexane/ethyl acetate 5:1 as eluent. Yield: 40 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0055] 3-Bromo-5-formyl-2-methoxythiophene.

5-Formyl-2-methoxythiophene<sup>1</sup> (2.13 g) was dissolved in dichloromethane (20 mL) at room temperature and N-bromosuccinimide (3.20 g) was added. The mixture was stirred over night. The reaction was diluted with ether (50 mL) and extracted with water. The organic phase was dried (MgSO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography (gradient elution: 0-25% EtOAc in hexanes). Yield: 2.53 g. The identity of the compound was confirmed by mass spectrometry and NMR.

[0056] Thio105

[0057] (Z)-3-Bromo-5-bromomethylenethiophen-2(5H)-one. Acetyl bromide (3.0 mL) was added to a solution of 3-bromo-5-formyl-2-methoxythiophene (882 mg) in dichloromethane (5 mL). The reaction was stirred for 48 h before it was diluted with ether (40 mL) and extracted with NaOH (1.0M, aq) and water. The organic phase was dried (MgSO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography (gradient elution: 0-20% EtOAc in hexanes). Yield: 301 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0058] Thio106 and Thio108

[0059] 5-Dibromomethylenethiophen-2(5H)-one and 3-bromo-5-dibromomethylenethiophen-2(5H)-one. Bromine (0.5 mL, 2M in CCl<sub>4</sub>) was added dropwise at 0° C. to a solution of (Z)-5-bromomethylenethiophen-2(5H)-one (40 mg) in CDCl<sub>3</sub> (1 mL). The mixture was stirred at room temperature for 4 h before another portion of bromine (0.2 mL) was added. Stirring was continued for 2 h before the mixture was evaporated and diisopropylethylamine (44 mg) added. The mixture was stirred for 2 h before diethyl ether was added and the organic phase washed with aqueous HCl (1M). The solution was dried (MgSO<sub>4</sub>) and evaporated. The crude product was purified by flash chromatography using hexane/ethyl acetate 8:1 as eluent. Yield 3-bromo-5-dibromomethylenethiophen-2(5H)-one: 16 mg. Yield 5-dibromomethylenethiophen-2(5H)-one: 2 mg. The identity of the compounds were confirmed by mass spectrometry and NMR.

[0060] Thio 107

[0061] (Z)-5-(2,2-Dibromoethylidene)thiophen-2(5H)-one. Acetyl bromide (1 mL) was added dropwise at 0° C. to a solution of 2-(2,2-dibromovinyl)-5-methoxythiophene (230 mg) in CDCl<sub>3</sub> (2.0 mL). The mixture was stirred at room temperature for 5 h before it was evaporated. Ether was added and the solution washed with aqueous NaHCO<sub>3</sub> and brine. The dried solution (MgSO<sub>4</sub>) was evaporated and the crude product was purified by flash chromatography using hexane/ethyl acetate 5:1 as eluent. Yield: 105 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0062] Thio109

[0063] (Z)-5-Benzylidenethiophen-2(5H)-one. Acetyl bromide (0.035 mL) was added at 0° C. to a solution of (5-methoxythiophen-2-yl)(phenyl)methanol<sup>3</sup> (103 mg) in CDCl<sub>3</sub> (1.0

mL.) The mixture was stirred at 0° C. for 30 min before diethyl ether was added and the solution washed with aqueous NaHCO<sub>3</sub>. The crude product was purified by flash chromatography using hexane/ethyl acetate 5:1 as eluent. Yield: 44 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0064] Thio110

[0065] (Z)-3-Bromo-5-benzylidenethiophen-2(5H)-one. Bromine (0.05 mL, 2M in CCl<sub>4</sub>) was added at 0° C. to a solution of (Z)-5-benzylidenethiophen-2(5H)-one (16 mg) in CDCl<sub>3</sub> (1 mL). The mixture was stirred for 24 h at room temperature before it was evaporated and dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 mL). Diisopropylethylamine (39 mg) was added and the mixture stirred for 2 h before washing with HCl (1 M). The crude product was purified by flash chromatography using hexane/ethyl acetate 5:1 as eluent. Yield: 16 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0066] Thio111 and Thio112

[0067] (E)- and (Z)-5-Bromo(phenyl)methylenethiophen-2(5H)-one. Acetyl bromide (1.5 mL) was added dropwise at room temperature to a solution of 5-benzoyl-2-methoxythiophene<sup>2</sup> (100 mg) in CH<sub>2</sub>Cl<sub>2</sub> (1.0 mL). The mixture was stirred at room temperature for 8 d and under reflux for 4 h before it was evaporated. The crude product was purified by flash chromatography using hexane/ethyl acetate 8:1 as eluent. Yield bromo(phenyl)methylenethiophen-2(5H)-one: 38 mg. Yield (Z)-5-bromo(phenyl)methylenethiophen-2(5H)-one: 17 mg. The identity of the compounds were confirmed by mass spectrometry and NMR.

[0068] Thio113

[0069] (Z)-3-Bromomethyl-5-bromomethylenethiophen-2(5H)-one. 3-Chloromethyl-5-formyl-2-methoxythiophene (400 mg) was dissolved in dichloromethane (4 mL) at room temperature and acetyl bromide (1.6 mL) was added. The mixture was stirred for 96 h at room temperature before it was diluted with ether (20 mL) and extracted with NaOH (1.0M, aq) and water. The organic phase was dried (MgSO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography (gradient elution: 0-20% EtOAc in hexanes). Yield: 264 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0070] Thio115


[0071] (Z)-3-Hydroxymethyl-5-bromomethylenethiophen-2(5H)-one. (Z)-3-Bromomethyl-5-bromomethylenethiophen-2(5H)-one (970 mg) was dissolved in 20 mL acetone/water (9:1) at room temperature in a round bottom flask covered with aluminium foil before silver triflate (2.62 g) was added. The mixture was stirred for 24 h at room temperature before it was diluted with ether (50 mL) and extracted with brine. The organic phase was dried (MgSO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography (gradient elution: 0-40% EtOAc in hexanes). Yield: 550 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0072] Thio202


[0073] 2-(2,2-Dibromovinyl)-5-methoxythiophene: Tetra-bromomethane (0.70 g) and triphenylphosphine (1.0 g) were successively added to a solution of 5-formyl-2-methoxythiophene<sup>1</sup> (282 mg) in dichloromethane (10 mL). After stirring for 5 min was another portion of triphenylphosphine (0.2 g) was added and the mixture stirred at 0° C. for 30 min. The mixture was then filtered through a short pad of silica gel and purified by flash chromatography using hexane/ethyl

acetate 10:1 as eluent. Yield: 230 mg. The identity of the compound was confirmed by mass spectrometry and NMR.


[0074] Thio301



[0075] Thio302



[0076] Thio304



[0077] Thio305

[0078] (Z)-5-((3,4-Dibromo-5-methoxythiophen-2-yl)methylene)thiophen-2(5H)-one. Bromine (0.035 mL, 2M in CCl<sub>4</sub>) was added at 0° C. to a solution of (Z)-5-((5-methoxythiophen-2-yl)methylene)thiophen-2(5H)-one<sup>4</sup> (12 mg) in CDCl<sub>3</sub> (1 mL). The mixture was stirred for 1 h at 0° C. before ether was added. The organic solution was washed with aqueous thiosulfate, dried (MgSO<sub>4</sub>) and evaporated. Yield: 15 mg. The identity of the compound was confirmed by NMR.

[0079] 3-Chloromethyl-5-formyl-2-methoxythiophene.

5-Formyl-2-methoxythiophene (141 mg) was dissolved in dichloromethane (1 mL) at 0° C. before chloromethyl ethyl ether (0.48 mL) was added followed by TiCl<sub>4</sub> (0.17 mL). The mixture was stirred for 2 hours at room temperature before was diluted with dichloromethane (20 mL) and extracted with water and brine. The organic phase was dried (MgSO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography (gradient elution: 0-25% EtOAc in hexanes). Yield: 83 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0080] (5-Bromomethylene-2-oxo-2,5-dihydrothiophen-3-yl)methyl acrylate and (5-chloromethylene-2-oxo-2,5-dihydrothiophen-3-yl)methyl acrylate. (Z)-3-Hydroxymethyl-5-bromomethylenethiophen-2(5H)-one (220 mg) was dissolved in dichloromethane (2 mL) before acryloyl chloride (180 mg) and triethylamine (0.15 g) were added. The mixture was stirred for 1 h at room temperature before it was diluted with ether (25 mL) and extracted with water. The organic phase was dried (MgSO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography (gradient elution: 0-20% EtOAc in hexanes) to give a 1:1 mixture of the title compounds. Yield: 213 mg. The identity of the compounds were confirmed by mass spectrometry and NMR.

[0081] Benzyl 3-(triethoxysilyl)propylcarbamate To benzyl alcohol (2.16 g) was (3-isocyanatopropyl)triethoxysilane (5.2 mL) added and the mixture stirred at 85° C. for 3 h. The reaction was dried in vacuo. The crude product was purified by flash chromatography (gradient elution: 0-10% EtOAc in hexanes). Yield: 6.53 g. The identity of the compound was confirmed by NMR.

[0082] (Z)-5-bromomethylene-2-oxo-2,5-dihydrothiophen-3-yl)methyl 3-(triethoxysilyl)propylcarbamate. A mixture of (Z)-3-Hydroxymethyl-5-bromomethylenethiophen-2(5H)-one (44 mg) and triethoxy(3-isocyanatopropyl)silane (248 mg) in dry toluene (1 mL) was heated at 60° C. over night. The solvent was evaporated off and the crude product was purified by flash chromatography using hexane/ethyl

acetate 2:1 as eluent. Yield 40 mg. The identity of the compound was confirmed by NMR.

## REFERENCES

- [0083] 1) Profft, Elmar, Justus Liebigs Annalen der Chemie (1959), 622 196-200.  
 [0084] 2) Pearson, D. E.; Buehler, Calvin A. Synthesis (1972), (10), 533-42.  
 [0085] 3) Lavrushin, V. F.; Nikitchenko, V. M.; Trusevich, N. D.; Pedchenko, N. F.; Kanate, B.; Pivnenko, N. S.; Pogonina, R. I. Khar'k. Gos. Univ. im. Gor'kogo, Kharkov, USSR. Editor(s): Gal'pern, G. D. Tezisy Dokl. Nauchn. Sess. Khim. Tekhnol. Org. Soedin. Sery Semistykh Neftei, 13th (1974), 182-3.  
 [0086] 4) H. Halvorsen, H. Hope and J. Skramstad, Synth. Commun. 37 (2007) 1167-1177

## Example 2a

[0087] This example describes the effect of thiophenones on biofilm formation by various bacteria. Biofilm formation was measured according to a static biofilm model and according to a shaking biofilm model and it was shown that the various thiophenones tested were found to inhibit or prevent biofilm formation.

[0088] According to the static biofilm model, a given thiophenone 200  $\mu\text{mol/L}$  was dissolved in 500  $\mu\text{l}$  absolute ethanol and applied to wells of a standard 24 well microtiter plate. The ethanol was evaporated from the wells in a laminar air sterile work bench at room temperature so as to leave a coating of the thiophenone in the well. A sample of bacteria was then added to the well and incubated for a given period of time. After incubation, the percentage of bacteria remaining was assessed by safranine staining of the biofilm. Bound safranine was released by acetic acid and optical density was measured in Synergy HT Multi-Detection Microplate Reader and compared to a control. The results are set out in Table 1 below.

[0089] According to the shaking biofilm model, the microtiter plates were shaken (200 rpm) in a Minitron Incubator Shaker during biofilm formation.

[0090] Table 1 shows the results of tests on various thiophene structures in this example. The structure and name of each thiophene is given, together with the bacteria tested. The percentage of bacteria remaining in the biofilm is shown, together with the time of incubation.

[0091] It may be concluded from these results that Thiophenone inhibits biofilm formation by various bacteria. The thiophenone effect is mediated through interference with quorum-sensing communication via AI-1 and AI-2. The thiophenone is more effective than furanone (FIGS. 1-4).

TABLE 1






Effect of thiophenones on biofilm formation by various bacteria						
Structure	Name		Static biofilm model		Shaking biofilm model	
			% Remaining	Time incubation	% Remaining	Time incubation
	Thio101	<i>S. epidermidis</i>	28	6 h	13	6 h
	(Z)-5-(bromomethylene)	<i>S. epidermidis</i>			15	20 h
	thiophen-2(5H)-one	<i>E. faecalis</i>			54	8 h
	Chemical Formula: C <sub>5</sub> H <sub>3</sub> BrOS	<i>E. faecalis</i>			56	18 h
	Molecular Weight: 191.05	<i>V. harveyi</i> coating	66	4 h		
		<i>V. harveyi</i> in medium	32	4 h		
		<i>V. harveyi</i> coating	20	4 h		
		<i>V. harveyi</i> in medium	9	4 h		
		Pseudo-alteromonas coating	52	4 h		
		Pseudo-alteromonas in medium	36	4 h		
	Thio102	<i>S. epidermidis</i>	95	6 h	91	6 h
	(E)-5-(bromomethylene)	<i>S. epidermidis</i>			103	20 h
						
	Thio103	<i>S. epidermidis</i>	32	6 h	92	6 h
	(Z)-5-(chloromethylene)	<i>S. epidermidis</i>			96	20 h
	thiophen-2(5H)-one					
						
	Chemical Formula: C <sub>5</sub> H <sub>3</sub> ClOS					
	Molecular Weight: 146.59					
	Thio104	<i>S. epidermidis</i>	34	6 h	106	6 h
	(Z)-5-(5-oxothiophen-2(5H)-ylidene)methyl acetate	<i>S. epidermidis</i>			103	20 h
	Chemical Formula: C <sub>7</sub> H <sub>6</sub> O <sub>3</sub> S					
	Molecular Weight: 170.19					

TABLE 1-continued










Effect of thiophenones on biofilm formation by various bacteria					
Structure	Name	Static biofilm model		Shaking biofilm model	
		% Remaining	Time incubation	% Remaining	Time incubation
	Thio105			20	6 h
	(Z)-3-bromo-5-(bromomethylene)thiophen-2(5H)-one Chemical Formula: C <sub>5</sub> H <sub>2</sub> Br <sub>2</sub> OS Molecular Weight: 269.94			18	20 h
	Thio106			11	6 h
	5-(dibromomethylene)thiophen-2(5H)-one Chemical Formula: C <sub>5</sub> H <sub>2</sub> Br <sub>2</sub> OS Molecular Weight: 269.94			27	20 h
	Thio107			77	6 h
	(Z)-5-(2,2-dibromoethylidene)thiophen-2(5H)-one Chemical Formula: C <sub>6</sub> H <sub>4</sub> Br <sub>2</sub> OS Molecular Weight: 283.97			93	20 h
	Thio108			1	6 h
	3-bromo-5-(dibromomethylene)thiophen-2(5H)-one Chemical Formula: C <sub>5</sub> HBr <sub>3</sub> OS Molecular Weight: 348.84			9	20 h
	Thio109			98	6 h
	(Z)-5-benzylidenethiophen-2(5H)-one Chemical Formula: C <sub>11</sub> H <sub>8</sub> OS Molecular Weight: 188.25			107	20 h
	Thio110			30	6 h
	(Z)-5-benzylidene-3-bromothiophen-2(5H)-one Chemical Formula: C <sub>11</sub> H <sub>7</sub> BrOS Molecular Weight: 267.14			27	20 h
	Thio111			63	6 h
	(E)-5-(bromo(phenyl)methylene)thiophen-2(5H)-one Chemical Formula: C <sub>11</sub> H <sub>7</sub> BrOS Molecular Weight: 267.14			50	20 h
	Thio112			63	6 h
	(Z)-5-(bromo(phenyl)methylene)thiophen-2(5H)-one Chemical Formula: C <sub>11</sub> H <sub>7</sub> BrOS Molecular Weight: 267.14			59	20 h
	Thio201	56	6 h	81	6 h
	5-methoxythiophene-2-carbaldehyde Chemical Formula: C <sub>6</sub> H <sub>6</sub> O <sub>2</sub> S Molecular Weight: 142.18			101	20 h

TABLE 1-continued









Effect of thiophenones on biofilm formation by various bacteria					
Structure	Name	Static biofilm model		Shaking biofilm model	
		% Remaining	Time incubation	% Remaining	Time incubation
	Thio202 2-(2,2-dibromovinyl)- 5-methoxythiophene Chemical Formula: C <sub>7</sub> H <sub>6</sub> Br <sub>2</sub> OS Molecular Weight: 297.99			94 103	6 h 20 h
	Thio301 (Z)-5-(thiophen-2-ylmethylene) thiophen-2(5H)-one Chemical Formula: C <sub>9</sub> H <sub>6</sub> OS <sub>2</sub> Molecular Weight: 194.27			80 98	6 h 20 h
	Thio302 (Z)-5-((5-bromothiophen-2-yl) methylene)thiophen-2(5H)-one Chemical Formula: C <sub>9</sub> H <sub>5</sub> BrOS <sub>2</sub> Molecular Weight: 273.17			98 101	6 h 20 h
	Thio303 (Z)-5-((5-oxothiophen-2(5H)- ylidene)methyl)thiophene- 2-carbaldehyde Chemical Formula: C <sub>10</sub> H <sub>6</sub> O <sub>2</sub> S <sub>2</sub> Molecular Weight: 222.28			95 91	6 h 20 h
	Thio304 (Z)-5-((5-methoxythiophen-2- yl)methylene)thiophen- 2(5H)-one Chemical Formula: C <sub>10</sub> H <sub>8</sub> O <sub>2</sub> S <sub>2</sub> Molecular Weight: 224.30			72 94	6 h 20 h
	Thio305 (Z)-5-((3,4-dibromo-5- methoxythiophen-2- yl)methylene)thiophen- 2(5H)-one Chemical Formula: C <sub>10</sub> H <sub>6</sub> Br <sub>2</sub> O <sub>2</sub> S <sub>2</sub> Molecular Weight: 382.09			66 101	6 h 20 h

TABLE 1-continued

Effect of thiophenones on biofilm formation by various bacteria						
Structure	Name		Static biofilm model		Shaking biofilm model	
			% Remaining	Time incubation	% Remaining	Time incubation
	Thio306 (Z)-5-(2,2'-bithiophen-5-ylmethylene)-thiophen-2(5H)-one Chemical Formula: C <sub>13</sub> H <sub>8</sub> OS <sub>3</sub> Molecular Weight: 276.40	<i>S. epidermidis</i>	67	6 h		
	Thi401 5-(hydroxymethyl)-5-((5-methoxythiophen-2-yl)methyl)thiophen-2(5H)-one Chemical Formula: C <sub>11</sub> H <sub>12</sub> O <sub>3</sub> S <sub>2</sub> Molecular Weight: 256.34	<i>S. epidermidis</i> <i>S. epidermidis</i>	51	6 h	80 94	6 h 20 h

## Example 2b

[0092] This example describes the effect of further thiophenes on biofilm formation and planktonic growth by various bacteria.

[0093] Planktonic growth was determined in "Low Binding Plates" in which the bacteria form minimal amounts of biofilm. The quantity is determined by optical density measurements at 600 nm.


[0094] Biofilm formation was measured in static cultures in wells of microtiter plates for *S. epidermidis*, or on "peggs" according to the Calgary method (The Calgary biofilm devices: New technology for rapid determination of antibiotic susceptibilities of bacterial biofilms. Ceri et al. J Clin Microbiol 1999;37:1771-1776) for *V. harveyi*. In both cases the safranin staining method was applied and optical density was measured at 530 nm for quantification of biofilm mass.

[0095] This compound was synthesized as follows:


[0096] Hünig's base (155 mg, 1.2 mmol) and DMAP (catalytic amount, ~10 mg) was dissolved in 2 mL DCM and added to a solution of succinic anhydride (120 mg, 1.2 mmol) and (Z)-5-(bromomethylene)-3-(hydroxymethyl)thiophen-2(5H)-one (0.22 g, 1.0 mmol) in 4 mL DCM at room temperature. The reaction mixture was stirred for 30 minutes, diluted with 25 mL DCM and washed with 3\*5 mL water. The combined aqueous phases were extracted with 2\*10 mL ether. The combined organic phases were dried over MgSO<sub>4</sub>, filtrated and the solvents were removed in vacuo. The residue was dissolved in a small amount of THF and ether(1:2), and the product was precipitated by addition of pentane. The solution was filtered, leaving a yellow solid. Yield: 220 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0097] Biofilm formation by *V. harveyi* was measured using the Calgary method. There was a 17% reduction at a thiophenone concentration of 5 µM, 70% reduction at 50 µM and 85% reduction at 100 µM. This represents a strong effect on *V. harveyi* biofilm reduction. However a weak effect on planktonic bacteria of 18% reduction at 100 µM was observed.

Thio401



Thio402






### Synthesis of 2'-methoxy-2,3'-bithiophene-5'-carbaldehyde

**[0098]** 4-bromo-5-methoxythiophene-2-carbaldehyde (0.22 g, 1 mmol), tributyl(thiophen-2-yl)stannane (0.75 g, 2 mmol),  $\text{PdCl}_2(\text{PhCN})_2$  (38 mg, 0.1 mmol) and  $\text{PPh}_3$  (79 mg, 0.3 mmol) were dissolved in N,N-dimethylformamide (3 mL) and stirred at 50° C. for 24 h. The reaction mixture was cooled to room temperature, diluted with 25 mL ether, and washed with 3\*10 mL water. The combined aqueous phases were extracted with 10 mL ether, and the combined organic phases were dried over  $\text{MgSO}_4$ , filtrated and the solvents removed in vacuo. The product was purified by flash column chromatography on silica (0-20% EtOAc in hexanes). Yield: 187 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

### Synthesis of (Z)-5-(bromomethylene)-3-(thiophen-2-yl)thiophen-2(5H)-one

**[0099]** Acetyl bromide (0.56 mL, 7.5 mmol) was added dropwise to a solution of 2'-methoxy-2,3'-bithiophene-5'-carbaldehyde (0.11 g, 0.5 mmol) in 6 mL DCM at 0° C. The mixture was heated to room temperature and stirred for 3 days. The mixture was diluted with 10 mL ether, washed with 10 mL NaOH (aq, 1M) and 2\*5 mL water. The combined aqueous phases were extracted with 25 mL DCM, and the combined organic phases were dried over  $\text{MgSO}_4$ , filtrated and the solvents were removed in vacuo. The product mixture was purified by flash column chromatography on silica (0-15% EtOAc in hexanes). Yield: 44 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

**[0100]** Using the Calgary method, a *V. harveyi* biofilm reduction of 40 to 50% was observed in the presence of the thiophenone at a concentration of from 40 to 100  $\mu\text{M}$ . This compound is therefore less effective than Thio401. A 30% reduction in planktonic bacteria was observed at 100  $\mu\text{M}$ .



(Z)-5-(iodomethylene)thiophen-2(5H)-one


**[0101]** This compound was synthesized as follows:

**[0102]** (Z)-5-(Bromomethylene)thiophen-2(5H)-one (103 mg) in acetone (2 mL) was added to a solution of sodium iodide (530 mg) in acetone (3 mL). The mixture was stirred for 3 days at room temperature before ether and water was added. The aqueous phase was extracted with ether and the combined organic phase was dried ( $\text{MgSO}_4$ ) and evaporated. The crude product was purified by flash chromatography (hexane/EtOAc 5:1). Yield 97 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

**[0103]** This compound showed a biofilm reduction of 46% at 5  $\mu\text{M}$ , 80% at 10  $\mu\text{M}$  and 95% at 15  $\mu\text{M}$  when tested against *V. harveyi* bacteria. This compound is therefore strongly effective against biofilm formation. It is also strongly effective on planktonic bacterial growth showing a 70 to 80% reduction at 15  $\mu\text{M}$  to 100  $\mu\text{M}$  for *V. harveyi*.

**[0104]** The same compound was tested against *S. epidermis* and found to be less effective. Biofilm reduction was 50% at

50  $\mu\text{M}$  and 64% at 100  $\mu\text{M}$ . Planktonic bacterial growth reduction was 50% at 50  $\mu\text{M}$  and 63% at 100  $\mu\text{M}$ .



(E)-5-(1-bromoethylidene)thiophen-2(5H)-one

Thio404


**[0105]** This compound was synthesized as follows:

**[0106]** Acetyl bromide (0.017 mL) was added at room temperature to a solution of 5-acetyl-2-methoxythiophene<sup>REF</sup> (21 mg) in  $\text{CDCl}_3$  (1.0 mL). The mixture was stirred at room temperature for 24 h before another portion of acetyl bromide (0.15 mL) was added. The mixture was stirred for 3 d before it was evaporated and purified by flash chromatography (hexane/EtOAc 5:1). Yield E-isomer: 11 mg. Yield Z-isomer: 5 mg. The identity of the compounds were confirmed by mass spectrometry and NMR.

**[0107]** REF

**[0108]** Sice, Jean, Journal of the American Chemical Society (1953), 75 3697-700

**[0109]** This compound was tested on *V. harveyi* for biofilm reduction and showed 70 to 80% reduction at 50 to 100  $\mu\text{M}$ , which may be considered a moderate effect. A relatively minor effect on planktonic growth was observed with 40 to 55% reduction at 50 to 100  $\mu\text{M}$ .



(Z)-5-(1-thiocyanatomethylene)thiophen-2(5H)-one

Thio405

**[0110]** This compound was synthesized as follows:

**[0111]** Ammonium isothiocyanate (34 mg) was added to a solution of (Z)-5-(1-bromomethylidene)thiophen-2(5H)-one (22 mg) in acetone. The mixture was stirred at room temperature for 1 h before ether and water was added. The aqueous phase was extracted with ether and the combined organic phase was dried ( $\text{MgSO}_4$ ) and evaporated. The identity of the compound was confirmed by mass spectrometry and NMR.


**[0112]** This compound had a strong effect on biofilm reduction and planktonic growth in *S. epidermis*. A biofilm reduction of 90 to 100% was observed at concentrations from 25  $\mu\text{M}$ . Planktonic growth was reduced by 70% at thiophenone concentrations from 25  $\mu\text{M}$ .

**[0113]** The compound was also found to have a strong effect on *V. harveyi* biofilm reduction showing a 90% reduction at 25  $\mu\text{M}$ . Planktonic reduction for *V. harveyi* was 50% at 25  $\mu\text{M}$ .

### Example 2c (Comparative Example)

**[0114]** This example follows the same methodology as Example 2b. However, the substituent groups on the thiophenones are unsuitable to render the compound capable of

blocking or interfering with quorum-sensing microbial communication or preventing or inhibiting biofilm formation.




Thio501

(Z)-5-((diethylamino)methylene)thiophen-2(5H)-one

[0115] This compound was synthesized as follows:

[0116] A mixture of diethyl amine (47 mg) in  $\text{CDCl}_3$  (1.5 mL) was slowly added to a solution of (Z)-5-(1-bromomethylidene)thiophen-2(5H)-one (46 mg) in  $\text{CDCl}_3$  (1.5 mL) at  $0^\circ\text{C}$ . The mixture was stirred for 2.5 h before it was evaporated. The crude product was purified by flash chromatography ( $\text{CHCl}_3/\text{MeOH}$  20:1). Yield: 58 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0117] No effect on *V. harveyi* biofilm or planktonic growth.




Thio502

(Z)-5-((triethylamino)methylene)thiophen-2(5H)-one, bromide salt

[0118] This compound was synthesized as follows:

[0119] A mixture of triethyl amine (10 mg) in  $\text{CDCl}_3$  (1 mL) was slowly added to a solution of (Z)-5-(1-bromomethylidene)thiophen-2(5H)-one (14 mg) in  $\text{CDCl}_3$  (1 mL) at  $0^\circ\text{C}$ . The mixture was stirred at room temperature for 24 h before it was evaporated. The identity of the compound was confirmed by NMR.

[0120] No effect on *S. epidermis* biofilm or planktonic growth.



Thio503


(Z)-5-((4-nitrophenoxy)methylene)thiophen-2(5H)-one

[0121] This compound was synthesized as follows:

[0122] A mixture of (Z)-5-(1-bromomethylidene)thiophen-2(5H)-one (20 mg), p-nitrophenol (58 mg) and triethyl amine (42 mg) in  $\text{CDCl}_3$  (1.5 mL) was stirred at room temperature for 18 h before it was evaporated. The crude

product was purified by flash chromatography ( $\text{CH}_2\text{Cl}_2$ ). Yield: 21 mg. The identity of the compound was confirmed by mass spectrometry and NMR.

[0123] No effect on *S. epidermis* biofilm or planktonic growth.



Thio504

4-(5-methoxythiophen-2-yl)dihydrothiophen-2(3H)-one

[0124] This compound was synthesized as follows:


[0125] Dry HCl was bubbled through a solution of 2-methoxythiophene (224 mg) in  $\text{CDCl}_3$  (2 mL) at  $0^\circ\text{C}$ . for ca 2 min. The mixture was left at room temperature over night, evaporated and purified by Flash chromatography. hexane/EtOAc 5:1. Yield 123 mg. The identity of the compound was confirmed by mass spectrometry and NMR

[0126] No effect on *V. harveyi* biofilm or planktonic growth.

### Example 3


[0127] This example relates to the synthesis of polymeric thiophenones and thiophenones containing functional groups for adhesion to surfaces.

[0128] Formation of a Co-Polymer from Compound 1 and Styrene:



[0129] 320 mg of compound 1, 2.88 g of styrene and 160 mg of AIBN (Azo Bis Isobutyronitrile) were added to 6 mL of toluene and degassed with argon for 30 minutes at rt. The solution was then stirred at  $70^\circ\text{C}$ . for 24 h., cooled to room temperature, and added to 25 mL pentane. The precipitate was washed two times with pentane and dried under high vacuum at room temperature overnight.

[0130] The copolymer formed in this way has the predicted general formula




[0131] in which  $n$  is an integer. The thiophene agent forms an end group or side chain.

#### Formation of a Styrene Polymer (Comparative Example)

[0132] 10 g styrene was added to 10 mL of toluene and 100 mg of AIBN. The solution was degassed with argon for 30 min at room temperature. The solution was stirred for 3 h at 70° C., cooled to room temperature, and added to 40 mL of pentane. The precipitate was washed two times with pentane and dried under high vacuum at room temperature overnight.

[0133] Formation of a Co-Polymer from Compound 2) and Tert-Butyl Acrylate:




2

X:Cl/Br, ca 1:1

[0134] 240 mg of compound 2, 2.16 g of tert-butyl acrylate and 120 mg of AIBN were added to 5 mL of toluene and degassed with argon for 30 minutes at room temperature. The solution was stirred at 70° C. for 24 h., cooled to room temperature, and added to 25 mL pentane. The precipitate was washed two times with pentane and dried under high vacuum at room temperature overnight.

[0135] The monomer units of the copolymer formed in this way may be linked together as shown below:



#### Formation of a Tert-Butyl Acrylate Polymer (Comparative Example)

[0136] 10 g tert-butyl acrylate was added to 10 mL of toluene and 100 mg of AIBN. The solution was degassed with argon for 30 min at room temperature. The solution was stirred for 1 h at 70° C., cooled to room temperature, and added to 40 mL of pentane. The residue was washed two times with pentane and dried under high vacuum at room temperature overnight.

#### Formation of Benzyl 3-(triethoxysilyl)propylcarbamate (Compound E)

[0137] To benzyl alcohol (2.16 g) was (3-isocyanatopropyl)triethoxysilane (5.2 mL) added and the mixture stirred at 85° C. for 3 h. The reaction was dried in vacuo. The crude product was purified by flash chromatography (gradient elution: 0-10% EtOAc in hexanes). Yield: 6.53 g. The identity of the compound was confirmed by NMR.

#### Formation of (Z)-(5-bromomethylene-2-oxo-2,5-dihydrothiopen-3-yl)methyl 3-(triethoxysilyl)propylcarbamate (Compound F):

[0138] A mixture of (Z)-3-Hydroxymethyl-5-bromomethylenethiophen-2(5H)-one (44 mg) and triethoxy(3-isocyanatopropyl)silane (248 mg) in dry toluene (1 mL) was heated at 60° C. over night. The solvent was evaporated off and the crude product was purified by flash chromatography using hexane/ethyl acetate 2:1 as eluent. Yield: 40 mg. The identity of the compound was confirmed by NMR

#### Example 4

[0139] This example relates to the effects of surface coatings on biofilm formation by bacteria.

[0140] The materials of Example 3 were tested in a static biofilm model as follows.

[0141] Triplicate samples of a coating composition were applied to a glass vial in accordance with Table 2 set out below where each vial is given a number. Vials 1 to 12 are each loaded with 13 mg of the indicated material, followed by the addition of 1.0 mL dichloromethane. The dichloromethane was slowly evaporated under atmospheric pressure to leave a coating on the internal walls of the receptacles. Vials 13 to 18 were each loaded by adding 10 mg of the indicated material pre-dissolved in 10 mL dichloromethane. These vials were heated in an oven at 125° C. for 24 hours. To remove any residues of the dichloromethane solvent, all vials were dried at high vacuum (0.5 mmHg) for 2 hours at room temperature.

TABLE 2

Glass Vial	Material
1, 2 and 3	Copolymer of A and B (~10% by weight of A and ~90% by weight of B)
4, 5 and 6	Polymer of B
7, 8 and 9	Polymer of C
10, 11 and 12	Copolymer of C and D (~10% by weight of D and ~90% by weight of C)
13, 14 and 15	E
16, 17 and 18	F

TABLE 2-continued







Glass Vial	Material
	  
	X = Cl and Br, approx. 1:1
	 
	A
	D
	E

TABLE 2-continued

Glass Vial	Material
	
	F

[0142] FIG. 2 shows the results of incubating bacteria *S. epidermidis* over a time period of 4 hours. The biofilm was assessed as described in Example 2.

[0143] These results show that thiophene polymers have inhibitory activity in relation to biofilm formation.

[0144] The results also show that thiophenes according to the invention may be covalently bound to a surface so as to inhibit biofilm formation. It is possible in the example of the copolymer of C and D that bacteria are killed in addition to the inhibition of biofilm formation. Similar results were obtained in a static biofilm model where *S. epidermidis* bacteria were incubated over a time period of 5 hours. The results are shown in Table 3.

TABLE 3







Effect of polymers on biofilm formation			
Structure	Strain	Static biofilm model	
		% remaining	incubation time
	<i>S. Epidermidis</i>	100	5 h
	<i>S. Epidermidis</i>	100	5 h
	<i>S. Epidermidis</i>	100	5 h
	<i>S. Epidermidis</i>	33	5 h

TABLE 3-continued

Effect of polymers on biofilm formation			
Structure	Strain	Static biofilm model	
		% remaining	incubation time
	<i>S. Epidermidis</i>	100	5 h
	<i>S. Epidermidis</i>	45	5 h

## Example 5

**[0145]** Coating and Biological Testing of Steel Samples with Compound F

**[0146]** In the previous Example, Compound F was attached to a glass substrate, where it decreased biofilm formation. The current experiment was performed to determine whether the compound would attach to steel substrates and decrease biofilm formation and thereby biologically induced corrosion in a marine environment.

**[0147]** Coating experiments were performed on stainless steel substrates (30×40 mm<sup>2</sup>) were received from Ole Øystein Knudsen. To determine a good procedure for coating the substrates, the following samples were prepared and investigated by X-ray photoelectron spectroscopy (XPS):

**[0148]** A) Coated with T<sub>B</sub>18210, heated to 120°, rinsed, 120° C.

**[0149]** B) Coated with T<sub>B</sub>18210, rinsed, heated to 120° C., rinsed, 120° C.

**[0150]** C) Rinsed substrate, heated to 120° C., rinsed, 120° C.

**[0151]** Coating Procedure

**[0152]** All substrates were rinsed in an ultrasonic bath for 1 min in dichloromethane, 1 min in acetone and finally 1 min in isopropanol. They were etched for 3 min in 20 wt % HNO<sub>3</sub> at room temperature, then rinsed in RO water and dried at ambient conditions.

**[0153]** A 0.01 M solution of Compound F was prepared by dissolving 0.02366 g in 5 ml toluene.

**[0154]** Two rinsed substrates (for samples A and B) were wiped with the Compound F solution for 1 min. A) was immediately put into an oven at 120° C. for 1 hour. B) was first rinsed in toluene to remove any excess T<sub>B</sub>18210, then placed in an oven at 120° C. for 1 hour. Sample C) was not coated, just heated to 120° C. for 1 hour.

**[0155]** After cooling, all samples were rinsed in toluene and heated at 120° C. for 1 hour.

**[0156]** Results

**[0157]** XPS showed that the Compound F is attached to sample A), but not on sample B). Samples of type B) were therefore not prepared for biological testing.

**[0158]** Samples for Biological Testing

**[0159]** Three samples of each of type A) and C) were prepared for testing of biofilm formation in a marine environment. For comparison, three Cu-covered samples (marked D) were prepared by painting rinsed substrates with "aqua-net" Cu paint from Steen/Hansen Maling. The painted samples were dried at ambient conditions over night.

**[0160]** Biological Testing

**[0161]** Biological testing of coated stainless steel substrates were performed in a static cultures of marine bacteria. The biofilm prevention efficiencies of the coatings were recorded as a) total enumeration of bacteria attached on the steel surfaces and b) most-probable number (MPN) viability testing of attached bacteria. The nature of the thiophenes (expecting to inhibit biofilm generation without being toxic to the bacteria) should account for inhibition of both total counts and viability counts when thiophene-coated surfaces were compared to non-treated controls.

**[0162]** Methods

**[0163]** Three samples of stainless steel substrates were prepared as described above, each of these in triplicate. A primary culture of marine bacteria were prepared by inoculation of 2 ml seawater (collected from 90 m depth through a pipeline system supplying seawater to SINTEF Sealab, Trondheim) in 100 ml Marine Broth 2216 (MB). The cultures was incubated at 20° C. with continuous agitation until significant increase in medium turbidity (cell density appr. 10<sup>9</sup> cells/ml). Nine 500 ml sterile bottles (PE) with wide necks were prepared with individual steel surfaces placed in 200 ml MB. Primary culture (1.0 ml) was inoculated to each bottle and the cultures with steel surfaces incubated at 20° C. for 14 days with careful continuous agitation, and with medium changes after day 3, 7, and 10. During medium changes 75% of the culture media in each bottle was replaced by fresh MB medium.

**[0164]** At the end of the experimental period (14 days) the steel surfaces were removed from the media and carefully washed in sterile and particle-free (filtered through 0.2 µm sterile filters) seawater. Washed steel surfaces were placed in 200 ml sterile and particle-free seawater and placed in an ultrasound bath for 15 minutes to desorb cells from the sur-

faces. Desorbed concentrations of bacteria were recorded by epifluorescence microscopy and by MPN-counts in MB medium.

[0165] For epifluorescence microscopy 10 ml desorbed samples were stained with 2  $\mu$ g 3'5-diamino phenylindol (DAPI) and incubated for 5-10 minutes. The stained samples were filtered through 0.2  $\mu$ m black polycarbonate filters and analysed in a fluorescence microscope at 1250 $\times$  magnification. For MPN-counts 10-fold serial dilutions of desorbed samples were prepared in sterile seawater and 0.2 ml of each dilution (undiluted to 10<sup>-9</sup> dilution) inoculated in triplicate in 2 ml MB medium in 24-well sterile tissue culture plates. The plates were incubated at 20° C. for 5 days and positive growth recorded as turbidity in the wells. Concentrations of viable bacteria in the desorbed samples were determined from MPN-tables with 95% confidence intervals.

#### [0166] Results

[0167] The results from the epifluorescence and MPN-counts are shown in FIG. 3.

[0168] FIG. 3 shows total (A) and viable (B) counts of desorbed bacteria from stainless steel surfaces coated with thiophene (Compound F), with no coating and with Cu coating. The results are calculated as concentrations of cells per cm<sup>2</sup> surface.

[0169] The inhibition of bacterial attachment (total counts) and growth (viable counts) are shown in FIG. 4.

[0170] FIG. 4 shows the inhibition of bacterial attachment (total counts) and growth (median values) for stainless steel surfaces coated with thiophene Compound F and with Cu.

[0171] The results of FIGS. 3 and 4 showed that the thiophene inhibited bacterial attachment and viability at a level comparable with Cu, showing appr. 80% inhibition of biofilm attachment and >99% inhibition of viable attached bacteria when median values were compared, although concentrations of viable bacteria differed significantly for the viable counts. While Cu is harmful in high concentrations the thiophene is regarded as a non-toxic chemical. It is important to emphasize that the bacterial concentrations in the surrounding environment was much higher than expected in normal seawater, and that experiments were conducted at temperatures much higher than normal in Norwegian seawaters.

#### Example 6

[0172] This example relates to the effects of thiophenones according to the invention on quorum-sensing communication between bacteria.


[0173] The effect of (Z)-5-(bromomethylene) thiophen-2 (5H)-one on AI-1 and AI-2 quorum-sensing by bacteria was tested in a bioluminescence assay as follows:

[0174] Inhibition of quorum sensing was assessed as the ability of the furanone to reduce bioluminescence induced by cell free bacterial culture supernatant containing either AI-1 or AI-2 signal molecules. *Vibrio harveyi* BB886 was used as an AI-1 reporter and *Vibrio harveyi* BB170 was used as reporter of AI-2 communication.


[0175] As a comparison, the corresponding furanone compound was also tested. It was found that the thiophenone compound was considerably more effective in inhibiting quorum-sensing by both AI-1 and AI-2.

[0176] The results are set out in FIGS. 5 and 6 for AI-1 and AI-2 quorum-sensing respectively.


1. An agent comprising the compound according to general formula (I):



wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> are each independently H or a substituent, and wherein at least one of R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> is halogen, cyano, cyanate, thiocyanate or C<sub>1</sub>-C<sub>6</sub> haloalkyl, with the proviso that the compound is neither nor




2. An agent for use in medicine, which agent comprises the compound of general formula (II):



wherein X is O, S, NH or NR', in which R' is an optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl group;

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are each independently H or a substituent, and wherein the compound is capable of blocking or interfering with quorum-sensing microbial communication.

3. An agent for use in medicine, which agent comprises the compound of general formula (II):




wherein X is O, S, NH or NR', in which R' is an optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl group;

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are each independently H or a substituent, and wherein the compound is capable of preventing or inhibiting biofilm formation.

4. The agent according to claim 2 or claim 3 wherein X is O.

5. An agent according to any preceding claim wherein each substituent is independently a halogen, cyano, cyanate, thiocyanate, alkyl, alkoxy, haloalkyl, alkyl ester, alkylsilyl, alkenyl, alkynyl, aryl, or arylalkyl, which may be substituted or unsubstituted, optionally interrupted by one or more heteroatoms, straight chain or branched chain.

6. An agent according to any preceding claim wherein each substituent is independently halogen, haloalkyl, alkoxy, alkyl ester, phenyl or



wherein R<sub>5</sub>, R<sub>6</sub> and R<sub>7</sub> are each independently H, Br, Cl, OMe, or CHO.

7. An agent according to any preceding claim wherein R<sub>4</sub> is a substituent.

8. An agent according to any preceding claim wherein at least one of R<sub>1</sub>, R<sub>3</sub> and R<sub>4</sub> is halogen or C<sub>1</sub>-C<sub>6</sub> haloalkyl.

9. An agent according to any preceding claim wherein at least one of R<sub>1</sub> and R<sub>4</sub> is halogen or C<sub>1</sub>-C<sub>6</sub> haloalkyl.

10. An agent according to any preceding claim wherein R<sub>1</sub> and R<sub>2</sub> are each independently H or halogen.

11. An agent according to any preceding claim wherein R<sub>3</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub> haloalkyl or phenyl.

12. An agent according to any preceding claim wherein R<sub>4</sub> is halogen, C<sub>1</sub>-C<sub>6</sub> haloalkyl or phenyl.

13. An agent according to any preceding claim wherein the halogen is Br.

14. An agent according to any preceding claim wherein R<sub>1</sub> is Br.

15. An agent according to any preceding claim wherein R<sub>4</sub> is Br.

16. An agent according to any preceding claim wherein R<sub>2</sub> is H.

17. An agent according to any preceding claim wherein R<sub>3</sub> is H.

18. An agent according to any of claims 1 to 13 wherein R<sub>1</sub> and R<sub>4</sub> are each Br and R<sub>2</sub> and R<sub>3</sub> are each H.

19. An agent according to any of claims 1 to 13 wherein R<sub>3</sub> and R<sub>4</sub> are each Br and R<sub>1</sub> and R<sub>2</sub> are each H.

20. An agent according to any of claims 1 to 13 wherein R<sub>1</sub>, R<sub>3</sub> and R<sub>4</sub> are each Br and R<sub>2</sub> is H.

21. An agent according to any of claims 1 to 13 wherein R<sub>1</sub> is —CH<sub>2</sub>—O—CO—(CH<sub>2</sub>)<sub>2</sub>—COOH and R<sub>4</sub> is Br.

22. An agent according to any of claims 1 to 13 wherein R<sub>1</sub> is thienyl and R<sub>4</sub> is Br.

23. An agent according to any of claims 1 to 13 wherein R<sub>3</sub> is methyl and R<sub>4</sub> is Br.



24. An agent according to claim 5, wherein R<sub>3</sub> is H and R<sub>4</sub> is thiocyanate.

25. A polymer which comprises a compound as defined in any preceding claim.



26. A polymer according to claim 25 which comprises one or more side chain functional groups comprising the compound.

27. A polymer according to claim 25 or claim 26 which comprises a polyacrylate, polymethacrylate, polycrotonate, polyvinyl alcohol, polyvinyl acetate, polystyrene, acrylonitrile or siloxane.

28. A polymer according to any of claims 25 to 27, which is obtainable by polymerising the compound according to general formula (III) with the compound according to general formula (IV):



29. A polymer according to any of claims 25 to 27, which is obtainable by polymerising the compound according to general formula (V) with the compound according to general formula (VI):



30. An agent or polymer for use in medicine, which comprises the compound as defined in any preceding claim or a pharmaceutically-acceptable salt, ester or prodrug thereof.

31. An agent or polymer according to any preceding claim for use as an antimicrobial.

32. An agent or polymer according to claim 31 for use in the prevention or treatment of microbial infection.

33. An agent or polymer according to claim 32 wherein the microbial infection is a bacterial or fungal infection.

34. An agent or polymer according to any one of claims 31 to 33, which interferes with quorum-sensing microbial communication so as to treat or prevent a condition mediated by microbes which are regulated by quorum-sensing communication.

35. A composition comprising an agent or polymer according to any preceding claim and one or more further antimicrobial agents as a combined preparation for simultaneous, separate or sequential use in the prevention or treatment of microbial infection.

**36.** A composition according to claim **35** wherein the antimicrobial agent is an antibiotic or an antifungal agent.

**37.** An agent, polymer or composition according to any of claims **31** to **36** for use in the treatment of periodontitis, gingivitis, dental caries, topical infections, respiratory infections, eye infections, ear infections or localised organ infection.


**38.** A coating composition comprising the agent or polymer according to any preceding claim.

**39.** A coating composition according to claim **38** which is capable of binding covalently to a surface.

**40.** A coating composition according to claim **38** or **39**, which comprises the agent is covalently linked to the group  $\text{Si}(\text{OR}_5)_3$ , wherein each  $\text{R}_5$  is independently substituted or unsubstituted  $\text{C}_1$ - $\text{C}_6$  hydrocarbyl.

**41.** A coating composition according to claim **40**, which comprises the compound according to general formula (VII):

(VII)



**42.** An antimicrobial surface cleanser comprising an agent or polymer as defined in any preceding claim.

**43.** An antimicrobial surface cleanser according to claim **42** which is a disinfectant or cleaning composition.

**44.** An antimicrobial surface cleanser according to claim **42** or claim **43** wherein the surface is an inanimate surface.

**45.** An antimicrobial surface cleanser according to claim **42** or claim **44** wherein the surface is the skin of a human or animal.

**46.** An anti-biofouling composition or coating comprising an agent or polymer as defined in any preceding claim.

**47.** An anti-biofouling composition according to claim **46** wherein the biofouling occurs on marine vessels, sub marine installations, pipelines, water pipes, industrial machines or installations, water systems, cooling towers, heat exchangers, chain lubrication systems, oil or gas platforms, fish-farming installations or surfaces, machines, tools or devices used in food production.

**48.** A personal care product comprising an agent or polymer as defined in any preceding claim which is a personal hygiene article, shampoo, soap, deodorant, dentifrice, mouthwash, contact lens solution or cosmetic composition.

**49.** A medical device or implant which is coated with the coating composition as defined in any of claims **38** to **41**.

**50.** A medical device or implant according to claim **49**, which is a catheter, artificial heart valve, surgical pin, pacemaker capsule, prosthetic joint, stent, shunt, endotracheal or gastrointestinal tube, surgical or dental instrument, surgical suture, dental implant, electrode, dialysis device or bandage.

\* \* \* \* \*