

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2008 (15.05.2008)

PCT

(10) International Publication Number
WO 2008/057938 A1

(51) International Patent Classification:

C07D 471/08 (2006.01) **A61K 31/407** (2006.01)
A61P 25/00 (2006.01) **C07D 487/04** (2006.01)

Matt, S. [US/US]; 2074 Craig Street, Winston-salem, North Carolina 27103 (US).

(21) International Application Number:

PCT/US2007/083330

(74) Agent: **MASSEY, JR., Carl B.**; Womble Carlyle Sandridge & Rice, PLLC, P.O. Box 7037, Atlanta, Georgia 30357 (US).

(22) International Filing Date:

1 November 2007 (01.11.2007)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/856,079 2 November 2006 (02.11.2006) US

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **TARGACEPT, INC.** [US/US]; 200 East First Street, Suite 300, Winston-salem, North Carolina 27101-4165 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MAZUROV, Anatoly** [UA/US]; 3704 Timberoak Drive, Greensboro, North Carolina 27410 (US). **MIAO, Lan** [US/US]; 226 March Ferry Road, Advance, North Carolina 27006 (US). **XIAO, Yun-de** [CN/US]; 2079 Waterford Village Drive, Clemmons, North Carolina 27012 (US). **HAMMOND, Philip, S.** [US/US]; 1162 Bradley Road, Pinnacle, North Carolina 27043 (US). **MILLER, Craig, H.** [US/US]; 1564 Sharon Road, Winston-salem, North Carolina 27103 (US). **AKIREDDY, Srinivasa, Rao** [IN/US]; 5829 Fox Point Circle, Winston-salem, North Carolina 27104 (US). **MURTHY, V. Srinivasa** [IN/US]; 595 Stonebridge Drive, Winston-salem, North Carolina 27104 (US). **WHITAKER, Regina C.** [US/US]; 3257 U.S. Hwy. 158, Mocksville, North Carolina 27028 (US). **BREINING, Scott R.** [US/US]; 2363 Rosewood Avenue, Winston-salem, North Carolina 27103 (US). **MELVIN,**

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

(54) Title: NICOTINIC ACETYLCHOLINE RECEPTORSUB-TYPE SELECTIVE AMIDES OF DIAZABICYCLOALKANES

(57) Abstract: Compounds, pharmaceutical compositions including the compounds, and methods of preparation and use thereof are disclosed. The compounds are amide compounds which can be prepared from certain heteroaryl carboxylic acids and certain diazabicycloalkanes. The compounds exhibit selectivity for, and bind with high affinity to, neuronal nicotinic receptors of the $\alpha 4 \beta 2$ subtype in the central nervous system (CNS). The compounds and compositions can be used to treat and/or prevent a wide variety of conditions or disorders, particularly CNS disorders. The compounds can: (i) alter the number of nicotinic cholinergic receptors of the brain of the patient, (ii) exhibit neuroprotective effects, and (iii) when employed in effective amounts, not result in appreciable adverse side effects (e.g. side effects such as significant increases in blood pressure and heart rate, significant negative effects upon the gastrointestinal tract, and significant effects upon skeletal muscle).

WO 2008/057938 A1

SUB-TYPE SELECTIVE AMIDES OF DIAZABICYCLOALKANES

Field of the Invention

The present invention relates to compounds that bind to and modulate the activity of neuronal nicotinic acetylcholine receptors, to processes for preparing these compounds, to pharmaceutical compositions containing these compounds and to methods of using these compounds for treating a wide variety of conditions and disorders, including those associated with dysfunction of the central nervous system (CNS).

10

Background of the Invention

The therapeutic potential of compounds that target neuronal nicotinic receptors (NNRs), also known as nicotinic acetylcholine receptors (nAChRs), has been the subject of several recent reviews (see Breining et al., *Ann. Rep. Med. Chem.* **40**: 3 (2005), Hogg and Bertrand, *Curr. Drug Targets: CNS Neurol. Disord.* **3**: 123 (2004), Suto and Zacharias, *Expert Opin. Ther. Targets* **8**: 61 (2004), Dani et al., *Bioorg. Med. Chem. Lett.* **14**: 1837 (2004), Bencherif and Schmitt, *Curr. Drug Targets: CNS Neurol. Disord.* **1**: 349 (2002)). Among the kinds of indications for which NNR ligands have been proposed as therapies are cognitive disorders, including Alzheimer's disease, attention deficit disorder and schizophrenia (Newhouse et al., *Curr. Opin. Pharmacol.* **4**: 36 (2004), Levin and Rezvani, *Curr. Drug Targets: CNS Neurol. Disord.* **1**: 423 (2002), Graham et al., *Curr. Drug Targets: CNS Neurol. Disord.* **1**: 387 (2002), Ripoll et al., *Curr. Med. Res. Opin.* **20**(7): 1057 (2004), and McEvoy and Allen, *Curr. Drug Targets: CNS Neurol. Disord.* **1**: 433 (2002)); pain and inflammation (Decker et al., *Curr. Top. Med. Chem.* **4**(3): 369 (2004), Vincler, *Expert Opin. Invest. Drugs* **14**(10): 1191 (2005), Jain, *Curr. Opin. Inv. Drugs* **5**: 76 (2004), Miao et al., *Neuroscience* **123**: 777 (2004)); depression and anxiety (Shytle et al., *Mol. Psychiatry* **7**: 525 (2002), Damaj et al., *Mol. Pharmacol.* **66**: 675 (2004), Shytle et al., *Depress. Anxiety* **16**: 89 (2002)); neurodegeneration (O'Neill et al., *Curr. Drug Targets: CNS Neurol. Disord.* **1**: 399 (2002), Takata et al., *J. Pharmacol. Exp. Ther.* **306**: 772 (2003), Marrero et al., *J. Pharmacol. Exp. Ther.* **309**: 16 (2004)); Parkinson's disease

(Jonnala and Buccafusco, *J. Neurosci. Res.* **66**: 565 (2001)); addiction (Dwoskin and Crooks, *Biochem. Pharmacol.* **63**: 89 (2002), Coe et al., *Bioorg. Med. Chem. Lett.* **15**(22): 4889 (2005)); obesity (Li et al., *Curr. Top. Med. Chem.* **3**: 899 (2003)); and Tourette's syndrome (Sacco et al., *J. Psychopharmacol.* **18**(4): 457
5 (2004), Young et al., *Clin. Ther.* **23**(4): 532 (2001)).

A limitation of some nicotinic compounds is that they are associated with various undesirable side effects, for example, by stimulating muscle and ganglionic receptors. It would be desirable to have compounds, compositions and methods for preventing and/or treating various conditions or disorders (e.g., CNS disorders),
10 including alleviating the symptoms of these disorders, where the compounds exhibit nicotinic pharmacology with a beneficial effect (e.g., upon the functioning of the CNS), but without significant associated side effects. It would further be highly desirable to provide compounds, compositions and methods that affect CNS function without significantly affecting those receptor subtypes which have the
15 potential to induce undesirable side effects (e.g., appreciable activity at cardiovascular and skeletal muscle sites). The present invention provides such compounds, compositions and methods.

Summary of the Invention

20 The present invention provides certain amide compounds which can be formed from certain heteroaryl carboxylic acids and certain diazabicycloalkanes, particularly 3,7-diazabicyclo[3.3.0]octane and 3,7-diazabicyclo[3.3.1]nonane. These amide compounds bind with high affinity to NNRs of the $\alpha 4\beta 2$ subtype, found in the CNS, and exhibit selectivity for the $\alpha 4\beta 2$ subtype over the $\alpha 7$ NNR
25 subtype, also found in the CNS. The present invention also relates to pharmaceutically acceptable salts prepared from these compounds and the pharmaceutical compositions thereof, which can be used for treating and/or preventing a wide variety of conditions or disorders, and particularly those disorders characterized by dysfunction of nicotinic cholinergic neurotransmission
30 or the degeneration of the nicotinic cholinergic neurons. Also provided are methods for treating and/or preventing disorders, such as CNS disorders, and also for treating certain conditions (e.g., alleviating pain and inflammation), in mammals in need of such treatment. The methods involve administering to a

subject a therapeutically effective amount of the compounds (including salts) or pharmaceutical compositions including such compounds. Further provided is a method for treatment of disorders selected from the group consisting of age-associated memory impairment, mild cognitive impairment, pre-senile dementia
5 (early onset Alzheimer's disease), senile dementia (dementia of the Alzheimer's type), Lewy body dementia, vascular dementia, Alzheimer's disease, stroke, AIDS dementia complex, attention deficit disorder, attention deficit hyperactivity disorder, dyslexia, schizophrenia, schizophreniform disorder, and schizoaffective disorder. Even further provided is a method for treatment of disorders selected
10 from the group consisting of the treatment of mild to moderate dementia of the Alzheimer's type, attention deficit disorder, mild cognitive impairment and age associated memory impairment.

The pharmaceutical compositions incorporate a compound of the present invention which, when employed in effective amounts, interacts with relevant
15 nicotinic receptor sites of a subject, and hence acts as a therapeutic agent to treat and prevent a wide variety of conditions and disorders. The pharmaceutical compositions provide therapeutic benefit to individuals suffering from such disorders and exhibiting clinical manifestations of such disorders, in that the compounds within those compositions, when employed in effective amounts, can
20 (i) exhibit nicotinic pharmacology and affect relevant nicotinic receptors sites (e.g., act as a pharmacological agonist to activate nicotinic receptors), and/or (ii) elicit neurotransmitter secretion, and hence prevent and suppress the symptoms associated with those diseases. In addition, the compounds have the potential to (i) increase the number of nicotinic cholinergic receptors of the brain of the patient,
25 (ii) exhibit neuroprotective effects, and/or (iii) when employed in effective amounts, to not cause appreciable adverse side effects (e.g., significant increases in blood pressure and heart rate, significant negative effects upon the gastro-intestinal tract, and significant effects upon skeletal muscle). The pharmaceutical compositions comprising the compounds of the invention, are believed to be safe
30 and effective with regards to prevention and treatment of a wide variety of conditions and disorders.

The foregoing and other aspects of the present invention are explained in detail in the detailed description and examples set forth below.

Brief Description of the Figures

Figure 1 is a chart showing the results of a study on object recognition in rats treated orally with N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane. The results are shown as a function of recognition index (%) versus dose (mg/kg).

Figure 2 is a chart showing the results of a study on object recognition in rats treated orally with N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane. The results are shown as a function of recognition index (%) versus dose (mg/kg).

Detailed Description

The subtype selective compounds, pharmaceutical compositions including these compounds, methods of preparing the compounds, and methods of treatment and/or prevention using the compounds are described in detail below.

The compounds and methods described herein will be better understood with reference to the following preferred embodiments. The following definitions will be useful in defining the scope of the invention:

In this specification, unless stated otherwise, the term "alkyl" includes both straight and branched chain alkyl groups. These may be, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, neo-pentyl, n-hexyl or i-hexyl. The term "C₁₋₄ alkyl" thus includes alkyl groups having 1 to 4 carbon atoms, including, but are not limited to, methyl, ethyl, n-propyl, i-propyl or *tert*-butyl.

In this specification, unless stated otherwise, the term "cycloalkyl" refers to an optionally substituted, partially or completely saturated monocyclic, bicyclic or bridged hydrocarbon ring system. The term "C₃₋₈ cycloalkyl" may be, but is not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

As used herein, heterocyclyl radicals contain from 3 to 10 members including one or more heteroatoms selected from oxygen, sulfur and nitrogen. Examples of suitable heterocyclyl moieties include, but are not limited to, piperidinyl, morpholinyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl,

isothiazolidinyl, thiazolidinyl, isoxazolidinyl, oxazolidinyl, piperazinyl, oxanyl (tetrahydropyranyl), and oxolanyl (tetrahydrofuranlyl).

As used herein, C₁₋₆ alkoxy radicals contain from 1 to 6 carbon atoms in a straight or branched chain, and also include C₃₋₆ cycloalkoxy radicals and alkoxy radicals that contain C₃₋₆ cycloalkyl moieties. Examples include, but are not limited to methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, isobutoxy, cyclopropylmethoxy, allyloxy or propargyloxy.

As used herein, "aromatic" refers to 3 to 10, preferably 5 and 6-membered ring aromatic and heteroaromatic rings.

As used herein, "aromatic group-containing species" refers to moieties that are or include an aromatic group. Accordingly, phenyl and benzyl moieties are included in this definition, as both are or include an aromatic group, and pyridinyl and pyrimidinyl are included in the definition, as both are heteroaromatic, a subset of aromatic.

As used herein, aryl radicals are selected from phenyl, naphthyl and indenyl.

As used herein, heteroaryl radicals contain from 3 to 10 members, preferably 5 or 6 members, including one or more heteroatoms selected from oxygen, sulfur and nitrogen. Examples of suitable 5-membered ring heteroaryl moieties include furanyl, thienyl, pyrrolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, tetrazolyl, triazolyl, and pyrazolyl. Examples of suitable 6-membered heteroaryl moieties include pyridinyl, pyrimidinyl, pyrazinyl, and pyridazinyl. Examples of 9-membered heteroaryl groups include benzimidazolyl, indolizinyl, indolyl, purinyl and indolinyl. Examples of 10-membered heteroaryl groups include quinolinyl and isoquinolinyl.

It will be appreciated that throughout the specification, the number and nature of substituents on rings in the compounds of the invention will be selected so as to avoid sterically undesirable combinations.

Certain compound names of the present invention were generated with the aid of computer software (ACDLabs 8.0/Name(IUPAC)).

Examples of suitable pharmaceutically acceptable salts include inorganic acid addition salts such as chloride, bromide, sulfate, phosphate, and nitrate; organic acid addition salts such as acetate, galactarate, propionate, succinate,

lactate, glycolate, malate, tartrate, citrate, maleate, fumarate, methanesulfonate, p-toluenesulfonate, and ascorbate; salts with acidic amino acid such as aspartate and glutamate; alkali metal salts such as sodium salt and potassium salt; alkaline earth metal salts such as magnesium salt and calcium salt; ammonium salt; organic basic
5 salts such as trimethylamine salt, triethylamine salt, pyridine salt, picoline salt, dicyclohexylamine salt, and N,N'-dibenzylethylenediamine salt; and salts with basic amino acid such as lysine salt and arginine salt. The salts may be in some cases hydrates or ethanol solvates. Representative salts are provided as described in U.S. Patent Nos. 5,597,919 to Dull et al., 5,616,716 to Dull et al. and 5,663,356
10 to Ruecroft et al.

The compounds of Formula I and pharmaceutically acceptable salts thereof may exist in solvated, for example hydrated, as well as unsolvated forms, and the present invention encompasses all such forms.

As used herein, an "agonist" is a substance that stimulates its binding
15 partner, typically a receptor. Stimulation is defined in the context of the particular assay, or may be apparent in the literature from a discussion herein that makes a comparison to a factor or substance that is accepted as an "agonist" or an "antagonist" of the particular binding partner under substantially similar circumstances as appreciated by those of skill in the art. Stimulation may be
20 defined with respect to an increase in a particular effect or function that is induced by interaction of the agonist or partial agonist with a binding partner and can include allosteric effects.

As used herein, an "antagonist" is a substance that inhibits its binding partner, typically a receptor. Inhibition is defined in the context of the particular
25 assay, or may be apparent in the literature from a discussion herein that makes a comparison to a factor or substance that is accepted as an "agonist" or an "antagonist" of the particular binding partner under substantially similar circumstances as appreciated by those of skill in the art. Inhibition may be defined with respect to a decrease in a particular effect or function that is induced by
30 interaction of the antagonist with a binding partner, and can include allosteric effects.

As used herein, a "partial agonist" or a "partial antagonist" is a substance that provides a level of stimulation or inhibition, respectively, to its binding partner

that is not fully or completely agonistic or antagonistic, respectively. It will be recognized that stimulation, and hence, inhibition is defined intrinsically for any substance or category of substances to be defined as agonists, antagonists, or partial agonists.


5 As used herein, "intrinsic activity" or "efficacy" relates to some measure of biological effectiveness of the binding partner complex. With regard to receptor pharmacology, the context in which intrinsic activity or efficacy should be defined will depend on the context of the binding partner (*e.g.*, receptor/ligand) complex and the consideration of an activity relevant to a particular biological outcome. For
10 example, in some circumstances, intrinsic activity may vary depending on the particular second messenger system involved. See Hoyer, D. and Boddeke, H., *Trends Pharmacol. Sci.* **14**(7): 270-5 (1993). Where such contextually specific evaluations are relevant, and how they might be relevant in the context of the present invention, will be apparent to one of ordinary skill in the art.

15 As used herein, modulation of a receptor includes agonism, partial agonism, antagonism, partial antagonism, or inverse agonism of a receptor.

As used herein, neurotransmitters whose release is mediated by the compounds described herein include, but are not limited to, acetylcholine, dopamine, norepinephrine, serotonin and glutamate, and the compounds described
20 herein function as modulators at the $\alpha 4\beta 2$ subtype of the CNS NNRs.

Compounds

The compounds described herein are amide compounds formed from certain heteroaryl carboxylic acids and certain diazabicycloalkanes. These
25 compounds can be represented as Formula I:



Formula I

wherein n has the value of 0 or 1, and Cy is a heteroaryl group chosen from the

group of 2-furanyl, 3-furanyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 1,3,4-oxadiazol-2-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 1,3,4-thiadiazol-2-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl and
 5 4-pyridinyl, which heteroaryl groups are optionally substituted with up to three non-hydrogen substituents independently selected from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₂₋₆ alkenyl, substituted C₂₋₆ alkenyl, C₂₋₆ alkynyl, substituted C₂₋₆ alkynyl, C₃₋₈ heterocyclyl, substituted C₃₋₈ heterocyclyl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₅₋₁₀ aryl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ aryl, substituted C₅₋₁₀
 10 heteroaryl, C₁₋₆ alkyl-C₅₋₁₀ aryl, C₁₋₆ alkyl-C₅₋₁₀ heteroaryl, substituted C₁₋₆ alkyl-C₅₋₁₀ aryl, substituted C₁₋₆ alkyl-C₅₋₁₀ heteroaryl, C₅₋₁₀ aryl-C₁₋₆ alkyl, C₅₋₁₀ heteroaryl-C₁₋₆ alkyl, substituted C₅₋₁₀ aryl-C₁₋₆ alkyl, substituted C₅₋₁₀ heteroaryl-C₁₋₆ alkyl, halo, -OR', -NR'R'', -CF₃, -CN, -NO₂, -C₂R', -SR', -N₃, -C(=O)NR'R'', -NR'C(=O)R'', -C(=O)R', -C(=O)OR', -OC(=O)R', -OC(=O)NR'R'', -NR'C(=O)OR'',
 15 -SO₂R', -SO₂NR'R'', and -NR'SO₂R'', where R' and R'' are independently selected from hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ heterocyclyl, C₅₋₁₀ aryl, C₅₋₁₀ heteroaryl or C₅₋₁₀ aryl-C₁₋₆ alkyl, or R' and R'' and the atoms to which they are attached together can form a C₃₋₈ heterocyclic ring, wherein the term "substituted", as applied to alkyl, alkenyl, alkynyl, heterocyclyl, cycloalkyl, aryl, heteroaryl,
 20 alkylaryl, alkylheteroaryl, arylalkyl and heteroarylalkyl, refers to substitution by one or more alkyl, aryl, heteroaryl, halo, -OR' and -NR'R'' groups, or pharmaceutically acceptable salts thereof.

One embodiment of the invention relates to compounds of Formula I wherein n has the value of 0 or 1, and Cy is a C₅₋₁₀ heteroaryl group chosen from
 25 the group of 2-furanyl or 3-furanyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 1,3,4-oxadiazol-2-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 1,3,4-thiadiazol-2-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl and 4-pyridinyl, which heteroaryl groups are optionally substituted
 30 with up to three non-hydrogen substituents independently selected from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, halo, and C₂₋₆ alkynyl substituted with phenyl.

In one embodiment n is 0. In another embodiment n is 1. In a further embodiment Cy is 2-furanyl. In yet another embodiment Cy is 2-furanyl

substituted with halo. In one embodiment Cy is 2-furanyl substituted with chlorine. In yet a further embodiment n is 0 and Cy is 2-furanyl optionally substituted with halo. In one embodiment n is 1 and Cy is 2-furanyl optionally substituted with halo. In yet another embodiment 2-furanyl is substituted on
5 position 5. In another embodiment R' and R" are independently selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, i-butyl, s-butyl or t-butyl. In a further embodiment R' and R" are independently selected from phenyl or benzyl.

In some cases, compounds of the present invention are chiral. The present invention includes all enantiomeric or diastereomeric forms of such compounds.

10 Representative compounds of the present invention include the following:

N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
15 N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(4-phenylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-(2-pyridinyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
20 N-(5-(phenylethynyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(furan-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(oxazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(oxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(oxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
25 N-(isoxazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isoxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-bromoisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-methoxyisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
30 N-(1,2,4-oxadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,2,4-oxadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,3,4-oxadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(thiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,

- N-(thiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(thiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isothiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isothiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
5 N-(isothiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,2,4-thiadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,2,4-thiadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,3,4-thiadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(pyridin-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
10 and pharmaceutically acceptable salts thereof.

Representative compounds of the present invention also include the following:

- N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(3-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
15 N-(5-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(3-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(3-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
20 N-(4-phenylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-(2-pyridinyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-(phenylethynyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(furan-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(oxazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
25 N-(oxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(oxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isoxazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isoxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
30 N-(3-bromoisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(3-methoxyisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(1,2,4-oxadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(1,2,4-oxadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,

N-(1,3,4-oxadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(thiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(thiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(thiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
5 N-(isothiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isothiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isothiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(1,2,4-thiadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(1,2,4-thiadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
10 N-(1,3,4-thiadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(pyridin-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
and pharmaceutically acceptable salts thereof.

One embodiment relates to compound N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane, or pharmaceutically acceptable salts thereof. Another
15 embodiment relates to compound N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane, or pharmaceutically acceptable salts thereof.

Compound Preparation

The compounds of the present invention can be prepared via the coupling
20 of mono-protected diazabicycle (i.e., one in which one of the two amine functional groups is rendered un-reactive by suitable derivatization) with a suitably functionalized heteroaryl acid chloride or other reactive carboxylic acid derivative.

There are numerous methods for preparing the mono-protected diazabicycles used to prepare the compounds of the present invention. Methods for
25 the synthesis of a suitably protected 3,7-diazabicyclo[3.3.0]octane are described in PCT WO 02/070523 to Colon-Cruz et al. and in U.S. application 2006/0019985 to Zhenkun et al., in which N-benzylmaleimide is condensed with either paraformaldehyde and N-benzylglycine or N-(methoxymethyl)-N-(trimethylsilylmethyl)benzylamine to produce 3,7-dibenzyl-3,7-
30 diazabicyclo[3.3.0]octane-2,4-dione (also known as 2,5-dibenzyltetrahydropyrrolo[3,4-c]pyrrole-1,3-dione). Subsequent transformation of this intermediate can follow several paths. In one instance, treatment with α -chloroethylchloroformate produces 3-benzyl-3,7-diazabicyclo[3.3.0]octane-2,4-

dione (also known as 2-benzyltetrahydropyrrolo[3,4-c]pyrrole-1,3-dione), which is then sequentially reduced (using borane-dimethylsulfide complex), converted into its N-(tert-butoxycarbonyl) derivative, and hydrogenated (to remove the second benzyl group). This produces N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane, which can be used in coupling with carboxylic acids, and their derivatives, to produce compounds of the present invention. Alternately, 3,7-dibenzyl-3,7-diazabicyclo[3.3.0]octane-2,4-dione can be reduced (with lithium aluminum hydride), partially hydrogenated (to remove one benzyl group), converted into its N-(tert-butoxycarbonyl) derivative, and hydrogenated (to remove the second benzyl group), to produce N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane. Other methods for installation and removal of the benzyl, tert-butoxycarbonyl, and other amine protecting groups are well known by those skilled in the art and are described further in T. W. Greene and P.G. M. Wuts, Protective Groups in Organic Synthesis, 3rd Edition, John Wiley & Sons, New York (1999).

An alternative preparation of N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane has been described in U.S. applications 2004/0186107 to Schrimpf et al. and 2005/0101602 to Basha et al., and involves the condensation of maleimide and N-(methoxymethyl)-N-(trimethylsilylmethyl)benzylamine to give 7-benzyl-3,7-diazabicyclo[3.3.0]octane-2,4-dione (also known as 5-benzyltetrahydropyrrolo[3,4-c]pyrrole-1,3-dione). Subsequent treatment with a reducing agent (e.g., lithium aluminum hydride) produces the 3-benzyl-3,7-diazabicyclo[3.3.0]octane, the free amine of which can be protected by a tert-butoxycarbonyl group, followed by removal of the benzyl protecting group by hydrogenolysis.

Maleate esters can be used as alternatives to maleimides in these condensation reactions. Thus, according to PCT WO 96/007656 to Schaus et al., condensation of N-benzylglycine with paraformaldehyde and dimethylmaleate will give N-benzyl-cis-3,4-pyrrolidinedicarboxylic acid dimethyl ester. This compound can then be reduced, for example, with lithium aluminum hydride, to give the diol, which can be further reacted with methanesulfonyl chloride in the presence of triethylamine to produce the corresponding dimesylate. Further treatment with ammonia and heat provides the N-benzyl protected 3,7-diazabicyclo[3.3.0]octane.

As described above, this can be converted into N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane.

Suitable derivatives of 3,7-diazabicyclo[3.3.1]nonane (bispidine) can be used to make compounds of the present invention. One such derivative is N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane, which can be made in a variety of ways. One synthesis proceeds through N-benzyl-N'-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane, described by Stead et al. in *Org. Lett.* 7: 4459 (2005). Thus the Mannich reaction between N-(tert-butoxycarbonyl)piperidin-4-one, benzylamine and paraformaldehyde affords N-benzyl-N'-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonan-9-one, which can be treated sequentially with *p*-toluenesulfonylhydrazide and sodium borohydride (to remove the carbonyl oxygen), giving N-benzyl-N'-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane. The benzyl group can be removed as described above to provide N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane. Alternative syntheses of diazabicyclo[3.3.1]nonanes, suitable for conversion to either N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane or another mono-protected derivative, have been described by Jeyaraman and Avila in *Chem. Rev.* **81**(2): 149-174 (1981) and in U.S. Patent 5,468,858 to Berlin et al.

One means of making amides of the present invention is to couple the either N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane or N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane with a suitably functionalized carboxylic acid and then remove the tert-butoxycarbonyl protecting group. Many such carboxylic acids are commercially available, and others can be easily prepared by procedures known to those skilled in the art. The condensation of an amine and a carboxylic acid, to produce an amide, typically requires the use of a suitable activating agent, such as N,N'-dicyclohexylcarbodiimide (DCC), (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), O-(benzotriazol-1-yl)-N,N,N',N'-bis(tetramethylene)uronium hexafluorophosphate (HBTU), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), or (1-ethyl-3-(3-dimethylaminopropyl)carbodiimide) (EDCI) with 1-hydroxybenzotriazole (HOBt).

Other activating agents are well known to those skilled in the art (for example, see Kiso and Yajima, *Peptides*, pp 39-91, Academic Press, San Diego, CA (1995)).

Alternatively, the amide bond can be formed by coupling a mono-protected diazabicyclo with a suitably functionalized acid chloride, which may be available
5 commercially or may be prepared by conversion of the suitably functionalized carboxylic acid. The acid chloride may be prepared by treatment of the appropriate carboxylic acid with, among other reagents, thionyl chloride or oxalyl chloride.

After amide formation, removal of the protecting group (e.g., the tert-butoxycarbonyl group) with acid, either aqueous or anhydrous, will afford the
10 compounds of the present invention.

Those skilled in the art of organic synthesis will appreciate that there exist multiple means of producing compounds of the present invention which are labeled with a radioisotope appropriate to various diagnostic uses. Thus, condensation of a ^{11}C - or ^{18}F -labeled heteroaromatic carboxylic acid with either N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane or N-(tert-butoxycarbonyl)-3,7-
15 diazabicyclo[3.3.1]nonane, using the methods described above, and subsequent removal of the tert-butoxycarbonyl group will produce a compound suitable for use in positron emission tomography.

20 **Methods of Treatment**

The compounds of the present invention are modulators of the $\alpha 4\beta 2$ NNR subtype, characteristic of the CNS, and can be used for preventing and/or treating various conditions or disorders, including those of the CNS, in subjects which have or are susceptible to such conditions or disorders, by modulation of $\alpha 4\beta 2$ NNRs.
25 The compounds have the ability to selectively bind to the $\alpha 4\beta 2$ NNRs and express nicotinic pharmacology (e.g., to act as agonists, partial agonists, antagonists and the like). For example, compounds of the present invention, when administered in effective amounts to patients in need thereof, provide some degree of prevention of the progression of the CNS disorder (i.e., providing protective effects),
30 amelioration of the symptoms of the CNS disorder, and/or amelioration of the reoccurrence of the CNS disorder.

The compounds of the present invention can be used to treat and/or prevent those types of conditions and disorders for which other types of nicotinic

compounds have been proposed as therapeutics. See, for example, the references previously listed in the "Background of the Invention" section, as well as Williams et al., *Drug News Perspec.* **7(4)**: 205 (1994), Arneric et al., *CNS Drug Rev.* **1(1)**: 1-26 (1995), Arneric et al., *Exp. Opin. Invest. Drugs* **5(1)**: 79-100 (1996), Bencherif et al., *J. Pharmacol. Exp. Ther.* **279**: 1413 (1996), Lippiello et al., *J. Pharmacol. Exp. Ther.* **279**: 1422 (1996), Damaj et al., *J. Pharmacol. Exp. Ther.* **291**: 390 (1999); Chiari et al., *Anesthesiology* **91**: 1447 (1999), Lavand'homme and Eisenbach, *Anesthesiology* **91**: 1455 (1999), Holladay et al., *J. Med. Chem.* **40(28)**: 4169-94 (1997), Bannon et al., *Science* **279**: 77 (1998), PCT WO 94/08992, PCT WO 96/31475, PCT WO 96/40682, and U.S. Patent Nos. 5,583,140 to Bencherif et al., 5,597,919 to Dull et al., 5,604,231 to Smith et al. and 5,852,041 to Cosford et al., the disclosures of which are incorporated herein by reference in their entirety.

The compounds and their pharmaceutical compositions are useful in the treatment and/or prevention of a variety of CNS disorders, including neurodegenerative disorders, neuropsychiatric disorders, neurologic disorders, and addictions. The compounds and their pharmaceutical compositions can be used to treat and/or prevent cognitive deficits (age-related and otherwise), attentional disorders and dementias (including those due to infectious agents or metabolic disturbances); to provide neuroprotection; to treat convulsions and multiple cerebral infarcts; to treat mood disorders, compulsions and addictive behaviors; to provide analgesia; to control inflammation (such as mediated by cytokines and nuclear factor kappa B) and treat inflammatory disorders; to provide pain relief; and to treat infections (as anti-infectious agents for treating bacterial, fungal and viral infections). Among the disorders, diseases and conditions that the compounds and pharmaceutical compositions of the present invention can be used to treat and/or prevent are: age-associated memory impairment, mild cognitive impairment, pre-senile dementia (early onset Alzheimer's disease), senile dementia (dementia of the Alzheimer's type), Lewy body dementia, HIV-dementia, vascular dementia, Alzheimer's disease, stroke, AIDS dementia complex, attention deficit disorder, attention deficit hyperactivity disorder, dyslexia, schizophrenia, schizophreniform disorder, schizoaffective disorder, Parkinsonism including Parkinson's disease, Pick's disease, Huntington's chorea, tardive dyskinesia, hyperkinesia, progressive supranuclear palsy, Creutzfeld-Jakob disease, multiple

sclerosis, amyotrophic lateral sclerosis, epilepsy, mania, anxiety, depression, panic disorders, bipolar disorders, generalized anxiety disorder, obsessive compulsive disorder, rage outbursts, Tourette's syndrome, autism, drug and alcohol addiction, tobacco addiction, obesity, cachexia, psoriasis, lupus, acute cholangitis, aphthous stomatitis, asthma, ulcerative colitis, inflammatory bowel disease, pouchitis, viral pneumonitis and arthritis (e.g., rheumatoid arthritis and osteoarthritis), endotoxaemia, sepsis, atherosclerosis, idiopathic pulmonary fibrosis and neoplasias.

It is advantageous that the treatment or prevention of diseases, disorders and conditions occurs without appreciable adverse side effects (e.g., significant increases in blood pressure and heart rate, significant negative effects upon the gastro-intestinal tract, and significant effects upon skeletal muscle). The compounds of the present invention, when employed in effective amounts, can modulate the activity of the $\alpha 4\beta 2$ NNRs without appreciable interaction with the nicotinic subtypes that characterize the human ganglia (as demonstrated by their lack of the ability of to elicit nicotinic function in adrenal chromaffin tissue) or skeletal muscle (as demonstrated by their lack of ability to elicit nicotinic function in cell preparations expressing muscle-type nicotinic receptors). Thus, these compounds are capable of treating and/or preventing diseases, disorders and conditions without eliciting significant side effects associated activity at ganglionic and neuromuscular sites. Thus, administration of the compounds provides a therapeutic window in which treatment of certain diseases, disorders and conditions is provided, and certain side effects are avoided. That is, an effective dose of the compound is sufficient to provide the desired effects upon the disease, disorder or condition, but is insufficient (i.e., is not at a high enough level) to provide undesirable side effects.

Thus, the present invention provides the use of a compound or Formula I, or a pharmaceutically acceptable salt thereof, for use in therapy (such as a therapy described above).

In yet another aspect the present invention provides the use of a compound or Formula I, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment of a CNS disorder (such as a disorder, disease or condition described above).

In a further aspect the invention provides the use of a compound of Formula I, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment mild to moderate dementia of the Alzheimer's type, attention deficit disorder, mild cognitive impairment and age associated
5 memory impairment.

Diagnostic Uses

The compounds can be used in diagnostic compositions, such as probes, particularly when they are modified to include appropriate labels. The probes can
10 be used, for example, to determine the relative number and/or function of specific receptors, particularly the $\alpha 4\beta 2$ receptor subtype. For this purpose the compounds of the present invention most preferably are labeled with a radioactive isotopic moiety such as ^{11}C , ^{18}F , ^{76}Br , ^{123}I or ^{125}I .

The administered compounds can be detected using known detection
15 methods appropriate for the label used. Examples of detection methods include position emission topography (PET) and single-photon emission computed tomography (SPECT). The radiolabels described above are useful in PET (*e.g.*, ^{11}C , ^{18}F or ^{76}Br) and SPECT (*e.g.*, ^{123}I) imaging, with half-lives of about 20.4 minutes for ^{11}C , about 109 minutes for ^{18}F , about 13 hours for ^{123}I , and about 16
20 hours for ^{76}Br . A high specific activity is desired to visualize the selected receptor subtypes at non-saturating concentrations. The administered doses typically are below the toxic range and provide high contrast images. The compounds are expected to be capable of administration in non-toxic levels. Determination of dose is carried out in a manner known to one skilled in the art of radiolabel
25 imaging. See, for example, U.S. Patent No. 5,969,144 to London *et al.*

The compounds can be administered using known techniques. See, for example, U.S. Patent No. 5,969,144 to London *et al.* The compounds can be administered in formulation compositions that incorporate other ingredients, such as those types of ingredients that are useful in formulating a diagnostic
30 composition. Compounds useful in accordance with carrying out the present invention most preferably are employed in forms of high purity. See, U.S. Patent No. 5,853,696 to Elmalch *et al.*

After the compounds are administered to a subject (*e.g.*, a human subject), the presence of that compound within the subject can be imaged and quantified by appropriate techniques in order to indicate the presence, quantity, and functionality of selected NNR subtypes. In addition to humans, the compounds can also be administered to animals, such as mice, rats, dogs, and monkeys. SPECT and PET imaging can be carried out using any appropriate technique and apparatus. See Villemagne *et al.*, In: Arneric *et al.* (Eds.) *Neuronal Nicotinic Receptors: Pharmacology and Therapeutic Opportunities*, 235-250 (1998) and U.S. Patent No. 5,853,696 to Elmalch *et al.* for a disclosure of representative imaging techniques.

The radiolabeled compounds bind with high affinity to selective NNR subtypes (*e.g.*, $\alpha 4\beta 2$) and preferably exhibit negligible non-specific binding to other nicotinic cholinergic receptor subtypes (*e.g.*, those receptor subtypes associated with muscle and ganglia). As such, the compounds can be used as agents for noninvasive imaging of nicotinic cholinergic receptor subtypes within the body of a subject, particularly within the brain for diagnosis associated with a variety of CNS diseases and disorders.

In one aspect, the diagnostic compositions can be used in a method to diagnose disease in a subject, such as a human patient. The method involves administering to that patient a detectably labeled compound as described herein, and detecting the binding of that compound to selected NNR subtypes (*e.g.*, $\alpha 4\beta 2$ receptor subtype). Those skilled in the art of using diagnostic tools, such as PET and SPECT, can use the radiolabeled compounds described herein to diagnose a wide variety of conditions and disorders, including conditions and disorders associated with dysfunction of the central and autonomic nervous systems. Such disorders include a wide variety of CNS diseases and disorders, including Alzheimer's disease, Parkinson's disease, and schizophrenia. These and other representative diseases and disorders that can be evaluated include those that are set forth in U.S. Patent No. 5,952,339 to Bencherif *et al.*

In another aspect, the diagnostic compositions can be used in a method to monitor selective nicotinic receptor subtypes of a subject, such as a human patient. The method involves administering a detectably labeled compound as described

herein to that patient and detecting the binding of that compound to selected nicotinic receptor subtypes (*e.g.*, the $\alpha 4\beta 2$ receptor subtype).

Pharmaceutical Compositions

5 According to one embodiment of the present invention there is provided a pharmaceutical composition comprising as active ingredient a therapeutically effective amount of a compound of the present invention, in association with one or more pharmaceutically acceptable diluents, excipients and/or inert carriers.

 The manner in which the compounds are administered can vary. The
10 compositions are preferably administered orally (*e.g.*, in liquid form within a solvent such as an aqueous or non-aqueous liquid, or within a solid carrier). Preferred compositions for oral administration include pills, tablets, capsules, caplets, syrups, and solutions, including hard gelatin capsules and time-release capsules. Compositions may be formulated in unit dose form, or in multiple or
15 subunit doses. Preferred compositions are in liquid or semisolid form. Compositions including a liquid pharmaceutically inert carrier such as water or other pharmaceutically compatible liquids or semisolids may be used. The use of such liquids and semisolids is well known to those of skill in the art.

 The compositions can also be administered via injection, *i.e.*, intravenously,
20 intramuscularly, subcutaneously, intraperitoneally, intraarterially, intrathecally, and intracerebroventricularly. Intravenous administration is a preferred method of injection. Suitable carriers for injection are well known to those of skill in the art, and include 5% dextrose solutions, saline, and phosphate buffered saline. The compounds can also be administered as an infusion or injection (*e.g.*, as a
25 suspension or as an emulsion in a pharmaceutically acceptable liquid or mixture of liquids).

 The formulations may also be administered using other means, for example, rectal administration. Formulations useful for rectal administration, such as suppositories, are well known to those of skill in the art. The compounds can also
30 be administered by inhalation (*e.g.*, in the form of an aerosol either nasally or using delivery articles of the type set forth in U.S. Patent No. 4,922,901 to Brooks et al., the disclosure of which is incorporated herein in its entirety); topically (*e.g.*, in lotion form); transdermally (*e.g.*, using a transdermal patch, using technology that

is commercially available from Novartis and Alza Corporation, or by powder injection); or by buccal, sublingual or intranasal absorption. Although it is possible to administer the compounds in the form of a bulk active chemical, it is preferred to present each compound in the form of a pharmaceutical composition or
5 formulation for efficient and effective administration.

Exemplary methods for administering such compounds will be apparent to the skilled artisan. The usefulness of these formulations may depend on the particular composition used and the particular subject receiving the treatment. For example, the compositions can be administered in the form of a tablet, a hard
10 gelatin capsule or as a time release capsule. These formulations may contain a liquid carrier that may be oily, aqueous, emulsified or contain certain solvents suitable to the mode of administration.

The administration of the pharmaceutical compositions described herein can be intermittent, or at a gradual, continuous, constant or controlled rate to a
15 warm-blooded animal, (e.g., a mammal such as a mouse, rat, cat, rabbit, dog, pig, cow, or monkey); but advantageously is preferably administered to a human being. In addition, the time of day and the number of times per day that the pharmaceutical composition is administered can vary.

The appropriate dose of the compound is that amount effective to prevent
20 occurrence of the symptoms of the disorder or to treat some symptoms of the disorder from which the patient suffers. By "effective amount", "therapeutic amount" or "effective dose" is meant that amount sufficient to elicit the desired pharmacological or therapeutic effects, thus resulting in effective prevention or treatment of the disorder. Thus, when treating a CNS disorder, an effective amount
25 of compound is an amount sufficient to pass across the blood-brain barrier of the subject, to bind to relevant receptor sites in the brain of the subject, and to modulate the activity of relevant nicotinic receptor subtypes (e.g., modulate neurotransmitter secretion, thus resulting in effective prevention or treatment of the disorder). Prevention of the disorder is manifested by delaying the onset of the
30 symptoms of the disorder. Treatment of the disorder is manifested by a decrease in the symptoms associated with the disorder or an amelioration of the reoccurrence of the symptoms of the disorder.

The effective dose can vary, depending upon factors such as the condition

of the patient, the severity of the symptoms of the disorder, and the manner in which the pharmaceutical composition is administered. For human patients, the effective dose of typical compounds generally requires administering the compound in an amount sufficient to modulate disease-relevant receptors to affect neurotransmitter (e.g., dopamine) release but the amount should be insufficient to induce effects on skeletal muscles and ganglia to any significant degree. The effective dose of compounds will of course differ from patient to patient but in general includes amounts starting where CNS effects or other desired therapeutic effects occur, but below the amount where muscular and ganglionic effects are observed.

Typically, to be administered in an effective dose, compounds require administering in an amount of less than 5 mg/kg of patient weight. Often, the compounds may be administered in an amount from less than about 1 mg/kg patient weight to less than about 100 µg/kg of patient weight, and occasionally between about 10 µg/kg to less than 100 µg/kg of patient weight. The foregoing effective doses typically represent that amount administered as a single dose, or as one or more doses administered over a 24 hours period. For human patients, the effective dose of the compounds may require administering the compound in an amount of at least about 1, but not more than about 1000, and often not more than about 500 mg/ 24 hr/ patient.

Compositions useful as diagnostics can be employed, as set forth in U.S. Patent Nos. 5,853,696 to Elmalch et al. and 5,969,144 to London et al., the contents of which are hereby incorporated by reference. The compounds also can be administered in formulation compositions that incorporate other ingredients, such as those types of ingredients that are useful in formulating a diagnostic composition.

The following examples are provided to illustrate the present invention, and should not be construed as limiting thereof. In these examples, all parts and percentages are by weight, unless otherwise noted.

Biological Assays

Example 1: Radioligand Binding at CNS nAChRs

α4β2 nAChR Subtype

Preparation of membranes from rat cortex: Rats (female, Sprague-Dawley), weighing 150-250 g, were maintained on a 12 h light/dark cycle and were allowed free access to water and food supplied by PMI Nutrition International, Inc. Animals were anesthetized with 70% CO₂, and then decapitated. Brains were removed and placed on an ice-cold platform. The cerebral cortex was removed and placed in 20 volumes (weight:volume) of ice-cold preparative buffer (137 mM NaCl, 10.7 mM KCl, 5.8 mM KH₂PO₄, 8 mM Na₂HPO₄, 20 mM HEPES (free acid), 5 mM iodoacetamide, 1.6 mM EDTA, pH 7.4); PMSF, dissolved in methanol to a final concentration of 100 µM, was added and the suspension was homogenized by Polytron. The homogenate was centrifuged at 18,000 x g for 20 min at 4°C and the resulting pellet was re-suspended in 20 volumes of ice-cold water. After 60 min incubation on ice, a new pellet was collected by centrifugation at 18,000 x g for 20 min at 4°C. The final pellet was re-suspended in 10 volumes of buffer and stored at -20°C.

Preparation of membranes from SH-EP1/human α4β2 clonal cells: Cell pellets from 40 150 mm culture dishes were pooled, and homogenized by Polytron (Kinematica GmbH, Switzerland) in 20 milliliters of ice-cold preparative buffer. The homogenate was centrifuged at 48,000 g for 20 minutes at 4°C. The resulting pellet was re-suspended in 20 mL of ice-cold preparative buffer and stored at -20°C.

On the day of the assay, the frozen membranes were thawed and spun at 48,000 x g for 20 min. The supernatant was decanted and discarded. The pellet was resuspended in Dulbecco's phosphate buffered saline (PBS, Life Technologies) pH 7.4 and homogenized with the Polytron for 6 seconds. Protein concentrations were determined using a Pierce BCA Protein Assay Kit, with bovine serum albumin as the standard (Pierce Chemical Company, Rockford, IL).

Membrane preparations (approximately 50 µg for human and 200-300 µg protein for rat α4β2) were incubated in PBS (50 µL and 100 µL respectively) in the presence of competitor compound (0.01 nM to 100 µM) and 5 nM [³H]nicotine for 2-3 hours on ice. Incubation was terminated by rapid filtration on a multi-manifold tissue harvester (Brandel, Gaithersburg, MD) using GF/B filters presoaked in 0.33% polyethyleneimine (w/v) to reduce non-specific binding. Tissue was rinsed

3 times in PBS, pH 7.4. Scintillation fluid was added to filters containing the washed tissue and allowed to equilibrate. Filters were then counted to determine radioactivity bound to the membranes by liquid scintillation counting (2200CA Tri-Carb LSC, Packard Instruments, 50% efficiency or Wallac Trilux 1450 MicroBeta, 40% efficiency, Perkin Elmer).

Data were expressed as disintegrations per minute (DPMs). Within each assay, each point had 2-3 replicates. The replicates for each point were averaged and plotted against the log of the drug concentration. IC_{50} , which is the concentration of the compound that produces 50% inhibition of binding, was determined by least squares non-linear regression. K_i values were calculated using the Cheng-Prusoff equation (1973):

$$K_i = IC_{50} / (1 + N/K_d)$$

where N is the concentration of [3H]nicotine and K_d is the affinity of nicotine (3 nM, determined in a separate experiment).

15

$\alpha 7$ nAChR Subtype

Rats (female, Sprague-Dawley), weighing 150-250 g, were maintained on a 12 h light/dark cycle and were allowed free access to water and food supplied by PMI Nutrition International, Inc. Animals were anesthetized with 70% CO_2 , then decapitated. Brains were removed and placed on an ice-cold platform. The hippocampus was removed and placed in 10 volumes (weight:volume) of ice-cold preparative buffer (137 mM NaCl, 10.7 mM KCl, 5.8 mM KH_2PO_4 , 8 mM Na_2HPO_4 , 20 mM HEPES (free acid), 5 mM iodoacetamide, 1.6 mM EDTA, pH 7.4); PMSF, dissolved in methanol to a final concentration of 100 μM , was added and the tissue suspension was homogenized by Polytron. The homogenate was centrifuged at 18,000 x g for 20 min at 4°C and the resulting pellet was re-suspended in 10 volumes of ice-cold water. After 60 min incubation on ice, a new pellet was collected by centrifugation at 18,000 x g for 20 min at 4°C. The final pellet was re-suspended in 10 volumes of buffer and stored at -20°C. On the day of the assay, tissue was thawed, centrifuged at 18,000 x g for 20 min, and then re-suspended in ice-cold PBS (Dulbecco's Phosphate Buffered Saline, 138 mM NaCl,

2.67 mM KCl, 1.47 mM KH₂PO₄, 8.1 mM Na₂HPO₄, 0.9 mM CaCl₂, 0.5 mM MgCl₂, Invitrogen/Gibco, pH 7.4) to a final concentration of approximately 2 mg protein/mL. Protein was determined by the method of Lowry et al., *J. Biol. Chem.* **193**: 265 (1951), using bovine serum albumin as the standard.

5 The binding of [³H]MLA was measured using a modification of the methods of Davies et al., *Neuropharmacol.* **38**: 679 (1999). [³H]MLA (Specific Activity = 25-35 Ci/mmol) was obtained from Tocris. The binding of [³H]MLA was determined using a 2 h incubation at 21°C. Incubations were conducted in 48-well micro-titre plates and contained about 200 µg of protein per well in a final
10 incubation volume of 300 µL. The incubation buffer was PBS and the final concentration of [³H]MLA was 5 nM. The binding reaction was terminated by filtration of the protein containing bound ligand onto glass fiber filters (GF/B, Brandel) using a Brandel Tissue Harvester at room temperature. Filters were soaked in de-ionized water containing 0.33% polyethyleneimine to reduce non-
15 specific binding. Each filter was washed with PBS (3 x 1 mL) at room temperature. Non-specific binding was determined by inclusion of 50 µM non-radioactive MLA in selected wells.

 The inhibition of [³H]MLA binding by test compounds was determined by including seven different concentrations of the test compound in selected wells.
20 Each concentration was replicated in triplicate. IC₅₀ values were estimated as the concentration of compound that inhibited 50 percent of specific [³H]MLA binding. Inhibition constants (K_i values), reported in nM, were calculated from the IC₅₀ values using the method of Cheng et al., *Biochem. Pharmacol.* **22**: 3099-3108 (1973).

25 **Example 2: Determination of Dopamine Release**

 Dopamine release was measured using striatal synaptosomes obtained from rat brain, according to the procedures set forth by Rapier et al., *J. Neurochem.* **54**: 937 (1990). Rats (female, Sprague-Dawley), weighing 150-250 g, were maintained on a 12 h light/dark cycle and were allowed free access to water and food supplied
30 by PMI Nutrition International, Inc. Animals were anesthetized with 70% CO₂, then decapitated. The brains were quickly removed and the striata dissected. Striatal tissue from each of 2 rats was pooled and homogenized in ice-cold 0.32 M sucrose (5 mL) containing 5 mM HEPES, pH 7.4, using a glass/glass homogenizer.

The tissue was then centrifuged at 1,000 x g for 10 min. The pellet was discarded and the supernatant was centrifuged at 12,000 x g for 20 min. The resulting pellet was re-suspended in perfusion buffer containing monoamine oxidase inhibitors (128 mM NaCl, 1.2 mM KH₂PO₄, 2.4 mM KCl, 3.2 mM CaCl₂, 1.2 mM MgSO₄,
5 25 mM HEPES, 1 mM ascorbic acid, 0.02 mM pargyline HCl and 10 mM glucose, pH 7.4) and centrifuged for 15 min at 25,000 x g. The final pellet was resuspended in perfusion buffer (1.4 mL) for immediate use.

The synaptosomal suspension was incubated for 10 min at 37°C to restore metabolic activity. [³H]Dopamine ([³H]DA, specific activity = 28.0 Ci/mmol,
10 NEN Research Products) was added at a final concentration of 0.1 μM and the suspension was incubated at 37°C for another 10 min. Aliquots of tissue (50 μL) and perfusion buffer (100 μL) were loaded into the suprafusion chambers of a Brandel Suprafusion System (series 2500, Gaithersburg, MD). Perfusion buffer (room temperature) was pumped into the chambers at a rate of 1.5 mL/min for a
15 wash period of 16 min. Test compound (10 μM) or nicotine (10 μM) was then applied in the perfusion stream for 48 sec. Fractions (24 sec each) were continuously collected from each chamber throughout the experiment to capture basal release and agonist-induced peak release and to re-establish the baseline after the agonist application. The perfusate was collected directly into scintillation vials,
20 to which scintillation fluid was added. [³H]DA released was quantified by scintillation counting. For each chamber, the integrated area of the peak was normalized to its baseline.

Release was expressed as a percentage of release obtained with an equal concentration of L-nicotine. Within each assay, each test compound was replicated
25 using 2-3 chambers; replicates were averaged. When appropriate, dose-response curves of test compound were determined. The maximal activation for individual compounds (E_{max}) was determined as a percentage of the maximal activation induced by L-nicotine. The compound concentration resulting in half maximal activation (EC₅₀) of specific ion flux was also defined.

30

Example 3: Selectivity vs. Peripheral nAChRs

Interaction at the Human Muscle nAChR Subtype

Activation of muscle-type nAChRs was established on the human clonal

line TE671/RD, which is derived from an embryonal rhabdomyosarcoma (Stratton et al., *Carcinogen* 10: 899 (1989)). These cells express receptors that have pharmacological (Lukas, *J. Pharmacol. Exp. Ther.* **251**: 175 (1989)), electrophysiological (Oswald et al., *Neurosci. Lett.* **96**: 207 (1989)), and molecular
5 biological profiles (Luther et al., *J. Neurosci.* **9**: 1082 (1989)) similar to the muscle-type nAChR.

TE671/RD cells were maintained in proliferative growth phase according to routine protocols (Bencherif et al., *Mol. Cell. Neurosci.* **2**: 52 (1991) and Bencherif et al., *J. Pharmacol. Exp. Ther.* **257**: 946 (1991)). Cells were cultured in
10 Dulbecco's modified Eagle's medium (Gibco/BRL) with 10% horse serum (Gibco/BRL), 5% fetal bovine serum (HyClone, Logan UT), 1mM sodium pyruvate, 4 mM L-Glutamine, and 50,000 units penicillin-streptomycin (Irvine Scientific). When cells were 80% confluent, they were plated to 12 well polystyrene plates (Costar). Experiments were conducted when the cells reached
15 100% confluency.

Nicotinic acetylcholine receptor (nAChR) function was assayed using $^{86}\text{Rb}^+$ efflux according to the method described by Lukas et al., *Anal. Biochem.* **175**: 212 (1988). On the day of the experiment, growth media was gently removed from the well and growth media containing ^{86}Rb chloride ($10^6 \mu\text{Ci/mL}$) was added to
20 each well. Cells were incubated at 37°C for a minimum of 3 h. After the loading period, excess $^{86}\text{Rb}^+$ was removed and the cells were washed twice with label-free Dulbecco's phosphate buffered saline (138 mM NaCl, 2.67 mM KCl, 1.47 mM KH_2PO_4 , 8.1 mM Na_2HPO_4 , 0.9 mM CaCl_2 , 0.5 mM MgCl_2 , Invitrogen/Gibco, pH. 7.4), taking care not to disturb the cells. Next, cells were exposed to either 100 μM
25 of test compound, 100 μM of L-nicotine (Acros Organics) or buffer alone for 4 min. Following the exposure period, the supernatant containing the released $^{86}\text{Rb}^+$ was removed and transferred to scintillation vials. Scintillation fluid was added and released radioactivity was measured by liquid scintillation counting.

Within each assay, each point had 2 replicates, which were averaged. The
30 amount of $^{86}\text{Rb}^+$ release was compared to both a positive control (100 μM L-nicotine) and a negative control (buffer alone) to determine the percent release relative to that of L-nicotine.

When appropriate, dose-response curves of test compound were

determined. The maximal activation for individual compounds (E_{max}) was determined as a percentage of the maximal activation induced by L-nicotine. The compound concentration resulting in half maximal activation (EC_{50}) of specific ion flux was also determined.

5 **Interaction at the Rat Ganglionic nAChR Subtype**

Activation of rat ganglion nAChRs was established on the pheochromocytoma clonal line PC12, which is a continuous clonal cell line of neural crest origin, derived from a tumor of the rat adrenal medulla. These cells express ganglion-like nAChRs (see Whiting et al., *Nature* **327**: 515 (1987); Lukas,
10 *J. Pharmacol. Exp. Ther.* **251**: 175 (1989); Whiting et al., *Mol. Brain Res.* **10**: 61 (1990)).

Rat PC12 cells were maintained in proliferative growth phase according to routine protocols (Bencherif et al., *Mol. Cell. Neurosci.* **2**: 52 (1991) and Bencherif et al., *J. Pharmacol. Exp. Ther.* **257**: 946 (1991)). Cells were cultured in
15 Dulbecco's modified Eagle's medium (Gibco/BRL) with 10% horse serum (Gibco/BRL), 5% fetal bovine serum (HyClone, Logan UT), 1mM sodium pyruvate, 4 mM L-Glutamine, and 50,000 units penicillin-streptomycin (Irvine Scientific). When cells were 80% confluent, they were plated to 12 well Nunc plates (Nunc) and coated with 0.03% poly-L-lysine (Sigma, dissolved in 100
20 mM boric acid). Experiments were conducted when the cells reached 80% confluency.

Nicotinic acetylcholine receptor (nAChR) function was assayed using $^{86}\text{Rb}^+$ efflux according to a method described by Lukas et al., *Anal. Biochem.* **175**: 212 (1988). On the day of the experiment, growth media was gently removed from the
25 well and growth media containing $^{86}\text{Rb}^+$ chloride ($10^6 \mu\text{Ci/mL}$) was added to each well. Cells were incubated at 37°C for a minimum of 3 h. After the loading period, excess $^{86}\text{Rb}^+$ was removed and the cells were washed twice with label-free Dulbecco's phosphate buffered saline (138 mM NaCl, 2.67 mM KCl, 1.47 mM KH_2PO_4 , 8.1 mM Na_2HPO_4 , 0.9 mM CaCl_2 , 0.5 mM MgCl_2 , Invitrogen/Gibco, pH.
30 7.4), taking care not to disturb the cells. Next, cells were exposed to either 100 μM of test compound, 100 μM of nicotine or buffer alone for 4 min. Following the exposure period, the supernatant containing the released $^{86}\text{Rb}^+$ was removed and transferred to scintillation vials. Scintillation fluid was added and released

radioactivity was measured by liquid scintillation counting

Within each assay, each point had 2 replicates, which were averaged. The amount of $^{86}\text{Rb}^+$ release was compared to both a positive control (100 μM nicotine) and a negative control (buffer alone) to determine the percent release relative to that of L-nicotine.

When appropriate, dose-response curves of test compound were determined. The maximal activation for individual compounds (E_{max}) was determined as a percentage of the maximal activation induced by L-nicotine. The compound concentration resulting in half maximal activation (EC_{50}) of specific ion flux was also determined.

Interaction at the Human Ganglionic nAChR Subtype

The cell line SH-SY5Y is a continuous line derived by sequential subcloning of the parental cell line, SK-N-SH, which was originally obtained from a human peripheral neuroblastoma. SH-SY5Y cells express a ganglion-like nAChR (Lukas et al., *Mol. Cell. Neurosci.* **4**: 1 (1993)).

Human SH-SY5Y cells were maintained in proliferative growth phase according to routine protocols (Bencherif et al., *Mol. Cell. Neurosci.* **2**: 52 (1991) and Bencherif et al., *J. Pharmacol. Exp. Ther.* **257**: 946 (1991)). Cells were cultured in Dulbecco's modified Eagle's medium (Gibco/BRL) with 10% horse serum (Gibco/BRL), 5% fetal bovine serum (HyClone, Logan UT), 1mM sodium pyruvate, 4 mM L-Glutamine, and 50,000 units penicillin-streptomycin (Irvine Scientific). When cells were 80% confluent, they were plated to 12 well polystyrene plates (Costar). Experiments were conducted when the cells reached 100% confluency.

Nicotinic acetylcholine receptor (nAChR) function was assayed using $^{86}\text{Rb}^+$ efflux according to a method described by Lukas et al., *Anal. Biochem.* **175**: 212 (1988). On the day of the experiment, growth media was gently removed from the well and growth media containing $^{86}\text{Rubidium}$ chloride ($10^6 \mu\text{Ci/mL}$) was added to each well. Cells were incubated at 37°C for a minimum of 3 h. After the loading period, excess $^{86}\text{Rb}^+$ was removed and the cells were washed twice with label-free Dulbecco's phosphate buffered saline (138 mM NaCl, 2.67 mM KCl, 1.47 mM KH_2PO_4 , 8.1 mM Na_2HPO_4 , 0.9 mM CaCl_2 , 0.5 mM MgCl_2 , Invitrogen/Gibco, pH 7.4), taking care not to disturb the cells. Next, cells were exposed to either 100 μM

of test compound, 100 μM of nicotine, or buffer alone for 4 min. Following the exposure period, the supernatant containing the released $^{86}\text{Rb}^+$ was removed and transferred to scintillation vials. Scintillation fluid was added and released radioactivity was measured by liquid scintillation counting

5 Within each assay, each point had 2 replicates, which were averaged. The amount of $^{86}\text{Rb}^+$ release was compared to both a positive control (100 μM nicotine) and a negative control (buffer alone) to determine the percent release relative to that of L-nicotine.

 When appropriate, dose-response curves of test compound were
10 determined. The maximal activation for individual compounds (E_{max}) was determined as a percentage of the maximal activation induced by L-nicotine. The compound concentration resulting in half maximal activation (EC_{50}) of specific ion flux was also defined.

15 **Example 4: Novel Object Recognition (NOR) Task**

 The novel object recognition (NOR) task was performed in accord with the description of Ennaceur and Delacour *Behav. Brain Res.* **100**: 85-92 (1988).

Synthetic Examples

20 **Example 5: Synthesis of N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane**

 Example 5 relates to the synthesis of N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane, which was prepared as described in U.S. applications 2004/0186107 to Schrimpf et al. and 2005/0101602 to Basha et al., according to
25 the following techniques:

5-Benzyltetrahydropyrrolo[3,4-c]pyrrole-1,3-dione (or 7-benzyl-3,7-diazabicyclo[3.3.0]octan-2,4-dione)

 Trifluoroacetic acid (TFA, 0.50 mL, 6.5 mmol) was added to a cold (0°C) solution of maleimide (6.27 g, 0.0646 mol) in dichloromethane (150 mL) under
30 nitrogen. A solution of N-(methoxymethyl)-N-(trimethylsilylmethyl)benzylamine (20 g, 0.084 mol) in dichloromethane (100 mL) was added drop-wise over 45 min. After the addition was complete, the mixture was warmed slowly to ambient temperature and stirred for 16 h. The mixture was concentrated and the resulting

residue was dissolved in dichloromethane (200 mL) and washed with saturated aqueous sodium bicarbonate (2 x 50 mL). The aqueous layer was separated and extracted with dichloromethane (2 x 75 mL). The combined dichloromethane extracts were washed with brine (50 mL), dried over anhydrous magnesium sulfate, filtered and concentrated to give 12.5 g (83.9% yield) of a light yellow, waxy solid (MS m/z 231 (M+H)).

2-Benzyl-octahydropyrrolo[3,4-c]pyrrole (or 3-benzyl-3,7-diazabicyclo[3.3.0]octane)

The crude 5-benzyltetrahydropyrrolo[3,4-c]pyrrole-1,3-dione (4.9 g, 0.021 mol) was dissolved in cold (0°C) dry tetrahydrofuran (THF) (50 mL) under nitrogen, and lithium aluminum hydride (63 mL of 1 M in THF, 0.063 mol) was added drop-wise over 30 min to the continuously cooled solution. The resulting mixture was stirred at ambient temperature for 30 min and then warmed to reflux for 4 h. The mixture was then cooled to 0°C and quenched by the slow addition of excess solid sodium sulfate decahydrate. The mixture was warmed to ambient temperature and stirred for 16 h. The solids were filtered and the residue was washed with ethyl acetate (3 x 100 mL). The combined filtrates were concentrated to give 4.2 g (99% yield) of a waxy solid (MS m/z 203 (M+H)).

5-Benzylhexahydropyrrolo[3,4-c]pyrrole-2-carboxylic acid tert-butyl ester (or N-benzyl-N'-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane)

The crude 2-benzyl-octahydropyrrolo[3,4-c]pyrrole (4.2 g, 0.021 mol) was dissolved in THF (50 mL). Di-t-butyl dicarbonate (5.5 g, 0.025 mol) and aqueous saturated NaHCO₃ (10 mL) were added, and the mixture was stirred at ambient temperature overnight. The reaction was quenched with water (10 mL), and ethyl acetate (30 mL) was added. The aqueous layer was extracted with ethyl acetate (2 x 20 mL), and the combined organic extracts were dried over anhydrous sodium sulfate and concentrated. Purification via silica gel column chromatography (1:1 hexanes/ ethyl acetate) gave 5.07 g (79.8% yield) of the title compound (MS m/z 303 (M+H)).

Hexahydropyrrolo[3,4-c]pyrrole-2-carboxylic acid tert-butyl ester (or N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.0]octane)

The 5-benzylhexahydropyrrolo[3,4-c]pyrrole-2-carboxylic acid tert-butyl

ester (5.07 g, 0.0168 mol) was dissolved in methanol (50 mL) and 20% Pd(OH)₂/C (wet) (~2 g) was added under a nitrogen atmosphere. The resulting mixture was warmed (45-50°C) and shaken for 2 h under 40 psi of hydrogen. The mixture was filtered and concentrated to give 3.49 g (97.7% yield) of the title compound (MS m/z 213 (M+H)).

Example 6: Synthesis of N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane

Example 6 relates to the synthesis of furan-2-yl(hexahydropyrrolo[3,4-c]pyrrol-2-yl)methanone (or N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane), which was prepared according to the following techniques, illustrative of the coupling reaction used to make heteroaromatic amides of 3,7-diazabicyclo[3.3.0]octane:

Furan-2-yl(hexahydropyrrolo[3,4-c]pyrrol-2-yl)methanone trifluoroacetate (or N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane trifluoroacetate)

Furan-2-carboxylic acid (0.037 g, 0.33 mmol) and triethylamine (0.125 mL, 0.99 mmol) were combined in dry dichloromethane (1 mL), and O-(benzotriazol-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate (HBTU; 0.125 g, 0.33 mmol) was added. A solution of hexahydropyrrolo[3,4-c]pyrrole-2-carboxylic acid tert-butyl ester (0.064 g, 0.30 mmol) in dichloromethane (0.5 mL) was added, and the mixture was stirred at ambient temperature overnight. The mixture was shaken with 10% aqueous sodium hydroxide, and the organic layer was separated. The aqueous layer was extracted with chloroform (2 x 2 mL). The combined organic extracts were washed with water (2 x 1 mL) and concentrated. The resulting residue was dissolved in dimethylformamide (DMF) (0.3 mL) and purified by HPLC (acetonitrile/water gradient). Fractions containing the desired material were pooled and concentrated, leaving the tert-butoxycarbonyl-protected product. This material was dissolved in a mixture of trifluoroacetic acid (0.5 mL) and dichloromethane (0.5 mL), and the mixture was stirred at ambient temperature for 1h. The volatiles were removed by rotary evaporation, followed by high vacuum treatment, to give 77 mg of an oil (80% yield) (¹H NMR (d₄-methanol, 300 MHz) 3.20 (m, 2H), 3.47-4.2 (m, 8H), 6.60 (t, 1H), 7.18 (d, 1H), 7.72 (d, 1H); MS m/z

207 (M+H)).

Example 7: Synthesis of N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane trifluoroacetate

5 Example 7 relates to the synthesis of 5-chlorofuran-2-yl(hexahydropyrrolo[3,4-c]pyrrol-2-yl)methanone trifluoroacetate (or N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane trifluoroacetate), which was prepared according to the following techniques, illustrative of the coupling reaction used to make heteroaromatic amides of 3,7-diazabicyclo[3.3.0]octane:

10 **5-Chlorofuran-2-carboxylic acid**

Aqueous sodium hydroxide (80 mL of 10%) was added to a solution of silver nitrate (8.0 g, 47 mmol) in water (20 mL). This suspension was stirred and slowly treated with 30% aqueous ammonium hydroxide until it became clear. A solution of 5-chlorofuran-2-carboxaldehyde (3.0 g, 23 mmol) (Aldrich Chemical)
15 in methanol (5 mL) was added, and the resulting mixture was stirred at ambient temperature for 30 min. The reaction mixture was filtered, and the filtrate was washed with ether (100 mL). The aqueous filtrate was then made acidic (~pH 3) by the addition of cold 20% sulfuric acid. The resulting mixture was extracted with ethyl acetate (3 x 100 mL). The extracts were washed with saturated aqueous
20 sodium chloride solution (100 mL), dried (anhydrous sodium sulfate) and concentrated under vacuum to give 3.2 g (95% yield) of white solid (mp 178-179°C). This reaction was easily scalable and was run multiple times at >10 g scale.

25 **N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane trifluoroacetate**

Oxalyl chloride (12.2 g, 95.8 mmol) containing a drop of DMF was added drop-wise to an ice-cooled solution of 5-chlorofuran-2-carboxylic acid (6.25 g, 47.9 mmol) in 200 mL of dichloromethane. After complete addition, the ice bath was removed and the reaction was warmed to ambient temperature over a 1 h
30 period. The volatiles were then removed under vacuum, and the residue was dissolved in THF (50 mL). This solution of the acid chloride was then added to an stirred, ice-cooled solution of hexahydropyrrolo[3,4-c]pyrrole-2-carboxylic acid tert-butyl ester (10.2 g, 47.9 mmol) and diisopropylethylamine (25 g, ~4

equivalents) in THF (200 mL). This mixture was stirred at ambient temperature for 16 h. The volatiles were then removed under vacuum, and the residue was partitioned between water (100 mL) and ether (300 mL). The ether layer and two ether extracts (100 mL) of the aqueous layer were concentrated on the rotary evaporator. The residue was column chromatographed on silica gel, eluting with a 0-60% ethyl acetate in hexane gradient. Concentration of selected fractions gave 13.9 g (85.3% yield) of pale yellow syrup. A portion of this material (12.9 g, 37.9 mmol) was dissolved in a mixture of dichloromethane and trifluoroacetic acid (100 mL each). This mixture was stirred at ambient temperature for 2 h and then concentrated under vacuum. The residue was partitioned between chloroform (200 mL) and 50% aqueous potassium carbonate (200 mL), and the aqueous layer was extracted with chloroform (3 x 200 mL). The combined chloroform layers were dried over anhydrous sodium sulfate and concentrated under vacuum, leaving 8.66 g (95% yield) of pale yellow solid (¹H NMR (d₄-methanol, 300 MHz) 3.15-3.35 (m, 4H), 3.50-4.20 (m, 6H), 6.51 (d, 1H), 7.17 (d, 1H); MS m/z 241 (M+H)).

Example 8: Synthesis of 3,7-diazabicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester

Example 8 relates to the synthesis of 3,7-diazabicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester (or N-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane), which was prepared according to the following techniques:

7-Benzyl-3,7-diazabicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester (or N-benzyl-N'-(tert-butoxycarbonyl)-3,7-diazabicyclo[3.3.1]nonane)

7-Benzyl-3,7-diazabicyclo [3.3.1] nonane-3-carboxylic acid tert-butyl ester was prepared according to procedures set forth by Stead et al. in *Org. Lett.* **7(20)**: 4459 (2005).

3,7-Diazabicyclo[3.3.1]-3-carboxylic acid tert-butyl ester

7-Benzyl-3,7-diazabicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester (0.49 g, 1.6 mmol) was dissolved in methanol (20 mL) and 20% Pd(OH)₂/C (wet) (~ 2 g) was added under a nitrogen atmosphere. This mixture was warmed to about 50°C and shaken for 2 h under 55 psi of hydrogen. The resulting mixture was

filtered and concentrated to give 0.32 g (94% yield) of the title compound (MS m/z 227 (M+H)).

Example 9: Synthesis of N-(furan-2-ylcarbonyl)-3,7-

5 diazabicyclo[3.3.1]nonane trifluoroacetate

Example 9 relates to the synthesis of (3,7-diazabicyclo[3.3.1]non-3-yl)-furan-2-ylmethanone trifluoroacetate (or N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane trifluoroacetate), which was prepared according to the following techniques, illustrative of the coupling reaction used to make
10 heteroaromatic amides of 3,7-diazabicyclo[3.3.1]nonane:

**3,7-Diazabicyclo[3.3.1]non-3-yl)-furan-2-yl methanone trifluoroacetate
(or N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane
trifluoroacetate)**

Furan-2-carboxylic acid (0.032 g, 0.29 mmol) was combined with
15 triethylamine (0.870 mmol, 0.121 mL) in dry dichloromethane (1 mL) and HBTU (0.11 g, 0.29 mmol) was added. A solution of 3,7-diazabicyclo[3.3.1]-3-carboxylic acid tert-butyl ester (0.059 g, 0.26 mmol) in dichloromethane (0.5 mL) was added, and the mixture was stirred at ambient temperature overnight. The mixture was treated with 10% aqueous sodium hydroxide and extracted with chloroform (2 x 2
20 mL). The combined organic extracts were washed with water (2 x 1 mL), and concentrated. The resulting residue was dissolved in DMF (0.3 mL) and purified by HPLC (acetonitrile/water gradient). Fractions containing the desired material were pooled and concentrated, leaving the tert-butoxycarbonyl-protected product. This material was dissolved in a mixture of trifluoroacetic acid (0.5 mL) and
25 dichloromethane (0.5 mL), and the mixture was stirred at ambient temperature for 1h. The volatiles were removed by rotary evaporation, followed by high vacuum treatment, to give 36 mg of an oil (41% yield) (¹H NMR (d₄-methanol, 300 MHz) 2.10 (bs, 2H), 2.35 (bs, 2H), 3.30-3.45 (m, 4H), 3.55 (m, 2H), 6.65 (m, 1H), 7.15 (d, 1H), and 7.75 (d, 1H). MS m/z 221 (M+H)).








Example 10: Synthesis of N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane trifluoroacetate









Example 10 relates to the synthesis of (3,7-diazabicyclo[3.3.1]non-3-yl)-5-chlorofuran-2-ylmethanone trifluoroacetate (or N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane trifluoroacetate), which was prepared by a process similar to that described in Example 9, according to the following techniques:









5-Chlorofuran-2-carboxylic acid (0.96 g, 6.5 mmol) was combined with triethylamine (21 mmol, 2.9 mL) in dry dichloromethane (10 mL), and HBTU (2.47 g, 65.1 mmol) was added. A solution of 3,7-diazabicyclo[3.3.1]-3-carboxylic acid tert-butyl ester (1.5 g, 66 mmol) in dichloromethane (5 mL) was added, and the mixture was stirred at ambient temperature overnight. The mixture was treated with 10% aqueous sodium hydroxide and extracted with chloroform (2 x 20 mL). The combined organic extracts were washed with water (2 x 10 mL), and concentrated. The resulting residue was purified by column chromatography on silica gel, eluting with an ethyl acetate in hexane gradient, to give the tert-butoxycarbonyl-protected product, as a viscous oil. This material was dissolved in a mixture of trifluoroacetic acid (20 mL) and dichloromethane (20 mL), and the mixture was stirred at ambient temperature for 1h. The volatiles were removed by rotary evaporation, followed by high vacuum treatment, to give 1.38 g (57.5% yield) of viscous yellow oil (^1H NMR (d_4 -methanol, 300 MHz) 2.00 (bs, 2H), 2.15 (bs, 2H), 3.15-3.35 (m, 6H), 4.25 (m, 2H), 6.53 (d, 1H) and 7.10 (d, 1H). MS m/z 255 (M+H)).







Example 11: Tabular Spectral and Receptor Binding Data








The above illustrated amide coupling procedures were utilized to make the compounds shown in Tables 1 and 2. In some cases, compounds were synthesized on a scale sufficient to obtain nuclear magnetic resonance (NMR) data. In other cases, compounds were produced on a smaller scale in various kinds of parallel synthesis apparatus and were (structurally) characterized by LCMS only.







Table 1					
Structure	Rat $\alpha 4\beta 2$ Ki	Human $\alpha 4\beta 2$ Ki	$\alpha 7$ Ki	MS: m/z (M+H)	^1H NMR: CD_3OD , 300 MHz
	20	33	13000	207	δ 7.72 (d, 1H), 7.18 (d, 1H), 6.60 (t, 1H), 3.47-4.2 (m, 8H), 3.2 (m, 2H)
	26	160	ND; failed HTS	218	δ 9.00 (d, 2H), 8.20 (m, 2H), 3.7-4.0 (m, 3H), 3.47-3.7 (m, 5H), 3.2 (m, 2H)
	54	73	12000	287	
	53	220	3100	207	
	35	31	2900	252	
	19	19	3500	238	
	1.5	1.5	12000	287	

	33	28	91000	241	δ 7.17 (d, J = 0.61 Hz, 1H), 6.51 (d, J = 0.73 Hz, 1H), 4.20-3.50 (m, 6H), 3.35-3.15 (m, 4H)
	44	54	25000	221	
	230	43	3200	307	
	130	28	23000	221	
	17	44	ND; failed HTS	288	
	29	18	ND; failed HTS	287	δ 7.80 (s, 1H), 7.2 (s, 1H), 3.5-4.2 (m, 8H), 3.20 (m, 2H)
	140	40	ND; failed HTS	267	
	54	24	ND; failed HTS	319	δ 7.50 (s, 1H), 7.17 (m, 2H), 6.78 (m, 1H), 6.60 (s, 1H), 3.95-3.76 (m, 4H), 3.70-3.58 (m, 2H), 3.32-3.08 (m, 4H)

	260	86	12000	283	
	110	95	ND; failed HTS	273	
	150	84	ND; failed HTS	222	
	15	9.2	210000	222	
	24	14	56000	222	
	32	31	ND; failed HTS	222	
	61	29	160000	221	δ 7.38 (d, J = 1.8 Hz, 1 H), 6.55 (d, J = 1.8 Hz, 1 H), 3.79 (m, 2 H), 3.45 (m, 2 H), 3.06 (m, 2 H), 2.89 (m, 2 H), 2.67 (m, 2 H), 2.37 (s, 3H)
	53	63	ND; failed HTS	250	

	353	100	ND; failed HTS	237	δ 7.55 (s, 1H), 6.7-6.6 (m, 1H), 3.9 (s, 3H), 3.85 (m, 2H), 3.75 (m, 2H), 3.65 (m, 2H), 3.15 (m, 4H)
	1000	45	ND; failed HTS	289	δ 6.48 (s, 1H), 3.5-4.0 (m, 6H), 3.3-3.1 (m, 4H), 2.35 (s, 3H)
	320	36	ND; failed HTS	275	δ 8.25 (s, 1H), 7.35 (s, 1H), 4.2-3.6 (m, 4H), 3.55 (dd, 2H), 3.3-3.2 (m, 4H)
	31	28	ND; failed HTS	286	δ 8.15 (s, 1H), 7.35 (s, 1H), 4.2-3.6 (m, 4H), 3.5 (dd, 2H), 3.4-3.2 (m, 4H)
	20	22	ND; failed HTS	299/301	
	140	51	ND; failed HTS	233	δ 7.58 (dd, J = 2, 0.5 Hz, 1H), 7.26 (dd, J = 18, 11 Hz, 1H), 6.86 (dd, J = 2, 0.5 Hz, 1H), 5.70 (dd, J = 18, 1.5 Hz, 1H), 5.32 (dd, J = 11, 1.5 Hz, 1H), 4.20-3.62 (m, 4H), 3.62-3.58 (m, 2H), 3.25-3.18 (m, 4H)

	43	21	ND; failed HTS	287	δ 7.81 (s, 1H), 7.08 (d, J = 1 Hz, 1H), 4.18-3.62 (m, 4H), 3.59 (m, 2H), 3.30-3.20 (m, 4H), 2.40 (t, J = 7 Hz, 2H), 1.60-1.44 (m, 4H), 0.95 (t, J = 7 Hz, 3H)
	2.8	2.4	ND; failed HTS	241	
	8.3	16	ND; failed HTS	225	
	20	4.9	ND; failed HTS	366	
	35	27	ND; failed HTS	225	δ 7.60 (d, J = 2 Hz, 1H), 7.59 (d, J = 2 Hz, 1H), 4.18-3.65 (m, 4H), 3.60 (m, 2H), 3.22 (m, 4H)
	25	14	ND; failed HTS	232	δ 7.85 (d, J = 2 Hz, 1H), 6.95 (d, J = 2 Hz, 1H), 4.25-3.63 (m, 4H), 3.60 (m, 2H), 3.24 (m, 4H)
	21	9.7	ND; failed HTS	241	δ 7.83 (d, J = 1 Hz, 1H), 7.16 (d, J = 0.7 Hz, 1H), 4.20-3.62 (m, 4H), 3.60 (m,

					2H), 3.28-3.19 (m, 4H)
	89	90	ND; failed HTS	208	δ 8.79 (d, J = 1.7 Hz, 1H), 6.80 (d, J = 1.7 Hz, 1H), 4.12 (dd, J = 12, 7 Hz, 1H), 3.99 (dd, J = 12, 3 Hz, 1H), 3.90 (dd, J = 13, 8 Hz, 1H), 3.75 (dd, J = 13, 4 Hz, 1H), 3.58 (m, 2H), 3.24 (m, 4H)
	36	21	ND; failed HTS	225	δ 7.80 (d, J = 1 Hz, 1H), 7.78 (d, J = 1 Hz, 1H), 4.21-3.62 (m, 4H), 3.58 (m, 2H), 3.21 (m, 4H)
	140	48	ND; failed HTS	231	
	50	40	ND; failed HTS	232	
	65	14	ND; failed HTS	239	δ 6.24 (s, 1H), 4.17-3.65 (m, 4H), 3.57 (m, 2H), 3.19 (m, 4H), 2.34 (s, 3H)
	51	130	ND; failed HTS	257	





































	43	69	ND; failed HTS	208.3	δ 8.52 (d, J = 2 Hz, 1H), 6.96 (d, J = 2 Hz, 1H), 4.15 (dd, J = 8, 12 Hz, 1H), 3.91 (m, 2H), 3.72 (dd, J = 4, 14 Hz, 1H), 3.60 (m, 2H), 3.33 - 3.23 (m, 4H).
---	----	----	----------------------	-------	---


Table 2					
Structure	Rat $\alpha 4\beta 2$ Ki	Human $\alpha 4\beta 2$ Ki	$\alpha 7$ Ki	MS: m/z (M+H)	^1H NMR: CD_3OD , 300 MHz
	31	9.9	15000	221	δ 7.75 (d, 1H), 7.15 (d, 1H), 6.65 (m, 1H), 4.55 (m, 2H), 3.55 (m, 2H), 3.3-3.45 (m, 4H), 2.35 (bs, 2H), 2.1 (bs, 2H)
	110	18	1700	232	
	79	17	ND; failed HTS	301	
	41	51	ND; failed HTS	301	

	100	42	ND; failed HTS	235	
	15	7.3	120000	255	δ 7.10 (d, 1H), 6.53 (d, 1H), 4.25 (m, 2H), 3.15-3.35 (m, 6H), 2.15 (bs, 2H), 2.0 (bs, 2H)
	33	180	ND; failed HTS	252	
	5.9	18	ND; failed HTS	302	
	87	210	ND; failed HTS	278	δ 7.18 (d, 1H), 6.36 (d, 1H), 4.56 (m, 2H), 3.58 (m, 2H), 3.40-3.24 (m, 4H), 2.30 (m, 2H), 2.18 (s, 3H), 2.05 (m, 2H)
	68	96	ND; failed HTS	236	
	20	64	ND; failed HTS	236	
	9.5	21	ND; failed HTS	236	

	29	47	ND; failed HTS	236	
	25	83	ND; failed HTS	262	
	96	29	ND; failed HTS	255	
	6.4	4.6	ND; failed HTS	239	
	86	62	ND; failed HTS	300	δ 8.19 (s, 1H), 7.3 (s, 1H), 4.45 (d, 2H), 3.6-3.2 (m, 6H), 2.27 (bs, 2H), 2.1-2.0 (m, 2H)
	6.3	23	ND; failed HTS	333	δ 8.22 (d, J = 1 Hz, 1H), 7.50 (d, J = 1 Hz, 1H), 7.27 (m, 2H), 6.90 (m, 1H), 4.55 (d, J = 13 Hz, 2H), 3.58 (m, 2H), 3.42-3.30 (m, 4H), 2.31 (bs, 2H), 2.06 (m, 2H)
	12	35	ND; failed HTS	287	

	4	1.6	ND; failed HTS	380	
	110	57	ND; failed HTS	239	δ 7.63 (dd, $J = 4, 2$ Hz, 1H), 6.62 (dd, $J = 2, 1$ Hz, 1H), 4.37 (d, $J = 14$ Hz, 2H), 3.50 (m, 2H), 3.34 (m, 4H), 2.28 (bs, 2H), 2.06 (m, 2H)
	110	31	ND; failed HTS	246	
	13	5.2	ND; failed HTS	255	
	77	10	ND; failed HTS	255	
	7.5	5.2	ND; failed HTS	313/315	(CDCl ₃) δ 6.95 (s, 1H), 4.60 (d, 2H), 3.55 (d, 2H), 3.38 (m, 2H), 3.25 (d, 2H), 2.35 (s, 3H), 2.25 (m, 2H), 2.05 (m, 2H)
	560	28	ND; failed HTS	251	

	28	26	ND; failed HTS	222	
	28	14	ND; failed HTS	239	
	220	47	ND; failed HTS	245	
	35	8.7	ND; failed HTS	246	
	500	62	ND; failed HTS	253/505	
	370	85	ND; failed HTS	222	δ 8.09 (d, 1H), 7.40 (d, 1H), 3.65-3.23 (m, 8H), 2.30 (m, 2H), 2.05 (m, 2H)
	27	23	ND; failed HTS	246	
	12	1.7	ND; failed HTS	301	
	65	31	ND; failed HTS	236	

	52	30	ND; failed HTS	222	δ 8.53 (d, 1H), 6.90 (d, 1H), 4.60-4.20 (m, 2H), 3.58-3.25 (m, 6H), 2.30 (m, 2H), 2.05 (m, 2H)
---	----	----	----------------------	-----	---

Summary of Biological Data


Compounds of Tables 1 and 2, representative of the present invention, exhibited inhibition constants (K_i values) at the rat and human $\alpha 4\beta 2$ subtypes in the ranges of 1 nM to 1000 nM and 1 nM to 220 nM respectively, indicating high
5 affinity for the $\alpha 4\beta 2$ subtype. K_i values at the $\alpha 7$ subtype vary within the range of 1700 nM to 210,000 nM (in many cases the compounds did not bind sufficiently in high through-put screening at the $\alpha 7$ subtype to warrant K_i determination). These same compounds exhibited relatively little functional activity at either the human muscle (1-25% of the maximal response to nicotine) or human ganglion (1-20% of
10 the maximal response to nicotine) subtypes.

Certain exemplified compounds were assessed in the NOR task. Thus, both N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane (Figure 1) and N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane (Figure 2) were active in
15 OR in rats, at 0.1 mg/kg and 0.3 mg/kg respectively. This provides evidence of the efficacy (and potency) of the compounds of the present invention in treating cognitive deficits, attentional disorders and dementias, and the potential of these compounds for human therapy.

The foregoing is illustrative of the present invention and is not to be construed as limiting thereof. The invention is defined by the following claims,
20 with equivalents of the claims to be included therein. All patents and publications referred to herein are incorporated by reference in their entirety, for all purposes.

Claims

1. Compounds represented as Formula I:



Formula I

wherein n has the value of 0 or 1, and Cy is a heteroaryl group chosen from the group of 2-furanyl, 3-furanyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 1,3,4-oxadiazol-2-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 1,3,4-thiadiazol-2-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl and 4-pyridinyl, which heteroaryl groups are optionally substituted with up to three non-hydrogen substituents independently selected from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₂₋₆ alkenyl, substituted C₂₋₆ alkenyl, C₂₋₆ alkynyl, substituted C₂₋₆ alkynyl, C₃₋₈ heterocyclyl, substituted C₃₋₈ heterocyclyl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₅₋₁₀ aryl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ aryl, substituted C₅₋₁₀ heteroaryl, C₁₋₆ alkyl-C₅₋₁₀ aryl, C₁₋₆ alkyl-C₅₋₁₀ heteroaryl, substituted C₁₋₆ alkyl-C₅₋₁₀ aryl, substituted C₁₋₆ alkyl-C₅₋₁₀ heteroaryl, C₅₋₁₀ aryl-C₁₋₆ alkyl, C₅₋₁₀ heteroaryl-C₁₋₆ alkyl, substituted C₅₋₁₀ aryl-C₁₋₆ alkyl, substituted C₅₋₁₀ heteroaryl-C₁₋₆ alkyl, halo, -OR', -NR'R'', -CF₃, -CN, -NO₂, -C₂R', -SR', -N₃, -C(=O)NR'R'', -NR'C(=O)R'', -C(=O)R', -C(=O)OR', -OC(=O)R', -OC(=O)NR'R'', -NR'C(=O)OR'', -SO₂R', -SO₂NR'R'', and -NR'SO₂R'', where R' and R'' are independently selected from hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ heterocyclyl, C₅₋₁₀ aryl, C₅₋₁₀ heteroaryl or C₅₋₁₀ aryl-C₁₋₆ alkyl, or R' and R'' and the atoms to which they are attached together can form a C₃₋₈ heterocyclic ring, wherein the term "substituted", as applied to alkyl, alkenyl, alkynyl, heterocyclyl, cycloalkyl, aryl, heteroaryl, alkylaryl, alkylheteroaryl, arylalkyl and heteroarylalkyl, refers to substitution by one or more alkyl, aryl, heteroaryl, halo, -OR' and -NR'R'' groups, or pharmaceutically acceptable salts thereof.

2. The compounds according to claim 1, wherein:

n has the value of 0 or 1, and Cy is a C₅₋₁₀ heteroaryl group chosen from the group of 2-furanyl or 3-furanyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 1,3,4-oxadiazol-2-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 1,3,4-thiadiazol-2-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl and 4-pyridinyl, which heteroaryl groups are optionally substituted with up to three non-hydrogen substituents independently selected from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, halo, and C₂₋₆ alkynyl substituted with phenyl.

3. The compound according to claim 1 or 2, wherein n is 0.

4. The compound according to any one of claims 1 to 3, wherein n is 0 and Cy is 2-furanyl.

5. The compound according to any one of claims 1 to 3, wherein Cy is 2-furanyl substituted with halo.

6. A compound selected from the group consisting of:

N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(4-phenylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-(2-pyridinyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(5-(phenylethynyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(furan-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(oxazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(oxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(oxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isoxazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isoxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(3-bromoisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,

N-(3-methoxyisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,2,4-oxadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,2,4-oxadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,3,4-oxadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(thiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(thiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(thiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isothiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isothiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(isothiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,2,4-thiadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,2,4-thiadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(1,3,4-thiadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane,
N-(pyridin-4-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane, and pharmaceutically acceptable salts thereof.

7. A compound selected from the group consisting of:

N-(furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(3-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-methylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(3-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(3-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-bromofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(4-phenylfuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-(2-pyridinyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(5-(phenylethynyl)furan-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(furan-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(oxazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(oxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(oxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isoxazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isoxazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
N-(isoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,

N-(3-bromoisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(3-methoxyisoxazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(1,2,4-oxadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(1,2,4-oxadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(1,3,4-oxadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(thiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(thiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(thiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(isothiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(isothiazol-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(isothiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(1,2,4-thiadiazol-3-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(1,2,4-thiadiazol-5-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(1,3,4-thiadiazol-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane,
 N-(pyridin-4-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane, and pharmaceutically acceptable salts thereof.

8. The compound N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.0]octane, or pharmaceutically acceptable salts thereof.

9. The compound N-(5-chlorofuran-2-ylcarbonyl)-3,7-diazabicyclo[3.3.1]nonane, or pharmaceutically acceptable salts thereof.

10. Use of the compound according to any one of claims 1 to 9, in the manufacture of a medicament.

11. A method for treatment of central nervous system disorders, comprising administering to a mammal in need of such treatment, a therapeutically effective amount of the compound according to any one of claims 1 to 9.

12. The method of claim 11, wherein the disorder is selected from the group consisting of age-associated memory impairment, mild cognitive impairment, pre-senile dementia (early onset Alzheimer's disease), senile dementia (dementia of the Alzheimer's type), Lewy body dementia, vascular dementia, Alzheimer's disease, stroke, AIDS dementia complex, attention deficit disorder, attention deficit hyperactivity disorder, dyslexia, schizophrenia, schizophreniform disorder and schizoaffective disorder.

13. The method of claim 11, wherein the disorder is selected from the group consisting of mild to moderate dementia of the Alzheimer's type, attention deficit disorder, mild cognitive impairment and age associated memory impairment.

14. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of a compound according to any one of claims 1 to 9, in association with one or more pharmaceutically acceptable diluents, excipients and/or inert carriers.

15. Use of the pharmaceutical composition according to claim 14 in the manufacture of a medicament for treatment of central nervous system disorders.

FIG. 1





FIG. 2



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2007/083330

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D471/08 A61P25/00 A61K31/407 C07D487/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2004/016616 A (ASTRAZENECA AB [SE]; ERNST GLEN [US]; PHILLIPS EIFION [US]; SCHMIESING) 26 February 2004 (2004-02-26) the whole document	1-15
A	WO 2004/076453 A (NEUROSEARCH AS [DK]; PETERS DAN [DK]; OLSEN GUNNAR M [DK]; NIELSEN ELS) 10 September 2004 (2004-09-10) the whole document	1-15
A,P	WO 2006/124748 A (LEXICON GENETICS INC [US]; BARBOSA JOSEPH [US]; DONG LI [US]; FINK CYN) 23 November 2006 (2006-11-23) the whole document	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

9 April 2008

Date of mailing of the international search report

18/04/2008

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lange, Tim

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2007/083330

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 11-13 are directed to a method of treatment of the human body, the search has been carried out and based on the alleged effects of the compound.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers allsearchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search reportcovers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2007/083330

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2004016616 A	26-02-2004	AU 2003248590 A1	03-03-2004
		BR 0313233 A	14-06-2005
		CA 2493246 A1	26-02-2004
		CN 1675215 A	28-09-2005
		EP 1539764 A1	15-06-2005
		JP 2005539030 T	22-12-2005
		KR 20060005334 A	17-01-2006
		MX PA05001584 A	25-04-2005
		US 2006052368 A1	09-03-2006
		ZA 200501227 A	26-07-2006
WO 2004076453 A	10-09-2004	AU 2004215658 A1	10-09-2004
		BR PI0407216 A	24-01-2006
		CA 2518675 A1	10-09-2004
		EP 1599476 A1	30-11-2005
		JP 2006519208 T	24-08-2006
		MX PA05006861 A	12-12-2005
WO 2006124748 A	23-11-2006	AU 2006247482 A1	23-11-2006
		CA 2608240 A1	23-11-2006
		EP 1893212 A2	05-03-2008