

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
10 May 2007 (10.05.2007)

PCT

(10) International Publication Number  
**WO 2007/051333 A1**

(51) International Patent Classification:

A61K 31/53 (2006.01) C07D 405/12 (2006.01)  
A61P 25/28 (2006.01)

CH-8057 Zürich (CH). **BARBERIS, Alcide** [CH/CH];  
Allenmoosstrasse 64, CH-8057 Zürich (CH).

(21) International Application Number:

PCT/CH2006/000607

(74) Agent: **E. BLUM & CO.**; Vorderberg 11, CH-8044 Zurich (CH).

(22) International Filing Date: 30 October 2006 (30.10.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

05023893.0 2 November 2005 (02.11.2005) EP  
60/734,384 8 November 2005 (08.11.2005) US

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **ON-CALIS AG** [CH/CH]; Wagistrasse 21, CH-8952 Schlieren (CH).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **LÜTHI, Urs** [CH/CH]; Hotzenweidstrasse 43, CH-8330 Pfäffikon (CH). **HUANG, Danzhi** [CN/CH]; Roswiesenstrasse 171, CH-8051 Zürich (CH). **KOLB, Peter** [AT/CH]; Schürbungert 14, CH-8057 Zürich (CH). **CECCHINI, Marco** [IT/CH]; Hubenstrasse 20, CH-8057 Zürich (CH). **MAJEUX, Nicolas** [CH/CH]; Auhofstrasse 24, CH-8051 Zürich (CH). **DEY, Fabian** [CH/CH]; Hanfrosee 21, CH-8055 Zürich (CH). **AUDÉTAT, Stephan Valentin** [CH/CH]; Meierwiesenstrasse 48, CH-8107 Buchs (CH). **CAFLISCH, Amedeo** [CH/CH]; Malvenstrasse 10,

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TRIAZINE BETA-SECRETASE INHIBITORS



(57) Abstract: It has been found that compounds of formula (I) wherein, in preferred embodiments, X<sub>1</sub> and X<sub>2</sub> are N-H or N-R<sub>3</sub> or N-R<sub>4</sub>, R<sub>N</sub> and R<sub>C</sub> are both H, R<sub>1</sub> and R<sub>3</sub> are independently from each other optionally substituted phenyl, or R<sub>1</sub> and R<sub>2</sub> and/or R<sub>3</sub> and R<sub>4</sub> form together with the N to which R<sub>2</sub> and/or R<sub>4</sub>, respectively, are bound a 5 or 6 membered heterocycle comprising 1 or 2 heteroatoms, and A is preferably halogen substituted phenyl or a preferably phenyl substituted furanyl are good β-secretase inhibitors and are useful for the treatment of Alzheimer's disease.

## Triazine $\beta$ -secretase inhibitors

### Field of the Invention

This invention relates to compounds acting as  
5 beta-secretase inhibitors.

### Background Art

Alzheimer's disease (AD) is the most common  
10 form of dementia among older people, and affects parts of  
the brain that control thought, memory and language.  
Susceptibility to Alzheimer's disease increases with age,  
but Alzheimer's disease is not a normal part of the ageing  
process.

15 A characteristic of this disease is the  
presence of extracellular senile plaque, the major  
component of which is the  $\beta$ -amyloid peptide ( $A\beta$ ). The  
hydrophobic, 39-43-amino-acid-long  $A\beta$  peptide is excised  
from the amyloid precursor protein (APP) by sequential  
20 cleavage by the so-called  $\beta$ - and  $\gamma$ -secretases.

Known genetic predispositions for AD mostly  
affect genes involved in  $A\beta$  generation or  $A\beta$  deposition.  
Since the  $A\beta$  peptide seems to play an important role in  
the pathogenesis of AD, current therapeutic strategies  
25 often focus on inhibition of  $A\beta$  deposition and generation.  
Inhibition of  $\beta$ -secretase activity represents an  
attractive option to achieve this goal.

Despite major efforts to identify novel  $\beta$ -  
secretase inhibitors by applying in vitro high-throughput  
30 screening (HTS) assays with purified soluble BACE-1  
fragments and fluorogenic peptide substrates, the best  
progress towards efficient BACE-1 inhibition has been  
achieved so far by the use of peptidic transition-state  
mimetic compounds. However, for efficient inhibition of  $\beta$ -  
35 secretase in cells, their molecular weight must be reduced  
and their structure modified so as to allow for permeation

of cellular membranes, the blood-brain barrier and for activity in the natural cellular environment.

There also exist some assays for identifying low molecular weight inhibitors of secretases that can  
5 block these membrane-bound enzymes at the natural location within intracellular compartments. Cell-based HTS assays, however, are generally faced with the problem that selection signals are often caused by compounds that interfere with cellular processes or pathways that are  
10 redundant with that of the target. For example, some compounds found by mammalian cell based assays impair the production of A $\beta$  through the increase of the pH in intracellular compartments, or they function through protein phosphorylation, or they simply catalyze  
15 polymerization of A $\beta$ , thus reducing the percentage of soluble peptide.

Some candidate compounds for inhibiting the production of A $\beta$  peptide in a biological system have been proposed in US 5,814,646 and US 5,624,937, and triazine  
20 compounds have already been suggested for the treatment of Alzheimer's disease (see GB 2 397 301, WO 2004/063196).

Nevertheless, there is still a need for potent  $\beta$ -secretase inhibitors that directly inhibit  $\beta$ -secretase.


25

### Disclosure of the Invention

Hence, it is a general object of the invention to provide compounds that directly act as  $\beta$ -secretase  
30 inhibitors.

Now, in order to implement these and still further objects of the invention, which will become more readily apparent as the description proceeds, the  $\beta$ -secretase inhibitors of the present invention are  
35 manifested by the following formula I

5



wherein

10  $X_1$  and  $X_2$  independently from each other are selected from O or S or N- $R_2$  or N- $R_4$ , and in particular  $X_1$  is N- $R_2$  and  $X_2$  is N- $R_4$ ,


$R_N$  and  $R_C$  are independently from each other chosen from small substituents such as -H, -OH, -CH=O,  $NH_2$ , preferably -H,

15  $R_1$  and  $R_3$  are independently from each other selected from the group comprising C1-C3 alkoxy, in particular C1-C3 alkoxy, 3- to 8-membered aliphatic or heteroaliphatic cycles, optionally substituted 5- or 6-membered heteroaryl or aryl, in particular optionally  
 20 substituted phenyl, with the substituents being independently from each other selected from 1 to 5, in particular 1 or 2, of C1-C6 alkyl, halogen, C1-C6 alkoxy, hydroxy and C2-C6 alkene, wherein preferred alkyls are C1-C3 alkyl, in particular methyl, the halogens are selected  
 25 from fluorine, chlorine, bromine and iodine, in particular from fluorine and chlorine, preferred alkoxy are C1-C3 alkyloxy, in particular methoxy, and preferred alkene are vinyl or allyl,

$R_2$  and  $R_4$  are independently from each other  
 30 selected from the group comprising hydrogen, C1-C3 alkyl, C1-C3 alkoxy, phenoxy, C1-C3 thioalkyl, wherein preferred C1-C3 alkyl is methyl, preferred C1-C3 alkoxy is methoxy, and preferred C1-C3 thioalkyl is -S-CH<sub>3</sub>,


or  $R_1$  and  $R_2$  and/or  $R_3$  and  $R_4$  form together  
 35 with the N to which  $R_2$  and/or  $R_4$ , respectively, are bound a 5 or 6 membered heterocycle comprising 1 or 2

heteroatoms, the second heteroatom being N or O, in particular



A is

10



15

wherein

R<sub>5</sub> is selected from the group comprising -H, -  
20 OH, -CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NH-CH<sub>2</sub>CH<sub>3</sub>,

R<sub>5'</sub> is selected from the group comprising -H, -  
-OH, -CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NH-CH<sub>2</sub>CH<sub>3</sub>, -O-  
CH<sub>2</sub>CH<sub>3</sub>, -O-phenyl, -CH<sub>2</sub>-phenyl, -NH-phenyl, -F, -Cl, -Br,  
-I, and much preferred is -H,

25 R<sub>6</sub> is selected from the group comprising -H, -  
F, -Cl, -Br, -I,


R<sub>7</sub> is selected from the group comprising -H, -  
OH, -CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NH-CH<sub>2</sub>CH<sub>3</sub>, -O-  
CH<sub>2</sub>CH<sub>3</sub>, -O-phenyl, -CH<sub>2</sub>-phenyl, -NH-phenyl,

30 R<sub>8</sub> is selected from the group comprising -H, -  
F, -Cl, -Br, -I, and wherein

R<sub>9</sub> is selected from the group comprising -H, -  
CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, phenyl, -CH<sub>2</sub>-phenyl, -O-phenyl,  
-NH-phenyl, Br,

35 R<sub>10</sub> is selected from the group comprising -H,  
-CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, phenyl, -CH<sub>2</sub>-phenyl, -O-phenyl,  
-NH-phenyl, Br,

$R_{11}$  is selected from the group comprising -  
 $\text{CH}_2\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{NH}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{NH}-\text{CH}_2\text{COOH}$ , -  
 $\text{O}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{O}-\text{CH}_2\text{COOH}$ ,  $-\text{S}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{S}-\text{CH}_2\text{COOH}$ ,  $-\text{Br}$ ,  
 5 are optionally substituted, in particular



wherein  $R_{12}$  is selected from the group  
 comprising  $-\text{H}$ , halogen selected from  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , in  
 15 particular  $-\text{Cl}$ , C1-C3 alkyl, in particular  $-\text{CH}_3$ ,  $-\text{OH}$ , C1-  
 C3 alkoxy, in particular methoxy,

$R_{13}$  is selected from the group comprising  $-\text{H}$ ,  
 $-\text{OH}$ ,  $-\text{NO}_2$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , in particular  $-\text{Cl}$ , C1-C3 alkyl,  
 in particular  $-\text{CH}_3$ , C1-C3 alkoxy, in particular methoxy,

20  $R_{14}$  is selected from the group comprising  $-\text{H}$ ,  
 $-\text{COOH}$ ,  $-\text{COOR}$  with R being selected from C1-C6 alkyl or C2-  
 C6 alkenyl,  $-\text{NO}_2$ ,  $-\text{Cl}$ ,  $-(\text{C}=\text{O})\text{CH}_3$ ,  $\text{SO}_2$ ,  
 or pharmaceutically acceptable salts thereof.

25 In preferred embodiments, at least one of the  
 below mentioned preferences is present, preferably two or  
 more of them, most preferred all of them.

(i)  $X_1$  is  $\text{N}-\text{R}_2$  and  $X_2$  is  $\text{N}-\text{R}_4$ ;

(ii) at least one of  $R_1$  and  $R_3$  are optionally  
 30 substituted phenyl with the substituents being independent  
 from each other selected from 1 or 2 of C1-C3 alkyl, in  
 particular methyl; halogen selected from chlorine, bromine  
 and iodine, in particular fluorine and chlorine; C1-C3  
 alkyloxy, in particular methoxy, and


35  $R_2$  and  $R_4$  are hydrogen;

(iii)  $R_1$  and  $R_2$  and/or  $R_3$  and  $R_4$  form together  
 with the N to which  $R_2$  and/or  $R_4$ , respectively, are bound

a 5 or 6 membered heterocycle comprising 1 or 2 heteroatoms, the second heteroatom being N or O, in particular piperidinyl, morpholinyl, pyrrolinyl;

(iv) at most one of  $R_1$  or  $R_3$  is C1-C3-alkoxy, in particular methoxy

(v) A is



wherein

$R_5$  is -H or -OH,


$R_5'$  is -H,

$R_6$  is selected from the group comprising -H, -Cl, -Br, -I, -OCH<sub>3</sub>,

$R_7$  is -H or -OH,

$R_8$  is selected from the group comprising H, Cl, Br, or I,

(vi) A is



wherein

$R_{12}$  is selected from the group comprising -H, -Cl, -CH<sub>3</sub>,

$R_{13}$  is selected from the group comprising -H, -OH, -Cl, -NO<sub>2</sub>,

$R_{14}$  is selected from the group comprising -H, -COOH, -NO<sub>2</sub>, -Cl;

(vii)  $R_N$  and  $R_C$  are both -H.

Much preferred are compounds wherein  $X_1$  is N- $R_2$ ,  $X_2$  is N- $R_4$ , and  $R_N$  and  $R_C$  are both -H, and which additionally have at least one of the preferences below,  
5 namely

(i)  $R_1$  and  $R_3$  are optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of methyl, fluorine, chlorine, and methoxy, and

10  $R_2$  and  $R_4$  are hydrogen;


(ii)  $R_1$  and  $R_2$ , and  $R_3$  and  $R_4$  form together with the N to which  $R_2$  and  $R_4$  are bound a heterocycle selected from the group consisting of piperidinyl, morpholinyl, pyrrolinyl;

15 (iii)  $R_1$  is optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of methyl, fluorine, chlorine, and methoxy, and  $R_2$  is hydrogen, and  $R_3$  and  $R_4$  form together with the N to which  $R_4$  is bound a heterocycle selected  
20 from the group consisting of piperidinyl, morpholinyl, pyrrolinyl;

(iv)  $R_1$  is as defined in (i) or (ii) and  $R_3$  is methoxy;

(v) A is

25




30

wherein at least one of  $R_6$  and  $R_8$  is Cl, Br, I, preferably both, and much preferred both are the same;

(vi) A is


35





wherein at least 3 of  $R_5$  to  $R_8$  are not hydrogen, preferably at least  $R_8$  and much preferred  $R_8$  and  $R_6$  are Cl, Br or I, in particular  $R_8$  and  $R_6$  are the same;

(vii) A is




wherein  $R_{14}$  is  $-\text{COOH}$ ;

(vii)  $R_{14}$  is  $-\text{COOH}$  and  $R_{13}$  is  $-\text{OH}$  or  $-\text{Cl}$ ;

(viii)  $R_{13}$  is  $-\text{OH}$  or  $-\text{Cl}$  or  $-\text{NO}_2$ .

In a further aspect, the invention relates to a compound of formula (I-B)



(I-B)

wherein

$X_1$  represents O, S or  $\text{N}-R_2$ ;

$X_2$  represents O, S or  $\text{N}-R_4$ ;

$R_N$  represents hydrogen, hydroxy, formyl or amino;


$R_C$  represents hydrogen, hydroxy, formyl or amino;

- R<sub>1</sub> represents C1-C6 alkyl, 3- to 8-membered aliphatic cycles or 3- to 8-membered heteroaliphatic cycles, optionally substituted 5- or 6-membered heteroaryl or optionally substituted aryl, in particular optionally substituted phenyl, with the substituents of the phenyl being independent from each other selected from 1 to 5, preferably from 1 or 2, of C1-C6 alkyl, halogen selected from fluorine, chlorine, bromine, iodine, C1-C6 alkyloxy, hydroxy, C1-C6 alkene;
- R<sub>2</sub> represents hydrogen, C1-C3 alkyl, C1-C3 alkoxy, phenoxy, C1-C3 thioalkyl;
- R<sub>3</sub> represents C1-C6 alkoxy, 3- to 8-membered aliphatic cycles or 3- to 8-membered heteroaliphatic cycles, optionally substituted 5- or 6-membered heteroaryl or optionally substituted aryl, in particular optionally substituted phenyl, with the substituents of the phenyl being independent from each other selected from 1 to 5, preferably from 1 or 2, of C1-C6 alkyl, fluorine, chlorine, bromine, iodine, C1-C6 alkyloxy, hydroxy, C1-C6 alkene; and
- R<sub>4</sub> represents hydrogen, C1-C3 alkyl, C1-C3 alkoxy, phenoxy, C1-C3 thioalkyl; or
- R<sub>1</sub> and R<sub>2</sub> and/or R<sub>3</sub> and R<sub>4</sub> form together with the N to which R<sub>2</sub> and/or R<sub>4</sub>, respectively, are bound a 5 or 6 membered heterocycle comprising 1 or 2 heteroatoms, the second heteroatom being N or O;
- R<sub>9</sub> represents -H, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, phenyl, benzyl, phenoxy, -NH-phenyl, Br;
- R<sub>10</sub> represents -H, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, phenyl, benzyl, phenoxy, -NH-phenyl, Br;
- R<sub>11</sub> represents -(C<sub>1</sub>-C<sub>3</sub>-alkyle)COOH, -NH-(C<sub>1</sub>-C<sub>3</sub>-alkyl)COOH, -O-(C<sub>1</sub>-C<sub>3</sub>-alkyle)COOH, -S-(C<sub>1</sub>-C<sub>3</sub>-alkyl)COOH, halogen, or a 5-membered or 6-membered heteroaromatic or aromatic ring, which ring is optionally substituted by 1 to 3

substituents, the substituents are selected from the group comprising halogene, -NO<sub>2</sub>, hydroxy, -SO<sub>2</sub>, -COOH, C1-C6 alkyl, C1-C6 alkoxy, -(C=O)-(C1-C6 alkyl), -COOR, wherein R is selected from  
 5 C1-C6 alkyl or C2-C6 alkenyl;  
 or pharmaceutically acceptable salts thereof.

Advantageous embodiments of compounds of formula (I-B) are  
 10 described as follows:


- (i) A compound of formula (I-B) wherein X<sub>1</sub> represents N-R<sub>2</sub> and X<sub>2</sub> represents N-R<sub>4</sub>.
- (ii) A compound of formula (I-B) wherein R<sub>N</sub> represents hydrogen.
- 15 (iii) A compound of formula (I-B) wherein
  - R<sub>1</sub> represents methyl, optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of methyl, fluorine, chlorine, methoxy;
  - 20 R<sub>2</sub> represents hydrogen, methoxy;
  - R<sub>3</sub> represents methoxy, optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of methyl, fluorine, chlorine, methoxy;
  - 25 R<sub>4</sub> represents hydrogen, methoxy;  
and / or
  - R<sub>1</sub> and R<sub>2</sub> and/or R<sub>3</sub> and R<sub>4</sub> form together with the N to which R<sub>2</sub> and/or R<sub>4</sub>, respectively, are bound a 5 or 6 membered heterocycle selected from  
 30 pyrrolidinyl, piperidinyl, morpholinyl.
- (iv) A compound of formula (I-B), wherein  
 R<sub>11</sub> represents



and  $R_{12}$ ,  $R_{13}$ , and  $R_{14}$  are as defined in claim 1.

(v) A compound of formula (I-B) wherein

$R_{11}$  represents  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{NH}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{NH}-\text{CH}_2\text{COOH}$ ,  $-\text{O}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{O}-\text{CH}_2\text{COOH}$ ,  $-\text{S}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{S}-\text{CH}_2\text{COOH}$ ,  $-\text{Br}$ , or



wherein

$R_{12}$  represents  $-\text{H}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ ,  $-\text{CH}_3$ ,  $-\text{OH}$ ,  $-\text{OCH}_3$ ;

$R_{13}$  represents  $-\text{H}$ ,  $-\text{OH}$ ,  $-\text{NO}_2$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ ,  $-\text{CH}_3$ ,  $\text{OCH}_3$  and

$R_{14}$  represents  $-\text{H}$ ,  $-\text{NO}_2$ ,  $\text{Cl}$ ,  $-(\text{C}=\text{O})-\text{CH}_3$ ,  $\text{SO}_2$ ,  $-\text{COOH}$ ,  $-\text{COOCH}_3$ ;

(vi) A compound of formula (I-B) wherein

$R_1$  represents optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of C1-C3 alkyl (in particular methyl), halogen (in particular fluorine and chlorine). C1-C3 alkyloxy (in particular methoxy)

$R_3$  represents optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of C1-C3 alkyl (in particular methyl), halogen (in particular fluorine and chlorine). C1-C3 alkyloxy (in particular methoxy).

(vii) A compound of formula (I-B), wherein  $R_2$  and  $R_4$  represent hydrogen.

(iix) A compound of formula (I-B), wherein  $R_1$  and  $R_2$  and/or  $R_3$  and  $R_4$  form together with the N to which  $R_2$  and/or  $R_4$ , respectively, are bound a 5 or 6 membered heterocycle comprising 1 or 2 heteroatoms, the second heteroatom being N or O.

(ix) A compound of formula (I-B), wherein  $R_1$  and  $R_2$ , and/or  $R_3$  and  $R_4$  form together with the N to which  $R_2$  and/or  $R_4$ , respectively, are bound a heterocycle

selected from the group of piperidinyl, morpholinyl, pyrrolinyl.

(x) A compound of formula (I-B), wherein

$R_{12}$  represents -H, -Cl, -CH<sub>3</sub>,


5  $R_{13}$  represents -H, -OH, -Cl, -NO<sub>2</sub>,

$R_{14}$  represents -H, -COOH, -NO<sub>2</sub>, Cl.

(xi) A compound of formula (I-B), wherein  $R_{14}$  represents -COOH.

10 (xii) A compound of formula (I-B), wherein  $R_{13}$  represents -OH or Cl.

In a further aspect, the invention relates to a compound of formula (I-C)



(I-C)

15 wherein

$R_1$  represents optionally substituted phenyl, with the substituents being independently from each other selected from 1 to 5, in particular 1 or 2, of C1-C6 alkyl, halogen, C1-C6 alkoxy, hydroxy and C2-C6 alkene, wherein preferred alkyls are C1-C3 alkyl, in particular methyl, the halogens are selected from fluorine, chlorine, bromine and iodine, in particular from fluorine and chlorine, preferred alkoxy are C1-C3 alkyloxy, in particular methoxy, and preferred alkene are vinyl or allyl and

25

$R_N$ ,  $R_C$ ,  $R_5$ ,  $R_5'$ ,  $R_6$ ,  $R_7$ ,  $R_8$  are as defined above.

Advantageous embodiments of compounds of formula (I-C) are described as follows:

- (i) A compound of formula (I-C) wherein  $R_C$  represents hydrogen.
- (ii) A compound of formula (I-C) wherein  $R_N$  represents hydrogen.
- 5 (iii) A compound of formula (I-C) wherein  $R_5$  represents hydrogen.
- (iv) A compound of formula (I-C) wherein  $R_5$  represents hydroxy.

10 It is to be understood that the various embodiments and preference as disclosed herein may be combined at will. Further, selected definitions may not apply.  
The invention further relates to compounds of formula (I) as pharmaceutical.


15

The invention further relates to compounds of formula (I-B) as pharmaceutical.

The invention further relates to compounds of  
20 formula (I-C) as pharmaceutical.

The invention further relates to a pharmaceutical composition comprising one or more compounds or pharmaceutically acceptable salts thereof as  
25 described herein, together with a pharmaceutically acceptable carrier and optionally one or more adjuvants.

The invention further relates to the  
30 manufacture of a compound of formula (I) comprising the step of reacting a compound of formula (II)




wherein the substituents are as defined above  
with a compound of formula (III)

5




wherein the substituents are as defined above,  
optionally in the presence of a solvent, such as an  
10 alcohol (e.g. methanol) and optionally in the presence of  
an reaction auxiliary, such as a water-removing agent  
(e.g. molecular sieve).

The invention further relates to the  
15 manufacture of a compound of formula (I-B) comprising the  
step of reacting a compound of formula (II-B)




20 wherein the substituents are as defined above  
with a compound of formula (III-B)



(III-B)


wherein the substituents are as defined above,  
 optionally in the presence of a solvent, such as an  
 5 alcohol (e.g. methanol) and optionally in the presence of  
 an reaction auxiliary, such as a water-removing agent  
 (e.g. molecular sieve).

The invention further relates to the  
 10 manufacture of a compound of formula (I-C) comprising the  
 step of reacting a compound of formula (II-C)



(II-C)

15 wherein the substituents are as defined above  
 with a compound of formula (III)



(III-C)



wherein the substituents are as defined above, optionally in the presence of a solvent, such as an alcohol (e.g. methanol) and optionally in the presence of an reaction auxiliary, such as a water-removing agent  
5 (e.g. molecular sieve).

The starting materials used in these reactions are known or available by using known manufacturing steps. The compounds of formula (II). (II-B) and (II-C) are  
10 available from the reaction of hydrazine with the corresponding chlorine-derivatives.

It has been found that compounds of the above formulas are efficient in inhibiting  $\beta$ -secretase activity.  
15 Thus, such compounds are suitable in the prophylaxis, treatment and/or delay of progression of diseases related to  $\beta$ -secretase activity or diseases related to the deposition of amyloid beta-protein, respectively. Such diseases include Alzheimer's disease (AD), Down's syndrome  
20 and advanced aging of brain of which AD is of particular importance.

Further, the invention relates to the use of a compound as described herein, or a pharmaceutically acceptable salt  
25 thereof, in the manufacture of a medicament for inhibiting the  $\beta$ -secretase activity or diseases related to the deposition of amyloid beta-protein (such as the production and/or the accumulation of amyloid beta-protein) mammals, in particular human beings. In particular, the invention  
30 relates to the use of a compound as described herein, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment of Alzheimer's disease and / or Down's syndrome and / or aging of brain.

Accordingly, the invention also relates to a method of prevention, treatment or delay of progression of a disease involving an abnormal  $\beta$ -secretase activity or a disease  
5 related to the deposition of amyloid beta-protein (such as the production and/or the accumulation of amyloid beta-protein), in a subject in need thereof comprising the step of administering an effective amount of a compound as described herein. In particular, the invention relates to  
10 method of prevention, treatment or delay of progression of a disease selected from the group consisting of Alzheimer's disease (AD) , Down's syndrome , aging of brain, in a subject in need thereof comprising the step of administering an effective amount of a compound as  
15 described herein.

AD, as used in this specification includes all stages of this disease, its mild, moderate and severe form.

20

In therapeutic applications, the compounds are administered to a host already suffering from the disease. The compounds will be administered in an amount sufficient to inhibit further deposition of senile plaques. The  
25 specific dose of compound(s) administered according to this invention to obtain therapeutic and/or prophylactic effects will, of course, be determined by the particular circumstances, such as the specific compound administered, the condition being treated, etc. A daily dose will  
30 contain a dosage level of from about 0.01 mg/kg to about 50 mg/kg of body weight of an active compound, preferably from about 0.05 mg/kg to about 20 mg/kg, for example from about 0.1 mg/kg to about 120 mg/kg.

The compound can be administered by a variety  
35 of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular and intranasal either as such, but preferable in a formulation comprising

carriers adjuvants etc. Suitable pharmaceutically acceptable solid and liquid carriers and/or pharmaceutically acceptable adjuvants, such as stabilizing agents, emulsifiers, etc. are known in the art.

5                   For example, a typical pharmaceutical composition for intramuscular injection would contain about one  $\mu\text{g}$  to one mg of the compound in from one to four milliliters of sterile buffered water. The typical pharmaceutical composition for intravenous infusion would  
10 contain about one to one hundred milligrams of the compound in from one hundred to five hundred milliliters of sterile Ringer's solution.


                  The pharmaceutical formulations are prepared by known procedures using known and readily available  
15 ingredients.

#### Modes for Carrying Out the Invention


                  Specific compounds and their  $\beta$ -secretase  
20 inhibiting effects are further described below.

                  The compounds investigated were the compounds with the following formulas:


202E09




201C08




201G11




201G06





206E02






205C07







5




10

21




5




10

250C04



250B04



5

The tests performed were

a) A $\beta$ 1-40 (Sw) bioassay, which measures the  
10 amount of the amyloid peptide A $\beta$ 1-40 in the supernatant of  
Swedish APP695 transgenic HEK293 cells in the presence of  
the various BACE inhibitors via ELISA (enzyme-linked  
immunosorbent assay). In the table, the inhibitory  
concentration that reduces A $\beta$ 1-40 secretion to 50 % is  
15 indicated (IC<sub>50</sub>), or the % reduction of A $\beta$ 1-40 secretion  
at the indicated concentration.

b) FRET assay, which measures the activity of  
recombinant BACE enzyme in the presence of the various  
BACE inhibitors via a FRET (fluorescence resonance energy  
20 transfer)-based readout. In the table, the inhibitory  
concentration that reduces the activity of BACE to 50 % is  
indicated (IC<sub>50</sub>), or the % reduction of the activity of  
BACE at the indicated concentration.

c) An additional in silico test was performed  
25 for the compounds listed in Tables 1 to 3. The compounds  
were docked with the FFLD approach (Budin et al., Biol.  
Chem. 382, 1365-1372, 2001) and their binding energy was  
evaluated with the LIECE method (Huang and Caflisch, J.  
Med. Chem. 47, 5791-5797, 2004). The affinity evaluated  
30 with LIECE is in the low micromolar range for most of  
these compounds.

Table 1:


Compound	LIECE Ki [ $\mu$ M]	A $\beta$ 1-40(sw) bioassay (cell-based)	BACE FRET assay (in vitro)
202E09	41.41	25% (5 $\mu$ M)	IC <sub>50</sub> 53.9 $\mu$ M
201C08	20.20	25% (5 $\mu$ M)	IC <sub>50</sub> 11.2 $\mu$ M
201G11	74.22	26% (5 $\mu$ M)	IC <sub>50</sub> 11.9 $\mu$ M
201G06	38.77	No (5 $\mu$ M)	IC <sub>50</sub> 25.5 $\mu$ M
206E02	49.35	27% (50 $\mu$ M)	IC <sub>50</sub> 20.6 $\mu$ M
205C07	30.89	No (50 $\mu$ M)	IC <sub>50</sub> 48.1 $\mu$ M
205A04	56.49	No (5 $\mu$ M)	IC <sub>50</sub> 81.8 $\mu$ M
251A07	13.00	not done	IC <sub>50</sub> 10.6 $\mu$ M
251C08	20.00	not done	IC <sub>50</sub> 24.4 $\mu$ M
251A09	14.00	not done	IC <sub>50</sub> 34.4 $\mu$ M
251G09	24.00	not done	IC <sub>50</sub> 169.7 $\mu$ M
251B07	12.00	not done	IC <sub>50</sub> 45.4 $\mu$ M
251C10	--	not done	IC <sub>50</sub> >100 $\mu$ M
251F09	15.00	not done	IC <sub>50</sub> 188.5 $\mu$ M
251A10	18.00	not done	IC <sub>50</sub> 123.7 $\mu$ M
250A05	6.64	IC <sub>50</sub> ~10-20 $\mu$ M	IC <sub>50</sub> 141.5 $\mu$ M
250B05	2.68	IC <sub>50</sub> ~10-20 $\mu$ M	IC <sub>50</sub> 30.9 $\mu$ M
250C04	12.82	0 (25 $\mu$ M)	IC <sub>50</sub> 7.1 $\mu$ M
250B04	12.61	IC <sub>50</sub> ~10-20 $\mu$ M	IC <sub>50</sub> 66.6 $\mu$ M

5 While there are shown and described presently preferred embodiments of the invention, it is to be distinctly understood that the invention is not limited thereto but may be otherwise variously embodied and practiced within the scope of the following claims.



Claims

1. A compound of formula (I-B)



(I-B)

wherein

$X_1$  represents O, S or N- $R_2$ ;

$X_2$  represents O, S or N- $R_4$ ;

$R_N$  represents hydrogen, hydroxy, formyl or amino;

$R_C$  represents hydrogen, hydroxy, formyl or amino;

$R_1$  represents C1-C6 alkyl, 3- to 8-membered aliphatic cycles or 3- to 8-membered heteroaliphatic cycles, optionally substituted 5- or 6-membered heteroaryl or optionally substituted aryl, in particular optionally substituted phenyl, with the substituents of the phenyl being independent from each other selected from 1 to 5, preferably from 1 or 2, of C1-C6 alkyl, halogen selected from fluorine, chlorine, bromine, iodine, C1-C6 alkyloxy, hydroxy, C1-C6 alkene;

$R_2$  represents hydrogen, C1-C3 alkyl, C1-C3 alkoxy, phenoxy, C1-C3 thioalkyl;

$R_3$  represents C1-C6 alkoxy, 3- to 8-membered aliphatic cycles or 3- to 8-membered heteroaliphatic cycles, optionally substituted 5- or 6-membered heteroaryl or optionally substituted aryl, in particular optionally substituted phenyl, with the substituents of the phenyl being independent from each other selected from 1 to 5, preferably from 1 or 2, of C1-C6 alkyl, fluorine,

chlorine, bromine, iodine, C1-C6 alkyloxy,  
hydroxy, C1-C6 alkene; and

R<sub>4</sub> represents hydrogen, C1-C3 alkyl, C1-C3 alkoxy,  
phenoxy, C1-C3 thioalkyl; or

5 R<sub>1</sub> and R<sub>2</sub> and/or R<sub>3</sub> and R<sub>4</sub> form together with the N  
to which R<sub>2</sub> and/or R<sub>4</sub>, respectively, are bound a 5  
or 6 membered heterocycle comprising 1 or 2  
heteroatoms, the second heteroatom being N or O;

10 R<sub>9</sub> represents -H, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, phenyl,  
benzyl, phenoxy, -NH-phenyl, Br;

R<sub>10</sub> represents -H, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, phenyl,  
benzyl, phenoxy, -NH-phenyl, Br;

15 R<sub>11</sub> represents -(C<sub>1</sub>-C<sub>3</sub>-alkyle)COOH, -NH-(C<sub>1</sub>-C<sub>3</sub>-  
alkyl)COOH, -O-(C<sub>1</sub>-C<sub>3</sub>-alkyle)COOH, -S-(C<sub>1</sub>-C<sub>3</sub>-  
alkyl)COOH, halogen, or a 5-membered or 6-  
membered heteroaromatic or aromatic ring, which  
ring is optionally substituted by 1 to 3  
substituents, the substituents are selected from  
the group comprising halogene, -NO<sub>2</sub>, hydroxy, -  
20 SO<sub>2</sub>, -COOH, C1-C6 alkyl, C1-C6 alkoxy, -(C=O)-(C1-C6 alkyl), -COOR, wherein R is selected from  
C1-C6 alkyl or C2-C6 alkenyl;

or pharmaceutically acceptable salts thereof.

25 2. The compound of formula (I-B) claim 1 wherein

X<sub>1</sub> represents N-R<sub>2</sub> and

X<sub>2</sub> represents N-R<sub>4</sub>.

30 3. The compound of formula (I-B) claim 1 or 2, wherein R<sub>N</sub>  
represents hydrogen.

4. The compound of formula (I-B) claim 1 or 2, wherein R<sub>C</sub>  
represents hydrogen.

35 5. The compound of formula (I-B) of anyone of the  
preceeding claims wherein

$R_1$  represents methyl, optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of methyl, fluorine, chlorine, methoxy;

5  $R_2$  represents hydrogen, methoxy;

$R_3$  represents methoxy, optionally substituted phenyl with the substituents being independent from each other selected from 1 or 2 of methyl, fluorine, chlorine, methoxy;


10  $R_4$  represents hydrogen, methoxy; and / or

$R_1$  and  $R_2$  and/or  $R_3$  and  $R_4$  form together with the N to which  $R_2$  and/or  $R_4$ , respectively, are bound a 5 or 6 membered heterocycle selected from pyrrolidinyl, piperidinyl, morpholinyl.

15

6. The compound of formula (I-B) of anyone of the preceding claims, wherein


$R_{11}$  represents



and  $R_{12}$ ,  $R_{13}$ , and  $R_{14}$  are as defined in claim 1.

25 7. The compound of formula (I-B) of anyone of the preceding claims, wherein

$R_{11}$  represents  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{NH}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{NH}-\text{CH}_2\text{COOH}$ ,  $-\text{O}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{O}-\text{CH}_2\text{COOH}$ ,  $-\text{S}-\text{CH}_2\text{CH}_2\text{COOH}$ ,  $-\text{S}-\text{CH}_2\text{COOH}$ ,  $-\text{Br}$ , or



wherein

$R_{12}$  represents  $-\text{H}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ ,  $-\text{CH}_3$ ,  $-\text{OH}$ ,  $-\text{OCH}_3$ ;

R<sub>13</sub> represents -H, -OH, -NO<sub>2</sub>, -Cl, -Br, -I, -CH<sub>3</sub>, OCH<sub>3</sub>  
and

R<sub>14</sub> represents -H, -NO<sub>2</sub>, Cl, -(C=O)-CH<sub>3</sub>, SO<sub>2</sub>, -COOH, -  
COOCH<sub>3</sub>

5

8. The compound of formula (I-B) according to anyone of  
the preceding claims, wherein

R<sub>1</sub> represents optionally substituted phenyl with the  
substituents being independent from each other  
selected from 1 or 2 of C1-C3 alkyl (in particular  
methyl), halogen (in particular fluorine and  
chlorine). C1-C3 alkyloxy (in particular methoxy)

R<sub>3</sub> represents optionally substituted phenyl with the  
substituents being independent from each other  
selected from 1 or 2 of C1-C3 alkyl (in particular  
methyl), halogen (in particular fluorine and  
chlorine). C1-C3 alkyloxy (in particular methoxy).

9. The compound of formula (I-B) according to anyone of  
the preceding claims, wherein R<sub>2</sub> and R<sub>4</sub> represent  
hydrogen.

10. The compound of anyone of the preceding claims,  
wherein R<sub>1</sub> and R<sub>2</sub> and/or R<sub>3</sub> and R<sub>4</sub> form together with  
the N to which R<sub>2</sub> and/or R<sub>4</sub>, respectively, are bound a  
5 or 6 membered heterocycle comprising 1 or 2  
heteroatoms, the second heteroatom being N or O.

11. The compound of anyone of the preceding claims,  
wherein R<sub>1</sub> and R<sub>2</sub>, and/or R<sub>3</sub> and R<sub>4</sub> form together with  
the N to which R<sub>2</sub> and/or R<sub>4</sub>, respectively, are bound a  
heterocycle selected from the group consisting of  
piperidinyl, morpholinyl, pyrrolinyl.

12. The compound of anyone of the preceding claims,  
wherein  
R<sub>12</sub> represents -H, -Cl, -CH<sub>3</sub>,

R<sub>13</sub> represents -H, -OH, -Cl, -NO<sub>2</sub>,  
R<sub>14</sub> represents -H, -COOH, -NO<sub>2</sub>, Cl.

13. The compound of anyone of the preceding claims as  
5 pharmaceutical.
14. Use of a compound of anyone of claims 1 - 12, or a  
pharmaceutically acceptable salt thereof, in the  
manufacture of a medicament for inhibiting the  
10 production and/or the accumulation of amyloid beta-  
protein in warm blooded mammals, in particular human  
beings, especially for the treatment of Alzheimer's  
disease and / or Down's syndrome and / or aging of  
brain.
- 15
15. A pharmaceutical composition comprising one or more  
compounds according to anyone of claims 1 to 12, or a  
pharmaceutically acceptable salt thereof, together  
with a pharmaceutically acceptable carrier and  
20 optionally one or more adjuvants.

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/CH2006/000607

## A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/53 A61P25/28 C07D405/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMCATS CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US; XP002422160 A1877/0078986 A0575/0026531 720397 A0879/0041217 A0879/0041217 A1570/0068441 720399 A1335/0060272 &amp; "AMBINTER STOCK SCREENING COLLECTION" 3 July 2005 (2005-07-03), AMBINTER , PARIS, F-75016, FRANCE</p> <p style="text-align: center;">----- -/--</p>	1-7,9,12

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

27 February 2007

Date of mailing of the international search report

15/03/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Damiani, Federica

## INTERNATIONAL SEARCH REPORT

International application No

PCT/CH2006/000607

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE REGISTRY Chemical Library,  Supplier: Interchim  6 March 2002 (2002-03-06),  XP002422161  retrieved from STN accession no.  RN398471-97-1  abstract</p>	1-7,9,12
X	<p>DATABASE REGISTRY Chemical Library,  Supplier: AsInEx  17 April 2001 (2001-04-17),  XP002422162  retrieved from STN accession no.  RN331655-12-0  abstract</p>	1-7,9,12
X	<p>DATABASE REGISTRY Chemical Library,  Supplier: Ambinter  28 February 2002 (2002-02-28),  XP002422163  retrieved from STN accession no.  RN396725-04-5  abstract</p>	1-7,9,12
X	<p>DATABASE REGISTRY Chemical Library,  Supplier: SPECS 29 May 2003 (2003-05-29),  XP002422164  retrieved from STN accession no.  RN521934-50-9  abstract</p>	1-7,9,12
X,P	<p>HUANG DANZHI ET AL: "In silico discovery  of beta-secretase inhibitors."  JOURNAL OF THE AMERICAN CHEMICAL SOCIETY  26 APR 2006,  vol. 128, no. 16,  26 April 2006 (2006-04-26), pages  5436-5443, XP002421783  ISSN: 0002-7863  the whole document</p>	1-14
X,P	<p>US 2006/223812 A1 (MANDELKOW ECKHARD [DE]  ET AL) 5 October 2006 (2006-10-05)  paragraph [0001]  table 1  RN 339113-32-5claims 1-30,42,43</p>	1-4,6,7, 10-15
A	<p>GB 2 397 301 A (* DE NOVO PHARMACEUTICALS  LTD) 21 July 2004 (2004-07-21)  cited in the application  page 1 - page 2  page 4 - page 5  claims 1-20</p>	1-15
	-/--	

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/CH2006/000607

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>US 6 335 339 B1 (ARENAS JAIME E ET AL)  1 January 2002 (2002-01-01)  abstract  figures 47,69,76,112,144  -----</p>	1-13,15



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/CH2006/000607

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 2006223812	A1	05-10-2006	NONE	
GB 2397301	A	21-07-2004	NONE	
US 6335339	B1	01-01-2002	NONE	