

WO 2011/103289 A2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 August 2011 (25.08.2011)

(10) International Publication Number
WO 2011/103289 A2

PCT

(51) International Patent Classification:
C07D 405/14 (2006.01) *A61K 31/506* (2006.01)
C07D 403/14 (2006.01) *A61K 31/5377* (2006.01)
C07D 401/14 (2006.01) *A61P 35/00* (2006.01)
C07D 413/14 (2006.01)

(74) Agents: **GORDON, Dana, M.** et al.; Patent Group, Foley Hoag LLP, Seaport West, 155 Seaport Boulevard, Boston, MA 02210-2600 (US).

(21) International Application Number:
PCT/US2011/025246

(22) International Filing Date:
17 February 2011 (17.02.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/305,326 17 February 2010 (17.02.2010) US

(71) Applicant (for all designated States except US): **JASCO PHARMACEUTICALS, LLC** [US/US]; 10-N Roessler Road, Woburn, MA 01801 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BALDINO, Carmen, M.** [US/US]; 4 Katarina Lane, Woburn, MA 01801 (US). **CASERTA, Justin, L.** [US/US]; 3 Newport Drive, Billerica, MA 01821 (US). **LEE, Chee-Seng** [MY/US]; 38 Clark Street, Unit 1L, Somerville, MA 02143 (US). **NICEWONGER, Robert, B.** [US/US]; 21 Fairview Avenue, Tyngsboro, MA 01879 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: IMIDAZOLE-2, 4-DIONE INHIBITORS OF CASEIN KINASE 1

(57) Abstract: Disclosed are compounds, pharmaceutical compositions containing those compounds, and uses of the compounds and compositions as modulators of casein kinase 1 (e.g., CK1 γ), the TGF β pathway and/or the Wnt pathway. Uses are also disclosed for the treatment or prevention of a range of therapeutic indications due at least in part to aberrant physiological activity of casein kinase 1 (e.g., CK1 γ), the TGF β pathway and/or the Wnt pathway.

Imidazole-2,4-Dione Inhibitors of Casein Kinase 1

RELATED APPLICATIONS

This application claims the benefit of priority to United States Provisional Patent
5 Application serial number 61/305,326, filed February 17, 2010.

BACKGROUND


Casein kinase 1 (CK1) is a family of evolutionarily conserved serine/threonine kinases including seven known members in vertebrates (CK1 α , - β , - γ 1, - γ 2, - γ 3, - δ and - ϵ). The CK1s contain a typical kinase domain followed by a C-terminal tail region, which has 10 been implicated in the regulation of CK1 localization, substrate selectivity and kinase activity. Myriad proteins have been found to be phosphorylated by CK1s, which are involved in a wide range of cellular functions including vesicular trafficking, DNA damage repair, cell cycle progression, cytokinesis and circadian rhythms (reviewed by Gross and Anderson (1998); Vielhaber and Virshup (2001); Knippschild *et al.* (2005)). Moreover, 15 CK1 family members (- α , - δ/ϵ and - γ) modulate the activities of major signaling pathways (for example, Wnt and Shh) through several mechanisms (Peters *et al.*, 1999; Liu *et al.*, 2002; Price and Kalderon, 2002; Davidson *et al.*, 2005; Zeng *et al.*, 2005 and reviewed by Price (2006)).

In mammals seven CK1 isoforms, namely CK1 α , β , $\gamma_{1,3}$, δ and ϵ , and several splice 20 variants have been described. They all contain a highly conserved kinase domain, a short N-terminal domain of 6 to 76 amino acids and a highly variable C-terminal domain of 24 to more than 200 amino acids. The constitutive phosphotransferase activity of CK1 isoforms is tightly controlled by several mechanisms. For example, the closely related isoforms CK1 δ and ϵ , which share a 98% identity at the amino acid level in their catalytic domain, 25 are regulated by autophosphorylation, dephosphorylation and proteolytic cleavage. Members of the CK1 family are found in the nucleus, the cytoplasm and in the plasma membrane. By phosphorylating many different substrates bearing either a canonical or non-canonical consensus sequence they modulate the activity of key regulator proteins involved in many cellular processes such as cell differentiation, cell proliferation, apoptosis, 30 circadian rhythm, chromosome segregation, and vesicle transport.

SUMMARY

One aspect of the present invention relates to compounds that inhibit casein kinase

1. For example, one embodiment relates to a compound of formula 1:



5 or a pharmaceutically acceptable salt thereof, wherein independently for each occurrence:

W and X are independently oxygen or sulfur;

Z¹, Z² and Z³ are independently C-R²¹ or N, provided that at least one of Z¹ and Z² is N;

R¹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aralkyl, heteroaralkyl, -COR⁶, and -C(O)OR⁶;

10 R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷);

R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶,

15 -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷); or R² and R³ are joined together to form an optionally substituted heterocyclic ring;


R⁴ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocyclylalkyl, aralkyl, heteroaryl, heteroaralkyl, halo, hydroxy, alkoxy, hydroxyalkyl, and alkoxyalkyl;

20 R⁵ is selected from the group consisting of aryl, heteroaryl, heterocyclyl, -N(R⁸)(R⁹), -N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and -N(R⁸)P(O)(OR⁹)(OR⁹);

p is 1, 2, 3, 4, 5, or 6;

- R^6 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;
- R^7 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;
- 5 R^8 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;
- R^9 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl; or R^8 and R^9 are joined together to form a heterocyclic ring;
- 10 R^{20} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, halo, haloalkyl, trifluoromethyl, fluoroalkyl, perfluoroalkyl, thio, cyano, hydroxy, methoxy, alkoxy, phenoxy, aryloxy, heteroaryloxy, carboxyl, alkoxycarbonyl, acyl, nitro, amino, alkylamino, arylamino, heteroarylamino, amido, acylamino, sulfate, sulfonate, sulfonyl, sulfoxido, sulfonamido,
- 15 sulfamoyl, $-[C(R^4)_2]_p-R^5$, $NR^{14}R^{15}$, OR^{16} , $O-[C(R^4)_2]_p-R^5$, $NR^{14}-[C(R^4)_2]_p-R^5$ and SR^{16} ;
- R^{21} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, halo, haloalkyl, thio, cyano, carboxyl, alkoxycarbonyl, acyl, nitro, amino, amido, acylamino, sulfate, sulfonate, sulfonamido, sulfamoyl, NR¹⁴R¹⁵, OR¹⁶, and SR¹⁶;
- 20 R^{14} and R^{15} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, $-[C(R^4)_2]_p-R^5$, $-COR^6$, $-C(O)OR^6$, $-SO_2(R^6)$, $-C(O)N(R^6)(R^7)$, $-SO_2N(R^6)(R^7)$, and $-P(O)(OR^6)(OR^7)$; or R^{14} and R^{15} are joined together to form an optionally substituted heterocyclic ring;
- 25 R^{16} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, $-[C(R^4)_2]_p-R^5$, $-COR^6$, and $-C(O)N(R^6)(R^7)$; and
- r is 0 or 1;
- wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl may be optionally substituted.

Another embodiment relates to a compound of formula 2:



or a pharmaceutically acceptable salt thereof, wherein independently for each occurrence:

- R¹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aralkyl,
5 heteroaralkyl, -COR⁶, and -C(O)OR⁶;
- R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl,
heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶,
-SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷);
- R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,
10 heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶,
-C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷); or R² and R³ are
joined together to form an optionally substituted heterocyclic ring;
- R⁴ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,
heterocyclylalkyl, aralkyl, heteroaryl, heteroaralkyl, halo, hydroxy, alkoxy, hydroxyalkyl,
15 and alkoxyalkyl;
- R⁵ is selected from the group consisting of aryl, heteroaryl, heterocyclyl, -N(R⁸)(R⁹),
-N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -
SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -
S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and -N(R⁸)P(O)(OR⁹)(OR⁹);
- 20 p is 1, 2, 3, 4, 5, or 6;
- R⁶ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,
heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;
- R⁷ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,
heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;

R⁸ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl;

R⁹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl; or R⁸ and R⁹ are

5 joined together to form a heterocyclic ring; and

R²⁰ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclalkyl, halo, haloalkyl, trifluoromethyl, fluoroalkyl, perfluoroalkyl, thio, cyano, hydroxy, methoxy, alkoxy, phenoxy, aryloxy, heteroaryloxy, carboxyl, alkoxycarbonyl, acyl, nitro, amino, alkylamino, 10 arylamino, heteroarylarnino, amido, acylamino, sulfate, sulfonate, sulfonyl, sulfoxido, sulfonamido, sulfamoyl, -[C(R⁴)₂]_p-R⁵, NR¹⁴R¹⁵, OR¹⁶, O-[C(R⁴)₂]_p-R⁵, NR¹⁴-[C(R⁴)₂]_p-R⁵ and SR¹⁶;

wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl may be optionally substituted.

15 Another embodiment relates to a pharmaceutical composition, comprising at least one pharmaceutically acceptable excipient; and any one of the aforementioned compounds.

Another embodiment relates to a method of inhibiting CK1 (e.g., CK1 γ , CK1 γ 1, CK1 γ 2, or CK1 γ 3) activity, comprising contacting CK1 (e.g., CK1 γ , CK1 γ 1, CK1 γ 2, or CK1 γ 3) with any one of the aforementioned compounds or 20 pharmaceutical compositions.

Another embodiment relates to treating or preventing a condition associated with aberrant CK1 (e.g., CK1 γ) activity, comprising the step of administering to a mammal in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

25 Another embodiment relates to a method of treating cancer, comprising administering to a mammal in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

Another embodiment relates to a method of treating Alzheimer's disease, comprising administering to a mammal in need thereof a therapeutically effective amount 30 of any one of the aforementioned compounds or compositions.

Another embodiment relates to a method of treating a Wnt dependent disease, comprising administering to a mammal in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

Another embodiment relates to a method of treating a TGF β dependent disease,
5 comprising administering to a mammal in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

Another embodiment relates to a method of treating or preventing inflammation, inflammatory diseases (e.g., osteoarthritis and rheumatoid arthritis), neurological conditions (e.g., Alzheimer's disease) and neurodegeneration, comprising administering to a mammal
10 in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

Another embodiment relates to a method of treating or preventing bone-related diseases and conditions, including osteoporosis and bone formation, or facilitating bone restoration, comprising administering to a mammal in need thereof a therapeutically
15 effective amount of any one of the aforementioned compounds or compositions.

Another embodiment relates to a method of treating or preventing hypoglycemia, metabolic syndrome and diabetes, comprising administering to a mammal in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

20 Another embodiment relates to a method of influencing apoptosis (e.g., increasing the rate of apoptosis in cancerous cells), comprising administering to a mammal in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

Another embodiment relates to a method of treating or preventing aberrant
25 embryonic development, comprising administering to a mammal in need thereof a therapeutically effective amount of any one of the aforementioned compounds or compositions.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 depicts the relative activity of CK1 γ 1(h) as a function of the concentration
30 of compound 5114.

Figure 2 depicts the relative activity of CK1 γ 2(h) as a function of the concentration of compound 5114.

Figure 3 depicts the relative activity of CK1 γ 3(h) as a function of the concentration of compound 5114.

5 **Figure 4** depicts the relative activity of CK1 γ 2(h) as a function of the concentration of compound 5132.

Figure 5 depicts the relative activity of CK1 γ 2(h) as a function of the concentration of compound 5124.

10 **Figure 6** depicts the relative activity of CK1 γ 2(h) as a function of the concentration of compound 5336.

Figure 7 depicts the relative activity of CK1y as a function of the concentration of compound 5347.

Figure 8 depicts the relative activity of CK1y as a function of the concentration of compound 5343.

15 **Figure 9** depicts the relative activity of CK1y as a function of the concentration of compound 5363.

Figure 10 depicts the relative activity of CK1y as a function of the concentration of compound 5378.

DETAILED DESCRIPTION

20 DEFINITIONS

The definitions of terms used herein are meant to incorporate the present state-of-the-art definitions recognized for each term in the chemical and pharmaceutical fields. Where appropriate, illustration is provided. The definitions apply to the terms as they are used throughout this specification, unless otherwise limited in specific instances, either 25 individually or as part of a larger group.

Where stereochemistry is not specifically indicated, all stereoisomers of the inventive compounds are included within the scope of the invention, as pure compounds as well as mixtures thereof. Unless otherwise indicated, individual enantiomers,

diastereomers, geometrical isomers, and combinations and mixtures thereof are all encompassed by the present invention. Polymorphic crystalline forms and solvates are also encompassed within the scope of this invention.

As used herein, the term "isolated" in connection with a compound of the present invention means the compound is not in a cell or organism and the compound is separated from some or all of the components that typically accompany it in nature.

As used herein, the term "pure" in connection with an isolated sample of a compound of the present invention means the isolated sample contains at least 60% by weight of the compound. Preferably, the isolated sample contains at least 70% by weight of the compound. More preferably, the isolated sample contains at least 80% by weight of the compound. Even more preferably, the isolated sample contains at least 90% by weight of the compound. Most preferably, the isolated sample contains at least 95% by weight of the compound. The purity of an isolated sample of a compound of the present invention may be assessed by a number of methods or a combination of them; e.g., thin-layer, preparative or flash chromatography, mass spectrometry, HPLC, NMR analysis, and the like.

The term "heteroatom" is art-recognized and refers to an atom of any element other than carbon or hydrogen. Illustrative heteroatoms include boron, nitrogen, oxygen, phosphorus, sulfur and selenium.

The term "alkyl" is art-recognized, and includes saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In certain embodiments, a straight chain or branched chain alkyl has about 30 or fewer carbon atoms in its backbone (e.g., C₁-C₃₀ for straight chain, C₃-C₃₀ for branched chain), and alternatively, about 20 or fewer. Likewise, cycloalkyls have from about 3 to about 10 carbon atoms in their ring structure, and alternatively about 5, 6 or 7 carbons in the ring structure.

Unless the number of carbons is otherwise specified, "lower alkyl" refers to an alkyl group, as defined above, but having from one to about ten carbons, alternatively from one to about six carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths.

The term “aralkyl” is art-recognized and refers to an alkyl group substituted with an aryl group (e.g., an aromatic or heteroaromatic group).

The terms “alkenyl” and “alkynyl” are art-recognized and refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, 5 but that contain at least one double or triple bond respectively.

The term “aryl” is art-recognized and refers to 5-, 6- and 7-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, naphthalene, anthracene, pyrene, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, triazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Those aryl 10 groups having heteroatoms in the ring structure may also be referred to as “aryl heterocycles” or “heteroaromatics.” The aromatic ring may be substituted at one or more ring positions with such substituents as described above, for example, halogen, azide, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, alkoxyl, amino, nitro, sulphydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, 15 sulfonamido, ketone, aldehyde, ester, heterocyclyl, aromatic or heteroaromatic moieties, -CF₃, -CN, or the like. The term “aryl” also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings (the rings are “fused rings”) wherein at least one of the rings is aromatic, e.g., the other cyclic rings may be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls.

20 The terms ortho, meta and para are art-recognized and refer to 1,2-, 1,3- and 1,4-disubstituted benzenes, respectively. For example, the names 1,2-dimethylbenzene and ortho-dimethylbenzene are synonymous.

The terms “heterocyclyl”, “heteroaryl”, or “heterocyclic group” are art-recognized and refer to 3- to about 10-membered ring structures, alternatively 3- to about 7-membered 25 rings, whose ring structures include one to four heteroatoms. Heterocycles may also be polycycles. Heterocyclyl groups include, for example, thiophene, thianthrene, furan, pyran, isobenzofuran, chromene, xanthene, phenoxyanthene, pyrrole, imidazole, pyrazole, isothiazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthyridine, 30 quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, pyrimidine, phenanthroline, phenazine, phenarsazine, phenothiazine, piperonyl, furazan, phenoxazine, pyrrolidine, oxolane, thiolane, oxazole, piperidine, piperazine,

morpholine, lactones, lactams such as azetidinones and pyrrolidinones, sultams, sultones, and the like. The heterocyclic ring may be substituted at one or more positions with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, 5 carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.


The term "optionally substituted" refers to a chemical group, such as alkyl, cycloalkyl aryl, and the like, wherein one or more hydrogen may be replaced with a substituent as described herein, including but not limited to halogen, azide, alkyl, aralkyl, 10 alkenyl, alkynyl, cycloalkyl, hydroxyl, alkoxy, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, sulfonamido, ketone, aldehyde, ester, heterocyclyl, aromatic or heteroaromatic moieties, -CF₃, -CN, or the like.

The terms "polycyclyl" or "polycyclic group" are art-recognized and refer to two or 15 more rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings, e.g., the rings are "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each 20 of the rings of the polycycle may be substituted with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.

The term "carbocycle" is art-recognized and refers to an aromatic or non-aromatic ring in which each atom of the ring is carbon.


25 The term "nitro" is art-recognized and refers to -NO₂; the term "halogen" is art-recognized and refers to -F, -Cl, -Br or -I; the term "sulfhydryl" is art-recognized and refers to -SH; the term "hydroxyl" means -OH; and the term "sulfonyl" is art-recognized and refers to -SO₂⁻. "Halide" designates the corresponding anion of the halogens, and 30 "pseudohalide" has the definition set forth on 560 of "Advanced Inorganic Chemistry" by Cotton and Wilkinson.

The terms "amine" and "amino" are art-recognized and refer to both unsubstituted and substituted amines, e.g., a moiety that may be represented by the general formulas:




wherein R₅₀, R₅₁ and R₅₂ each independently represent a hydrogen, an alkyl, an alkenyl, -(CH₂)_m-R₆₁, or R₅₀ and R₅₁, taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure; R₆₁ represents an 5 aryl, a cycloalkyl, a cycloalkenyl, a heterocycle or a polycycle; and m is zero or an integer in the range of 1 to 8. In other embodiments, R₅₀ and R₅₁ (and optionally R₅₂) each independently represent a hydrogen, an alkyl, an alkenyl, or -(CH₂)_m-R₆₁. Thus, the term "alkylamine" includes an amine group, as defined above, having a substituted or unsubstituted alkyl attached thereto, i.e., at least one of R₅₀ and R₅₁ is an alkyl group.

10 The term "acylamino" is art-recognized and refers to a moiety that may be represented by the general formula:



wherein R₅₀ is as defined above, and R₅₄ represents a hydrogen, an alkyl, an alkenyl or -(CH₂)_m-R₆₁, where m and R₆₁ are as defined above.

15 The term "amido" is art recognized as an amino-substituted carbonyl and includes a moiety that may be represented by the general formula:




wherein R₅₀ and R₅₁ are as defined above. Certain embodiments of the amide in the present invention will not include imides which may be unstable.

20 The term "alkylthio" refers to an alkyl group, as defined above, having a sulfur radical attached thereto. In certain embodiments, the "alkylthio" moiety is represented by

one of -S-alkyl, -S-alkenyl, -S-alkynyl, and -S-(CH₂)_m-R61, wherein m and R61 are defined above. Representative alkylthio groups include methylthio, ethyl thio, and the like.

The term "carboxyl" is art recognized and includes such moieties as may be represented by the general formulas:

5




wherein X50 is a bond or represents an oxygen or a sulfur, and R55 and R56 represents a hydrogen, an alkyl, an alkenyl, -(CH₂)_m-R61 or a pharmaceutically acceptable salt, R56 represents a hydrogen, an alkyl, an alkenyl or -(CH₂)_m-R61, where m and R61 are defined above. Where X50 is an oxygen and R55 or R56 is not hydrogen, the formula represents an ester. Where X50 is an oxygen, and R55 is as defined above, the moiety is referred to herein as a carboxyl group, and particularly when R55 is a hydrogen, the formula represents a carboxylic acid. Where X50 is an oxygen, and R56 is hydrogen, the formula represents a formate. In general, where the oxygen atom of the above formula is replaced by sulfur, the formula represents a thiocarbonyl group. Where X50 is a sulfur and R55 or R56 is not hydrogen, the formula represents a thioester. Where X50 is a sulfur and R55 is hydrogen, the formula represents a thiocarboxylic acid. Where X50 is a sulfur and R56 is hydrogen, the formula represents a thiocarbonate. On the other hand, where X50 is a bond, and R55 is not hydrogen, the above formula represents a ketone group. Where X50 is a bond, and R55 is hydrogen, the above formula represents an aldehyde group.

20 The term "carbamoyl" refers to -O(C=O)NRR', where R and R' are independently H, aliphatic groups, aryl groups or heteroaryl groups.

The term "oxo" refers to a carbonyl oxygen (=O).

The terms "oxime" and "oxime ether" are art-recognized and refer to moieties that may be represented by the general formula:


25



wherein R75 is hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, or $-(CH_2)_m-R61$. The moiety is an “oxime” when R is H; and it is an “oxime ether” when R is alkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, or $-(CH_2)_m-R61$.


The terms “alkoxyl” or “alkoxy” are art-recognized and refer to an alkyl group, as defined above, having an oxygen radical attached thereto. Representative alkoxy groups include methoxy, ethoxy, propyloxy, tert-butoxy and the like. An “ether” is two hydrocarbons covalently linked by an oxygen. Accordingly, the substituent of an alkyl that renders that alkyl an ether is or resembles an alkoxy, such as may be represented by one of $-O\text{-alkyl}$, $-O\text{-alkenyl}$, $-O\text{-alkynyl}$, $-O\text{--}(CH_2)_m\text{-R61}$, where m and R61 are described above.

10 The term “sulfonate” is art recognized and refers to a moiety that may be represented by the general formula:




in which R57 is an electron pair, hydrogen, alkyl, cycloalkyl, or aryl.

15 The term “sulfate” is art recognized and includes a moiety that may be represented by the general formula:



in which R57 is as defined above.


The term “sulfonamido” is art recognized and includes a moiety that may be represented by the general formula:



20


in which R50 and R56 are as defined above.

The term “sulfamoyl” is art-recognized and refers to a moiety that may be represented by the general formula:




in which R50 and R51 are as defined above.

- 5 The term “sulfonyl” is art-recognized and refers to a moiety that may be represented by the general formula:




in which R58 is one of the following: hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl or heteroaryl.

- 10 The term “sulfoxido” is art-recognized and refers to a moiety that may be represented by the general formula:




in which R58 is defined above.

- 15 The term “phosphoryl” is art-recognized and may in general be represented by the formula:




wherein Q50 represents S or O, and R59 represents hydrogen, a lower alkyl or an aryl.

When used to substitute, e.g., an alkyl, the phosphoryl group of the phosphorylalkyl may be represented by the general formulas:



wherein Q50 and R59, each independently, are defined above, and Q51 represents O, S or N. When Q50 is S, the phosphoryl moiety is a “phosphorothioate”.


5 The term “phosphoramidite” is art-recognized and may be represented in the general formulas:



wherein Q51, R50, R51 and R59 are as defined above.

The term “phosphonamidite” is art-recognized and may be represented in the general formulas:

10



wherein Q51, R50, R51 and R59 are as defined above, and R60 represents a lower alkyl or an aryl.

15

Analogous substitutions may be made to alkenyl and alkynyl groups to produce, for example, aminoalkenyls, aminoalkynyls, amidoalkenyls, amidoalkynyls, iminoalkenyls, iminoalkynyls, thioalkenyls, thioalkynyls, carbonyl-substituted alkenyls or alkynyls.

The definition of each expression, e.g., alkyl, m, n, and the like, when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

20 The terms triflyl, tosyl, mesyl, and nonaflyl are art-recognized and refer to trifluoromethanesulfonyl, *p*-toluenesulfonyl, methanesulfonyl, and

nonafluorobutanesulfonyl groups, respectively. The terms triflate, tosylate, mesylate, and nonaflate are art-recognized and refer to trifluoromethanesulfonate ester, *p*-toluenesulfonate ester, methanesulfonate ester, and nonafluorobutanesulfonate ester functional groups and molecules that contain said groups, respectively.

5 The abbreviations Me, Et, Ph, Tf, Nf, Ts, and Ms represent methyl, ethyl, phenyl, trifluoromethanesulfonyl, nonafluorobutanesulfonyl, *p*-toluenesulfonyl and methanesulfonyl, respectively. A more comprehensive list of the abbreviations utilized by organic chemists of ordinary skill in the art appears in the first issue of each volume of the Journal of Organic Chemistry; this list is typically presented in a table entitled Standard List
10 of Abbreviations.

Certain compounds contained in compositions of the present invention may exist in particular geometric or stereoisomeric forms. In addition, polymers of the present invention may also be optically active. The present invention contemplates all such compounds, including cis- and trans-isomers, *R*- and *S*-enantiomers, diastereomers, (*D*)-isomers, (*L*)-
15 isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

If, for instance, a particular enantiomer of compound of the present invention is
20 desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed
25 by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

It will be understood that "substitution" or "substituted with" includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted
30 atom and the substituent, and that the substitution results in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, or other reaction.

The term "substituted" is also contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. Illustrative substituents include, for example, those described herein above. The permissible substituents may be one or more and the same or different for appropriate organic compounds. For purposes of this invention, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valences of the heteroatoms. This invention is not intended to be limited in any manner by the permissible substituents of organic compounds.

The phrase "protecting group" as used herein means temporary substituents which protect a potentially reactive functional group from undesired chemical transformations. Examples of such protecting groups include esters of carboxylic acids, silyl ethers of alcohols, and acetals and ketals of aldehydes and ketones, respectively. Examples of nitrogen protecting groups include an amide (-NRC(=O)R) or a urethane (-NRC(=O)OR), for example, as: a methyl amide (-NHC(=O)CH₃); a benzyloxy amide (-NHC(=O)OCH₂C₆H₅NHCbz); as a t-butoxy amide (-NHC(=O)OC(CH₃)₃, -NHBOC); a 2-biphenyl-2-propoxy amide (-NHC(=O)OC(CH₃)₂C₆H₄C₆H₅NHBoc), as a 9-fluorenylmethoxy amide (-NHFmoc), as a 6-nitroveratryloxy amide (-NHNvoc), as a 2-trimethylsilylethoxy amide (-NHTeoc), as a 2,2,2-trichloroethoxy amide (-NHTroc), as an allyloxy amide (-NHALloc), as a 2-(phenylsulfonyl)ethoxy amide (-NHPsec); or, in suitable cases (e.g., cyclic amines), as a nitroxide radical. The field of protecting group chemistry has been reviewed (Greene, T.W.; Wuts, P.G.M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991). Protected forms of the inventive compounds are included within the scope of this invention.

For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 67th Ed., 1986-87, inside cover.

The term "pharmaceutically acceptable salt" or "salt" refers to a salt of one or more compounds. Suitable pharmaceutically acceptable salts of compounds include acid addition salts, such as those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and also those formed with organic acids such as maleic acid. For example, acids

commonly employed to form pharmaceutically acceptable salts include inorganic acids such as hydrogen bisulfide, hydrochloric, hydrobromic, hydroiodic, sulfuric and phosphoric acid, as well as organic acids such as para-toluenesulfonic, salicylic, tartaric, bitartaric, ascorbic, maleic, besylic, fumaric, gluconic, glucuronic, formic, glutamic, methanesulfonic, 5 ethanesulfonic, benzenesulfonic, lactic, oxalic, para-bromophenylsulfonic, carbonic, succinic, citric, benzoic and acetic acid, and related inorganic and organic acids. Such pharmaceutically acceptable salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, 10 acrylate, formate, isobutyrate, caprate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, terephthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, β-hydroxybutyrate, glycolate, maleate, tartrate, 15 methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like.

Where the compounds carry one or more acidic moieties, pharmaceutically acceptable salts may be formed by treatment of a solution of the compound with a solution of a pharmaceutically acceptable base. Suitable bases for forming pharmaceutically acceptable salts with acidic functional groups include, but are not limited to, hydroxides and carbonates of alkali metals such as sodium, potassium, and lithium; alkaline earth metal such as calcium and magnesium; and other metals, such as aluminum and zinc. Suitable bases also include ammonia, and organic amines, such as unsubstituted or hydroxy-substituted mono-, di-, or trialkylamines; dicyclohexylamine; tributyl amine; pyridine; N-20 methyl,N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-hydroxy-lower alkyl amines), such as mono-, bis-, or tris-(2-hydroxyethyl)amine, 2-hydroxy-tert-butylamine, or tris-(hydroxymethyl)methylamine, N,N-di alkyl-N-(hydroxy alkyl)-amines, such as N,N-dimethyl-N-(2-hydroxyethyl)amine, or tri-(2-hydroxyethyl)amine; N-methyl-D-glucamine; and amino acids such as arginine, lysine, and the like.

30 Certain compounds of the invention and their salts may exist in more than one crystalline form (i.e., polymorph); the present invention includes each of the crystal forms and mixtures thereof.

Certain compounds of the invention and their salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate and mixtures thereof.

Certain compounds of the invention may contain one or more chiral centers, and 5 exist in different optically active forms. When compounds of the invention contain one chiral center, the compounds exist in two enantiomeric forms and the present invention includes both enantiomers and mixtures of enantiomers, such as racemic mixtures thereof. The enantiomers may be resolved by methods known to those skilled in the art; for example, enantiomers may be resolved by formation of diastereoisomeric salts which may 10 be separated, for example, by crystallization; formation of diastereoisomeric derivatives or complexes which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example, via enzymatic esterification; or gas-liquid or liquid chromatography in a chiral environment, for example, on a chiral support; suitable include chiral supports (e.g., silica 15 with a bound chiral ligand) or in the presence of a chiral solvent. Where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be used to liberate the desired purified enantiomer. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using 20 optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation.

When a compound of the invention contains more than one chiral center, it may exist in diastereoisomeric forms. The diastereoisomeric compounds may be separated by methods known to those skilled in the art (for example, chromatography or crystallization) and the individual enantiomers may be separated as described above. The present invention 25 includes the various diastereoisomers of compounds of the invention, and mixtures thereof. Compounds of the invention may exist in different tautomeric forms or as different geometric isomers, and the present invention includes each tautomer and/or geometric isomer of compounds of the invention, and mixtures thereof. For example, any olefins present in the compounds may exist as either the E- or Z- geometric isomers, or a mixture 30 thereof unless stated otherwise. Compounds of the invention may exist in zwitterionic form. The present invention includes each zwitterionic form of compounds of the invention, and mixtures thereof.

As used herein the term "pro-drug" refers to an agent which is converted into the parent drug in vivo by some physiological chemical process (e.g., a prodrug on being brought to the physiological pH is converted to the desired drug form). Pro-drugs are often useful because, in some situations, they may be easier to administer than the parent drug.

5 They may, for instance, be bioavailable by oral administration whereas the parent drug is not. The prodrug may also have improved solubility in pharmacological compositions over the parent drug. An example, without limitation, of a pro-drug would be a compound of the present invention wherein it is administered as an ester (the "pro-drug") to facilitate transmittal across a cell membrane where water solubility is not beneficial, but then it is

10 metabolically hydrolyzed to the carboxylic acid once inside the cell where water solubility is beneficial. Pro-drugs have many useful properties. For example, a pro-drug may be more water soluble than the ultimate drug, thereby facilitating intravenous administration of the drug. A pro-drug may also have a higher level of oral bioavailability than the ultimate drug. After administration, the prodrug is enzymatically or chemically cleaved to deliver

15 the ultimate drug in the blood or tissue.

Exemplary pro-drugs release an amine of a compound of the invention wherein the free hydrogen of an amine or alcohol is replaced by (C₁-C₆)alkanoyloxymethyl, 1-((C₁-C₆)alkanoyloxy)ethyl, 1-methyl-1-((C₁-C₆)alkanoyloxy)ethyl, (C₁-C₆)alkoxycarbonyloxymethyl, N-(C₁-C₆)alkoxycarbonylamino-methyl, succinoyl, (C₁-C₆)alkanoyl, α-

20 amino(C₁-C₄)alkanoyl, arylactyl and α-aminoacyl, or α-aminoacyl-α-aminoacyl wherein said α-aminoacyl moieties are independently any of the naturally occurring L-amino acids found in proteins, -P(O)(OH)₂, -P(O)(O(C₁-C₆)alkyl)₂ or glycosyl (the radical resulting from detachment of the hydroxyl of the hemiacetal of a carbohydrate).

Other exemplary pro-drugs upon cleavage release a corresponding free acid, and

25 such hydrolyzable ester-forming residues of the compounds of this invention include but are not limited to carboxylic acid substituents (e.g., -(CH₂)C(O)OH or a moiety that contains a carboxylic acid) wherein the free hydrogen is replaced by (C₁-C₄)alkyl, (C₂-C₁₂)alkanoyloxymethyl, (C₄-C₉)1-(alkanoyloxy)ethyl, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-

30 (alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-

(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁-C₂)alkylamino(C₂-C₃)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C₁-C₂)alkyl, N,N-di(C₁-C₂)-alkylcarbamoyl-(C₁-C₂)alkyl and piperidino-, pyrrolidino- or morpholino(C₂-C₃)alkyl.

5 The term "subject" as used herein, refers to an animal, typically a mammal or a human, that will be or has been the object of treatment, observation, and/or experiment. When the term is used in conjunction with administration of a compound or drug, then the subject has been the object of treatment, observation, and/or administration of the compound or drug.

10 The terms "co-administration" and "co-administering" refer to both concurrent administration (administration of two or more therapeutic agents at the same time) and time varied administration (administration of one or more therapeutic agents at a time different from that of the administration of an additional therapeutic agent or agents), as long as the therapeutic agents are present in the patient to some extent at the same time.


15 The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits a biological or medicinal response in a cell culture, tissue system, animal, or human that is being sought by a researcher, veterinarian, clinician, or physician, which includes alleviation of the symptoms of the disease, condition, or disorder being treated.

20 The term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product that results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

25 The term "pharmaceutically acceptable carrier" refers to a medium that is used to prepare a desired dosage form of a compound. A pharmaceutically acceptable carrier can include one or more solvents, diluents, or other liquid vehicles; dispersion or suspension aids; surface active agents; isotonic agents; thickening or emulsifying agents; preservatives; solid binders; lubricants; and the like. Remington's Pharmaceutical Sciences, Fifteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1975) and Handbook of Pharmaceutical Excipients, Third Edition, A. H. Kibbe ed. (American Pharmaceutical Assoc. 2000), disclose various carriers used in formulating pharmaceutical compositions and known techniques for the preparation thereof.

COMPOUNDS

One embodiment of the invention relates to a compound of formula 1:



or a pharmaceutically acceptable salt thereof, wherein independently for each occurrence:


- 5 W and X are independently oxygen or sulfur;
- Z¹, Z² and Z³ are independently C-R²¹ or N, provided that at least one of Z¹ and Z² is N;
- R¹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aralkyl, heteroaralkyl, -COR⁶, and -C(O)OR⁶;
- R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷);
- R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷); or R² and R³ are joined together to form an optionally substituted heterocyclic ring;
- R⁴ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocyclylalkyl, aralkyl, heteroaryl, heteroaralkyl, halo, hydroxy, alkoxy, hydroxyalkyl, and alkoxyalkyl;
- R⁵ is selected from the group consisting of aryl, heteroaryl, heterocyclyl, -N(R⁸)(R⁹), -N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and -N(R⁸)P(O)(OR⁹)(OR⁹);
- p is 1, 2, 3, 4, 5, or 6;

- R^6 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;
- R^7 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;
- 5 R^8 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;
- R^9 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl; or R^8 and R^9 are joined together to form a heterocyclic ring;
- 10 R^{20} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, halo, haloalkyl, trifluoromethyl, fluoroalkyl, perfluoroalkyl, thio, cyano, hydroxy, methoxy, alkoxy, phenoxy, aryloxy, heteroaryloxy, carboxyl, alkoxycarbonyl, acyl, nitro, amino, alkylamino, arylamino, heteroaryl amino, amido, acylamino, sulfate, sulfonate, sulfonyl, sulfoxido, sulfonamido, 15 sulfamoyl, $-[C(R^4)_2]_p-R^5$, $NR^{14}R^{15}$, OR^{16} , $O-[C(R^4)_2]_p-R^5$, $NR^{14}-[C(R^4)_2]_p-R^5$ and SR^{16} ;
- R^{21} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, halo, haloalkyl, thio, cyano, carboxyl, alkoxycarbonyl, acyl, nitro, amino, amido, acylamino, sulfate, sulfonate, sulfonamido, sulfoxido, sulfamoyl, $NR^{14}R^{15}$, OR^{16} , and SR^{16} ;
- 20 R^{14} and R^{15} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, $-[C(R^4)_2]_p-R^5$, $-COR^6$, $-C(O)OR^6$, $-SO_2(R^6)$, $-C(O)N(R^6)(R^7)$, $-SO_2N(R^6)(R^7)$, and $-P(O)(OR^6)(OR^7)$; or R^{14} and R^{15} are joined together to form an optionally substituted heterocyclic ring;
- 25 R^{16} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, $-[C(R^4)_2]_p-R^5$, $-COR^6$, and $-C(O)N(R^6)(R^7)$; and
- r is 0 or 1;
- wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl may be optionally substituted.

- In another embodiment, W and X are oxygen.
- In another embodiment, Z¹ and Z² are nitrogen; and Z³ is C-R²¹.
- In another embodiment, Z¹, Z² and Z³ are nitrogen.
- In another embodiment, Z¹ is nitrogen; and Z² and Z³ are each C-R²¹.
- 5 In another embodiment, Z² is nitrogen; and Z¹ and Z³ are each C-R²¹.
- In another embodiment, R¹ is hydrogen, alkyl, or aralkyl.
- In another embodiment, W and X are oxygen, Z¹ and Z² are each nitrogen, Z³ is C-R²¹, and R¹ is hydrogen, alkyl, or aralkyl.
- In another embodiment, R¹ is hydrogen.
- 10 In another embodiment, R² and R³ are joined together to form an optionally substituted heterocyclic ring.
- In another embodiment, the optionally substituted heterocyclic ring is selected from the group consisting of piperazinyl, homopiperizinyl, pyrrolidinyl, piperidinyl, homopiperidinyl, morpholinyl, and quinolinyl.
- 15 In another embodiment, R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), and R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl may be optionally substituted.
- 20 In another embodiment, R² is -[C(R⁴)₂]p-R⁵, and R³ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl may be optionally substituted.
- 25 In another embodiment, R⁵ is aryl or heteroaryl, each of which may be optionally substituted.
- In another embodiment, R⁵ is -N(R⁸)(R⁹).
- In another embodiment, R⁴ is hydrogen.

In another embodiment, r is 0.

Another aspect of the invention relates to a compound of formula 2:



- 5 or a pharmaceutically acceptable salt thereof, wherein independently for each occurrence:
 - R¹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aralkyl, heteroaralkyl, -COR⁶, and -C(O)OR⁶;
 - R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶,
- 10 10 -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷);
- R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷); or R² and R³ are joined together to form an optionally substituted heterocyclic ring;
- 15 R⁴ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocyclylalkyl, aralkyl, heteroaryl, heteroaralkyl, halo, hydroxy, alkoxy, hydroxyalkyl, and alkoxyalkyl;
- R⁵ is selected from the group consisting of aryl, heteroaryl, heterocyclyl, -N(R⁸)(R⁹), -N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -
- 20 20 SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and -N(R⁸)P(O)(OR⁹)(OR⁹);
- p is 1, 2, 3, 4, 5, or 6;
- R⁶ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;

R^7 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl;

R^8 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl;

- 5 R^9 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl; or R^8 and R^9 are joined together to form a heterocyclic ring; and

R^{20} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclalkyl, halo, haloalkyl, 10 trifluoromethyl, fluoroalkyl, perfluoroalkyl, thio, cyano, hydroxy, methoxy, alkoxy, phenoxy, aryloxy, heteroaryloxy, carboxyl, alkoxycarbonyl, acyl, nitro, amino, alkylamino, arylamino, heteroaryl amino, amido, acylamino, sulfate, sulfonate, sulfonyl, sulfoxido, sulfonamido, sulfamoyl, $-[C(R^4)_2]_p-R^5$, $NR^{14}R^{15}$, OR^{16} , $O-[C(R^4)_2]_p-R^5$, $NR^{14}-[C(R^4)_2]_p-R^5$ and SR^{16} ;

15


wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl may be optionally substituted.

In another embodiment, R^1 is hydrogen, alkyl, or aralkyl.

In another embodiment, R^1 is hydrogen, methyl, or benzyl.

- 20 In another embodiment, R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring.

In another embodiment, R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring selected from the group consisting of :



- 25 wherein, independently for each occurrence:

R^{10} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclalkyl, $-[C(R^4)_2]_p-R^5$, $-COR^6$, $-C(O)OR^6$, $-SO_2(R^6)$, $-C(O)N(R^6)(R^7)$, $-SO_2N(R^6)(R^7)$; and $-P(O)(OR^6)(OR^7)$;


R^{11} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, halo, haloalkyl, thio, cyano, alkylthio, nitro, $-N(R^8)(R^9)$, $-N(R^8)COR^9$, $-N(R^8)C(O)OR^9$, $-N(R^8)SO_2(R^9)$, $-CON(R^8)(R^9)$, $-OC(O)N(R^8)-(R^9)$, $-SO_2N(R^8)(R^9)$, $-OC(O)OR^8$, $-COOR^9$, $-C(O)N(OH)(R^8)$, $-OS(O)_2OR^8$, $-S(O)_2OR^8$, $-S(O)_2R^8$, $-OR^8$, $-COR^8$, $-OP(O)(OR^8)(OR^8)$, $-P(O)(OR^8)(OR^8)$, $-N(R^8)P(O)(OR^9)(OR^9)$ and $-[C(R^4)_2]_p-R^5$;

and $-[C(R^4)_2]_p-R^5$; and

10 n is 0, 1, 2, or 3;

wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclalkyl may be optionally substituted.


In another embodiment, R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring of the formula:



In another embodiment,, wherein n is 0.

In another embodiment, R^{10} is hydrogen, alkyl, aryl, heteroaryl, $-COR^6$, $-C(O)OR^6$, or $-SO_2(R^6)$, or $-[C(R^4)_2]_p-R^5$.

20 In another embodiment, R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring of the formula:



In another embodiment, n is 0 or 1.

In another embodiment, R^{11} is alkyl, aryl, heteroaryl, heterocyclyl, $-N(R^8)(R^9)$, $-N(R^8)COR^9$, $-N(R^8)C(O)OR^9$, $-N(R^8)SO_2(R^9)$ and $-[C(R^4)_2]_p-R^5$.

In another embodiment, R¹¹ is -[C(R⁴)₂]_p-R⁵, wherein p is 1 or 2, R⁴ is hydrogen, and R⁵ is selected from the group consisting of aryl, heteroaryl, heterocyclyl, -N(R⁸)(R⁹), -N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, and -N(R⁸)SO₂(R⁹).

5 In another embodiment, R² and R³ are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -[C(R⁴)₂]_p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl may be optionally substituted.

10 In another embodiment, R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷); and R³ is -[C(R⁴)₂]_p-R⁵.

In another embodiment, R² is optionally substituted alkyl.

In another embodiment, R³ is aryl, heteroaryl, heterocyclyl, each of which may be optionally substituted.

In another embodiment, at least one of R² and R³ is -[C(R⁴)₂]_p-R⁵.

15 In another embodiment, R⁴ is hydrogen.


In another embodiment, R⁵ is selected from the group consisting of aryl, heteroaryl, -N(R⁸)(R⁹), -N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and 20 -N(R⁸)P(O)(OR⁹)(OR⁹).


In another embodiment, R⁵ is -N(R⁸)(R⁹).


In another embodiment, R⁵ is aryl or heteroaryl.


In another embodiment, R²⁰ is hydrogen.


Another aspect of the invention relates to a compound, or a pharmaceutically acceptable salt thereof, selected from the group consisting of:













Any one of the aforementioned compounds may exist as the *E*-geometric isomer, the *Z*-geometric isomer, or mixtures thereof. For example, in one embodiment, " " in the aforementioned structures represents the *E*-isomer of the particular compound. In 5 another embodiment, " " represents the *Z*-isomer of the particular compound. In yet another embodiment, " " represents a mixture of *E* and *Z* isomers of the particular compound.

In another embodiment, any one of the aforementioned compounds is an inhibitor of CK1 γ 1, CK1 γ 2, or CK1 γ 3.

10 In another embodiment, any one of the aforementioned compounds is an inhibitor of the Wnt pathway.

In another embodiment, any one of the aforementioned compounds is an inhibitor of the TGF β pathway.


15 In some embodiments, the compound has an IC₅₀ of less than 5000 nM for CK1 γ 1, CK1 γ 2, or CK1 γ 3.

In some embodiments, the compound has an IC₅₀ of less than 1000 nM for CK1 γ 1, CK1 γ 2, or CK1 γ 3.

In some embodiments, the compound has an IC₅₀ of less than 500 nM for CK1 γ 1, CK1 γ 2, or CK1 γ 3.


20 GENERAL SYNTHETIC SCHEMES

The general synthetic schemes that were utilized to prepare compounds disclosed in this application are described below. For example, compounds of the invention may be prepared as shown in Scheme I:




Scheme I.

Alternatively, the compounds of the present disclosure can be made as shown in
5 Scheme II:




Scheme II.

Yet another method of making the compounds disclosed herein is depicted in Scheme III:



Scheme III.

Another method of making the compounds disclosed herein is depicted in Scheme IV:




5

Scheme IV.

PROPHETIC EMBODIMENTS

10 Certain compounds of the invention could be made in accordance with the above schemes by reacting an amine (Reactant A) with the hydantoin core (Reactant B). Non-limiting prophetic examples of Reactant A and Reactant B are shown in Table 1 and Table 2, respectively.

Table 1: Reactant A Prophetic Examples.


Reactant A # 1
Structure

Molecular Weight

162.232


Molecular Formula
C₁₀H₁₄N₂
Chemical name

1-phenylpiperazine


Reactant A # 2




Molecular Weight
163.22
Molecular Formula
 $C_9H_{13}N_3$
Chemical name
1-(pyridin-3-yl)piperazine




Molecular Weight
164.208
Molecular Formula
 $C_8H_{12}N_4$
Chemical name
5-(piperazin-1-yl)pyrimidine




Molecular Weight
164.208
Molecular Formula
 $C_8H_{12}N_4$
Chemical name
2-(piperazin-1-yl)pyrimidine



Molecular Weight
205.256
Molecular Formula
 $C_{11}H_{15}N_3O$
Chemical name
N-phenylpiperazine-1-carboxamide




Molecular Weight
197.32
Molecular Formula
 $C_{11}H_{23}N_3$
Chemical name
1-(1-ethylpiperidin-4-yl)piperazine



Molecular Weight
177.246
Molecular Formula
 $C_{10}H_{15}N_3$
Chemical name
1-(pyridin-4-yl)-1,4-diazepane


Reactant A # 8

Structure**Molecular Weight**

217.267

Molecular FormulaC₁₂H₁₅N₃O**Chemical name**


2-(1,4-diazepan-1-yl)benzo[d]oxazole

Reactant A # 9**Structure****Molecular Weight**

219.283

Molecular FormulaC₁₂H₁₇N₃O**Chemical name**


N-phenyl-1,4-diazepane-1-carboxamide

Reactant A # 10**Structure****Molecular Weight**

261.366

Molecular FormulaC₁₄H₂₃N₅**Chemical name**

1-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepane

Reactant A # 11**Structure****Molecular Weight**


130.231

Molecular FormulaC₇H₁₈N₂**Chemical name**

N1,N1-diethyl-N2-methylethane-1,2-diamine

Reactant A # 12

Structure



Molecular Weight

251.305

Molecular Formula


 $C_{11}H_{13}N_3O_2S$

Chemical name

N-(2-aminoethyl)isoquinoline-5-sulfonamide

Reactant A # 13

Structure



Molecular Weight

164.204

Molecular Formula


 $C_9H_{12}N_2O$

Chemical name

N-(2-aminoethyl)benzamide

Reactant A # 14

Structure



Molecular Weight

189.214

Molecular Formula


 $C_{10}H_{11}N_3O$

Chemical name

2-(2-aminoethyl)quinazolin-4(3H)-one

Reactant A # 15

Structure



Molecular Weight

190.242

Molecular Formula


 $C_{11}H_{14}N_2O$

Chemical name


phenyl(piperazin-1-yl)methanone

Reactant A # 16

Structure




Molecular Weight 226.295
 Molecular Formula C₁₀H₁₄N₂O₂S
 Chemical name 1-(phenylsulfonyl)piperazine
Reactant A # 17
 Structure




Molecular Weight 216.279
 Molecular Formula C₁₃H₁₆N₂O
 Chemical name (hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(phenyl)methanone
Reactant A # 18

Structure




Molecular Weight 252.333
 Molecular Formula C₁₂H₁₆N₂O₂S
 Chemical name 2-(phenylsulfonyl)octahydropyrrolo[3,4-c]pyrrole
Reactant A # 19

Structure




Molecular Weight 231.294
 Molecular Formula C₁₃H₁₇N₃O
 Chemical name N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide
Reactant A # 20

Structure




Molecular Weight 217.267
 Molecular Formula C₁₂H₁₅N₃O
 Chemical name N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide
Reactant A # 21

Structure**Molecular Weight**

202.252

Molecular Formula $C_{12}H_{14}N_2O$ **Chemical name**

phenyl(2,6-diazaspiro[3.3]heptan-2-yl)methanone


Reactant A # 22**Structure****Molecular Weight**

238.306

Molecular Formula $C_{11}H_{14}N_2O_2S$ **Chemical name**

2-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptane


Table 2: Reactant B Prophetic Examples.

Reactant B # 1**Structure****Molecular Weight**

309.708

Molecular Formula $C_{12}H_{12}ClN_5O_3$ **Chemical name**


(Z)-5-((2-chloro-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione

Reactant B # 2**Structure****Molecular Weight**

322.75

Molecular Formula $C_{13}H_{15}ClN_6O_2$ **Chemical name**


(Z)-5-((2-chloro-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

Reactant B # 3*Structure**Molecular Weight*

343.768

Molecular Formula $C_{16}H_{14}ClN_5O_2$ *Chemical_name*


(Z)-5-((6-(benzyl(methyl)amino)-2-chloropyrimidin-4-yl)methylene)imidazolidine-2,4-dione

Reactant B # 4*Structure**Molecular Weight*

297.698

Molecular Formula $C_{11}H_{12}ClN_5O_3$ *Chemical_name*


(Z)-5-((2-chloro-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

Reactant B # 5*Structure**Molecular Weight*

329.741





Molecular Formula $C_{15}H_{12}ClN_5O_2$ *Chemical_name*

(Z)-5-((2-chloro-6-(methyl(phenyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione


Reactant B # 6*Structure**Molecular Weight*

320.688


Molecular Formula C₁₃H₉ClN₄O₄
Chemical_name (Z)-5-((2-chloro-6-(furan-2-ylmethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

Reactant B # 7*Structure**Molecular Weight* 316.699*Molecular Formula* C₁₄H₉ClN₄O₃*Chemical_name* (Z)-5-((2-chloro-6-phenoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione**Reactant B # 8***Structure**Molecular Weight* 330.726*Molecular Formula* C₁₅H₁₁ClN₄O₃*Chemical_name* (Z)-5-((6-(benzyloxy)-2-chloropyrimidin-4-yl)methylene)imidazolidine-2,4-dione**Reactant B # 9***Structure**Molecular Weight* 298.682*Molecular Formula* C₁₁H₁₁ClN₄O₄*Chemical_name* (Z)-5-((2-chloro-6-(2-methoxyethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione**Reactant B # 10***Structure*

Molecular Weight 311.724
Molecular Formula C₁₂H₁₄ClN₅O₃
Chemical_name (Z)-5-((2-chloro-6-(2-(dimethylamino)ethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

Reactant B # 11*Structure*

Molecular Weight 224.604
Molecular Formula C₈H₅ClN₄O₂
Chemical_name (Z)-5-((2-chloropyrimidin-4-yl)methylene)imidazolidine-2,4-dione

Reactant B # 12*Structure*

Molecular Weight 225.592
Molecular Formula C₇H₄ClN₅O₂
Chemical_name (Z)-5-((4-chloro-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione

Additional prophetic embodiments of the invention that may be made in accordance with the above reaction schemes using Reactants A and B are listed in Table 3. The 5 geometric isomers listed in Table 3 are believed to reflect the actual geometry of the prophetic compounds if they were to be made; however, final structural assignments may only be made if the compounds are synthesized and subjected to appropriate 2D NMR experiments. Further, although the compounds are listed as the "Z" geometric isomer, both the E and Z geometric isomers and mixtures thereof are contemplated.

10 Table 3: Additional prophetic embodiments of the invention.

No	Chemical Name	Formula	Mol Weight	Reactant	
				A	B
1	(Z)-5-((6-morpholino-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C ₂₂ H ₂₅ N ₇ O ₃	435.479	1	1
2	(Z)-5-((6-morpholino-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C ₂₁ H ₂₄ N ₈ O ₃	436.467	2	1

3	(Z)-5-((6-morpholino-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H23N9O3	437.455	3	1
4	(Z)-5-((6-morpholino-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H23N9O3	437.455	4	1
5	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-morpholinopyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C23H26N8O4	478.504	5	1
6	(Z)-5-((2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H34N8O3	470.568	6	1
7	(Z)-5-((6-morpholino-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H26N8O3	450.494	7	1
8	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H26N8O4	490.514	8	1
9	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-morpholinopyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C24H28N8O4	492.53	9	1
10	(Z)-5-((2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H34N10O3	534.613	10	1
11	(Z)-5-((2-(diethylaminoethyl)(methyl)amino)-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C19H29N7O3	403.479	11	1
12	(Z)-N-(2-((4-(2,4-dioxoimidazolidin-5-ylidene)methyl)-6-morpholinopyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C23H24N8O5S	524.552	12	1
13	(Z)-N-(2-((4-(2,4-dioxoimidazolidin-5-ylidene)methyl)-6-morpholinopyrimidin-2-yl)amino)ethyl)benzamide	C21H23N7O4	437.452	13	1
14	(Z)-5-((6-morpholino-2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H22N8O4	462.461	14	1
15	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H25N7O4	463.489	15	1
16	(Z)-5-((6-morpholino-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H25N7O5S	499.543	16	1
17	(Z)-5-((2-(5-benzoylhexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H27N7O4	489.526	17	1
18	(Z)-5-((6-morpholino-2-(5-(phenylsulfonyl)hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H27N7O5S	525.58	18	1

19	(Z)-5-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-morpholinopyrimidin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C25H28N8O4	504.541	19	1
20	(Z)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-morpholinopyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C24H26N8O4	490.514	20	1
21	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H25N7O4	475.5	21	1
22	(Z)-5-((6-morpholino-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H25N7O5S	511.554	22	1
23	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H28N8O2	448.521	1	2
24	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H27N9O2	449.509	2	2
25	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H26N10O2	450.497	3	2
26	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H26N10O2	450.497	4	2
27	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(4-methylpiperazin-1-yl)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C24H29N9O3	491.546	5	2
28	(Z)-5-((2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H37N9O2	483.61	6	2
29	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H29N9O2	463.535	7	2
30	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H29N9O3	503.556	8	2
31	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(4-methylpiperazin-1-yl)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C25H31N9O3	505.572	9	2
32	(Z)-5-((2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H37N11O2	547.655	10	2
33	(Z)-5-((2-((2-(diethylamino)ethyl)(methyl)amino)-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H32N8O2	416.52	11	2

34	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(4-methylpiperazin-1-yl)pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C24H27N9O4S	537.594	12	2
35	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(4-methylpiperazin-1-yl)pyrimidin-2-yl)amino)ethyl)benzamide	C22H26N8O3	450.494	13	2
36	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-((4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H25N9O3	475.503	14	2
37	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H28N8O3	476.531	15	2
38	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H28N8O4S	512.585	16	2
39	(Z)-5-((2-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H30N8O3	502.568	17	2
40	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(5-(phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H30N8O4S	538.622	18	2
41	(Z)-5-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(4-methylpiperazin-1-yl)pyrimidin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C26H31N9O3	517.583	19	2
42	(Z)-6-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(4-methylpiperazin-1-yl)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C25H29N9O3	503.556	20	2
43	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H28N8O3	488.542	21	2
44	(Z)-5-((6-(4-methylpiperazin-1-yl)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H28N8O4S	524.595	22	2
45	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H27N7O2	469.538	1	3
46	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H26N8O2	470.526	2	3
47	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H25N9O2	471.514	3	3
48	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H25N9O2	471.514	4	3

49	(Z)-4-(4-(benzyl(methyl)amino)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C27H28N8O3	512.563	5	3
50	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H36N8O2	504.627	6	3
51	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H28N8O2	484.553	7	3
52	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-(benzyl(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C28H28N8O3	524.574	8	3
53	(Z)-4-(4-(benzyl(methyl)amino)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C28H30N8O3	526.59	9	3
54	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C30H36N10O2	568.673	10	3
55	(Z)-5-((6-(benzyl(methyl)amino)-2-((2-(diethylamino)ethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H31N7O2	437.538	11	3
56	(Z)-N-(2-((4-(benzyl(methyl)amino)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C27H26N8O4S	558.612	12	3
57	(Z)-N-(2-((4-(benzyl(methyl)amino)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)amino)ethyl)benzamide	C25H25N7O3	471.511	13	3
58	(Z)-5-((6-(benzyl(methyl)amino)-2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H24N8O3	496.521	14	3
59	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-(benzyl(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H27N7O3	497.548	15	3
60	(Z)-5-((6-(benzyl(methyl)amino)-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H27N7O4S	533.602	16	3
61	(Z)-5-((2-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-(benzyl(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C29H29N7O3	523.586	17	3
62	(Z)-5-((6-(benzyl(methyl)amino)-2-(5-(phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C28H29N7O4S	559.639	18	3
63	(Z)-5-(4-(benzyl(methyl)amino)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C29H30N8O3	538.6	19	3

64	(Z)-6-(4-(benzyl(methyl)amino)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C28H28N8O3	524.574	20	3
65	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-(benzyl(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C28H27N7O3	509.559	21	3
66	(Z)-5-((6-(benzyl(methyl)amino)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H27N7O4S	545.613	22	3
67	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H25N7O3	423.468	1	4
68	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H24N8O3	424.456	2	4
69	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C19H23N9O3	425.444	3	4
70	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C19H23N9O3	425.444	4	4
71	(Z)-4-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C22H26N8O4	466.493	5	4
72	(Z)-5-((2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H34N8O3	458.557	6	4
73	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H26N8O3	438.483	7	4
74	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H26N8O4	478.504	8	4
75	(Z)-4-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C23H28N8O4	480.52	9	4
76	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H34N10O3	522.603	10	4
77	(Z)-5-((2-((2-(diethylamino)ethyl)(methyl)amino)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C18H29N7O3	391.468	11	4
78	(Z)-N-(2-((4-(2,4-dioxoimidazolidin-5-ylidene)methyl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-2-	C22H24N8O5S	512.542	12	4

	yl)amino)ethyl)isoquinoline-5-sulfonamide				
79	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-2-yl)amino)ethyl)benzamide	C20H23N7O4	425.441	13	4
80	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H22N8O4	450.451	14	4
81	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H25N7O4	451.478	15	4
82	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(4-(phenylsulfonylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H25N7O5S	487.532	16	4
83	(Z)-5-((2-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H27N7O4	477.516	17	4
84	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(5-(phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H27N7O5S	513.569	18	4
85	(Z)-5-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C24H28N8O4	492.53	19	4
86	(Z)-6-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C23H26N8O4	478.504	20	4
87	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-((2-hydroxyethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H25N7O4	463.489	21	4
88	(Z)-5-((6-((2-hydroxyethyl)(methyl)amino)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H25N7O5S	499.543	22	4
89	(Z)-5-((6-(methyl(phenyl)amino)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H25N7O2	455.512	1	5
90	(Z)-5-((6-(methyl(phenyl)amino)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H24N8O2	456.5	2	5
91	(Z)-5-((6-(methyl(phenyl)amino)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H23N9O2	457.488	3	5

92	(Z)-5-((6-(methyl(phenyl)amino)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H23N9O2	457.488	4	5
93	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(methyl(phenyl)amino)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C26H26N8O3	498.536	5	5
94	(Z)-5-((2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-6-(methyl(phenyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H34N8O2	490.601	6	5
95	(Z)-5-((6-(methyl(phenyl)amino)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H26N8O2	470.526	7	5
96	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-(methyl(phenyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H26N8O3	510.547	8	5
97	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(methyl(phenyl)amino)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C27H28N8O3	512.563	9	5
98	(Z)-5-((6-(methyl(phenyl)amino)-2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C29H34N10O2	554.646	10	5
99	(Z)-5-((2-(diethylaminoethyl)(methyl)amino)-6-(methyl(phenyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H29N7O2	423.511	11	5
100	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(methyl(phenyl)amino)pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C26H24N8O4S	544.585	12	5
101	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(methyl(phenyl)amino)pyrimidin-2-yl)amino)ethyl)benzamide	C24H23N7O3	457.485	13	5
102	(Z)-5-((6-(methyl(phenyl)amino)-2-(2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H22N8O3	482.494	14	5
103	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-(methyl(phenyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H25N7O3	483.522	15	5
104	(Z)-5-((6-(methyl(phenyl)amino)-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H25N7O4S	519.575	16	5
105	(Z)-5-((2-(5-benzoylhexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-(methyl(phenyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C28H27N7O3	509.559	17	5

106	(Z)-5-((6-(methyl(phenyl)amino)-2-(5-(phenylsulfonyl)hexahdropyrrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H27N7O4S	545.613	18	5
107	(Z)-5-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(methyl(phenyl)amino)pyrimidin-2-yl)-N-phenylhexahdropyrrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C28H28N8O3	524.574	19	5
108	(Z)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(methyl(phenyl)amino)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C27H26N8O3	510.547	20	5
109	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-(methyl(phenyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H25N7O3	495.533	21	5
110	(Z)-5-((6-(methyl(phenyl)amino)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H25N7O4S	531.586	22	5
111	(Z)-5-((6-(furan-2-ylmethoxy)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H22N6O4	446.459	1	6
112	(Z)-5-((6-(furan-2-ylmethoxy)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H21N7O4	447.447	2	6
113	(Z)-5-((6-(furan-2-ylmethoxy)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H20N8O4	448.435	3	6
114	(Z)-5-((6-(furan-2-ylmethoxy)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H20N8O4	448.435	4	6
115	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(furan-2-ylmethoxy)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C24H23N7O5	489.483	5	6
116	(Z)-5-((2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-6-(furan-2-ylmethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H31N7O4	481.547	6	6
117	(Z)-5-((6-(furan-2-ylmethoxy)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H23N7O4	461.473	7	6
118	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-(furan-2-ylmethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H23N7O5	501.494	8	6
119	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(furan-2-ylmethoxy)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C25H25N7O5	503.51	9	6
120	(Z)-5-((6-(furan-2-ylmethoxy)-2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-	C27H31N9O4	545.593	10	6

	dione				
121	(Z)-5-((2-((2-(diethylamino)ethyl)(methyl)amino)-6-(furan-2-ylmethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H26N6O4	414.458	11	6
122	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(furan-2-ylmethoxy)pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C24H21N7O6S	535.532	12	6
123	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(furan-2-ylmethoxy)pyrimidin-2-yl)amino)ethyl)benzamide	C22H20N6O5	448.431	13	6
124	(Z)-5-((6-(furan-2-ylmethoxy)-2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H19N7O5	473.441	14	6
125	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-(furan-2-ylmethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H22N6O5	474.469	15	6
126	(Z)-5-((6-(furan-2-ylmethoxy)-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H22N6O6S	510.522	16	6
127	(Z)-5-((2-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-(furan-2-ylmethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H24N6O5	500.506	17	6
128	(Z)-5-((6-(furan-2-ylmethoxy)-2-(5-(phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H24N6O6S	536.56	18	6
129	(Z)-5-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(furan-2-ylmethoxy)pyrimidin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C26H25N7O5	515.521	19	6
130	(Z)-6-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(furan-2-ylmethoxy)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C25H23N7O5	501.494	20	6
131	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-(furan-2-ylmethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H22N6O5	486.479	21	6
132	(Z)-5-((6-(furan-2-ylmethoxy)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H22N6O6S	522.533	22	6
133	(Z)-5-((6-phenoxy-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H22N6O3	442.47	1	7
134	(Z)-5-((6-phenoxy-2-(4-pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-	C23H21N7O3	443.458	2	7

	2,4-dione				
135	(Z)-5-((6-phenoxy-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H20N8O3	444.446	3	7
136	(Z)-5-((6-phenoxy-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H20N8O3	444.446	4	7
137	(Z)-4-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-phenoxy(pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C25H23N7O4	485.495	5	7
138	(Z)-5-((2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-6-phenoxy(pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H31N7O3	477.559	6	7
139	(Z)-5-((6-phenoxy-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H23N7O3	457.485	7	7
140	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-phenoxy(pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H23N7O4	497.505	8	7
141	(Z)-4-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-phenoxy(pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C26H25N7O4	499.521	9	7
142	(Z)-5-((2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)-6-phenoxy(pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C28H31N9O3	541.604	10	7
143	(Z)-5-((2-((2-(diethylamino)ethyl)(methyl)amino)-6-phenoxy(pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H26N6O3	410.47	11	7
144	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-phenoxy(pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C25H21N7O5S	531.543	12	7
145	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-phenoxy(pyrimidin-2-yl)amino)ethyl)benzamide	C23H20N6O4	444.443	13	7
146	(Z)-5-((2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)-6-phenoxy(pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H19N7O4	469.452	14	7
147	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-phenoxy(pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H22N6O4	470.48	15	7
148	(Z)-5-((6-phenoxy-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H22N6O5S	506.534	16	7
149	(Z)-5-((2-(5-benzoylhexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-phenoxy(pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H24N6O4	496.517	17	7
150	(Z)-5-((6-phenoxy-2-(5-(phenylsulfonyl)hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H24N6O5S	532.571	18	7

151	(Z)-5-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-phenoxy pyrimidin-2-yl)-N-phenylhexahydro pyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C27H25N7O4	511.532	19	7	
152	(Z)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-phenoxy pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C26H23N7O4	497.505	20	7	
153	(Z)-5-((2-(6-benzoyl)-2,6-diazaspiro[3.3]heptan-2-yl)-6-phenoxy pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H22N6O4	482.491	21	7	
154	(Z)-5-((6-phenoxy-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H22N6O5S	518.544	22	7	
155	(Z)-5-((6-(benzyloxy)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H24N6O3	456.496	1	8	
156	(Z)-5-((6-(benzyloxy)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H23N7O3	457.485	2	8	
157	(Z)-5-((6-(benzyloxy)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H22N8O3	458.473	3	8	
158	(Z)-5-((6-(benzyloxy)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H22N8O3	458.473	4	8	
159	(Z)-4-(4-(benzyloxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C26H25N7O4	499.521	5	8	
160	(Z)-5-((6-(benzyloxy)-2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H33N7O3	491.585	6	8	
161	(Z)-5-((6-(benzyloxy)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H25N7O3	471.511	7	8	
162	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-(benzyloxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H25N7O4	511.532	8	8	
163	(Z)-4-(4-(benzyloxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C27H27N7O4	513.548	9	8	
164	(Z)-5-((6-(benzyloxy)-2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C29H33N9O3	555.631	10	8	
165	(Z)-5-((6-(benzyloxy)-2-((2-(diethylamino)ethyl)(methyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H28N6O3	424.496	11	8	
166	(Z)-N-(2-((4-(benzyloxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C26H23N7O5S	545.57	12	8	
167	(Z)-N-(2-((4-(benzyloxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-	C24H22N6O4	458.469	13	8	

	2-yl)amino)ethyl)benzamide				
168	(Z)-5-((6-(benzyloxy)-2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H21N7O4	483.479	14	8
169	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-(benzyloxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H24N6O4	484.507	15	8
170	(Z)-5-((6-(benzyloxy)-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H24N6O5S	520.56	16	8
171	(Z)-5-((2-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-(benzyloxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C28H26N6O4	510.544	17	8
172	(Z)-5-((6-(benzyloxy)-2-(5-(phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H26N6O5S	546.598	18	8
173	(Z)-5-(4-(benzyloxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C28H27N7O4	525.558	19	8
174	(Z)-6-(4-(benzyloxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C27H25N7O4	511.532	20	8
175	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-(benzyloxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C27H24N6O4	496.517	21	8
176	(Z)-5-((6-(benzyloxy)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H24N6O5S	532.571	22	8
177	(Z)-5-((6-(2-methoxyethoxy)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H24N6O4	424.453	1	9
178	(Z)-5-((6-(2-methoxyethoxy)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H23N7O4	425.441	2	9
179	(Z)-5-((6-(2-methoxyethoxy)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C19H22N8O4	426.429	3	9
180	(Z)-5-((6-(2-methoxyethoxy)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C19H22N8O4	426.429	4	9
181	(Z)-4-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(2-methoxyethoxy)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C22H25N7O5	467.478	5	9
182	(Z)-5-((2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-6-(2-methoxyethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H33N7O4	459.542	6	9
183	(Z)-5-((6-(2-methoxyethoxy)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H25N7O4	439.468	7	9

184	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-(2-methoxyethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H25N7O5	479.489	8	9
185	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(2-methoxyethoxy)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C23H27N7O5	481.504	9	9
186	(Z)-5-((6-(2-methoxyethoxy)-2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H33N9O4	523.587	10	9
187	(Z)-5-((2-((2-(diethylamino)ethyl)(methyl)amino)-6-(2-methoxyethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C18H28N6O4	392.453	11	9
188	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(2-methoxyethoxy)pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C22H23N7O6S	513.526	12	9
189	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(2-methoxyethoxy)pyrimidin-2-yl)amino)ethyl)benzamide	C20H22N6O5	426.426	13	9
190	(Z)-5-((6-(2-methoxyethoxy)-2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H21N7O5	451.435	14	9
191	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-(2-methoxyethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H24N6O5	452.463	15	9
192	(Z)-5-((6-(2-methoxyethoxy)-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H24N6O6S	488.517	16	9
193	(Z)-5-((2-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-(2-methoxyethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H26N6O5	478.5	17	9
194	(Z)-5-((6-(2-methoxyethoxy)-2-(5-(phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H26N6O6S	514.554	18	9
195	(Z)-5-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(2-methoxyethoxy)pyrimidin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C24H27N7O5	493.515	19	9
196	(Z)-6-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-6-(2-methoxyethoxy)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C23H25N7O5	479.489	20	9
197	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-(2-methoxyethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H24N6O5	464.474	21	9
198	(Z)-5-((6-(2-methoxyethoxy)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H24N6O6S	500.528	22	9

199	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H27N7O3	437.495	1	10
200	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C21H26N8O3	438.483	2	10
201	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H25N9O3	439.471	3	10
202	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C20H25N9O3	439.471	4	10
203	(Z)-4-(4-(2-(dimethylamino)ethoxy)-6-((2,4-dioxoimidazolin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenylpiperazine-1-carboxamide	C23H28N8O4	480.52	5	10
204	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H36N8O3	472.584	6	10
205	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H28N8O3	452.51	7	10
206	(Z)-5-((2-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-6-(2-(dimethylamino)ethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H28N8O4	492.53	8	10
207	(Z)-4-(4-(2-(dimethylamino)ethoxy)-6-((2,4-dioxoimidazolin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C24H30N8O4	494.546	9	10
208	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C26H36N10O3	536.629	10	10
209	(Z)-5-((2-((2-(diethylamino)ethyl)(methyl)amino)-6-(2-(dimethylamino)ethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C19H31N7O3	405.495	11	10
210	(Z)-N-(2-((4-(2-(dimethylamino)ethoxy)-6-((2,4-dioxoimidazolin-5-ylidene)methyl)pyrimidin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C23H26N8O5S	526.568	12	10
211	(Z)-N-(2-((4-(2-(dimethylamino)ethoxy)-6-((2,4-dioxoimidazolin-5-ylidene)methyl)pyrimidin-2-yl)amino)ethyl)benzamide	C21H25N7O4	439.468	13	10
212	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H24N8O4	464.477	14	10
213	(Z)-5-((2-(4-benzoylpiperazin-1-yl)-6-(2-(dimethylamino)ethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H27N7O4	465.505	15	10
214	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C22H27N7O5S	501.559	16	10

215	(Z)-5-((2-(5-benzoylhexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)-6-(2-(dimethylamino)ethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C25H29N7O4	491.542	17	10	
216	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(5-(phenylsulfonyl)hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H29N7O5S	527.596	18	10	
217	(Z)-5-(4-(2-(dimethylamino)ethoxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenylhexahdropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C25H30N8O4	506.557	19	10	
218	(Z)-6-(4-(2-(dimethylamino)ethoxy)-6-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C24H28N8O4	492.53	20	10	
219	(Z)-5-((2-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-6-(2-(dimethylamino)ethoxy)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C24H27N7O4	477.516	21	10	
220	(Z)-5-((6-(2-(dimethylamino)ethoxy)-2-(6-(phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione	C23H27N7O5S	513.569	22	10	
221	(Z)-5-((4-(4-phenylpiperazin-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C18H18N6O2	350.375	1	11	
222	(Z)-5-((4-(4-(pyridin-3-yl)piperazin-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C17H17N7O2	351.363	2	11	
223	(Z)-5-((4-(4-(pyrimidin-5-yl)piperazin-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C16H16N8O2	352.351	3	11	
224	(Z)-5-((4-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C16H16N8O2	352.351	4	11	
225	(Z)-4-(2-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-4-yl)-N-phenylpiperazine-1-carboxamide	C19H19N7O3	393.399	5	11	
226	(Z)-5-((4-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C19H27N7O2	385.463	6	11	
227	(Z)-5-((4-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C18H19N7O2	365.389	7	11	
228	(Z)-5-((4-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C20H19N7O3	405.41	8	11	
229	(Z)-4-(2-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-4-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C20H21N7O3	407.426	9	11	
230	(Z)-5-((4-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C22H27N9O2	449.509	10	11	

231	(Z)-5-((4-((2-(diethylamino)ethyl)(methyl)amino)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C15H22N6O2	318.374	11	11
232	(Z)-N-(2-((2-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-4-yl)amino)ethyl)isoquinoline-5-sulfonamide	C19H17N7O4S	439.448	12	11
233	(Z)-N-(2-((2-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-4-yl)amino)ethyl)benzamide	C17H16N6O3	352.347	13	11
234	(Z)-5-((4-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C18H15N7O3	377.357	14	11
235	(Z)-5-((4-(4-benzoylpiperazin-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C19H18N6O3	378.385	15	11
236	(Z)-5-((4-(4-(phenylsulfonyl)piperazin-1-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C18H18N6O4S	414.438	16	11
237	(Z)-5-((4-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C21H20N6O3	404.422	17	11
238	(Z)-5-((4-(5-phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C20H20N6O4S	440.476	18	11
239	(Z)-5-(2-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-4-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C21H21N7O3	419.437	19	11
240	(Z)-6-(2-((2,4-dioxoimidazolidin-5-ylidene)methyl)pyrimidin-4-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C20H19N7O3	405.41	20	11
241	(Z)-5-((4-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C20H18N6O3	390.395	21	11
242	(Z)-5-((4-(6-phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-2-yl)methylene)imidazolidine-2,4-dione	C19H18N6O4S	426.449	22	11
243	(Z)-5-((4-(4-phenylpiperazin-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C17H17N7O2	351.363	1	12
244	(Z)-5-((4-(4-(pyridin-3-yl)piperazin-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C16H16N8O2	352.351	2	12
245	(Z)-5-((4-(4-(pyrimidin-5-yl)piperazin-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C15H15N9O2	353.339	3	12
246	(Z)-5-((4-(4-(pyrimidin-2-yl)piperazin-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C15H15N9O2	353.339	4	12
247	(Z)-4-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-1,3,5-triazin-2-yl)-N-phenylpiperazine-1-carboxamide	C18H18N8O3	394.387	5	12
248	(Z)-5-((4-(4-(1-ethylpiperidin-4-yl)piperazin-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C18H26N8O2	386.451	6	12

249	(Z)-5-((4-(4-(pyridin-4-yl)-1,4-diazepan-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C17H18N8O2	366.377	7	12
250	(Z)-5-((4-(4-(benzo[d]oxazol-2-yl)-1,4-diazepan-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C19H18N8O3	406.398	8	12
251	(Z)-4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-1,3,5-triazin-2-yl)-N-phenyl-1,4-diazepane-1-carboxamide	C19H20N8O3	408.414	9	12
252	(Z)-5-((4-(4-(6-methyl-2-(pyrrolidin-1-yl)pyrimidin-4-yl)-1,4-diazepan-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C21H26N10O2	450.497	10	12
253	(Z)-5-((4-((2-(diethylamino)ethyl)(methyl)amino)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C14H21N7O2	319.362	11	12
254	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-1,3,5-triazin-2-yl)amino)ethyl)isoquinoline-5-sulfonamide	C18H16N8O4S	440.436	12	12
255	(Z)-N-(2-((4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-1,3,5-triazin-2-yl)amino)ethyl)benzamide	C16H15N7O3	353.335	13	12
256	(Z)-5-((4-((2-(4-oxo-3,4-dihydroquinazolin-2-yl)ethyl)amino)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C17H14N8O3	378.345	14	12
257	(Z)-5-((4-(4-benzoylpiperazin-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C18H17N7O3	379.373	15	12
258	(Z)-5-((4-(4-(phenylsulfonyl)piperazin-1-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C17H17N7O4S	415.426	16	12
259	(Z)-5-((4-(5-benzoylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C20H19N7O3	405.41	17	12
260	(Z)-5-((4-(5-phenylsulfonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C19H19N7O4S	441.464	18	12
261	(Z)-5-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-1,3,5-triazin-2-yl)-N-phenylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide	C20H20N8O3	420.425	19	12
262	(Z)-6-(4-((2,4-dioxoimidazolidin-5-ylidene)methyl)-1,3,5-triazin-2-yl)-N-phenyl-2,6-diazaspiro[3.3]heptane-2-carboxamide	C19H18N8O3	406.398	20	12
263	(Z)-5-((4-(6-benzoyl-2,6-diazaspiro[3.3]heptan-2-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C19H17N7O3	391.383	21	12
264	(Z)-5-((4-(6-phenylsulfonyl)-2,6-diazaspiro[3.3]heptan-2-yl)-1,3,5-triazin-2-yl)methylene)imidazolidine-2,4-dione	C18H17N7O4S	427.437	22	12

In addition, it may be convenient or desirable to prepare, purify, and/or handle the active compound in a chemically protected form. The term "chemically protected form," as used herein, pertains to a compound in which one or more reactive functional groups are protected from undesirable chemical reactions (i.e., they have been modified with a 5 protecting group).

By protecting a reactive functional group, reactions involving other unprotected reactive functional groups can be performed without affecting the protected group; the protecting group may be removed, usually in a subsequent step, without substantially affecting the remainder of the molecule. See, for example, Protective Groups in Organic 10 Synthesis (T. Green and P. Wuts, Wiley, 1991), and Protective Groups in Organic Synthesis (T. Green and P. Wuts; 3rd Edition; John Wiley and Sons, 1999).

For example, a hydroxy group may be protected as an ether (-OR) or an ester (-OC(=O)R), for example, as: a t-butyl ether; a benzyl, benzhydryl (diphenylmethyl), or trityl (triphenylmethyl) ether; a trimethylsilyl or t-butyldimethylsilyl ether; or an acetyl 15 ester (-OC(=O)CH₃, -OAc).

For example, an aldehyde or ketone group may be protected as an acetal or ketal, respectively, in which the carbonyl group (C(=O)) is converted to a diether (C(OR)₂), by reaction with, for example, a primary alcohol. The aldehyde or ketone group is readily regenerated by hydrolysis using a large excess of water in the presence of acid.

20 For example, an amine group may be protected, for example, as an amide (-NRC(=O)R) or a urethane (-NRC(=O)OR), for example, as: a methyl amide (-NHC(=O)CH₃); a benzyloxy amide (-NHC(=O)OCH₂C₆H₅NHCbz); as a t-butoxy amide (-NHC(=O)OC(CH₃)₃, -NHBOC); a 2-biphenyl-2-propoxy amide (-NHC(=O)OC(CH₃)₂C₆H₄C₆H₅NHBoc), as a 9-fluorenylmethoxy amide (-NHFmoc), as a 6-nitroveratryloxy amide (-NHNvoc), as a 2-trimethylsilylethoxy amide (-NHTeoc), as a 25 2,2,2-trichloroethoxy amide (-NHTroc), as an allyloxy amide (-NHAlloc), as a 2-(phenylsulfonyl)ethoxy amide (-NHPsec); or, in suitable cases (e.g., cyclic amines), as a nitroxide radical.

For example, a carboxylic acid group may be protected as an ester or an amide, for 30 example, as: a benzyl ester; a t-butyl ester; a methyl ester; or a methyl amide.

For example, a thiol group may be protected as a thioether (-SR), for example, as: a benzyl thioether; or an acetamidomethyl ether (-SCH₂NHC(=O)CH₃).

PHARMACEUTICAL COMPOSITIONS

One or more compounds of this invention can be administered to a mammal by themselves or in pharmaceutical compositions where they are mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a disease or condition as described herein. Mixtures of these compounds can also be administered to the patient as a simple mixture or in suitable formulated pharmaceutical compositions. For example, one aspect of the invention relates to pharmaceutical composition comprising a therapeutically effective dose 10 of a compound of formula I, or a pharmaceutically acceptable salt, solvate, enantiomer or stereoisomer thereof; and a pharmaceutically acceptable diluent or carrier.

Techniques for formulation and administration of the compounds of the instant application may be found in references well known to one of ordinary skill in the art, such as "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest 15 edition.

Suitable routes of administration may, for example, include oral, eyedrop, rectal, transmucosal, topical, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

20 Alternatively, one may administer a compound in a local rather than a systemic manner, for example, via injection of the compound directly into an edematous site, often in a depot or sustained release formulation.

Furthermore, one may administer a compound in a targeted drug delivery system, for example, in a liposome coated with endothelial-cell-specific antibody.

25 The pharmaceutical compositions of the present invention may be manufactured, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus 30 may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active

compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, 5 or physiological saline buffer. For transmucosal administration, penetrants are used in the formulation appropriate to the barrier to be permeated. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such 10 carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by combining the active compound with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, 15 to obtain tablets or dragee cores. Suitable excipients include fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the 20 cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginato.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl 25 pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as 30 glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may

be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

5 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of pressurized aerosol the dosage unit may be determined
10 by providing a valve to deliver a metered amount. Capsules and cartridges of e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds can be formulated for parenteral administration by injection, e.g., bolus injection or continuous infusion. Formulations for injection may be presented in unit
15 dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions
20 of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl
25 cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for reconstitution before use with a suitable vehicle, e.g., sterile pyrogen-free water.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be
5 formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly or by intramuscular injection). Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives (for example, as a sparingly soluble salt).

10 Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed. Additionally, the compounds may be delivered using a sustained-release system, such as semi-permeable matrices of solid hydrophobic polymers containing the therapeutic
15 agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

20 The pharmaceutical compositions may also comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers, such as polyethylene glycols.

METHODS OF TREATMENT

25 Provided herein are methods of modulating the activity of CK1 and subtypes thereof, the Wnt pathway, and/or the TGF β pathway. Also provided herein are methods of treating or preventing conditions and diseases the course of which can be influenced by modulating the activity of CK1 (e.g., CK1 γ), the Wnt pathway, and/or the TGF β pathway. Such methods typically comprise administering to a subject in need thereof a
30 therapeutically effective amount of a compound or composition of the invention.

Various diseases, such as cancers, inflammation, and inflammatory diseases (e.g., osteoarthritis and rheumatoid arthritis), and neurological conditions (e.g., Alzheimer's disease) and neurodegeneration can be treated by administration of modulators of CK1 (e.g., CK1 γ), the Wnt pathway and/or the TGF β pathway. Bone-related diseases and 5 conditions, including osteoporosis and bone formation, also can be treated by administration of modulators of CK1 (e.g., CK1 γ), the Wnt pathway and/or the TGF β pathway. Bone restoration can be facilitated by administration of modulators of CK1 (e.g., CK1 γ), the Wnt pathway and/or the TGF β pathway. Additional conditions that can be treated by administration of modulators of CK1 (e.g., CK1 γ), the Wnt pathway and/or the 10 TGF β pathway include hypoglycemia, metabolic syndrome and diabetes. Modulators of CK1 (e.g., CK1 γ), the Wnt pathway and/or the TGF β pathway are also useful for influencing apoptosis (e.g., increasing the rate of apoptosis in cancerous cells). Modulators of CK1 (e.g., CK1 γ), the Wnt pathway and/or the TGF β pathway are also useful in treatment or prevention of aberrant embryonic development.

15 Based at least on the fact that increased CK1 γ has been found to be associated with certain cancers, a method for treating cancer in a subject comprises administering to the subject in need thereof a therapeutically effective amount of a compound that inhibits CK1 γ .

CK1 γ inhibiting compounds can be used for modulating cell proliferation, generally. 20 Accordingly, diseases that may be treated include hyperproliferative diseases, such as benign cell growth and malignant cell growth.

Exemplary cancers that may be treated include leukemias, e.g., acute lymphoid leukemia and myeloid leukemia, and carcinomas, such as colorectal carcinoma and hepatocarcinoma. Other cancers include Acute Lymphoblastic Leukemia; Acute 25 Lymphoblastic Leukemia; Acute Myeloid Leukemia; Acute Myeloid Leukemia; Adrenocortical Carcinoma Adrenocortical Carcinoma; AIDS-Related Cancers; AIDS-Related Lymphoma; Anal Cancer; Astrocytoma, Childhood Cerebellar; Astrocytoma, Childhood Cerebral; Basal Cell Carcinoma, see Skin Cancer (non-Melanoma); Bile Duct Cancer, Extrahepatic; Bladder Cancer; Bladder Cancer; Bone Cancer, 30 osteosarcoma/Malignant Fibrous Histiocytoma; Brain Stem Glioma; Brain Tumor; Brain Tumor, Brain Stem Glioma; Brain Tumor, Cerebellar Astrocytoma; Brain Tumor, Cerebral

Astrocytoma/Malignant Glioma; Brain Tumor, Ependymoma; Brain Tumor, Medulloblastoma; Brain Tumor, Supratentorial Primitive Neuroectodermal Tumors; Brain Tumor, Visual Pathway and Hypothalamic Glioma; Brain Tumor; Breast Cancer; Breast Cancer and Pregnancy; Breast Cancer; Breast Cancer, Male; Bronchial Adenomas/Carcinoids; Burkitt's Lymphoma; Carcinoid Tumor; Carcinoid Tumor, Gastrointestinal; Carcinoma of Unknown Primary; Central Nervous System Lymphoma, Primary; Cerebellar Astrocytoma; Cerebral Astrocytoma/Malignant Glioma; Cervical Cancer; Childhood Cancers; Chronic Lymphocytic Leukemia; Chronic Myelogenous Leukemia; Chronic Myeloproliferative Disorders; Colon Cancer; Colorectal Cancer;

Cutaneous T-Cell Lymphoma, see Mycosis Fungoides and Sezary Syndrome; Endometrial Cancer; Ependymoma; Esophageal Cancer; Esophageal Cancer; Ewing's Family of Tumors; Extracranial Germ Cell Tumor; Extragonadal Germ Cell Tumor; Extrahepatic Bile Duct Cancer; Eye Cancer, Intraocular Melanoma; Eye Cancer, Retinoblastoma; Gallbladder Cancer; Gastric (Stomach) Cancer; Gastric (Stomach) Cancer; Gastrointestinal Carcinosarcoma; Germ Cell Tumor, Extracranial; Germ Cell Tumor, Extragonadal; Germ Cell Tumor, Ovarian; Gestational Trophoblastic Tumor; Glioma; Glioma, Childhood Brain Stem; Glioma, Childhood Cerebral Astrocytoma; Glioma, Childhood Visual Pathway and Hypothalamic; Hairy Cell Leukemia; Head and Neck Cancer; Hematologic (Blood) Cancer, Hepatocellular (Liver) Cancer, Adult (Primary); Hepatocellular (Liver) Cancer, Childhood (Primary); Hodgkin's Lymphoma; Hodgkin's Lymphoma; Hodgkin's Lymphoma During Pregnancy; Hypopharyngeal Cancer; Hypothalamic and Visual Pathway Glioma; Intraocular Melanoma; Islet Cell Carcinoma (Endocrine Pancreas); Kaposi's Sarcoma; Kidney (Renal Cell) Cancer; Kidney Cancer; Laryngeal Cancer; Laryngeal Cancer; Leukemia, Acute Lymphoblastic; Leukemia, Acute Lymphoblastic; Leukemia, Acute Myeloid; Leukemia, Acute Myeloid; Leukemia, Chronic Lymphocytic; Leukemia; Chronic Myelogenous; Leukemia, Hairy Cell; Lip and Oral Cavity Cancer; Liver Cancer, Adult (Primary); Liver Cancer, Childhood (Primary); Lung Cancer, Non-Small Cell; Lung Cancer, Small Cell; Lymphoma, AIDS-Related; Lymphoma, Burkitt's; Lymphoma, Cutaneous T-Cell, see Mycosis Fungoides and Sezary Syndrome; Lymphoma, Hodgkin's; Lymphoma, Hodgkin's; Lymphoma, Hodgkin's During Pregnancy; Lymphoma, Non-Hodgkin's; Lymphoma, Non-Hodgkin's; Lymphoma, Non-Hodgkin's During Pregnancy; Lymphoma, Primary Central Nervous System; Macroglobulinemia, Waldenstrom's; Malignant Fibrous Histiocytoma of Bone/Osteosarcoma; Medulloblastoma; Melanoma;

Melanoma, Intraocular (Eye); Merkel Cell Carcinoma; Mesothelioma, Adult Malignant; Mesothelioma; Metastatic Squamous Neck Cancer with Occult Primary; Multiple Endocrine Neoplasia Syndrome; Multiple Myeloma/Plasma Cell Neoplasm' Mycosis Fungoides; Myelodysplastic Syndromes; Myelodysplastic/Myeloproliferative Diseases;

5 Myelogenous Leukemia, Chronic; Myeloid Leukemia, Adult Acute; Myeloid Leukemia, Childhood Acute; Myeloma, Multiple; Myeloproliferative Disorders, Chronic; Nasal Cavity and Paranasal Sinus Cancer; Nasopharyngeal Cancer; Nasopharyngeal Cancer; Neuroblastoma; Non-Hodgkin's Lymphoma; Non-Hodgkin's Lymphoma; Non-Hodgkin's Lymphoma During Pregnancy; Non-Small Cell Lung Cancer; Oral Cancer; Oral Cavity

10 Cancer, Lip and; Oropharyngeal Cancer; Osteosarcoma/Malignant Fibrous Histiocytoma of Bone; Ovarian Cancer; Ovarian Epithelial Cancer; Ovarian Germ Cell Tumor; Ovarian Low Malignant Potential Tumor; Pancreatic Cancer; Pancreatic Cancer; Pancreatic Cancer, Islet Cell; Paranasal Sinus and Nasal Cavity Cancer; Parathyroid Cancer; Penile Cancer; Pheochromocytoma; Pineoblastoma and Supratentorial Primitive Neuroectodermal Tumors;

15 Pituitary Tumor; Plasma Cell Neoplasm/Multiple Myeloma; Pleuropulmonary Blastoma; Pregnancy and Breast Cancer; Pregnancy and Hodgkin's Lymphoma; Pregnancy and Non-Hodgkin's Lymphoma; Primary Central Nervous System Lymphoma; Prostate Cancer; Rectal Cancer; Renal Cell (Kidney) Cancer; Renal Cell (Kidney) Cancer; Renal Pelvis and Ureter, Transitional Cell Cancer; Retinoblastoma; Rhabdomyosarcoma; Salivary Gland

20 Cancer; Salivary Gland Cancer; Sarcoma, Ewing's Family of Tumors; Sarcoma, Kaposi's; Sarcoma, Soft Tissue; Sarcoma, Soft Tissue; Sarcoma, Uterine; Sezary Syndrome; Skin Cancer (non-Melanoma); Skin Cancer; Skin Cancer (Melanoma); Skin Carcinoma, Merkel Cell; Small Cell Lung Cancer; Small Intestine Cancer; Soft Tissue Sarcoma; Soft Tissue Sarcoma; Squamous Cell Carcinoma, see Skin Cancer (non-Melanoma); Squamous Neck

25 Cancer with Occult Primary, Metastatic; Stomach (Gastric) Cancer; Stomach (Gastric) Cancer; Supratentorial Primitive Neuroectodermal Tumors; T-Cell Lymphoma, Cutaneous, see Mycosis Fungoides and Sezary Syndrome; Testicular Cancer; Thymoma; Thymoma and Thymic Carcinoma; Thyroid Cancer; Thyroid Cancer; Transitional Cell Cancer of the Renal Pelvis and Ureter; Trophoblastic Tumor, Gestational; Unknown Primary Site,

30 Carcinoma of; Unknown Primary Site, Cancer of; Unusual Cancers of Childhood; Ureter and Renal Pelvis, Transitional Cell Cancer; Urethral Cancer; Uterine Cancer, Endometrial; Uterine Sarcoma; Vaginal Cancer; Visual Pathway and Hypothalamic Glioma; Vulvar Cancer; Waldenstrom's Macroglobulinemia; Wilms' Tumor; and Women's Cancers.

Neurologic diseases that may be treated include epilepsy, schizophrenia, bipolar disorder or other psychological and/or psychiatric disorders, neuropathies, skeletal muscle atrophy, and neurodegenerative diseases, e.g., a neurodegenerative disease. Exemplary neurodegenerative diseases include: Alzheimer's, Amyotrophic Lateral Sclerosis (ALS),
5 and Parkinson's disease. Another class of neurodegenerative diseases includes diseases caused at least in part by aggregation of poly-glutamine. Diseases of this class include: Huntington's Disease, Spinalbulbar Muscular Atrophy (SBMA or Kennedy's Disease) Dentatorubropallidoluysian Atrophy (DRPLA), Spinocerebellar Ataxia 1 (SCA1),
10 Spinocerebellar Ataxia 2 (SCA2), Machado-Joseph Disease (MJD; SCA3), Spinocerebellar Ataxia 6 (SCA6), Spinocerebellar Ataxia 7 (SCA7), and Spinocerebellar Ataxia 12 (SCA12).

Any other disease in which the Wnt pathway, TGF β pathway, or CK1 γ plays a role may be treatable or preventable using compounds and methods described herein.

DOSAGE

15 As used herein, a "therapeutically effective amount" or "therapeutically effective dose" is an amount of a compound of the invention or a combination of two or more such compounds, which inhibits, totally or partially, the progression of the condition or alleviates, at least partially, one or more symptoms of the condition. A therapeutically effective amount can also be an amount which is prophylactically effective. The amount
20 which is therapeutically effective will depend upon the patient's size and gender, the condition to be treated, the severity of the condition and the result sought. For a given patient, a therapeutically effective amount may be determined by methods known to those of skill in the art.

A therapeutically effective dose refers to that amount of the compound that results
25 in amelioration of symptoms in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the maximum tolerated dose (MTD) and the ED₅₀ (effective dose for 50% maximal response). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between
30 MTD and ED₅₀. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little

or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. In the treatment of crises, the administration of an acute bolus or an infusion approaching the MTD may be required to obtain a rapid response.

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the CK1 γ modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using the MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90% until the desired amelioration of symptoms is achieved. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.


KITS

The compounds and compositions of the invention (e.g., compounds and compositions of formula I) may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labelled for treatment of an indicated condition. Instructions for use may also be provided.

EXEMPLIFICATION


The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit 5 the invention. The geometric isomers depicted below are believed to be correct, but final structural assignment will be made via 2-D NMR experiments. Although the exemplary compounds described below are believed to be the Z-geometric isomers, the E-geometric isomers and mixtures of the E-and Z-isomers are also contemplated by the present disclosure.

10 EXAMPLE 1



(E)-4-(dimethylamino)-1,1-dimethoxybut-3-en-2-one (1): 1,1-dimethoxy-N,N-dimethylmethanamine (100 g, 839 mmol, 1.02 equiv.) and 1,1-dimethoxypropan-2-one (97 g, 821 mmol) were added and stirred at 110°C for 3 hours. The produced methanol was removed by a Dean-Stark apparatus. After the solution was cooled to room temperature, the remaining volatile materials were removed *in vacuo* to provide 130 g of the crude product, (E)-4-(dimethylamino)-1,1-dimethoxybut-3-en-2-one (1) (130 g, 143 g theoretical, 91%). LC-MS *m/z* 283 (M+1). Reference: WO20060097341A1, pg 67.


20 EXAMPLE 2



sodium 4-(dimethoxymethyl)pyrimidine-2-thiolate (2): A solution of thiourea (64.7 g, 850 mmol, 1.13 equiv.), sodium methanolate (95%, 40.5 g, 751 mmol, 1.0 equiv.) in methanol (500 mL, 1.5 M) was stirred at room temperature for 30 minutes. A solution of (E)-4-(dimethylamino)-1,1-dimethoxybut-3-en-2-one (1) (130 g, 751 mmol) in methanol


(200 mL) was added and the reaction stirred at room temperature for 2 h. The crude sodium 4-(dimethoxymethyl)pyrimidine-2-thiolate (**2**) was used directly in the next step without further purification. LC-MS *m/z* 209 (M+1). Reference: WO2006009734A1, pg 67.

5 EXAMPLE 3



4-(dimethoxymethyl)-2-(methylthio)pyrimidine (3): Iodomethane (128 g, 902 mmol, 1.20 equiv.) was added carefully to the crude solution of **sodium 4-(dimethoxymethyl)pyrimidine-2-thiolate (2)** (156 g, 751 mmol) in methanol (700 mL, 1.1 M) while maintaining the reaction temperature below 28°C using an ice-water bath for cooling. The resulting mixture was stirred at room temperature for 16 h. After removal of the solvent under reduced pressure, the residue was diluted with water (300 mL) and extracted with ethyl acetate (2 x 150 mL). The combined organic layer was concentrated under reduced pressure and the crude residue purified by passing through a short silica gel pad and washing with diethyl ether (200 mL) to afford **4-(dimethoxymethyl)-2-(methylthio)pyrimidine (3)** as a brown oil (53.7 g, 150 g theoretical, 35.7%). LC-MS *m/z* 201 (M+1). Reference: WO20060097341A1, pg 67.


20 EXAMPLE 4



2-(methylthio)pyrimidine-4-carbaldehyde (4): 4-(dimethoxymethyl)-2-(methylthio)pyrimidine (3) (53.7 g, 268 mmol) was added carefully to 1.2 N aqueous HCl (300 mL, 268 mmol, 1.0 equiv.) and stirred at 60°C for 3 hours. The reaction mixture was 25 then cooled to room temperature and neutralized by the slow addition of solid sodium bicarbonate. The crude mixture was extracted with diethyl ether (3 x 150 mL) and the combined organic layer was concentrated under reduced pressure to afford 2-


(methylthio)pyrimidine-4-carbaldehyde (4) as a yellow solid (14.2 g, 41.5 g theoretical, 34%). LC-MS *m/z* 155 (M+1). Reference: W2006009734A1, pg 67.

EXAMPLE 5




5 **(Z)-5-((2-(methylthio)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (5a):** A 40 mL round bottomed vial was charged with **2-(methylthio)pyrimidine-4-carbaldehyde (4)** (771 mg, 5 mmol), imidazolidine-2,4-dione (500 mg, 5 mmol, 1.0 equiv.), and piperidine (400 µL, 4 mmol, 0.8 equiv.) in ethanol (20 mL, 0.25 M). The reaction mixture was heated to 80°C and shaken for 20 h. The resulting yellow precipitate was isolated by filtration and 10 washed with ethanol (1 x 20 mL) and dried *in vacuo* to afford **(Z)-5-((2-(methylthio)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (5a)** as a yellow solid (565 mg, 1.18 mg theoretical, 48%). LC-MS *m/z* 237 (M+1). The same procedure can be used to prepare **5b** or **5c** by replacing imidazolidine-2,4-dione with 1-benzylimidazolidine-2,4-dione or 1-methylimidazolidine-2,4-dione respectively.

15 EXAMPLE 6




(Z)-5-((2-(methylsulfonyl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (6a): A mixture of **(Z)-5-((2-(methylthio)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (5)** (3.5 g, 14.83 mmol) in THF(100 mL, 0.15 M) was treated with a solution of oxone (27.35 g, 44.5 mmol, 3.0 equiv.) in water(175 mL). The resulting mixture was stirred at room temperature for 48 h. The resulting precipitate was filtered and washed with water (20 mL) and diethyl ether (20 mL) to afford **(Z)-5-((2-(methylsulfonyl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (6)** as a solid (2.48 g, 3.97 g theoretical, 62%). LC-MS *m/z* 269 (M+1). The same procedure can be used with the N-benzyl (**6b**) and N-methyl (**6c**) analogs in comparable yields.

EXAMPLE 7




General displacement procedure 1: 2 dram round bottomed vials were charged with (*Z*)-5-((2-(methylsulfonyl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (**6a**) (30 mg, 0.1118 mmol), DMSO (0.5 mL, 0.2 M), diisopropylethylamine (50 μ L, 0.306 mmol, 2.7 equiv.), and the appropriate amine (0.1118 mmol, 1.0 equiv.). The reaction mixture was heated to 120°C and shaken for 16 h. The solvent was removed under reduced pressure (Genevac HT-4) and the crude residues were purified using reverse phase HPLC (MS-triggered fraction collection) with an acetonitrile/water gradient and trifluoroacetic acid as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac (HT-4) to provide the desired products (**7a**).

EXAMPLE 8




15 General displacement procedure 2: 2 dram round bottomed vials were charged with (*Z*)-1-benzyl-5-((2-(methylsulfonyl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (**6b**) (30 mg, 0.084 mmol), DMSO (0.5 mL, 0.16 M), diisopropylethylamine (50 μ L, 0.306 mmol, 3.6 equiv.), and the appropriate amine (0.084 mmol, 3.6 equiv.). The reaction mixture was heated to 120°C and shaken for 16 h. The solvent was removed under reduced pressure (Genevac HT-4) and the crude residues were purified using reverse phase HPLC (MS-triggered fraction collection) with an acetonitrile/water gradient and trifluoroacetic acid as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac (HT-4) to provide the desired products (**7b**).

EXAMPLE 9




General displacement procedure 3: 2 dram round bottomed vials were charged with (*Z*)-1-methyl-5-((2-(methylsulfonyl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (**6c**) (25 mg, 0.089 mmol), DMSO (0.5mL, 0.16 M), diisopropylethylamine (50 μ L, 0.177 mmol, 3.4 equiv.), and the appropriate amine (0.089 mmol, 1.0 equiv.). The reaction mixture was heated to 120°C and shaken for 16 h. The solvent was removed under reduced pressure (Genevac HT-4) and the crude residues were purified using reverse phase HPLC (MS-triggered fraction collection) with an acetonitrile/water gradient and trifluoroacetic acid as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac (HT-4)) to provide the desired products (**7c**).

EXAMPLE 10




(*Z*)-5-((2-(4-(benzo[d][1,3]dioxol-5-ylmethyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (25.1 mg, 36.4 mg theoretical, 69%). LC-MS m/z 409.4 (M+1).

EXAMPLE 11




(*Z*)-5-((2-(4-(p-tolyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (14.3 mg, 32.5 mg theoretical, 44%). LC-MS m/z 365.4 (M+1).

EXAMPLE 12




(Z)-5-((2-(methyl(2-(pyridin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 5 1 (24.4 mg, 29 mg theoretical, 84.1%). LC-MS m/z 325.3 (M+1).

EXAMPLE 13




(Z)-5-((2-(4-isopropylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione 10 was prepared using general displacement procedure 1 (21.9 mg, 28.2 mg theoretical, 77.7%). LC-MS m/z 317.4 (M+1).

EXAMPLE 14




(Z)-5-((2-(3,4-dihydroisoquinolin-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione 15 was prepared using general displacement procedure 1 (11.6 mg, 28.7 mg theoretical, 40.4%). LC-MS m/z 322.3 (M+1).

EXAMPLE 15



(Z)-5-((2-(4-(pyridin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (6.1 mg, 31.4 mg theoretical, 19.4%). LC-MS m/z 352.4 (M+1).


EXAMPLE 16



5

(Z)-5-((2-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (25.7 mg, 25.7 mg theoretical, 100%). LC-MS m/z 289.3 (M+1).


EXAMPLE 17



10

(Z)-methyl 1-((2,5-dioxoimazolidin-4-ylidene)methyl)pyrrolidin-2-yl)pyrimidin-2-yl)pyrrolidin-2-carboxylate was prepared using general displacement procedure 1 (14.4 mg, 28.3 mg theoretical, 50.9%). LC-MS m/z 318.3 (M+1).

EXAMPLE 18




15

(Z)-5-((2-(4-propylpiperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (7.9 mg, 28.2 mg theoretical, 28%). LC-MS m/z 316.4 (M+1).


20

EXAMPLE 19




(**Z**)-5-((2-(4-morpholinopiperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (30.8 mg, 32 mg theoretical, 96.3%). LC-MS m/z 359.4 (M+1).

EXAMPLE 20




(**Z**)-5-((2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (4.3 mg, 31.5 mg theoretical, 13.7%). LC-MS m/z 353.4 (M+1).

EXAMPLE 21




(**Z**)-5-((2-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (14 mg, 24.6 mg theoretical, 56.9%). LC-MS m/z 276.3 (M+1).

EXAMPLE 22



(Z)-5-((2-(piperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (16.8 mg, 24.4 mg theoretical, 68.9%). LC-MS m/z 274.3 (M+1).


EXAMPLE 23



5

(Z)-5-((2-(pyrrolidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (2.9 mg, 23.2 mg theoretical, 12.5%). LC-MS m/z 260.3 (M+1).


EXAMPLE 24



10

(Z)-5-((2-(4-(pyrrolidin-1-yl)piperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (26.7 mg, 30.6 mg theoretical, 87.3%). LC-MS m/z 343.4 (M+1).

EXAMPLE 25




15

(Z)-5-((2-(4-(2-morpholino-2-oxoethyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 1 (35.8 mg, 35.8 mg theoretical, 100%). LC-MS m/z 402.4 (M+1).

20

EXAMPLE 26




(Z)-5-((2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

was prepared using general displacement procedure 1 (10.6 mg, 31.3 mg theoretical,

5 LC-MS m/z 351.4 (M+1).

EXAMPLE 27




(Z)-5-((2-(methyl(phenethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

was prepared using general displacement procedure 1 (15.8 mg, 28.9 mg theoretical,

10 LC-MS m/z 324.3 (M+1).

EXAMPLE 28




(Z)-5-((2-(4-(pyridin-4-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-

dione was prepared using general displacement procedure 1 (15.6 mg, 31.4 mg theoretical,


15 LC-MS m/z 352.4 (M+1).

EXAMPLE 29




(Z)-1-methyl-5-((2-(4-(p-tolyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (9.9 mg, 33.8 mg theoretical, 29.3%). LC-MS m/z 379.4 (M+1).

EXAMPLE 30




(Z)-1-methyl-5-((2-(methyl(2-(pyridin-2-yl)ethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (10.3 mg, 30.2 mg theoretical, 34.1%). LC-MS m/z 339.4 (M+1).

EXAMPLE 31




(Z)-5-((2-(4-isopropylpiperazin-1-yl)pyrimidin-4-yl)methylene)-1-methylimidazolidine-2,4-dione was prepared using general displacement procedure 3 (10.4 mg, 29.5 mg theoretical, 35.3%). LC-MS m/z 331.4 (M+1).

EXAMPLE 32




(Z)-5-((2-(3,4-dihydroisoquinolin-2(1H)-yl)pyrimidin-4-yl)methylene)-1-methylimidazolidine-2,4-dione was prepared using general displacement procedure 3 (10.9 mg, 29.9 mg theoretical, 36.5%). LC-MS m/z 336.4 (M+1).

EXAMPLE 33




(Z)-1-methyl-5-((2-(4-(pyridin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (15.7 mg, 32.6 mg theoretical, 48.2%). LC-MS m/z 366.4 (M+1).

EXAMPLE 34




(Z)-1-methyl-5-((2-(4-methylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (14.9 mg, 27 mg theoretical, 55.2%). LC-MS m/z 303.3 (M+1).

EXAMPLE 35




(Z)-1-methyl-5-((2-(4-propylpiperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (5.8 mg, 29.4 mg theoretical, 19.7%). LC-MS m/z 330.4 (M+1).

EXAMPLE 36



(Z)-1-methyl-5-((2-(4-morpholinopiperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (16 mg, 33.2 mg theoretical, 48.2%). LC-MS m/z 373.4 (M+1).


EXAMPLE 37



5

(Z)-tert-butyl (1-(4-((3-methyl-2,5-dioxoimazolidin-4-ylidene)methyl)pyrimidin-2-yl)pyrrolidin-3-yl)carbamate was prepared using general displacement procedure 3 (19 mg, 34.7 mg theoretical, 54.8%). LC-MS m/z 389.4 (M+1).


EXAMPLE 38



10

(Z)-1-methyl-5-((2-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (17.1 mg, 32.7 mg theoretical, 52.3%). LC-MS m/z 367.4 (M+1).

EXAMPLE 39




15

(Z)-1-methyl-5-((2-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (17.8 mg, 25.8 mg theoretical, 69%). LC-MS m/z 290.3 (M+1).

20

EXAMPLE 40




(Z)-1-methyl-5-((2-(piperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

was prepared using general displacement procedure 3 (11.5 mg, 25.7 mg theoretical,

5 44.7%). LC-MS m/z 288.3 (M+1).

EXAMPLE 41




(Z)-1-methyl-5-((2-(pyrrolidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione

was prepared using general displacement procedure 3 (13.5 mg, 24.4 mg theoretical,

10 55.3%). LC-MS m/z 274.3 (M+1).

EXAMPLE 42




(Z)-1-methyl-5-((2-(4-(pyrrolidin-1-yl)piperidin-1-yl)pyrimidin-4-

yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure

15 3 (11.9 mg, 31.8 mg theoretical, 37.4%). LC-MS m/z 357.4 (M+1).

EXAMPLE 43




(Z)-tert-butyl 4-((3-methyl-2,5-dioxoimidazolidin-4-ylidene)methyl)pyrimidin-2-

yl)piperazine-1-carboxylate was prepared using general displacement procedure 3 (5.4


20 mg, 34.7 mg theoretical, 15.6%). LC-MS m/z 389.4 (M+1).

EXAMPLE 44




(*Z*)-tert-butyl 4-((3-methyl-2,5-dioxoimidazolidin-4-ylidene)methyl)pyrimidin-2-yl-1,4-diazepane-1-carboxylate was prepared using general displacement procedure 3 (15.3 mg, 35.9 mg theoretical, 42.6%). LC-MS m/z 403.4 (M+1).

EXAMPLE 45




(*Z*)-1-methyl-5-((2-(4-(2-morpholino-2-oxoethyl)piperazin-1-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (18.6 mg, 37.1 mg theoretical, 50.1%). LC-MS m/z 416.4 (M+1).

EXAMPLE 46



(*Z*)-1-methyl-5-((2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (15.8 mg, 32.5 mg theoretical, 48.6%). LC-MS m/z 365.4 (M+1).

EXAMPLE 47



(Z)-1-methyl-5-((2-(methyl(phenethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 3 (10.6 mg, 30.1 mg theoretical, 35.2%). LC-MS m/z 338.4 (M+1).


EXAMPLE 48



5

(Z)-tert-butyl (1-((3-methyl-2,5-dioxoimidazolidin-4-ylidene)methyl)pyrimidin-2-yl)piperidin-4-ylcarbamate was prepared using general displacement procedure 3 (17.6 mg, 35.9 mg theoretical, 49%). LC-MS m/z 403.4 (M+1).


EXAMPLE 49



10

(Z)-tert-butyl ((1-((3-methyl-2,5-dioxoimidazolidin-4-ylidene)methyl)pyrimidin-2-yl)piperidin-3-yl)carbamate was prepared using general displacement procedure 3 (17.3 mg, 37.2 mg theoretical, 46.5%). LC-MS m/z 417.5 (M+1).

EXAMPLE 50




15

(Z)-5-((2-(4-(benzo[d][1,3]dioxol-5-ylmethyl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 2 (21 mg, 44.5 mg theoretical, 47.2%). LC-MS m/z 499.5 (M+1).

20

EXAMPLE 51




(Z)-1-benzyl-5-((2-(3,4-dihydroisoquinolin-2(1H)-yl)yl)methylene)imidazolidine-2,4-dione

was prepared using general displacement procedure

5 2 (8.2 mg, 36.7 mg theoretical, 22.3%). LC-MS m/z 412.5 (M+1).

EXAMPLE 52




(Z)-tert-butyl (1-(4-((3-benzyl-2,5-dioxoimazolidin-4-ylidene)methyl)pyrimidin-2-

yl)pyrrolidin-3-yl)carbamate

was prepared using general displacement procedure 2 (41.4

10 mg, 41.4 mg theoretical, 100%). LC-MS m/z 465.5 (M+1).

EXAMPLE 53




(Z)-1-benzyl-5-((2-morpholinopyrimidin-4-yl)methylene)imidazolidine-2,4-dione

was prepared using general displacement procedure 2 (11.1 mg, 32.6 mg theoretical, 34%). LC-


15 MS m/z 366.4 (M+1).

EXAMPLE 54




(*Z*)-tert-butyl 4-((3-benzyl-2,5-dioxoimidazolidin-4-ylidene)methyl)pyrimidin-2-yl-1,4-diazepane-1-carboxylate was prepared using general displacement procedure 2 (12.1 mg, 42.7 mg theoretical, 28.3%). LC-MS m/z 479.5 (M+1).

EXAMPLE 55




(*Z*)-1-benzyl-5-((2-(4-phenylpiperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 2 (7.4 mg, 39.3 mg theoretical, 18.8%). LC-MS m/z 441.5 (M+1).

EXAMPLE 56




(*Z*)-1-benzyl-5-((2-(methyl(phenethyl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 2 (15.9 mg, 36.9 mg theoretical, 43.1%). LC-MS m/z 414.5 (M+1).

EXAMPLE 57




(Z)-1-benzyl-5-((2-(4-(pyridin-4-yl)piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione was prepared using general displacement procedure 5 2 (9.9 mg, 39.4 mg theoretical, 25.1%). LC-MS m/z 442.5 (M+1).

EXAMPLE 58




(Z)-tert-butyl ((1-(4-((3-benzyl-2,5-dioxoimazolidin-4-ylidene)methyl)piperidin-3-yl)methyl)carbamate was prepared using general displacement procedure 10 2 (19.4 mg, 43.9 mg theoretical, 44.2%). LC-MS m/z 493.6 (M+1).

EXAMPLE 59




(Z)-5-((2-(piperazin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: The boc-protected compound was prepared using general displacement procedure 1. The purified 15 boc-protected compound was treated with dichloromethane (1.0 mL), hydrochloric acid in methanol (500 µL, 1.25 M) and shaken at 50°C for 16 h. The reaction mixture was then concentrated under reduced pressure (Genevac HT-4) to provide the desired compound (5 mg, 24.5 mg theoretical, 20.4%). LC-MS m/z 275.3 (M+1).

EXAMPLE 60



(Z)-5-((2-(1,4-diazepan-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: The boc-protected compound was prepared using general displacement procedure 1. The purified boc-protected compound was treated with dichloromethane (1.0 mL), hydrochloric acid in methanol (500 μ L, 1.25 M) and shaken at 50°C for 16 h. The reaction mixture was then concentrated under reduced pressure (Genevac HT-4) to provide the desired compound (2 mg, 25.7 mg theoretical, 7.8%). LC-MS m/z 289.3 (M+1).


EXAMPLE 61



10

(Z)-5-((2-(3-aminopyrrolidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: The boc-protected compound was prepared using general displacement procedure 1. The purified boc-protected compound was treated with dichloromethane (1.0 mL), hydrochloric acid in methanol (500 μ L, 1.25 M) and shaken at 50°C for 16 h. The reaction mixture was then concentrated under reduced pressure (Genevac HT-4) to provide the desired compound (3.6 mg, 24.5 mg theoretical, 14.7%). LC-MS m/z 275.3 (M+1).


EXAMPLE 62



(Z)-5-((2-(4-aminopiperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: The boc-protected compound was prepared using general displacement procedure 1. The purified boc-protected compound was treated with dichloromethane (1.0 mL), hydrochloric acid in methanol (500 μ L, 1.25M) and shaken at 50°C for 16 h. The reaction mixture was


then concentrated under reduced pressure (Genevac HT-4) to provide the desired compound (6.1 mg, 25.7 mg theoretical, 23.7%). LC-MS m/z 289.3 (M+1).

EXAMPLE 63




5 **(Z)-5-((2-(3-(aminomethyl)piperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione:** The boc-protected compound was prepared using general displacement procedure 1. The purified boc-protected compound was treated with dichloromethane (1.0 mL), hydrochloric acid in methanol (500 µL, 1.25 M) and shaken at 50°C for 16 h. The reaction mixture was then concentrated under reduced pressure (Genevac HT-4) to provide the
10 desired compound (4.7 mg, 27 mg theoretical, 17.4%). LC-MS m/z 303.3 (M+1).

EXAMPLE 64




15 **(Z)-5-((2-(7-amino-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione:** The boc-protected compound was prepared using general displacement procedure 1 and 1,2,3,4-tetrahydroisoquinolin-7-amine. The boc-protected compound was treated with dichloroethane (1.0 mL), trifluoroacetic acid (1.0 mL)
and shaken at RT for 16 h. The reaction mixture was then concentrated under reduced pressure (Genevac HT-4) and the crude residues were purified using reverse phase HPLC (MS-triggered fraction collection) with a methanol/water gradient and trifluoroacetic acid
20 as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac (HT-4) to provide the desired product (11 mg, 37.6 mg theoretical, 29.2 %). LC-MS m/z 337 (M+1).

EXAMPLE 65




(Z)-5-((2-(4-(aminomethyl)piperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: The boc-protected compound was prepared using general displacement procedure 1 and tert-butyl (piperidin-4-ylmethyl)carbamate. The boc-protected compound was treated with dichloroethane (1.0 mL), trifluoroacetic acid (1.0 mL) and shaken at RT for 16 h. The 5 reaction mixture was then concentrated under reduced pressure (Genevac HT-4) and the crude residues were purified using reverse phase HPLC (MS-triggered fraction collection) with a methanol/water gradient and trifluoroacetic acid as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac (HT-4)) to provide the desired product (37.5 mg, 63.9 mg theoretical, 58.7 %). LC-MS m/z 303 (M+1).

10 EXAMPLE 66



(Z)-5-((2-(5-amino-3,4-dihydroisoquinolin-2(1H)-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: The boc-protected compound was prepared using general displacement procedure 1 and 1,2,3,4-tetrahydroisoquinolin-5-amine. The boc- 15 protected compound was then treated with dichloroethane (1.0 mL), trifluoroacetic acid (1.0 mL) and shaken at RT for 16 h. The reaction mixture was then concentrated under reduced pressure (Genevac HT-4) and the crude residues were purified using reverse phase HPLC (MS-triggered fraction collection) with a methanol/water gradient and trifluoroacetic acid as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac (HT-4)) to provide the desired product (35.1 mg, 37.6 mg theoretical, 93 %). LC- 20 MS m/z 337 (M+1).

EXAMPLE 67




(S,Z)-5-((2-(methyl(piperidin-3-yl)amino)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: The boc-protected compound was prepared using general displacement procedure 1 and (R)-tert-butyl 3-(methylamino)piperidine-1-carboxylate. The boc-protected compound 25

was then treated with dichloroethane (1.0 mL), trifluoroacetic acid (1.0 mL) and shaken at RT for 16 h. The reaction mixture was then concentrated under reduced pressure (Genevac HT-4) and the crude residues were purified using reverse phase HPLC (MS-triggered fraction collection) with a methanol/water gradient and trifluoroacetic acid as the modifier.

- 5 The pure fractions were then concentrated under reduced pressure (Genevac (HT-4) to provide the desired product (16.1 mg, 18.7 mg theoretical, 86 %). LC-MS m/z 303 (M+1).


EXAMPLE 68



(Z)-5-((2-(4-((quinolin-2-ylmethyl)amino)methyl)piperidin-1-yl)pyrimidin-4-

- 10 **yl)methylene)imidazolidine-2,4-dione:** A 2 dram round bottomed vial was charged with (Z)-5-((2-(4-(aminomethyl)piperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (17 mg, 0.056 mmol), quinoline-2-carbaldehyde (8.84 mg, 0.056 mmol, 1 equiv.), diisopropylethylamine (0.1 mL, 0.573 mmol, 10 equiv.), and DMSO (1 mL). The reaction mixture was shaken for 1 h at room temperature then treated with sodium 15 triacetoxyhydroborate (23.83 mg, 0.112 mmol, 2 equiv.) and DCM (0.5 mL). The reaction mixture was shaken overnight. The sample was then filtered through a 0.45 micron syringe filter. The DMSO solution was purified using reverse phase HPLC (MS-triggered fraction collection) with an acetonitrile/water gradient and trifluoroacetic acid as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac HT-4) to provide 20 the desired product (3.2 mg, 24.9 mg theoretical, 12.8 %). LC-MS m/z 444 (M+1).

EXAMPLE 69



(Z)-5-((2-(4-(((6-(thiophen-3-yl)pyridin-2-yl)methyl)amino)methyl)piperidin-1-

yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione: A 2 dram round bottomed vial was charged with (Z)-5-((2-(4-(aminomethyl)piperidin-1-yl)pyrimidin-4-yl)methylene)imidazolidine-2,4-dione (17 mg, 0.056 mmol), 6-(thiophen-3-yl)picinaldehyde (10.64 mg, 0.056 mmol, 1 equiv.), diisopropylethylamine (0.1 mL, 0.573 mmol, 10 equiv.), and DMSO (1 mL). The reaction mixture was shaken for 1 h at room temperature then treated with sodium triacetoxyhydroborate (23.83 mg, 0.112 mmol, 2 equiv.) and DCM (0.5 mL). The reaction mixture was shaken overnight. The sample was then filtered through a 0.45 micron syringe filter. The DMSO solution was purified using reverse phase HPLC (MS-triggered fraction collection) with an acetonitrile/water gradient and triflouoroacetic acid as the modifier. The pure fractions were then concentrated under reduced pressure (Genevac HT-4) to provide the desired product (9.8 mg, 26.7mg theoretical, 36 %). LC-MS m/z 476 (M+1).

EXAMPLE 70

Protocols for Kinase Activity Screening for CK1 γ 1 (h), CK1 γ 2 (h), CK1 γ 3 (h), CK1 δ

15 **(h) and CK1 (y):** Kinase dilution buffer composition: 20 mM MOPS, 1 mM EDTA, 0.01% Brij-35, 5% Glycerol, 0.1% b-mercaptoethanol, 1 mg/mL BSA.

Table 4: Kinase assay ATP concentration within 15 μ M of K_M.

Kinase	K _M (μ M)
CK1 γ 1 (h)	15
CK1 γ 2 (h)	10
CK1 γ 3 (h)	10
CK1 δ (h)	70
CK1 (y)	45

In a final reaction volume of 25 μ L, the compound of interest (at the desired concentration) and the appropriate kinase (5-10 mU) were incubated with 8 mM MOPS pH 7.0, 0.2 mM 20 EDTA, 200 μ M KRRRLS(p)VASLPGL, 10 mM magnesium acetate and [γ -33P-ATP] (specific activity approx. 500 cpm/pmol, concentration as required). The reaction was initiated by the addition of the MgATP mix. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of 5 μ L of a 3% phosphoric acid solution. 10 μ L of the reaction mixture was then spotted onto a P30 filtermat; and washed 25 three times for 5 minutes in 75 mM phosphoric acid, and once in methanol prior to drying and scintillation counting. Tables 5, 6 and 7 show the relative activity of CK1 γ 1(h), CK1 γ 2(h), and CK1 γ 3(h), at varying concentrations of compound 5114.

Table 5. Relative activity of CK1 γ 1(h) at varying concentration of compound 5114.

ATP Concentration: 15 μM

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5114 @ 0.001μM	10040	9047	105	/	/
5114 @ 0.003μM	10254	9261	108	/	/
5114 @ 0.01μM	9722	8729	102	/	/
5114 @ 0.03μM	8374	7381	86	/	/
5114 @ 0.1μM	6373	5380	63	/	/
5114 @ 0.3μM	4497	3504	41	/	/
5114 @ 1μM	3259	2266	26	/	/
5114 @ 3μM	3658	2665	31	/	/
5114 @ 10μM	4178	3185	37	/	/
CONTROL	9207	8582	96	100	6
	9195		96		
	9668		101		
	10233		108		
BLANK	908	/	/	/	/
	1079		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Table 6. Relative activity of CK1γ2(h) at varying concentration of compound 5114.

ATP Concentration: 10 μM

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5114 @ 0.001μM	7649	7041	98	/	/
5114 @ 0.003μM	7313	6705	94	/	/
5114 @ 0.01μM	5658	5050	71	/	/
5114 @ 0.03μM	4532	3924	55	/	/
5114 @ 0.1μM	2530	1922	27	/	/
5114 @ 0.3μM	1328	720	10	/	/
5114 @ 1μM	746	138	2	/	/
5114 @ 3μM	839	231	3	/	/
5114 @ 10μM	998	390	5	/	/
CONTROL	8112	7156	105	100	10
	7521		97		
	8510		110		
	6912		88		
BLANK	582	/	/	/	/
	634		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Table 7. Relative activity of CK1 γ 3(h) at varying concentration of compound 5114.ATP Concentration: 10 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5114 @ 0.001 μ M	9610	9389	88	/	/
5114 @ 0.003 μ M	8980	8759	82	/	/
5114 @ 0.01 μ M	8181	7960	75	/	/
5114 @ 0.03 μ M	8141	7920	75	/	/
5114 @ 0.1 μ M	7310	7089	67	/	/
5114 @ 0.3 μ M	6004	5783	54	/	/
5114 @ 1 μ M	3320	3099	29	/	/
5114 @ 3 μ M	2477	2256	21	/	/
5114 @ 10 μ M	3459	3238	30	/	/
CONTROL	11003	10623	101	100	2
	10644		98		
	11024		102		
	10708		99		
BLANK	190	/	/	/	/
	253		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Tables 8-10 show the relative activity of CK1 γ 2(h) at varying concentration of compounds 5132, 5124, and 5336.

5 Table 8. Relative activity of CK1 γ 2(h) at varying concentration of compound 5132.

ATP Concentration: 10 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5132 @ 0.001 μ M	9557	8769	97	/	/
5132 @ 0.003 μ M	9228	8440	93	/	/
5132 @ 0.01 μ M	8663	7875	87	/	/
5132 @ 0.03 μ M	7659	6871	76	/	/
5132 @ 0.1 μ M	5435	4647	51	/	/
5132 @ 0.3 μ M	2963	2175	24	/	/
5132 @ 1 μ M	1453	665	7	/	/
5132 @ 3 μ M	929	141	2	/	/
5132 @ 10 μ M	747	-41	0	/	/
CONTROL	9907	9040	101	100	1
	9751		99		
	9916		101		
	9736		99		
BLANK	741	/	/	/	/
	835		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Table 9. Relative activity of CK1 γ 2(h) at varying concentration of compound 5124.

ATP Concentration: 10 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5124 @ 0.001 μ M	9126	8301	97	/	/
5124 @ 0.003 μ M	9311	8486	100	/	/
5124 @ 0.01 μ M	9458	8633	101	/	/
5124 @ 0.03 μ M	8389	7564	89	/	/
5124 @ 0.1 μ M	7007	6182	73	/	/
5124 @ 0.3 μ M	5630	4805	56	/	/
5124 @ 1 μ M	2827	2002	23	/	/
5124 @ 3 μ M	1644	819	10	/	/
5124 @ 10 μ M	1122	297	3	/	/
CONTROL	9210	8521	98	100	2
	9429		101		
	9560		103		
	9186		98		
BLANK	806	/	/	/	/
	844		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Table 10. Relative activity of CK1 γ 2(h) at varying concentration of compound 5336.

ATP Concentration: 10 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5336 @ 0.001 μ M	9228	8450	95	/	/
5336 @ 0.003 μ M	9511	8733	98	/	/
5336 @ 0.01 μ M	9197	8419	94	/	/
5336 @ 0.03 μ M	9091	8313	93	/	/
5336 @ 0.1 μ M	8538	7760	87	/	/
5336 @ 0.3 μ M	8060	7282	81	/	/
5336 @ 1 μ M	5306	4528	51	/	/
5336 @ 3 μ M	4439	3661	41	/	/
5336 @ 10 μ M	2029	1251	14	/	/
CONTROL	9855	8938	102	100	2
	9697		100		
	9853		102		
	9459		97		
BLANK	750	/	/	/	/
	806		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Tables 11-14 show the relative activity of CK1(y) at varying concentrations of compounds 5347, 5343, 5363 and 5378

Table 11. Relative activity of CK1(y) at varying concentration of compound 5347.

ATP Concentration: 45 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5347 @ 0.001 μ M	7648	6881	109	/	/
5347 @ 0.003 μ M	7504	6737	107	/	/
5347 @ 0.01 μ M	5664	4897	78	/	/
5347 @ 0.03 μ M	5743	4976	79	/	/
5347 @ 0.1 μ M	5564	4797	76	/	/
5347 @ 0.3 μ M	5163	4396	70	/	/
5347 @ 1 μ M	5328	4561	72	/	/
5347 @ 3 μ M	5835	5068	80	/	/
5347 @ 10 μ M	5163	4396	70	/	/
CONTROL	7344	6308	104	100	9
	6256		87		
	7622		109		
	7079		100		
BLANK	699	/	/	/	/
	835		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

5

Table 12. Relative activity of CK1(y) at varying concentration of compound 5343.

ATP Concentration: 45 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5343 @ 0.001 μ M	7851	7158	98	/	/
5343 @ 0.003 μ M	7723	7030	96	/	/
5343 @ 0.01 μ M	7857	7164	98	/	/
5343 @ 0.03 μ M	7773	7080	97	/	/
5343 @ 0.1 μ M	7344	6651	91	/	/
5343 @ 0.3 μ M	7239	6546	89	/	/
5343 @ 1 μ M	6649	5956	81	/	/
5343 @ 3 μ M	5523	4830	66	/	/
5343 @ 10 μ M	5370	4677	64	/	/
CONTROL	8329	7328	104	100	3
	7923		99		
	8055		100		

	7778		97		
BLANK	651	/	/	/	/
	736		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Table 13. Relative activity of CK1(y) at varying concentration of compound 5363.

ATP Concentration: 45 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5363 @ 0.001 μ M	5899	5159	108	/	/
5363 @ 0.003 μ M	6009	5269	111	/	/
5363 @ 0.01 μ M	6326	5586	117	/	/
5363 @ 0.03 μ M	6257	5517	116	/	/
5363 @ 0.1 μ M	5857	5117	107	/	/
5363 @ 0.3 μ M	6019	5279	111	/	/
5363 @ 1 μ M	5675	4935	104	/	/
5363 @ 3 μ M	6071	5331	112	/	/
5363 @ 10 μ M	5837	5097	107	/	/
CONTROL	5646	4764	103	100	6
	5081		91		
	5723		105		
	5566		101		
BLANK	696	/	/	/	/
	785		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Table 14. Relative activity of CK1(y) at varying concentration of compound 5378.

ATP Concentration: 45 μ M

Sample	Counts	Mean (Counts - Blanks)	Activity (% Control)	Mean	SD*
5378 @ 0.001 μ M	5556	4772	101	/	/
5378 @ 0.003 μ M	6095	5311	113	/	/
5378 @ 0.01 μ M	6043	5259	112	/	/
5378 @ 0.03 μ M	6327	5543	118	/	/
5378 @ 0.1 μ M	6481	5697	121	/	/
5378 @ 0.3 μ M	6601	5817	124	/	/
5378 @ 1 μ M	5981	5197	111	/	/
5378 @ 3 μ M	5886	5102	108	/	/
5378 @ 10 μ M	5553	4769	101	/	/
CONTROL	5836	4702	107	100	6

	5392		98		
	5532		101		
	5187		94		
BLANK	711	/	/	/	/
	858		/		

* NB. Where n = 2, the value reported here is actually range / $\sqrt{2}$

Table 15 summarizes the activity of several casein kinases at 10 μmol concentration of several compounds.

Table 15.

	CK1 γ 1(h)	CK1 γ 2(h)	CK1 γ 3(h)	CK1 δ 1(h)	CK1y
5113	58	25	46	84	64
5117	64	50	70	101	55
5121	37	19	25	85	81
5126	46	43	40	85	99
5132	19	7	31	82	111
5114	24	15	47	88	115
5118	41	17	28	83	99
5122	86	60	104	103	105
5127	56	52	83	89	115
5133	54	47	48	92	106
5115	81	93	81	98	107
5119	78	71	61	100	104
5124	12	4	26	75	114
5128	61	29	53	100	95
5116	92	98	67	98	32
5120	92	104	72	89	94
5125	64	28	68	103	108
5131	70	85	52	94	54
5336	31	8	47	101	90
5337	92	109	73	107	49
5338	93	96	105	106	92
5339	76	51	58	103	36
5340	40	24	51	94	80
5345	73	62	99	107	88
5349	86	83	82	107	113
5353	83	122	71	110	76
5358	80	74	75	105	98
5341	83	100	87	107	107
5346	41	16	61	97	114
5350	66	76	40	94	93
5354	28	11	56	97	101
5343	67	54	61	104	4
5347	80	85	91	103	1

5351	50	10	53	102	2
5355	81	74	69	102	88
5344	84	98	86	104	92
5348	85	80	102	96	31
5352	105	96	86	108	1
5357	81	70	78	105	18
5359	76	81	54	102	85
5376	99	91	76	99	91
5382	87	78	58	100	93
5363	60	67	29	83	0
5378	66	68	37	91	1
5369	94	78	74	87	1
5379	89	71	44	96	92
5371	106	92	88	98	81
5380	105	96	86	96	107

The estimated IC₅₀ values for several compounds are provided in Table 16.

Table 16: Estimated IC₅₀ values.

Compound	Kinase	IC50 (nM)
5114	CK1γ1(h)	165
5114	CK1γ2(h)	31
5114	CK1γ3(h)	307
5132	CK1γ2(h)	114
5124	CK1γ2(h)	349
5336	CK1γ2(h)	1,600
5347	CK1	>10,000
5343	CK1	>10,000
5363	CK1	>10,000
5378	CK1	>10,000

- 5 The relative activity of the kinase as a function of the concentration of several compounds is depicted in Figures 1-10.

EXAMPLE 71: Cell proliferation studies

Cell proliferation studies are summarized in Tables 17 and 18:

10 **Table 17**

No.	CK1γ2†	PAMPA (Pe)	HCT-116‡	HCT-116 *	A549‡	DU145‡	DU145*	MDA-MB-468‡	MDA-MB-468*
5114	31	14	0.8	20.64	5.4	38.8	31.75		x
5124	349	x	4.6	x	-0.7	6.1	x		x
5132	114	x	4.5	x	-1.2	23.2	59.6		x
5336	1,600	x	3.5	x	-0.7	11.3	x		x
5343	10,000	x	3.3	x	-0.9	-1.2	x		x

5347	10,000	x	5.7	x	-2.6	12.7	x		x
5363	10,000	x	5.7	x	-4.3	-18.4	x		x
A	x	x	x	1.29	x	x	1.28	x	3.34
B	x	x	x	x	x	x	x	X	xx

A: Gemcitabine control

B: Sorafenib control

†: IC50 (μ M)‡: % at 10 μ M

5 *: 50 % inhibition concentration

Table 18

No.	HCC 1954 ‡	Caco- 2*	PC-3‡	PC-3*	OVCAR- 3‡	OVCAR- 3*	LNCaP‡	LNCaP*	Jurkat‡
5114	-8.1	x	3.6	x	4.3	64.3	-0.2	51.52	43.4
5124	-17.3	x	-7.1	x	12.5	x	9.6	x	21.3
5132	6.7	x	-18.9	x	-5.8	x	-0.4	73.53	-23.8
5336	-4.9	x	-15.1	x	-15.1	x	0.1	x	28.1
5343	-7.2	x	-9.7	x	-24.2	x	-8.5	x	18.9
5347	0.2	x	-5.5	x	-6.9	x	-12.5	x	27.3
5363	-8.9	x	-2.6	x	-57.4	x	-12.2	x	41.2
A	x	x	x	41.32	x	11.2	x	1.97	3.2
B	x	8.78	x	x	x	x	x	x	x

A: Gemcitabine control

B: Sorafenib control

10 ‡: % at 10 μ M

*: 50 % inhibition concentration

Inhibition of Caco-2 cellsCells: Caco-2cells, ATCC Passage 33, *Mycoplasma* free.

15 Medium: DMEM Medium(GIBCO Cat#11995073) supplemented with 10% fetal bovine serum (Hyclone Cat#SH30396.03).

Seeding: 3,000 cells/well (100 μ L) into 96-wells plate, incubated overnight at 37°C in a humidified, 5% CO₂ atmosphere.Treatment: Test compounds and Sorafenib were prepared in a serial dilution (1:2) by 20 culture medium. Fifty microliters (50 μ L) of diluted compounds were added into each well. The final concentrations of the positive control and test compounds were shown in the graphs. The cells were incubated for 72 hours after addition of the test compounds.MTS: 15 uL of MTS solution(Promega Cat#G5430) was added into each well and incubated with cells for 4 hours at 37°C in a humidified, 5% CO₂ atmosphere.

25 Measurement: Absorbance at 490nm was measured using MD Spectramax Plus384 spectrophotometer. The regression curves were calculated by the GraphPad Prism 5 using "log(inhibitor) vs response--variable slope".

Calculation: % of inhibition=(ODzero ctrl-OD compound)/ODzero ctrl*100. Results are shown in table 19.

Table 19.

Compound ID	IC50 NO.	Concentration (μM)	OD value		% of inhibition	
			1	2	1	2
5114	5	0.000	0.557	0.584	0.00	0.00
		0.400	0.557	0.527	-0.04	9.74
		0.800	0.536	0.567	3.77	2.96
		1.600	0.557	0.619	0.02	-5.97
		3.200	0.627	0.596	-12.57	-2.05
		6.250	0.521	0.633	6.43	-8.41
		12.500	0.467	0.606	16.16	-3.66
		25.000	0.452	0.544	18.94	6.95
		50.000	0.414	0.463	25.73	20.73
		100.000	0.420	0.461	24.58	21.01
5124	6	0.000	0.608	0.640	0.00	0.00
		0.400	0.558	0.626	1.63	2.19
		0.800	0.489	0.553	3.14	13.46
		1.600	0.619	0.586	-1.88	8.30
		3.200	0.606	0.657	0.33	-2.77
		6.250	0.543	0.555	2.47	13.29
		12.500	0.597	0.522	1.73	18.37
		25.000	0.570	0.557	6.17	12.93
		50.000	0.532	0.507	12.52	20.03
		100.000	0.428	0.446	29.67	30.30
5132	7	0.000	0.600	0.619	0.00	0.00
		0.400	0.578	0.573	3.63	7.44
		0.800	0.535	0.618	7.53	0.18
		1.600	0.549	0.646	8.60	-4.33
		3.200	0.607	0.616	-1.22	0.48
		6.250	0.591	0.622	1.52	-0.39
		12.500	0.572	0.622	4.77	-0.39

		25.000	0.593	0.663	1.13	-7.09
		50.000	0.581	0.604	3.13	2.42
		100.000	0.592	0.595	1.37	3.89
Sorafenib	8	0.000	0.619	0.697	0.000	0.000
		0.400	0.618	0.695	0.049	0.315
		0.800	0.557	0.654	9.975	6.237
		1.600	0.530	0.618	14.292	11.442
		3.200	0.513	0.632	17.008	9.392
		6.250	0.496	0.534	19.886	23.457
		12.500	0.144	0.166	76.762	76.265
		25.000	(0.003)	0.010	100.463	98.503
		50.000	(0.015)	0.007	102.371	99.020
		100.000	0.099	0.023	84.053	96.668

Inhibition of DU145 cells

Cells: DU145 cells, ATCC Passage unknown, *Mycoplasma* free.

Medium: DMEM Medium(GIBCO Cat#11995073) supplemented with 10% fetal bovine serum(Hyclone Cat#SH30396.03).

Seeding: 750 cells/well (100 μ L) into 96-wells plate, incubated overnight at 37°C in a humidified, 5% CO₂ atmosphere.

Treatment: Test compounds and Gemcitabine were prepared in a serial dilution (1:2) by culture medium. Fifty microliters (50 μ L) of serially diluted solutions were added into each well. The final concentrations of the positive control and test compounds were shown in the graphs. The cells were incubated for 72 hours after addition of the test compounds..

MTS: 15 μ L of MTS solution(Promega Cat#G5430) was added into each well and incubated with cells for 4 hours at 37°C in a humidified, 5% CO₂ atmosphere.

Measurement: Absorbance at 490nm was measured using MD Spectramax Plus384 spectrophotometer. The regression curves were calculated by the GraphPad Prism 5 using "log(inhibitor) vs response--variable slope".

Calculation: % of inhibition=(ODzero ctrl-OD compound)/ODzero ctrl*100
Results are shown in Table 20.

Table 20

Compound ID	IC50 NO.	Concentration (uM)	OD value		% of inhibition	
			1	2	1	2
5114	5	0.000	1.066	1.001	0.00	0.00
		0.400	1.060	0.981	0.53	1.99
		0.800	1.122	0.873	-5.24	12.76
		1.600	0.887	0.966	16.79	3.54
		3.200	0.967	0.937	9.32	6.41
		6.250	0.819	0.851	23.21	15.02
		12.500	0.670	0.793	37.17	20.75
		25.000	0.461	0.481	56.73	51.92
		50.000	0.420	0.442	60.60	55.86
		100.000	0.327	0.336	69.34	66.47
5124	6	0.000	1.006	1.011	0.00	0.00
		0.400	1.018	1.023	-1.21	-1.28
		0.800	0.957	0.943	4.86	6.68
		1.600	0.985	0.910	2.10	9.96
		3.200	0.967	0.861	3.86	14.79
		6.250	0.892	0.880	11.37	12.96
		12.500	0.927	0.976	7.89	3.41
		25.000	0.932	0.956	7.32	5.35
		50.000	0.948	0.845	5.73	16.39
		100.000	0.791	0.768	21.41	24.06
5132	7	0.000	1.217	1.070	0.00	0.00
		0.400	0.993	0.981	18.37	8.34
		0.800	1.080	1.020	11.22	4.67
		1.600	1.000	0.902	17.86	15.71
		3.200	1.043	0.943	14.27	11.95
		6.250	1.016	0.948	16.49	11.42
		12.500	0.904	0.904	25.67	15.54
		25.000	0.695	0.723	42.88	32.45
		50.000	0.591	0.631	51.42	41.09
		100.000	0.555	0.494	54.41	53.82

Gemcitabine	8	0.000	1.1367875	1.042	0.000	0.000
		0.400	0.9392875	0.880	17.374	15.551
		0.800	0.7368875	0.699	35.178	32.934
		1.600	0.4662875	0.477	58.982	54.261
		3.200	0.0513875	0.039	95.480	96.223
		6.250	0.0359875	0.039	96.834	96.223
		12.500	0.0385875	0.028	96.606	97.297
		25.000	0.0345875	0.040	96.957	96.203
		50.000	0.0299875	0.012	97.362	98.870
		100.000	0.0235875	0.029	97.925	97.172

Inhibition of HCT116 cells

Cells: HCT116 cells, ATCC Passage unknown, *Mycoplasma* free.

Medium: DMEM Medium(GIBCO Cat#11995073) supplemented with 10% fetal bovine serum(Hyclone Cat#SH30396.03).

Seeding: 750 cells/well (100 µL) into 96-wells plate, incubated overnight at 37°C in a humidified, 5% CO₂ atmosphere.

Treatment: Test compounds and Gemcitabine were prepared in a serial dilution (1:2) by culture medium. Fifty microliters (50 uL) of serially diluted solutions were added into each well. The final concentrations of the positive control and test compounds were shown in the graphs. The cells were incubated for 72 hours after addition of the test compounds.

MTS: 15 uL of MTS solution (Promega Cat#G5430) was added into each well and incubated with cells for 4 hours at 37°C in a humidified, 5% CO₂ atmosphere.

Measurement: Absorbance at 490 nm was measured using MD Spectramax Plus384 spectrophotometer. The regression curves were calculated by the GraphPad Prism 5 using "log(inhibitor) vs response--variable slope".

Calculation: % of inhibition=(ODzero ctrl-OD compound)/ODzero ctrl*100.

Results are shown in Table 21.

Table 21.

Compound ID	IC50 NO.	Concentration (uM)	OD value		% of inhibition	
			1	2	1	2
5114	5	0.000	0.766	0.964	0.00	0.00
		0.400	0.707	0.936	7.74	2.98

		0.800	0.671	0.972	12.43	-0.80
		1.600	0.709	0.862	7.49	10.57
		3.200	0.654	0.816	14.72	15.37
		6.250	0.747	0.888	2.53	7.91
		12.500	0.664	0.803	13.36	16.69
		25.000	0.217	0.270	71.62	72.02
		50.000	0.145	0.163	81.12	83.12
		100.000	0.141	0.139	81.56	85.60
5124	6	0.000	0.812	0.904	0.00	0.00
		0.400	0.760	0.882	6.37	2.40
		0.800	0.758	1.016	6.59	-12.40
		1.600	0.761	0.865	6.19	4.34
		3.200	0.731	0.844	9.98	6.58
		6.250	0.768	0.894	5.41	1.08
		12.500	0.763	0.879	6.05	2.73
		25.000	0.722	0.796	11.05	11.89
		50.000	0.644	0.834	20.69	7.77
		100.000	0.635	0.766	21.74	15.28
5132	7	0.000	0.798	0.860	0.00	0.00
		0.400	0.750	0.920	5.97	-6.93
		0.800	0.859	0.878	-7.71	-2.01
		1.600	0.799	0.872	-0.15	-1.36
		3.200	0.834	0.791	-4.53	8.02
		6.250	0.822	0.740	-3.03	13.95
		12.500	0.801	0.747	-0.40	13.15
		25.000	0.732	0.663	8.27	22.98
		50.000	0.648	0.628	18.74	27.03
		100.000	0.580	0.630	27.24	26.79
Gemcitabine	8	0.000	0.985	0.968	0.00	0.00
		0.400	0.799	0.766	18.84	20.88
		0.800	0.619	0.723	37.19	25.28
		1.600	0.452	0.535	54.10	44.71

	3.200	0.083	0.083	91.60	91.43
	6.250	0.076	0.065	92.29	93.27
	12.500	0.065	0.063	93.39	93.46
	25.000	0.075	0.061	92.39	93.66
	50.000	0.036	0.046	96.36	95.22
	100.000	0.037	0.037	96.28	96.22

Inhibition of PC-3 cells:

Cells: PC-3 cells, ATCC Passage unknown, *Mycoplasma* free.

Medium: DMEM Medium(GIBCO Cat#11995073) supplemented with 10% fetal bovine serum(Hyclone Cat#SH30396.03).

Seeding: 3,000 cells/well (100ul) into 96-wells plate, incubated overnight at 37°C in a humidified, 5% CO₂ atmosphere.

Treatment: Test compounds and Gemcitabine were prepared in a serial dilution (1:2) by culture medium. Fifty microliters (50 uL) of serially diluted solutions were added into each well. The final concentrations of the positive control and test compounds were shown in the graphs. The cells were incubated for 72 hours after addition of the test compounds.

MTS: 15 uL of MTS solution(Promega Cat#G5430) was added into each well and incubated with cells for 4 hours at 37°C in a humidified, 5% CO₂ atmosphere.

Measurement: Absorbance at 490nm was measured using MD Spectramax Plus384 spectrophotometer. The regression curves were calculated by the GraphPad Prism 5 using "log(inhibitor) vs response--variable slope".

Calculation: % of inhibition=(ODzero ctrl-OD compound)/ODzero ctrl*100.

Results are shown in table 22.

Table 22.

Compound ID	IC50 NO.	Concentration (uM)	OD value		% of inhibition	
			1	2	1	2
5114	5	0.000	0.776	0.848	0.00	0.00
		0.400	0.817	0.814	-5.35	3.99
		0.800	0.782	0.786	-0.89	7.32
		1.600	0.721	0.736	7.03	13.20
		3.200	0.735	0.726	5.17	14.31
		6.250	0.685	0.735	11.63	13.27
		12.500	0.761	0.738	1.84	12.92
		25.000	0.683	0.650	11.91	23.26
		50.000	0.562	0.562	27.57	33.69
		100.000	0.391	0.492	49.54	41.90
5124	6	0.000	0.811	0.675	0.00	0.00
		0.400	0.829	0.630	-2.24	6.67
		0.800	0.836	0.728	-3.14	-7.90
		1.600	0.725	0.682	10.56	-1.11
		3.200	0.813	0.718	-0.25	-6.45
		6.250	0.698	0.743	13.97	-10.18
		12.500	0.819	0.695	-0.94	-3.05
		25.000	0.756	0.703	6.75	-4.18
		50.000	0.692	0.655	14.71	2.96
		100.000	0.839	0.721	-3.43	-6.83
5132	7	0.000	0.854	0.743	0.00	0.00
		0.400	0.817	0.680	4.29	8.53
		0.800	0.850	0.759	0.42	-2.17
		1.600	0.827	0.745	3.11	-0.23
		3.200	0.842	0.728	1.36	2.07
		6.250	0.794	0.783	7.03	-5.42
		12.500	0.815	0.787	4.58	-5.89
		25.000	0.771	0.768	9.66	-3.36
		50.000	0.793	0.694	7.11	6.64

		100.000	0.893	0.771	-4.53	-3.77
Gemcitabine	8	0.000	0.903	0.883	0.00	0.00
		0.400	0.884	0.866	2.10	1.92
		0.800	0.860	0.803	4.80	9.05
		1.600	0.794	0.778	12.00	11.91
		3.200	0.726	0.756	19.70	14.35
		6.250	0.686	0.692	24.10	21.66
		12.500	0.617	0.628	31.70	28.93
		25.000	0.572	0.583	36.70	33.96
		50.000	0.458	0.488	49.30	44.46
		100.000	0.246	0.214	72.80	75.81

Inhibition of LNCaP cells

Cells: LNCaP, ATCC Passage unknown, *Mycoplasma* free.

Medium: RPMI-1640 Medium(GIBCO Cat#22400121) supplemented with 10% fetal

5 bovine serum(Hyclone Cat#SH30396.03).

Seeding: 3,000 cells/well (100 uL) into 96-wells plate, incubated overnight at 37°C in a humidified, 5% CO₂ atmosphere.

Treatment: Test compounds and Gemcitabine were prepared in a serial dilution (1:2) by culture medium. Fifty microliters (50 uL) of serially diluted solutions were added into each 10 well. The final concentrations of the positive control and test compounds were shown in the graphs. The cells were incubated for 72 hours after addition of the test compounds.

MTS: 15 uL of MTS solution(Promega Cat#G5430) was added into each well and incubated with cells for 4 hours at 37°C in a humidified, 5% CO₂ atmosphere.

Measurement: Absorbance at 490nm was measured using MD Spectramax Plus384 15 spectrophotometer. The regression curves were calculated by the GraphPad Prism 5 using "log(inhibitor) vs response--variable slope".

Calculation: % of inhibition +(AVE zero ctrl - AVE compound)/AVE zero ctrl * 100.

Results are shown in Table 23:

Table 23.

Compound ID	IC50 NO.	Concentration (uM)	OD value		% of inhibition	
			1	2	1	2
5114	5	0.000	0.630	0.687	0.00	0.00
		0.400	0.609	0.688	3.33	-0.15
		0.800	0.541	0.593	14.12	13.69
		1.600	0.568	0.594	9.84	13.54
		3.200	0.558	0.638	11.43	7.13
		6.250	0.560	0.593	11.11	13.69
		12.500	0.537	0.602	14.76	12.37
		25.000	0.471	0.496	25.23	27.81
		50.000	0.306	0.386	51.42	43.82
		100.000	0.164	0.195	73.95	71.63
5124	6	0.000	0.615	0.710	0.00	0.00
		0.400	0.604	0.687	1.79	3.24
		0.800	0.523	0.602	14.96	15.21
		1.600	0.593	0.643	3.58	9.44
		3.200	0.538	0.636	12.52	10.42
		6.250	0.556	0.614	9.59	13.52
		12.500	0.556	0.636	9.59	10.42
		25.000	0.577	0.588	6.18	17.19
		50.000	0.592	0.641	3.74	9.72
		100.000	0.615	0.651	0.00	8.31
5132	7	0.000	0.693	0.752	0.00	0.00
		0.400	0.668	0.753	3.61	-0.13
		0.800	0.645	0.679	6.93	9.71
		1.600	0.675	0.724	2.60	3.72
		3.200	0.653	0.706	5.77	6.12
		6.250	0.639	0.705	7.79	6.25
		12.500	0.658	0.712	5.05	5.32
		25.000	0.678	0.681	2.16	9.44
		50.000	0.600	0.625	13.42	16.89

		100.000	0.322	0.333	53.53	55.73
Gemcitabine	8	0.000	0.663	0.766	0.00	0.00
		0.400	0.578	0.662	12.82	13.58
		0.800	0.459	0.434	30.76	43.35
		1.600	0.448	0.472	32.42	38.39
		3.200	0.322	0.337	51.42	56.01
		6.250	0.182	0.190	72.54	75.21
		12.500	0.166	0.173	74.95	77.43
		25.000	0.101	0.177	84.75	76.91
		50.000	0.124	0.202	81.28	73.64
		100.000	0.112	0.197	83.09	74.29

Inhibition of MDA-MB-468 cells

Cells: MDA-MB-468 cells, ATCC Passage unknown, *Mycoplasma* free.

Medium: RPMI-1640 Medium(GIBCO Cat#22400121) supplemented with 10% fetal

5 bovine serum(Hyclone Cat#SH30396.03).

Seeding: 2,000 cells/well (100ul) into 96-wells plate, incubated overnight at 37°C in a humidified, 5% CO₂ atmosphere.

10 **Treatment:** Test compounds and Gemcitabine were prepared in a serial dilution (1:2) by culture medium. Fifty microliters (50 uL) of serially diluted solutions were added into each well. The final concentrations of the positive control and test compounds were shown in the graphs. The cells were incubated for 72 hours after addition of the test compounds.

MTS: 15 uL of MTS solution(Promega Cat#G5430) was added into each well and incubated with cells for 4 hours at 37°C in a humidified, 5% CO₂ atmosphere.

15 **Measurement:** Absorbance at 490nm was measured using MD Spectramax Plus384 spectrophotometer. The regression curves were calculated by the GraphPad Prism 5 using "log(inhibitor) vs response--variable slope".

Calculation: % of inhibition=(ODzero ctrl-OD compound)/ODzero ctrl*100

Results are shown in Table 24:

Table 24.

Compound ID	IC50 NO.	Concentration (μM)	OD value		% of inhibition	
			1	2	1	2
5114	5	0.000	0.381	0.391	0.00	0.00
		0.400	0.352	0.332	7.61	15.08
		0.800	0.351	0.317	7.88	18.91
		1.600	0.342	0.373	10.24	4.60
		3.200	0.303	0.335	20.48	14.31
		6.250	0.282	0.290	25.99	25.81
		12.500	0.292	0.295	23.37	24.53
		25.000	0.257	0.249	32.56	36.28
		50.000	0.250	0.267	34.39	31.68
		100.000	0.252	0.246	33.87	37.05
5124	6	0.000	0.347	0.343	0.00	0.00
		0.400	0.317	0.347	8.65	-1.16
		0.800	0.326	0.306	6.05	10.78
		1.600	0.374	0.350	-7.78	-2.04
		3.200	0.281	0.358	19.03	-4.37
		6.250	0.303	0.294	12.68	14.27
		12.500	0.308	0.325	11.24	5.24
		25.000	0.268	0.268	22.77	21.84
		50.000	0.280	0.285	19.32	16.89
		100.000	0.254	0.233	26.81	32.03
5132	7	0.000	0.378	0.378	0.00	0.00
		0.400	0.315	0.353	16.67	6.61
		0.800	0.341	0.332	9.79	12.16
		1.600	0.315	0.345	16.67	8.72
		3.200	0.315	0.337	16.67	10.84
		6.250	0.318	0.312	15.88	17.44
		12.500	0.293	0.324	22.49	14.27
		25.000	0.291	0.328	23.02	13.21
		50.000	0.360	0.402	4.76	-6.34
		100.000	0.430	0.385	-13.76	-1.85

Gemcitabine	8	0.000	0.333	0.341	0.00	0.00
		0.400	0.313	0.337	6.01	1.17
		0.800	0.275	0.310	17.42	9.08
		1.600	0.201	0.264	39.65	22.56
		3.200	0.151	0.192	54.68	43.65
		6.250	0.101	0.105	69.70	69.13
		12.500	0.066	0.053	80.21	84.36
		25.000	0.033	0.035	90.12	89.64
		50.000	0.064	0.064	80.81	81.14
		100.000	0.034	0.007	89.82	97.84

Inhibition of Ovar 3 cells**Cells:** Ovar-3 cells, ATCC Passage 4, *Mycoplasma* free.**Medium:** RPMI-1640 Medium(GIBCO Cat#22400121) supplemented with 10% fetal bovine serum(Hyclone Cat#SH30396.03).**Seeding:** 3,000 cells/well (100 uL) were seeded into 96-wells plate, incubated overnight at 37°C in a humidified, 5% CO₂ atmosphere.**Treatment:** Test compounds and Gemcitabine were prepared in a serial dilution (1:2) by culture medium. Fifty microliters (50 uL) of serially diluted solutions were added into each well. The final concentrations of the positive control and test compounds were shown in the graphs. The cells were incubated for 72 hours after addition of the test compounds.**MTS:** 15 uL of MTS solution(Promega Cat#G5430) was added into each well and incubated with cells for 4 hours at 37°C in a humidified, 5% CO₂ atmosphere.**Measurement:** Absorbance at 490nm was measured using MD Spectramax Plus384 spectrophotometer. The regression curves were calculated by the GraphPad Prism 5 using "log(inhibitor) vs response--variable slope".**Calculation:** % of inhibition=(ODzero ctrl-OD compound)/ODzero ctrl*100

Results are shown in Table 25:

Table 25.

Compound ID	IC50 NO.	Concentration (μM)	OD value		% of inhibition	
			1	2	1	2
5114	5	0.000	0.660	0.586	0.00	0.00
		0.400	0.707	0.553	-7.13	5.71

		0.800	0.587	0.505	11.07	13.83
		1.600	0.581	0.584	11.87	0.34
		3.200	0.565	0.571	14.40	2.56
		6.250	0.540	0.519	18.19	11.44
		12.500	0.530	0.485	19.71	17.25
		25.000	0.345	0.345	47.75	41.15
		50.000	0.345	0.272	47.75	53.62
		100.000	0.339	0.281	48.66	52.08
5124	6	0.000	0.605	0.567	0.00	0.00
		0.400	0.609	0.553	-0.79	2.45
		0.800	0.557	0.471	7.94	16.94
		1.600	0.464	0.521	23.32	8.12
		3.200	0.533	0.480	11.91	15.35
		6.250	0.439	0.423	27.46	25.41
		12.500	0.497	0.453	17.86	20.12
		25.000	0.364	0.346	39.86	39.00
		50.000	0.353	0.384	41.68	32.30
		100.000	0.263	0.351	56.56	38.12
5132	7	0.000	0.560	0.552	0.00	0.00
		0.400	0.575	0.451	-2.68	18.31
		0.800	0.557	0.467	0.54	15.41
		1.600	0.492	0.511	12.15	7.43
		3.200	0.551	0.526	1.61	4.71
		6.250	0.464	0.485	17.15	12.15
		12.500	0.645	0.550	-15.19	0.36
		25.000	0.542	0.563	3.22	-1.99
		50.000	0.600	0.574	-7.15	-3.99
		100.000	0.567	0.531	-1.26	3.81
Gemcitabine	8	0.000	0.600	0.548	0.00	0.00
		0.400	0.591	0.575	1.50	-4.93
		0.800	0.490	0.460	18.34	16.07
		1.600	0.440	0.453	26.64	17.35

		3.200	0.411	0.432	31.52	21.18
		6.250	0.363	0.388	39.52	29.22
		12.500	0.257	0.234	57.20	57.34
		25.000	0.040	0.009	93.39	98.43
		50.000	0.020	0.010	96.73	98.09
		100.000	0.012	0.011	97.97	98.02

A summary of the cell line data, including IC50's, 50 % viability concentration and maximal inhibition % is provided in Table 26:

Table 26.

cell line	compound ID	IC50(µM)	50% viability conc.	Maximal 5 inhibition %
Caco-2	5114	22.1	N.D.	22.8
	5124	N.D.	397.5	N.D.
	5132	N.D.	N.D.	N.D.
	Sorafenib	9.367	8.767	100
DU145	5114	N.D.	31.75	N.D.
	5124	N.D.	N.D.	N.D. 10
	5132	N.D.	59.6	N.D.
	Gemcitabine	1.167	1.28	97.55
HCT116	5114	13.6	20.64	83.58
	5124	N.D.	N.D.	N.D.
	5132	N.D.	N.D.	N.D.
	Gemcitabine	1.42	1.293	96.25
PC-3	5114	N.D.	143.7	N.D. 15
	5124	N.D.	N.D.	N.D.
	5132	N.D.	N.D.	N.D.
	Gemcitabine	N.D.	41.32	N.D.
LNCaP	5114	N.D.	51.52	N.D.
	5124	N.D.	N.D.	N.D.
	5132	N.D.	73.53	N.D. 20
	Gemcitabine	2.38	1.967	78.69
MDA-MB-468	5114	3.704	N.D.	38.46
	5124	17.48	N.D.	29.42
	5132	N.D.	N.D.	N.D.
	Gemcitabine	2.526	3.34	93.83
Ovcar3	5114	13.97	64.28	50.37
	5124	N.D.	115.9	N.D. 25
	5132	N.D.	N.D.	N.D.
	Gemcitabine	9.871	11.18	98

EXAMPLE 72

In Vitro ADME assays of PAMPA and human and rat hepatic microsomal stability.

The generic gradient HPLC and MS method summarized in Table 27 was used for the analysis of compound 5114.

5 **Table 27. HPLC conditions.**

Instrument	Applied Biosystems API 4000 mass spectrometer		
Ionization Mode	Electrospray, positive ions		
MRM	5114: 365.2 → 232.1		
Column	ACE 2 C18, 2.1x50mm, 3µm Part Number ACE-111-0502		
Eluent A	2 mM ammonium acetate, 0.1% formic acid in 95:5 water:methanol		
Eluent B	2 mM ammonium acetate, 0.1% formic acid in 95:5 methanol:water		
Pump Gradient Program	Time (min)	%A	%B
	0	75	25
	0.50	75	25
	1.00	0	100
	2.00	0	100
	2.10	75	25
	2.50	75	25
Flow (mL/min)	0.5		
Column Temperature	Ambient		
Injection Volume	3-30		
Sample Temperature	Ambient		
Run Time (min)	2.5		

- Parallel artificial membrane permeability assays (PAMPA) were performed with compound 5114. The target concentration in the assay was 10 µM, prepared by diluting (1000-fold) the 10 mM stock solutions in DMSO into PBS, pH 7.4. The final DMSO concentration was 10 0.1%. The 10 µM solutions were added, 300 µL, to wells in the donor plate. The receiver plate, which contained 200 µL of PBS, pH 7.4 per well, was placed in the donor plate and the assembly was incubated for 5 hours at ambient temperature. At the end of the incubation period the plates were separated and the compound concentrations in each solution were determined by LC/MS/MS. The assay was performed in triplicate.
- 15 Dexamethasone and verapamil were used as reference compounds. The permeability, P_c , and mass retention, R , of each compound were calculated using the following equations, and the results are summarized in Table 28. The results for dexamethsone and verapamil were consistent with historical data.

$$P_e = \frac{-\ln[1 - C_{A(t)}/C_E] \times 10^7}{A \times (1/V_D + 1/V_A) \times t}$$

$$R = 1 - \frac{C_{D(t)}V_D + C_{A(t)}V_A}{C_0V_D}$$

Where:

- C0 is the initial concentration in the donor well (μM)
- CD(t) is the concentration in the donor well after incubation (μM)
- 5 CA(t) is the concentration in the acceptor well after incubation (μM)
- VD is the volume in the donor well (0.3 mL)
- VA is the volume in the acceptor well (0.2 mL)
- CE is $(CD(t)VD + CA(t)VA)/(VD + VA)$
- A is the filter area (0.3 cm²)
- 10 t is the incubation time (18,000 s).

Table 28: PAMPA Assay data summary.

Compound	Permeability P_e (nm/s)	Mass Retention R (%)
5114	14	0
Verapamil	75	20
Dexamethasone	9.0	9

15

Hepatic microsomal assays were performed with compound 5114 in human and rat (Sprague-Dawley). Protein concentrations of 0.4 (human) and 0.2 mg/mL (rat) with an NADPH regenerating cofactor system (2.6 mM NADP+, 6.6 mM glucose-6-phosphate, 0.8 U/mL glucose-6-phosphate dehydrogenase, and 6.6 mM magnesium chloride) were used.

- 20 A 100 μM 20% DMSO/80% acetonitrile working stock of each of the compounds was diluted 100 fold resulting in 1 μM compound/1% final organic reaction concentrations. Time points were removed at 0 and 60 minutes. At each time point, 100 μL of the incubation suspension was added to 200 μL of acetonitrile containing internal standard (tolbutamide), followed by centrifugation at 3,220 rcf for 10 minutes. Two hundred (200)
- 25 μL of the resulting supernatants were removed, dried under nitrogen and reconstituted in 100 μL of 2 mM ammonium acetate, 0.1% formic acid in 50% methanol prior to analysis by LC/MS/MS. Testosterone and dexamethasone were used as reference compounds. Table 29 summarizes the results. The results for testosterone and dexamethasone were consistent with historical data.

Table 29: Hepatic microsomal stability summary

Compound	% remaining after incubation	
	Rat Microsomes	Human Microsomes
5114	6.4	71
Testosterone	0.6	42
Dexamethasone	91	85

Materials used are summarized in Table 30.

Table 30: Materials.

Material	Supplier	Part No.	Lot No.
Testosterone	Sigma	T1500	087K1440
Dexamethasone	Sigma	D1756	096K1805
Verapamil	Aldrich	381195	12731MA
Tolbutamide	Sigma	T0891	076K1277
PBS	Sigma	P3813	096K8204
Ammonium acetate	J.T. Baker	0599-08	E49H15
Formic acid	Acros Organics	147930250	AO266198
Acetonitrile	EMD	AX0145-1	49099
DMSO	Alfa Aeser	32434	D04R008
Isopropanol	J.T. Baker	9827-03	C38H23
Methanol	EMD	MX0486-1	49178
0.5 M Potassium Phosphate pH 7.4	BD Gentest	451201	06123
PAMPA plate	BD Gentest	353015	431256
Human microsomes	BD Gentest	452161	18888
Rat microsomes	BD Gentest	452501	21027
NADPH Regeneration System Solution A	BD Gentest	451220	51893
NADPH Regeneration System Solution B	BD Gentest	451220	47758
water	House DI (Barnstead Nanopure)		

5

LC/MS equipment:

Chromatograph: Shimadzu LC-20 AD

Autosampler: CTC HTS PAL

MS: API 4000

Software System: Analyst Software, Version 1.4.2.

EXAMPLE 73

Protocols for Kinase Activity Screening: Kinase screening was performed by Millipore UK Ltd. Kinase dilution buffer composition: 20 mM MOPS, 1 mM EDTA, 0.01% Brij-35, 5% Glycerol, 0.1% β -mercaptoethanol, 1 mg/mL BSA.

The percent kinase activity of different kinases is measured after treatment with various compounds listed in Table 31.

Table 31: Percent Kinase Activity of Enzyme When Treated with 300 nM of Compound

Example	CK1g2	CK1	CK2	Pim-1	Pim-2	Pim-3
64	75	81	65	83	81	97
65	96	97	94	92	78	76
66	77	98	78	91	86	96
67	111	104	94	98	99	114
68	84	97	109	75	69	63
69	76	92	106	46	86	59

10

EXAMPLE 74

Selected Cell Proliferation Inhibition Data

LNCaP cells

Cells: LNCaP, ATCC Passage unknown, *Mycoplasma* free.

15 **Medium:** RPMI-1640 Medium (GIBCO Cat#31800-022) supplemented with 10% fetal bovine serum (Hyclone Cat#SH30396.03).

Seeding: 3,000 cells/well (100 μ L) into 96-well plates, incubated overnight at 37°C in a humidified 5% CO₂ atmosphere.

DU145 cells

20 **Cells:** DU145 cells, ATCC Passage unknown, *Mycoplasma* free.

Medium: DMEM Medium (GIBCO Cat#11995073) supplemented with 10% fetal bovine serum (Hyclone Cat#SH30396.03).

Seeding: 750 cells/well (100 µL) into 96-well plates, incubated overnight at 37°C in a humidified 5% CO₂ atmosphere.

5 **K562 cells**

Cells: K562, ATCC Passage unknown, *Mycoplasma* free.

Medium: RPMI-1640 Medium (GIBCO Cat#31800-022) supplemented with 10% fetal bovine serum (Hyclone Cat#SH30396.03).

10 **Seeding:** 3,000 cells/well (100 µL) into 96-well plates, incubated overnight at 37°C in a humidified 5% CO₂ atmosphere.

MV4-11

Cells: MV4-11, ATCC Passage unknown, *Mycoplasma* free.

Medium: IMDM Medium (GIBCO Cat#31800-02231800-022) supplemented with 10% fetal bovine serum (Hyclone Cat#SH30396.03).

15 **Seeding:** 3,000 cells/well (100 µL) into 96-well plates, incubated overnight at 37°C in a humidified 5% CO₂ atmosphere.

RPMI-8226

Cells: **RPMI-8226**, 3,000 cells/well (100 µL) into 96-well plates, incubated overnight at 37°C in a humidified 5% CO₂ atmosphere.

20 **Medium:** RPMI-1640 Medium (GIBCO Cat#31800-022) supplemented with 10% fetal bovine serum (Hyclone Cat#SH30396.03).

Seeding: 3,000 cells/well (100 µL) into 96-well plates, incubated overnight at 37°C in a humidified 5% CO₂ atmosphere.

25 **Treatment:** Test compounds were first diluted 333-fold in the medium. Fifty microliters (50 µL) of diluted compounds were added into each well (i.e., another 3-fold dilution). The final concentration of test compounds was 10 µM. The final concentrations of the positive control (Gemcitabine, also 50 µL) added in each well) is shown in Figure 48. The cells were incubated for 72 hours after addition of the test compounds.

MTS: Added 20 µL of MTS solution (Promega Cat #G5430) into each well and incubated for 4 hours.

Measurement: Absorbance at 490 nm using MD Spectramax Plus 384 spectrophotometer.

Calculation: % of inhibition + (AVE zero ctrl - AVE compound)/AVE zero ctrl * 100.

- 5 Table 32 shows IC50 values (µM) for the inhibition of specific cell lines using various compounds.

Table 32: Cell Proliferation Data (MTS)

COMPOUND	DU145	LNCaP	K562	MV4-11	RPMI-8226
10641					
10642	10.07			13.35	
10643	20.34	15.38	47.83	6.76	7.47
10647					

INCORPORATION BY REFERENCE

- 10 All of the U.S. patents and U.S. published patent applications cited herein are hereby incorporated by reference.


EQUIVALENTS

While several embodiments of the present invention have been described and illustrated herein, those of ordinary skill in the art will readily envision a variety of other 15 means and/or structures for performing the functions and/or obtaining the results and/or one or more of the advantages described herein, and each of such variations and/or modifications is deemed to be within the scope of the present invention. More generally, those skilled in the art will readily appreciate that all parameters, dimensions, materials, and configurations described herein are meant to be exemplary and that the actual parameters, 20 dimensions, materials, and/or configurations will depend upon the specific application or applications for which the teachings of the present invention is/are used. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. It is, therefore, to be understood that the foregoing embodiments are presented by way of 25 example only and that, within the scope of the appended claims and equivalents thereto, the

invention may be practiced otherwise than as specifically described and claimed. The present invention is directed to each individual feature, system, article, material, kit, and/or method described herein. In addition, any combination of two or more such features, systems, articles, materials, kits, and/or methods, if such features, systems, articles,
5 materials, kits, and/or methods are not mutually inconsistent, is included within the scope of the present invention.

We Claim:

1. A compound of formula 1:



- 5 or a pharmaceutically acceptable salt thereof, wherein independently for each occurrence:
- W and X are independently oxygen or sulfur;
- Z¹, Z² and Z³ are independently C-R²¹ or N, provided that at least one of Z¹ and Z² is N;
- R¹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aralkyl, heteroaralkyl, -COR⁶, and -C(O)OR⁶;
- 10 R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷);
- R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷); or R² and R³ are joined together to form an optionally substituted heterocyclic ring;
- 15 R⁴ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocyclylalkyl, aralkyl, heteroaryl, heteroaralkyl, halo, hydroxy, alkoxy, hydroxyalkyl, and alkoxyalkyl;
- R⁵ is selected from the group consisting of aryl, heteroaryl, heterocyclyl, -N(R⁸)(R⁹), -N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and -N(R⁸)P(O)(OR⁹)(OR⁹);
- 20 p is 1, 2, 3, 4, 5, or 6;

R⁶ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;

R⁷ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;

5 R⁸ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;

R⁹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl; or R⁸ and R⁹ are joined together to form a heterocyclic ring;

10 R²⁰ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, halo, haloalkyl, trifluoromethyl, fluoroalkyl, perfluoroalkyl, thio, cyano, hydroxy, methoxy, alkoxy, phenoxy, aryloxy, heteroaryloxy, carboxyl, alkoxycarbonyl, acyl, nitro, amino, alkylamino, arylamino, heteroarylamino, amido, acylamino, sulfate, sulfonate, sulfonyl, sulfoxido, sulfonamido, 15 sulfamoyl, -[C(R⁴)₂]_p-R⁵, NR¹⁴R¹⁵, OR¹⁶, O-[C(R⁴)₂]_p-R⁵, NR¹⁴-[C(R⁴)₂]_p-R⁵ and SR¹⁶;

R²¹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, halo, haloalkyl, thio, cyano, carboxyl, alkoxycarbonyl, acyl, nitro, amino, amido, acylamino, sulfate, sulfonate, sulfonyl, sulfoxido, sulfonamido, sulfamoyl, NR¹⁴R¹⁵, OR¹⁶, and SR¹⁶;

20 R¹⁴ and R¹⁵ are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]_p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), and -P(O)(OR⁶)(OR⁷); or R¹⁴ and R¹⁵ are joined together to form an optionally substituted heterocyclic ring;


25 R¹⁶ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]_p-R⁵, -COR⁶, and -C(O)N(R⁶)(R⁷); and

r is 0 or 1;

wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, 30 heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl may be optionally substituted.

2. The compound of claim 1, wherein W and X are oxygen.
3. The compound of claim 1, wherein Z¹ and Z² are nitrogen; and Z³ is C-R²¹.
4. The compound of claim 1, wherein Z¹, Z² and Z³ are nitrogen.
5. The compound of claim 1, wherein Z¹ is nitrogen; and Z² and Z³ are each C-R²¹.
- 5 6. The compound of claim 1, wherein Z² is nitrogen; and Z¹ and Z³ are each C-R²¹.
7. The compound of claim 1, wherein R¹ is hydrogen, alkyl, or aralkyl.
8. The compound of claim 1, wherein W and X are oxygen, Z¹ and Z² are each nitrogen, Z³ is C-R²¹, and R¹ is hydrogen, alkyl, or aralkyl.
9. The compound of claim 8, wherein R¹ is hydrogen.
- 10 10. The compound of any one of claims 1 to 9, wherein R² and R³ are joined together to form an optionally substituted heterocyclic ring.
11. The compound of claim 10, wherein the optionally substituted heterocyclic ring is selected from the group consisting of piperazinyl, homopiperizinyl, pyrrolidinyl, piperidinyl, homopiperidinyl, morpholinyl, and quinolinyl.
- 15 12. The compound of any one of claims 1 to 9, wherein R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), and R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl may be optionally substituted.
- 20 13. The compound of claim 12, wherein R² is -[C(R⁴)₂]p-R⁵, and R³ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl may be optionally substituted.
- 25 14. The compound of claim 13, wherein R⁵ is aryl or heteroaryl, each of which may be optionally substituted.
15. The compound of claim 13, wherein R⁵ is -N(R⁸)(R⁹).
16. The compound of claim 13, wherein R⁴ is hydrogen.

17. A compound of formula 2:



or a pharmaceutically acceptable salt thereof, wherein independently for each occurrence:

R¹ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aralkyl,

5 heteroaralkyl, -COR⁶, and -C(O)OR⁶;

R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷);

R³ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,

10 heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), -SO₂N(R⁶)(R⁷), -P(O)(OR⁶)(OR⁷); or R² and R³ are joined together to form an optionally substituted heterocyclic ring;

R⁴ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocyclylalkyl, aralkyl, heteroaryl, heteroaralkyl, halo, hydroxy, alkoxy, hydroxyalkyl,

15 and alkoxyalkyl;

R⁵ is selected from the group consisting of aryl, heteroaryl, heterocyclyl, -N(R⁸)(R⁹),

-N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -

SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -

S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and -N(R⁸)P(O)(OR⁹)(OR⁹);

20 p is 1, 2, 3, 4, 5, or 6;

R⁶ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,

heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;

R⁷ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,

heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;


R^8 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl;

R^9 is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl; or R^8 and R^9 are joined together to form a heterocyclic ring; and

R^{20} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, halo, haloalkyl, trifluoromethyl, fluoroalkyl, perfluoroalkyl, thio, cyano, hydroxy, methoxy, alkoxy, phenoxy, aryloxy, heteroaryloxy, carboxyl, alkoxycarbonyl, acyl, nitro, amino, alkylamino, arylamino, heteroaryl amino, amido, acylamino, sulfate, sulfonate, sulfonyl, sulfoxido, sulfonamido, sulfamoyl, $-[C(R^4)_2]_p-R^5$, $NR^{14}R^{15}$, OR^{16} , $O-[C(R^4)_2]_p-R^5$, $NR^{14}-[C(R^4)_2]_p-R^5$ and SR^{16} ;

wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl may be optionally substituted.

- 15 18. The compound of claim 17, wherein R^1 is hydrogen, alkyl, or aralkyl.
19. The compound of claim 18, wherein R^1 is hydrogen, methyl, or benzyl.
20. The compound of any one of claims 17 to 19, wherein R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring.
21. The compound of claim 20, wherein R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring selected from the group consisting of :



wherein, independently for each occurrence:

R^{10} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, heterocyclylalkyl, $-[C(R^4)_2]_p-R^5$, $-COR^6$,


25 $-C(O)OR^6$, $-SO_2(R^6)$, $-C(O)N(R^6)(R^7)$, $-SO_2N(R^6)(R^7)$; and $-P(O)(OR^6)(OR^7)$;

R^{11} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, halo, haloalkyl, thio, cyano, alkylthio, nitro, $-N(R^8)(R^9)$, $-N(R^8)COR^9$, $-N(R^8)C(O)OR^9$, $-N(R^8)SO_2(R^9)$, $-CON(R^8)(R^9)$, $-OC(O)N(R^8)-(R^9)$, $-SO_2N(R^8)(R^9)$, $-OC(O)OR^8$, $-COOR^9$, $-C(O)N(OH)(R^8)$, $-OS(O)_2OR^8$, $-S(O)_2OR^8$, $-S(O)_2R^8$, $-OR^8$, $-COR^8$, 5 $-OP(O)(OR^8)(OR^8)$, $-P(O)(OR^8)(OR^8)$, $-N(R^8)P(O)(OR^9)(OR^9)$ and $-[C(R^4)_2]_p-R^5$; and $-[C(R^4)_2]_p-R^5$; and

n is 0, 1, 2, or 3;

wherein any one of the aforementioned alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, heteroaralkyl, and heterocyclylalkyl may be optionally substituted.


- 10 22. The compound of claim 21, wherein R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring of the formula:



23. The compound of claim 22, wherein n is 0.

- 15 24. The compound of claim 22, wherein R^{10} is hydrogen, alkyl, aryl, heteroaryl, $-COR^6$, $-C(O)OR^6$, or $-SO_2(R^6)$, or $-[C(R^4)_2]_p-R^5$.

25. The compound of any one of claims 17 to 19, wherein R^2 and R^3 are joined together to form an optionally substituted heterocyclic ring of the formula:





26. The compound of claim 25, wherein n is 0 or 1.


- 20 27. The compound of claim 25, wherein R^{11} is alkyl, aryl, heteroaryl, heterocyclyl, $-N(R^8)(R^9)$, $-N(R^8)COR^9$, $-N(R^8)C(O)OR^9$, $-N(R^8)SO_2(R^9)$ and $-[C(R^4)_2]_p-R^5$.


28. The compound of claim 26, wherein R^{11} is $-[C(R^4)_2]_p-R^5$, wherein p is 1 or 2, R^4 is hydrogen, and R^5 is selected from the group consisting of aryl, heteroaryl, heterocyclyl, $-N(R^8)(R^9)$, $-N(R^8)COR^9$, $-N(R^8)C(O)OR^9$, and $-N(R^8)SO_2(R^9)$.


29. The compound of any one of claims 17 to 19, wherein R² and R³ are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -[C(R⁴)₂]p-R⁵, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷), wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl may be optionally substituted.
- 5
30. The compound of claim 26, wherein R² is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -COR⁶, -C(O)OR⁶, -SO₂(R⁶), -C(O)N(R⁶)(R⁷), and -SO₂N(R⁶)(R⁷); and R³ is -[C(R⁴)₂]p-R⁵.
31. The compound of claim 30, wherein R² is optionally substituted alkyl.
- 10 32. The compound of claim 30, wherein R³ is aryl, heteroaryl, heterocyclyl, each of which may be optionally substituted.
33. The compound of claim 29, wherein at least one of R² and R³ is -[C(R⁴)₂]p-R⁵.
34. The compound of claim 33, wherein R⁴ is hydrogen.
35. The compound of claim 33, wherein R⁵ is selected from the group consisting of aryl, heteroaryl, -N(R⁸)(R⁹), -N(R⁸)COR⁹, -N(R⁸)C(O)OR⁹, -N(R⁸)SO₂(R⁹), -CON(R⁸)(R⁹), -OC(O)N(R⁸)-(R⁹), -SO₂N(R⁸)(R⁹), -OC(O)OR⁸, -COOR⁹, -C(O)N(OH)(R⁸), -OS(O)₂OR⁸, -S(O)₂OR⁸, -S(O)₂R⁸, -OR⁸, -COR⁸, -OP(O)(OR⁸)(OR⁸), -P(O)(OR⁸)(OR⁸) and -N(R⁸)P(O)(OR⁹)(OR⁹).
- 15
36. The compound of claim 35, wherein R⁵ is -N(R⁸)(R⁹).
- 20 37. The compound of claim 35, wherein R⁵ is aryl or heteroaryl.
38. A compound, or a pharmaceutically acceptable salt thereof, selected from the group consisting of:














and .

39. The compound of any one of claims 1-38, wherein the compound is an inhibitor of CK1 γ , CK1 γ 1, CK1 γ 2, or CK1 γ 3.

40. The compound of any one of claims 1-38, wherein the compound is an inhibitor of
5 the Wnt pathway.

41. The compound of any one of claims 1-38, wherein the compound is an inhibitor of the TGF β pathway.

42. The compound of any one of claims 1-38, wherein the compound has an IC₅₀ of less than 5000 nM for CK1 γ , CK1 γ 1, CK1 γ 2, or CK1 γ 3.

10 43. The compound of claim 42, wherein the compound has an IC₅₀ of less than 1000 nM for CK1 γ , CK1 γ 1, CK1 γ 2, or CK1 γ 3.

44. The compound of claim 42, wherein the compound has an IC₅₀ of less than 500 nM for CK1 γ , CK1 γ 1, CK1 γ 2, or CK1 γ 3.

15 45. A pharmaceutical composition, comprising at least one pharmaceutically acceptable excipient; and a compound of any one of claims 1-44.

46. A method of inhibiting CK1 γ activity, comprising contacting CK1 γ 1, CK1 γ 2, or CK1 γ 3 with a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

20 47. A method for treating or preventing a condition associated with aberrant CK1 γ activity, comprising the step of administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

25 48. A method of treating cancer, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

49. The method of claim 48, wherein the cancer is a cancer of the hematopoietic system, immune system, endocrine system, pulmonary system, gastrointestinal system, musculoskeletal system, reproductive system, central nervous system or urologic system.

50. The method of claim 48, wherein the cancer is located in the mammal's myeloid tissues, lymphoid tissues, pancreatic tissues, thyroid tissues, lung tissues, colon tissues, rectal tissues, anal tissues, liver tissues, blood, skin, bone, ovarian tissues, uterine tissues, cervical tissues, breast, prostate, testicular tissues, brain, brainstem, meningeal tissues, kidney or bladder.

51. The method of claim 48, wherein the cancer is breast cancer, colon cancer, multiple myeloma, prostate cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, leukemia, renal cell carcinoma, renal cancer, malignant melanoma, pancreatic cancer, lung cancer, colorectal carcinoma, brain cancer, head and neck cancer, hematologic (blood) cancer, bladder cancer, thyroid cancer, ovarian cancer, cervical cancer or myelodysplastic syndrome.

15 52. A method of treating Alzheimer's disease, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

53. A method of treating a Wnt dependent disease, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

54. A method of treating a TGF β dependent disease, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

25 55. A method of treating or preventing inflammation, inflammatory diseases (e.g., osteoarthritis and rheumatoid arthritis), neurological conditions (e.g., Alzheimer's disease) and neurodegeneration, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

56. A method of treating or preventing bone-related diseases and conditions, including 30 osteoporosis and bone formation, or facilitating bone restoration, comprising administering

to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.



5 57. A method of treating or preventing hypoglycemia, metabolic syndrome and diabetes, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

10 58. A method of influencing apoptosis (e.g., increasing the rate of apoptosis in cancerous cells), comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.



59. A method of treating or preventing aberrant embryonic development, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of claims 1-44 or a pharmaceutical composition of claim 45.

60. The compound of any one of claims 1-16, wherein r is 0.

15 61. The compound of any one of claims 17-37, wherein R²⁰ is hydrogen.

Figure 1**Figure 2**

2/5

Figure 3**Figure 4**

3/5






Figure 5**Figure 6**

Figure 7**5347 v CK1(y)****Figure 8****5343 v CK1(y)**

5/5

Figure 9**Figure 10**