



(11) **EP 2 601 962 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
12.06.2013 Bulletin 2013/24

(51) Int Cl.:
A61K 38/00 ^(2006.01) **A61K 31/337** ^(2006.01)
A61K 38/17 ^(2006.01) **A61P 35/00** ^(2006.01)

(21) Application number: **12196316.9**

(22) Date of filing: **03.10.2008**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT
RO SE SI SK TR**

(30) Priority: **05.10.2007 EP 07291214**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
08836333.8 / 2 205 257

(71) Applicant: **Immutep
91400 Orsay (FR)**

(72) Inventor: **Triebel, Frédéric
F-78000 Versailles (FR)**

(74) Representative: **Thornton, Neil
Reddie & Grose LLP
16 Theobalds Road
London
WC1X 8PL (GB)**

Remarks:

- This application was filed on 10-12-2012 as a divisional application to the application mentioned under INID code 62.
- Claims filed after the date of receipt of the divisional application (Rule 68(4) EPC).

(54) **Use of Recombinant LAG-3 or the Derivatives Thereof for Eliciting Monocyte Immune Response**

(57) The present invention relates to the use of a re-combinant LAG-3 or derivatives thereof in order to boost a monocyte-mediated immune response, in particular to elicit an increase in the number of monocytes in blood.

This finds use in the development of novel therapeutic agents for the treatment of an infectious disease or cancer.

EP 2 601 962 A1

Description

TECHNICAL FIELD

[0001] The present invention relates to the use of recombinant LAG-3 or derivatives thereof in order to boost a monocyte-mediated immune response.

[0002] It enables an increase in monocyte numbers in blood.

[0003] It finds many applications in particular in the development of novel therapeutic agents in cancer immunotherapy.

[0004] In the description which follows, the references between brackets [] refer to the attached reference list.

STATE OF THE ART

[0005] The lymphocyte activation gene 3 (*hlag-3*) expressed in human CD4⁺ and CD8⁺ activated T cells as well as in activated NK cells encodes a 503 amino acids type I membrane protein (LAG-3) with four extracellular immunoglobulin superfamily (IgSF) domains [1]. A murine lymphocyte activation gene 3 (*mlag-3*) was cloned and approximatively 70% of homology was found with *hlag-3*, with the same proline rich motif in the intracytoplasmic tail.

[0006] LAG-3 (CD223), described as being a natural high affinity ligand for MHC class II, is known to induce maturation of monocyte-derived dendritic cells *in vitro*, and is used as an immunotherapy adjuvant to induce CD4 T helper type 1 responses and CD8 T cell responses *in vivo* [2]. Further information regarding LAG-3 and its use as an immunostimulant may be found in TRIEBEL et al. [1], TRIEBEL et al. [3], and HUARD et al. [4].

[0007] Some forms of soluble LAG-3 can bind to MHC class II molecules and can induce dendritic cells to mature and migrate to secondary lymphoid organs where they can prime naïve CD4-helper and CD8-cytotoxic T cells leading to tumour rejection [5].

[0008] More recently a recombinant soluble human LAG-3Ig fusion protein (IMP321) was shown to activate a large range of effector cells in both innate and acquired immune responses, for example inducing monocytes-macrophages to secrete cytokines/chemokines [6].

[0009] Monocytes are produced by the bone marrow from haematopoietic stem cell precursors called monoblasts. They constitute between three to eight percent of the leukocytes in the blood. Monocytes circulate in the bloodstream for about 24 hours (half-life of 8 hours) and then typically move into tissues throughout the body. In the tissues, monocytes mature into macrophages, epithelioid cells or antigen-presenting cells (APCs, for example dendritic cells). Monocytes are responsible for phagocytosis (ingestion) of foreign substances in the body. Monocytes can perform phagocytosis using intermediary (opsonising) proteins such as antibodies or complement that coat the pathogen, as well as by binding to

the pathogen directly via pattern-recognition receptors that recognize pathogens. Monocytes are also capable of killing infected host cells via antibody, termed antibody-dependent cell-mediated cytotoxicity (ADCC). Due to their secretion and phagocytosis properties, monocytes-macrophages occur in aspecific and specific immune response.

[0010] The study of membrane markers allows the identification of monocyte populations, mature or not, dystrophic or not. The molecules present on monocyte membranes, mature or not, are almost always non specific but correspond to the following activities :

- receptor for the Fc fragment of IgG (CD16, CD32, CD64),
- receptor for the Fc fragment of IgE (CD23),
- receptor for complement fractions (CD11b, CD21/CD35),
- leukocyte adhesion proteins (CD11 a, CD11c),
- protein facilitating binding to LPS of Gram - bacteria (CD14),
- protein with tyrosine phosphatase activity (CD45).

DISCLOSURE OF THE INVENTION

[0011] The authors of the present invention have now discovered, entirely unexpectedly, that human LAG-3 or derivatives thereof when inoculated into patients with highly malignant tumors, for example patients with metastatic breast cancer (MBC) or metastatic renal clear-cell carcinoma (MRCC), induced a potent immunity which is monocyte dependent.

[0012] Said induced immunity manifests itself by a significant increase in blood monocyte counts.

[0013] This result was achieved by means of plural administration of LAG-3 or derivatives thereof to patients receiving immunotherapy or chemo-immunotherapy.

[0014] This result is rather surprising since binding to, and activation of, monocytes is not expected to be followed by monocyte expansion. Indeed monocytes are end-of-differentiation hematopoietic cells and can not proliferate. They can stay in the blood as monocytes or differentiate toward either macrophages or dendritic cells under the influence of different cytokines, until they die. Thus it is believed, without limitation to the following theory, that the mechanism of action involved may be a proliferative signal directed to hematopoietic precursor cells (before the promonocyte stage) residing in the bone marrow, or an increase in the half-life or residence time of mature circulating monocytes.

[0015] Therefore the present invention relates to the use of a recombinant LAG-3 protein or derivative thereof that elicits monocyte mediated immune response, for the manufacture of a medicament inducing an increase in monocyte numbers for the treatment of an infectious disease or cancer.

[0016] By "derivatives of LAG-3", in the sense of the present invention, is meant mutants, variants and frag-

ments of LAG-3 provided that they maintain the ability of LAG-3 to bind MHC class II molecules.

[0017] Thus, the following forms of LAG-3 may be used :

- the whole LAG-3 protein,
- a soluble polypeptide fragment thereof consisting of at least one of the four immunoglobulin extracellular domains, namely the soluble part of LAG-3 comprised of the extracellular region stretching from the amino acid 23 to the amino acid 448 of the LAG-3 sequence disclosed in French patent Application FR 90 00 126,
- a fragment of LAG-3 consisting of substantially all of the first and second domains,
- a fragment of LAG-3 consisting of substantially all of the first and second domains or all of the four domains, such as defined in WO 95/30750,
- a mutant form of soluble LAG-3 or a fragment thereof comprising the D1 and D2 extracellular domains and consisting of :

* a substitution of an amino acid at one of the following positions :

position 73 where ARG is substituted with GLU,
position 75 where ARG is substituted with ALA or GLU,
position 76 where ARG is substituted with GLU,
or a combination of two or more of those substitutions,

* a substitution of an amino acid at one of the following positions :

position 30 where ASP is substituted with ALA,
position 56 where HIS is substituted with ALA,
position 77 where TYR is substituted with PHE,
position 88 where ARG is substituted with ALA,
position 103 where ARG is substituted with ALA,
position 109 where ASP is substituted with GLU,
position 115 where ARG is substituted with ALA,
or a deletion of the region comprised between the position 54 and the position 66,
or a combination of two or more of those substitutions.

[0018] Those mutants are described by HUARD et al.

(Proc. Natl. Acad. Sci. USA, 11 : 5744-5749, 1997).

- a physiological variant of LAG-3 comprised of the soluble 52kDa protein containing D1, D2 and D3.
- a recombinant soluble human LAG-3Ig fusion protein (IMP321), a 200-kDa dimer produced in Chinese hamster ovary cells transfected with a plasmid encoding for the extracellular domain of hLAG-3 fused to the human IgG1 Fc.

[0019] By « medicament », in the sense of the present invention, is meant an effective plurality of doses of a recombinant LAG-3 protein or derivative thereof.

[0020] By « effective plurality of doses of a recombinant LAG-3 protein or derivative thereof », in the sense of the present invention, is meant a formulation that allows administration of one dose of a recombinant LAG-3 protein or derivative thereof every one to several weeks for at least 12 weeks, and preferably for at least 24 weeks, separated by 13-day ± 2 days administration-free intervals. Advantageously, the administration is an every-two-week schedule.

[0021] By « one dose of a recombinant LAG-3 protein or derivative thereof », in the sense of the present invention, is meant a formulation that allows one administration in the range of 0.25-30 mg, preferably 1-6.25 mg, more preferably 6-30 mg, and for example about 1.25 mg of recombinant LAG-3 protein or derivative thereof to a patient in need thereof having a body mass index (weight/height²) in the range of 18-30 kg/m².

[0022] The recombinant LAG-3 or derivatives thereof are administered in a free form, for example in a soluble form by inoculating them systemically, for example as a subcutaneous, intramuscular or intravenous injection, preferably as a subcutaneous injection.

[0023] Said recombinant LAG-3 or derivatives thereof may also be formulated so as to allow administration with a compound having anti-cancer or anti-infectious disease immunotherapeutic or chemotherapeutic properties.

[0024] By "administration with a compound", in the sense of the present invention, is meant an administration of a recombinant LAG-3 or derivative thereof before, with, or subsequent to, the administration of said compound.

[0025] By "compound having anti-cancer or anti-infectious disease chemotherapeutic properties", in the sense of the present invention, is meant for example a chemotherapy agent selected from the group consisting of taxanes (paclitaxel, docetaxel), gemcitabine and anthracyclines (doxorubicine) or an anti-viral agent such as ribavirin.

[0026] In a particular embodiment of the invention, recombinant LAG-3 protein or derivative thereof is administered to patients after the first administration of the cytotoxic compound having anti-cancer or anti-infectious disease chemotherapeutic properties.

[0027] Advantageously, recombinant LAG-3 protein or

derivative thereof is administered to patients is administered 12 to 96 hours after the administration of the cytotoxic compound having anti-cancer or anti-infectious disease chemotherapeutical properties.

[0028] In another embodiment, recombinant LAG-3 protein or derivative thereof is administered to patients is administered one or two days after the first administration of the compound having anti-cancer or anti-infectious disease chemotherapeutical properties.

[0029] In another particular embodiment of the invention, recombinant LAG-3 protein or derivative and the cytotoxic compound having anti-cancer or anti-infectious disease chemotherapeutical properties are administered simultaneously, separately or sequentially.

[0030] Advantageously, in this particular embodiment of the invention, recombinant LAG-3 protein or derivative thereof is administered at least six times, for example seven times, ten times, twelve times or more.

[0031] Advantageously, in this particular embodiment of the invention, recombinant LAG-3 protein or derivative thereof is administered on an every-two-week schedule.

[0032] Advantageously, recombinant LAG-3 protein or derivative thereof is administered at a dose comprised between 0.25 to 30 mg, eventually at a dose comprised between 6 to 30 mg, eventually at a dose comprised between 8 to 25 mg, eventually between 12 and 20 mg.

[0033] By "compound having anti-cancer or anti-infectious disease immunotherapeutical properties", in the sense of the present invention, is also meant for example a compound selected from the group consisting of therapeutic antibodies killing tumour cells through ADCC (antibody-dependent cell cytotoxicity), and mixtures thereof, and preferably from the group consisting of rituximab, cetuximab, edrecolomab, and trastuzumab.

[0034] In a particular embodiment of the invention, recombinant LAG-3 protein or derivative thereof and therapeutic antibodies are administered to patients simultaneously, separately or sequentially.

[0035] Advantageously, in a particular embodiment of the invention, recombinant LAG-3 protein or derivative thereof is administered to patients the same day as therapeutic antibodies.

[0036] The present invention also relates to kit-of-parts, i.e. a combined preparation, containing recombinant LAG-3 protein or derivative thereof and a therapeutic antibody for simultaneous, separate or sequential use.

[0037] Advantageously, the kit-of-parts contains recombinant LAG-3 protein or derivative thereof and a therapeutic antibody selected from the group consisting of rituximab, cetuximab, edrecolomab, and trastuzumab.

[0038] Preferentially, the kit-of-part of the invention contain recombinant LAG-3 protein or derivative thereof and rituximab.

[0039] In the kit-of-parts of the invention, recombinant LAG-3 protein or derivative thereof and a therapeutic antibody form a functional unity, because of a synergistic cytotoxic effect between the two components. This effect

is a new joint effect, because the two components administered alone does not have the same effect as when administered as a combined preparation.

[0040] The present invention also relates to kit-of-parts, i.e. a combined preparation, containing recombinant LAG-3 protein or derivative thereof and a compound having anti-cancer or anti-infectious disease chemotherapeutical properties for simultaneous, separate or sequential use.

[0041] Advantageously, the kit-of-parts contains recombinant LAG-3 protein or derivative thereof and a compound having anti-cancer or anti-infectious disease chemotherapeutical properties selected from the group consisting of taxanes (paclitaxel, docetaxel), gemcitabine and anthracyclines (doxorubicine).

[0042] The present invention also relates to a method for treating a pathological condition involving a monocyte mediated immune response, comprising administering the medicament as above defined to a patient in need thereof.

[0043] By "pathological condition involving a monocyte mediated immune response", in the sense of the present invention, is meant viral infectious diseases, parasitic infectious diseases, bacterial infectious diseases, and cancer.

[0044] Various aspects of the invention are defined in the following numbered paragraphs.

1. Use of a recombinant LAG-3 protein or derivative thereof that elicits a monocyte-mediated immune response, for the manufacture of a medicament inducing an increase in the number of monocytes for the treatment of an infectious disease or cancer.

2. Use according to paragraph 1, wherein the medicament consists of an effective plurality of doses of a recombinant LAG-3 protein or derivative thereof.

3. Use according to paragraph 2, wherein said plurality of doses of a recombinant LAG-3 protein or derivative thereof is formulated so as to allow administration of one dose every one to several weeks for at least 12 weeks, and preferably for at least 24 weeks, separated by 13-day ± 2 days administration-free intervals.

4. Use according to either one of paragraphs 2 or 3, wherein said one dose of a recombinant LAG-3 protein or derivative thereof is formulated so as to allow subcutaneous or intravenous administration in the range of 0.25-30 mg, and preferably 6-30 mg, of a recombinant LAG-3 protein or derivative thereof to a patient in need thereof having a body mass index (weight/height²) in the range of 18-30 kg/m².

5. Use according to any one of paragraphs 2 to 4, wherein said doses of a recombinant LAG-3 protein or derivative thereof are formulated so as to allow administration with a compound having anti-cancer or anti-infectious disease immunotherapeutical or chemotherapeutical properties.

6. Use according to paragraph 5, wherein one dose

of a recombinant LAG-3 protein or derivative thereof comprises 0.25-30 mg, and preferably 6-30 mg, of said recombinant LAG-3 protein or derivative thereof.

7. Use according to either one of paragraphs 5 or 6, wherein said compound is selected from the group consisting of chemotherapy agents.

8. Use according to paragraph 7, wherein said chemotherapy agents are selected from the group consisting of taxanes, for example paclitaxel and docetaxel, and anthracyclines, for example doxorubicine, and gemcitabine.

9. Use according to either one of paragraph 5 or 6, wherein said compound is selected from the group consisting of therapeutic antibodies killing tumour cells through ADCC (antibody-dependent cell-mediated cytotoxicity), and mixtures thereof.

10. Use according to paragraph 9, wherein said therapeutic antibodies are from the group consisting of rituximab, cetuximab, edrecolomab and trastuzumab.

11. Kit-of-parts containing recombinant LAG-3 protein or derivative thereof and a therapeutic antibody as mentioned in paragraphs 9 or 10, for simultaneous, separate or sequential use.

12. Kit-of-parts containing recombinant LAG-3 protein or derivative thereof and a compound having anti-cancer chemotherapeutical properties as mentioned in paragraphs 7 or 8, for simultaneous, separate or sequential use.

[0045] Other advantages may also appear to one skilled in the art from the non-limitative examples given below, and illustrated by the enclosed figures.

BRIEF DESCRIPTION OF THE FIGURES

[0046]

- Figure 1 represents fluorescence-activated cell sorting (FACS) analysis of monocytes (i.e. CD14⁺CD45⁺ cells) in PBMCs from metastatic breast carcinoma patients.
- Figure 2 represents fluorescence-activated cell sorting (FACS) analysis of monocytes (i.e. CD14⁺CD45⁺ cells) in fresh whole blood from metastatic breast carcinoma patients.
- Figure 3 represents fluorescence-activated cell sorting (FACS) analysis of monocytes (i.e. CD14⁺CD45⁺ cells) in fresh whole blood from metastatic renal cell cancer patients.
- Figure 4 represents fluorescence-activated cell sorting (FACS) analysis of monocytes (i.e. CD14⁺CD45⁺ cells) in fresh whole blood from metastatic breast carcinoma patients.
- Figure 5 represents the pharmacokinetic profiles of IMP321 measured by ELISA in the plasma of metastatic renal cell cancer patients.

- Figure 6 represents the flow cytometry analysis of PBMC cultured in different conditions with rituximab and/or IMP321.

EXAMPLES

Example 1 : Monocytes increase in metastatic breast cancer (MBC) patients using low IMP321 dose

[0047] Five MBC patients, receiving chemotherapy known to induce tumour cell apoptosis, each received one subcutaneous IMP321 dose of 0.25 mg 1-2 days after chemotherapy every other week, for 24 weeks, separated by 14-day administration-free intervals.

[0048] Blood samples were collected in heparinated lithium tubes (Vacutainer; BD Biosciences) from each patient, 14 days after the last IMP321 injection (i.e. looking at lasting immunomodulatory effects of the product), at 3 months (Day 85) and 6 months (Day 170). PBMCs were isolated on Ficoll-Paque gradient (Pharmacia) using LeucoSep tubes (Greiner Bio-One), and used immediately.

[0049] The increase in number of monocytes was analysed by fluorescence-activated cell sorting (FACS) in said fresh PBMC samples (because monocytes are sensitive to freezing), and compared with the monocyte counts carried out on fresh PBMC samples collected before IMP321 administration (Day 1).

[0050] The results are represented in Figure 1.

[0051] The results showed a 2.5-fold (at 3 months, Day 85) and a 3.5-fold (at 6 months, Day 170) mean increase of monocyte counts at this low IMP321 dose clinical protocol.

[0052] In order to confirm the above results, a more direct and probably more accurate approach was carried out, which was to quantify directly *ex-vivo* the number of monocytes in whole blood (i.e. without prior purification of PBMCs on Ficoll gradient) by first measuring the exact volume of blood to be analyzed with diluted fluorescent beads and then counting the number of CD14⁺ cells (i.e. monocytes) in the gated CD45⁺ (leukocytes) cells present in this whole blood volume.

[0053] The results are represented in Figure 2.

[0054] The results showed a 4.4-fold mean increase at Day 170 (2.8-fold at Day 85) when IMP321 was given at low dose (0.25 mg) for a long period of time, 6 months, with 12 injections, showing strong and direct stimulation of the targeted MHC class II⁺ monocyte-like hematopoietic cells.

Example 2: Monocytes increase in metastatic renal clear-cell carcinoma (MRCC) patients using high IMP321 dose

[0055] Three MRCC patients each received one subcutaneous IMP321 dose of 6.25 mg every other week, for 12 weeks, separated by 14-day administration-free intervals.

[0056] Blood samples were collected as described above from each patient, 14 days after the last IMP321 injection (i.e. looking at lasting immunomodulatory effects of the product), at 2 months (Day 57) and 3 months (Day 85), and used immediately.

[0057] The expansion of CD14⁺ CD45⁺ cells was analysed by FACS in fresh blood samples (because monocytes are sensitive to freezing), and compared with the monocyte counts carried out on fresh blood samples collected before IMP321 administration (Day 1).

[0058] The results are represented in Figure 3.

[0059] The results showed a 2-fold (at 3 months, Day 85) mean increase of monocyte counts with this high IMP321 dose clinical protocol where patients received only 6 injections.

Example 3 : Monocytes increase in metastatic breast carcinoma patients receiving paclitaxel and IMP321 doses

[0060] Patients receiving as a first line chemotherapy for metastatic breast carcinoma 6 cycles of paclitaxel (80 mg/m² given i.v.) on days 1, 8, and 15 of a 28 day cycle, received 1-30 mg s.c. (sub-cutaneous) IMP321 on days 2 and 16 of each 28-day cycle. Alternatively, IMP321 was administered at days 3 or 17.

[0061] Accordingly, each patient received a standard 6-month course of weekly paclitaxel with 12 s.c. injections of IMP321, each injection being given one to two days after paclitaxel administration on an every-two-week schedule.

[0062] The increase in absolute monocyte counts per microliter of fresh blood was analysed by fluorescence-activated cell sorting (FACS), 14 days after the last injection, at 3 months (Day 85) and 6 months (Day 170) compared to day 1.

[0063] The results obtained in patients injected with a low dose IMP321 (1.25 mg) are represented in Figure 4.

[0064] These data showed that doses of 1.25 mg in most if not all patients (Figure 4) induce an expansion of the monocyte subset pool in the blood.

[0065] It is predicted that the optimal dose regimen for IMP321 will be between 6 and 30 mg/injection.

[0066] These doses have been shown to be safe and give an acceptable systemic exposure based on the results of pharmacokinetics data obtained in metastatic renal cell cancer patients (Figure 5). A blood concentration of IMP321 superior to 1 ng/ml for at least 24 hours after s.c. injection could be obtained in patients injected by IMP321 doses of more than 6 mg (Figure 5).

Example 4: Treatment of advanced pancreas cancer patients receiving gemcitabine and IMP321 doses

[0067] Patients, receiving as a first line chemotherapy for advanced pancreas cancer (or patients not eligible for surgical removal of the tumor) 6 cycles of standard gemcitabine (1 gm/m² given i.v. over 30 min) on days 1,

8, and 15 of a 28 day cycle, receive in addition 6 to 30 mg s.c. IMP321 on days 2 and 16 of each 28-day cycle. Alternatively, IMP321 is administered at days 3 or 17.

[0068] Accordingly, each patient receives a standard 6-month course of gemcitabine with 12 s.c. injections of IMP321, each injection being given one to two days after gemcitabine administration on an every-two-week schedule.

[0069] The number of monocytes is analysed by fluorescence-activated cell sorting (FACS) as in example 1.

Example 5: Induction of increased rituximab-mediated ADCC by low doses IMP321

[0070] PBMCs are first incubated for 40 hours with IL-2 (100 U/ml), with or without IMP321 (at the concentrations 0 µg/ml, 0.03 µg/ml or 0.1 µg/ml). PBMCs are then incubated with increasing concentrations of rituximab (0, 0.5 and 5 µg/ml) in the presence of target cells (i.e. human CD20⁺ Raji B cells).

[0071] Raji cells were first labeled with CFSE (carboxy-fluorescein succinimidyl ester), incubated in medium with rituximab at 0, 0.5 or 5 µg/ml and cocultured with effector cells at an effector-target ratio of 25:1 for 6 hours at 37°C.

[0072] The cells were then incubated with 7-AAD (7-amino-actinomycin-D) for 15 min on ice and analyzed by flow cytometry to determine the percentage of dead CFSE⁺ 7-AAD⁺ Raji target cells (i.e. % of cytotoxicity).

[0073] The results are presented in Figure 6.

[0074] Increasing the concentration of rituximab increased the percentage of cytotoxicity, clearly showing a dose-dependent ADCC activity.

[0075] When 0.03 or 0.1 µg/ml IMP321 is added, the percentage of cytotoxicity greatly increased. For instance, a 30 % cytotoxicity is observed with 0.5 µg/ml rituximab in the presence of 0.1 µg/ml IMP321 which is superior to the 25 % cytotoxicity value obtained with 5 µg/ml rituximab in the absence of IMP321.

[0076] Thus, adding 0.1 µg/ml IMP321 potentializes 10-15 fold the activity of rituximab because a superior cytotoxicity is obtained with 10 time less antibody when a low dose IMP321 (0.1 µg/ml) is added.

[0077] These data show the synergistic effect between rituximab and IMP321.

Reference List

[0078]

[1] TRIEBEL et al., J. Exp. Med., 171 : 1393-1405, 1990

[2] BRIGNONE et al., J. Immune Based Ther Immunotherapies, 5 : 5, 2007

[3] TRIEBEL et al., Trends Immunol., 24 : 619-622, 2003

[4] HUARD et al., Proc. Natl. Acad. Sci. USA, 94 : 5744-5749, 1997

[5] PRIGENT et al., Eur. J. Immunol., 29 : 3867-3876, 1999

[6] BRIGNONE et al., J. Immunol., 179: 4202-4211, 2007

tein, or derivative thereof that maintains the ability of LAG-3 to bind MHC class II molecules, that elicits a monocyte-mediated immune response.

- 5 8. A multiple-dose formulation according to claim 7, wherein the recombinant LAG-3 protein or derivative thereof consists of the recombinant soluble human LAG-3Ig fusion protein IMP321.

10

Claims

1. Recombinant LAG-3 protein, or derivative thereof that maintains the ability of LAG-3 to bind MHC class II molecules, that elicits a monocyte-mediated immune response, for use as an anti-cancer medicament inducing an increase in the number of monocytes, wherein said medicament consists of an effective plurality of doses of the recombinant LAG-3 protein or derivative thereof. 15 20
2. Recombinant LAG-3 protein or derivative thereof for use according to claim 1, wherein one dose of a recombinant LAG-3 protein or derivative thereof comprises a dose of said recombinant LAG-3 protein or derivative thereof which is a molar equivalent of 0.25-30 mg, preferably 6-30 mg, of the recombinant soluble human LAG-3Ig fusion protein IMP321. 25
3. Recombinant LAG-3 protein or derivative thereof for use according to claim 1 or 2, wherein the recombinant LAG-3 protein or derivative thereof consists of the recombinant soluble human LAG-3Ig fusion protein IMP321. 30 35
4. Recombinant LAG-3 protein or derivative thereof for use according to any one of claims 1 to 3, wherein said plurality of doses of a recombinant LAG-3 protein or derivative thereof are to be administered as follows: one dose every one to several weeks for at least 12 weeks, and preferably for at least 24 weeks, separated by 13-day \pm 2 days administration-free intervals. 40
5. Recombinant LAG-3 protein or derivative thereof for use according to any one of claims 1 to 4, wherein the recombinant LAG-3 protein or derivative thereof is to be administered systemically, preferably as a subcutaneous, intramuscular, or intravenous injection. 45 50
6. Recombinant LAG-3 protein or derivative thereof for use according to any one of claims 1 to 5, wherein the medicament is to be administered to a human patient in need thereof. 55
7. A multiple-dose formulation, which comprises an effective plurality of doses of a recombinant LAG-3 pro-




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6



EUROPEAN SEARCH REPORT

Application Number
EP 12 19 6316

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|--|--|---|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (IPC) |
| X | US 2002/192195 A1 (TRIEBEL FREDERIC [FR]) 19 December 2002 (2002-12-19) * page 1, paragraphs 1,10 * * page 2, paragraph 16 * * claims 7,8 * * page 2, paragraph 9 * * page 4, paragraph 8 * | 1-8 | INV. A61K38/00 A61K31/337 A61K38/17 ADD. A61P35/00 |
| A | "Principles of cancer therapy" In: Anonymous: "The Merck Manual of Diagnosis and Therapy, 18th Edition", 2006, Merck Research Laboratories, Whitehouse Station, NJ, XP002695786, ISBN: 0911910182 pages 1157-1171, * the whole document * | 1-8 | |
| A | WO 98/23748 A (APPLIED RESEARCH SYSTEMS [NL]; BIFFONI MAURO [IT]; PAPOIAN RUBEN [IT]) 4 June 1998 (1998-06-04) * page 9, line 1 - line 3 * | 1-8 | |
| A | WO 98/23741 A (ROUSSY INST GUSTAVE [FR]; INST NAT SANTE RECH MED [FR]; APPLIED RESEAR) 4 June 1998 (1998-06-04) * page 4, line 8 - line 24 * * page 11, line 28 - line 30; claims 1,11 * * page 12, line 12 - line 19 * | 1-8 | |
| A,P | BRIGNONE,C. ET AL.: "IMP321 (sLAG-3) safety and T cell response potentiation using an influenza vaccine as a model antigen: A single-blind phase study", VACCINE, vol. 25, 2007, pages 4641-4650, XP022095366, * page 4649, paragraph 3 * | 1-8 | |
| The present search report has been drawn up for all claims | | | |
| Place of search Munich | | Date of completion of the search 2 May 2013 | Examiner Rodrigo-Simón, Ana |
| CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document | | T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document | |

EPO FORM 1503 03/82 (P04C01)



EUROPEAN SEARCH REPORT

Application Number
EP 12 19 6316

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|---|---|---|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (IPC) |
| A | PRIGENT P ET AL: "Lymphocyte activation gene-3 induces tumor regression and antitumor immune responses", EUROPEAN JOURNAL OF IMMUNOLOGY, WILEY - V C H VERLAG GMBH & CO. KGAA, DE, vol. 29, no. 12, 1 December 1999 (1999-12-01), pages 3867-3876, XP002291775, ISSN: 0014-2980, DOI: 10.1002/(SICI)1521-4141(199912)29:12<3867:AID-IMMU3867>3.3.CO;2-5 * abstract * | 1-8 | |
| E L | WO 2009/044273 A2 (IMMUTEP [FR]; TRIEBEL FREDERIC [FR]) 9 April 2009 (2009-04-09) * the whole document * | 1-8 | |
| | | | TECHNICAL FIELDS SEARCHED (IPC) |
| | | | |
| The present search report has been drawn up for all claims | | | |
| Place of search Munich | | Date of completion of the search 2 May 2013 | Examiner Rodrigo-Simón, Ana |
| CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document | | T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document | |

1
EPO FORM 1503 03.02 (P04031)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 12 19 6316

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

02-05-2013

| Patent document cited in search report | | Publication date | Patent family member(s) | Publication date |
|---|----|---------------------|----------------------------|---------------------|
| US 2002192195 | A1 | 19-12-2002 | AT 324455 T | 15-05-2006 |
| | | | AT 478956 T | 15-09-2010 |
| | | | AU 753738 B2 | 24-10-2002 |
| | | | AU 9254798 A | 16-02-1999 |
| | | | BR 9811037 A | 01-08-2000 |
| | | | CA 2293805 A1 | 04-02-1999 |
| | | | CA 2732570 A1 | 04-02-1999 |
| | | | CN 1265149 A | 30-08-2000 |
| | | | DE 69834330 T2 | 29-03-2007 |
| | | | DK 1002108 T3 | 21-08-2006 |
| | | | EE 200000039 A | 16-10-2000 |
| | | | EP 0893507 A1 | 27-01-1999 |
| | | | EP 1002108 A2 | 24-05-2000 |
| | | | EP 1698701 A1 | 06-09-2006 |
| | | | ES 2262242 T3 | 16-11-2006 |
| | | | HK 1030014 A1 | 10-06-2005 |
| | | | IL 134212 A | 29-03-2012 |
| | | | JP 4723722 B2 | 13-07-2011 |
| | | | JP 2001510806 A | 07-08-2001 |
| | | | NO 20000378 A | 25-01-2000 |
| | | | PT 1002108 E | 29-09-2006 |
| | | | UA 75322 C2 | 16-10-2000 |
| | | | US 6410509 B1 | 25-06-2002 |
| | | | US 2002192195 A1 | 19-12-2002 |
| | | | WO 9904810 A2 | 04-02-1999 |
| ----- | | | | |
| WO 9823748 | A | 04-06-1998 | AR 018489 A1 | 28-11-2001 |
| | | | AT 271607 T | 15-08-2004 |
| | | | AU 733825 B2 | 24-05-2001 |
| | | | AU 1066797 A | 22-06-1998 |
| | | | BR 9612821 A | 11-04-2000 |
| | | | CA 2272130 A1 | 04-06-1998 |
| | | | DE 69632967 D1 | 26-08-2004 |
| | | | DE 69632967 T2 | 11-08-2005 |
| | | | EP 0941329 A1 | 15-09-1999 |
| | | | ES 2225901 T3 | 16-03-2005 |
| | | | HK 1023599 A1 | 05-08-2005 |
| | | | IL 130168 A | 05-09-2006 |
| | | | JP 3920928 B2 | 30-05-2007 |
| | | | JP 2001505770 A | 08-05-2001 |
| | | | KR 20000053291 A | 25-08-2000 |
| | | | NO 992569 A | 27-05-1999 |
| | | | PT 941329 E | 30-11-2004 |
| | | | UA 73270 C2 | 15-12-2004 |
| | | | US 2002081282 A1 | 27-06-2002 |
| | | | WO 9823748 A1 | 04-06-1998 |

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 12 19 6316

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

02-05-2013

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| | | ZA 9710737 A | 30-07-1998 |
| WO 9823741 A | 04-06-1998 | AT 320488 T | 15-04-2006 |
| | | AU 728911 B2 | 18-01-2001 |
| | | AU 5229498 A | 22-06-1998 |
| | | CA 2273080 A1 | 04-06-1998 |
| | | DE 69735502 T2 | 14-12-2006 |
| | | EP 0942973 A1 | 22-09-1999 |
| | | IL 130123 A | 24-07-2007 |
| | | JP 2000516101 A | 05-12-2000 |
| | | US 6482925 B1 | 19-11-2002 |
| | | WO 9823741 A1 | 04-06-1998 |
| WO 2009044273 A2 | 09-04-2009 | AU 2008306576 A1 | 09-04-2009 |
| | | CN 101873864 A | 27-10-2010 |
| | | EP 2044949 A1 | 08-04-2009 |
| | | EP 2205257 A2 | 14-07-2010 |
| | | JP 2010540616 A | 24-12-2010 |
| | | US 2011008331 A1 | 13-01-2011 |
| | | WO 2009044273 A2 | 09-04-2009 |

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- FR 9000126 [0017]
- WO 9530750 A [0017]

Non-patent literature cited in the description

- HUARD et al. *Proc. Natl. Acad. Sci. USA*, 1997, vol. 11, 5744-5749 [0018]
- TRIEBEL et al. *J. Exp. Med.*, 1990, vol. 171, 1393-1405 [0078]
- BRIGNONE et al. *J. Immune Based Ther Immunotherapies*, 2007, vol. 5, 5 [0078]
- TRIEBEL et al. *Trends Immunol.*, 2003, vol. 24, 619-622 [0078]
- HUARD et al. *Proc. Natl. Acad. Sci. USA*, 1997, vol. 94, 5744-5749 [0078]
- PRIGENT et al. *Eur. J. Immunol.*, 1999, vol. 29, 3867-3876 [0078]
- BRIGNONE et al. *J. Immunol.*, 2007, vol. 179, 4202-4211 [0078]