



US008829013B1

(12) **United States Patent**  
**Rodgers et al.**(10) **Patent No.:** **US 8,829,013 B1**(45) **Date of Patent:** **\*Sep. 9, 2014**(54) **SALTS OF THE JANUS KINASE INHIBITOR  
(R)-3-(4-(7H-PYRROLO[2,3-D]PYRIMIDIN-  
4-YL)-1H-PYRAZOL-1-YL)-3-  
CYCLOPENTYLPROPANENITRILE**(71) Applicant: **Incyte Corporation**, Wilmington, DE  
(US)(72) Inventors: **James D. Rodgers**, Landenberg, PA  
(US); **Hui-Yin Li**, Hockessin, DE (US)(73) Assignee: **Incyte Corporation**, Wilmington, DE  
(US)(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.This patent is subject to a terminal dis-  
claimer.(21) Appl. No.: **14/256,311**(22) Filed: **Apr. 18, 2014****Related U.S. Application Data**(63) Continuation of application No. 14/097,588, filed on  
Dec. 5, 2013, now Pat. No. 8,722,693, which is a  
continuation of application No. 12/137,892, filed on  
Jun. 12, 2008, now abandoned.(60) Provisional application No. 60/943,705, filed on Jun.  
13, 2007.(51) **Int. Cl.****A61K 31/519** (2006.01)**A61P 35/00** (2006.01)**C07D 487/04** (2006.01)(52) **U.S. Cl.**CPC ..... **C07D 487/04** (2013.01); **C07B 2200/13**  
(2013.01)USPC ..... **514/265.1**; 544/280(58) **Field of Classification Search**CPC ..... **C07D 487/04**; **A61K 31/519**

USPC ..... 544/280; 514/265.1

See application file for complete search history.

(56) **References Cited****U.S. PATENT DOCUMENTS**

2,985,589 A 5/1961 Broughton et al.  
 4,402,832 A 9/1983 Gerhold  
 4,498,991 A 2/1985 Oroskar  
 4,512,984 A 4/1985 Seufert et al.  
 4,548,990 A 10/1985 Mueller et al.  
 5,510,101 A 4/1996 Stroppolo  
 5,521,184 A 5/1996 Zimmermann  
 5,630,943 A 5/1997 Grill  
 5,856,326 A 1/1999 Anthony  
 5,919,779 A 7/1999 Proudfoot  
 6,060,038 A 5/2000 Burns  
 6,136,198 A 10/2000 Adam et al.  
 6,217,895 B1 4/2001 Guo  
 6,335,342 B1 1/2002 Longo et al.  
 6,375,839 B1 4/2002 Adam et al.  
 6,413,419 B1 7/2002 Adam et al.  
 6,486,322 B1 11/2002 Longo et al.  
 6,548,078 B2 4/2003 Guo  
 6,569,443 B1 5/2003 Dawson

6,579,882 B2 6/2003 Bhatia et al.  
 6,635,762 B1 10/2003 Blumenkopf et al.  
 6,712,973 B2 3/2004 Adam et al.  
 6,852,727 B2 2/2005 Goulet et al.  
 6,953,776 B2 10/2005 Di Napoli  
 7,005,436 B2 2/2006 Lloyd et al.  
 7,265,108 B2 9/2007 Ozaki  
 7,335,667 B2 2/2008 Rodgers  
 7,358,255 B2 4/2008 Nakamura  
 7,517,870 B2 4/2009 Auricchio  
 7,598,257 B2 10/2009 Rodgers  
 7,750,007 B2 7/2010 Bearss et al.  
 7,834,022 B2 11/2010 Rodgers  
 8,053,433 B2 11/2011 Rodgers  
 8,158,616 B2 4/2012 Rodgers  
 8,309,718 B2 11/2012 Li  
 8,410,265 B2 4/2013 Zhou  
 8,415,362 B2 4/2013 Rodgers  
 8,420,629 B2 4/2013 Rodgers  
 8,445,488 B2 5/2013 Rodgers  
 8,486,902 B2 7/2013 Rodgers  
 8,513,270 B2 8/2013 Arvanitis  
 8,563,541 B2 10/2013 Arvanitis  
 8,604,043 B2 12/2013 Li  
 8,722,693 B2 5/2014 Rodgers  
 2002/0111353 A1 8/2002 Ledebor et al.  
 2003/0100756 A1 5/2003 Adams et al.  
 2003/0144309 A1 7/2003 Choon-Moon  
 2003/0165576 A1 9/2003 Fujii et al.  
 2004/0009222 A1 1/2004 Chou et al.

(Continued)

**FOREIGN PATENT DOCUMENTS**

DE 30 36 390 A1 5/1982  
 JP 7010876 A 1/1995

(Continued)

**OTHER PUBLICATIONS**Mathers, "Evaporation from the ocular surface", Exp Eye Res, 2004,  
vol. 78, pp. 389-394.McNamara, et al., "Fluorometry in contact lens research: The next  
step", Optom Vis Sci, 1998, vol. 75, pp. 316-322.Mengher, et al., "Non-invasive tear film break-up time: sensitivity  
and specificity", Acta Ophthalmol (Copenh), 1986, vol. 64, No. 4, pp.  
441-444.Mesa, et al. "INCB018424, A Selective JAK 1/2 Inhibitor, Signifi-  
cantly Improves the Compromised Nutritional Status and Frank  
Cachexia in Patients with Myelofibrosis (MF)" Poster #1760 at the  
American Society of Hematology Annual Meeting (ASH), Dec. 6,  
2008 (19 pages).

(Continued)

*Primary Examiner* — Susanna Moore(74) *Attorney, Agent, or Firm* — Lathrop & Gage LLP; Brian  
C. Trinquet(57) **ABSTRACT**

The present invention provides salt forms of (R)-3-(4-(7H-  
pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclo-  
pentylpropanenitrile that are useful in the modulation of  
Janus kinase activity and are useful in the treatment of dis-  
eases related to activity of Janus kinases including, for  
example, immune-related diseases, skin disorders, myeloid  
proliferative disorders, cancer, and other diseases.

**14 Claims, No Drawings**

(56)

**References Cited**

## U.S. PATENT DOCUMENTS

2004/0009983	A1	1/2004	Cox et al.	WO	97/02266	A1	1/1997
2004/0029857	A1	2/2004	Hale et al.	WO	97/36587	A1	10/1997
2004/0077654	A1	4/2004	Bouillot	WO	97/38664	A1	10/1997
2004/0198737	A1	10/2004	Cox et al.	WO	97/45412	A1	12/1997
2004/0204404	A1	10/2004	Zelle	WO	98/44797	A1	10/1998
2004/0214928	A1	10/2004	Aronov	WO	98/51391	A1	11/1998
2004/0235862	A1	11/2004	Burns	WO	99/00654	A2	1/1999
2005/0014966	A1	1/2005	Tabé	WO	W09900654	A2	1/1999
2005/0054568	A1	3/2005	Ling	WO	99/62908	A1	12/1999
2005/0153989	A1	7/2005	Grotzfeld et al.	WO	99/65908	A1	12/1999
2006/0004010	A1	1/2006	Habashita et al.	WO	99/65909	A1	12/1999
2006/0079511	A1	4/2006	Liu et al.	WO	00/09495	A1	2/2000
2006/0106020	A1	5/2006	Rodgers et al.	WO	00/51614	A1	9/2000
2006/0106027	A1	5/2006	Furet et al.	WO	00/53595	A1	9/2000
2006/0128803	A1	6/2006	Klimko	WO	00/63168	A1	10/2000
2006/0135537	A1	6/2006	Knegtel et al.	WO	01/14402	A1	3/2001
2006/0183761	A1	8/2006	Ledeboer et al.	WO	01/42246	A2	6/2001
2006/0183906	A1	8/2006	Rodgers et al.	WO	01/64655	A1	9/2001
2006/0223864	A1	10/2006	Biju	WO	01/81345	A1	11/2001
2006/0293311	A1	12/2006	Li et al.	WO	01/98344	A2	12/2001
2007/0135461	A1	6/2007	Rodgers et al.	WO	02/00196	A2	1/2002
2007/0135466	A1	6/2007	Ledeboer et al.	WO	02/00661	A1	1/2002
2007/0149506	A1	6/2007	Arvanitis et al.	WO	02/46184	A1	6/2002
2007/0149561	A1	6/2007	Dhanak et al.	WO	02/055084	A1	7/2002
2007/0191405	A1	8/2007	Noronha	WO	02/055496	A1	7/2002
2007/0208053	A1	9/2007	Arnold et al.	WO	02/060492	A1	8/2002
2007/0259904	A1	11/2007	Noronha	WO	02/092573	A2	11/2002
2008/0021026	A1	1/2008	Borchardt et al.	WO	02/096909	A1	12/2002
2008/0085898	A1	4/2008	Lu	WO	03/000695	A1	1/2003
2008/0096852	A1	4/2008	Yanni	WO	03/011285	A1	2/2003
2008/0119496	A1	5/2008	Ohlmeyer	WO	03/024967	A2	3/2003
2008/0161346	A1	7/2008	Cheng	WO	03/037347	A1	5/2003
2008/0188500	A1	8/2008	Arvanitis et al.	WO	03/048162	A1	6/2003
2008/0194468	A1	8/2008	Bodor	WO	03/099771	A2	12/2003
2008/0207584	A1	8/2008	Habashita et al.	WO	03/099796	A1	12/2003
2008/0280876	A1	11/2008	Hobson et al.	WO	2004/003026	A1	1/2004
2008/0312258	A1	12/2008	Rodgers et al.	WO	2004/005281	A1	1/2004
2008/0312259	A1	12/2008	Rodgers et al.	WO	2004/005282	A1	1/2004
2009/0018156	A1	1/2009	Tang et al.	WO	2004/026406	A1	4/2004
2009/0076070	A1	3/2009	Harada et al.	WO	2004/041814	A1	5/2004
2009/0088445	A1	4/2009	Ledeboer et al.	WO	2004/046120	A2	6/2004
2009/0131403	A1	5/2009	Kusuda	WO	2004/047843	A1	6/2004
2009/0181959	A1	7/2009	Rodgers et al.	WO	2004/056786	A2	7/2004
2009/0197869	A1	8/2009	Arvanitis et al.	WO	2004/072063	A1	8/2004
2009/0203637	A1	8/2009	Hocek et al.	WO	2004/080980	A1	9/2004
2009/0215766	A1	8/2009	Rodgers et al.	WO	2004/092154	A1	10/2004
2009/0233903	A1	9/2009	Rodgers et al.	WO	2004/099204	A1	11/2004
2009/0318405	A1	12/2009	Li et al.	WO	2004/099205	A1	11/2004
2010/0022522	A1	1/2010	Rodgers et al.	WO	2005/005988	A2	1/2005
2010/0069381	A1	3/2010	Itoh et al.	WO	2005/013986	A1	2/2005
2010/0113416	A1	5/2010	Friedman et al.	WO	2005/020921	A2	3/2005
2010/0190981	A1	7/2010	Zhou et al.	WO	2005/026129	A1	3/2005
2010/0210627	A1	8/2010	Mao et al.	WO	2005/028444	A1	3/2005
2010/0298334	A1	11/2010	Rodgers et al.	WO	2005/049033	A1	5/2005
2010/0298355	A1	11/2010	Li et al.	WO	2005/051393	A1	6/2005
2011/0059951	A1	3/2011	Rodgers et al.	WO	2005/060972	A2	7/2005
2011/0082159	A1	4/2011	Rodgers et al.	WO	2005/061463	A1	7/2005
2011/0086810	A1	4/2011	Rodgers et al.	WO	2005/062795	A2	7/2005
2011/0086835	A1	4/2011	Rodgers et al.	WO	2005/089502	A2	9/2005
2011/0207754	A1	8/2011	Li et al.	WO	2005/095400	A1	10/2005
2011/0223210	A1	9/2011	Rodgers et al.	WO	2005/105146	A1	11/2005
2011/0224157	A1	9/2011	Rodgers et al.	WO	2005/105814	A1	11/2005
2011/0224190	A1	9/2011	Huang et al.	WO	2005/105988	A2	11/2005
2011/0288107	A1	11/2011	Parikh et al.	WO	2005/110410	A2	11/2005
2012/0014989	A1	1/2012	Rodgers	WO	2005/117909	A2	12/2005
2012/0149681	A1	6/2012	Rodgers	WO	2005/121130	A2	12/2005
2012/0149682	A1	6/2012	Rodgers et al.	WO	2005/123719	A1	12/2005
2012/0214825	A1	8/2012	Vannucchi et al.	WO	2006/004984	A1	1/2006
				WO	2006/013114	A1	2/2006
				WO	2006/022459	A1	3/2006
				WO	2006/039718	A2	4/2006
				WO	2006/046023	A1	5/2006
				WO	2006/046024	A1	5/2006
				WO	2006/052913	A1	5/2006
				WO	2006/056399	A2	6/2006
				WO	2006/067445	A2	6/2006
				WO	2006/069080	A2	6/2006
				WO	2006/077499	A1	7/2006
				WO	2006/096270	A1	9/2006

## FOREIGN PATENT DOCUMENTS

JP 2003155285 A 5/2003  
WO 96/30343 A1 10/1996  
WO 97/02262 A1 1/1997

(56)

**References Cited**

## FOREIGN PATENT DOCUMENTS

WO 2006/101783 A2 9/2006  
 WO 2006/108103 A1 10/2006  
 WO 2006/127587 A1 11/2006  
 WO 2006/129199 A1 12/2006  
 WO 2006/136823 A1 12/2006  
 WO 2007/002433 A1 1/2007  
 WO 2007/025090 A2 3/2007  
 WO 2007/041130 A2 4/2007  
 WO 2007/043677 A1 4/2007  
 WO 2007/044894 A2 4/2007  
 WO 2007/049041 A1 5/2007  
 WO 2007/062459 A1 6/2007  
 WO 2007/070514 A1 6/2007  
 WO 2007/076423 A2 7/2007  
 WO 2007/077949 A1 7/2007  
 WO 2007/084557 A2 7/2007  
 WO 2007/090141 A2 8/2007  
 WO 2007/090748 A1 8/2007  
 WO 2007/117494 A1 10/2007  
 WO 2008/043031 A1 10/2007  
 WO 2007/129195 A2 11/2007  
 WO 2007/140222 A2 12/2007  
 WO 2008/013925 A2 1/2008  
 WO 2008/028937 A1 3/2008  
 WO 2008/035376 A2 3/2008  
 WO 2008/043031 A1 4/2008  
 WO 2008/058126 A2 5/2008  
 WO 2008/064157 A1 5/2008  
 WO 2008/067119 A2 6/2008  
 WO 2008/077712 A1 7/2008  
 WO 2008/079291 A2 7/2008  
 WO 2008/079292 A1 7/2008  
 WO 2008/082198 A1 7/2008  
 WO 2008/082839 A2 7/2008  
 WO 2008/082840 A1 7/2008  
 WO 2008/106692 A1 9/2008  
 WO 2008/124323 A1 10/2008  
 WO 2008/139161 A1 11/2008  
 WO 2008/145681 A2 12/2008  
 WO 2008/145688 A2 12/2008  
 WO 2008/157207 A2 12/2008  
 WO 2008/157208 A2 12/2008  
 WO 2009/016460 A2 2/2009  
 WO 2009/049028 A1 4/2009  
 WO 2009/064486 A2 5/2009  
 WO 2009/064835 A1 5/2009  
 WO 2009/071577 A1 6/2009  
 WO 2009/100130 A1 8/2009  
 WO 2009/109576 A1 9/2009  
 WO 2009/114512 A1 9/2009  
 WO 2009/115572 A2 9/2009  
 WO 2009/158687 A1 12/2009  
 WO 2010/000978 A1 1/2010  
 WO 2010/001169 A2 1/2010  
 WO 2010/020905 A1 2/2010  
 WO 2010/022076 A1 2/2010  
 WO 2010/022081 A1 2/2010  
 WO 2010/026121 A1 3/2010  
 WO 2010/026122 A1 3/2010  
 WO 2010/026124 A1 3/2010  
 WO 2009/049028 A1 4/2010  
 WO 2010/081692 A1 7/2010  
 WO 2010/083283 A2 7/2010  
 WO 2010/135621 A1 11/2010  
 WO 2010/135650 A1 11/2010  
 WO 2011/025685 A1 3/2011  
 WO 2011/028685 A1 3/2011  
 WO 2011/029802 A1 3/2011  
 WO 2011/031554 A2 3/2011  
 WO 2011/035900 A1 3/2011  
 WO 2011/044481 A1 4/2011  
 WO 2011/057784 A1 5/2011  
 WO 2011/069141 A2 6/2011  
 WO 2011/112662 A1 9/2011  
 WO 2011/130146 A1 10/2011

WO 2011/144338 A1 11/2011  
 WO 2012/003457 A1 1/2012  
 WO 2012/068440 A1 5/2012  
 WO 2012/068450 A1 5/2012

## OTHER PUBLICATIONS

Mesa, et al., "Evaluating the serial use of the myelofibrosis symptom assessment form for measuring symptomatic improvement: Performance in 87 myelofibrosis patients on a JAK1 and JAK2 inhibitor (INCB018424) clinical trial", *Cancer*, 2011, vol. 117, No. 21, pp. 4869-4877.

Mesa, et al., "Emerging drugs for the therapy of primary and post essential thrombocythemia, post polycythemia vera myelofibrosis", *Expert Opinion on Emerging Drugs England*, 2009, vol. 14, No. 3, pp. 471-479.

Meydan, et al., "Inhibition of acute lymphoblastic leukaemia by a Jak-2 inhibitor", *Nature*, 1996, vol. 379, No. 6566, pp. 645-648.

Miethchen, "Micelle-activated reactions. I. Micelle-activated iodination and partial dehalogenation of pyrazoles and 1,2,4-triazoles", *Journal F. prakt. Chemie, Band 331, Heft 5, S. 799-805 (1989)* (1 page abstract also provided).

Minegishi, et al., "Human Tyrosine Kinase 2 Deficiency Reveals Its Requisite Roles in Multiple Cytokine Signals Involved in Innate and Acquired Immunity", *Immunity* 2006, vol. 25, pp. 745-755.

Mishchenko, et al., "Treatment options for hydroxyurea-refractory disease complications in myeloproliferative neoplasms: JAK2 inhibitors, radiotherapy, splenectomy and transjugular intrahepatic portosystemic shunt", *Eur J Haematol*. 2010, vol. 85, No. 3, pp. 192-199.

Mishima, et al., "Determination of tear volume and tear flow", *Invest Ophthalmol*, 1966, vol. 5, pp. 264-276.

Mishima, "Some physiological aspects of the precorneal tear film", *Arch Ophthalmol*, 1965, vol. 73, pp. 233-241.

Mitsunobu, "The Use of Diethyl Axodicarboxylate and Triphenylphosphine in Synthesis and Transformation of Natural Products." *Synthesis* 1981, No. 1, pp. 1-28 (1981).

Miyata, et al., "Stereospecific nucleophilic addition reactions to olefins", *J. Org. Chem.* 1991, vol. 56, pp. 6556-6564.

Miyaura et al., "Palladium-Catalyzed Cross-Coupling Reactions of Organoboron Compounds," *Chem. Rev.*, 1995, vol. 95, pp. 2457-2483.

Miyoshi et al., "Interleukin-8 concentrations in conjunctival epithelium brush cytology samples correlate with neutrophil, eosinophil infiltration, and corneal damage", *Cornea*, 2001, vol. 20, pp. 743-747.

Moreland, et al. "A Randomized Placebo-Controlled Study of INCB018424, a Selective Janus Kinase 1&2 (JAK 1&2) Inhibitor in Rheumatoid Arthritis (RA)" Presentation at the American College of Rheumatology meeting, Oct. 26, 2008. (20 pages).

Moriarty, et al., "The synthesis and SAR of 2-amino-pyrrolo[2,3-d]pyrimidines: A new class of Aurora-A kinase inhibitors", *Bioorganic and Medicinal Chemistry Letters*, 2006 vol. 16, No. 22, pp. 5778-5783.

Mosby's Dictionary of Medicine, Nursing, & Health Professions, sicca complex, 2009, Elsevier, printed from [http://www.credoreference.com/entry/ehsmosbymed/sicca\\_complex](http://www.credoreference.com/entry/ehsmosbymed/sicca_complex), 2 pages.

Mullighan, et al, "JAK mutations in high-risk childhood acute lymphoblastic leukemia", *Proc Natl Acad Sci USA*. 2009, vol. 106, pp. 9414-9418.

Naka, "The paradigm of IL-6: from basic science to medicine", *Arthritis Res*. 2002, vol. 4, Suppl. 3, pp. S233-S242.

Nally, et al., "Ocular discomfort and tear film break-up time in dry eye patients: A correlation", *Invest Ophthalmol Vis Sci*, 2000, vol. 41, No. 4, pp. 1436 (Poster Presentation).

Naqvi, et al., "A potential role of ruxolitinib in leukemia", *Expert Opinion on Investigational Drugs*, 2011, vol. 20, No. 8, pp. 1159-1166.

National Cancer Institute, "FDA Approval for Ruxolitinib Phosphate", <http://www.cancer.gov/cancertopics/druginfo/fda-ruxolitinibphosphate> posted Nov. 18, 2011 (3 pages).

Naus, et al., "6-(Het)ary1-7-Deazapurine Ribonucleosides as Novel Potent Cytostatic Agents", *J. Med. Chem.*, 2010, vol. 53, No. 1, pp. 460-470.

(56)

**References Cited**

## OTHER PUBLICATIONS

- Neidle, "Chapter 18: Failure modes in anticancer drug discovery and development" in *Cancer Drug Discovery and Design* (Elsevier/Academic Press, 2008), pp. 427-431.
- Nelson, et al., "Tear film osmolality determination: an evaluation of potential errors in measurement" *Curr Eye Res*, 1986, vol. 5, No. 9, pp. 677-681.
- Nicholoff, et al., "Recent Insights into the immunopathogenesis of psoriasis provide new therapeutic opportunities", *J. Clin. Invest.*, 2004, vol. 113, pp. 1664-1675.
- Nichols, et al., "The lack of association between signs and symptoms in patients with dry eye disease", *Cornea*, 2004, vol. 23, No. 8, pp. 762-770.
- Nichols, et al., "The repeatability of clinical measurements of dry eye", *Cornea*, 2004, vol. 23, No. 3, pp. 272-285.
- Nitta, et al., "Peptide-Titanium Complex as Catalyst for Asymmetric Addition of Hydrogen Cyanide to Aldehyde", *J. Am. Chem. Soc.*, 1992, Vol. 114, pp. 7969-7975.
- Norn, M., "Quantitative tear ferning. Clinical investigations", *Acta Ophthalmol* (Copenh), 1994, vol. 72, No. 3, pp. 369-372.
- Notice of Allowance and Fee(s) Due dated Sep. 21, 2007 in connection with U.S. Appl. No. 11/313,394 (6 pages).
- Notice of Hearing and Preliminary Report for EP Patent 1966202, dated Mar. 18, 2013 (7 pages).
- Office Action (Non-final) dated Aug. 22, 2007 in connection with U.S. Appl. No. 11/115,702 (9 pages).
- Office Action (Non-final) dated Dec. 3, 2007 in connection with U.S. Appl. No. 11/524,641 (13 pages).
- Office Action (Non-final) dated Feb. 25, 2009 for U.S. Appl. No. 12/137,892 (13 pgs.).
- Office Action (Final) dated Feb. 7, 2008 for U.S. Appl. No. 11/115,702 (5 pages).
- Office Action (Final) dated Nov. 30, 2009 for U.S. Appl. No. 12/137,892 (9 pgs.).
- Office Action (Non-final) dated Apr. 20, 2007 in connection with U.S. Appl. No. 11/313,394 (16 pages).
- Office Action received for European Application No. 06 839 328.9 (Jan. 22, 2009) (5 pages).
- Office Action received for Japanese Application No. 2008-545733 dated Oct. 11, 2011 (5 pages).
- Office Action received for New Zealand Application No. 569015 dated Feb. 24, 2010 (2 pages).
- Office Action received for Singapore Application No. 2008-04386-1 (Aug. 24, 2010).
- Office Action received for Vietnamese Patent Application No. 1-2011-03188 dated Mar. 8, 2012 as translated by foreign associate (10 pages).
- Office Action, Canadian Patent Office, Application No. 2,632,466, dated May 8, 2012 (3 pages).
- Office Action, Eurasian Patent Office, prepared Feb. 5, 2010.
- Office Action, European Patent Office, Application No. 06 839 328.9 mailed Oct. 21, 2010.
- Office Action, European Patent Office, mailed Nov. 6, 2009.
- Office Action, Mexico, Patent Appl. No. MX/a/2008/007635 as received from foreign associate dated Jun. 15, 2010 (1 page).
- STN Search conducted Nov. 5, 2010 (5 pages).
- STN Search conducted Nov. 9, 2010 (43 pages).
- STN Search, Nov. 12, 2009 (180 pages).
- STN Search, Oct. 20, 2009 (601 pages).
- STN Search, Sep. 20, 2009 (864 pages).
- Sullivan, et al., "4th International Conference on the Lacrimal Gland, Tear Film & Ocular Surface and Dry Eye Syndromes, Nov. 20, 2004" (2 pages).
- Takahashi, et al., "Solvent-Free Reaction Using Pmospwonium Salts: Chlorination of Hydroxyheteroaromatics and dehydration of Primary Amides", *Heterocycles* 2006, vol. 68, pp. 1973-1979.
- Takano, et al., "Inflammatory cells in brush cytology samples correlate with the severity of corneal lesions in atopic keratoconjunctivitis", *Br J Ophthalmol*, 2004, vol. 88, pp. 1504-1505.
- Tan, et al., "Racemization processes at a quaternary carbon center in the context of the asymmetric Michael reaction", *Tetrahedron Lett.*, 2001, vol. 42, No. 30, pp. 5021-5023.
- Tang, et al., "Knowledge-based design of 7-azaindoles as selective B-Raf inhibitors", *Bioorganic & Medicinal Chemistry Letters* 2008, vol. 18, No. 16, pp. 4610-4614.
- Tasian, et al., "Understanding the biology of CRLF2-overexpressing acute lymphoblastic leukemia", *Critical Reviews in Oncogenesis*, 2011, vol. 16, No. 1-2, pp. 13-24.
- Tefferi, et al., "The Clinical Phenotype of Myelofibrosis Encompasses a Chronic Inflammatory State that is Favorably Altered by INCB018424, A Selective Inhibitor of JAK1/2" Poster #2804 at the American Society of Hematology Annual Meeting (ASH), Dec. 7, 2008, (18 pages).
- Tefferi, "Primary myelofibrosis: 2012 update on diagnosis, risk stratification, and management", *American Journal of Hematology*, 2011, vol. 86, No. 12, pp. 1017-1026.
- Tefferi, et al., "Serious adverse events during ruxolitinib treatment discontinuation in patients with myelofibrosis", *Mayo Clinic Proceedings*, 2011, vol. 86, No. 12, pp. 1188-1191.
- Tiffany, et al., "Meniscometry using the Tearscope-plus (ARVO abstract). *Invest Ophthalmol Vis Sci*, 2001, vol. 42, No. S37 (1 page).
- Tiffany, "Refractive index of meibomian and other lipids", *Curr Eye Res*, 1986, vol. 5, pp. 887-889.
- Toyonaga, "Blockade of constitutively activated Janus kinase/signal transducer and activator of transcription-3 pathway inhibits growth of human pancreatic cancer", *Cancer Lett.* 2003, vol. 201, No. 1, pp. 107-116.
- Tsubota, et al., "Brush cytology for the evaluation of dry-eye", *Nippon Ganka Gakkai Zasshi*, 1990a, vol. 94, pp. 224-230 (in Japanese with English abstract).
- Tsubota et al., "Conjunctival brush cytology", *Acta Cytol*, 1990, vol. 34, No. 2, pp. 233-235.
- Tsubota et al., "Detection by brush cytology of mast cells and eosinophils in allergic and vernal conjunctivitis", *Cornea*, 1991, vol. 10, No. 6, pp. 525-531.
- Ueda et al., "1,2-Benzisoxazol-3-yl Diphenyl Phosphate: A New, Reactive Activating Agent for the Synthesis of Amides, Esters, and Peptides via Condensation", *J. Org. Chem.* 1985, vol. 50, pp. 760-763.
- van Best et al., "Measurement of basal tear turnover using a standardized protocol", *Graefe's Arch Clin Exp Ophthalmol*, 1995, vol. 233, pp. 1-7.
- van Bijsterveld, "Diagnostic tests in the sicca syndrome", *Arch Ophthalmol*, 1969, vol. 82, pp. 10-14.
- Vannucchi, et al., "The mTOR Inhibitor, RAD001, Inhibits the Growth of Cells From Patients with Myeloproliferative Neoplasms", *Blood: ASH Annual Meeting Abstracts*, 51st Annual Meeting of the American Society of Hematology, 2009, vol. 114, No. 22 (2 pages).
- Vannucchi, et al., "Inhibitors of PI3K/Akt and/or mTOR Inhibit the Growth of Cells of Myeloproliferative Neoplasms and Synergize with JAK2 Inhibitor and Interferon", *Blood*, 2011, vol. 118, No. 21, pp. 1638-1639, XP008150742 ASH Annual Meeting Abstract 3835 American Society of Hematology.
- Vannucchi, et al., "RAD001, An Inhibitor of mTOR, Shows Clinical Activity in a Phase I/II Study in Patients with Primary Myelofibrosis (PMF) and Post Polycythemia Vera/Essential Thrombocythemia Myelofibrosis (PPV/PET MF)", *Blood, ASH Annual Meeting Abstracts* 307, 2009, vol. 114, No. 22 (2 pages).
- Vasilevsky, et al., "Ethyl Vinyl Ether—an Agent for Protection of the Pyrazole NH-Fragment. A Convenient Method for the Preparation of N-Unsubstituted 6-Alkynylpyrazoles", *Heterocycles*, 2003, vol. 60, No. 4, pp. 879-886.
- Verma, et al., "Jak family of kinases in cancer", *Cancer and Metastasis Reviews*, 2003, vol. 22, No. 4, pp. 423-434.
- Verstovsek, "Therapeutic Potential of JAK2 Inhibitors", *Hematology Am Soc Hematol Educ Program*, 2009, pp. 636-642.
- Verstovsek, et al. "The JAK Inhibitor INCB018424 Demonstrates Durable and Marked Clinical Responses in Primary Myelofibrosis (PMF) and Post-Polycythemia/Essential Thrombocythemia Myelofibrosis (Post-PV/ET-MF)" Poster #1762 at the American Society of Hematology Annual Meeting (ASH), Dec. 6, 2008 (19 pages).

(56)

## References Cited

## OTHER PUBLICATIONS

- Verstovsek, et al., "Characterization of JAKS V617F Allele Burden in Advanced Myelofibrosis (MF) Patients: No Change in V617F:WT JAK2 Ratio in Patients with High Allele Burdens despite Profound Clinical Improvement Following Treatment with the JAKL Inhibitor, INCB018424," 50th ASH Annual Meeting and Exposition, Abstract No. 2802 (2008).
- Vitali, et al., "The European Community Study Group on diagnostic criteria for Sjogren's syndrome. Sensitivity and specificity of tests for ocular and oral involvement in Sjogren's syndrome." *Ann Rheum Dis*, 1994, vol. 53, No. 10, pp. 637-647.
- Wagh, et al., "Polymers used in ocular dosage form and drug delivery systems", *Asian J. Pharm.*, 2008, pp. 12-17.
- WebMD. "Diabetes Health Center." Available at: < [http://diabetes.webmd.com/guide/diabetestreatment\\_care](http://diabetes.webmd.com/guide/diabetestreatment_care) >. 3 pages, downloaded Jan. 6, 2014.
- Webster's New World Medical Dictionary, Sjogren's syndrome, 2003, Wiley Publishing, printed from [http://www.credoreference.com/entry/webstermed/sjogren\\_s\\_syndrome](http://www.credoreference.com/entry/webstermed/sjogren_s_syndrome), 2 pages.
- Weiss, et al., "Evaluation of a Series of Naphthamides as Potent, Orally Active Vascular Endothelial Growth Factor Receptor-2 Tyrosine Kinase Inhibitors", *J. Med Chem.*, 2008, vol. 51, pp. 1668-1680.
- Welch et al., "An approach to a more standardized method of evaluating tear film break-up time", *Invest Ophthalmol Vis Sci*, 2003; 2485/B324 (abstract only—2 pages).
- White et al., "Human basic tear fluid osmolality. I. Importance of sample collection strategy", *Acta Ophthalmol (Copenh)*, 1993, vol. 71, No. 4, pp. 524-529.
- Williams et al., "Carbohydrate Chemistry: Recent Advances", *Chem. Rev.* 1981, vol. 81, pp. 589-636.
- Williams, et al. "Initial Efficacy of INCB018424, a selective Janus Kinase1& 2 (JAK1&2) Inhibitor in Rheumatoid Arthritis (RA)," European League Against Rheumatism (EULAR) meeting presentation and abstract (Jun. 11-14, 2008, Paris, France). *Annals Rheum Dis* 67SII:62, 2008.
- Wolf, et al., "Burger's Medicinal Chemistry and Drug Discovery", 5th Ed. Part I, pp. 975-977 (1995).
- Xiaoyang et al., "Knockdown of STAT3 Expression by RNA Interference Inhibits the Induction of Breast Tumors in Immunocompetent Mice", *Cancer Res* 2005, vol. 65, pp. 2532.
- Yamaoka et al., "Janus kinase (JAK) inhibitors in rheumatoid arthritis", *Current Rheumatology Reviews*, 2011, vol. 7, No. 4, pp. 306-312.
- Yang et al., "Constitutive NF- $\kappa$ B activation confers interleukin 6 (IL6) independence and resistance to dexamethasone and Janus kinase inhibitor INCB018424 in murine plasmacytoma cells", *Journal of Biological Chemistry*, 2011, vol. 286, No. 32, pp. 27988-27997.
- Yao, et al. "Glucocorticoid-Induced Bone Loss in Mice Can Be Reversed by the Actions of Parathyroid Hormone and Risedronate on Different Pathways for Bone Formation and Mineralization", *Arthritis and Rheumatism*, 2008, vol. 58, pp. 3485-3497.
- Yao, et al., "Glucocorticoid Excess in Mice Results in Early Activation of Osteoclastogenesis and Adipogenesis and Prolonged Suppression of Osteogenesis", *Arthritis and Rheumatism*, 2008, vol. 58, No. 6, pp. 1674-1686.
- Yokoi et al., "A newly developed video-meibography system featuring a newly designed probe", *Jpn J Ophthalmol*, 2007, vol. 51, pp. 53-56.
- Yokoi et al., "Assessment of meibomian gland function in dry eye using meibometry", *Arch Ophthalmol*, 1999, vol. 117, pp. 723-729.
- Yokoi et al., "Correlation of tear lipid layer interference patterns with the diagnosis and severity of dry eye", *Am J Ophthalmol*, 1996, vol. 122, pp. 818-824.
- Yokoi et al., "Non-invasive methods of assessing the tear film", *Exp Eye Res*, 2004, vol. 78, pp. 399-407.
- Yu, et al., "Constitutive activation of the Janus kinase-STAT pathway in T lymphoma overexpressing the Lck protein tyrosine kinase", *J Immunol*. 1997, vol. 159, No. 11, pp. 5206-5210.
- Zheng, et al., "Discovery of INCB108201PF-4178903, a potent, selective, and orally bioavailable dual CCR2 and CCR5 antagonist", *Bioorganic & Medicinal Chemistry Letters* 2011, vol. 21, pp. 1442-1445.
- Zoppellaro, et al., "A Multifunctional High-Spin Biradical Pyrazolylbipyridine-bisnitronyl nitroxide", *Org. Lett.* 2004, Vol. , No. 26, pp. 4929-4932.
- Manjula, et al. "The Protein Kinase Complement of the Human Genome", *Science*, 2002, vol. 298, Iss. 5600, pp. 1912-1916 and 1933-1934.
- Minegishi, et al. "Human Tyrosine Kinase 2 Deficiency Reveals Its Requisite Roles in Multiple Cytokine Signals Involved in Innate and Acquired Immunity" *Immunity* 2006, vol. 25, pp. 745-755.
- James, et al., "A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera", *Nature*, 2005, vol. 434, No. 7037, pp. 1144-1148.
- Kawamura, et al., "Molecular cloning of L-JAK, a Janus family protein-tyrosine kinase expressed in natural killer cells and activated leukocytes", *Proc Natl Acad Sci USA*, 1994, vol. 91, No. 14, pp. 6374-6378.
- Kharas, et al., "ABL Oncogenes and Phosphoinositide 3-Kinase: Mechanism of Activation and Downstream Effectors.", *Cancer Res.*, 2005, vol. 65, No. 6, pp. 2047-2053.
- Kruh et al., "The complete coding sequence of arg defines the Abelson subfamily of cytoplasmic tyrosine kinases.", *Proc. Natl. Acad. Sci.*, 1990, vol. 87, pp. 5802-5806.
- Kubinyi, H. "QSAR: Hansch Analysis and Related Approaches," *Methods and Principles in Medicinal Chemistry*, Manhold, R. ed. Weinheim, NY, 1993.
- Kudelacz, et al. "The JAK-3 inhibitor CP-690550 is a potent anti-inflammatory agent in a murine model of pulmonary eosinophilia", *European Journal of Pharmacology* 2008, vol. 582, pp. 154-161.
- Letter translation of Office Action, Chilean Application No. 3496-2006 as received from the foreign associate (Jul. 5, 2010) (4 pages).
- Levine, et al., "Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis", *Cancer Cell*, 2005, vol. 7, pp. 387-397.
- Levy, et al., INCB18424 Discussion presentation at the American Society of Hematology, 49th Annual Meeting and Exposition, Atlanta, GA. Abstract #558, Dec. 10, 2007 (25 pages).
- Madhusudan et al., "Tyrosine kinase inhibitors in cancer therapy", *Clin Biochem.*, 2004, vol. 37, No. 7, pp. 618-635.
- Manning, et al., "The Protein Kinase Complement of the Human Genome", *Science*. 2002, vol. 298, No. 5600, pp. 1912-1916 and 1933-1934.
- Methods in Molecular Biology*: vol. 225, *Inflammation Protocols.*, Winyard, P.G. and Willoughby, D.A., Humana Press, 2003.
- Milici, A.J., et al., "Cartilage preservation by inhibition of Janus kinase 3 in two rodent models of rheumatoid arthritis", *Arthritis Research & Therapy*, 2008, vol. 10, pp. R14 (<http://arthritis-research.com/content/10/1/R14>) (9 pages).
- Nakagawara, Akira, "Trk receptor tyrosine kinases: A bridge between cancer and neural development." *Cancer Letters*, 2001, vol. 169, pp. 107-114.
- Neubauer, H., et al., "Jak2 Deficiency Defines an Essential Developmental Checkpoint in Definitive Hematopoiesis", *Cell*, 1998, vol. 93, No. 3, pp. 397-409.
- Nishio, et al., "Tyrosine kinase-dependent modulation by interferon- $\alpha$  of the ATP-sensitive K<sup>+</sup> current in rabbit ventricular myocytes", *FEBS Letters*, 1999, vol. 445, pp. 87-91.
- Palmer, et al., "Multiple roles of ephrins in morphogenesis, neuronal networking, and brain function." *Genes & Dev.*, 2003, vol. 17, pp. 1429-1450.
- Parganas, et al., "Jak2 is Essential for Signaling through a Variety of Cytokine Receptors", *Cell*, 1998, vol. 93, No. 3, pp. 385-395.
- Park et al., "Homogeneous Proximity Tyrosine Kinase Assays: Scintillation Proximity Assay versus Homogeneous Time-Resolved Fluorescence", *Analytical Biochemistry*, 1999, vol. 269, pp. 94-104.
- Patani, et al., "Bioisosterism: A Rational Approach in Drug Design", *Chem. Rev.*, 1996, vol. 96, pp. 3147-3176.
- Pernis, et al., "JAK-STAT signaling in asthma." *J Clin Invest*, 2002, vol. 109, No. 10, pp. 1279-1283 (2002).

(56)

## References Cited

## OTHER PUBLICATIONS

- Pirard, B. et al., "Classification of Kinase Inhibitors Using BCUT Descriptors", *J. Chem. Inf. Comput. Sci.*, 2000, vol. 40, pp. 1431-1440.
- Press Release dated Sep. 18, 2008: "Incyte's Topical JAK Inhibitor Demonstrates Positive Proof-of-Concept Results in Patients with Mild to Moderate Psoriasis" (4 pages).
- Punwani et al., Poster/presentation, "Initial Efficacy and Safety of Topical INCYB018424 Cream, a Selective Janus Kinase 1&2 (JAK 1&2) Inhibitor in Psoriasis" 17th Congress of the European Academy of Dermatology and Venereology, Paris, France, Sep. 17, 2008 (15 pages).
- Quesada et al., "One-pot conversion of activated alcohols into 1,1-dibromoalkenes and terminal alkynes using tandem oxidation processes with manganese dioxide", *Tetrahedron*, 2006, vol. 62, pp. 6673-6680.
- Ravin, "Preformulation", *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., 1985, Chapter 76, pp. 1409-1423.
- Rodig, et al., "Disruption of the Jak1 gene demonstrates obligatory and nonredundant roles of the Jaks in cytokine-induced biologic responses." *Cell*, 1998, vol. 93, No. 3, pp. 373-383.
- Roudebush, et al., "Pharmacologic manipulation of a four day marine delayed type hyper sensitivity model", *Agents Actions*, 1993, vol. 38, No. 1-2, pp. 116-121.
- Rousvoal, et al., "Janus kinase 3 inhibition with CP-690,550 prevents allograft vasculopathy", *Transpl Int.*, 2006, vol. 19, No. 12, pp. 1014-1021.
- Saemann, et al., "Prevention of CD40-triggered dendritic cell maturation and induction of T-cell hyporeactivity by targeting of Janus kinase 3", *Am J Transplant*, 2003, vol. 3, No. 11, pp. 1341-1349.
- Schindler, et al., "Hormones and Signaling: Cytokines and Stat Signaling", *Adv Pharmacol.* 2000; 47:113-74.
- Scott, et al., "Jaks, STATs, Cytokines, and Sepsis", *Clin Diagn Lab Immunol*, 2002, vol. 9, No. 6, pp. 1153-1159.
- Seto, et al., "Enhanced Th2 cell-mediated allergic inflammation in Tyk2-deficient mice", *J Immunol*, 2003, vol. 170, No. 2, pp. 1077-1083.
- Shah, et al., "Multiple BCR-ABL kinase domain mutations confer polyclonal resistance to the tyrosine kinase inhibitor imatinib (STI571) in chronic phase and blast crisis chronic myeloid leukemia." *Cancer Cell*, 2002, vol. 2, pp. 117-125.
- Sriram, et al., "Induction of gp130-related Cytokines and Activation of JAK2/STAT3 Pathway in Astrocytes Precedes Up-regulation of Glial Fibrillary Acidic Protein in the 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine Model of Neurodegeneration", *J. Biol. Chem.*, 2004, vol. 279, No. 19, pp. 19936-19947.
- Staerk, et al., "JAK1 and Tyk2 activation by the homologous polycythemia vera JAK2 V617F mutation: cross-talk with IGF1 receptor", *J Biol Chem.*, 2005, vol. 280, pp. 41893-41899.
- Takemoto, et al., "Proliferation of adult T cell leukemia/lymphoma cells is associated with the constitutive activation of JAK/STAT proteins." *Proc Natl Acad Sci USA*, 1997, vol. 94, No. 25, pp. 13897-13902.
- Thompson, et al., "Photochemical Preparation of a Pyridone Containing Tetracycline: A Jak Protein Kinase Inhibitor", *Bioorganic & Medicinal Chemistry Letters*, 12 (2002) 1219-1223.
- Verstovsek, et al., "The selective Janus kinase (JAK) inhibitor, INCB018424, shows efficacy in phase I/II trial in patients with primary myelofibrosis (PMF) and post polycythemia vera/essential thrombocythemia myelofibrosis (Post-PV/ET MF)" Abstract #0444, presented Saturday, Jun. 14, 2008 at the European Hematology Association, 13th Congress, Jun. 12-15, Copenhagen, Denmark (2 pages).
- Verstovsek, et al. INCB18424, an Oral, Selective JAK2 Inhibitor, Shows Significant Clinical Activity in a Phase I/II Study in Patient with Primary Myelofibrosis (PMF) and Post Polycythemia Vera/Essential Thrombocythemia Myelofibrosis (Post-PV/ET MF), presentation at the American Society of Hematology 49th Annual Meeting and Exposition, Dec. 10, 2007 (16 pages).
- Wu, et al., One-Pot Two-Step Microwave-Assisted Reaction in Construction 4,5-Disubstituted Pyrazolopyrimidines *Organic Letters*, 2003, vol. 5, No. 20, pp. 3587-3590.
- Zou, et al., "Signaling Pathways Activated by Oncogenic Forms of Abl Tyrosine Kinase." *Journal of Biological Chemistry*, 1997, vol. 274, No. 26, pp. 18141-18144.
- International Search Report and Written Opinion for International Appln. No. PCT/US2008/066662 dated Dec. 23, 2008 (11 pages).
- International Search Report and Written Opinion for International Appln. No. PCT/US2006/047369 dated Apr. 24, 2007 (16 pages).
- Manjula, et al., "Rapid Method of Converting Primary Amides to Nitriles and Nitriles to Primary Amides by ZnCl2 using Microwaves under Different Reaction Conditions", *Syn. Commun.* 2007, vol. 37, pp. 1545-1550.
- U.S. Appl. No. 12/137,892, filed Jun. 12, 2008, James D. Rodgers.
- U.S. Appl. No. 14/097,598, filed Dec. 5, 2013, James D. Rodgers.
- U.S. Appl. No. 14/256,311, filed Apr. 18, 2014, James D. Rodgers.
- U.S. Appl. No. 14/256,383, filed Apr. 18, 2014, James D. Rodgers.
- U.S. Appl. No. 14/270,915, filed May 6, 2014, James D. Rodgers.
- U.S. Appl. No. 13/761,742, filed Feb. 7, 2013, Jiacheng Zhou.
- U.S. Appl. No. 13/761,771, filed Feb. 7, 2013, Jiacheng Zhou.
- U.S. Appl. No. 13/761,830, filed Feb. 7, 2013, Jiacheng Zhou.
- U.S. Appl. No. 13/917,124, filed Jun. 13, 2013, James D. Rodgers.
- U.S. Appl. No. 13/030,682, filed Feb. 18, 2011, Yun-Long Li.
- U.S. Appl. No. 13/399,274, filed Feb. 17, 2012, Alessandro M. Vannucchi.
- U.S. Appl. No. 13/571,525, filed Aug. 10, 2012, Alessandro M. Vannucchi.
- U.S. Appl. No. 11/115,702, filed Apr. 27, 2005, James D. Rodgers.
- U.S. Appl. No. 11/524,641, filed Sep. 21, 2006, Argyrios G. Arvanitis.
- U.S. Appl. No. 12/187,061, filed Aug. 6, 2008, James D. Rodgers.
- U.S. Appl. No. 12/418,132, filed Apr. 3, 2009, Argyrios G. Arvanitis.
- U.S. Appl. No. 12/571,834, filed Oct. 1, 2009, Paul A. Friedman.
- U.S. Appl. No. 12/784,916, filed May 21, 2010, James D. Rodgers.
- U.S. Appl. No. 12/872,925, filed Aug. 31, 2010, James D. Rodgers.
- U.S. Appl. No. 13/043,986, filed Mar. 9, 2011, Taisheng Huang.
- U.S. Appl. No. 13/076,176, filed Mar. 30, 2011, James D. Rodgers.
- U.S. Appl. No. 13/076,220, filed Mar. 30, 2011, James D. Rodgers.
- U.S. Appl. No. 13/112,370, filed May 20, 2011, Bhavnish Parikh.
- U.S. Appl. No. 13/300,094, filed Nov. 18, 2011, James D. Rodgers.
- U.S. Appl. No. 13/300,137, filed Nov. 18, 2011, James D. Rodgers.
- U.S. Appl. No. 13/526,957, filed Jun. 19, 2012, Wenqing Yao.
- U.S. Appl. No. 13/588,776, filed Aug. 17, 2012, James D. Rodgers.
- U.S. Appl. No. 13/564,271, filed Aug. 1, 2012, Paul A. Friedman.
- U.S. Appl. No. 13/605,331, filed Sep. 6, 2012, Jiacheng Zhou.
- U.S. Appl. No. 13/754,533, filed Jan. 30, 2013, James D. Rodgers.
- U.S. Appl. No. 13/853,475, filed Mar. 29, 2013, James D. Rodgers.
- Office Action, Mexico, Patent Appl. No. MX/a/2008/007635 as received from foreign associate dated Nov. 13, 2009 (4 pages).
- Office Action/Examination Report received for Pakistan Application No. 211/2009 dated Jan. 18, 2010 (1 page).
- Oguz, et al., "The height and radius of the tear meniscus and methods for examining these parameters", *Cornea*, 2000, vol. 19, pp. 497-500.
- Opposition for EP Patent 1966202, filed on Jun. 21, 2012 (30 pages).
- Opposition for India Patent Application No. 2365/KOLNP/2008 dated Nov. 12, 2012 (received by Applicants from Indian associate on Apr. 17, 2013) 37 pages.
- Opposition, Costa Rica, translation from Foreign Associate Dated Jun. 13, 2012, 6 pages.
- Opposition, Ecuador Patent Office, mailed Nov. 18, 2008 1 page letter from Foreign Associate enclosing the translation (5 pages) of the Opposition.
- Ortmann, et al., "Janus kinases and signal transducers and activators of transcription: their roles in cytokine signaling, development and immunoregulation." *Arthritis Res*, 2(1): 16-32 (2000).
- Ostojic, et al., "Ruxolitinib for the treatment of myelofibrosis", *Drugs of Today*, 2011, vol. 47, No. 11, pp. 817-827.
- Ousler, et al., "Factors that influence the inter-blink interval (IBI) as measured by the ocular protection index (OPI)", *Invest Ophthalmol Vis Sci* 2001; 43: E-abstract 56 (Poster presentation) ARVO (2002) 2 pages, downloaded from <http://abstracts.iov.s.org/cgi/content/abstract/43/12/56?maxtoshow> on Aug. 14, 2009.

(56)

## References Cited

## OTHER PUBLICATIONS

- Pardanani, "JAK2 inhibitor therapy in myeloproliferative disorders: rationale, preclinical studies and ongoing clinical trials JAK2 inhibitor therapy in MPD", *Leukemia* 2008, vol. 22, pp. 23-30.
- Patrick, "An Introduction to medicinal chemistry" Oxford University Press Inc., New York (31 pages) (cited in Opposition from India dated Nov. 12, 2012).
- Pearce et al., "Spatial location studies on the chemical composition of human tear ferns", *Ophthalmic Physiol Opt*, 2000, vol. 20, No. 4, pp. 306-313.
- Pearce, et al., "An improved fluorophotometric method for tear turn-over assessment", *Optom Vis Sci*, 2001, vol. 78No. 1, pp. 30-36.
- Pensyl, et al., "The repeatability of tear mucus ferning grading", *Optom Vis Sci*, 1998, vol. 75, No. 8, pp. 600-604.
- Peters, et al., "Functional Significance of Tie2 Signaling in the Adult Vasculature", 2004, © The Endocrine Society.
- Pflugfelder, et al., "Evaluation of subjective assessments and objective diagnostic tests for diagnosing tear-film disorders known to cause ocular irritation", *Cornea*, 1998, vol. 17, No. 1, pp. 38-56.
- Pillonel, "Evaluation of phenylaminopyrimidines as antifungal protein kinase inhibitors", *Pest Management Science*, 2005, vol. 61, pp. 1069-1076.
- Pisella, et al., Flow cytometric analysis of conjunctival epithelium in ocular rosacea and keratoconjunctivitis sicca. *Ophthalmology*, 2000, vol. 107, pp. 1841-1849.
- Pisella, et al., Conjunctival proinflammatory and proapoptotic effects of latanoprost, preserved timolol and unpreserved timolol: an ex vivo in vitro study. *Invest Ophthalmol Vis Sci*, 2004, vol. 45, pp. 1360-1368.
- Portnaya, et al., "Azomethine dyes. IV. Indoaniline dyes derived from heterocyclic N-substituted 1-hydroxy-2-naphthamid", *Ts Vses Nauchn Issled Kinofotoinst*, 1960, No. 40, pp. 106-108 (with English abstract 20 pages total).
- Prezent, et al., "Boron chelates as intermediates in the synthesis of new functionalized pyridines and pyrimidines from a, a-dioxoketene aminals", *Proceedings of the International Conference on the Chemistry of Boron*, 2003, vol. 11 (abstract only—1 page).
- Quintas-Cardama, et al., "Preclinical characterization of the selective JAK1/2 inhibitor INCB018424: therapeutic implications for the treatment of myeloproliferative neoplasms", *Blood First Edition Paper*, pre-published online Feb. 3, 2010, American Society of Hematology.
- Ren et al., "Compounds and Compositions as Protein Kinase Inhibitors," U.S. Appl. No. 60/578,491, filed Jun. 10, 2004 (56 pages).
- Response and Amendment dated Aug. 25, 2009 to non-final Office Action for U.S. Appl. No. 12/137,892 (34 pgs.).
- Response and Amendment in Reply to Action of Apr. 20, 2007 filed Jul. 17, 2007 for U.S. Appl. No. 11/313,394 (39 pages).
- Response to Action of Aug. 22, 2007 dated Nov. 19, 2007, U.S. Appl. No. 11/115,702 (7 pages).
- Response to Restriction Requirement dated May 29, 2007, U.S. Appl. No. 11/115,702 (8 pages).
- Restriction Requirement dated Mar. 6, 2007 in connection with U.S. Appl. No. 11/115,702 (8 pages).
- Roberts, et al., "Trends in the Risks and Benefits to Patients with Cancer Participating in Phase I Clinical Trials", *JAMA* 2004, vol. 292, No. 17, pp. 2130-2140.
- Robin et al., In vivo transillumination biomicroscopy and photography of meibomian gland dysfunction. *Ophthalmology*, 1985, vol. 92, pp. 1423-1426.
- Rolando et al., "Tear mucus crystallization in children with cystic fibrosis", *Ophthalmologica*, 1988, vol. 197, No. 4, pp. 202-206.
- Rolando et al., "Tear mucus ferning test in keratoconjunctivitis sicca", Holly FJ, Lamberts DW, MacKeen DL (eds.): *The precocular tear film in health, disease, and contact lens wear*, 1st Intern Tear Film Symposium. Lubbock (Texas, USA), Dry Eye Institute, 1986, pp. 203-210.
- Rolando et al., "The effect of hyperosmolarity on tear mucus ferning", *Fortschr Ophthalmol*, 1986, vol. 83, pp. 644-646.
- Rolando, et al., *The Ocular Surface and Tear Film and Their Dysfunction in Dry Eye Disease*, Survey of Ophthalmology, 2001, vol. 45, Suppl. 2, pp. S203-S210.
- Rolando, "Tear mucus ferning test in normal and keratoconjunctivitis sicca eyes." *Chibret Int J Ophthalmol*, 1984, vol. 2, No. 4, pp. 32-41.
- Saettone, et al., "Ocular inserts for topical delivery," *Advanced Drug Delivery Reviews*, 1995, vol. 16, pp. 95-106.
- Samanta, et al., "Janus kinase 2: a critical target in chronic myelogenous leukemia", *Cancer Res.* 2006, vol. 66, No. 13, pp. 6468-6472.
- Sawadam et al., "Increased Lipophilicity and Subsequent Cell Partitioning Decrease Passive Transcellular Diffusion of Novel, Highly Lipophilic Antioxidants", *The Journal of Pharmacology and Experimental Therapeutics*, 1999, vol. 3, No. 288, pp. 1317-1326, p. 1321, compound 26.
- Schrader, et al., "Animal Models of Dry Eye," *Developmental Ophthalmology*, Karger 2008, vol. 41, pp. 298-312.
- Seefeld, et al., "Discovery of 5-pyrrolopyridinyl-2-thiophenecarboxamides as potent AKT kinase", *Bioorganic & Medicinal Chemistry Letters*, 2009, vol. 19, No. 8, pp. 2244-2248.
- Seela, et al., "Synthesis of Pyrrolo[2,3-d]pyrimidine 2', 3'-Dideoxyribonucleosides Related to 2',3'-Dideoxyadenosine and 2',3'-Dideoxyguanosine and Inhibitory Activity of 5'-Triphosphates on HIV-1 Reverse Transcriptase", *Helvetica Chimica, Acta*, 1991, vol. 74, No. 3, pp. 554-564.
- Seki, "STAT3 and MAPK in human lung cancer tissues and suppression of oncogenic growth by JAB and dominant negative STAT3", *Int J Oncol*. 2004, vol. 24, No. 4, pp. 931-934.
- Shi, et al., "The pharmacokinetics, pharmacodynamics, and safety of orally dosed INCB018424 phosphate in healthy volunteers", *Journal of Clinical Pharmacology*, 2011, vol. 51, No. 12, pp. 1644-1654.
- Shimazaki et al., "Meibomian gland dysfunction in patients with Sjogren syndrome", *Ophthalmology*, 1998, vol. 105, No. 8, pp. 1485-1488.
- Smith et al., "Basic pathogenic mechanisms operating in experimental model acute anterior uveitis," *Immunology and Cell Biology*, 1998, vol. 76, pp. 497-512.
- State Intellectual Property Office, PR China, Office Action, dated Sep. 3, 2010 Pat. Appl. No. 200680052750.7 (8 pages).
- Stirewalt et al., "Predictors of relapse and overall survival in Philadelphia chromosome-positive acute lymphoblastic leukemia after transplantation", *Biol Blood Marrow Transplant*. 2003, vol. 9, No. 3, pp. 206-212.
- STN Search conducted Aug. 30, 2010 (17 pages).
- STN Search conducted Jun. 24, 2011 (24 pages).
- 26th Annual JPMorgan Healthcare Conference presentation dated Jan. 8, 2008 (28 pages).
- Abstract of Chilean patent application No. 3496-06 published in Official Gazette of the Republic of Chile (Jun. 1, 2007) and publication (2 pages).
- Anderson et al., "Biochemical characterization of GSK1070916, a potent and selective inhibitor of Aurora B and Aurora C kinases with an extremely long residence time", *Biochem. J.*, 2009, vol. 420, No. 2, pp. 259-265.
- Baudouin et al., "Flow cytometry in impression cytology specimens. A new method for evaluation of conjunctival inflammation", *Invest Ophthalmol Vis Sci*, 1997, vol. 38, pp. 1458-1464.
- Bell, Malcolm, and Zalay, Andrew, "Synthesis of Substituted 3-Amino[6, 5-b] triazinoindoles." *Journal of Heterocyclic Chemistry*, 1975, vol. 12, No. 5, pp. 1001-1004.
- Berge, et al., "Pharmaceutical salts", *J. Pharma. Science*, 1977, vol. 66, No. 1, pp. 1-19.
- Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.
- Blume-Jensen, et al., "Oncogenic kinase signaling", *Nature* 2001, vol. 411, No. 6835, pp. 355-365.
- Bolen, "Nonreceptor tyrosine protein kinases", *Oncogene*, 1993, vol. 8, No. 8, pp. 2025-2031.
- Bollrath et al., "gp130-Mediated Stat3 Activation in Enterocytes Regulates Cell Survival and Cell-Cycle Progression during Colitis-Associated Tumorigenesis," *Cancer Cell*, 2009, vol. 15, pp. 91-102.

(56)

## References Cited

## OTHER PUBLICATIONS

- Borie, et al., "Combined Use of the Jak3 Inhibitor CP-690, 550 with Mycophenolate Mofetil to Prevent Kidney Allograft Rejection in Nonhuman Primates", *Transplantation*, 2005, vol. 80, No. 12, pp. 1756-1764.
- Boudny, et al., "JAK/STAT signaling pathways and cancer", *Neoplasia*, 2002, vol. 49, pp. 349-355.
- Bowman, et al., "STATs in oncogenesis", *Oncogene*, 2000, vol. 19, pp. 2474-2488.
- Bromberg et al., "Inflammation and Cancer: IL-6 and STAT3 Complete the Link," *Cancer Cell*, 2009, vol. 15, pp. 79-80.
- Burger, et al., "Gp130 and ras mediated signaling in human plasma cell line IN/a-6: a cytokine-regulated tumor model for plasmacytoma", *Hematol J.*, 2001, vol. 2, pp. 42-53.
- Candotti, et al., "Molecular aspects of primary immuno-deficiencies: lessons from cytokine and other signaling pathways.", *J Clin Invest*, 2002, Vol. 109, No. 10, pp. 1261-1269.
- Candotti, et al., "Structural and functional basis for JAK3-deficient severe combined immunodeficiency.", *Blood*, 1997, vol. 90, No. 10, pp. 3996-4003.
- Carey and Sundberg, *Advanced Organic Chemistry, Part B: Reactions and Synthesis, Oxidations*, 4th ed., Kluwer Academic/Plenum Publishers: New York, 2001, pp. 747-757.
- Cetkovic-Cvrlje, et al., "Targeting JAK3 with JANEX-1 for prevention of autoimmune type 1 diabetes in NOD mice.", *Clin Immunol*, 2003, vol. 106, No. 3, pp. 213-225.
- Chalandon, "Targeting mutated protein tyrosine kinases and their signaling pathways in hematologic malignancies", *Haematologica*, 2005, vol. 90, No. 7, pp. 949-968.
- Changelian, et al., "Prevention of Organ Allograft Rejection by a Specific Janus Kinase 3 Inhibitor", *Science*, 2003, vol. 302, pp. 875-878.
- Chen, et al., "Stat3 Activation in Human Endometrial and Cervical Cancer", *British Journal of Cancer*, 2007, vol. 96, pp. 591-599.
- Conklyn, M. et al., "The JAK3 inhibitor CP0690550 selectively reduces NK and CD8+ cell numbers in cynomolgus monkey blood following chronic oral dosing", *Journal of Leukocyte Biology*, 2004, vol. 76, pp. 1248-1255.
- Current Protocols in Immunology*, vol. 3., Coligan, J.E. et al, Wiley Press (1988).
- Daniels et al., "Imatinib mesylate inhibits the profibrogenic activity of TGF- $\beta$  and prevents bleomycin-mediated lung fibrosis." *J. Clin. Invest.*, 2004, vol. 114, No. 9, pp. 1308-1316.
- De Vos, et al., "JAK2 tyrosine kinase inhibitor tyrphostin AG490 downregulates the mitogen-activated protein kinase (MAPK) and signal transducer and activator of transcription (STAT) pathways and induces apoptosis in myeloma cells.", *Br J Haematol*, 2000, vol. 109, No. 4, pp. 823-828.
- Deuse, et al., "Novel Immunosuppression: R348, a JAK3- and Syk-Inhibitor Attenuates Acute Cardiac Allograft Rejection", *Transplantation*, 2008, vol. 85, No. 6, pp. 885-892.
- Doleschall, et al., "Thermal and Acid Catalysed Degradations of 3-alkylthio-6,7-dihydro[1,2,4]triazino[1,6-c] quinazolin-5-ium-1-olates", *Tetrahedron*, 1974, vol. 30, pp. 3997-4012.
- Dudley, et al., "AVEGF/JAK2/STAT5 axis may partially mediate endothelial cell tolerance to hypoxia", *Biochem. J.* 2005, vol. 390(Pt 2), pp. 427-436.
- Fridman, J. et al., "Selective JAK Inhibition is Efficacious against Multiple Myeloma Cells and Reverses the Protective Effects of Cytokine and Stromal Cell Support" Abstract #0956, presented Sunday, Jun. 15, 2008 at the European Hematology Association, 13th Congress, Jun. 12-15, Copenhagen, Denmark (1 page).
- Fridman, Jordan et al., "Discovery and Preclinical Characterization of INCB018424, a Selective JAK2 Inhibitor for the Treatment of Myeloproliferative Disorders" poster presented at the American Society of Hematology, 49th Annual Meeting and Exposition, GA. Abstract #3538, poster #757, Dec. 10, 2007 (1 page).
- Fridman, Jordan et al., "Efficacy and Tolerability of Novel JAK Inhibitors in Animal Models of Rheumatoid Arthritis" poster presented at the ACR/ARHP (American College of Rheumatology/Association of Rheumatology Health Professionals) Scientific Meeting 2007, Boston, MA. Nov. 10, 2007. Abstract 1771, poster 285 (1 page).
- Fridman, Jordan, et al., "Discovery and Preclinical Development of Selective JAK Inhibitors for the Treatment of Hematological Malignancies" poster presented at European Hematology Association, 12th Congress, Vienna, Austria. Abstract 0324, Jun. 8, 2007 (1 page).
- Fridman, Jordan, et al., "Discovery and Preclinical Development of Selective JAK Inhibitors for the Treatment of Myeloproliferative Disorders" poster presented at the 4th International Congress on Myeloproliferative Diseases and Myelodysplastic Syndromes, New York, NY. Nov. 8-10, 2007. Poster 0009 (1 page).
- Gaertner, "Cyclization of 1-Alkylamino-3-halo-2-alkanols to 1-Alkyl-3-azetidinols," *J. Org. Chem.*, 1967, vol. 32, pp. 2972-2976.
- Gooseman, et al., "The intramolecular b-fluorine . . . ammonium interaction in 4- and 8-membered rings", *Chem. Commun*, 2006, vol. 30, pp. 3190-3192.
- Gorre, M.E. et al., "Clinical Resistance to STI-571 Cancer Therapy Caused by BCR-ABL Gene Mutation or Amplification." *Science*, 2001, vol. 293, pp. 876.
- Gotlieb, Alice, Presentation at the 2008 American Academy of Dermatology, 66th Annual Meeting, San Antonio, TX. Feb. 1, 2008, symposium-303 (12 pp.).
- Gottlieb, A.B., et al., "Psoriasis: Emerging Therapeutic Strategies", *Nat Rev Drug Disc.*, 2005, vol. 4, pp. 19-34.
- Grabbe, et al., "Immunoregulatory mechanisms involved in elicitation of allergic-contact hypersensitivity", *Immunol Today*, 1998, vol. 19, No. 1, pp. 37-44 (only 1 page provided and marked "best available copy").
- Green, T.W. and Wuts, P.G.M., *Protective Groups in Organic Synthesis*, 3rd. Ed., Wiley & Sons, Inc., New York (1999).
- Hamze' et al., "Synthesis of Various 3-Substituted 1,2,4-Oxadiazole-Containing Chiral  $\beta$ - and  $\alpha$ -Amino Acids from Fmoc-Protected Aspartic Acid," *J. Org. Chem.*, 2003, vol. 68, No. 19, pp. 7316-7321.
- Hardwicke, et al., "GSK1070916, a potent Aurora B/C kinase inhibitor with broad antitumor activity in tissue culture cells and human tumor xenograft models", *Molecular Cancer Therapeutics* 2009, vol. 8, No. 7, pp. 1808-1817.
- Helal et al., "Stereoselective Synthesis of cis-1,3-Disubstituted Cyclobutyl Kinase Inhibitors," *Organic Letters*, 2004, vol. 6, No. 11, pp. 1853-1856.
- Higuchi, et al., "Pro-drugs as Novel Delivery Systems," vol. 14 of the A.C.S. Symposium Series (1975).
- International Preliminary Report on Patentability for International Appln. No. PCT/US2008/066662 dated Dec. 17, 2009 (7 pgs.).
- International Search Report and Written Opinion for International Appln. No. PCT/US2008/066662 dated Dec. 23, 2008 (11 pgs.).
- International Search Report and Written Opinion for PCT/US2006/047369, 16 pages (Apr. 24, 2007).
- Ishizaki, et al., "Pharmacological Properties of Y-27632, a Specific Inhibitor of Rho-Associated Kinases", *Molecular Pharmacology*, 2000, Vol. 57, pp. 976-983.
- Itagaki, et al., "Expedient Synthesis of Potent Cannabinoid Receptor Agonist (-)-CP55,940", *Organic Letters*, 2005, vol. 7, No. 19, pp. 4181-4183.
- Abe, et al., "Effective Methods for Introducing Some Aryl and Heteroaryl Substituent onto 1-Azaazulene Nuclei", *Heterocycles*, 2005, vol. 66, pp. 229-240.
- Abelson, et al., "Alternate reference values for tear film break-up time in normal and dry eye populations, Lacrimal Gland, Tear Film, and Dry Eye Syndromes 3 Part B", *Adv Exp Med Biol*, 2002, vol. 506, pp. 1121-1125.
- Abelson, et al., "Dry eye syndrome: diagnosis, clinical trials, and pharmaceutical treatment-improving clinical trials". *Lacrimal Gland, Tear Film, and Dry Eye Syndromes 3 Part B*, *Adv Exp Med Biol*, 2002, vol. 506, pp. 1079-1086.
- Aho, et al., Expression of human pim family genes is selectively up-regulated by cytokines promoting T helper type 1, but not T helper type 2, cell differentiation, *Immunology* 2005, vol. 116, pp. 82-88.
- Albach, et al., "Diagnosis of keratoconjunctivitis sicca in rheumatoid arthritis. The value of various tests", *Ophthalmologie*, 1994, vol. 91, No. 2, pp. 229-234—in German (with English abstract/summary contained therein).



(56)

## References Cited

## OTHER PUBLICATIONS

- Bachmann, et al., "The serine/threonine kinase Pim-1," *The International Journal of Biochemistry and Cell Biology* 2005, vol. 37, pp. 726-730.
- "Prodrugs" from Modern Pharmaceuticals, Banker and Rhodes, Ed.; Marcel Dekker, Inc.: New York, NY: pp. 596 (1996).
- Barabino, et al., "Tear film and ocular surface tests in animal models of dry eye; uses and limitations", *Experimental Eye Research*, 2004, vol. 79, pp. 613-621.
- Barr, et al., "Corneal scarring in the Collaborative Longitudinal Evaluation of Keratoconus (CLEK) Study: baseline prevalence and repeatability of detection", *Cornea*, 1999, vol. 18, No. 1, pp. 34-46.
- Baytel, et al., "The human Pim-2 proto-oncogene and its testicular expression" *Biochimica et Biophysica Acta* 1998, vol. 1442, pp. 274-285.
- Begley, et al., "Use of the dry eye questionnaire to measure symptoms of ocular irritation in patients with aqueous tear deficient dry eye", *Cornea*, 2002, vol. 21, pp. 664-670.
- Beyer, "Über die Synthese von 2-Methylmercapto-1,3,4-thiadiazinen und deren Umlagerung in Pyrazolderivate (The synthesis of 2-methylthio-1,3,4-thiadiazines and their rearrangement to pyrazole derivatives)", *Chem. Berichte Jahrg.*, 1959, vol. 92, pp. 2593-2599 (abstract provided).
- Bhovi, et al., "1,3-Dipolar Cycloaddition Reaction: Synthesis and Antimicrobial Activity of Some New 3-Ethoxycarbonyl-5-Methoxy-6-Bromo-2-Triazolylmethylindoles", *Indian Journal of Heterocyclic Chemistry*, 2004, vol. 14, pp. 15-18.
- Bosworth, JAK1/JAK2 Inhibitor Ruxolitinib Is a Rising Star, *Clinical Oncology*, 2011, vol. 06, No. 04 (3 pages).
- Bourcier, et al., "Expression of CD40 and CD40 ligand in the human conjunctival epithelium", *Invest Ophthalmol Vis Sci*, 2000, vol. 41, pp. 120-126.
- Brignole, et al., "Expression of Fas-Fas Ligand Antigens and Apoptotic Marker APO2-7 by the Human Conjunctival Epithelium. Positive correlation with class II HLA DR expression in inflammatory Ocular Surface Disorders", *Exp Eye Res*, 1998, vol. 67, pp. 687-697.
- Brignole, et al., "Flow cytometric analysis of inflammatory markers in conjunctival epithelial cells of patients with dry eyes", *Invest Ophthalmol Vis Sci*, 2000, vol. 41, pp. 1356-1363.
- Brignole, et al., "Flow cytometric analysis of inflammatory markers in KCS: 6-month treatment with topical cyclosporin A", *Invest Ophthalmol Vis Sci*, 2001, vol. 42, pp. 90-95.
- Brignole, et al., "Flow cytometry in conjunctival impression cytology: a new tool for exploring ocular surface pathologies", *Exp Eye Res*, 2004, vol. 78, pp. 473-481.
- Bron, et al., "Grading of corneal and conjunctival staining in the context of other dry eye tests", *Cornea*, 2003, vol. 22, No. 7, pp. 640-650.
- Bron, et al., "Methodologies to Diagnose and Monitor Dry Eye Disease: Report of the Diagnostic Methodology Subcommittee of the International Dry Eye Workshop (2007)", *The Ocular Surface*, 2007, vol. 5, No. 2, pp. 108-152.
- Burger, et al., "Janus kinase inhibitor INCB20 has antiproliferative and apoptotic effects on human myeloma cells in vitro and in vivo", *Mol. Cancer Ther.* 2009, vol. 8, No. 1, pp. 26-35.
- Campas-Moya, "Ruxolitinib. Tyrosine-protein kinase JAK1/2 inhibitor, treatment of myelofibrosis, treatment of myeloproliferative neoplasms, treatment of psoriasis", *Drugs of the Future*, 2010, vol. 35, No. 6, pp. 457-465.
- Carey and Sundberg, *Advanced Organic Chemistry, Part B: Reactions and Synthesis*, 4th ed., Kluwer Academic/Plenum Publishers: New York, pp. 111-119 (2001).
- Cermak, et al., "Is complete androgen insensitivity syndrome associated with alterations in the meibomian gland and ocular surface", *Cornea*, 2003, vol. 22, pp. 516-521.
- Chauhan, et al., "Autoimmunity in Dry Eye due to resistance of Th17 to Treg Suppression", *J. Immunology*, 2009, vol. 182, No. 3, pp. 1247-1252.
- Chew, et al., "An instrument for quantifying meibomian lipid on the lid margin: the Meibometer", *Curr Eye Res*, 1993a, vol. 12, pp. 247-254.
- Chew, et al., "The casual level of meibomian lipids in humans", *Current Eye Research*, 1993b, vol. 12, pp. 255-259.
- Cho, et al., "Review of the tear break-up time and a closer look at the tear break-up time of Hong Kong Chinese", *Optom Vis Sci*, 1993, vol. 70, No. 1, pp. 30-38.
- Choi Ha-Soon, et al., "Design and synthesis of 7H-pyrrolo[2,3-d]pyrimidines as focal adhesion kinase inhibitors. Part 1", *Bioorg. & Med. Chem. Lett.*, 2006, vol. 16, No. 8, pp. 2173-2176.
- Chu-Moyer, et al., "Preparation of the Four Regioisomeric 2-(Methylthio)oxazopyridines: Useful Synthons for Elaboration to 2-(Amino substituted)oxazopyridines", *J. Org. Chem.* 1995, vol. 60, No. 17, pp. 5721-5725.
- Cilloni, et al., "Emerging drugs for chronic myeloid leukemia", *Expert Opinion on Emerging Drugs*, 2010, vol. 15, No. 2, pp. 175-184.
- Coligan, et al, Wiley Press; *Methods in Molecular Biology*: vol. 225, *Inflammation Protocols.*, Winyard, P.G. and Willoughby, D.A., Humana Press (2003).
- Communication dated Jan. 22, 2009 for European Appln. No. 06839328.9 (5 pgs.).
- Craig, et al., "Tear lipid layer structure and stability following expression of the meibomian glands.", *Ophthalmic Physiol Opt*, 1995, vol. 15, No. 6, pp. 569-574.
- Danjo, et al., "Observation of precorneal tear film in patients with Sjogren's syndrome", *Acta Ophthalmol Scand*, 1995, vol. 73, pp. 501-505.
- De Paiva, et al., "IL-17 disrupts corneal barrier following desiccating stress", *Mucosal Immunol.* 2009, vol. 2, No. 3, pp. 243-253.
- Deng Jun, et al., "Rh-catalyzed asymmetric hydrogenation of gamma-phthalimido-substituted esters: an efficient enantioselective synthesis of beta-aryl-gamma-amino acids", *Org. Lett.* 2007, vol. 9, No. 23, pp. 4825-4827.
- Doane, "An instrument for in vivo tear film interferometry", *Optom Vis Sci*, 1989, vol. 66, pp. 383-388.
- Eghtedar, "Phase II Study of the JAK2 Inhibitor, INCB018424, in Patients with Refractory Leukemias Including Post-Myeloproliferative Disorder Acute Myeloid Leukemia", *American Society of Hematology (ASH) annual meeting in Orlando, FL (Dec. 6, 2010), Abstract/poster 509.*
- Eimmahl, et al., "Therapeutic applications of viscous and injectable poly(ortho esters)", *Adv. Drug. Deliv. Rev.* 2001, vol. 53, pp. 45-73.
- Eliason, et al., "Staining of the conjunctiva and conjunctival tear film", *Br J Ophthalmol*, 1990, vol. 74, pp. 519-522.
- Expert Scientific Group on Phase One Clinical Trials Final Report, Nov. 30, 2006, pp. C1, C35-C38.
- Fabrizio, et al., "Ocular inserts for topical delivery", *Advanced Drug Delivery Reviews* 1998, vol. 16, pp. 95-106.
- Farrell, et al., "A classification for dry eyes following comparison of tear thinning time with Schirmer tear test", *Acta Ophthalmol (Copenh)*, 1992, vol. 70, No. 3, pp. 357-360.
- Farrell, et al., "A clinical procedure to predict the value of temporary occlusion therapy in keratoconjunctivitis sicca" *Ophthalmol Physiol Opt*, 2003, vol. 23, pp. 1-8.
- Farris, "Tear osmolarity—a new gold standard?" *Adv Exp Med Biol*, 1994, vol. 350, pp. 495-503.
- Flex, et al., "Somatically acquired JAK1 mutations in adult acute lymphoblastic leukemia", *J. Exp Med.* 2008, vol. 205, pp. 751-758.
- Fonseca, et al., "Interleukin-6 as a key player in systemic inflammation and joint destruction", *Autoimmunity Reviews*, 2009, vol. 8, pp. 538-542.
- Fridman, et al., "Preclinical evaluation of local JAK1 and JAK2 inhibition in cutaneous inflammation", *Journal of Investigative Dermatology*, 2011, vol. 131, No. 9, pp. 1838-1844.
- International Search Report for PCT/US2010/047252 mailed Nov. 17, 2010 (4 pages).
- International Search Report for PCT/US2010/052011 mailed Nov. 30, 2010 (3 pages).

(56)

## References Cited

## OTHER PUBLICATIONS

- Iranpoor, et al., "A Rapid and Facile Conversion of Primary Amides and Aldoximes to Nitriles and Ketoximes to Amides with Triphenylphosphine and N-Chlorosuccinimide", *G Syn. Commun* 2002, vol. 32, pp. 2535-2541.
- Janes, et al., "Effective and selective targeting of leukemia cells using a TORC1/2 kinase inhibitor.", *Nature Medicine* 2010, vol. 16, No. 2, pp. 205-213.
- Jee, et al., "Overview: animal models of osteopenia and osteoporosis", *J Musculoskel. Neuron, Interact.*, 2001, vol. 1, No. 3, pp. 193-207.
- Jester, et al., "In vivo biomicroscopy and photography of meibomian glands in a rabbit model of meibomian gland dysfunction", *Invest Ophthalmol Vis Sci*, 1982, vol. 22, pp. 660-667.
- Johnson, et al., "The effect of instilled fluorescein solution volume on the values and repeatability of TBUT measurements", *Cornea*, 2005, vol. 24, pp. 811-817.
- Kaercher, T., "Ocular symptoms and signs in patients with ectodermal dysplasia syndromes", *Grafes Arch Clin Exp Ophthalmol*, 2004, pp. 495-500.
- Kamb, *Nature Reviews Drug Discovery* 2005, vol. 4, pp. 161-165.
- Kaushansky, K., "Lineage-Specific Hematopoietic Growth Factors", *NEJM* 2006, vol. 354, pp. 2034-2045.
- Kim, et al., "Zinc-Modified Cyanoborohydride as a Selective Reducing Agent", *J. Org. Chem.* 1985, vol. 50, pp. 1927-1932.
- King-Smith et al., "Three interferometric methods for measuring the thickness of layers of the tear film", *Optom Vis Sci*, 1999, vol. 76, pp. 19-32.
- Kiss, Robert, "Recent developments on JAK2 inhibitors: A patent review", *Expert Opinion on Therapeutic Patents*, 2010, vol. 20, No. 4, pp. 471-495.
- Kojima et al., "A new noninvasive tear stability analysis system for the assessment of dry eyes", *Invest Ophthalmol Vis Sci*, 2004, vol. 45, No. 5, pp. 369-374.
- Kola, *Nature Reviews Drug Discovery* 2004, vol. 3, pp. 711-715.
- Komuro et al., "Assessment of meibomian gland function by a newly developed laser meibometer", *Adv Exp Med Biol*, 2002, vol. 506, pp. 517-520.
- Korb et al., "The effect of two novel lubricant eye drops on tear film lipid layer thickness in subjects with dry eye symptoms", *Optom Vis Sci*, 2005, vol. 82, pp. 594-601.
- Korb, et al., "Increase in tear film lipid layer thickness following treatment of meibomian gland dysfunction", *Adv Exp Med Biol*, 1994, vol. 350, pp. 293-298.
- Korolev, et al., "Pd-EDTA as an efficient catalyst for Suzuki-Miyaura reactions in water", *Tet. Lett.* 2004, vol. 46, pp. 5751-5754.
- Kortylewski, et al., "Regulation of the IL-23 and IL-12 balance by Stat3 signaling in the tumor microenvironment", *Cancer Cell*, 2009, Vol. 15, pp. 114-123.
- Kumar, , "Kinase drug discovery approaches in chronic myeloproliferative disorders", *Oncogene*, 2009, vol. 28, No. 24, pp. 2305-2323.
- Kuo, et al., "Pd-EDTA as an efficient catalyst for Suzuki-Miyaura reactions in water", *Chem Commun* 2007, pp. 301-303.
- Kuppens et al., "Basal tear turnover and topical timolol in glaucoma patients and healthy controls by Fluorophotometry", *Invest Ophthalmol Vis Sci*, 1992, vol. 33, pp. 3442-3448.
- Lai, et al., "Mechanistic Study on the Inactivation of General Acyl-CoA Dehydrogenase by a Metabolite of Hypoglycin A", *J. Am. Chem. Soc.* 1991, vol. 113, pp. 7388-7397.
- Lam, et al, "Tear Cytokine Profiles in Dysfunctional Tear Syndrome", *Am J Ophthalmol.*, 2009, vol. 147, No. 2, pp. 198-205.
- Larock, R., "Comprehensive Organic Transformations", Wiley-VCH, 2nd Ed. (1999) pp. 1949-1950, 1958-1959, 1976, and 1983-1985.
- Leaf, *Health Administrator* 2005, vol. XVII, No. 1, pp. 172-183.
- Lemp, "Report of National Eye Institute/Industry Workshop on clinical trials in dry eyes", *CLAO J*, 1995, vol. 21, pp. 221-232.
- Lemp, et al., "Corneal desiccation despite normal tear volume", *Ann Ophthalmol*, 1970, No. 2, pp. 258-261 & 284.
- Lemp, et al., "The Definition and Classification of Dry Eye Disease: Report of the Definition and Classification Subcommittee of the International Dry Eye WorkShop", *The Ocular Surface*, 2007, vol. 5, No. 2, pp. 75-92.
- Levitzi, "Tyrosine kinases as targets for cancer therapy", *Eur. J. Cancer* 2002, vol. 38, Suppl. 5, pp. S11-S18.
- Levy, et al. "INCB018424 A Selective Janus Kinase 1/2 Inhibitor" Presentation at the 50th American Society of Hematology Annual Meeting (ASH), Dec. 8, 2008.
- Li, et al., "Pim-3, a proto-oncogene with serine/threonine kinase activity, is aberrantly expressed in human pancreatic cancer and phosphorylates Bad-mediated apoptosis in human pancreatic cell lines" *Cancer Research* 2006, vol. 66, No. 13, pp. 6741-6747.
- Lin, "Constitutive Activation of JAK3/STAT3 in Colon Carcinoma Tumors and Cell Lines", *Am J Pathol.* 2005, vol. 167, No. 4, pp. 969-980.
- Lin, et al., "Enantioselective synthesis of Janus kinase inhibitor INCB018424 via an organocatalytic aza-Michael reaction," *Organic Letters*, 2009, vol. 11, No. 9, pp. 1999-2002.
- Liu, et al., "Combined Inhibition of Janus Kinase 1/2 for the Treatment of JAK2V617F-Driven Neoplasms: Selective Effects on Mutant Cells and Improvements in Measures of Disease Severity", *Clin. Cancer Res.* 2009, vol. 15, No. 22, pp. 6891-6900.
- Macchi, et al., "Mutations of Jak-3 gene in patients with autosomal severe combined immune deficiency (SCID)", *Nature* 1995, vol. 377, pp. 65-68.
- Madden, et al. Comparative study of two non-invasive tear film stability techniques. *Curr Eye Res*, 1994, vol. 13, No. 4, pp. 263-269.
- Maffioli, et al., "Mild and Reversible Dehydration of Primary Amides with PdC12 in Aqueous Acetonitrile", *Org. Lett.*, 2005, vol. 7, No. 23, pp. 5237-5239.
- Main, et al., "High throughput synthesis of diverse 2,5-disubstituted indoles using titanium carbenoids bearing boronate functionality", *Tetrahedron*, 2007, vol. 64, No. 5, pp. 901-914.
- Mainstone, et al., "Tear meniscus measurement in the diagnosis of dry eye", *Curr Eye Res*, 1996, vol. 15, pp. 653-661.
- Smolen, et al, "Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (Option study): a double-blind, placebo-controlled, randomized trial", *Lancet* 2008, vol. 371, pp. 987, 2008.
- Mancini, et al., "RAD 001 (everolimus) prevents mTOR and Akt late re-activation in response to imatinib in chronic myeloid leukemia.", *J. Cellular Biochemistry*, 2010, vol. 109, No. 2, pp. 320-328.
- March, Jerry, *Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 3rd ed., John Wiley & Sons: New York, pp. 845-855 (1985).
- Marquardt et al., "Modification of tear film break-up time test for increased reliability" in Holly ed. *The Preocular Tear Film in Health, Disease and Contact Lens Wear*. Lubbock, Texas: Dry Eye Institute, 1986, pp. 57-63.
- Mariyama, et al., "Effect of environmental conditions on tear dynamics in soft contact lens wearers", *Invest Ophthalmol Vis Sci*, 2004, vol. 45, No. 8, pp. 2563-2568.
- Mathers, et al., "Assessment of the tear film with tandem scanning confocal microscopy", *Cornea*, 1997, vol. 16, pp. 162-168.
- Mathers, et al., "Tear film changes associated with normal aging", *Cornea*, 1996, vol. 15, pp. 229-334.
- Mathers, et al., "Tear flow and evaporation in patients with and without dry eye", *Ophthalmology*, 1996, vol. 103, pp. 664-669.
- Mathers, et al., "Video imaging of the meibomian gland", *Arch Ophthalmol*, 1994, vol. 112, pp. 448-449.
- Fujiwara, et al., "Evaluation of human conjunctival epithelium by a combination of brush cytology and flow cytometry: an approach to the quantitative technique", *Diagn Cytopathol*, 1997, vol. 17, pp. 456-460.
- Fujii, et al., "Aberrant expression of serine/threonine kinase Pim-3 in hepatocellular carcinoma development and its role in the proliferation of human hepatoma cell lines" *International Journal of Cancer* 2005, vol. 114, pp. 209-218.
- Fukagawa, et al., "Histological evaluation of brush cytology of rabbit conjunctiva", *Nippon Ganka Gakkai Zasshi*, 1993, vol. 97, pp. 1173-1178 (contains English abstract within the article).

(56)

**References Cited****OTHER PUBLICATIONS**

- Ghelardi, et al., "A Mucoadhesive Polymer Extracted from Tamarind Seed Improves the Intraocular Penetration and Efficacy of Rifaximin in Topical Treatment of Experimental Bacterial Keratitis", *Antimicrob. Agents Chemother.* 2004, vol. 48, pp. 3396-3401.
- Glasson, et al., "Differences in clinical parameters and tear film of tolerant and intolerant contact lens wearers", *Invest Ophthalmol Vis Sci.* 2003, vol. 44, pp. 5116-5124.
- Glattfeld, "Improvements in the Preparation of DL-Threonic and DL-Erythronic Acids", *J. Am. Chem. Soc.* 1940, vol. 62, pp. 974-977.
- Gobbels, et al., "Tear secretion in dry eyes as assessed by objective fluorophotometry", *Ger J Ophthalmol.* 1992, vol. 1, pp. 350-353.
- Golding, et al., "X-ray and scanning electron microscopic analysis of the structural composition of tear ferns", *Cornea* 1994, vol. 13, No. 1, pp. 58-66.
- Gomtsyan, et al., "Design, synthesis, and structure-activity relationship of 6-alkynylpyrimidines as potent adenosine kinase inhibitors", *J. Med. Chem.* 2002, vol. 45, No. 17, pp. 3639-3648.
- Goto, et al. Color mapping of tear lipid layer thickness distribution from the image analysis in DR-1 tear lipid layer interference images (ARVO abstract). ARVO 2004.
- Goto, et al., "Computer-synthesis of an interference color chart of human tear lipid layer by a colorimetric approach", *Invest Ophthalmol Vis Sci.* 2003, vol. 44, pp. 4693-4697.
- Goto, et al., "Differentiation of lipid tear deficiency dry eye by kinetic analysis of tear interference images", *Arch Ophthalmol.* 2003, vol. 121, pp. 173-180.
- Goto et al., "Evaluation of the tear film stability after laser in situ keratomileusis using the tear film stability analysis system", *Am J Ophthalmol.* 2004b, vol. 137, No. 1, pp. 116-120.
- Goto, et al., "Tear Film Stability Analysis System: Introducing a new application for videokeratography", *Cornea*, 2004a, vol. 3, No. 8, pp. S65-S70.
- Goto, et al., Kinetic analysis of tear interference images in aqueous tear deficiency dry eye before and after punctal occlusion. *Invest Ophthalmol Vis Sci.* 2003, vol. 44, pp. 1897-1905.
- Gregory, et al., "Clinical and laboratory features of myelofibrosis and limitations of current therapies", *Clinical Advances in Hematology and Oncology*, 2011, vol. 9, No. 9, pp. 1-3.
- Grivennikov, et al., "IL-6 and STAT3 are required for survival of intestinal epithelial cells and the development of colitis-associated cancer", *Cancer Cell*, 2009, vol. 15, pp. 103-111.
- Groneberg, et al., "Animal models of allergic and inflammatory conjunctivitis," *Allergy*, 2003, vol. 58, pp. 1101-1113.
- Guillon, "Tear film photography and contact lens wear", *J Br Contact Lens Assoc.* 1982, vol. 5, pp. 84-87.
- Gura, "Systems for identifying new drugs are often faulty", *Science*, 1997, vol. 278, No. 5340, pp. 1041-1042.
- Guschin, et al., "A major role for the protein tyrosine kinase JAK1 in the JAK/STAT signal transduction pathway in response to interleukin-6", *Embo J* 1995, vol. 14, pp. 1421-1429.
- Holly et al., "Lacrimation kinetics in Humans as determined by a novel technique", in Holly FJ (ed). *The precocular tear film*. Lubbock TX, Lubbock Dry Eye Institute, 1986, pp. 76-88.
- Hong, et al., "Total Synthesis of Onnamide A", *J. Am. Chem. Soc.*, 1991, vol. 113, pp. 9693-9694.
- Huang, "Inhibition of STAT3 activity with AG490 decreases the invasion of human pancreatic cancer cells in vitro", *Cancer Sci.* 2006, vol. 97, No. 12, pp. 1417-1423.
- Huttel, et al., "Lithium pyrazole compounds", *Liebigs Ann. Chem. Bd.*, 1959, vol. 625, pp. 55-65 (abstract provided).
- International Search Report for PCT/US2010/035783 mailed Aug. 23, 2010 (4 pages).
- International Preliminary Report on Patentability (with Written Opinion) dated Mar. 6, 2012 for International Appln. No. PCT/US2010/047252 (7 pgs.).
- International Preliminary Report on Patentability (with Written Opinion) dated Nov. 22, 2011 for International Appln. No. PCT/US2010/035728 (8 pgs.).
- International Preliminary Report on Patentability (with Written Opinion) dated Nov. 22, 2011 for International Appln. No. PCT/US2010/035783 (5 pgs.).
- International Preliminary Report on Patentability for PCT/US2008/66658 mailed Dec. 17, 2009 (7 pages).
- International Preliminary Report on Patentability for PCT/US2009/036635 mailed Sep. 14, 2010 (6 pages).
- International Preliminary Report on Patentability for PCT/US2009/059203 mailed Apr. 5, 2011 (6 pages).
- International Preliminary Report on Patentability for PCT/US2010/021003 mailed Jul. 19, 2011 (11 pages).
- International Preliminary Report on Patentability for PCT/US2010/052011 mailed Apr. 11, 2012 (4 pages).
- International Preliminary Report on Patentability for PCT/US2011/025433 mailed Aug. 21, 2012 (7 pages).
- International Preliminary Report on Patentability for PCT/US2011/027665 mailed Sep. 11, 2012 (7 pages).
- International Search Report and Written Opinion dated Feb. 9, 2010 for International Appln. No. PCT/US2009/059203 (10 pages).
- International Search Report and Written Opinion for International Appln. No. PCT/US2005/046207 dated May 15, 2007 (6 pages).
- International Search Report and Written Opinion for International Appln. No. PCT/US2009/036635 dated Jun. 3, 2009 14 pages.
- International Search Report and Written Opinion for PCT/US2008/083319, 29 pages mailed Mar. 13, 2009.
- International Search Report and Written Opinion for PCT/US2011/025433, 12 pages (mailed Jul. 20, 2011).
- International Search Report and Written Opinion for PCT/US2011/027665 mailed Jun. 27, 2011 (14 pages).
- International Search Report and Written Opinion for PCT/US2011/037291, 11 pages (Apr. 19, 2012).
- International Search Report and Written Opinion for PCT/US2011/061351 mailed Feb. 17, 2012 (12 pages).
- International Search Report and Written Opinion for PCT/US2011/061374 mailed Mar. 27, 2012 (10 pages).
- International Search Report and Written Opinion for PCT/US2012/025581, 16 pages (mailed Apr. 26, 2011).
- International Search Report for PCT/US2008/66658 mailed Dec. 23, 2008 (4 pages).
- International Search Report for PCT/US2010/021003 mailed Aug. 16, 2010 (8 pages).
- International Search Report for PCT/US2010/035728 mailed Jul. 8, 2010 (3 pages).

1

**SALTS OF THE JANUS KINASE INHIBITOR  
(R)-3-(4-(7H-PYRROLO[2,3-D]PYRIMIDIN-  
4-YL)-1H-PYRAZOL-1-YL)-3-  
CYCLOPENTYLPROPANENITRILE**

**CROSS-REFERENCE TO RELATED  
APPLICATIONS**

This application is a continuation of U.S. application Ser. No. 14/097,588, filed on Dec. 5, 2013, which is a continuation of U.S. application Ser. No. 12/137,892, filed Jun. 12, 2008, which claims the benefit of U.S. Ser. No. 60/943,705, filed Jun. 13, 2007, the disclosure of each of which is incorporated herein by reference in their entireties.

**FIELD OF THE INVENTION**

The present invention provides salt forms of (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile that are useful in the modulation of Janus kinase activity and are useful in the treatment of diseases related to activity of Janus kinases including, for example, immune-related diseases, skin disorders, myeloid proliferative disorders, cancer, and other diseases.

**BACKGROUND OF THE INVENTION**


Protein kinases (PKs) are a group of enzymes that regulate diverse, important biological processes including cell growth, survival and differentiation, organ formation and morphogenesis, neovascularization, tissue repair and regeneration, among others. Protein kinases exert their physiological functions through catalyzing the phosphorylation of proteins (or substrates) and thereby modulating the cellular activities of the substrates in various biological contexts. In addition to the functions in normal tissues/organs, many protein kinases also play more specialized roles in a host of human diseases including cancer. A subset of protein kinases (also referred to as oncogenic protein kinases), when dysregulated, can cause tumor formation and growth, and further contribute to tumor maintenance and progression (Blume-Jensen P et al, Nature 2001, 411(6835):355-365). Thus far, oncogenic protein kinases represent one of the largest and most attractive groups of protein targets for cancer intervention and drug development.

The Janus Kinase (JAK) family plays a role in the cytokine-dependent regulation of proliferation and function of cells involved in immune response. Currently, there are four known mammalian JAK family members: JAK1 (also known as Janus kinase-1), JAK2 (also known as Janus kinase-2), JAK3 (also known as Janus kinase, leukocyte; JAKL; L-JAK and Janus kinase-3) and TYK2 (also known as protein-tyrosine kinase 2). The JAK proteins range in size from 120 to 140 kDa and comprise seven conserved JAK homology (JH) domains; one of these is a functional catalytic kinase domain, and another is a pseudokinase domain potentially serving a regulatory function and/or serving as a docking site for STATs (Scott, Godshall et al. 2002, supra).

Blocking signal transduction at the level of the JAK kinases holds promise for developing treatments for human cancers. Inhibition of the JAK kinases is also envisioned to have therapeutic benefits in patients suffering from skin immune disorders such as psoriasis, and skin sensitization. Accordingly, inhibitors of Janus kinases or related kinases are widely sought and several publications report effective classes of compounds. For example, certain JAK inhibitors, including (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-

2

yl)-3-cyclopentylpropanenitrile depicted below, are reported in U.S. Ser. No. 11/637,545, filed Dec. 12, 2006.



Thus, new or improved forms of existing Janus kinase inhibitors are continually needed for developing new, improved, and more effective pharmaceutical formulations for the treatment of cancer and other diseases. The salt forms and methods described herein are directed toward these needs and other ends.

**SUMMARY OF THE INVENTION**

The present invention provides, inter alia, salts selected from:

- (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile maleic acid salt;
- (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile sulfuric acid salt; and
- (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile phosphoric acid salt.

The present invention further provides methods of preparing a salt of the invention comprising combining (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile with maleic acid, sulfuric acid, or phosphoric acid.

The present invention further provides compositions comprising a salt form of the invention and at least one pharmaceutically acceptable carrier.

The present invention further provides methods of modulating an activity of JAK comprising contacting JAK with a salt of the invention.

The present invention further provides methods of treating a disease in a patient, wherein the disease is associated with JAK activity, comprising administering to the patient a therapeutically effective amount of a salt of the invention.

The present invention further provides methods of treating cancer, skin disorders, or inflammation in a patient, comprising administering to the patient a therapeutically effective amount of a salt of the invention.

**DETAILED DESCRIPTION**

The present invention provides, inter alia, salts of the JAK inhibitor (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile selected from the maleic acid salt, sulfuric acid salt, and phosphoric acid salt. These salts modulate the activity of one or more JAKs and are useful, for example, in the treatment of diseases associated with JAK expression or activity.

The salts of the invention have numerous advantageous properties over the free base form and other salt forms. In

particular, these salts were highly crystalline which would facilitate the preparation of pharmaceutical formulations and improve general handling, manipulation, and storage of the active ingredient. The salts of the invention also have superior aqueous solubility, rate of dissolution, chemical stability (with a longer shelf life), compatibility with excipients, and reproducibility compared with the free base form.

In some embodiments, the salts of the invention are substantially isolated. By "substantially isolated" is meant that the salt is at least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the salt of the invention. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the salt.

Salts of the invention also include all isotopes of atoms occurring in the salts. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include tritium and deuterium.

Salts of the invention can be prepared using known techniques. Conventionally, a salt form is prepared by combining in solution the free base compound and an acid containing the anion of the salt form desired, and then isolating the solid salt product from the reaction solution (e.g., by crystallization, precipitation, evaporation, etc.). Other salt-forming techniques can be employed.

#### Methods of Use

Salts of the invention can modulate activity of one or more Janus kinases (JAKs). The term "modulate" is meant to refer to an ability to increase or decrease the activity of one or more members of the JAK family of kinases. Accordingly, compounds of the invention can be used in methods of modulating a JAK by contacting the JAK with any one or more of the compounds or compositions described herein. In some embodiments, salts of the present invention can act as inhibitors of one or more JAKs. In some embodiments, compounds of the present invention can act to stimulate the activity of one or more JAKs. In further embodiments, the compounds of the invention can be used to modulate activity of a JAK in an individual in need of modulation of the receptor by administering a modulating amount of a salt of the invention.

JAKs to which the present salts bind and/or modulate include any member of the JAK family. In some embodiments, the JAK is JAK1, JAK2, JAK3 or TYK2. In some embodiments, the JAK is JAK1 or JAK2. In some embodiments, the JAK is JAK2. In some embodiments, the JAK is JAK3.

The salts of the invention can be selective. By "selective" is meant that the compound binds to or inhibits a JAK with greater affinity or potency, respectively, compared to at least one other JAK. In some embodiments, the compounds of the invention are selective inhibitors of JAK1 or JAK2 over JAK3 and/or TYK2. In some embodiments, the salts of the invention are selective inhibitors of JAK2 (e.g., over JAK1, JAK3 and TYK2). Without wishing to be bound by theory, because inhibitors of JAK3 can lead to immunosuppressive effects, a compound which is selective for JAK2 over JAK3 and which is useful in the treatment of cancer (such as multiple myeloma, for example) can offer the additional advantage of having fewer immunosuppressive side effects. Selectivity can be at least about 5-fold, 10-fold, at least about 20-fold, at least about 50-fold, at least about 100-fold, at least about 200-fold, at least about 500-fold or at least about 1000-fold. Selectivity can be measured by methods routine in the art. In some embodiments, selectivity can be tested at the  $K_m$  of each

enzyme. In some embodiments, selectivity of salts of the invention for JAK2 over JAK3 can be determined by the cellular ATP concentration.

Another aspect of the present invention pertains to methods of treating a JAK-associated disease or disorder in an individual (e.g., patient) by administering to the individual in need of such treatment a therapeutically effective amount or dose of a salt of the present invention or a pharmaceutical composition thereof. A JAK-associated disease can include any disease, disorder or condition that is directly or indirectly linked to expression or activity of the JAK, including overexpression and/or abnormal activity levels. A JAK-associated disease can also include any disease, disorder or condition that can be prevented, ameliorated, or cured by modulating JAK activity.

Examples of JAK-associated diseases include diseases involving the immune system including, for example, organ transplant rejection (e.g., allograft rejection and graft versus host disease).

Further examples of JAK-associated diseases include autoimmune diseases such as multiple sclerosis, rheumatoid arthritis, juvenile arthritis, type I diabetes, lupus, psoriasis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, myasthenia gravis, immunoglobulin nephropathies, autoimmune thyroid disorders, and the like. In some embodiments, the autoimmune disease is an autoimmune bullous skin disorder such as pemphigus vulgaris (PV) or bullous pemphigoid (BP).

Further examples of JAK-associated diseases include allergic conditions such as asthma, food allergies, atopic dermatitis and rhinitis. Further examples of JAK-associated diseases include viral diseases such as Epstein Barr Virus (EBV), Hepatitis B, Hepatitis C, HIV, HTLV 1, Varicella-Zoster Virus (VZV) and Human Papilloma Virus (HPV).

Further examples of JAK-associated diseases or conditions include skin disorders such as psoriasis (for example, psoriasis vulgaris), atopic dermatitis, skin rash, skin irritation, skin sensitization (e.g., contact dermatitis or allergic contact dermatitis). For example, certain substances including some pharmaceuticals when topically applied can cause skin sensitization. In some embodiments, co-administration or sequential administration of at least one JAK inhibitor of the invention together with the agent causing unwanted sensitization can be helpful in treating such unwanted sensitization or dermatitis. In some embodiments, the skin disorder is treated by topical administration of at least one JAK inhibitor of the invention.

In further embodiments, the JAK-associated disease is cancer including those characterized by solid tumors (e.g., prostate cancer, renal cancer, hepatic cancer, pancreatic cancer, gastric cancer, breast cancer, lung cancer, cancers of the head and neck, thyroid cancer, glioblastoma, Kaposi's sarcoma, Castleman's disease, melanoma etc.), hematological cancers (e.g., lymphoma, leukemia such as acute lymphoblastic leukemia, acute myelogenous leukemia (AML), or multiple myeloma), and skin cancer such as cutaneous T-cell lymphoma (CTCL) and cutaneous B-cell lymphoma. Example cutaneous T-cell lymphomas include Sezary syndrome and mycosis fungoides.

JAK-associated diseases can further include those characterized by expression of a mutant JAK2 such as those having at least one mutation in the pseudo-kinase domain (e.g., JAK2V617F).

JAK-associated diseases can further include myeloproliferative disorders (MPDs) such as polycythemia vera (PV), essential thrombocythemia (ET), myeloid metaplasia with myelofibrosis (MMM), chronic myelogenous leukemia

(CML), chronic myelomonocytic leukemia (CMML), hyper-eosinophilic syndrome (HES), systemic mast cell disease (SMCD), and the like.

Further JAK-associated diseases include inflammation and inflammatory diseases. Example inflammatory diseases include inflammatory diseases of the eye (e.g., iritis, uveitis, scleritis, conjunctivitis, or related disease), inflammatory diseases of the respiratory tract (e.g., the upper respiratory tract including the nose and sinuses such as rhinitis or sinusitis or the lower respiratory tract including bronchitis, chronic obstructive pulmonary disease, and the like), inflammatory myopathy such as myocarditis, and other inflammatory diseases. Other inflammatory diseases treatable by the compounds of the invention include systemic inflammatory response syndrome (SIRS) and septic shock.

The JAK inhibitors described herein can further be used to treat ischemia reperfusion injuries or a disease or condition related to an inflammatory ischemic event such as stroke or cardiac arrest. The JAK inhibitors described herein can further be used to treat anorexia, cachexia, or fatigue such as that resulting from or associated with cancer. The JAK inhibitors described herein can further be used to treat restenosis, sclerodermitis, or fibrosis. The JAK inhibitors described herein can further be used to treat conditions associated with hypoxia or astrogliosis such as, for example, diabetic retinopathy, cancer, or neurodegeneration. See, e.g., Dudley, A. C. et al. *Biochem. J.* 2005, 390(Pt 2):427-36 and Sriram, K. et al. *J. Biol. Chem.* 2004, 279(19):19936-47. Epub 2004 Mar. 2.

The JAK inhibitors described herein can further be used to treat gout and increased prostate size due to, e.g., benign prostatic hypertrophy or benign prostatic hyperplasia.

As used herein, the term "contacting" refers to the bringing together of indicated moieties in an in vitro system or an in vivo system. For example, "contacting" a JAK with a salt of the invention includes the administration of a salt of the present invention to an individual or patient, such as a human, having a JAK, as well as, for example, introducing a salt of the invention into a sample containing a cellular or purified preparation containing the JAK.

As used herein, the term "individual" or "patient," used interchangeably, refers to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

As used herein, the phrase "therapeutically effective amount" refers to the amount of active salt or pharmaceutical agent that elicits the biological or medicinal response that is being sought in a tissue, system, animal, individual or human by a researcher, veterinarian, medical doctor or other clinician.

As used herein, the term "treating" or "treatment" refers to one or more of (1) preventing the disease; for example, preventing a disease, condition or disorder in an individual who may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease; (2) inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder; and (3) ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology) such as decreasing the severity of disease. Combination Therapies

One or more additional pharmaceutical agents such as, for example, chemotherapeutics, anti-inflammatory agents, ste-

roids, immunosuppressants, as well as Bcr-Abl, Flt-3, RAF and FAK kinase inhibitors such as, for example, those described in WO 2006/056399, or other agents can be used in combination with the salts of the present invention for treatment of JAK-associated diseases, disorders or conditions. The one or more additional pharmaceutical agents can be administered to a patient simultaneously or sequentially.

Example chemotherapeutics include proteasome inhibitors (e.g., bortezomib), thalidomide, revlimid, and DNA-damaging agents such as melphalan, doxorubicin, cyclophosphamide, vincristine, etoposide, carmustine, and the like.

Example steroids include corticosteroids such as dexamethasone or prednisone.

Example Bcr-Abl inhibitors include the compounds, and pharmaceutically acceptable salts thereof, of the genera and species disclosed in U.S. Pat. No. 5,521,184, WO 04/005281, EP2005/009967, EP2005/010408, and U.S. Ser. No. 60/578,491.

Example suitable Flt-3 inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 03/037347, WO 03/099771, and WO 04/046120.

Example suitable RAF inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 00/09495 and WO 05/028444.

Example suitable FAK inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 04/080980, WO 04/056786, WO 03/024967, WO 01/064655, WO 00/053595, and WO 01/014402.

In some embodiments, the salt forms of the invention can be used in combination with other kinase inhibitors such as imatinib, particularly for the treatment of patients resistant to imatinib or other kinases.

In some embodiments, one or more salt forms of the invention can be used in combination with a chemotherapeutic in the treatment of cancer, such as multiple myeloma, and may improve the treatment response as compared to the response to the chemotherapeutic agent alone, without exacerbation of its toxic effects. Examples of additional pharmaceutical agents used in the treatment of multiple myeloma, for example, can include, without limitation, melphalan, melphalan plus prednisone [MP], doxorubicin, dexamethasone, and Velcade (bortezomib). Further additional agents used in the treatment of multiple myeloma include Bcr-Abl, Flt-3, RAF and FAK kinase inhibitors. Additive or synergistic effects are desirable outcomes of combining a JAK inhibitor of the present invention with an additional agent. Furthermore, resistance of multiple myeloma cells to agents such as dexamethasone may be reversible upon treatment with a JAK inhibitor of the present invention. The agents can be combined with the present compounds in a single or continuous dosage form, or the agents can be administered simultaneously or sequentially as separate dosage forms.

In some embodiments, a corticosteroid such as dexamethasone is administered to a patient in combination with at least one JAK inhibitor where the dexamethasone is administered intermittently as opposed to continuously.

In some further embodiments, combinations of one or more JAK inhibitors of the invention with other therapeutic agents can be administered to a patient prior to, during, and/or after a bone marrow transplant or stem cell transplant.

Pharmaceutical Formulations and Dosage Forms

When employed as pharmaceuticals, the salts of the invention can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated.

Administration may be topical (including transdermal, epidermal, ophthalmic and to mucous membranes including intranasal, vaginal and rectal delivery), pulmonary (e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal, intramuscular or injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Parenteral administration can be in the form of a single bolus dose, or may be, for example, by a continuous perfusion pump. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

This invention also includes pharmaceutical compositions which contain, as the active ingredient, one or more of the compounds of the invention above in combination with one or more pharmaceutically acceptable carriers (excipients). In making the compositions of the invention, the active ingredient is typically mixed with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, the active compound can be milled to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it can be milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size can be adjusted by milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

The compounds of the invention may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other formulation types. Finely divided (nanoparticulate) preparations of the compounds of the invention can be prepared by processes known in the art, for example see International Patent Application No. WO 2002/000196.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

The compositions can be formulated in a unit dosage form, each dosage containing from about 5 to about 1000 mg (1 g), more usually about 100 to about 500 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and

other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

The active compound can be effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It will be understood, however, that the amount of the compound actually administered will usually be determined by a physician, according to the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, the active ingredient is typically dispersed evenly throughout the composition so that the composition can be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, about 0.1 to about 1000 mg of the active ingredient of the present invention.

The tablets or pills of the present invention can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the compounds and compositions of the present invention can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions or suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described supra. In some embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions can be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device can be attached to a face mask tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an appropriate manner.

The amount of salt or composition administered to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions can be administered to a patient already suffering from a disease in an amount sufficient to cure or at least partially arrest the

symptoms of the disease and its complications. Effective doses will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the disease, the age, weight and general condition of the patient, and the like.

The compositions administered to a patient can be in the form of pharmaceutical compositions described above. These compositions can be sterilized by conventional sterilization techniques, or may be sterile filtered. Aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

The therapeutic dosage of the salts of the present invention can vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. The proportion or concentration of a salt of the invention in a pharmaceutical composition can vary depending upon a number of factors including dosage, chemical characteristics (e.g., hydrophobicity), and the route of administration. For example, the salts of the invention can be provided in an aqueous physiological buffer solution containing about 0.1 to about 10% w/v of the compound for parenteral administration. Some typical dose ranges are from about 1 µg/kg to about 1 g/kg of body weight per day. In some embodiments, the dose range is from about 0.01 mg/kg to about 100 mg/kg of body weight per day. The dosage is likely to depend on such variables as the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, formulation of the excipient, and its route of administration. Effective doses can be extrapolated from dose-response curves derived from in vitro or animal model test systems.

The compositions of the invention can further include one or more additional pharmaceutical agents such as a chemotherapeutic, steroid, anti-inflammatory compound, or immunosuppressant, examples of which are listed hereinabove.

#### Labeled Compounds and Assay Methods

Another aspect of the present invention relates to labeled salts of the invention (radio-labeled, fluorescent-labeled, etc.) that would be useful not only in imaging techniques but also in assays, both in vitro and in vivo, for localizing and quantitating JAK in tissue samples, including human, and for identifying JAK ligands by inhibition binding of a labeled compound. Accordingly, the present invention includes JAK assays that contain such labeled compounds.

The present invention further includes isotopically-labeled salts of the invention. An "isotopically" or "radio-labeled" compound is a salt of the invention where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (i.e., naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present invention include but are not limited to <sup>2</sup>H (also written as D for deuterium), <sup>3</sup>H (also written as T for tritium), <sup>11</sup>C, <sup>13</sup>C, <sup>14</sup>C, <sup>13</sup>N, <sup>15</sup>N, <sup>15</sup>O, <sup>17</sup>O, <sup>18</sup>O, <sup>18</sup>F, <sup>35</sup>S, <sup>36</sup>Cl, <sup>82</sup>Br, <sup>75</sup>Br, <sup>76</sup>Br, <sup>77</sup>Br, <sup>123</sup>I, <sup>124</sup>I, <sup>125</sup>I and <sup>131</sup>I. The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for in vitro metalloprotease labeling and competition assays, compounds that incorporate <sup>3</sup>H, <sup>14</sup>C, <sup>82</sup>Br, <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S or will generally be most useful.

For radio-imaging applications <sup>11</sup>C, <sup>18</sup>F, <sup>125</sup>I, <sup>123</sup>I, <sup>124</sup>I, <sup>131</sup>I, <sup>75</sup>Br, <sup>76</sup>Br, or <sup>77</sup>Br, or will generally be most useful.

It is understood that a "radio-labeled" or "labeled compound" is a salt that has incorporated at least one radionuclide. In some embodiments the radionuclide is selected from the group consisting of <sup>3</sup>H, <sup>14</sup>C, <sup>125</sup>I, <sup>35</sup>S and <sup>82</sup>Br.

The present invention can further include synthetic methods for incorporating radio-isotopes into compounds of the invention. Synthetic methods for incorporating radio-isotopes into organic compounds are well known in the art, and a person of ordinary skill in the art will readily recognize the methods applicable for the compounds of invention.

A labeled salt of the invention can be used in a screening assay to identify/evaluate compounds. For example, a newly synthesized or identified compound (i.e., test compound) which is labeled can be evaluated for its ability to bind a JAK by monitoring its concentration variation when contacting with the JAK, through tracking of the labeling. For example, a test compound (labeled) can be evaluated for its ability to reduce binding of another compound which is known to bind to a JAK (i.e., standard compound). Accordingly, the ability of a test compound to compete with the standard compound for binding to the JAK directly correlates to its binding affinity. Conversely, in some other screening assays, the standard compound is labeled and test compounds are unlabeled. Accordingly, the concentration of the labeled standard compound is monitored in order to evaluate the competition between the standard compound and the test compound, and the relative binding affinity of the test compound is thus ascertained.

#### Kits

The present invention also includes pharmaceutical kits useful, for example, in the treatment or prevention of JAK-associated diseases or disorders, such as cancer, inflammation, or skin disorders, which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a salt of the invention. Such kits can further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, can also be included in the kit.

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of noncritical parameters which can be changed or modified to yield essentially the same results.

## EXAMPLES

### Example 1

Preparation of (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile maleic acid salt

To a test tube was added (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile (153.7 mg, 0.5 mmol) and maleic acid (61.7 mg) followed by isopropyl alcohol (IPA) (4 mL). The resulting mixture was heated to clear, cooled to room temperature, and then stirred for another 2.5 hours. The precipitate was collected by filtra-



## 11

tion and the cake was washed with 0.8 mL of cold IPA. The cake was dried under vacuum to constant weight to provide the final salt product (173 mg).

The maleic acid salt was shown to be a 1:1 salt by  $^1\text{H}$  NMR and crystallinity was confirmed by X-ray powder diffraction (XRPD). Differential scanning calorimetry (DSC) gave a sharp melting peak at about 175.96° C. (onset at 175.67° C.). The product showed only slight weight loss up to 150° C. by thermogravimetric analysis (TGA).

## Example 2

Preparation of (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile phosphoric acid salt

To a test tube was added (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile (153.5 mg) and phosphoric acid (56.6 mg) followed by isopropyl alcohol (IPA) (5.75 mL). The resulting mixture was heated to clear, cooled to room temperature, and then stirred for another 2 hours. The precipitate was collected by filtration and the cake was washed with 0.6 mL of cold IPA. The cake was dried under vacuum to constant weight to provide the final salt product (171.7 mg).

The phosphoric acid salt was shown to be a 1:1 salt by  $^1\text{H}$  NMR and crystallinity was confirmed by X-ray powder diffraction (XRPD). Differential scanning calorimetry (DSC) gave a sharp melting peak at about 198.66° C. The product showed little weight loss up to 200° C. by TGA.

## Example 3

Preparation of (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile sulfuric acid salt

To a test tube was added (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile (153.0 mg) and sulfuric acid (56.1 mg) followed by acetonitrile (7.0 mL). The resulting mixture was heated to clear, cooled to room temperature, and then stirred for another 2 hours. The precipitate was collected by filtration and the cake was washed with 0.8 mL of cold acetonitrile. The cake was dried under vacuum to constant weight to provide the final salt product (180 mg).

The sulfuric acid salt was shown to be a 1:1 salt by  $^1\text{H}$  NMR and crystallinity was confirmed by X-ray powder diffraction (XRPD). Differential scanning calorimetry (DSC) gave a sharp melting peak at about 186.78° C. The product showed little weight loss up to 175° C. by TGA.

## Example A

## In Vitro JAK Kinase Assay

Inhibitory activity of test compounds on JAK targets can be tested according to the following in vitro assay described in Park et al., *Analytical Biochemistry* 1999, 269, 94-104. The catalytic domains of human JAK1 (a.a. 837-1142), Jak2 (a.a. 828-1132) and Jak3 (a.a. 781-1124) with an N-terminal His tag are expressed using baculovirus in insect cells and purified. The catalytic activity of JAK1, JAK2 or JAK3 is assayed by measuring the phosphorylation of a biotinylated peptide.

## 12

The phosphorylated peptide was detected by homogenous time resolved fluorescence (HTRF).  $\text{IC}_{50}$ s of compounds are measured for each kinase in the reactions that contain the enzyme, ATP and 500 nM peptide in 50 mM Tris (pH 7.8) buffer with 100 mM NaCl, 5 mM DTT, and 0.1 mg/mL (0.01%) BSA. The ATP concentration in the reactions is 90  $\mu\text{M}$  for Jak1, 30  $\mu\text{M}$  for Jak2 and 3  $\mu\text{M}$  for Jak3. Reactions are carried out at room temperature for 1 hr and then stopped with 20  $\mu\text{L}$  45 mM EDTA, 300 nM SA-APC, 6 nM Eu-Py20 in assay buffer (Perkin Elmer, Boston, Mass.). Binding to the Europium labeled antibody takes place for 40 minutes and HTRF signal is measured on a Fusion plate reader (Perkin Elmer, Boston, Mass.). Both the phosphoric acid salt of the invention, and the corresponding free base compound, were found to have  $\text{IC}_{50}$  values of less than 50 nM for each of JAK1, JAK2, and JAK3.

Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. Each reference cited in the present application is incorporated herein by reference in its entirety.

What is claimed is:

1. A method of treating a myeloproliferative disorder in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a compound that is (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile phosphoric acid salt.

2. The method of claim 1, wherein the myeloproliferative disorder is polycythemia vera (PV).

3. The method of claim 1, wherein the myeloproliferative disorder is essential thrombocythemia (ET).

4. The method of claim 1, wherein the myeloproliferative disorder is myeloid metaplasia with myelofibrosis (MMM).

5. The method of claim 1, wherein (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile phosphoric acid salt is a crystalline salt.

6. The method of claim 5, wherein said crystalline salt is 1:1 (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile:phosphoric acid salt.

7. A method of treating a myeloproliferative disorder in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a pharmaceutical composition comprising (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile phosphoric acid salt and a pharmaceutically acceptable carrier.

8. The method of claim 7, wherein the myeloproliferative disorder is polycythemia vera (PV).

9. The method of claim 7, wherein the myeloproliferative disorder is essential thrombocythemia (ET).

10. The method of claim 7, wherein the myeloproliferative disorder is myeloid metaplasia with myelofibrosis (MMM).

11. The method of claim 7, wherein the pharmaceutical composition is suitable for oral administration.

12. The method of claim 7, wherein the pharmaceutical composition is in tablet form.

13. The method of claim 7, wherein (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile phosphoric acid salt is a crystalline salt.

14. The method of claim 13, wherein said crystalline salt is 1:1 (R)-3-(4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl)-3-cyclopentylpropanenitrile:phosphoric acid salt.

\* \* \* \* \*