

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
17 April 2014 (17.04.2014)

(10) International Publication Number
WO 2014/058427 A1

(51) International Patent Classification:

A61K 9/107 (2006.01) *A61K 31/575* (2006.01)
A61K 9/12 (2006.01) *A61P 17/06* (2006.01)

(21) International Application Number:

PCT/US2012/059701

(22) International Filing Date:

11 October 2012 (11.10.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant (for all designated States except US): **ANA-PLASI PHARMACEUTICALS LLC** [US/US]; 38-25 Astoria Blvd., Astoria, NY 11103 (US).

(72) Inventor; and

(71) Applicant (for US only): **YUHAS, Edward, Richard** [US/US]; 40 Cambridge Avenue, Yonkers, NY 10707 (US).

(74) Agent: **ENDRES, Martin, P.**; Florek & Endres PLLC, 1156 Avenue Of The Americas, Suite 600, New York, NY 10036 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: METHOD AND COMPOSITIONS FOR TREATING PSORIASIS

(57) Abstract: Method and composition for treating psoriasis and other skin disorders.



WO 2014/058427 A1

METHOD AND COMPOSITIONS FOR TREATING PSORIASIS

TECHNICAL FIELD OF THE INVENTION:

The present invention relates to a method and compositions for treating psoriasis
5 and other similar skin disorders such as eczema, sunburn, and wound and scar care. The
method comprises the topical administration of a composition comprising a sterol ester.

BACKGROUND OF THE INVENTION:

Psoriasis is a skin disease that results when skin cells reproduce at a faster than
10 normal rate. Psoriasis produces red patches, sometimes referred to as plaques, and dry
grayish-white or silvery scales on a patient's skin.

A number of different treatments are currently employed to treat psoriasis
outbreaks. One of the more common treatments is the topical application of
corticosteroids to the affected area to reduce the inflammation. The patient may develop
15 a resistance over time to corticosteroids and therefore their use is often limited.

Other common treatments involve the topical application of anthralin, also known
as dithranol, the topical application of a vitamin D such as calcipotriene or calcitriol, or
the topical application of a vitamin A such as tazarotene. It is believed that anthralin,
vitamin D and vitamin A slow the patient's production of new skin cells. Anthralin can
20 stain a patient's skin or clothing and does not work well on very active psoriasis
outbreaks. The topical applications of vitamins D and/or A may make a patient sensitive
to sun light.

Coal tar has also been used to treat psoriasis. The coal tar can be applied directly to the affected area or it can be added to bath water. Although helpful, coal tar has a strong and unpleasant odor.

The topical application of salicylic acid has also been employed with some psoriasis patients. Salicylic acid is believed to function as a peeling agent that will cause the outer layer of skin to shed. Salicylic acid may cause irritation and/or hair loss if left on the skin for extended periods of time.

Some psoriasis patients have also used ultraviolet light to treat psoriasis. The ultraviolet light can involve the use of controlled amounts of natural sunlight on the exposed area or artificial sources directed to the affected area.

Other patients have combined oral administration of psoralen, also known as psoralene, with ultraviolet light treatments.

Still other psoriasis patients have used orally administered retinoids, i.e., very concentrated forms of vitamin A, or drugs that slow down the patient's immune system such as methotrexate or cyclosporine.

Psoriasis patients are also advised to keep the skin moist by using moisturizing creams, lotions and soaps to help reduce the redness and itching associated with the psoriasis outbreaks or to use bath oils and salts.

Although there are many potential treatment options for psoriasis, none of the above mentioned methods are 100% successful or without adverse potentials.

It is an object of the present invention to provide a method for treating psoriasis and to provide compositions that are useful in treating psoriasis without unpleasant odors or irritation to the patient's skin.

SUMMARY OF THE INVENTION:

The present invention accomplishes the above objectives and others by providing a method of treating psoriasis comprising the topical application of a composition

5 comprising a sterol ester, preferably C₁₀-C₃₀ carboxylic acid sterol esters, to the psoriatic area of a patient. The application of the composition will occur as needed but should occur at least once daily and preferably immediately or shortly after, i.e., within 5, 10 or 15 minutes, washing and drying the affected area. The application of the composition to the psoriatic area should continue periodically until the psoriasis plaques and scales have
10 been removed.

The composition may be applied to the psoriatic area in an unoccluded or occluded manner. In one embodiment of the invention the composition should remain in contact with the psoriatic area for at least 20 minutes or longer, preferably at least 30 minutes or longer.

15 The composition may be a gel, cream, paste, lotion, ointment, salve, serum, spray, aerosol, mousse or foam. Preferably the composition is a serum, aerosol, mousse or foam.

In one embodiment of the present invention, the composition should comprise 25% or more of a sterol ester, preferably a C₁₀-C₃₀ carboxylic acid sterol ester, based
20 upon the total weight of the composition less solvent, preferably 30 % or more based upon the total weight of the composition less solvent and most preferably 40% or more based upon the total weight of the composition less solvent.

In an alternative embodiment of the present invention, the composition of the present invention will comprises a mixture of a sterol ester, preferably a C₁₀-C₃₀ carboxylic acid sterol ester, and a penetration enhancer.

It is also believed that the compositions of the present invention are also useful in
5 treating skin disorders and afflictions such as eczema, sunburn, and wound and scar care

DETAILED DESCRIPTION OF THE INVENTION:

As used herein, the term “psoriasis” refers to a dermatological condition that is believed to be caused by the patient’s body producing too many skin cells. The condition
10 is associated with red patches, thick white/silvery scales and often swelling on the patient’s skin, typically on the arms, scalp, ears and pubic area. It is believed that psoriasis cannot be cured but may be treated.

As used herein, the terms “treat” or “treating” refers to providing relief of one or more of the conditions associated with psoriasis or diminishing or lessening any one or
15 more of the conditions associated with psoriasis.

As used herein, the terms “enhancement,” “penetration enhancement” and “permeation enhancement” mean an increase in the permeability of a biological membrane (e.g., skin or mucosa) to a drug or biologically active substance, so as to increase the rate at which the drug or biologically active substance permeates through the
20 membrane. “Permeation enhancer,” “enhancer,” “penetration enhancer” or similar terms mean a material that achieves such permeation enhancement.

As used herein, “transdermal” means delivery of a drug or biologically active substance by passage into and through the skin or mucosal tissue. Hence, the terms

“transdermal” and “transmucosal” are used interchangeably unless specifically stated otherwise. Likewise, the terms “skin,” “derma,” “epidermis,” “mucosa” and the like will also be used interchangeably unless specifically stated otherwise.

As used herein, the term “topical” refers to outer skin or derma of a patient.

5 Hence, the phrase “topical application” refers to the application of a composition of the present invention and its various embodiments to the outer surface of a patient’s skin or derma.

As used herein, the terms “occlude,” “occluded,” “occlusive” and the like refer to a transdermal formulation that is applied to the skin with the use of a supporting or
10 otherwise associated structure. In other words, a topical formulation may be applied to the skin of a patient with the aid of a structure, such as a backing member, bandage or cover. A matrix patch is an example of an occluded device. Conversely, “unoccluded” and “non-occluded,” which may be used interchangeably, refer to a transdermal formulation that is applied to the skin without the use of a support, backing member,
15 cover or otherwise associated structure. In other words, the transdermal formulation is applied to the skin in a free form, which is sufficient to effect transdermal delivery of the drug or biologically active substance without the use of structures, such as a backing member, etc. A gel formulation is an example of a non-occluded composition; other non-occluded compositions include ointments, lotions, pastes, mousses, aerosols and creams.

20 The terms “combination” and “mixture” are used interchangeably and synonymously. These terms should always be interpreted as over-inclusive rather than under-inclusive where appropriate.

Concentration, weight percent and other numerical data may be presented herein in a range format. It is to be understood that such range format is used merely for convenience and brevity and should be interpreted flexibly to include not only the numerical values explicitly recited as the limits of the range, but also to include all the individual numerical values or sub-ranges encompassed within that range as if each numerical value and sub-range is explicitly recited. For example, a percent range of 1% to 20% should be interpreted to include not only the explicitly recited percent limits of 1% and 20% but also to include individual percentages such as 1.5%, 3%, 4.75%, 8.34% and sub-ranges such as 1% to 5%, 10% to 15%, 4.7% to 11.9% etc.

The present invention is a method for treating psoriasis and a composition that is useful for treating psoriasis.

The method comprises the topical application of a composition to the psoriatic area of a patient suffering from a psoriasis outbreak. The composition comprises a sterol ester, preferably a C₁₀-C₃₀ carboxylic acid sterol ester. The composition may be a gel, cream, paste, lotion, ointment, salve, serum, spray, aerosol, mousse or foam.

The method further comprises the steps of washing the psoriatic area, drying the psoriatic area and optionally exfoliating the psoriatic area prior to the application of the composition. The method or topical application of the composition should occur at least once a day or twice a day directly onto the psoriatic area. Three, 4, 5 or more applications of the composition to the psoriatic area may occur per day depending upon the severity of the psoriasis outbreak and location on the patient's body.

Depending upon the form in which the composition is applied, i.e., gel, serum or cream, after the application of the composition to the psoriatic area, the composition and

psoriatic area may be covered with a protective cloth, bandage, gauze or wrap to prevent transfer of the composition to a patient's clothing, furniture or third parties. The protective cloth, bandage, gauze or wrap may also be applied to protect the psoriatic area from dirt, germs and bacteria.

5 The method of the present invention is repeated every day or portion of the day as needed until the psoriasis outbreak has resided or ended.

 The composition of the present invention comprises a sterol ester, preferably a C₁₀-C₃₀ carboxylic acid sterol ester, and most preferably a C₁₀-C₃₀ carboxylic acid cholesterol/lanosterol mixture commercially available from Croda Chemicals Europe,
10 Ltd., of East Yorkshire, England under the trade name SUPER STEROL ESTER[®]. It is believed that C₁₀-C₃₀ cholesterol/lanosterol ester mixture sold by Croda Chemicals is derived from wool wax by a process described in Koresawa et al., U.S. Patent No. 4,138,416, which is incorporated in its entirety herein by reference. Additional examples of sterol esters that may be used in the composition of the present invention include but
15 are not limited to stearate, palmitate, acetate, lanolate, macadamiate, nonanoate, oleate, butyrate, hydroxystearate, isostearate, sulfate, isostearate carbonate of cholesterol and other sterols.

 The amount of sterol ester in the composition of the present invention should comprise 25% or more of the sterol ester based upon the weight of the composition less
20 solvent, preferably 30 % or more based upon the total weight of the composition less solvent and most preferably 40% or more based upon the total weight of the composition less solvent. Because numerous excipients in a transdermal formulation may have the ability to impart multiple functions depending upon the concentration and manner in

which they are used, when determining the amount of sterol ester less the amount of solvent present in a composition prepared in accordance with the present invention, the solvents present in the composition but excluded from sterol ester weight percent calculations, are compounds that are liquids at room temperature and have a boiling point
5 below 150°C under normal atmospheric conditions, preferably a boiling point below 125°C and most preferably a boiling point below 105°C. Examples of typical solvents that should be excluded from the sterol weight percent calculations are water, low molecular weight alcohols such as C₁-C₆ branched or straight chain alcohols, e.g., methanol, ethanol and isopropanol, low molecular weight ketones such as C₁-C₆
10 branched or straight chain ketones, e.g., acetone, aromatic compounds and low molecular weight alkanes such as C₁-C₁₀ branched or straight chain alkanes.

The compositions of the present invention may further comprise a penetration enhancer. The amount of penetration enhancer employed in the compositions of the present invention will vary depending upon the specific composition embodiment or
15 transdermal formulation, i.e., serum, cream or foam, and the specific penetration enhancer selected. Typically, the amount of penetration enhancer employed in the compositions of the present invention should be about 0.001 wt% to about 25 wt% , preferably about 0.005 wt% to about 15 wt% and most preferably about 0.01wt% to about 10 wt% based upon the total weight of the composition. Examples of penetration
20 enhancers that may be used in compositions of the present invention include, but are not limited to, fatty acids, fatty acid esters, fatty alcohols, fatty acid esters of lactic acid or glycolic acid, glycerol tri-, di- and monoesters, triacetin, short chain alcohols, amine oxides and mixtures thereof. Particular examples of permeation enhancers include oleyl

alcohol, lauryl alcohol, isopropyl myristate, oleyl oleate, levulinic acid, ethanol, glycerol monooleate, methyl laurate, sorbitain monooleate, triacetin, aloe vera oil, benzothonium chloride, cetyl dimethylamine oxide, cetyl alcohol, cetyl lactate, cocamidopropyl betaine, cocoamine oxide diethanolamine, dimethyloctylamine oxide, 2-

5 dodecoxyethyldimethylamine oxide, dimethyl-decylamine oxide, dimethylhexadecylamine oxide, dimethyl-tetradecylamine oxide, dimethyl isosorbide, dipropylene glycol, ethyl hexyl lactate, glycolic acid, 3-dodecoxy-2-hydroxypropyl di(3-hydroxypropyl)amine oxide, lactic acid, lauramine oxide, lauryl betaine, lauryl lactate, lauryl laurate, isopropyl palmitate, macrogol 15 hydroxystearate (Solutol HS 15),

10 menthol, menthyl lactate, myristyl alcohol, myristal lactate, octyldodecanol, octyl salicylate, oleamine oxide, oleic acid, oleyl betaine, oleyl di(2-hydroxyethyl) amine oxide, PEG 1000, pentadecalactone, propylene glycol, salicylic acid, stearyl alcohol, stearyl lactate, 3,6,9-trioxaheptadecyldiethylamine oxide, di(2-hydroxyethyl)-tetradecylamine oxide, triethanolamine triacetate and combinations thereof. Other permeation enhancers

15 useful with the present invention may be found in U.S. Patent Application Publication No. 2007/0269379, which is incorporated in its entirety herein by reference. Preferred permeation enhancers include oleyl alcohol, lauryl alcohol, isopropyl myristate, oleyl oleate, levulinic acid, glycerol monooleate, methyl laurate, sorbitain monooleate, triacetin, cetyl alcohol, cetyl lactate, dimethyl isosorbide, dipropylene glycol, ethyl hexyl

20 lactate, glycolic acid, lauramine oxide, lauryl betaine, lauryl lactate, lauryl laurate, isopropyl palmitate, myristyl alcohol, myristal lactate, octyl salicylate, oleamine oxide, oleic acid, oleyl betaine, salicylic acid, stearyl alcohol, stearyl lactate, triethanolamine triacetate and combinations thereof.

Depending upon the specific composition embodiment or transdermal formulation, i.e., serum, cream or foam, the composition of the present invention may also include further additives such as solvents, film forming/polymeric agents, viscosity increasing agents, emulsifiers, antioxidants, preservatives, pH adjusting agents,

5 propellants and combinations of the forgoing.

The compositions of the present invention may include any suitable solvent. Preferably, the solvent may include water and/or one or more organic compounds, e.g., esters, terpenes, alcohols, ketones, aldehydes, fatty acids, partially or fully esterified fatty acids, wherein the structures are cyclic, non-cyclic (e.g., alkyl), alicyclic (i.e., a bridged

10 ring compound), or aromatic, as well as organic compounds having combinations of these functional groups. Specific examples of solvents that may be employed are water, methanol, ethanol, isopropyl alcohol, acetone, hexane, butyl alcohol, ethyl acetate, polyethylene glycol, propylene glycol, ethylene glycol, triethylene glycol, glycerin, 1,3-propane diol, 2-methyl-1,3-propane diol, glycerol ricinoleate, mineral oil, peanut oil, corn

15 oil, cottonseed oil, sesame oil or a combination thereof. The solvent may be employed in any suitable amount. Typically, the solvent can be present in the composition in about 1.0 wt% to about 95.0 wt% based upon the total weight of the composition, preferably about 3.0 wt% to about 85 wt% based upon the total weight of the composition and most preferably about 5.0 wt% to about 75 wt% of the total weight of the composition.

20 The compositions of the present invention also may include a film-forming/polymeric agent. Although applicants do not wish to be bound by an particular theory, it is believed that the presence of a film-forming/polymeric agent in the compositions prepared in accordance with the present invention allow the compositions

to spread more easily over the skin and to form a protective barrier coating allowing the sterol ester to penetrate the skin of the psoriatic area and/or remain in contact with the skin of the psoriatic area. The film-forming/polymeric agent may also enhance the adherence of the composition to the patient's skin and improve the composition's

5 resistance to wash off or rub off.

Film-forming/polymeric agents are preferably soluble or miscible with the sterol ester and/or penetration enhancer. The composition of the present invention typically comprises from about 0.001 wt% to about 25 wt% , preferably about 0.005 wt% to about 15 wt% and most preferably about 0.010 wt% to about 10 wt% based upon the total

10 weight of the composition. Some examples of film-forming/polymeric agents that may be used in compositions of the present invention are polyalkenes, oleophilic copolymers of vinylpyrrolidone, acrylic copolymers, polyethylene glycol derivative, polyolefins, polyurethanes and mixtures thereof.

Examples of polyalkenes that may be included in the compositions of the present

15 invention are polyethylenes having a molecular weight ranging from about 300 to about 3000 (available as PERFORMALENE[®] from New Phase Technologies, Piscataway, NJ); polyisobutylenes (available as VISTANEX[™] from Exxon Chemical Company, Houston, TX); polyisobutenes (available as PRESERSE[™] from Sumitomo Corp.); polydecenes (SILKFLO[™] available from Amoco); and hydrogenated polyisobutenes (PANALANE[®]

20 available from Lipo Chemicals, Inc., Paterson, NJ).

Oleophilic copolymers of vinylpyrrolidone suitable for use in the compositions of the present invention may be copolymers of polyvinylpyrrolidone (PVP) and long chain alpha olefins, including, but not limited to, PVP/eicosene copolymers (GANEX[®] V-220

and V-220F), and tricontanyl PVP copolymers (GANEX[®]) available from Ashland, formerly International Specialty Products, Wayne, NJ.

Examples of acrylic copolymers that may be used in the compositions of the present invention include acrylic copolymers having long (C₈-C₃₀) alkyl chains to enhance their oleophilicity, such as acrylate/octylacrylamide copolymers (available as DERMACRYL[®] from Akzo Nobel). An example of a polyethylene glycol derivative that may be used as a film forming agent in compositions of the present invention is a polyethylene glycol derivative of Beeswax (ESTOL[®] E04BW-3752, E06BW-3753 or E03BW-3751 formerly available from Unichema, Wilmington, DE and currently available from Croda under the trade name CITHROL[®]). Examples of polyolefins that may be used as a film forming agent in compositions of the present invention are fatty acid ester/fatty acid anhydride grafted polyolefins wherein the esters and anhydrides are derived from C₁₂-C₂₂ fatty acid moieties, for example, C₃₀-C₃₈ olefin/isopropyl maleate/maleic anhydride copolymer (PERFORMA[™] V1608, available from New Phase Technologies, Piscataway, NJ).

A preferred group of film forming/polymeric agents that may be used in the compositions of the present invention include polyurethanes derived from isophorone diisocyanate such as those described in U.S. Patent Nos. 5,051,260 and 6,613,866 and sold by Alzo International Inc. under the tradename POLYDERM[®], polyisobutene/polybutene, hydrogenated polydecene and hydrogenated C₆-C₁₄ olefin polymers sold by ExxonMobil Chemical company under the tradename PURESYN[®]. The following table identifies a few of the preferred film forming/polymeric agents:

Commercial Name	Chemical Description	Origin
POLYDERM [®] PPI-BZ	Benzyl Alcohol-Ethylene	Alzo International Inc.

		Glycol/IPDI Copolymer	
5	POLYDERM [®] PPI-CA-15	Di-PEG-15 Cocamine/IPDI Copolymer	Alzo International Inc.
	POLYDERM [®] PPI-CO	Castor Oil/IPDI Copolymer	Alzo International Inc.
	POLYDERM [®] PPI-CO-H	Hydrogenated Castor Oil/IPDI Copolymer	Alzo International Inc.
10	POLYDERM [®] PPI-CO-40	PEG-40 Hydrogenated Castor Oil/IPDI Copolymer	Alzo International Inc.
15	POLYDERM [®] PPI-CO-200	PEG-200 Hydrogenated Castor Oil/IPDI Copolymer	Alzo International Inc.
	POLYDERM [®] PPI-DGDIS	Diglycerol Diisostearate/IPDI Copolymer	Alzo International Inc.
20	POLYDERM [®] PPI-GH	Glycereth-7 Hydroxystearate/IPDI Copolymer	Alzo International Inc.
	POLYDERM [®] PPI-PE	Diethylene Glycol Adipate/IPDI Copolymer	Alzo International Inc.
25	POLYDERM [®] PPI-SA	Di-2 PEG Soyamine/IPDI Copolymer	Alzo International Inc.
	POLYDERM [®] PPI-SI	Dimethiconol/IPDI Copolymer	Alzo International Inc.
30	POLYDERM [®] PPI-SI-50	Dimethiconol/IPDI Copolymer 50%	Alzo International Inc.
	POLYDERM [®] PPI-SI/SA	Dimethiconol-PEG-2 Soyamine/IPDI Copolymer	Alzo International Inc.
35	POLYDERM [®] PPI-SI-WI	Dimethicone Copolyol/IPDI Copolymer water insoluble	Alzo International Inc.
40	POLYDERM [®] PPI-SI-WS	Dimethicone Copolyol/IPDI Copolymer water soluble	Alzo International Inc.
	AVALURE [®] UR 450	PPG-17/IPDI/DMPA Copolymer	Noveon
45	POLYSYNLANE [®] Gel	Hydrogenated Polyisobutene (and) Butylene/Ethylene/Styrene Copolymer (and) Ethylene/Propylene/Styrene Copolymer	Collaborative Group
50	POLYFIX [®] JPN	Hydrogenated Polyisobutene	Collaborative Group

	(and) Polybutene	
	POLYSYNLANE [®] (HV)	Hydrogenated Polyisobutene Collaborative Group
5	PANALENE [®] H300E	Lipo Chemicals
	PANALENE [®] L-14E	Lipo Chemicals
10	PURESYN [®] 2, 4, 6, 8, 10, 40, 100, 150, 300 1000, 3000	Polydecene or Hydrogenated Polydecene ExxonMobil Chemicals

The preferred film forming/polymeric agents are water-insoluble, oleophilic and/or water-resistant.

15 The compositions of the present invention may also contain viscosity enhancing agents that thicken, gel or harden the composition. The composition of the present invention typically comprises from about 0.001 wt% to about 50 wt% of the viscosity enhancing agent, preferably about 0.005 wt% to about 40 wt% and most preferably about 0.01 wt% to about 25 wt% based upon the total weight of the composition. Exemplary

20 viscosity enhancing agents include organic materials such as natural or synthetic waxes, C₁₂-C₆₀ alcohols, C₁₂-C₆₀ acids, alpha-hydroxy fatty acids, polyhydroxy fatty acid esters, polyhydroxy fatty acid amides, and inorganic/organic materials such as metal ester complexes containing zinc, calcium, aluminum or magnesium, fumed silicas, and organoclays. Additional viscosity enhancing agents include polyol polyesters, glyceryl

25 esters, polyglyceryl esters and polysiloxanes that are a solid or semi-solid at ambient temperature.

Specific examples of viscosity enhancing agents that may be included in the compositions of the present invention include C₁₂-C₆₀ alcohols, preferably C₁₆-C₂₂ fatty alcohols, such as cetyl alcohol, stearyl alcohol, behenyl alcohol and mixtures thereof.

Other suitable viscosity enhancing agents include C₁₂-C₆₀ acids, preferably C₁₆-C₂₂ fatty acids, such as palmitic acid, stearic acid, behenic acid, oleic acid, linoleic acid, myristic acid, ricinoleic acid, eucic acid, lauric acid, isostearic acid and mixtures thereof. Further suitable viscosity enhancing agents that may be used herein are alpha-hydroxy fatty acids, including 12-hydroxystearic acid, 12-hydroxylauric acid, 16-hydroxyhexadecanoic acid and mixtures thereof. Additional examples of suitable fatty acids are further described in Klofta et al., U.S. Patent No. 7,449,613, Hofrichter, et al., U.S. Patent No. 5,429,816 and Motley, U.S. Patent. No. 5,552,136, disclosure of each is incorporated in its entirety herein by reference.

Waxes are also suitable for use as viscosity enhancing agents in compositions of the present invention. Natural waxes may include, but are not limited to, carnauba, ozokerite, beeswax, candelilla, paraffin, ceresin, esparto, ouricuri, rezowax and other known mined and mineral waxes. Synthetic waxes may include, but are not limited to, paraffin waxes and microcrystalline waxes.

Additional viscosity enhancing agents that may be used herein include polyhydroxy fatty acid esters, polyhydroxy fatty acid amides and mixtures thereof. Preferred esters and amides will have three or more free hydroxy groups on the polyhydroxy moiety and are typically nonionic in character. Because of the possible skin sensitivity of those using articles to which the composition is applied, these esters and amides should also be relatively mild and non-irritating to the skin. Suitable polyhydroxy fatty acid esters and polyhydroxy fatty acid amides are disclosed in Roe et al., U.S. Patent No. 5,643,588, the disclosure of which is incorporated in its entirety herein by reference.

Still further viscosity enhancing agents that may be included in the compositions of the present invention are gelling agents. Gelling agents are materials that can swell or expand when in contact with water. Examples of gelling agents that may be used in the present invention include swellable polymers, also known as osmopolymers or hydrogels.

5 The swellable polymer can be non-cross-linked or lightly cross-linked. The cross-links can be covalent or ionic bonds with the polymer possessing the ability to swell in the presence of fluid, and when cross-linked it will not be dissolved in the fluid. The polymer can be of plant, animal or synthetic origin. Polymeric materials useful for the present purpose include polyhydroalkylcellulose having a molecular weight greater than
10 50,000, such as hydroxyl propylmethylcellulose (METHOCEL[®] K 100M available from Dow Chemical); poly(hydroxyalkylmethacrylate) having a molecular weight of from 5,000 to 5,000,000; poly(vinylpyrrolidone) having a molecular weight of from 100,000 to 3,000,000; anionic and cationic hydrogels; poly(electrolyte) complexes; poly(vinylalcohol) having a low acetate residual; a swellable mixture of agar and
15 carboxymethyl cellulose; a swellable composition comprising methyl cellulose mixed with a sparingly cross-linked agar; a polyether having a molecular weight of from 10,000 to 6,000,000; a water-swallowable copolymer produced by a dispersion of a finely divided copolymer of maleic anhydride with styrene, ethylene, propylene, or isobutylene; a water-swallowable polymer of N-vinyl lactams and the like.

20 Other gelling agents useful in the present invention include pectin having a molecular weight ranging from 30,000 to 300,000; polysaccharides such as agar, acacia, karaya, tragacanth, algin and guar; CARBOPOL[®], an acrylic acid polymer, a carboxyvinyl polymer, sometimes referred to as carboxypolymethylene, a polymer of

acrylic acid cross-linked with a polyallyl ether of sucrose, as described in U.S. Patent Nos. 2,798,053 and 2,909,462 and available as CARBOPOL[®] 934, 940 and 941, and its salt derivatives; polyacrylamides; water-swellaable indene maleic anhydride polymers; GOOD-RITE[®] polyacrylic acid having a molecular weight of 80,000 to 200,000;

5 POLYOX[®] polyethylene oxide polymers having a molecular weight of 100,000 to 7,000,000; starch graft copolymers; AQUA-KEEP[®] acrylate polymers with water absorbability of about 400 times its original weight; diesters of polyglucan; a mixture of cross-linked polyvinyl alcohol and poly(N-vinyl-2-pyrrolidone); poly(ethylene glycol) having a molecular weight of 4,000 to 100,000. Representative polymers possessing

10 gelling properties are described in U.S. Patent. Nos. 6,419,954, 4,915,949, 4,327,725, 4,207,893 and in *Handbook of Common Polymers*, by Scott and Roff, published by Cleveland Rubber Company, Cleveland, OH.

Examples of inorganic viscosity enhancing agents that may be included in the compositions of the present invention include treated and untreated fumed silicas such as

15 those available from Cabot Corp., Tuscola, IL under the trade designations CAB-O-SIL M5 and MS-55. Exemplary surface-treated fumed silicas are also available from Cabot Corp., Tuscola, IL under the trade designations TS-720 and TS-610.

Suitable clays such as hectorite and smectite may also be used as viscosity enhancing agents in compositions of the present invention.

20 Hydrogenated vegetable oils such as cocoa butter, shea butter and mixtures thereof may also be used as viscosity enhancing agents in compositions of the present invention.

Suitable petroleum-based emollients may also be used as viscosity enhancing agents in compositions of the present invention. Examples of suitable petroleum-based emollients that may be used include petrolatums, i.e., hydrocarbons or mixtures of hydrocarbons; particularly preferred are hydrocarbons having chain lengths of from C₁₀ to C₁₀₀. Petroleum-based emollients within this chain length range include mineral oil and petrolatum. Mineral oil usually refers to less viscous mixtures of hydrocarbons having from 10 to 30 carbon atoms, though the hydrocarbon molecular weight distribution may vary. Since the lower molecular weight hydrocarbons can cause irritation in some individuals, mineral oils having a small percentage of lower molecular weight hydrocarbons are preferred. Petrolatum usually refers to more viscous mixtures of hydrocarbons of higher molecular weight hydrocarbons. Petrolatum and mineral oil are preferred skin conditioning agents for compositions of the present invention due to their ability to protect the skin from harmful or irritating stimuli. Petrolatum is particularly preferred because of its good barrier properties.

The compositions of the present invention may also contain humectants. The compositions of the present invention typically comprises from about 0.001 wt% to about 30 wt% of a humectant, preferably about 0.005 wt% to about 20 wt% and most preferably about 0.01wt% to about 10 wt% based upon the total weight of the composition. Examples of compounds that may be used as humectants in compositions of the present invention are esters of polyhydroxy alcohols. This type of ester may include glyceryl esters including glycerides and derivatized glycerides, polyglyceryl esters, and glycolic esters. Glyceryl esters are derived from glycerin, its derivatives and one or more carboxylic acid moieties. Non-limiting examples include various C₁-C₃₀

mono-, di- or tri-esters of glycerin and derivatives thereof, such as mono-, di-, tri-glycerides, acetoglycerides, and ethoxylated glycerides. Exemplary glyceryl esters include glyceryl behenate, glyceryl oleate, glyceryl stearate, glyceryl palmitate, glyceryl distearate, glyceryl dipalmitate and the like. Polyglyceryl esters having C₁₂-C₂₂ acid
5 moieties are also suitable for use herein. Non-limiting examples include polyglyceryl-4 isostearate, polyglyceryl-3 oleate, diglyceryl monooleate, tetraglyceryl monooleate and the like. Glycolic esters are derived from C₂-C₆ glycols, including ethylene glycol, propylene glycol, butylene glycol, hexylene glycol and derivatives thereof, and one or more carboxylic acid moieties having C₁-C₃₀ chains. Specific examples of glycolic esters
10 include polyethylene glycols (PEGs), such as PEG-2, PEG-3, PEG-30 and PEG-50, and polypropylene glycols (PPGs), such as PPG-9, PPG-12, PPG-15, PPG-17, PPG-20, PPG-26, PPG-30 and PPG-34.

The compositions of the present invention may also contain emulsifiers or dispersing agents such as anionic, cationic and nonionic surfactants. The compositions of
15 the present invention typically comprises from about 0.001 wt% to about 15 wt% of an emulsifier or dispersing agent, preferably about 0.005 wt% to about 10 wt% and most preferably about 0.01wt% to about 5 wt% based upon the total weight of the composition. Nonionic surfactants are preferred because of their low level of irritation to skin. Typical nonionic surfactants are monoglycerides such as glyceryl monostearate and the like;
20 sorbitan aliphatic esters such as sorbitan monolaurate and the like; sucrose aliphatic esters; polyoxyethylene aliphatic esters such as polyoxyethylene stearate; and polyoxyethylene higher alcohol ethers such as polyoxyethylene cetyl ether, polyoxyethylene oleyl ether, polyoxyethylene fatty ethers and the like.

The compositions of the present invention may also contain an antioxidant to minimize or prevent the oxidation process and enhance the shelf life of the composition. The compositions of the present invention typically comprises from about 0.001 wt% to about 25 wt% of an anti-oxidant, preferably about 0.005 wt% to about 15 wt% and most
5 preferably about 0.01wt% to about 10 wt% based upon the total weight of the composition. Antioxidants useful herein should preferably be mild and non-irritating. Antioxidants from natural sources are preferred, such as Vitamin E and its derivatives, including tocopherol, tocopherol acetate, mixed tocopherols (available as COVI-OX T-50 or T-70 from Henkel Corp, Ambler, PA), and the like or butylated hydroxytoluene,
10 butylated hydroxyanisole, sodium pyrosulfite, acetone sodium bisulfate and the like. Some of these antioxidants are also useful as skin antioxidants, which minimizes the wrinkles and dullness of the skin and provides a more youthful looking and firmer textured skin.

The compositions of the present invention may also contain a preservative to
15 prevent bacterial growth and odors thereof, particularly in compositions having a relatively high water content. The compositions of the present invention typically comprise from about 0.001 wt% to about 10 wt% of a preservative, preferably about 0.005 wt% to about 5 wt% and most preferably about 0.01wt% to about 2.5 wt% based upon the total weight of the composition. Suitable preservatives include propyl paraben,
20 methyl paraben, benzyl alcohol, benzalkonium chloride, tribasic calcium phosphate, phenoxyethanol, or acids such as citric, tartaric, maleic, lactic, malic, benzoic, salicylic and the like.

The compositions of the present invention may include an acid or base to adjust the pH of the composition to the desired or optimal range. Examples of compounds typically used to adjust the pH of topical compositions include oleic acid, hydrochloric acid, citric acid, lactic acid, tartaric acid, glacial acetic acid, sodium hydroxide or the like.

5 Depending upon the form in which the composition is applied, i.e., gel, serum or cream, and the location, the desired final pH value of the composition may vary, however, it is generally preferred that the composition range from a pH of about 5.0 to about 8.5, preferably about 6 to about 8.0 and most preferably about 6.5 to about 7.5.

In order to increase the stability of the compositions of the present invention, it
10 may be necessary to add a chelating agent. The chelating agents may include ethylenediaminetetraacetic acid (EDTA) and its derivatives, thioglycolic acid, thiolactic acid, thioglycerol and the like.

A fragrance may also be added to composition of the present invention if desired.

If the composition of the present invention is an aerosol, foam or mouse, the
15 composition will require a propellant for dispensing the composition from the container. The propellant may be any type of propellant commonly used in the cosmetic/pharmaceutical industry such as nitrogen, carbon dioxide, dimethyl ether, hydrocarbons, i.e., methane, ethane, propane, butanes and pentanes, halogenated hydrocarbons, i.e., CH_2ClF , $\text{CClF}_2\text{CHClF}$, CF_3CHClF , $\text{CHF}_2\text{CClF}_2$, CHClFCHF_2 ,
20 $\text{CF}_3\text{CH}_2\text{Cl}$, CClF_2CH_3 , CHF_2CHF_2 , $\text{CF}_3\text{CH}_2\text{F}$ (HFC 134a), CHF_2CH_3 (HFC 152a), CF_3CHF_2 (HFC 227), CF_3CF_3 and $\text{CF}_3\text{CF}_2\text{CF}_3$. Some of the more commonly used hydrocarbon propellants are A-46 (15.2% propane/84.8% isobutene); and NP-46 (25.9% propane/74.1% n-butane), NIP-46 (21.9% propane/31.3% isobutene/46.8% n-butane).

The amount of propellant will depend on the type of container for the composition of the present invention, the amount of the composition in the container, the amount of composition to be dispensed per actuation and the form in which the composition will be dispensed, i.e., mist or foam. The optimization of the propellant and container are within
5 the ability of the skilled artisan and examples can be found in Wai-Chiu So et al., U.S. Patent No. 6,946,120 and Remington, Science and Practice of Pharmacy, 21st ed., pp. 1000-1017 which are incorporated in their entirety herein by reference. The propellant is generally not included in the calculation of the weight percentages of the composition prepared in accordance with the present invention because it is merely part of the
10 dispensing device and typically does not remain part of the composition once the composition is dispensed and applied to the patient's psoriatic area.

The aerosols, foams and mousses of the present invention will include a solvent, preferably water and/or a lower alcohol, i.e., C₁-C₆ alcohols such as methanol, ethanol, isopropanol or mixtures thereof. The aerosols, foams or mousses may also comprise a
15 co-solvent selected from one or more of the group consisting of aromatic and polyhydric alcohols such as 1,3-butylene glycol, propylene glycol, polyethylene glycol 400, hexylene glycol and dipropylene glycol or glycerol. When the co-solvent is present, it may be present in amounts of approximately 10% by weight or less, preferably approximately 5% by weight or less based upon the total weight of the composition.

20 The composition of the present invention may also include an anti-inflammatory drug, a topical anesthetic or a combination thereof. Typical topical anesthetics include, but are not limited to, lidocaine, xylocaine, buprenorphine and fentanyl. Suitable topical anesthetics are known to those of skill in the art and are disclosed, e.g., in Goodman

Gilman, Alfred; Goodman, Louis S.; Gilman, Alfred; Goodman and Gilman's The Pharmacological Basis of Therapeutics, Ninth Edition, pp. 331-347.

Any suitable topical corticosteroid can be employed as an anti-inflammatory drug. Suitable corticosteroids are known to those of skill in the art and are disclosed in, e.g.,

- 5 Goodman Gilman, Alfred; Goodman, Louis S.; Gilman, Alfred; Goodman and Gilman's The Pharmacological Basis of Therapeutics, Ninth Edition, pp.1459-1483. Suitable exemplary corticosteroids include cortisol (hydrocortisone); tetrahydrocortisol; prednisone (cortisone); prednisolone (cortisol); 6 α -methylprednisolone; fludrocortisone (9 α -fluorocortisol); 11-desoxycortisol; cortisone (11-dehydrocortisol); corticosterone;
- 10 triamcinolone (9 α -fluoro-16 α -hydroxyprednisolone); paramethasone (6 α -fluoro-16 α -methylprednisolone); betamethasone (9 α -fluoro-16 β -methylprednisolone); dexamethasone (9 α -fluoro-16 α -methylprednisolone); desoxycorticosterone acetate (doca acetate, percorten acetate); desoxycorticosterone pivalate (percorten pivalate); fludrocortisone acetate (florine acetate); cortisol (hydrocortisone) (cortef, hydrocortone);
- 15 cortisol acetate (cortef acetate, hydrocortone acetate); cortisol cypionate (cortef); cortisol sodium phosphate (hydrocortone phosphate); cortisol sodium succinate (solu-cortef); beclomethasone dipropionate (vanceril); betamethasone (celestone); betamethasone sodium phosphate and acetate (celestone soluspan); betamethasone dipropionate (diprosone); betamethasone valerate (valisone); betamethasone benzoate (benisone, flurodate);
- 20 cortisone acetate (cortone acetate); dexamethasone (decadron, gammacorten); dexamethasone sodium phosphate (decadron phosphate, hexadrol phosphate); dexamethasone acetate (decadron-L.A.); fuprednisolone (alphadrol); meprednisone (betapar); methylprednisolone (medrol); methylprednisolone acetate (depo-medrol,

medrol acetate); methylprednisolone sodium succinate (solu-medrol); paramethasone acetate (haldrone); prednisolone (delta-cortef); prednisolone acetate (meticortelone acetate); prednisolone sodium phosphate (hydeltrasol); prednisolone sodium succinate (meticortelone soluble); prednisolone tebutate (hydelta-T.B.A.); prednisone (deltasone, 5 paracort); triamcinolone (aristocort, kenacort); triamcinolone acetonide (aristoderm, kenalog); triamcinolone diacetate (aristocort diacetate, kienacort diacetate); triamcinolone hexacetonide (aristospan); desonide (tridesilon); desoximetasone (topicort); flumethasone pivalate (locorten); fluocinolone acetonide (fluonid, synalar); fluocinonide (lidex, topsyn); fluorometholone (oxylone); flurandrenolide (cordran); halcinonide (halog); and 10 medrysone (HMS liquifilm, medrocort).

Preferably, the amount of a suitable topical anti-inflammatory or topical anesthetic can be present in about 0.1 wt% to about 99.9 wt% of the composition. Typically, the amount of anesthetic and/or anti-inflammatory present will depend upon the specific anesthetic and anti-inflammatory employed in the composition. In some 15 embodiments of the present invention, the anesthetic and/or anti-inflammatory can be up to about 10 wt%, up to about 5 wt%, up to about 2 wt%, up to about 1 wt% or up to about 0.1 wt% of the composition. Additionally, the nature and amount of the anesthetic and/or anti-inflammatory present in the composition should comply with any State and/or Federal guidelines that regulate the use of such compounds (e.g., FDA regulations).

20 The compositions of the present invention may also include known psoriasis treatment agents such as anthralin, also known as dithranol, vitamin D such as calcipotriene or calcitriol, and vitamin A such as tazarotene. Some of the known psoriasis treatment agents are described in greater detail in Goodman Gilman, Alfred;

Goodman, Louis S.; Gilman, Alfred; Goodman and Gilman's The Pharmacological Basis of Therapeutics, Ninth Edition, pp.1591-1613.

The following tables provide a summary overview of a few representative compositions that can be prepared in accordance with the present invention and is not
5 intended to limit the present invention:

Serum

Ingredient	Preferred Weight %	Most Preferred Weight %
Sterol Ester	75-99%	80-95%
Penetration Enhancer	0.01-15%	0.1-10%
Film-Forming/Polymeric Agent	0.1-10%	0.5-5%
Antioxidant	0-10%	0.5-5%
Preservative	0-10%	0.5-5%
Vitamin(s)	0-25%	0-15%

Salve

Ingredient	Preferred Weight %	Most Preferred Weight %
Sterol Ester	40-90%	60-85%
Penetration Enhancer	0.01-15%	0.1-10%
Film-Forming/Polymeric Agent	0.1-10%	0.5-5%
Viscosity Enhancing Agent	5-40%	10-25%
Antioxidant	0-10%	0.5-5%
Preservative	0-10%	0.5-5%
Vitamin(s)	0-25%	0-15%

Cream/Lotion/Gel

Ingredient	Preferred Weight %	Most Preferred Weight %
Sterol Ester	(30-80%)*	(40-70%)*
Penetration Enhancer	0.01- 15%	0.1-10%
Film-Forming/Polymeric Agent	0.1-10%	0.5-5%
Viscosity Enhancing Agent	1-40%	3-20%
Solvent	5-85%	10-75%
Emulsifier	0.1-20%	1-10%
Antioxidant	0-10%	0.5-5%
Preservative	0-10%	0.5-5%

*weight percent based upon amount of sterol ester less solvent

Aerosol/Mousse/Foam

Ingredient	Preferred Weight %	Most Preferred Weight %
Sterol Ester	(30-75%)*	(35-60%)*
Penetration Enhancer	0.001- 10%	0.002-5%
Film-Forming/Polymeric Agent	0.001-10%	0.005-5%
Viscosity Enhancing Agent	0.001-30%	0.005-20%
Solvent	50-95%	60-90%
Emulsifier	0.1-20%	1-10%
Propellant	0.5-25%	1-10%
Antioxidant	0-10%	0.001-5%

Preservative	0-10%	0.001-5%
--------------	-------	----------

*weight percent based upon amount of sterol ester less solvent

EXAMPLES

The following are provided by way of example only and are by no means intended to be limiting.

5 EXAMPLE 1

A serum with the following composition is prepared:

	SUPER STEROL LIQUID	91.5 wt%
	Shea Butter	1.0 wt%
10	Lauryl Laurate	2.5 wt%
	POLYDERM PPI-SA	1.0 wt%
	Oleic Acid	1.0 wt%
	Vitamin-A Palmitate	1.0 wt%
	Vitamin –E Acetate	1.0 wt%
15	<u>Phenoxyethanol</u>	<u>1.0 wt%</u>

The serum is prepared by adding the SUPER STEROL LIQUID, shea butter, POLYDERM PPI-SA, oleic acid and lauryl laurate to a suitable mixer and heating the mixture at 75-85°C for about 30 minutes until a clear solution is obtained. The SUPER
20 STEROL LIQUID is commercially available from Croda Chemicals Europe, Ltd., of East Yorkshire, England and is a mixture of C₁₀-C₃₀ cholesterol and lanosterol esters. POLYDERM PPI-SA is a Di-PEG-2 Soyamine/IPDI copolymer commercially available from Alzo International Inc.

The resulting composition is removed from the heat and the stirring is continued.
25 Once the temperature of the composition is 50°C or lower, vitamin A palmitate, vitamin E acetate and phenoxyethanol are added to the composition while stirring is maintained.

Once the composition obtains room temperature, the serum composition is packaged in a pump bottle.

EXAMPLE 2

A salve with the following composition is prepared:

5	SUPER STEROL SOLID	75 wt%
	Shea Butter	15 wt%
	Lauryl Laurate	2.5 wt%
	POLYDERM PPI-SA	2.5 wt%
10	Oleic Acid	1.0 wt%
	Vitamin-A Palmitate	1.5 wt%
	Vitamin –E Acetate	1.5 wt%
	Phenoxyethanol	1.0 wt%

15 The above salve is prepared by mixing the above ingredients in a suitable mixer by the procedure outlined in Example 1 and filling the slave into a suitable tube for dispensing by the patient.

EXAMPLE 3

A cream is with the following composition is prepared:

20	SUPER STEROL LIQUID	18.3* wt%
	Shea Butter	0.2 wt%
	Lauryl Laurate	0.5 wt%
	POLYDERM PPI-SA	0.2 wt%
25	Oleic Acid	0.2 wt%
	Vitamin-A Palmitate	0.2 wt%
	Vitamin –E Acetate	0.2 wt%
	Phenoxyethanol	0.2wt%
	POLAWAX™	10.0 wt%
30	Water	68.5 wt%
	Cetyl Alcohol	1.4 wt %
	DOWICIL™-200	0.1 wt%

*58.1% less the solvent (water)

The cream is prepared by adding the serum of Example 1, the POLAWAX™, water and cetyl alcohol to a suitable mixer and heating the mixture at 75-85°C for about 20-30 minutes. POLAWAX™ is a higher fatty alcohol self-emulsifying wax commercially available from Croda Chemicals Europe, Ltd., of East Yorkshire, England.

5 DOWICIL™-200 is a 1-3-chloroallyl-3,5,7-triaza-1-azoniaadamantane chloride preservative/anti-microbial compound commercially available from Dow Chemical Co. The resulting composition is removed from the heat and the stirring is continued. Once the solution obtains room temperature, the DOWICIL™-200 is added and the cream is packaged in a tube, jar, bottle or other suitable container.

EXAMPLE 4

A foam with the following composition is prepared:

	SUPER STEROL LIQUID	9.15* wt%
	Shea Butter	0.01 wt%
15	Lauryl Laurate	0.025 wt%
	POLYDERM PPI-SA	0.01 wt%
	Oleic Acid	0.01 wt%
	Vitamin-A Palmitate	0.01 wt%
	Vitamin -E Acetate	0.01 wt%
20	Phenoxyethanol	0.01wt%
	Glycerox 767	10.0 wt%
	Water	80.0 wt%

*45.75% less solvent (water

25 The above foam composition is prepared by adding the serum of Example 1, GLYCEROX 767 and water to a suitable mixer and heating the mixture at 75°C for about 10 minutes. GLYCEROX 767 is a PEG-6 capric/caprylic glycerides commercially available from Croda Chemicals Europe, Ltd., of East Yorkshire, England. The resulting composition is removed from the heat and stirred until it obtains room temperature. The

composition is filled into a sealed aerosol container along with A-46 propellant (15.2% propane/84.8% isobutene) commercially available from Diversified CPC International. The final aerosol container comprises approximately 96.5% of the above foam composition and 3.5% of the propellant with a pressure of about 46 PSIG @ 75°F.

5

EXAMPLE 5

The serum prepared in Example 1 was applied 14 patients suffering from psoriasis. The treatment regimen and results are summarized in the TABLE 1.

TABLE 1

<u>Subject</u>	<u>Age</u>	<u>Sex</u>	<u>Affected Area</u>	<u>Tx Duration</u>	<u>Tx</u>	<u>Previous Treatment</u>	<u>Result</u>
01	13 y.o.	F	- right knee - behind right knee - behind left knee	1 day	One treatment	- AQUAPHOR [®] (OTC) used for dry skin - hydrocortisone 1%	Complete resolution.
02	41 y.o.	M	- upper back - lower back - chest - small area on face - stomach.	10 months	Pt. applied serum, exfoliator, and wash 3x / wk at home	Pt. declined monoclonal antibody tx at Mayo Clinic and other options at the Rockefeller University to try the serum.	Pt. recovered 100% but still has residual pre-existing scars.
03	41 y.o.	M	- left forearm - left elbow - left wrist - left hand - right forearm - stomach - back of leg	14 months	Pt. applied serum everyday at home in addition to application on each clinical visit	Pt. declined i.v. clinical trial tx at Mayo Clinic to try the serum.	Left and right forearm, left hand, and back of right leg improved 100%. Stomach, left wrist and left elbow improved 30-40%.
04	56 y.o.	M	- left knee - right knee - back of right foot	1 day	One treatment	none	Complete resolution.
05	46 y.o.	M	- left knee	1 day	One treatment	- coco butter	Complete resolution.
06	Tx at 6 m.o.	M	- right foot - right leg - left foot - left leg - left buttock	1 day	One treatment	- ocean salt water	Complete resolution.

TABLE 1 (cont.)

<u>Subject</u>	<u>Age</u>	<u>Sex</u>	<u>Affected Area</u>	<u>Tx Duration</u>	<u>Tx</u>	<u>Previous Treatment</u>	<u>Result</u>
07	33 y.o.	M	- right ankle - left arm near elbow	1 month	Applied serum 1x / day	none	Complete resolution of symptoms. Partial resolution of physical appearance of lesions.
08	56 y.o.	F	- left hand - left wrist - left forearm - left elbow - right hand - right wrist - right elbow - heel of left foot - heel of right foot - bottom of left foot	7 months	Applied serum 1x / day	- HUMIRA [®] - SORIATANE [®] - light therapy - laser treatment	80-100% resolution involving all lesions.
09	63 y.o.	M	- left knee - left forearm near elbow - right knee - right forearm near elbow	6 months	Applied serum 1x / day	none	90-100% resolution involving all lesions.
10	53 y.o.	F	- left knuckles - left elbow - left ankle - right knuckles - right elbow	1 day	One treatment	- VASELINE [®]	Reported benefit, however, pt. was unable to follow-up due to transportation issues.

TABLE 1 (cont.)

<u>Subject</u>	<u>Age</u>	<u>Sex</u>	<u>Affected Area</u>	<u>Tx Duration</u>	<u>Tx</u>	<u>Previous Treatment</u>	<u>Result</u>
11	60 y.o.	F	<ul style="list-style-type: none"> - left hand - left arm - left foot - left ankle - left leg - right hand - right arm - right foot - right ankle - right leg - upper back - lower back 	1 day	One treatment	<ul style="list-style-type: none"> - topical triamcinolone acetate 0.1% 2x / day - steroid injections - prednisone - topical corticosteroid - lotion 	80% resolution.
12	39 y.o.	F	<ul style="list-style-type: none"> - right elbow - right arm - right knee - right shin - left elbow - left forearm near elbow - left knee - left shin - stomach 	3 weeks	Applied serum 1x / day for 3 wks	- artificial tanning with UV lights	80% resolution of symptoms. 50% resolution of physical appearance.
13	48 y.o.	M	- facial	3-6 months	Applied serum 1x / day	none	Complete resolution.
14	37 y.o.	M	- upper torso/neck	1-3 months	Applied serum 1x / day	none	Complete resolution.

EXAMPLE 6

One male patient has reported relief of sunburn pain after a single application of the serum prepared in Example 1. The serum was applied to the sunburnt area in the evening prior to bedtime and the patient reported relief when he awoke in the morning.

5 While certain preferred and alternative embodiments of the present invention have been set forth for purposes of disclosing the invention, modifications to the disclosed embodiments may occur to those who are skilled in the art. Accordingly, the appended claims are intended to cover all embodiments of the invention and modifications thereof which do not depart from the spirit and scope of the invention.

10 All documents, patents and other literature referred to herein are incorporated by reference in their entirety.

The term “comprising” as used in the following claims is an open-ended transitional term that is intended to include additional elements not specifically recited in the claims.

The term “consisting essentially of” as used in the following claims is a partially closed
15 transitional phrase and is intended to include the recited elements plus any unspecified elements that do not materially affect the basic and novel characteristics of the claims. For example, an adhesive laminate (the outermost layer of the applied patch) embossed or printed with indicia would still be included in the meaning of “consisting essentially of”, even if not specifically recited. The term “consisting of” as used in the following claims is
20 intended to indicate that the claims are restricted to the recited elements.

It should be noted that it is envisioned that any feature or element that is positively identified in this document may also be specifically excluded as a feature or element of an embodiment of the present invention.

CLAIMS:

1. A composition for topical administration comprising:
 - (a) 25% or more of a sterol ester based upon the weight of the composition less any solvent that may be present with a boiling point of 150°C or less;
 - 5 (b) a penetration enhancer; and
 - (c) optionally a further additive selected from the group consisting of solvents, film-forming/polymeric agents, viscosity increasing agents, emulsifiers, antioxidants, pH adjusting agents, preservatives, vitamins, propellants and combinations of the foregoing.
- 10 2. The composition of claim 1 wherein the sterol ester comprises 30% or more based upon the total weight of the composition less any solvent that may be present with a boiling point of 150°C or less.
3. The composition of claim 1 wherein the sterol ester comprises 40% or more based upon the total weight of the composition less any solvent that may be
15 present with a boiling point of 150°C or less.
4. The composition of claim 1 wherein the sterol ester is a C₁₀-C₃₀ carboxylic acid sterol ester.
5. The composition of claim 4 wherein the sterol ester is a C₁₀-C₃₀ carboxylic acid cholesterol/lanosterol mixture.
- 20 6. The composition of claim 1 which is a gel, cream, lotion, salve, serum, aerosol, mousse or foam and comprises a film-forming agent.

7. The composition of claim 1 which is a gel, cream, lotion, salve, serum, aerosol, mousse or foam and comprises a film-forming agent and a viscosity increasing agent.
8. The composition of claim 1 which is a gel, cream, lotion, aerosol, mousse or foam
5 and comprises a film-forming agent, a viscosity increasing agent and a solvent.
9. The composition of claim 1 which is an aerosol, mousse or foam and comprises a film-forming agent, a viscosity increasing agent, a solvent and a propellant.
10. A method for treating psoriasis comprising the topical administration of a composition to a psoriatic area of a patient at least once a day wherein the
10 composition comprises:
- (a) 25% or more of a sterol ester based upon the weight of the composition less any solvent that may be present with a boiling point of 150°C or less;
 - (b) a penetration enhancer; and
 - (c) optionally a further additive selected from the group consisting of solvents,
15 film-forming/polymeric agents, viscosity increasing agents, emulsifiers, antioxidants, pH adjusting agents, preservatives, vitamins, propellants and combinations of the foregoing.
11. The method of claim 10 wherein the sterol ester comprises 30% or more based upon the total weight of the composition less any solvent that may be present with
20 a boiling point of 150°C or less.

12. The method of claim 10 wherein the sterol ester comprises 40% or more based upon the total weight of the composition less any solvent that may be present with a boiling point of 150°C or less.
13. The method of claim 10 wherein the sterol ester is a C₁₀-C₃₀ carboxylic acid sterol ester.
14. The method of claim 13 wherein the sterol ester is a C₁₀-C₃₀ carboxylic acid cholesterol/lanosterol mixture.
15. The method of claim 10 wherein the composition is a gel, cream, lotion, salve, serum, aerosol, mousse or foam and comprises a film-forming agent.
16. The method of claim 10 wherein the composition is a gel, cream, lotion, salve, serum, aerosol, mousse or foam and comprises a film-forming agent and a viscosity increasing agent.
17. The method of claim 10 wherein the composition is a gel, cream, lotion, aerosol, mousse or foam and comprises a film-forming agent, a viscosity increasing agent and a solvent.
18. The method of claim 10 wherein the composition is an aerosol, mousse or foam and comprises a film-forming agent, a viscosity increasing agent, a solvent and a propellant.
19. The method of claim 10 wherein the composition is applied to the psoriatic area more than once a day.
20. The method of claim 10 further comprising the step of washing and drying the psoriatic area prior to application of the composition.

21. The method of claim 10 further comprising the step of covering the composition and psoriatic area with a protective cloth, bandage, gauze or wrap after application of the composition.

22. A composition for topical administration comprising:

- 5 (a) 25% or more of a sterol ester based upon the weight of the composition less any solvent that may be present with a boiling point of 150°C or less;
- (b) a solvent;
- (c) a propellant;
- (d) a penetration enhancer;
- 10 (e) a viscosity increasing agent; and
- (f) optionally a further additive selected from the group consisting of a co-solvent, film-forming/polymeric agents, emulsifiers, antioxidants, pH adjusting agents, preservatives, vitamins and combinations of the foregoing wherein the composition is an aerosol, mousse or foam.

15 23. The composition of claim 22 wherein the sterol ester comprises 30% or more based upon the total weight of the composition less any solvent that may be present with a boiling point of 150°C or less.

24. The composition of claim 22 wherein the sterol ester comprises 40% or more based upon the total weight of the composition less any solvent that may be
20 present with a boiling point of 150°C or less.

25. The composition of claim 22 wherein the sterol ester is a C₁₀-C₃₀ carboxylic acid sterol ester.

26. The composition of claim 22 wherein the sterol ester is a C₁₀-C₃₀ carboxylic acid cholesterol/lanosterol mixture.

INTERNATIONAL SEARCH REPORT		International application No. <div style="text-align: center;">PCT/US 2012/059701</div>	
A. CLASSIFICATION OF SUBJECT MATTER		<i>A61K 9/107 (2006.01)</i> <i>A61K 9/12 (2006.01)</i> <i>A61K 31/575 (2006.01)</i> <i>A61P 17/06 (2006.01)</i>	
According to International Patent Classification (IPC) or to both national classification and IPC			
B. FIELDS SEARCHED			
Minimum documentation searched (classification system followed by classification symbols)			
A61K 9/107, 9/12, 31/56, 31/575, A61P 17/00, 17/06			
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched			
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)			
PatSearch (RUPTO internal), EAPATIS, Esp@cenet, WIPO, RUPTO, USPTO, Scopus, Scirus			
C. DOCUMENTS CONSIDERED TO BE RELEVANT			
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	
X	US 6153209 A (THE PROCTER & GAMBLE COMPANY) 28.11.2000, claims 10, 11, 14, 15, 24, col. 18, lines 30, 31, col. 27, lines 14-15, col. 28, line 16	1-9, 22-26	
X Y	US 2007/0110685 A1 (BENJAMIN A. AUSPITZ et al.) 17.05.2007, paragraphs [0002], [0003], [0044], [0158] - [0160], [0164]	10-19 20, 21	
Y	RU 2091065 C1 (IVANOV ANATOLIY ALEKSEEVICH) 27.09.1997, p. 4, lines 13-16	20	
Y	RU 2093190 C1 (LABORATOIRES D' HYGIENE ET DE DIETETIQUE L.H.D.) 20.10.1997, p. 3, line 29	21	
Y	RU 2123334 C1 (TIKHOOKEANSKY NAUCHNO-ISSLEDOVATELSKY RYBOKHOZIAISTVENNY TSENTR et al.) 20.12.1998, claim 1	21	
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.			
* Special categories of cited documents:		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"A" document defining the general state of the art which is not considered to be of particular relevance		"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"E" earlier document but published on or after the international filing date		"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)		"&" document member of the same patent family	
"O" document referring to an oral disclosure, use, exhibition or other means			
"P" document published prior to the international filing date but later than the priority date claimed			
Date of the actual completion of the international search <div style="text-align: center;">06 May 2013 (06.05.2013)</div>		Date of mailing of the international search report <div style="text-align: center;">04 July 2013 (04.07.2013)</div>	
Name and mailing address of the ISA/ FIPS Russia, 123995, Moscow, G-59, GSP-5, Berezhkovskaya nab., 30-1		Authorized officer <div style="text-align: center;">A. Kilinskaya</div>	
Facsimile No. +7 (499) 243-33-37		Telephone No. (495)531-64-81	