

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 November 2011 (17.11.2011)

(10) International Publication Number
WO 2011/143106 A1

(51) International Patent Classification:

C07D 471/04 (2006.01) *A61P 29/00* (2006.01)
A61K 31/437 (2006.01) *A61P 37/08* (2006.01)

Market, Seattle, WA 98107 (US). **STEVENS, Kirk, L.** [US/US]; 24102 Lockwood Road, Bothell, WA 98021 (US). **VAN VELDHUIZEN, Josh** [US/US]; 417 N. Bowdoin Pl., Seattle, WA 98103 (US).

(21) International Application Number:

PCT/US2011/035741

(74) Agents: **ECK, Steven, R.** et al.; Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, CA 94404 (US).

(22) International Filing Date:

9 May 2011 (09.05.2011)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (for all designated States except US):
GILEAD SCIENCES, INC. [US/US]; 333 Lakeside Drive, Foster City, CA 94404 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and



(75) Inventors/Applicants (for US only): **BAKER, William, R.** [US/US]; 4759 171 Street SE, Bellevue, WA 98006 (US). **CAI, Shaopei** [US/US]; 3240 29th Ave. W, Seattle, WA 98199 (US). **KAPLAN, Joshua, Aaron** [US/US]; 2253 Gilman Drive W., Seattle, WA 98119 (US). **KIM, Musong** [US/US]; 2610 194th St. SE, Bothell, WA 98012 (US). **PHILLIPS, Gary** [US/US]; 22711-SE 51st Street, Issaquah, WA 98029 (US). **PURVIS, II, Lafe, J.** [US/US]; 906 Dexter Avenue N. Apt. #L-120, Seattle, WA 98109 (US). **STASIAK, Marcin** [US/US]; 358 NW

Published:

— with international search report (Art. 21(3))

[Continued on next page]

(54) Title: BI - FUNCTIONAL PYRAZOLOPYRIDINE COMPOUNDS



(57) Abstract: invention provides compounds of the formula: wherein R¹ and R² together with the phenyl to which they are bound may form a bicyclic, fused heterocyclic ring and all other variables are as defined herein, their use in pulmonary inflammation or bronchoconstriction therapy and compositions comprising and processes for preparing the same are provided.

WO 2011/143106 A1



-
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

BI-FUNCTIONAL PYRAZOLOPYRIDINE COMPOUNDS

Field of the Invention

The present invention relates to novel anti-inflammatory, bronchodilator
5 compounds, compositions containing the same, therapeutic methods and uses
for the same and processes for preparing the same.


Background of the Invention

The chronic inflammatory processes underlie respiratory diseases such as
10 Chronic Obstructive Pulmonary Disease (COPD) and asthma. These diseases
involve active inflammation in the bronchial airways, parenchyma and
pulmonary vasculature of the lungs. The inflammatory process in such
diseases are characterized by increased numbers of activated immune cells
such as macrophages, neutrophils, eosinophils and lymphocytes and the
15 release of a range of pro-inflammatory signaling molecules, namely cytokines
and chemokines from immune and resident lung cells. The pathogenesis of
these diseases is different but chronic inflammation is an underlying driving
mechanism to both. COPD is strongly associated with exposure to noxious
particles and gases from the external environment such as cigarette smoking
20 and exposure to wood burning fires and is characterized by oxidative stress
and an imbalance of harmful tissue proteinases with anti-proteases. These
processes can lead to distinctive pathologies such as goblet metaplasia and
mucus hypersecretion which cause bronchitis, alveolar wall destruction leading
to emphysema and inappropriate tissue repair and smooth muscle thickening
25 causing small airways remodeling (reviewed by Molfino & Jeffery, *Pulm.
Pharmacol. Ther.* 2007;20:462-72). In asthma, allergic immune mechanisms
underlie the chronic inflammatory processes which contributes to airway
hyperresponsiveness and structural changes in the bronchial airway, termed
remodeling, such as airway smooth muscle thickening and goblet cell
30 hyperplasia. (reviewed by Hamid & Tulic, *Annu. Rev. Physiol.* 2009; 71:489-
507).

Bronchodilator medications that can improve lung function and improve expiratory flow are used as a standard of care for symptom relief in the treatment of respiratory diseases. Inhaled long-acting β_2 adrenoceptor agonists (LABA) such as salmeterol or formoterol, or inhaled long-acting 5 muscarinic receptor antagonists (LAMA) such as tiotropium are commonly prescribed to provide symptom relief.

Inflammation is a central process underlying many respiratory diseases and treatments that are anti-inflammatory may be efficacious and have the potential 10 to impact disease progression. The phosphodiesterase-4 (PDE4) enzyme is a ubiquitously expressed enzyme that is responsible for catalyzing the hydrolysis of cyclic adenosine monophosphate (cAMP). Inhibition of the enzymatic activity of PDE4 with use of selective inhibitors elevates cellular levels of cAMP and this has anti-inflammatory effects in multiple immune and resident pulmonary 15 cell types (Spina, *Brit. J. Pharmacol.* 2008;155:308-15). Use of the oral PDE4 inhibitor roflumilast has demonstrated anti-inflammatory activity clinically showing a reduction of exacerbations and modest increases in lung function in COPD patients (Rabe *et al.*, *Lancet* 2005; 366:563-71; Calverly *et al.*, *Am. J. Respir. Crit. Care Med.* 2007; 176:154-61). Additionally, roflumilast improves 20 lung function in severe and symptomatic patients with COPD treated with salmeterol or tiotropium but remains dose-limited due to side-effects including nausea, head-ache, diarrhea, and weight loss (Fabbri *et al.*, *Lancet* 2009; 374:695-703; Calverly *et al.*, *Lancet* 2009; 374:685-94). Forest Research Institute's product Daxas (roflumilast) has been approved as a once daily oral 25 PDE4 inhibitor for the treatment of COPD. While it was accepted that Daxas demonstrated consistent evidence of efficacy concerns over a number of adverse event signals led the committee to deny approval based on the overall poor risk-to-benefit ratio. Topical delivery of a PDE4 inhibitor could therefore provide efficacious anti-inflammatory activity in the lungs whilst reducing the 30 potential for side-effects by limiting its exposure to the systemic circulation. Additionally, direct topical delivery may allow for higher local concentrations of the PDE4 inhibitor than could be achieved through oral dosing, and thus potential for further improvement in anti-inflammatory efficacy.



PCT Publication No. WO2005/090348 to GSK relates to compounds of formula (I) and pharmaceutically acceptable salts thereof as PDE4 Inhibitors:



5 wherein (*inter alia*):



W is Ar, -CR⁴R⁵Ar or a group (y) or (y1) wherein:

Ar is



10 wherein A, B, D, E, and F are C-R⁶, N or N+-O- and R⁶ is *inter alia* H, halo, C₁₋₆alkyl, ... R⁷R⁸N-S(O)₂-, ... R⁷R⁸N-CO-, ... R⁷R⁸N-, ... -CH₂-NR⁷R⁸, -CH₂-CH₂-NR⁷R⁸, ... -CH₂-CO-NR⁷R⁸, ... and all other variables are as defined therein.

15 PCT Publication No. WO2007/023110 to Hoffmann-La Roche relates to P38 Map Kinase Inhibitors of formula Ia or Ib:



wherein (*inter alia*):

R¹ is aryl or heteroaryl;

R⁴ is H, C₁₋₆alkyl, hydroxyl, amino, hetero-C₁₋₆alkyl, hetero-C₁₋₆alkoxy, hetero-C₁₋₆alkylamino, heterocyclyl, heterocyclyl-C₁₋₆alkyl, hydroxyC₃₋₆cycloalkyl, C₃₋₆cycloalkyl-C₁₋₆alkyl, C₁₋₆alkylsulfonyl, C₁₋₆alkylsulfonamido, aryl, heteroaryl, aryl-C₁₋₆alkyl, heteroaryl-C₁₋₆alkyl, C₁₋₆alkoxy, heteroaryl-C₁₋₆alkoxy, -(CHR^b)_r-C(=O)-R^c, -(CHR^b)_r-O-C(=O)-R^c, -(CHR^b)_r-NH-C(=O)-R^c or -SO₂-R^c;


X and Y are N or one of X and Y is N and the other is CR^d;

W is a bond, O, S(O)_m, CH₂, or NR^f;


A is O, CH₂, S(O)_m, C(=O), NR^h, or CH(OR^h);

and all other variables are as defined therein.

PCT Publication No. WO2008/015416 to Glaxo Group Limited relates to PDE4 inhibitors of formula (I):




wherein Ar is



wherein:


Q¹ is NH or NMe and Q² is -C(O)-, -S(O)₂-, -C(O)NH or -C(O)NMe-, or
 Q¹ is a bond or -O- and Q² is a bond, or
 Q¹ is -C(O)- and Q² is -NH or NMe, or
 Q¹ is -S(O)₂-, and Q² is -NH, NMe, or a bond;
 L is (CH₂)_n or -(CH₂)_m-O-(CH₂)_m-;
 R⁵ is H, methyl, ethyl, n-propyl, isopropyl, -CH₂CH₂OH, -CH₂CH(Me)OH, -CH₂CH₂CH₂OH, -CH₂CH₂OMe, or -CH₂CH₂CH₂OMe;
 and
 R⁶ is C₁₋₄alkyl or C₁₋₄alkyl substituted by on OH or OC₁₋₃alkyl;
 and all other variables are as defined therein.

PCT Publication No. 2008/015437 to Glaxo Group Limited relates to PDE4 inhibitors of formula (I):



5 wherein:





Ar is



wherein:

- Q^1 is NH or NMe and Q^2 is $-C(O)-$, $-S(O)_2-$, $-C(O)NH$ or $-C(O)NMe-$, or
- 10 Q^1 is a bond or $-O-$ and Q^2 is a bond, or
- Q^1 is $-C(O)-$ and Q^2 is -NH or NMe, or
- Q^1 is $-S(O)_2-$, and Q^2 is $-NH$, NMe, or a bond;
- Q^3 is a bond NH or NMe;
- L is $(CH_2)_n$ or $-(CH_2)_m-O-(CH_2)_m-$;
- 15 R^5 is H, methyl, ethyl, n-propyl, isopropyl, $-CH_2CH_2OH$, $-CH_2CH(Me)OH$,
 $-CH_2CH_2CH_2OH$, $-CH_2CH_2OMe$, or $-CH_2CH_2CH_2OMe$; and
- R^6 is C_{1-4} alkyl or C_{1-4} alkyl substituted by one OH or OC_{1-3} alkyl;
and all other variables are as defined therein.

- 20 PCT Publication Nos. WO2009/100166, WO2009/100167, WO2009/100169
and WO2009/0197871, all to Glaxo Group Limited relate to dual
pharmacophores-PDE4-Muscarinic antagonists of the following Formulas (I):




5 wherein all variables are as defined in each publication.

Conventional therapeutic agents for the treatment of inflammatory respiratory conditions suffer from limited efficacy and undesired side-effect profiles.

Accordingly, there remains a need in the art for new drugs designed to treat
10 respiratory conditions including inflammatory respiratory conditions such as asthma, COPD, chronic bronchitis, bronchiectasis, cystic fibrosis, etc.

Summary of the Invention


15 As one aspect, the present invention provides compounds Formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

X is selected from:



5

R¹ is CH₂OH, CH₂CH₂OH, N(H)C(O)H, N(H)S(O₂)CH₃, and R² is H;

or R¹ and R² together with the phenyl to which they are bound form a bicyclic,

10 fused heterocyclic ring having 9 or 10 ring atoms wherein 1 or 2 ring atoms are selected from N, O and S and the remaining ring atoms are C, wherein said bicyclic fused heterocyclic ring is optionally substituted with one, two or three additional substituents independently selected from alkyl, oxo and OH;

R³ is selected from C₄₋₁₂alkylene, C₄₋₁₂alkenylene, C₄₋₁₂alkynylene, R⁴-O-R⁴, R⁴-N(R⁸)-R⁴, C₃₋₆cycloalkylene, R⁴-C₃₋₆cycloalkylene,

15 C₃₋₆cycloalkylene-R⁴, R⁴-C₃₋₆cycloalkylene-R⁴, C₆₋₁₀arylene,

R⁴-C₆₋₁₀arylene, C₆₋₁₀arylene-R⁴, R⁴-C₆₋₁₀arylene-R⁴,

R⁴-C₆₋₁₀arylene-O-R⁴, R⁴-C₆₋₁₀arylene-N(R⁸)-R⁴,

R⁴-C₆₋₁₀arylene-C₆₋₁₀arylene, Het, R⁴-Het, R⁴-Het-C₆₋₁₀arylene, Het-R⁴,

R⁴-Het-R⁴, R⁴-O-Het R⁴-C₆₋₁₀arylene-O-Het, R⁴-C₆₋₁₀arylene-C(O)-Het,

20 and R⁴-C₆₋₁₀arylene-N(R⁸)-Het;

wherein said alkylene, alkenylene, alkynylene, cycloalkylene, or arylene is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸;

Het is 5-6 membered saturated or unsaturated heterocycle wherein 1

25 or 2 ring atoms are selected from N, O and S, and wherein said

heterocyclene is optionally substituted with 1, 2 or 3 substituents selected from halo, alkyl, alkoxy, oxo and OH;


R^4 is C_{1-10} alkylene, C_{2-10} alkenylene, or C_{2-10} alkynylene wherein each R^4 is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR^8 ; with the proviso that the total number of carbon atoms in the C_{1-10} alkylene, C_{2-10} alkenylene, or C_{2-10} alkynylene chains of two R^4 groups in any definition of R^3 is not greater than 12;

Y is $C(O)$, $C(O)N(R^8)CH_2$, $N(R^8)C(O)$, $O-C(O)N(R^8)CH_2$, $N(R^8)C(O)N(R^8)CH_2$,
10 or $SO_2N(R^8)CH_2$;

R^5 is alkyl;

R^6 is H or alkyl;

R^7 is selected from unsubstituted C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, and a heterocyclic group selected from formulas (i), (ii), and (iii):



15


wherein Z is O, S, $S(O)_2$, NH or $N-R^{7a}$, and

R^{7a} is selected from alkyl, $C(O)alkyl$, $C(O)NH_2$, $C(O)N(H)alkyl$, and
 $C(O)N(alkyl)_2$; and

R^8 is H or alkyl.

20

In one embodiment, the present invention provides compounds of Formula I':



and pharmaceutically acceptable salts thereof, wherein all variables are as

25 defined above.

As another aspect, the present invention provides a composition comprising a compound of Formula I or I' or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier, diluent or excipient. In one embodiment, the composition is suitable for inhalation.

5

As another aspect, the present invention provides a method comprising administering to a human, an effective amount of a compound of Formula I or I' or a pharmaceutically acceptable salt thereof.

- 10 As another aspect, the present invention provides a method for treating pulmonary inflammation or bronchoconstriction in a human in need thereof. The method comprises administering to the human an effective amount of a compound of Formula I or I' or a pharmaceutically acceptable salt thereof.
- 15 As another aspect, the present invention provides a method for treating a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associated bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human in need thereof. The method comprises administering to the human an effective amount of a compound of Formula I or I' or a pharmaceutically acceptable salt thereof. In one embodiment, the
- 20
- 25

- present invention provides a method for treating chronic obstructive pulmonary disease (COPD) or asthma in a human in need thereof using a compound of Formula I or I' or a pharmaceutically acceptable salt thereof.

- 30 As another aspect, the present invention provides a compound of Formula I or I' or a pharmaceutically acceptable salt thereof for use as a medicament.

As another aspect, the present invention provides a compound of Formula I or I' or a pharmaceutically acceptable salt thereof for use in a method of treating of pulmonary inflammation or bronchoconstriction in a human.

- 5 As another aspect, the present invention provides a compound of Formula I or I' or a pharmaceutically acceptable salt thereof for use in a method of treating a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis,
- 10 chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associate bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human. In one embodiment, the present invention provides a compound of Formula I or I' or a pharmaceutically acceptable salt
- 15 thereof for use in a method of treating a disease associated with chronic obstructive pulmonary disease (COPD) or asthma, in a human.

As another aspect, the present invention provides the use of a compound of Formula I or I' or a pharmaceutically acceptable salt thereof for the

- 20 manufacture of a medicament for the treatment of pulmonary inflammation or bronchoconstriction in a human.

As another aspect, the present invention provides the use of a compound of Formula I or I' or a pharmaceutically acceptable salt thereof for the

- 25 manufacture of a medicament for the treatment of a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis,
- 30 transplant-associate bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human.

As another aspect, the present invention provides a composition comprising a compound of Formula I or I' or a pharmaceutically acceptable salt thereof for use in the preparation of a medicament for the treatment of pulmonary inflammation or bronchoconstriction in a human.


5

As another aspect, the present invention provides a composition comprising a compound of Formula I or I' or a pharmaceutically acceptable salt thereof for use in the preparation of a medicament for the treatment of a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associated bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human.

20

Detailed Description of the Invention


One aspect provided are compounds Formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

X is selected from:



25 R¹ is CH₂OH, CH₂CH₂OH, N(H)C(O)H, N(H)S(O₂)CH₃, and R² is H;

or R¹ and R² together with the phenyl to which they are bound form a bicyclic, fused heterocyclic ring having 9 or 10 ring atoms wherein 1 or 2 ring atoms are selected from N, O and S and the remaining ring atoms are C, wherein said bicyclic fused heterocyclic ring is optionally substituted with one, two or three additional substituents independently selected from alkyl, oxo and OH;

R³ is selected from C₄₋₁₂alkylene, C₄₋₁₂alkenylene, C₄₋₁₂alkynylene, R⁴-O-R⁴, R⁴-N(R⁸)-R⁴, C₃₋₆cycloalkylene, R⁴-C₃₋₆cycloalkylene, C₃₋₆cycloalkylene-R⁴, R⁴-C₃₋₆cycloalkylene-R⁴, phenylene, R⁴-phenylene, phenylene-R⁴, R⁴-phenylene-R⁴, R⁴-phenylene-O-R⁴, 10 R⁴-phenylene-N(R⁸)-R⁴, R⁴-phenylene-phenylene, Het, R⁴-Het, R⁴-Het-phenylene, Het-R⁴, R⁴-Het-R⁴, R⁴-O-Het R⁴-phenylene-O-Het, R⁴-phenylene-C(O)-Het, and R⁴-phenylene-N(R⁸)-Het, wherein said alkylene, alkenylene, alkynylene, cycloalkylene, or arylene is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸;

Het is 5-6 membered saturated or unsaturated heterocyclene wherein 1 or 2 ring atoms are selected from N, O and S, and wherein said heterocyclene is optionally substituted with 1, 2 or 3 substituents selected from halo, alkyl, alkoxy, oxo and OH;


R⁴ is C₁₋₁₀alkylene, C₂₋₁₀alkenylene, or C₂₋₁₀alkynylene wherein each R⁴ is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸; with the proviso that the total number of carbon atoms in the C₁₋₁₀alkylene, C₂₋₁₀alkenylene, or C₂₋₁₀alkynylene chains of two R⁴ groups in any definition of R³ is not greater than 12;

Y is C(O), C(O)N(R⁸)CH₂, N(R⁸)C(O), O-C(O)N(R⁸)CH₂, N(R⁸)C(O)N(R⁸)CH₂, or SO₂N(R⁸)CH₂;

R⁵ is alkyl;

30 R⁶ is H or alkyl;

R⁷ is selected from unsubstituted C₃₋₆cycloalkyl, substituted C₃₋₆cycloalkyl, and a heterocyclic group selected from formulas (i), (ii), and (iii):



wherein Z is O, S, S(O)₂, NH or N-R^{7a}, and


R^{7a} is selected from alkyl, C(O)alkyl, C(O)NH₂, C(O)N(H)alkyl, and

C(O)N(alkyl)₂; and


- 5 R⁸ is H or alkyl.

Also provided within the descriptions of each group of compounds herein, and the pharmaceutically acceptable salts thereof, is another set of embodiments wherein R¹ and R² together with the phenyl to which they are bound form a


- 10 bicyclic, fused heterocyclic ring selected from;



Other embodiments provide compounds of the Formula II(a) and Formula II(b):




X is selected from:



5

R^1 is $-CH_2-OH$, $-NH-CHO$, or $-NH-SO_2-C_1-C_3$ alkyl, and R^2 is H; or R^1 and R^2 together with the phenyl ring to which they are bound for a bicyclic, fused heterocyclic ring of the formula:



10

R^4 is C_{1-10} alkylene, C_{2-10} alkenylene, or C_{2-10} alkynylene, wherein each R^4 is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR^8 ;


15 R^5 is alkyl;

R^6 is H or alkyl, and

R⁸ is H or alkyl; or a pharmaceutically acceptable salt thereof.


5

Another embodiment provides compounds of the formula:




wherein

10 X is selected from:



15 R¹ is -CH₂-OH, -NH-CHO, or -NH-SO₂-C₁-C₃ alkyl, and R² is H; or R¹ and R² together with the phenyl ring to which they are bound for a bicyclic, fused heterocyclic ring of the formula:



20 R⁴ is C₁₋₁₀ alkylene, C₂₋₁₀ alkenylene, or C₂₋₁₀ alkynylene, wherein each R⁴ is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸;



R⁵ is alkyl;

R⁶ is H or alkyl, and



25


R⁸ is H or alkyl; or a pharmaceutically acceptable salt thereof.

Additional embodiments are provided by compounds of Formulas IV, V, VI, VII, and VII:




5





wherein, in the compounds of each embodiment,


X is a substituted phenyl ring selected from:



5

R¹ is -CH₂-OH, -NH-CHO, or -NH-SO₂-CH₃, and R² is H; or R¹ and R² together with the phenyl ring to which they are bound for a bicyclic, fused heterocyclic ring of the formula:

10



15


R⁵ is alkyl;

R⁶ is H or alkyl, and

20 R⁸ is H or alkyl; or a pharmaceutically acceptable salt thereof.

Within each group of compounds described herein, or the pharmaceutically acceptable salts thereof, there is also a further embodiment wherein the moiety indicated by the formula:

25



A single molecule for inhaled use which has bifunctional activity as both a long-
 5 acting β_2 adrenoceptor agonist and a PDE4 inhibitor could provide both
 symptom control through bronchodilation and anti-inflammatory activity. Such
 compound would also have the potential to provide additive or synergistic anti-
 inflammatory activity through the complementary interaction of both molecular
 signaling pathways. Beta agonists when binding to a receptor through the
 10 action of G proteins will increase adenylate cyclase activity which causes
 elevation of cellular cyclic AMP. Inhibition of the PDE4 enzyme also serves to
 maintain cellular cAMP levels through inhibition of the enzyme responsible for
 its breakdown. An inhaled molecule that has both β_2 agonist and PDE4
 inhibitory activity may provide additive effects and potentially synergistic anti-
 15 inflammatory activity, and thus could be dose-sparing. Deposition of a single
 bi-functional compound in the lung microenvironment should also maximize the
 opportunity for this molecular interaction to occur compared to a mixture of the
 single agents dosed together. High lung to systemic exposure levels through
 topical delivery allied with long lung retention times will dramatically reduce the
 20 opportunity for side-effects mediated through exposure via the systemic
 circulation to other tissues and organs.

As used herein, the following terms are defined as indicated.

- 25 "A compound of the invention" means a compound of Formula I or a salt,
 particularly a pharmaceutically acceptable salt thereof.
 "A compound of Formula I" means a compound having the structural formula
 designated herein as Formula I or I' (as compounds of Formula I' are a subset

- of compounds of Formula I). Compounds of Formula I include solvates and hydrates (i.e., adducts of a compound of Formula I with a solvent). In those embodiments wherein a compound of Formula I includes one or more chiral centers, the phrase is intended to encompass racemic mixtures, each individual
- 5 stereoisomer including optical isomers (enantiomers and diastereomers) and geometric isomers (cis-/trans-isomerism) and mixtures of stereoisomers. In addition, compounds of Formula I also include tautomers of the depicted formula(s).
- 10 “Alkyl” is a linear or branched hydrocarbon chain of 1 to 8 carbon atoms (i.e., C₁₋₈ alkyl), or typically, 1 to 6 carbon atoms (i.e., C₁₋₆ alkyl), unless the number of carbon atoms is otherwise specified. When the compound of Formula I includes more than one alkyl, the alkyls may be the same or different. Examples of suitable alkyl groups include, but are not limited to, methyl (“Me”),
- 15 ethyl (“Et”), 1-propyl (n-propyl), isopropyl, n-butyl, isobutyl (2-methyl-1-propyl), sec-butyl (2-butyl), tert-butyl (-C(CH₃)₃), n-pentyl, 2-pentyl, 3-pentyl, hexyl, octyl, and the like.
- “Alkenyl” is a linear or branched hydrocarbon chain with at least one site of
- 20 unsaturation, i.e. a carbon-carbon double bond, and having from 2 to 8 carbon atoms (i.e., C₂₋₈ alkenyl), or typically, 2 to 6 carbon atoms (i.e., C₂₋₆ alkenyl) unless the number of carbon atoms is otherwise specified. When the compound of Formula I includes more than one alkenyl, the alkenyls may be the same or different. Examples of suitable alkenyl groups include, but are not
- 25 limited to, ethylene or vinyl (-CH=CH₂), allyl (-CH₂CH=CH₂), 5-hexenyl (-CH₂CH₂CH₂CH₂CH=CH₂), and the like.
- “Alkynyl” is a linear or branched hydrocarbon chain having at least one carbon-carbon triple bond, and optionally also one or more carbon-carbon double
- 30 bonds, and having from 2 to 8 carbon atoms (i.e., C₂₋₈ alkynyl), or more typically 2 to 6 carbon atoms (i.e., C₂₋₆ alkynyl) unless the number of carbon atoms is otherwise specified. When the compound of Formula I includes more than one alkynyl, the alkynyls may be the same or different. Examples of

suitable alkynyl groups include, but are not limited to, acetylenyl (-C≡CH), propargyl (-CH₂C≡CH), and the like.

“Alkylene” refers to a saturated, linear or branched divalent hydrocarbon radical having from 1 to 12 carbon atoms (“C₁₋₁₂ alkylene”), unless the number of carbon atoms is otherwise specified. When the compound of Formula I includes more than one alkylene, the alkynes may be the same or different. Typical alkylene radicals include, but are not limited to, methylene (-CH₂-), ethylene (-CH(CH₃)- or -CH₂CH₂-), propylene (e.g., -CH(CH₂CH₃)-, -CH₂CH(CH₃)- or -CH₂CH₂CH₂-), butylene (e.g., -CH₂CH₂CH₂CH₂-, -C(CH₃)₂CH₂-), and the like. In one embodiment, alkylene is linear. In one embodiment, alkylene is branched.

“Alkenylene” refers to an unsaturated, linear or branched divalent hydrocarbon radical having at least one carbon-carbon double bond, and having from 2 to 12 carbon atoms (“C₂₋₁₂ alkenylene”), unless the number of carbon atoms is otherwise specified. When the compound of Formula I includes more than one alkenylene the alkenylenes may be the same or different. Typical alkenylene radicals include, but are not limited to, 1,2-ethylene (-CH=CH-) and (-CH₂CH=CHCH₂CH₂-). In one embodiment, alkenylene is linear.

20

“Alkynylene” refers to an unsaturated, linear or branched divalent hydrocarbon radical having at least one carbon-carbon triple bond and optionally also one or more carbon-carbon double bonds, and having 2 to 12 carbon atoms (“C₂₋₁₂ alkynylene”), unless the number of carbon atoms is otherwise specified. When the compound of Formula I includes more than one alkenylene the alkenylenes may be the same or different. Typical alkynylene radicals include, but are not limited to, 1,2-ethynylene (-C≡C-) and (-CH₂C≡CCH₂CH₂-). In one embodiment, alkynylene is linear.

30 “Alkoxy” refers to O-alkyl, wherein “alkyl” is as defined above.

“Halo” or “halogen” are synonymous and refer to fluoro, chloro, bromo, and iodo. In one embodiment, halo is fluoro, chloro or bromo.

"Haloalkyl" is linear or branched hydrocarbon chain of 1 to 8 carbon atoms (*i.e.*, C₁₋₈haloalkyl), or typically, 1 to 6 carbon atoms (*i.e.*, C₁₋₆haloalkyl), unless the number of carbon atoms is otherwise specified, substituted by one or more

- 5 halogens, fluoro, chloro, bromo and iodo. Haloalkyl include perhaloalkyls such as trifluoromethyl. When the compound of Formula I includes more than one haloalkyl, the haloalkyls may be the same or different. Examples of suitable haloalkyl groups include, but are not limited to, fluoromethyl, chloromethyl, trifluoromethyl, dichloromethyl, dichloroethyl, and the like.

10

"Oxo" as used herein refers to the group =O attached directly to a carbon atom of a hydrocarbon ring or a C, N or S of a heterocyclic ring to result in oxides, -N-oxides, sulfones and sulfoxides.

- 15 "Cycloalkyl" refers to a monocyclic, saturated or partially unsaturated, non-aromatic ring having from 3 to 6 carbon atoms, (C₃₋₆cycloalkyl) unless a different number of carbon atoms is specified. Examples of specific cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. Cycloalkyl also includes cycloalkyl groups optionally substituted with 1 or 2 substituents, 20 which substituents are the same or different and are selected from halo, alkyl, hydroxyl, O-alkyl, oxo, amino (*e.g.*, NH₂), alkylamino (*e.g.*, N(H)alkyl), dialkylamino (*e.g.*, N(alkyl)₂), C(O)NH₂, C(O)N(H)alkyl, and C(O)N(alkyl)₂, or any subset thereof.


- 25 "Cycloalkylene" refers to a divalent, monocyclic, saturated or partially unsaturated, non-aromatic ring having from 3 to 6 carbon atoms, (C₃₋₆cycloalkylene) unless a different number of carbon atoms is specified. When the compound of Formula I includes more than one cycloalkylene, the cycloalkylene groups may be the same or different. Examples of specific 30 cycloalkylene groups include cyclopropylene, cyclobutylene, cyclopentylene, and cyclohexylene. Cycloalkylene also includes cycloalkyl groups optionally substituted with 1 or 2 substituents, which substituents are the same or different and are selected from halo, alkyl, hydroxyl, O-alkyl, oxo, amino (*e.g.*, NH₂),

alkylamino (e.g., N(H)alkyl), and dialkylamino (e.g., N(alkyl)₂), or any subset thereof. In one embodiment, the cycloalkylene is unsubstituted.

- “Arylene” refers to a divalent, monocyclic or fused bicyclic, aromatic ring having
- 5 from 6 to 10 carbon atoms, (C₆₋₁₀arylene) unless a different number of carbon atoms is specified. When the compound of Formula I includes more than one arylene, the arylene groups may be the same or different. Examples of specific arylene groups include phenylene and naphthylene. Arylene also includes arylene groups optionally substituted with 1 or 2 substituents, which substituents
- 10 are the same or different and are selected from halo, alkyl, hydroxyl, O-alkyl, amino (e.g., NH₂), alkylamino (e.g., N(H)alkyl), and dialkylamino (e.g., N(alkyl)₂), or any subset thereof. In one embodiment, arylene is phenylene. In one embodiment, arylene is unsubstituted phenylene.
- 15 “Heterocyclic group” or “heterocycle” are synonymous and refer to monocyclic and fused bicyclic, saturated or partially unsaturated, or aromatic rings having 5, 6, 9 or 10 ring atoms wherein 1, 2, 3, or 4 ring atoms is/are a heteroatom independently selected from N, O and S and all remaining ring atoms are C. In one embodiment, the heterocyclic group has 5, 6, 9 or 10 rings atoms wherein
- 20 1, 2 or 3 ring atoms is/are a heteroatom independently selected from N, O and S. In all embodiments wherein the heterocyclic group includes 2 or more heteroatoms (N, O and S) the heteroatoms may be the same or different. In all embodiments wherein the compound of Formula I includes 2 or more heterocyclic groups, the heterocyclic groups may be the same or different.
- 25 Examples of heterocyclic groups include but are not limited to furanyl, tetrahydrofuranyl, thiophenyl, tetrahydrothiophenyl, sulfur oxidized tetrahydrothiophenyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, dioxolanyl, oxazolidinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, imidazolinyl, imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, triazolyl, tetrazolyl, pyranyl, dihydropyranol, tetrahydropyranol, pyridyl, dihydropyridyl, piperidyl, dioxanyl, morpholinyl, dithianyl, thiomorpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, triazinyl, indolizinyl, indolyl, isoindolyl, oxindolyl, indolinyl, benzofuranyl, dihydrobenzofuranyl, isobenzofuranyl, benzothienyl, indazolyl,

- benzimidazolyl, benzoxazolinyl, benzoxazolyl, benzisoxazolyl, benzthiazolyl, benzotriazolyl, benzopyranyl, purinyl, quinolizinyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, octahydroisoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl,
 5 naphthyridinyl, pteridinyl, thianaphthalenyl and the like. Heterocyclic groups may be bound through any available ring carbon or ring heteroatom, such as N.

"Heterocyclene" refers to a bivalent heterocyclic group as defined herein. For example, heterocyclenes include:



10

Preferably, the heterocyclene groups in the compounds of Formula I are monocyclic, saturated or partially unsaturated rings having 5 or 6 ring atoms wherein 1, 2, or 3 ring atoms is/are a heteroatom independently selected from N, O and S and all remaining ring atoms are C.

15


The term "optionally substituted" in reference to a particular moiety of the compound of Formula I (e.g., an optionally substituted alkylene) refers to that moiety having no substituents, and that moiety having the specified number of substituents; typically up to 4 substituents unless otherwise indicated. Unless
 20 otherwise indicated, when the term "substituted" is used in conjunction with groups which have multiple available sites for substitution or two or more

moieties capable of substitution, the substituents can be attached to any available C or heteroatom.

Throughout the description and examples, compounds are named using
5 standard IUPAC naming principles where possible.

In some chemical structure representations where carbon atoms do not have a sufficient number of attached variables or bonds depicted to produce a valence of four, the remaining carbon substituents needed to provide a valence of four
10 are understood to be hydrogen. Similarly, in some chemical structures where a bond is drawn without specifying the terminal group, such bond is indicative of a methyl (Me, -CH₃) group, as is conventional in the art.

In one aspect, the invention comprises compounds of Formula I:



15

or a pharmaceutically acceptable salt thereof,

wherein:

R¹ is CH₂OH, CH₂CH₂OH, N(H)C(O)H, N(H)S(O₂)CH₃, and R² is H;

or R¹ and R² together with the phenyl to which they are bound form a bicyclic,

20 fused heterocyclic ring having 9 or 10 ring atoms wherein 1 or 2 ring atoms are selected from N, O and S and the remaining ring atoms are C, wherein said bicyclic fused heterocyclic ring is optionally substituted with one, two or three additional substituents independently selected from alkyl, oxo and OH;

25 R³ is selected from C₄₋₁₂alkylene, C₄₋₁₂alkenylene, C₄₋₁₂alkynylene, R⁴-O-R⁴, R⁴-N(R⁸)-R⁴, C₃₋₆cycloalkylene, R⁴-C₃₋₆cycloalkylene, C₃₋₆cycloalkylene-R⁴, R⁴-C₃₋₆cycloalkylene-R⁴, C₆₋₁₀arylene, R⁴-C₆₋₁₀arylene, C₆₋₁₀arylene-R⁴, R⁴-C₆₋₁₀arylene-R⁴, R⁴-C₆₋₁₀arylene-O-R⁴, R⁴-C₆₋₁₀arylene-N(R⁸)-R⁴,

R^4 -C₆₋₁₀arylene-C₆₋₁₀arylene, Het, R^4 -Het, Het- R^4 , R^4 -Het- R^4 , R^4 -O-Het

R^4 -C₆₋₁₀arylene-O-Het, R^4 -C₆₋₁₀arylene-C(O)-Het, and

R^4 -C₆₋₁₀arylene-N(R^8)-Het,

wherein said alkylene, alkenylene, alkynylene, cycloalkylene, or arylene

5 is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸;

Het is 5-6 membered saturated or unsaturated heterocyclene wherein 1

or 2 ring atoms are selected from N, O and S, and wherein said

heterocyclene is optionally substituted with 1, 2 or 3 substituents

10 selected from halo, alkyl, alkoxy, oxo and OH;


R^4 is C₁₋₁₀alkylene, C₂₋₁₀alkenylene, or C₂₋₁₀alkynylene wherein each R^4 is
optionally substituted with 1, 2 or 3 substituents selected from halo, oxo,
and OR⁸; with the proviso that the total number of carbon atoms in the
C₁₋₁₀alkylene, C₂₋₁₀alkenylene, or C₂₋₁₀alkynylene chains of two R^4

15 groups in any definition of R^3 is not greater than 12; Y is C(O),
C(O)N(R^8)CH₂, N(R^8)C(O), O-C(O)N(R^8)CH₂, N(R^8)C(O)N(R^8)CH₂, or
SO₂N(R^8)CH₂;

R^5 is alkyl;

R^6 is H or alkyl;

20 R^7 is selected from unsubstituted C₃₋₆cycloalkyl, substituted C₃₋₆cycloalkyl, and
a heterocyclic group selected from formulas (i), (ii), and (iii):



wherein Z is O, S, S(O)₂, NH or N-R^{7a}, and

R^{7a} is selected from alkyl, C(O)alkyl, C(O)NH₂, C(O)N(H)alkyl, and


25 C(O)N(alkyl)₂; and

R^8 is H or alkyl.

In one embodiment, the compounds of the invention are defined wherein R¹ is CH₂OH, N(H)C(O)H, or N(H)S(O₂)CH₃, and R² is H. In one particular

30 embodiment the compounds of the invention are defined wherein R¹ is CH₂OH and R² is H.

In one embodiment, the compounds of the invention are defined wherein R¹ and R² together with the phenyl to which they are bound form a bicyclic, fused heterocyclic ring having 9 or 10 ring atoms wherein 1 or 2 ring atoms are selected from N, O and S, and the bicyclic fused heterocyclic ring is optionally substituted with one, two or three additional substituents independently selected from alkyl, oxo and OH. The phrase "one, two or three additional substituents" refers to one, two or three substituents in addition to the -OH indicated in Formula I as being bound to the same phenyl ring as R¹ and R². In one embodiment, wherein R¹ and R² together with the phenyl to which they are bound form a bicyclic, fused heterocyclic ring having 9 or 10 ring atoms wherein 1 or 2 ring atoms are selected from N, O and S, and the bicyclic fused heterocyclic ring is optionally substituted with one additional substituent selected from alkyl, oxo and OH. In one such embodiment, the compounds of the invention are defined wherein R¹ and R² together with the phenyl to which they are bound form



In one embodiment, the compounds of the invention are defined wherein R³ is selected from C₄₋₁₂alkylene, C₄₋₁₂alkenylene, C₄₋₁₂alkynylene, R⁴-O-R⁴, and R⁴-N(R⁸)-R⁴, wherein said alkylene, alkenylene or alkynylene are each optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸. The alkylene, alkenylene or alkynylene groups of R³ may be linear or branched. In one embodiment R³ is selected from C₅₋₈alkylene, C₅₋₈alkenylene, C₅₋₈alkynylene, R⁴-O-R⁴, and R⁴-N(R⁸)-R⁴, wherein each R⁴ is C₁₋₄alkylene, C₂₋₄alkenylene, or C₂₋₄alkynylene each alkylene, alkenylene and alkynylene being linear or branched and optionally substituted with 1 or 2 substituents selected from halo, oxo, and OR⁸. As noted above, there is a proviso that the total number of carbon atoms in the C₁₋₁₀alkylene, C₂₋₁₀alkenylene, or C₂₋


- 10 alkynylene chains of two R⁴ groups in any definition of R³ is not greater than 12. For instance, when R³ is -R⁴-O-R⁴-, if the first R⁴ group is an ethylene (-CH₂-CH₂-) chain, the maximum number of carbon atoms in the second R⁴ group in that R³ member would be ten.
- 5 The alkylene, alkenylene or alkynylene groups of R⁴ may also be linear or branched. In one embodiment R³ is defined such that each alkylene, alkenylene and alkynylene and each group R⁴ is linear or branched, but unsubstituted by halo, oxo, and OR⁸. In one particular embodiment R³ is unsubstituted, linear or branched C₅₋₈alkylene. In one preferred embodiment 10 R³ is unsubstituted, linear C₅alkylene. In one preferred embodiment R³ is unsubstituted, linear C₆alkylene. In one preferred embodiment R³ is unsubstituted, linear C₇alkylene. In one particular embodiment R³ is unsubstituted, linear C₅alkynylene.
- 15 In one embodiment, the compounds of the invention are defined wherein R³ is selected from C₃₋₆cycloalkylene, R⁴-C₃₋₆cycloalkylene, C₃₋₆cycloalkylene-R⁴, R⁴-C₃₋₆cycloalkylene-R⁴, C₆₋₁₀arylene, R⁴-C₆₋₁₀arylene, C₆₋₁₀arylene-R⁴, R⁴-C₆₋₁₀arylene-R⁴, R⁴-C₆₋₁₀arylene-O-R⁴, R⁴-C₆₋₁₀arylene-N(R⁸)-R⁴, R⁴-C₆₋₁₀arylene-C₆₋₁₀arylene, Het, R⁴-Het, Het-R⁴, R⁴-Het-R⁴, R⁴-O-Het 20 R⁴-C₆₋₁₀arylene-O-Het, R⁴-C₆₋₁₀arylene-C(O)-Het, and R⁴-C₆₋₁₀arylene-N(R⁸)-Het. In one particular embodiment, R³ is selected from C₆₋₁₀arylene, R⁴-C₆₋₁₀arylene, R⁴-C₆₋₁₀arylene-R⁴, Het, R⁴-Het, and R⁴-Het-R⁴. In one particular embodiment, R³ is selected from R⁴-C₆₋₁₀arylene, Het, R⁴-Het, and R⁴-Het-R⁴. In one particular embodiment, R³ is selected from 25 R⁴-phenylene, R⁴-phenylene-R⁴, Het, R⁴-Het, and R⁴-Het-R⁴. In one particular embodiment, R³ is selected from R⁴-phenylene, Het, and R⁴-Het. In one particular embodiment, R³ is selected from R⁴-phenylene, Het, and R⁴-Het, and R⁴ is unsubstituted, linear or branched C₁₋₆alkylene, C₃₋₆alkenylene, or C₃₋₆alkynylene. In one preferred embodiment, R³ is selected from 30 R⁴-phenylene, and R⁴-Het, and R⁴ is unsubstituted, linear or branched C₁₋₆alkylene, C₃₋₆alkenylene, or C₃₋₆alkynylene.

In one preferred embodiment, R³ is R⁴-C₆₋₁₀arylene, particularly R⁴-phenylene.

In one such particular embodiment, R⁴ is unsubstituted, linear or branched C₁₋₆alkylene, C₃₋₆alkenylene, or C₃₋₆alkynylene, more particularly R⁴ is

unsubstituted, linear or branched C₁₋₆alkylene. In one preferred embodiment,

- 5 R³ is R⁴-phenylene and R⁴ is selected from -CH₂-, -(CH₂)₅-, -CH(CH₃)CH₂-, and -C(CH₃)₂CH₂-. In one preferred embodiment, R³ is selected from



In one embodiment, Het is a 6-membered saturated heterocyclene wherein 1

- 10 ring atom is N, and one ring atom is selected from C, N, O and S, wherein the heterocyclene is optionally substituted once with halo, alkyl, alkoxy, oxo or OH.

In one particular embodiment, Het is a 6-membered saturated heterocyclene wherein 1 ring atom is N, one ring atoms is selected from C, N, O and S, and all other ring atoms are C and wherein the heterocyclene is optionally

- 15 substituted once with halo (particularly Cl), alkyl, alkoxy (particularly OCH₃), oxo or OH. In one particular embodiment, Het is unsubstituted heterocyclene.

In one preferred embodiment, Het is unsubstituted, 6-membered saturated heterocyclene wherein 1 or 2 ring atom(s) is/are N, and all other ring atoms are C.


20

In those embodiments wherein R³ includes a Het moiety, R³ may be bound to Y through any suitable carbon or heteroatom. However, the selection of variables R³ and Y should be made in view of each other in order to avoid embodiments which are clearly unstable or inoperative based upon the knowledge of those

- 25 skilled in the art of organic chemistry. For example, when R³ is Het and Het is a nitrogen-containing heterocyclene which is bound to Y through N, one skilled in the art will appreciate that Y is not, for example N(R⁸)C(O) or N(R⁸)C(O)N(R⁸) as such embodiments would result in a N-N bond.

- 30 In one embodiment, the compounds of the invention are defined wherein Y is C(O), C(O)N(R⁸)CH₂, N(R⁸)C(O), or O-C(O)N(R⁸)CH₂. In one particular

embodiment Y is C(O) or C(O)N(R⁸)CH₂. In one preferred embodiment Y is C(O)N(R⁸)CH₂. In one preferred embodiment Y is C(O)N(H)CH₂. Thus in one preferred embodiment, the present invention provides compounds of Formula I':



5


and pharmaceutically acceptable salts thereof, wherein all variables are as defined herein, including all particular and preferred embodiments of variables as described herein.

- 10 When Y is bound to a ring of R³ (e.g., arylene such as phenylene, or heterocyclene), Y may be bound in the ortho, meta or para positions of the ring of R³. In one embodiment Y is bound in the meta or para position of the ring of R³.
- 15 In one embodiment, the compounds of the invention are defined wherein R⁵ is alkyl, particularly, C₁₋₃alkyl, more particularly -CH₂CH₃.

In one embodiment, the compounds of the invention are defined wherein R⁶ is alkyl, particularly, C₁₋₃alkyl, more particularly -CH₂CH₃. In one embodiment, R⁶ is H.

In one embodiment, R⁷ is selected from an unsubstituted C₃₋₆cycloalkyl, substituted C₃₋₆cycloalkyl. Substituted cycloalkyl refers to cycloalkyl substituted halo, alkyl, hydroxyl, O-alkyl, oxo, NH₂, N(H)alkyl, N(alkyl)₂, C(O)NH₂, C(O)N(H)alkyl, and C(O)N(alkyl)₂.

In one embodiment, R⁷ is a heterocyclic group selected from formulas (i), (ii), and (iii):




wherein Z is O, S, S(O)₂, NH or N-R^{7a}, and

R^{7a} is selected from alkyl, C(O)alkyl, C(O)NH₂, C(O)N(H)alkyl, and C(O)N(alkyl)₂.

5


In one embodiment, R⁷ is a heterocyclic group of formula (ii):



(ii) , wherein Z is as defined above.

In one embodiment, R⁷ is a heterocyclic group of formula (ii), wherein Z is O, S,

10 S(O)₂, NH or N-R^{7a}, and R^{7a} is selected from methyl, ethyl, n-propyl, isopropyl, C(O)methyl, C(O)NH₂, C(O)N(H)methyl, and C(O)N(methyl)₂.



In one embodiment, R⁷ is




In one preferred embodiment, R⁷ is

15


In one embodiment, the compounds of the invention are defined wherein R⁸ is H or C₁₋₄alkyl; more particularly H or C₁₋₃alkyl. In one embodiment R⁸ is H or methyl.

20 Specific examples of compounds of Formula I set forth in the examples which follow. Preferred compounds of Formula I are selected from:

N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[(R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]propyl]benzamide




(R)-N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]methyl]benzamide




5


(R)-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]methyl]benzamide



- 10 (R)-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[[5-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]pentyl]benzamide



- 15 (R)-N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-8-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]octanamide



and pharmaceutically acceptable salts thereof.

The compounds of Formula I, may be in the form of a free base or a salt, particularly a pharmaceutically acceptable salt. For a review of
5 pharmaceutically acceptable salts see Berge et al., *J. Pharma Sci.* (1977) 66:1-19.

Pharmaceutically acceptable salts formed from inorganic or organic acids include for example, hydrochloride, hydrobromide, hydroiodide, sulfate,
10 bisulfate, nitrate, sulfamate, phosphate, hydrogen phosphate, acetate, trifluoroacetate, maleate, malate, fumarate, lactate, tartrate, citrate, formate, gluconate, succinate, pyruvate, tannate, ascorbate, palmitate, salicylate, stearate, phthalate, alginate, polyglutamate, oxalate, oxaloacetate, saccharate, benzoate, alkyl or aryl sulfonates (e.g., methanesulfonate, ethanesulfonate,
15 benzenesulfonate, p-toluenesulfonate or naphthalenesulfonate) and isothionate; complexes formed with amino acids such as lysine, arginine, glutamic acid, glycine, serine, threonine, alanine, isoleucine, leucine and the like. The compounds of the invention may also be in the form of salts formed from elemental anions such as chlorine, bromine or iodine.

20 For therapeutic use, salts of active ingredients of the compounds of Formula I will be pharmaceutically acceptable, i.e. they will be salts derived from a pharmaceutically acceptable acid. However, salts of acids which are not pharmaceutically acceptable may also find use, for example, in the preparation
25 or purification of a pharmaceutically acceptable compound. Trifluoroacetate salts, for example, may find such use. All salts, whether or not derived from a pharmaceutically acceptable acid, are within the scope of the present invention. In one embodiment, the compounds of Formula I are in the form of the trifluoroacetate salt. In one embodiment, the compounds of Formula I are in
30 the form of the hydrochloride salt.

The term "chiral" refers to molecules which have the property of non-superimposability of the mirror image partner, while the term "achiral" refers to molecules which are superimposable on their mirror image partner.

- 5 The term "stereoisomers" refers to compounds which have identical chemical constitution, but differ with regard to the arrangement of the atoms or groups in space. "Diastereomer" refers to a stereoisomer with two or more centers of chirality and whose molecules are not mirror images of one another.
- 10 Diastereomers have different physical properties, e.g. melting points, boiling points, spectral properties, and reactivities. Mixtures of diastereomers may separate under high resolution analytical procedures such as electrophoresis and chromatography. "Enantiomers" refer to two stereoisomers of a compound which are non-superimposable mirror images of one another.
- 15 Stereochemical definitions and conventions used herein generally follow S. P. Parker, Ed., McGRAW-HILL DICTIONARY OF CHEMICAL TERMS (1984) McGraw-Hill Book Company, New York; and Eliel, E. and Wilen, S., STEREOCHEMISTRY OF ORGANIC COMPOUNDS (1994) John Wiley & Sons, Inc., New York.
- 20 Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane of plane-polarized light. In describing an optically active compound, the prefixes D and L or R and S are used to denote the absolute configuration of the molecule about its chiral center(s). A specific stereoisomer may also be referred to as an enantiomer, and a mixture of such
- 25 isomers is often called an enantiomeric mixture. A 50:50 mixture of enantiomers is referred to as a racemic mixture or a racemate, which may occur where there has been no stereoselection or stereospecificity in a chemical reaction or process. The terms "racemic mixture" and "racemate" refer to an equimolar mixture of two enantiomeric species.
- 30 The term "tautomers" refers to a type of stereoisomers in which migration of a hydrogen atom results in two or more structures. The compounds of Formula I may exist in different tautomeric forms. One skilled in the art will recognize that


amidines, amides, guanidines, ureas, thioureas, heterocycles and the like can exist in tautomeric forms. All possible tautomeric forms of the amidines, amides, guanidines, ureas, thioureas, heterocycles and the like of all of the embodiments of Formula I are within the scope of the instant invention.

- 5 Tautomers exist in equilibrium and thus the depiction of a single tautomer in the formulas provided will be understood by those skilled in the art to refer equally to all possible tautomers.

It is to be noted that all enantiomers, diastereomers, and racemic mixtures, 10 tautomers, polymorphs, pseudopolymorphs of compounds within the scope of Formula I and pharmaceutically acceptable salts thereof are embraced by the present invention. All mixtures of such enantiomers and diastereomers, including enantiomerically enriched mixtures and diastereomerically enriched mixtures are within the scope of the present invention. Enantiomerically enriched mixtures are mixtures of enantiomers wherein the ratio of the specified enantiomer to the alternative enantiomer is greater than 50:50. More particularly, an enantiomerically enriched mixture comprises at least about 75% of the specified enantiomer, and preferably at least about 85% of the specified enantiomer. In one embodiment, the enantiomerically enriched mixture is 15 substantially free of the other enantiomer. Similarly, diastereomerically enriched mixtures are mixtures of diastereomers wherein the amount of the specified diastereomer is greater than the amount of each alternative diastereomer. More particularly, a diastereomerically enriched mixture comprises at least about 75% of the specified diastereomer, and preferably at 20 least about 85% of the specified diastereomer. In one embodiment, the diastereomerically enriched mixture is substantially free of all other diastereomers. . The term "substantially free of" will be understood by those skilled in the art to indicate less than a 5% presence of other diastereomers, preferably less than 1%, more preferably less than 0.1%. In other 25 embodiments no other diastereomers will be present or the amount of any other diastereomers present will be below the level of detection. Stereoisomers may be separated by techniques known in the art, including high performance liquid chromatography (HPLC) and crystallization of chiral salts.


For illustrative purposes, specific examples of enantiomers within the scope of the present invention include:

- 5 N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[[**(R)**-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]propyl]benzamide




and

- 10 N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[[**S**-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]propyl]benzamide



- 15 In one embodiment, the present invention provides an enantiomerically enriched mixture comprising (**R**)-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[5-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]pentyl]benzamide



or a pharmaceutically acceptable salt thereof, as the predominant isomer.

20

A compound of Formula I and pharmaceutically acceptable salts thereof may exist as different polymorphs or pseudopolymorphs. As used herein, crystalline polymorphism means the ability of a crystalline compound to exist in different

crystal structures. The crystalline polymorphism may result from differences in crystal packing (packing polymorphism) or differences in packing between different conformers of the same molecule (conformational polymorphism). As used herein, crystalline pseudopolymorphism also includes the ability of a
5 hydrate or solvate of a compound to exist in different crystal structures. The pseudopolymorphs of the instant invention may exist due to differences in crystal packing (packing pseudopolymorphism) or due to differences in packing between different conformers of the same molecule (conformational pseudopolymorphism). The instant invention comprises all polymorphs and
10 pseudopolymorphs of the compounds of Formula I and pharmaceutically acceptable salts thereof.

A compound of Formula I and pharmaceutically acceptable salts thereof may also exist as an amorphous solid. As used herein, an amorphous solid is a
15 solid in which there is no long-range order of the positions of the atoms in the solid. This definition applies as well when the crystal size is two nanometers or less. Additives, including solvents, may be used to create the amorphous forms of the instant invention. The instant invention comprises all amorphous forms of the compounds of Formula I and pharmaceutically acceptable salts
20 thereof.

The compounds of the invention may also be in the form of prodrugs. More specifically, the compounds may be present in the form of *in vivo* cleavable esters of the compounds of Formula I and salts of such esters. Examples of
25 suitable esters include acetate, pivalate, tartrate, maleate, succinate and the like.

USES

The compounds of the invention exhibit bifunctional activity as a dual
30 pharmacophore phosphodiesterase 4 inhibitor (PDE4i), and beta agonists. Without being bound by any particular theory, it is believed that the compounds of the invention may function *in vivo* by reducing pulmonary inflammation (by elevation of cytosolic levels of 3',5'-cyclic adenosine monophosphate (cAMP)

through inhibition of the PDE4 enzyme and potentially other pro-inflammatory mechanisms) and inducing bronchodilation (by the beta adrenergic receptor agonist moiety). There may also be further positive cooperative anti-inflammatory effects through simultaneous interaction of downstream signaling pathways via modulation of both targets with the same cell.

5 Local delivery of a single bi-functional compound which has dual activity as a PDE4i and beta agonist offers advantages over combination and conjunctive therapies. In particular, such bi-functional compounds may provide cooperative
10 anti-inflammatory or bronchodilator effects though simultaneous modulation of the same pathways. Utilizing the bi-functional compounds of the present invention allows co-disposition in the same microenvironment which cannot be ensured with the individual drug compounds of combination or conjunctive therapy. In addition, such bi-functional compounds may provide reduced off-target effects leading to decreased risk of adverse events which may be associated with individual PDE4i or beta agonist compounds. If desired however, the dual active compounds of the present invention may nevertheless be combined with other pharmaceutical and non-pharmaceutical therapies which are conventionally employed in the treatment of respiratory diseases.
15 20 Further detail regarding combination therapies utilizing the compounds of the present invention are described below.

As a consequence, the compounds of the invention are useful as medicaments, particularly for the treatment of clinical conditions for which a PDE4i or beta
25 agonist may be indicated. Such conditions include the treatment of pulmonary inflammation or bronchoconstriction and a variety of respiratory diseases. For a review of potential therapeutic activities of PDE4i in the treatment of respiratory diseases see e.g., Kroegel & Foerster, Phosphodiesterase-4 inhibitors as a novel approach for the treatment of respiratory disease:
30 cilomilast, *Expert Opin. Investig. Drugs* (2007) 16:109-124; Dastidar et al., Therapeutic benefit of PDE4 Inhibitors in inflammatory diseases, *Curr Opin Investig Drugs* (2007) 8:364-372; Krymskaya, et al., Phosphodiesterases regulate airway smooth muscle function in health and disease, *Curr. Top. Dev.*

Biol. (2007) 79:61-74; and Spina, PDE4 inhibitors: current status, *Brit. J. Pharmacol.* 2008;155:308-15.

- In particular, the compounds of the invention are useful in methods of treating a variety of respiratory conditions such as diseases associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), including acute exacerbations of COPD, asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, and transplant-associated bronchiolitis, including lung- and bone marrow-transplant associated bronchiolitis, in a human in need thereof. The compounds of the invention may also be useful for treating ventilator-associated tracheobronchitis and/or preventing ventilator-associated pneumonia in ventilated patients. With respect to the treatment of acute exacerbations of COPD, the compounds of the invention are useful for reducing the frequency, severity or duration of acute exacerbation of COPD and/or for reducing the frequency, severity or duration of one or more symptoms of acute exacerbation of COPD.
- All therapeutic uses and methods described herein are carried out by the step of administering an effective amount of a compound of the invention (a compound of Formula I or a pharmaceutically acceptable salt thereof) to a subject (typically mammal and preferably human) in need of treatment.
- In one aspect, the present invention provides a method for the treatment of a condition in a mammal, such as a human, for which a PDE 4i or a beta agonist is indicated.
- The terms "treating" and "treatment", as used herein refers to reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition or one or more symptoms of such disorder or condition.

In one embodiment the invention provides a method for the treatment of a

respiratory disease. In one embodiment the invention provides a method for the treatment of a disease associated with reversible or irreversible airway obstruction in a mammal, particularly a human, in need thereof. In one particular embodiment the present invention provides a method for the 5 treatment of COPD in a mammal, particularly a human in need thereof. In one particular embodiment the present invention provides a method for reducing the frequency, severity or duration of acute exacerbation of COPD or for the treatment of one or more symptoms of acute exacerbation of COPD in a mammal, particularly a human in need thereof. In one embodiment the 10 invention provides a method for the treatment of asthma in a mammal, particularly a human, in need thereof. In one embodiment the invention provides a method for the treatment of bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis) in a mammal, particularly a human, in need thereof. In one embodiment the invention provides a method for the 15 treatment of bronchitis, including acute and chronic bronchitis in a mammal, particularly a human, in need thereof. In one embodiment the invention provides a method for the treatment of post-viral cough in a mammal, particularly a human, in need thereof. In one embodiment the invention provides a method for the treatment of cystic fibrosis in a mammal, particularly 20 a human, in need thereof. In one embodiment the invention provides a method for the treatment of emphysema in a mammal, particularly a human in need thereof. In one embodiment the invention provides a method for the treatment of pneumonia in a mammal, particularly a human in need thereof. In one embodiment the invention provides a method for the treatment of 25 panbronchiolitis in a mammal, particularly a human in need thereof. In one embodiment the invention provides a method for the treatment of transplant-associated bronchiolitis, including lung- and bone marrow-transplant associated bronchiolitis in a mammal, particularly a human in need thereof. In one embodiment the invention provides a method for treating ventilator-associated 30 tracheobronchitis and/or preventing ventilator-associated pneumonia in a ventilated human in need thereof. In one embodiment the invention provides a method for treating sinusitis in a mammal, particularly a human in need thereof.

There is also provided a compound of the invention for use in medical therapy, particularly for use in the treatment of condition in a mammal, such as a human, for which a PDE4i or beta agonist is indicated. In one embodiment the invention provides a method for the treatment of a respiratory disease. In one 5 embodiment there is provided a compound of the invention for use in the treatment of a disease associated with reversible or irreversible airway obstruction in a mammal, particularly a human, in need thereof. In one particular embodiment there is provided a compound of the invention for use in the treatment of chronic obstructive pulmonary disease (COPD) in a mammal, 10 particularly a human in need thereof. In one embodiment, there is provided a compound of the invention for use in reducing the frequency, severity or duration of acute exacerbation of COPD or for the treatment of one or more symptoms of acute exacerbation of COPD, in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound of the 15 invention for use in the treatment of asthma in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound for use in the treatment of bronchiectasis, including bronchiectasis due to conditions other than cystic fibrosis, in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound for use in the treatment of 20 bronchitis, including acute bronchitis and chronic bronchitis, in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound for use in the treatment of post-viral cough, in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound for use in the treatment of cystic fibrosis in a mammal, particularly a 25 human in need thereof. In one embodiment there is provided a compound of the invention for use in the treatment of emphysema in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound of the invention for use in the treatment of pneumonia in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound of 30 the invention for use in the treatment of panbronchiolitis or transplant-associated bronchiolitis, including lung- and bone marrow-transplant associated bronchiolitis in a mammal, particularly a human, in need thereof. In one embodiment there is provided a compound of the invention for use in the

treatment of ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia in a ventilated human in need thereof. In one embodiment there is provided a compound of the invention for use in the treatment of sinusitis in a mammal, particularly a human, in need thereof.

5

The present invention also provides the use of a compound of the invention in the manufacture of a medicament for the treatment of a condition in a mammal, such as a human, for which a PDE4i or beta agonist is indicated. In one embodiment is provided the use of a compound of the invention in the 10 manufacture of a medicament for the treatment of a respiratory disease. In one embodiment is provided the use of a compound of the invention in the manufacture of a medicament for the treatment of diseases associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), acute exacerbations of COPD, asthma, bronchiectasis 15 (including bronchiectasis due to conditions other than cystic fibrosis), bronchitis (including acute bronchitis and chronic bronchitis), post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associated bronchiolitis, (including lung- and bone marrow-transplant associated bronchiolitis), ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia or treating sinusitis.

The term "effective amount", as used herein, is an amount of compound of the invention which is sufficient in the subject to which it is administered, to elicit the biological or medical response of a cell culture, tissue, system, or mammal 25 (including human) that is being sought, for instance by a researcher or clinician. The term also includes within its scope, amounts effective to enhance normal physiological function. In one embodiment, the effective amount is the amount needed to provide a desired level of drug in the secretions and tissues of the airways and lungs, or alternatively, in the bloodstream of a subject to be treated 30 to give an anticipated physiological response or desired biological effect when such a composition is administered by inhalation. For example, an effective amount of a compound of the invention for the treatment of a condition for which a PDE4i or beta agonist is indicated is sufficient in the subject to which it

is administered to treat the particular condition. In one embodiment an effective amount is an amount of a compound of the invention which is sufficient for the treatment of COPD or cystic fibrosis in a human.

- 5 The precise effective amount of the compounds of the invention will depend on a number of factors including but not limited to the species, age and weight of the subject being treated, the precise condition requiring treatment and its severity, the bioavailability, potency, and other properties of the specific compound being administered, the nature of the formulation, the route of administration, and the delivery device, and will ultimately be at the discretion of the attendant physician or veterinarian. Further guidance with respect to appropriate dose may be found in considering conventional dosing of other PDE4i's such as cilomilast or roflumilast and other beta agonist's such as salmeterol, with due consideration also being given to any differences in
- 10 potency between those compounds and the compounds of the present invention and that bi-functional nature of the compounds of the present invention.
- 15

- An estimated dose administered topically to the airway surfaces of a subject (e.g., by inhalation) of a compound of the invention for treatment of a 70 kg human may be in the range of from about 20 to about 1000 µg. The selection of the specific dose for a patient will be determined by the attendant physician, clinician or veterinarian of ordinary skill in the art based upon a number of factors including those noted above. In one particular embodiment the dose of
- 25 a compound of the invention for the treatment of a 70 kg human will be in the range of from about 50 to about 750 µg. In one preferred embodiment, the dose of a compound of the invention for the treatment of a 70 kg human will be in the range of from about 50 to about 750 µg. The foregoing suggested doses may be adjusted using conventional dose calculations if the compound is
 - 30 administered via a different route. Determination of an appropriate dose for administration by other routes is within the skill of those in the art in light of the foregoing description and the general knowledge in the art.

Delivery of an effective amount of a compound of the invention may entail delivery of a single dosage form or multiple unit doses which may be delivered contemporaneously or separate in time over a designated period, such as 24 hours. A dose of a compound of the invention (alone or in the form of a

- 5 composition comprising the same) may be administered from one to ten times per day. Typically, a compound of the invention (alone or in the form of a composition comprising the same) will be administered four, three, two, or most preferably once per day (24 hours).

10 COMPOSITIONS

While it is possible for a compound of the invention to be administered alone, it is preferable to present it in the form of a composition, particularly a pharmaceutical composition (formulation). Thus, in another aspect, the invention provides compositions, and particularly pharmaceutical compositions 15 (such as an inhalable pharmaceutical composition) comprising a compound of the invention as an active ingredient, and a pharmaceutically acceptable excipient, diluent or carrier. The term "active ingredient" as employed herein refers to any compound of the invention or combination of two or more compounds of the invention in a pharmaceutical composition. The 20 pharmaceutically acceptable excipient(s), diluent(s) or carrier(s) must be acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Generally, the pharmaceutically acceptable excipient(s), diluent(s) or carrier(s) employed in the pharmaceutical formulation are "non-toxic" meaning that it/they is/are 25 deemed safe for consumption in the amount delivered in the formulation and "inert" meaning that it/they does/do not appreciable react with or result in an undesired effect on the therapeutic activity of the active ingredient(s). Pharmaceutically acceptable excipients, diluents and carriers are conventional in the art and may be selected using conventional techniques, based upon the 30 desired route of administration. See, REMINGTON'S, PHARMACEUTICAL SCIENCES, Lippincott Williams & Wilkins; 21st Ed (May 1, 2005). Preferably, the pharmaceutically acceptable excipient(s), diluent(s) or carrier(s) are Generally Regarded As Safe (GRAS) according to the FDA.

- Pharmaceutical compositions according to the invention include those suitable for topical administration and administration to the respiratory tract, including the nasal cavities and sinuses, oral and extrathoracic airways, and the lungs, including by use of aerosols which may be delivered by means of various types
- 5 of dry powder inhalers, pressurized metered dose inhalers, softmist inhalers, nebulizers, or insufflators. The most suitable route of administration may depend upon, several factors including the patient and the condition or disorder being treated.
- 10 The formulations may be presented in unit dosage form or in bulk form as for example in the case of formulations to be metered by an inhaler and may be prepared by any of the methods well known in the art of pharmacy. Generally, the methods include the step of bringing the active ingredient into association with the carrier, diluent or excipient and optionally one or more accessory
- 15 ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with one or more liquid carriers, diluents or excipients or finely divided solid carriers, diluents or excipients, or both, and then, if necessary, shaping the product into the desired formulation.
- 20 In one preferred embodiment, the composition is an inhalable pharmaceutical composition which is suitable for inhalation and delivery to the endobronchial space. Typically, such composition is in the form of an aerosol comprising particles for delivery using a nebulizer, pressurized metered dose inhaler (MDI),
- 25 softmist inhaler, or dry powder inhaler (DPI). The aerosol formulation used in the methods of the present invention may be a liquid (e.g., solution) suitable for administration by a nebulizer, softmist inhaler, or MDI, or a dry powder suitable for administration by an MDI or DPI.
- 30 Aerosols used to administer medicaments to the respiratory tract are typically polydisperse; that is they are comprised of particles of many different sizes. The particle size distribution is typically described by the Mass Median Aerodynamic Diameter (MMAD) and the Geometric Standard Deviation (GSD).

- For optimum drug delivery to the endobronchial space the MMAD is in the range from about 1 to about 10 μm and preferably from about 1 to about 5 μm , and the GSD is less than 3, and preferably less than about 2. Aerosols having a MMAD above 10 μm are generally too large when inhaled to reach the lungs.
- 5 Aerosols with a GSD greater than about 3 are not preferred for lung delivery as they deliver a high percentage of the medicament to the oral cavity. To achieve these particle sizes in powder formulation, the particles of the active ingredient may be size reduced using conventional techniques such as micronisation or spray drying. Non-limiting examples of other processes or techniques that can
- 10 be used to produce respirable particles include spray drying, precipitation, supercritical fluid, and freeze drying. The desired fraction may be separated out by air classification or sieving. In one embodiment, the particles will be crystalline. For liquid formulations, the particle size is determined by the selection of a particular model of nebulizer, softmist inhaler, or MDI.
- 15 Aerosol particle size distributions are determined using devices well known in the art. For example a multi-stage Anderson cascade impactor or other suitable method such as those specifically cited within the US Pharmacopoeia Chapter 601 as characterizing devices for aerosols emitted from metered-dose and dry powder inhalers.
- 20 Dry powder compositions for topical delivery to the lung by inhalation may be formulated without excipient or carrier and instead including only the active ingredients in a dry powder form having a suitable particle size for inhalation. Dry powder compositions may also contain a mix of the active ingredient and a
- 25 suitable powder base (carrier/diluent/excipient substance) such as mono-, di- or poly-saccharides (e.g., lactose or starch). Lactose is typically the preferred excipient for dry powder formulations. When a solid excipient such as lactose is employed, generally the particle size of the excipient will be much greater than the active ingredient to aid the dispersion of the formulation in the inhaler.
- 30 Non-limiting examples of dry powder inhalers include reservoir multi-dose inhalers, pre-metered multi-dose inhalers, capsule-based inhalers and single-dose disposable inhalers. A reservoir inhaler contains a large number of doses

- (e.g. 60) in one container. Prior to inhalation, the patient actuates the inhaler which causes the inhaler to meter one dose of medicament from the reservoir and prepare it for inhalation. Examples of reservoir DPIs include but are not limited to the Turbohaler® by AstraZeneca and the ClickHaler® by Vectura.
- 5 In a pre-metered multi-dose inhaler, each individual dose has been manufactured in a separate container, and actuation of the inhaler prior to inhalation causes a new dose of drug to be released from its container and prepared for inhalation. Examples of multidose DPI inhalers include but are not limited to Diskus® by GSK, Gyrohaler® by Vectura, and Prohaler® by Valois.
- 10 During inhalation, the inspiratory flow of the patient accelerates the powder out of the device and into the oral cavity. For a capsule inhaler, the formulation is in a capsule and stored outside the inhaler. The patient puts a capsule in the inhaler, actuates the inhaler (punctures the capsule), then inhales. Examples include the Rotohaler™ (GlaxoSmithKline), Spinhaler™ (Novartis),
- 15 HandiHaler™ (IB), TurboSpin™ (PH&T). With single-dose disposable inhalers, the patient actuates the inhaler to prepare it for inhalation, inhales, then disposes of the inhaler and packaging. Examples include the Twincer™ (U Groningen), OneDose™ (GFE), and Manta Inhaler™ (Manta Devices).
- 20 Generally, dry powder inhalers utilize turbulent flow characteristics of the powder path to cause the excipient-drug aggregates to disperse, and the particles of active ingredient are deposited in the lungs. However, certain dry powder inhalers utilize a cyclone dispersion chamber to produce particles of the desired respirable size. In a cyclone dispersion chamber, the drug enters a
- 25 coin shaped dispersion chamber tangentially so that the air path and drug move along the outer circular wall. As the drug formulation moves along this circular wall it bounces around and agglomerates are broken apart by impact forces. The air path spirals towards the center of the chamber exiting vertically. Particles that have small enough aerodynamic sizes can follow the air path and exit the chamber. In effect, the dispersion chamber works like a small jet mill.
- 30 Depending on the specifics of the formulation, large lactose particles may be added to the formulation to aid in the dispersion through impact with the API particles.

The Twincer™ single-dose disposable inhaler appears to operate using a coin-shaped cyclone dispersion chamber referred to as an “air classifier.” See, U.S. Published Patent Application No. 2006/0237010 to Rijksuniversiteit Groningen.

- 5 Papers published by the University of Groningen, have stated that a 60 mg dose of pure micronized colistin sulfomethate could be effectively delivered as an inhalable dry powder utilizing this technology.

In preferred embodiments, the aerosol formulation is delivered as a dry powder
10 using a dry powder inhaler wherein the particles emitted from the inhaler have an MMAD in the range of about 1 μm to about 5 μm and a GSD about less than 2.

Examples of suitable dry powder inhalers and dry powder dispersion devices
15 for use in the delivery of compounds and compositions according to the present invention include but are not limited to those disclosed in US7520278; US7322354; US7246617; US7231920; US7219665; US7207330; US6880555; US5,522,385; US6845772; US6637431; US6329034; US5,458,135; US4,805,811; and U.S. Published Patent Application No. 2006/0237010.

20 In one embodiment, the pharmaceutical formulation according to the invention is a dry powder for inhalation which is formulated for delivery by a Diskus®-type device. The Diskus® device comprises an elongate strip formed from a base sheet having a plurality of recesses spaced along its length and a lid sheet
25 hermetically but peelably sealed thereto to define one or more containers, each container having therein an inhalable formulation containing a predetermined amount of active ingredient either alone or in admixture with one or more carriers or excipients (e.g., lactose) and/or other therapeutically active agents. Preferably, the strip is sufficiently flexible to be wound into a roll. The lid sheet
30 and base sheet will preferably have leading end portions which are not sealed to one another and at least one of the leading end portions is constructed to be attached to a winding means. Also, preferably the hermetic seal between the base and lid sheets extends over their whole width. To prepare the dose for

inhalation, the lid sheet may preferably be peeled from the base sheet in a longitudinal direction from a first end of the base sheet.

In one embodiment, the pharmaceutical formulation according to the invention

5 is a dry powder for inhalation which is formulated for delivery using a single-dose disposable inhaler, and particularly the Twincer™ inhaler. The Twincer™ inhaler comprises a foil laminate blister with one or more recesses and a lid sheet hermetically but peelably sealed thereto to define a plurality of containers. Each container has therein an inhalable formulation containing a

10 predetermined amount of active ingredient(s) either alone or in admixture with one or more carriers or excipients (e.g., lactose). The lid sheet will preferably have a leading end portion which is constructed to project from the body of the inhaler. The patient would operate the device and thereby administer the aerosol formulation by 1) removing the outer packaging overwrap, 2) pulling the

15 foil tab to uncover the drug in the blister and 3) inhaling the drug from the blister.

In another embodiment, the pharmaceutical formulation according to the invention is a dry powder for inhalation wherein the dry powder is formulated into microparticles as described in PCT Publication No. WO2009/015286 or

20 WO2007/114881, both to NexBio. Such microparticles are generally formed by adding a counterion to a solution containing a compound of the invention in a solvent, adding an antisolvent to the solution; and gradually cooling the solution to a temperature below about 25°C, to form a composition containing microparticles comprising the compound. The microparticles comprising the

25 compound may then be separated from the solution by any suitable means such as sedimentation, filtration or lyophilization. Suitable counterions, solvents and antisolvents for preparing microparticles of the compounds of the invention are described in WO2009/015286.

30 In another embodiment, a pharmaceutical composition according to the invention is delivered as a dry powder using a metered dose inhaler. Non-limiting examples of metered dose inhalers and devices include those disclosed in US5,261,538; US5,544,647; US5,622,163; US4,955,371; US3,565,070;

US3,361306 and US6,116,234 and US7,108,159. In a preferred embodiment, a compound of the invention is delivered as a dry powder using a metered dose inhaler wherein the emitted particles have an MMAD that is in the range of about 1 μm to about 5 μm and a GSD that is less than about 2.

5

Liquid aerosol formulations for delivery to the endobronchial space or lung by inhalation may for example be formulated as aqueous solutions or suspensions or as aerosols delivered from pressurized packs, such as metered dose inhalers, with the use of suitable liquefied propellants, softmist inhalers, or

10 nebulizers. Such aerosol compositions suitable for inhalation can be either a suspension or a solution and generally contain the active ingredient(s) together with a pharmaceutically acceptable carrier or diluent (e.g., water (distilled or sterile), saline, hypertonic saline, or ethanol) and optionally one or more other therapeutically active agents.

15

Aerosol compositions for delivery by pressurized metered dose inhalers typically further comprise a pharmaceutically acceptable propellant. Examples of such propellants include fluorocarbon or hydrogen-containing chlorofluorocarbon or mixtures thereof, particularly hydrofluoroalkanes, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, especially 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoro-n-propane or a mixture thereof. The aerosol composition may be excipient free or may optionally contain additional formulation excipients well known in the art such as surfactants e.g., oleic acid or lecithin and cosolvents e.g., ethanol.

20 25 Pressurized formulations will generally be retained in a canister (e.g., an aluminum canister) closed with a valve (e.g., a metering valve) and fitted into an actuator provided with a mouthpiece.

In another embodiment, a pharmaceutical composition according to the
30 invention is delivered as a liquid using a metered dose inhaler. Non-limiting examples of metered dose inhalers and devices include those disclosed in US Patent Nos. 6,253,762, 6,413,497, 7,601,336, 7,481,995, 6,743,413, and 7,105,152. In a preferred embodiment, a compound of the invention is

delivered as a dry powder using a metered dose inhaler wherein the emitted particles have an MMAD that is in the range of about 1 μ m to about 5 μ m and a GSD that is less than about 2.

- 5 In one embodiment the aerosol formulation is suitable for aerosolization by a jet nebulizer, or ultrasonic nebulizer including static and vibrating porous plate nebulizers. Liquid aerosol formulations for nebulization may be generated by solubilizing or reconstituting a solid particle formulation or may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali,
- 10 buffer salts, and isotonicity adjusting agents. They may be sterilized by in-process techniques such as filtration, or terminal processes such as heating in an autoclave or gamma irradiation. They may also be presented in non-sterile form.
- 15 Patients can be sensitive to the pH, osmolality, and ionic content of a nebulized solution. Therefore these parameters should be adjusted to be compatible with the active ingredient and tolerable to patients. The most preferred solution or suspension of active ingredient will contain a chloride concentration >30 mM at pH 4.5-8.0 and an osmolality of from about 800-1600 mOsm/kg. The pH of the solution can be controlled by either titration with common acids (hydrochloric acid or sulfuric acid, for example) or bases (sodium hydroxide, for example) or via the use of buffers. Commonly used buffers include citrate buffers, acetate buffers, and phosphate buffers. Buffer strengths can range from 2 mM to 50 mM.
- 20
- 25 Such formulations may be administered using commercially available nebulizers or other atomizer that can break the formulation into particles or droplets suitable for deposition in the respiratory tract. Non-limiting examples of nebulizers which may be employed for the aerosol delivery of a composition of the invention include pneumatic jet nebulizers, vented or breath enhanced jet nebulizers, or ultrasonic nebulizers including static or vibrating porous plate nebulizers.
- 30

A jet nebulizer utilizes a high velocity stream of air blasting up through a column of water to generate droplets. Particles unsuitable for inhalation impact on walls or aerodynamic baffles. A vented or breath enhanced nebulizer works in essentially the same way as a jet nebulizer except that inhaled air passes

5 through the primary droplet generation area to increase the output rate of the nebulizer while the patient inhales.

In an ultrasonic nebulizer, vibration of a piezoelectric crystal creates surface instabilities in the drug reservoir that cause droplets to be formed. In porous

10 plate nebulizers pressure fields generated by sonic energy force liquid through the mesh pores where it breaks into droplets by Rayleigh breakup. The sonic energy may be supplied by a vibrating horn or plate driven by a piezoelectric crystal, or by the mesh itself vibrating. Non-limiting examples of atomizers include any single or twin fluid atomizer or nozzle that produces droplets of an

15 appropriate size. A single fluid atomizer works by forcing a liquid through one or more holes, where the jet of liquid breaks up into droplets. Twin fluid atomizers work by either forcing both a gas and liquid through one or more holes, or by impinging a jet of liquid against another jet of either liquid or gas.

20 The choice of nebulizer which aerosolizes the aerosol formulation is important in the administration of the active ingredient(s). Different nebulizers have differing efficiencies based their design and operation principle and are sensitive to the physical and chemical properties of the formulation. For example, two formulations with different surface tensions may have different

25 particle size distributions. Additionally, formulation properties such as pH, osmolality, and permeant ion content can affect tolerability of the medication, so preferred embodiments conform to certain ranges of these properties.

In a preferred embodiment, the formulation for nebulization is delivered to the

30 endobronchial space as an aerosol having an MMAD between about 1 μm and about 5 μm and a GSD less than 2 using an appropriate nebulizer. To be optimally effective and to avoid upper respiratory and systemic side effects, the aerosol should not have a MMAD greater than about 5 μm and should not have

a GSD greater than about 2. If an aerosol has an MMAD larger than about 5 μm or a GSD greater than about 2, a large percentage of the dose may be deposited in the upper airways decreasing the amount of drug delivered to the site of inflammation and bronchoconstriction in the lower respiratory tract. If the 5 MMAD of the aerosol is smaller than about 1 μm then a large percentage of the particles may remain suspended in the inhaled air and may then be exhaled during expiration.

10 The compounds of the invention may also be administered by transbronchoscopic lavage.

Compositions designed for nasal administration include aerosols, solutions, suspensions, sprays, mists and drops. Aerosolable formulations for nasal administration may be formulated in much the same ways as aerosolable 15 formulations for inhalation with the condition that particles of non-respirable size will be preferred in formulations for nasal administration. Typically, particles of about 5 microns in size, up to the size of visible droplets may be employed. Thus, for nasal administration, a particle size in the range of 10-500 μm may be used to ensure retention in the nasal cavity.

20 In another aspect, the invention provides a method for treating a respiratory disease in a human in need thereof, comprising administering to the human a pharmaceutical composition comprising a compound of the invention, wherein the compound is administered in an effective amount. In one preferred embodiment, the method comprises administering the pharmaceutical 25 composition as an inhalable composition comprising from about 20 to about 1000 μg of a compound of the invention.

In another aspect, the invention provides a method of treating any one of: a 30 disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia,

- panbronchiolitis, transplant-associate bronchiolitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human in need thereof, comprising administering to the human a pharmaceutical composition comprising a compound of the invention, wherein
- 5 the compound is administered in an effective amount. In one preferred embodiment, the method comprises administering the pharmaceutical composition as an inhalable composition comprising from about 20 to about 1000 micrograms of a compound of the invention.
- 10 Preferred unit dosage formulations for the compounds of the invention are those containing an effective amount of the active ingredient or an appropriate fraction thereof.
- It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question.
- 15 The compositions of the present invention may be formulated for immediate, controlled or sustained release as desired for the particular condition being treated and the desired route of administration. Because the free base of a compound is generally less soluble in aqueous solutions than the salt, compositions comprising a free base of a compound of Formula I may be employed to provide more sustained release of active agent delivered by inhalation to the lungs. An active agent present in the lungs in particulate form which has not dissolved into solution is not available to induce a physiological response, but serves as a depot of bioavailable drug which gradually dissolves into solution. As another example, a formulation may employ both a free base and salt form of a compound of the invention to provide both immediate release and sustained release of the active ingredient for dissolution into the mucus
- 20 secretions of, for example, the nose.
- 25
- 30

COMBINATIONS

The compounds of the invention may be formulated and/or used in combination with other therapeutically active agents. Examples of other therapeutically active agents which may be formulated or used in combination with the

- 5 compounds of the invention include but are not limited to anti-inflammatory agents, anticholinergic agents, peroxisome proliferator-activated receptor (PPAR) gamma agonists, PPAR delta agonists, epithelial sodium channel blockers (ENaC blockers), kinase inhibitors (e.g. p38 MAPK, PI3K, JNK, ERK, IKK2), antiinfective agents, and antihistamines.

10

The present invention thus provides, as another aspect, a composition comprising an effective amount of a compound of the invention and one or more other therapeutically active agents selected from anti-inflammatory agents, anticholinergic agents, P2Y2 receptor agonists, PPAR gamma agonists,

- 15 PPAR delta agonists, ENaC blockers, kinase inhibitors (e.g. p38 MAPK, PI3K, JNK, ERK, IKK2), antiinfective agents, and antihistamines. Use of the compounds of the invention in combination with one or more other therapeutically active agents may lower the dose of the compound of the invention that is required to treat the respiratory disease, thereby reducing the potential for undesired side-effects attributable to systemically absorbed beta 20 agonists.

Suitable anti-inflammatory agents for use in combination with the compounds of the invention include corticosteroids and non-steroidal anti-inflammatory drugs

- 25 (NSAIDs), particularly phosphodiesterase (PDE) inhibitors. Examples of corticosteroids for use in the present invention include oral or inhaled corticosteroids or prodrugs thereof. Specific examples include but are not limited to ciclesonide, desisobutyryl-ciclesonide, budesonide, flunisolide, mometasone and esters thereof (e.g., mometasone furoate), fluticasone 30 propionate, fluticasone furoate, beclomethasone, methyl prednisolone, prednisolone, dexamethasone, 6 α ,9 α -difluoro-17 α -[(2-furanylcarbonyl)oxy]-11 β -hydroxy-16 α -methyl-3-oxo-androsta-1,4-diene-17 β -carbothioic acid S-fluoromethyl ester, 6 α ,9 α -difluoro-11 β -hydroxy-16 α -methyl-3-oxo-17 α -

- propionyloxy-androsta-1,4-diene-17 β -carbothioic acid S-(2-oxo-tetrahydro-furan-3S-yl) ester, beclomethasone esters (e.g., the 17-propionate ester or the 17,21-dipropionate ester, fluoromethyl ester, triamcinolone acetonide, rofleponide, or any combination or subset thereof. Preferred corticosteroids for 5 formulation or use in combination with the compounds of the invention are selected from ciclesonide, desisobutyryl-ciclesonide, budesonide, mometasone, fluticasone propionate, and fluticasone furoate, or any combination or subset thereof.
- 10 NSAIDs for use in the present invention include but are not limited to sodium cromoglycate, nedocromil sodium, phosphodiesterase (PDE) inhibitors (e.g., theophylline, aminophylline, PDE4 inhibitors, mixed PDE3/PDE4 inhibitors or mixed PDE4/PDE7 inhibitors), leukotriene antagonists, inhibitors of leukotriene synthesis (e.g., 5 LO and FLAP inhibitors), nitric oxide synthase (iNOS) 15 inhibitors, protease inhibitors (e.g., tryptase inhibitors, neutrophil elastase inhibitors, and metalloprotease inhibitors) β 2-integrin antagonists and adenosine receptor agonists or antagonists (e.g., adenosine 2a agonists), cytokine antagonists (e.g., chemokine antagonists) or inhibitors of cytokine synthesis (e.g., prostaglandin D2 (CRTh2) receptor antagonists).
- 20 The PDE4 inhibitor, mixed PDE3/PDE4 inhibitor or mixed PDE4/PDE7 inhibitor may be any compound that is known to inhibit the PDE4 enzyme or which is discovered to act as a PDE4 inhibitor, and which are selective PDE4 inhibitors (i.e., compounds which do not appreciably inhibit other members of the PDE 25 family). Examples of specific PDE4 inhibitors for formulation and use in combination with the compounds of the present invention include but are not limited to roflumilast, pumafentrine, arofylline, cilomilast, tofimilast, oglemilast, tolafentrine, piclamilast, ibudilast, apremilast, 2-[4-[6,7-diethoxy-2,3-bis(hydroxymethyl)-1-naphthalenyl]-2-pyridinyl]-4-(3-pyridinyl)-1(2H)- 30 phthalazinone (T2585), N-(3,5-dichloro-4-pyridinyl)-1-[(4-fluorophenyl)methyl]-5-hydroxy- α -oxo-1H-indole-3-acetamide (AWD-12-281, 4-[(2R)-2-[3-(cyclopentyloxy)-4-methoxyphenyl]-2-phenylethyl]-pyridine (CDP-840), 2-[4-[[[2-(1,3-benzodioxol-5-yloxy)-3-pyridinyl]carbonyl]amino]methyl]-3-

- fluorophenoxy]-(2R)-propanoic acid (CP-671305), N-(4,6-dimethyl-2-pyrimidinyl)-4-[4,5,6,7-tetrahydro-2-(4-methoxy-3-methylphenyl)-5-(4-methyl-1-piperazinyl)-1H-indol-1-yl]benzenesulfonamide, (2E)-2-butenedioate (YM-393059), 9-[(2-fluorophenyl)methyl]-N-methyl-2-(trifluoromethyl)-9H-purin-6-amine (NCS-613), N-(2,5-dichloro-3-pyridinyl)-8-methoxy-5-quinolinecarboxamide (D-4418), N-[(3R)-9-amino-3,4,6,7-tetrahydro-4-oxo-1-phenylpyrrolo[3,2,1-][1,4]benzo-diazepin-3-yl]-3H-purin-6-amine (PD-168787), 3-[[3-(cyclopentyloxy)-4-methoxyphenyl]methyl]-N-ethyl-8-(1-methylethyl)-3H-purin-6-amine hydrochloride (V-11294A), N-(3,5-dichloro-1-oxido-4-pyridinyl)-8-methoxy-2-(trifluoromethyl)-5-quinolinecarboxamide (Sch351591), 5-[3-(cyclopentyloxy)-4-methoxyphenyl]-3-[(3-methylphenyl)methyl]-(3S,5S)-2-piperidinone (HT-0712), 5-[2-[(1R,4R)-4-amino-1-(3-(cyclopentyloxy)-4-methoxyphenyl)cyclohexyl]ethynyl]pyrimidine-2-amine, 6-[3-(dimethylcarbamoyl)phenylsulfonyl]-4-(3-methoxyphenylamino)-8-methylquinoline-3-carboxamide (GSK-256066), cis-[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol], and 4-[6,7-diethoxy-2,3-bis(hydroxymethyl)-1-naphthalenyl]-1-(2-methoxyethyl)-2(1H)-pyridinone (T-440), and any combination or subset thereof.
- Leukotriene antagonists and inhibitors of leukotriene synthesis include zafirlukast, montelukast sodium, zileuton, and pranlukast.

Anticholinergic agents for formulation or use in combination with the compounds of the invention include but are not limited to muscarinic receptor antagonists, particularly including pan antagonists and antagonists of the M₃ receptors. Exemplary compounds include the alkaloids of the belladonna plants, such as atropine, scopolamine, homatropine, hyoscyamine, and the various forms including salts thereof (e.g., anhydrous atropine, atropine sulfate, atropine oxide or HCl, methylatropine nitrate, homatropine hydrobromide, homatropine methyl bromide, hyoscyamine hydrobromide, hyoscyamine sulfate, scopolamine hydrobromide, scopolamine methyl bromide), or any combination or subset thereof.

Additional anticholinergics for formulation and use in combination with the methantheline, propantheline bromide, anisotropine methyl bromide or Valpin 50, aclidinium bromide, glycopyrrolate (Robinul), isopropamide iodide, mepenzolate bromide, tridihexethyl chloride, hexocyclium methylsulfate, 5 cyclopentolate HCl, tropicamide, trihexyphenidyl CCl, pirenzepine, telenzepine, and methocramine, or any combination or subset thereof.

Preferred anticholinergics for formulation and use in combination with the compounds of the invention include ipratropium (bromide), oxitropium 10 (bromide) and tiotropium (bromide), or any combination or subset thereof.

Examples of ENaC receptor blockers for formulation and use in combination with the compounds of the invention include but are not limited to amiloride and derivatives thereof such as those compounds described in US Patent No. 15 6858615, and PCT Publication Nos. WO2003/070182, WO2004/073629, WO2005/018644, WO2006/022935, WO2007/018640, and WO2007/146869, all to Parion Sciences, Inc.

Examples of kinase inhibitors include inhibitors of NFkB, PI3K 20 (phosphatidylinositol 3-kinase) (CAL-263 (oral), Trial trove and Calistoga web site), p38-MAP kinase (SB-681323 (oral); Singh *et al.*, *J Clin Pharmacol.* 2010 Jan;50(1):94-100).

Antiinfective agents for formulation and use in combination with the compounds 25 of the invention include antivirals and antibiotics. Examples of suitable antivirals include Tamiflu® and Relenza®. Examples of suitable antibiotics include but are not limited to aztreonam (arginine or lysine), fosfomycin, and aminoglycosides such as tobramycin, or any combination or subset thereof.

30 Antihistamines (i.e., H1-receptor antagonists) for formulation and use in combination with the compounds of the invention include but are not limited to: ethanolamines such as diphenhydramine HCl, carboxamine maleate, doxylamine, clemastine fumarate and dimenhydrinate;

- ethylenediamines such as pyrilamine maleate (metpyramine), tripeleannamine HCl, tripeleannamine citrate, and antazoline; alkylamines such as pheniramine, chloropheniramine, bromopheniramine, dexchlorpheniramine, triprolidine and acrivastine;
- 5 pyridines such as methapyriline, piperazines such as hydroxyzine HCl, hydroxyzine pamoate, cyclizine HCl, cyclizine lactate, meclizine HCl and cetirizine HCl; piperidines such as astemisole, levocabastine HCl, loratadine, descarboethoxy loratadine, terfenadine, and fexofenadine HCl;
- 10 tri- and tetracyclics such as promethazine, chlorpromethazine trimeprazine and azatadine; and azelastine HCl, or any combination or subset thereof.

In the above-described methods of treatment and uses, a compound of the invention may be employed alone, or in combination with one or more other therapeutically active agents. Typically, any therapeutically active agent that has a therapeutic effect in the disease or condition being treated with the compound of the invention may be utilized in combination with the compounds of the invention, provided that the particular therapeutically active agent is compatible with therapy employing a compound of the invention. Typical therapeutically active agents which are suitable for use in combination with the compounds of the invention include agents described above.

In one preferred embodiment, the compounds of the invention are used in combination with one or more anti-inflammatory agents, particularly PDE4i or an inhaled corticosteroid. In one preferred embodiment, the compounds of the invention are used in combination with one or more anticholinergics, particularly muscarinic (M3) receptor antagonists.

30 In another aspect, the invention provides methods for treatment and uses as described above, which comprise administering an effective amount of a compound of the invention and at least one other therapeutically active agent. The compounds of the invention and at least one additional therapeutically

active agent may be employed in combination concomitantly or sequentially in any therapeutically appropriate combination. The administration of a compound of the invention with one or more other therapeutically active agents may be by administration concomitantly in 1) a unitary pharmaceutical composition, such as the compositions described above, or 2) separate pharmaceutical compositions each including one or more of the component active ingredients. The components of the combination may be administered separately in a sequential manner wherein the compound of the invention is administered first and the other therapeutically active agent is administered second or vice versa.

When a compound of the invention is used in combination with another therapeutically active agent, the dose of each compound may differ from that when the compound of the invention is used alone. Appropriate doses will be readily determined by one of ordinary skill in the art. The appropriate dose of the compound of the invention, the other therapeutically active agent(s) and the relative timings of administration will be selected in order to achieve the desired combined therapeutic effect, and are within the expertise and discretion of the attendant physician, clinician or veterinarian.

20

SYNTHETIC PROCESSES

The present invention also provides processes for preparing the compounds of the invention and to the synthetic intermediates useful in such processes, as described in detail below.

25

Certain abbreviations and acronyms are used in describing the synthetic processes and experimental details. Although most of these would be understood by one skilled in the art, the following table contains a list of many of these abbreviations and acronyms.

Abbreviation	Meaning
AIBN	azobisisobutyronitrile
Bn	benzyl
Boc	tert-butoxycarbonyl

Abbreviation	Meaning
(Boc) ₂ O	di-tert-butyl dicarbonate
BOP	benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate
Cbz	Carbobenzyloxy
DCC	N,N'-dicyclohexylcarbodiimide
DCE	dichloroethane
DCM	dichloromethane
DIEA	N,N-diisopropylethylamine
DME	dimethoxyethane
DMF	dimethylformamide
DMSO	dimethylsulfoxide
Et	ethyl
EtOAc	ethyl acetate
EtOH	ethanol
Et ₃ N and TEA	triethylamine
ESI	electrospray ionization
g	gram(s)
h	hour(s)
H ₂	hydrogen gas
HATU	2-(1H-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate
HPLC	High performance liquid chromatography
iPrOH	Isopropyl alcohol
LAH	lithium aluminum hydride
M	Molar
mg	milligram(s)
Me	methyl
MeOH	methanol
m/z or m/e	mass to charge ratio
MH ⁺	mass plus 1
MH ⁻	mass minus 1


Abbreviation	Meaning
MIC	minimal inhibitory concentration
min	minute(s)
mL	milliliter(s)
mmol	millimole(s)
MS or ms	mass spectrum
MsCl	methanesulfonyl chloride, mesyl chloride
Ms	methanesulfonate; mesylate
N	Normal
NaBH(OAc) ₃	sodium triacetoxy borohydride
NaCNBH ₃	sodium cyanoborohydride
NaN ₃	sodium azide
PDC	Pyridinium dichromate
Pd(OH) ₂ /C	Palladium hydroxide on carbon
Ph	phenyl
PMP	1,2,2,6,6-pentamethylpiperidine
PPh ₃	triphenylphosphine
PtO ₂	Platinum oxide
Py	Pyridyl or pyridine
rt or r.t.	room temperature (aka ambient temperature)
t-Bu	tert-butyl
TBAF	tetrabutylammoniumfluoride
TBS	tert-butyldimethylsilyl
TBSCl	tert-butyldimethylsilyl chloride
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TLC or tlc	thin layer chromatography
δ	parts per million down field from tetramethylsilane

In the following synthetic processes, it may be desirable for the preparation of certain compounds of the invention to install protecting groups on reactive sites of the intermediate. One skilled in the art will readily be able to determine the

desirability of using protecting groups, suitable protecting groups to employ based on the compounds and reaction conditions and methods for the installation and removal of such protecting groups. Suitable protecting groups include TBS, Bn, and Boc. Conventional techniques for installing and removing such protecting groups may be employed in the instant reaction as well.

A general procedure to prepare compounds of the invention is shown in Scheme 1 below.

10

Scheme 1

wherein:

Y is C(O), C(O)NHCH₂, N(R⁷)C(O)NHCH₂, OC(O)NHCH₂, or SO₂NHCH₂;

15

PG is a suitable protecting group, such as H or TBS; and all other variables are as defined above.

Generally, one process for preparing compounds of the invention comprises the steps of

- a) reductive alkylation of a compound **3** or a salt thereof with a compound **4** or a salt thereof to prepare compound **5** or a salt thereof; and
- 5 b) optionally deprotecting the compound **5** or a salt thereof, to prepare a compound of Formula I or a salt thereof.


Coupling of compound **1** with an acid under standard conditions, such as for example, HATU couplings, mixed anhydrides, DCC couplings, and the like,

- 10 gives the amide compound **2**. Coupling of Compound **1** with a phosgene equivalent, such as carbonyldiimidazole or 4-nitrophenylochloroformate at low temperature, between -78°C and 0°C, gives an activated species, which can be subsequently reacted with an appropriately substituted alcohol or amine at higher temperature, between rt and 100°C, to give compound **2** with a
- 15 carbamate or urea respectively. Compound **1** is known in the literature.

- Compound **2** is oxidized under standard conditions (Dess-Martin, PDC, Swern) to give the corresponding carbonyl compound **3**. Alternatively, compound **3** may be formed directly from compound **1** under standard conditions, such as for example, HATU coupling, mixed anhydride, DCC coupling, and the like, if
- 20 the appropriately substituted carbonyl containing acid is available. Alternatively, compounds **2** or **3** may be formed directly if the appropriately substituted acid chloride or sulfonyl chloride is available.

- Compound **4**, which is known in the literature, is coupled with compound **3** under reductive alkylation conditions, such as NaCNBH₃ or NaBH(OAc)₃, in an appropriate solvent, such as an alcohol or DMF, to give the corresponding compound **5**. Compound **5** is converted to the compounds of the invention by removal of any protecting group. A TBS protecting group was often used and in those cases, deprotection was accomplished using conventional techniques,
- 30 such as deprotection with TBAF.

In another embodiment, compounds of the invention may be prepared by displacement of a leaving group as shown in Scheme 2.

Scheme 2

wherein:


- Y is C(O)NHCH₂, N(R⁸)C(O)NHCH₂, OC(O)NHCH₂, or SO₂NHCH₂;
- 5 PG is a suitable protecting group, such as H or TBS;
- LG is a suitable leaving group such as bromide, chloride, iodide, OMs, or O-triflate;
- and all other variables are as defined above.

- 10 Generally, this process for preparing compounds of the invention comprises the steps of
- coupling a compound 6 or a salt thereof with a compound 4 or a salt thereof by alkylation to prepare a compound 5 or a salt thereof; and
 - optionally deprotecting the compound 5 or a salt thereof, to prepare a compound of Formula I or a salt thereof.
- 15

More specifically, according to this embodiment, the alcohol in compound 2 may be converted to a suitable leaving group under standard conditions to give compound 6. For example, conversion of the alcohol of compound 2 to a mesylate may occur through treatment of compound 2 with MsCl and an appropriate base, such as TEA or pyridine, in an appropriate solvent such as CH₂Cl₂ at rt. Alternatively, conversion of the alcohol of compound 2 to a bromide may occur under standard conditions such as CBr₄ and PPh₃. Alternatively, compound 6 may be formed directly under standard conditions,

by coupling compound **1** with the appropriately substituted acid that contains a leaving group. Compound **6** is then coupled with compound **4** at elevated temperatures, such as about 50 to about 150°C, in an appropriate solvent such as DMSO or DMF with an appropriate base, such as K₂CO₃, DIEA, or PMP, to give compound **5**. Compound **5** may then be deprotected to provide compounds of Formula I as described above in Scheme 1.

Scheme 3



10 wherein:

Y is C(O)NHCH₂, N(R⁸)C(O)NHCH₂, OC(O)NHCH₂, or SO₂NHCH₂;

PG is a suitable protecting group, such as H or TBS;

PG¹ is a suitable protecting group such as Cbz or Boc;

and all other variables are as defined above.

15

Generally, this process for preparing compounds of the invention comprises the steps of

- a) coupling a compound **8** or a salt thereof with a compound **9** or a salt thereof to prepare a compound **5** or a salt thereof; and
- 20 b) optionally deprotecting the compound **5** or a salt thereof, to prepare a compound of Formula I or a salt thereof.


Additionally, according to this embodiment, compound **1** may be coupled to an appropriately protected amino acid, which are commercially available or known,

- to produce compound **7**. Appropriate protecting groups include Boc or Cbz.
- Coupling of Compound **1** with a phosgene equivalent, such as carbonyldiimidazole or 4-nitrophenyloxchloroformate at low temperature, between -78°C and 0°C, gives an activated species, which can be
- 5 subsequently reacted with an appropriately substituted alcohol or amine at higher temperature, between rt and 100°C, to give compound **7** as a carbamate or urea respectively. Alternatively, compounds **7** may be formed directly if the appropriately substituted acid chloride or sulfonyl chloride is available. The protecting group may be removed under standard conditions to give compound
- 10 **8**.
- 8**. Reaction of compound **8** with the bromide compound **9** at elevated temperatures, such as about 50 to about 150 °C, neat or in an appropriate solvent such as DMSO or DMF, produces compound **5**.

Intermediate compounds of formula **7** wherein Y is N(R⁸)C(O) may be prepared

15 from the corresponding compound **10** according to the following Scheme 4. These intermediate compounds **7** may then be used as described above to prepare compounds of Formula I wherein Y is N(R⁸)C(O).

Scheme 4




- 20 wherein:
- Y is N(R⁸)C(O);
- PG¹ is a suitable protecting group such as Cbz or Boc;
- and all other variables are as defined above.

- 25 Coupling of compound **10** with an monoprotected diamine under standard conditions, such as for example, HATU couplings, mixed anhydrides, DCC couplings, and the like, gives the amide compound **7**. Compound **10** is known in the literature.

Compound **10** may also be used to prepare compound **2** wherein Y is N(R⁸)C(O) as shown in Scheme 5 below.

Scheme 5



5 wherein:

Y is C(O) or N(R⁸)C(O);

and all other variables are as defined above.

Coupling of compound **10** with an amino-alcohol under standard conditions,
10 such as for example, HATU couplings, mixed anhydrides, DCC couplings, and
the like, gives the amide compound **2**.


The order of steps in the foregoing reactions is not critical to the practice of the
present invention and the steps may be carried out in any suitable order
15 according to the knowledge of those skilled in the art, to provide the
compounds of formula I.

The foregoing detailed description may be further understood from the following
examples, which are presented for the purposes of illustration only and are not
20 intended to limit the scope of the invention. The invention is defined solely by
the claims which follow. In the following examples, compounds are named
using standard IUPAC naming principles where possible. The naming
convention employed for the novel compounds are exemplified by Examples
below.

25

EXAMPLES


Intermediate 1: (R)-5-[2-Azido-1-[(tert-butyldimethylsilyl)oxy]ethyl]-8-(benzyloxy)quinolin-2(1H)-one



NaN₃ (266 mg, 4.1 mmol) was added to a stirred solution of (R)-8-(benzyloxy)-5-[2-bromo-1-[(tert-butyldimethylsilyl)oxy]ethyl]quinolin-2(1H)-one (1g, 2.05 mmol) in DMF (20 mL) at rt and warmed to 80 °C for 3h. The resulting solution 5 was poured into H₂O (80 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with H₂O (2 x 100 mL), brine (100 mL), dried over Na₂SO₄ (s), and concentrated to give the title compound (1.37 g) as a yellow solid. The compound was used with no further purification. ES/MS calcd. for C₂₄H₃₁N₄O₃Si⁺ 451.2, found m/z = 451.3 (M+H)⁺.


10

Intermediate 2: (R)-5-[2-Amino-1-[(tert-butyldimethylsilyl)oxy]ethyl]-8-hydroxyquinolin-2(1H)-one



Intermediate 1 (1.37g) was dissolved in MeOH (20 mL) and Pd(OH)₂/C (20 % 15 w/w, 288 mg, 0.41 mmol) was added. Nitrogen gas was bubbled through the solution for 5 min. The resulting suspension was attached to a balloon filled with H₂ and stirred over night. The reaction mixture was filtered through celite and concentrated to give a brown oil (1.208 g). Chromatography (9:1, CH₂Cl₂/MeOH, 0.1 % Et₃N) afforded the title compound (597 mg, 87 % 2 steps) 20 as a light yellow solid. ES/MS calcd. for C₁₇H₂₇N₂O₃Si⁺ 335.2, found m/z = 335.2 (M+H)⁺.


Intermediate 3: (R)-5-(2-Azido-1-hydroxyethyl)-8-(benzyloxy)quinolin-2(1H)-one



25 TBAF (1.0 M in THF, 0.443 mL, 0.443 mmol) was added to a stirring solution of Intermediate 1 (200 mg, 0.443 mmol) in THF (4 mL) at rt. The resulting mixture was stirred over night then concentrated. Chromatography (1:3,

Hexanes/EtOAc) afforded the title compound (137 mg, 92 %) as an off-white solid. ES/MS calcd. for $C_{18}H_{17}N_4O_3^+$ 337.1, found $m/z = 337.2 (M+H)^+$.


Intermediate 4: (R)-5-(2-Amino-1-hydroxyethyl)-8-hydroxyquinolin-2(1H)-one



5


The title compound was synthesized in a manner analogous to that described for Intermediate 2, using Intermediate 3 as a substrate. ES/MS calcd. for $C_{11}H_{13}N_2O_3^+$ 221.1, found $m/z = 221.1 (M+H)^+$.

10 Intermediate 5: (R)-1-[4-(Benzylxy)-3-[(tert-
butyldimethylsilyl)oxy]methyl]phenyl]-2-bromoethanol




TBSCl (5 g, 33.2 mmol) and imidazole (3.7 g, 55.4 mmol) were added to a stirring solution of (R)-4-(2-bromo-1-hydroxyethyl)-2-(hydroxymethyl)phenol (10 g, 27.7 mmol) in CH_2Cl_2 (200 mL) at rt. The resulting suspension was stirred for 1 h then quenched with H_2O (200 mL). The aqueous layer was extracted with CH_2Cl_2 (3 x 100 mL). The combined organic layer was washed with brine (300 mL), dried over Na_2SO_4 (s), and concentrated to give title compound (13.9 g) as a clear oil. The compound was used with no further purification. ES/MS calcd. for $C_{22}H_{31}BrNaO_3Si^+$ 473.1, found $m/z = 473.1 (M+Na)^+$.

15 Intermediate 6: (R)-[1-[4-(Benzylxy)-3-[(tert-
butyldimethylsilyl)oxy]methyl]phenyl]-2-bromoethoxy](tert-butyl)dimethylsilane




20 The title compound was synthesized in a manner analogous to that described for Intermediate 5. Excess TBSCl and a higher reaction temperature were used.

Intermediate 7: (R)-2-Azido-1-[4-(benzyloxy)-3-[(tert-butylidimethylsilyl)oxy]methyl]phenyl]ethanol



The title compound was synthesized in a manner analogous to that described
 5 for Intermediate 1, using Intermediate 5 in place of (R)-8-(benzyloxy)-5-[2-bromo-1-[(tert-butylidimethylsilyl)oxy]ethyl]quinolin-2(1H)-one. ES/MS calcd. for C₂₂H₃₁N₃NaO₃Si⁺ 436.2, found m/z = 436.2 (M+Na)⁺.


10 Intermediate 8: (R)-4-(2-Amino-1-hydroxyethyl)-2-[(tert-butylidimethylsilyl)oxy]methyl]phenol



The title compound was synthesized in a manner analogous to that described in Intermediate 2, using Intermediate 7 as a substrate. ES/MS calcd. for C₁₅H₂₇NNaO₃Si⁺ 320.2, found m/z = 320.2 (M+Na)⁺.

15


Intermediate 9: (R)-N-[2-(Benzylxy)-5-(2-bromo-1-hydroxyethyl)phenyl]formamide



(R)-1-[4-(Benzylxy)-3-nitrophenyl]-2-bromoethanol (0.2 g, 5.7 mmol) in
 20 THF:toluene (1:1, 5 mL) was reacted with PtO₂ (1% w/w) on a Parr shaker at 45 psi at rt overnight. The next morning the PtO₂ was removed by filtration over celite. The filtered solution was cooled to 0 °C and a solution of acetic anhydride (0.161 mL, 0.569 mmol) and formic acid (0.043 mL, 1.140 mmol) was added dropwise to a mixing solution of the aniline. The reaction was
 25 allowed to proceed at 0 °C for 30 min then warmed to rt and reacted for another 2 h. The reaction mixture was concentrated to near dryness then water was added. The aqueous layer was extracted with EtOAc (3x 25 mL). The combined organic layers were washed with NaHCO₃, brine, dried over Na₂SO₄

(s), and concentrated to give a residue. Chromatography (1:1 Hexanes/EtOAc) gives the title compound (156 mg, 78% 2 steps). ES/MS calcd. $C_{16}H_{17}BrNO_3^+$ 350.0, found $m/z = 350$ ($M+H$)⁺.


- 5 Intermediate 10: (R)-N-[5-(2-Azido-1-hydroxyethyl)-2-(benzyloxy)phenyl]formamide



The title compound was synthesized in a manner analogous to that described for Intermediate 1, using Intermediate 9 as substrate. ES/MS calcd.


- 10 $C_{16}H_{17}N_4O_3^+$ 313.1, found $m/z = 313$ ($M+H$)⁺.

- Intermediate 11: (R)-N-[5-[2-Azido-1-[(tert-butyldimethylsilyl)oxy]ethyl]-2-(benzyloxy)phenyl]formamide



- 15 The title compound was synthesized in a manner analogous to that described for Intermediate 5, using Intermediate 10 in place of (R)-4-(2-bromo-1-hydroxyethyl)-2-(hydroxymethyl)phenol. ES/MS calcd. $C_{22}H_{31}N_4O_3Si^+$ 427.2, found $m/z = 427$ ($M+H$)⁺.


- 20 Intermediate 12: (R)-N-[5-[2-Amino-1-[(tert-butyldimethylsilyl)oxy]ethyl]-2-hydroxyphenyl]formamide



The title compound was synthesized in a manner analogous to that described for Intermediate 2, using Intermediate 11 in place of Intermediate 1. ES/MS

- 25 calcd. $C_{15}H_{27}N_2O_3Si^+$ 311.2, found $m/z = 311$ ($M+H$)⁺

- Intermediate 13: (R)-1-[3-Amino-4-(benzyloxy)phenyl]-2-bromoethanol



(R)-1-[4-(benzyloxy)-3-nitrophenyl]-2-bromoethanol (0.200g, 0.569 mmol) in 1:1 THF:toluene (5 mL) was reacted with 1% (w/w) PtO₂ on a Parr shaker at 45 psi at rt overnight. The next morning the PtO₂ was removed by filtration over celite.


- 5 The product was concentrated to give the title compound. ES/MS calcd. for C₁₅H₁₇BrNO₂⁺ 322.0, found m/z = 322 (M+H)⁺.

Intermediate 14: (R)-1-[3-Amino-4-(benzyloxy)phenyl]-2-azidoethanol




- 10 The title compound was synthesized in a manner analogous to that described for Intermediate 1, using Intermediate 13 in place of (R)-8-(benzyloxy)-5-[2-bromo-1-[(tert-butyldimethylsilyl)oxy]ethyl]quinolin-2(1H)-one. ES/MS calcd. for C₁₅H₁₇N₄O₂⁺ 285.1, found m/z = 285 (M+H)⁺.

- 15 Intermediate 15: of (R)-8-(BenzylOxy)-5-[2-bromo-1-[(tert-butyldimethylsilyl)oxy]ethyl]quinolin-2(1H)-one



- The title compound was synthesized in a manner analogous to that described for Intermediate 5, using Intermediate 14 in place of (R)-4-(2-bromo-1-hydroxyethyl)-2-(hydroxymethyl)phenol. ES/MS calcd. for C₂₁H₃₁N₄O₂Si⁺ 399.2, found m/z = 399 (M+H)⁺.


Intermediate 16: (R)-N-[5-[2-Azido-1-[(tert-butyldimethylsilyl)oxy]ethyl]-2-(benzyloxy)phenyl]methanesulfonamide



Methanesulfonyl chloride (0.044 mL, 0.569 mmol) was added to a stirring solution of Intermediate 15 (275 mg, 0.569 mmol) in pyridine (10 mL) at 0°C. The resulting mixture was warmed to rt and monitored for completeness by LC/MS. An additional 1 equivalent of MsCl was added after 1 h followed by an 5 additional 0.5 equivalent after another hour for a total of 2.5 equivalents. After an additional 1 h, H₂O (50 mL) was added and stirred at rt for 2 h. The reaction was extracted with CH₂Cl₂ (4x 25 mL). The combined organic layers were washed with satd. NaHCO₃, brine, dried over Na₂SO₄ (s), and concentrated. Chromatography (1:1 Hexanes/EtOAc) afforded the title compound (197 mg, 10 72%, 3 steps). ES/MS calcd. for C₂₂H₃₃N₄O₄SSi⁺ 477.2, found m/z = 477 (M+H)⁺


Intermediate 17: (R)-N-[5-[2-Amino-1-[(tert-butyldimethylsilyl)oxy]ethyl]-2-hydroxyphenyl]methanesulfonamide

15



20

Intermediate 18: 3-(2-Hydroxy-2-methylpropyl)benzoic acid




CH₃MgBr (3.0 M in THF, 11.2 mL, 33.7 mmol) was added slowly to a stirring solution of 3-(2-oxopropyl)benzoic acid (2 g, 11.2 mmol) in THF (100 mL) at 0 °C. The resulting suspension was allowed to warm to rt then stirred overnight. The reaction mixture was diluted with H₂O (50 mL) then quenched with 5 N HCl (pH < 3). The aqueous layer was extracted with EtOAc (3 X 50 mL). The combined organic layers were washed with H₂O (100 mL), brine (100 mL), dried over Na₂SO₄ (s), and concentrated to give a yellow oil. Chromatography

(9:1, CH₂Cl₂/MeOH) afforded the title compound (1.36 g, 62 %) as a white solid. ES/MS calcd. for C₁₁H₁₄NaO₃⁺ 217.1, found m/z = 217.2 (M+Na)⁺.

Intermediate 19: methyl 3-(2-Amino-2-methylpropyl)benzoate


5



Chloroacetonitrile (0.845 mL, 13.4 mmol) and conc. H₂SO₄ (3 mL, 55.2 mmol) were added to a stirring solution of Intermediate 18 (1.3 g, 0.69 mmol) in glacial acetic acid (3 mL) at 0 °C. The resulting mixture was allowed to warm to rt then stirred for an additional 2 h. The reaction mixture was poured into ice and 10 made basic with K₂CO₃ (s) (pH ~ 4). The aqueous layer was extracted with EtOAc (3 X 50 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄ (s), and concentrated to give an off white solid. The solid was redissolved in glacial acetic acid (20 mL) and thiourea (904 mg, 11.9 mmol) was added. The reaction mixture was warmed to 95 °C for 3 h, cooled, 15 then concentrated. The residue was dissolved in MeOH (20 mL) and HCl (g) was bubbled through for 5 min. The resulting solution was refluxed overnight, cooled, and then concentrated. The residue was diluted with Et₂O (100 mL) and washed with H₂O (100 mL). The aqueous layer was made basic with satd. NaHCO₃ (pH > 10) and then extracted with CHCl₃ (3 x 50 mL). The combined 20 organic layers were washed with brine (100 mL), dried over Na₂SO₄ (s), and concentrated to the title compound (1.26 g) as a clear oil. The compound was used with no further purification. ES/MS calcd. for C₁₂H₁₈NO₂⁺ 208.1, found m/z = 208.2 (M+H)⁺.

25

Intermediate 20: Methyl 4-(5-hydroxypent-1-yn-1-yl)benzoate




Methyl 4-iodobenzoate (13.1 g, 60 mmol), PdCl₂(PPh₃)₂ (2.0 g, 3 mmol), and CuI (855 mg, 4.5 mmol) were added to TEA (60 mL). The mixture is degassed while stirring by bubbling N₂ vigorously into the solution. Pent-4-yn-1-ol (5.5 g, 30 66 mmol) was added by syringe. The mixture was heated at 90 °C overnight. After cooling, the crude mixture was concentrated by rotary evaporation and

pumped on the high vacuum for 1 h. The oil was dissolved in DCM and adsorbed onto Celite followed by purification by chromatography (gradient 10 - 100% EtOAc/Hexanes) to give the title compound as a light yellow powder (9 g, 69%). ¹H NMR (400 MHz, CDCl₃) δ 7.95 (dd, 2H, J = 9.0, 1.5 Hz), 7.44 (dd, 5 2H, J = 9.0, 1.5 Hz), 3.90 (s, 3H), 3.85 (m, 2H), 2.57 (t, 2H, J = 5.5 Hz), 1.87 (m, 2H), 1.42 (t, 1H, J = 5.5 Hz); ES/MS calcd. for C₁₃H₁₅O₃⁺ 219.1, found m/z = 219 (M+H)⁺.

Intermediate 21: 4-(5-Hydroxypent-1-yn-1-yl)benzoic acid

10



LiOH (2 g, 83 mmol) in H₂O (30 mL) was added dropwise to a stirring solution of Intermediate 20 (2 g, 9.2 mmol) in THF (30 mL). The resulting mixture was stirred at 42 °C for 4 h. After cooling, water (100 mL) and EtOAc (200 mL) were added. The mixture was extracted with EtOAc. The organic layer and the H₂O layer were independently neutralized with 1N HCl. The aqueous layer was extracted with EtOAc and the combined organic layers dried over MgSO₄(s). The material was concentrated by rotary evaporation and the crude solid was purified by trituration with EtOAc and hexanes to give the title compound as a white powder (1.75 g, 93%). ¹H NMR (400 MHz, DMSO-d₆) δ 13.05 (brs, 1H), 7.88 (dd, 2H, J = 8.5, 1.5 Hz), 7.47 (dd, 2H, J = 8.5, 1.5 Hz), 4.55 (brs, 1H), 3.50 (t, 2H, J = 5.5 Hz), 2.50 (m, 2H), 1.68 (m, 2H); ES/MS calcd. for C₁₂H₁₁O₃⁺ 205.1, found m/z = 205 (M+H)⁺.


25 Intermediate 22: 1-(5-chloropent-1-yn-1-yl)-4-nitrobenzene

A mixture of 4-iodonitrobenzene (25 g, 100 mmol), dichlorobis(triphenylphosphine)palladium(II) (0.30 g, 0.43 mmol) and copper (I) iodide (0.18 g, 0.95 mmol) in tetrahydrofuran (300 mL) and triethylamine (150 mL) was degassed under Argon for 10 minutes. The mixture was then heated to 55 °C and treated with 5-chloropentyne (12 mL, 110 mmol) via syringe. After 30 minutes of stirring, additional quantities of

dichlorobis(triphenylphosphine)palladium(II) (0.10 g) and 5-chloropentyne (1 mL) were added. After another 30 minutes of stirring, the mixture was allowed to cool to room temperature and then was filtered through a pad of Celite diatomaceous earth. The filtrate was concentrated to dryness under reduced pressure to provide 1-(5-chloropent-1-ynyl)-4-nitrobenzene, which was carried on without further purification. ES/MS calcd. for $C_{11}H_{11}ClNO_2^+$ 224.1, found m/z = 224.2 ($M+H$)⁺.

Intermediate 23: 1-(5-bromopent-1-yn-1-yl)-4-nitrobenzene

10



15


A solution of crude Intermediate 22 (approximately 200 mmol) in 3-pentanone (1 L) was treated with lithium bromide (10 eq, 174 g, 2 moles). The mixture was heated to reflux for 30 minutes, followed by concentration to dryness under reduced pressure. The residue was partitioned between ethyl acetate and water. The aqueous phase was extracted with ethyl acetate. The combined organics was washed with a 1 % aqueous hydrochloric acid solution and with saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and concentrated to dryness under reduced pressure. The residue was taken up again in 3-pentanone (1 L) and heated to reflux overnight in the presence of lithium bromide (250 g). The mixture was allowed to cool to room temperature and concentrated to dryness under reduced pressure. The residue was partitioned between ethyl acetate and water. The organic phase was washed with saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness under reduced pressure. The crude product was carried on without further purification. ES/MS calcd. for $C_{11}H_{11}BrNO_2^+$ 268.0, found m/z = 268.1 ($M+H$)⁺.

20

25

Intermediate 24: 4-(5-bromopent-1-yn-1-yl)aniline


30



A mixture of crude Intermediate 23 (200 mmol) was dissolved in a mixture of *N*-methylpyrrolidine/dichloromethane (1:1, 800 mL) and treated with tin (II)

chloride dehydrate (218, 960 mmol) in 30 g portions. The exothermic reaction mixture was cooled in an ice-water bath. Following completion of the tin chloride addition, the cooling bath was removed and the mixture was allowed to regain room temperature. After 45 minutes of stirring, the mixture was
5 quenched by adding it portion-wise to a mixture of ice and concentrated ammonium hydroxide solution. The slurry was filtered through a fritted glass funnel, washing with dichloromethane. The filtrates were concentrated under reduced pressure and diluted with diethyl ether. This organic phase was washed four times with water and once with a saturated aqueous sodium
10 chloride solution, dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness under reduced pressure. The residue was taken up in ethyl acetate and filtered through a ceramic filter. The filtrate was concentrated and the residue was triturated with benzene and filtered. The concentration-trituration cycle was performed five times. After the final iteration, the
15 concentrated filtrate was purified by automated flash silica gel chromatography (hexanes/ethyl acetate). The desire fractions were concentrated to a lesser volume and then treated dropwise with methanesulfonic acid until the precipitation of further material appeared to cease. The solid was collected by Büchner filtration, washed with diethyl ether and ethyl acetate, and dried under
20 house vacuum to provide the methanesulfonic acid salt of the title material as an off-white solid. ES/MS calcd. for $C_{11}H_{13}BrN^+$ 240.0, found $m/z = 240.1$ $(M+H)^+$.

Intermediate 25: N-(4-(5-chloropent-1-yn-1-yl)phenyl)-2,2,2-trifluoroacetamide




25

A mixture of 2,2,2-trifluoro-N-(4-iodophenyl)acetamide (3.2 g, 10 mmol, prepared according to Melissaris, A.P. and Litt, M.H. J Org Chem 1994, 59, 5818-5821), dichlorobis(triphenylphosphine)palladium(II) (0.14 g, 0.2 mmol), triphenylphosphine (19 mg, 0.07 mmol), and copper (I) iodide (0.019 g, 0.1 mmol) in triethylamine (50 mL) was degassed under Argon for 10 minutes while stirring in a 55 °C oil bath. The mixture was then treated with 5-chloropentyne (1.1 mL, 10 mmol). After four hours of heating, the mixture was allowed to cool


to room temperature and was then filtered through a pad of Celite diatomaceous earth. The filtrate was concentrated under reduced pressure to provide the crude title compound, which was carried forward without further purification. ES/MS calcd. for $C_{14}H_{14}ClF_3NO^+$: 290.1, found m/z = 290.1
 5 $(M+H)^+$

Intermediate 26: N-(4-(5-bromopent-1-yn-1-yl)phenyl)-2,2,2-trifluoroacetamide



A solution of Intermediate 25 (approximately 10 mmol) in 3-pentanone (200 mL) was treated with lithium bromide (10 eq, 100 mmol). The mixture was heated to reflux for 16 hours, followed by concentration to dryness under reduced pressure. The residue was taken up in ethyl acetate and washed with water. The concentrated organic phase was taken up again in 3-pentanone (200 mL) and heated to reflux for four hours in the presence of lithium bromide
 10 (10 eq, 100 mmol). The mixture was concentrated to dryness under reduced pressure. The residue was partitioned between ethyl acetate and water. The organic phase was washed with saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness under reduced pressure. The crude product was purified via automated flash
 15 silica gel chromatography, using a 40g SiliSep flash column (hexanes/ethyl acetate). Concentration of the desired fractions under reduced pressure provided the title compound as an off-white solid. ES/MS calcd. for
 20 $C_{13}H_{12}BrF_3NO^+$ 334.0, found m/z = 334.1 ($M+H$)⁺.
 25

Intermediate 27: 1-(4-((6-bromohexyl)oxy)butyl)-4-nitrobenzene




A stirred mixture of 4-(4-nitrophenyl)butan-1-ol (2.0 g, 10 mmol), tetra-*n*-butylammonium hydrogen sulfate (0.17 g, 0.5 mmol), and 1,6-dibromohexane (3.2 mL, 20 mmol) in dichloromethane (10 mL) was treated with aqueous
 30 sodium hydroxide solution (10 M, 1 mL). The reaction mixture was stirred for 6

days at room temperature. The organic and aqueous phases were separated. The aqueous phase was extracted thrice with dichloromethane. The combined organics were washed with water, dried over anhydrous magnesium sulfate, filtered, and concentrated under reduced pressure. The crude residue was 5 purified via automated flash silica gel chromatography, using a 40g SiliSep flash column (hexanes/ethyl acetate). Concentration of the desired fractions under reduced pressure provided the title compound as a pale straw-colored oil (0.72 g, 20 %). ES/MS calcd. for $C_{16}H_{25}BrNO_3^+$ 358.1, found m/z = 358.2 ($M+H$)⁺.


10

Intermediate 28: 3-(((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)amino)-4-ethoxycyclobut-3-ene-1,2-dione




A suspension of 5-(aminomethyl)-1,6-diethyl-N-(tetrahydro-2H-pyran-4-yl)-1H-pyrazolo[3,4-b]pyridin-4-amine dihydrochloride (0.60 g, 1.6 mmol) in ethanol (8 mL) was treated with *N,N*-diisopropylethylamine (DIEA, 0.57 mL, 3.3 mmol), giving a clear, homogeneous mixture, which was then cooled in an ice-water bath. 3,4-Diethoxycyclobut-3-ene-1,2-dione (TCI America, 0.41 g, 0.36 mmol) was then added dropwise via syringe. Following completion of the addition, the 15 mixture was allowed to regain room temperature and stir overnight. The precipitated solid was collected by Büchner filtration, washed with ethanol, and dried under house vacuum to provide the title compound as a white solid. 20 ES/MS calcd. for $C_{22}H_{30}N_5O_4^+$ 428.2, found m/z = 428.3 ($M+H$)⁺.

25 Intermediate 29: N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-(2-oxopropyl)benzamide



HATU (202 mg, 0.531 mmol) and DIEA (0.294 mL, 1.69 mmol) were added to a stirring solution 3-(2-oxopropyl)benzoic acid (86 mg, 0.483 mmol) in DMF (5 mL) at rt. After 5 min, 5-(aminomethyl)-1,6-diethyl-N-(tetrahydro-2H-pyran-4-yl)-1H-pyrazolo[3,4-b]pyridin-4-amine (200 mg, 0.531 mmol) was added and 5 the resulting solution was stirred an additional 2h. The reaction mixture was poured into H₂O (50 mL) and the aqueous layer was extracted with EtOAc (3 x 30 mL). The combined organic layers were washed with H₂O (2 x 100 mL), brine (100 mL), dried over Na₂SO₄ (s), and concentrated to give a yellow oil 10 (376 mg). Chromatography (9:1, CH₂Cl₂/MeOH, 0.1 % Et₃N) afforded the title compound (212 mg, 95 %) as a yellow solid. ES/MS calcd. for C₂₆H₃₄N₅O₃⁺ 464.3, found m/z = 464.3 (M+H)⁺.

Intermediate 30: N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-formylbenzamide




15

The title compound was synthesized in a manner analogous to that described for Intermediate 29, using 3-formylbenzoic acid in place of 3-(2-oxopropyl)benzoic acid. ES/MS calcd. for C₂₄H₃₀N₅O₃⁺ 436.2, found m/z = 436.2(M+H)⁺.

20


Intermediate 31: N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-formylbenzamide



25


The title compound was synthesized in a manner analogous to that described for Intermediate 29, using 4-formylbenzoic acid in place of 3-(2-oxopropyl)benzoic acid. ES/MS calcd. for C₂₄H₃₀N₅O₃⁺ 436.2, found m/z = 436.2(M+H)⁺.

Intermediate 32: N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-(5-hydroxypent-1-yn-1-yl)benzamide




- 5 The title compound was synthesized in a manner analogous to that described for Intermediate 29, using Intermediate 21 in place of 3-(2-oxopropyl)benzoic acid. ES/MS calcd. for $C_{28}H_{36}N_5O_3^+$ 490.3, found $m/z = 490.4$ ($M+H$) $^+$.

Intermediate 33: 8-Bromo-N-[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]octanamide



- 10 The title compound was synthesized in a manner analogous to that described for Intermediate 29, using 8-bromo-octanoic acid in place of 3-(2-oxopropyl)benzoic acid. ES/MS calcd. for $C_{24}H_{39}BrN_5O_2^+$ 508.2, found $m/z = 508.2$ ($M+H$) $^+$.


Intermediate 34: N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-(5-hydroxypentyl)benzamide



- 15 Pd(OH)₂ (200mg) was added to a solution of Intermediate 32 (200 mg, 0.40 mmol) in THF/MeOH (5 mL each). The solution was treated with H₂ at ambient pressure for 3 h. The reaction mixture was then filtered through a plug of celite. The plug was washed with MeOH (50 mL) and the solvent was concentrated *in*


vacuo to afford the title compound as a pale yellow solid (200 mg). ES/MS calcd. for $C_{28}H_{40}N_5O_3^+$ 494.3, found $m/z = 494.4 (M+H)^+$.

- 5 Intermediate 35: 1-(4-(5-bromopent-1-yn-1-yl)phenyl)-3-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)urea



A suspension of Intermediate 24 methanesulfonic acid salt (1.4 g, 4.2 mmol) in dichloromethane (50 mL) was treated with triethylamine (8.8 mL, 63 mmol) and then the mixture was cooled in an ice-water bath and treated with triphosgene (0.93 g, 3.2 mmol), followed by 5-(aminomethyl)-1,6-diethyl-N-(tetrahydro-2H-pyran-4-yl)-1H-pyrazolo[3,4-b]pyridin-4-amine dihydrochloride (1.3 g, 3.4 mmol). The mixture was quenched by the addition of methanol and concentrated under reduced pressure. The residue was partitioned between ethyl acetate and water. The aqueous phase was extracted twice with ethyl acetate. The combined organics were washed with a 5 % aqueous potassium hydrogen sulfate solution and saturated aqueous sodium chloride solution and then concentrated under reduced pressure. The crude residue was purified via automated flash silica gel chromatography, using a SiliSep flash column (dichloromethane/methanol/ammonium hydroxide). Concentration of the desired fractions under reduced pressure provided the title compound as a pale tan foam. ES/MS calcd. for $C_{28}H_{36}BrN_6O_2^+$ 567.2, found $m/z = 567.2 (M+H)^+$.

- 15 Intermediate 36: N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-(5-oxopentyl)benzamide




20 Dess Martin Reagent (432mg, 1.02 mmol) was added to a solution of Intermediate 34 (250 mg, 0.51 mmol) in THF (5 mL) and DCM (5 mL). The

resultant mixture was stirred at rt for 2 h. The solvent was removed *in vacuo* and the crude solid was purified by column chromatography on silica gel eluting with 0 -10% DCM – MeOH to afford the title compound as a white solid (250mg). ES/MS calcd. for $C_{28}H_{38}N_5O_3^+$ 492.3, found $m/z = 492.3$ ($M+H$)⁺.


5

Intermediate 37: (R)-N-(4-(5-((2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pent-1-yn-1-yl)phenyl)-2,2,2-trifluoroacetamide



- 10 A mixture of Intermediate 26 (0.33 g, 1.0 mmol) and Intermediate 2 (0.40 g, 1.2 mmol) in DMF (5 mL) was treated with *N,N*-diisopropylethylamine (DIEA, 0.25 mL, 1.5 mmol) and catalytic potassium iodide. The mixture was heated overnight in a 55 °C oil bath. The mixture was concentrated under reduced pressure and purified via automated flash silica gel chromatography, using a 15 12g Isco RediSep flash column (dichloromethane/methanol/ammonium hydroxide). Concentration of the desired fractions under reduced pressure provided the title compound (0.53 g, 90 %). ES/MS calcd. for $C_{30}H_{37}F_3N_3O_4Si^+$ 588.3, found $m/z = 588.4$ ($M+H$)⁺.

- 20 Intermediate 38: (R)-tert-butyl (5-(4-aminophenyl)pent-4-yn-1-yl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)carbamate




- 25 A solution of Intermediate 37 (approximately 0.35 g, 0.60 mmol) in dichloromethane (5 mL) was treated successively with triethylamine (0.42 g,

3.0 mmol), di-*tert*-butyldicarbonate (0.39 g, 1.8 mmol), and *N,N*-dimethylaminopyridine (DMAP, 50 mg). After two hours the mixture was concentrated to dryness under reduced pressure and taken up in methanol/water (2:1, 5 mL). Tetrahydrofuran was added dropwise until a homogeneous mixture was obtained, which was then treated with potassium carbonate (approximately 500 mg). The mixture was left stirring overnight at room temperature. It was concentrated to an aqueous suspension under reduced pressure and then adjusted to pH 7 with the addition of a 5 % aqueous potassium hydrogen sulfate solution. The aqueous mixture was extracted 5 thrice with dichloromethane. The combined extracts were concentrated to a reddish-brown foam, which was then purified by automated flash silica gel chromatography, using a 12g Silicycle SiliSep flash column (dichloromethane/methanol/ammonium hydroxide). Concentration of the desired fractions under reduced pressure provided the title compound (0.11 g, 10 30 %) as a pale tan foam. ES/MS calcd. for $C_{33}H_{46}N_3O_5Si^+$ 592.3, found *m/z* = 592.4 ($M+H$)⁺.

15

Intermediate 39: (R)-8-hydroxy-5-(2,2,3,3-tetramethyl-18-(4-nitrophenyl)-4,14-dioxa-7-aza-3-silaoctadecan-5-yl)quinolin-2(1H)-one


20



The title compound was prepared in a manner analogous to that described for Intermediate 37, using Intermediate 27 in place of Intermediate 26. ES/MS calcd. for $C_{33}H_{50}N_3O_6Si^+$ 612.4, found *m/z* = 612.5 ($M+H$)⁺.


25

Intermediate 40: (R)-*tert*-butyl (2-(8-((*tert*-butoxycarbonyl)oxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(6-(4-(4-nitrophenyl)butoxy)hexyl)carbamate



A solution of Intermediate 39 (0.53 g, 0.87 mmol) in dichloromethane (5 mL) was treated with di-*tert*-butyldicarbonate (0.57 g, 2.6 mmol), triethylamine (0.61 mL, 4.4 mmol) and *N,N*-dimethylaminopyridine (DMAP, 20 mg). The reaction mixture was stirred overnight at room temperature and then concentrated to dryness under reduced pressure. The residue was purified via automated flash silica gel chromatography, using a 25 g SiliSep flash column (hexanes/ethyl acetate). Concentration of the desired fractions under reduced pressure provided the title compound as clear, colorless oil, which turned into a white foam under reduced pressure. ES/MS calcd. for C₄₃H₆₆N₃O₁₀Si⁺ 812.5, found m/z = 812.6 (M+H)⁺.


Intermediate 41: (R)-tert-butyl (2-(2,8-bis((tert-butoxycarbonyl)oxy)quinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(6-(4-(4-nitrophenyl)butoxy)hexyl)carbamate



A solution of Intermediate 40 (approximately 150 mg) in dichloromethane (5 mL) was treated with di-*tert*-butyldicarbonate and triethylamine (both q.s.) and catalytic *N,N*-dimethylaminopyridine (DMAP) at room temperature. Upon LC/MS analysis of a complete tris-Boc protection, the mixture was concentrated to dryness under reduced pressure, and the residue was purified by automated flash silica gel chromatography, using a 25g SiliSep flash column (hexanes/ethyl acetate). Concentration of the desired fractions under reduced pressure provided the title compound (175 mg) as sticky colorless foam. ES/MS calcd. for C₄₈H₇₃N₃NaO₁₂Si⁺ 934.5, found m/z = 934.5 (M+Na)⁺.


Intermediate 42: (R)-tert-butyl (6-(4-(4-aminophenyl)butoxy)hexyl)(2-(2,8-bis((tert-butoxycarbonyl)oxy)quinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)carbamate

5




Intermediate 41 (175 mg, 0.19 mmol) was dissolved in methanol (4 mL) and ethyl acetate (1 mL) and treated with palladium on carbon (10 % w/w, wet Degussa-type, catalytic amount). The suspension was stirred under a balloon of hydrogen gas for 45 minutes. The mixture was then filtered through a pad of 10 Celite diatomaceous earth, and the filtrate was concentrated to dryness under reduced pressure to provide the title material. ES/MS calcd. for $C_{48}H_{76}N_3O_{10}Si^+$ 882.5, found $m/z = 882.5$ ($M+H$)⁺.

15 Intermediate 43: 3-[2-[(R)-2-[3-[(tert-Butyldimethylsilyl)oxy]methyl]-4-hydroxyphenyl]-2-hydroxyethyl]amino]propyl]-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide




Glacial acetic acid (0.024 mL, 0.421 mmol) and Intermediate 8 (125 mg, 0.421 mmol) were added to a stirring solution of Intermediate 29 (130 mg, 0.28 mmol) 20 in DMF (3 mL) at rt. The resulting solution was stirred for 2 h. $NaBH(OAc)_3$ (178 mg, 0.84 mmol) was added in portions. The reaction mixture was stirred overnight then poured into satd. $NaHCO_3$ (40 mL). The precipitate was filtered, washed with H_2O (50 mL), and dried to give the title compound (118 mg) as an off-white solid. The compound was used with no further purification. ES/MS calcd. for $C_{41}H_{61}N_6O_5Si^+$ 745.5, found $m/z = 745.5$ ($M+H$)⁺.

Intermediate 44: (R)-4-[5-[2-[(tert-Butyldimethylsilyl)oxy]methyl]-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]pentyl-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide




5 NaCNBH₃ (160 mg, 2.55 mmol) was added to a solution of Intermediate 2 (240 mg, 0.61 mmol), Intermediate 36 (250 mg, 0.51 mmol), and acetic acid (0.032 mL, 0.561 mmol) in THF (5 mL). The reaction mixture was stirred at rt for 1 h. H₂O (2 mL) was added followed by satd. NaHCO₃ solution (20 mL) and EtOAc (100 mL). The organic layer was separated, dried over MgSO₄ (s), filtered, and concentrated *in vacuo*. Purification by column chromatography on silica gel eluting with a gradient of 0 – 100% DCM – DCM/MeOH/Ammonia (80/20/1) afforded the title compound as a yellow solid (200 mg). ES/MS calcd. for C₄₅H₆₄N₇O₅Si⁺ 810.5, found m/z = 810.5 (M+H)⁺.

15 Intermediate 45: (R)-3-[[2-(3-[(tert-Butyldimethylsilyl)oxy]methyl)-4-hydroxyphenyl]-2-hydroxyethyl]amino]methyl]-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide



The title compound was synthesized in a manner analogous to that described, 20 using Intermediate 30 in place of Intermediate 29. ES/MS calcd. for C₃₉H₅₆N₆NaO₅Si⁺ 739.4, found m/z = 739.4 (M+Na)⁺.


25 Intermediate 46: (R)-4-[[2-(3-[(tert-Butyldimethylsilyl)oxy]methyl)-4-hydroxyphenyl]-2-hydroxyethyl]amino]methyl]-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide



The title compound was synthesized in a manner analogous to that described for Intermediate 43, using Intermediate 31 in place of Intermediate 29. ES/MS calcd. for $C_{39}H_{56}N_6NaO_5Si^+$ 739.4, found $m/z = 739.4$ ($M+Na$)⁺.

5


Intermediate 47: (R)-Methyl 3-[2-[(2-[4-(benzyloxy)-3-[(tert-butyldimethylsilyl)oxy]methyl]phenyl]-2-[(tert-butyldimethylsilyl)oxy]ethyl]amino]-2-methylpropyl]benzoate



10 Intermediate 6 (1.6 g, 2.89 mmol) and Intermediate 19 (900 mg, 4.3 mmol) were heated neat at 95 °C for 3 days. The reaction mixture was cooled then chromatography (3:1 Hexanes/EtOAc, 0.1 % Et₃N) afforded the title compound (730 mg, 37 %). ES/MS calcd. for $C_{40}H_{62}NO_5Si_2^+$ 692.4, found $m/z = 692.4$ ($M+H$)⁺.


15

Intermediate 48: (R)-Methyl 3-[2-[(2-[8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl]-2-[(tert-butyldimethylsilyl)oxy]ethyl]amino]-2-methylpropyl]benzoate




The title compound was synthesized in a manner analogous to that described
20 for Intermediate 47, using (R)-8-(benzyloxy)-5-[2-bromo-1-[(tert-butyldimethylsilyl)oxy]ethyl]quinolin-2(1H)-one in place of Intermediate 6. ES/MS calcd. for $C_{36}H_{47}N_2O_5Si^+$ 615.3, found $m/z = 615.3$ ($M+H$)⁺.

25 Intermediate 49: (R)-3-[2-[(2-[4-(BenzylOxy)-3-[(tert-butyldimethylsilyl)oxy]methyl]phenyl]-2-[(tert-butyldimethylsilyl)oxy]ethyl]amino]-2-methylpropyl]benzoic acid




LiOH (242 mg, 10.1 mmol) was added to a stirring solution of Intermediate 47 in THF/MeOH/H₂O (3:1:1) (10 mL) at rt. The resulting mixture was stirred for 3 days then quenched with 1 N HCl (pH < 3). The aqueous layer was extracted 5 with EtOAc (3 x 30 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ (s), and concentrated to give the title compound (730 mg). ES/MS calcd. for C₃₉H₆₀NO₅Si₂⁺ 678.4, found m/z = 678.5 (M+H)⁺.

10 Intermediate 50: (R)-3-[2-[(2-[8-(Benzyl)oxy]-2-oxo-1,2-dihydroquinolin-5-yl)oxy]ethyl]amino]-2-methylpropyl]benzoic acid



The title compound was synthesized in a manner analogous to that described for Intermediate 49, using Intermediate 48 as a substrate. ES/MS calcd. for C₃₅H₄₅N₂O₅Si⁺ 601.3, found m/z = 601.3 (M+H)⁺.
15


Intermediate 51: (R)-3-[2-[(2-[4-(Benzyl)oxy]-3-[(tert-butyldimethylsilyl)oxy]methyl]phenyl]-2-[(tert-butyldimethylsilyl)oxy]ethyl]amino]-2-methylpropyl]-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide



20 The title compound was synthesized in a manner analogous to that described for Intermediate 29, using Intermediate 49 in place of 3-(2-oxopropyl)benzoic acid. ES/MS calcd. for C₅₅H₈₃N₆O₅Si₂⁺ 963.6, found m/z = 963.6 (M+H)⁺.


25 Intermediate 52: (R)-3-[2-[(2-[8-(Benzyl)oxy]-2-oxo-1,2-dihydroquinolin-5-yl)oxy]ethyl]amino]-2-methylpropyl]-N-[[1,6-diethyl-4-

[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide



The title compound was synthesized in a manner analogous to that described
5 for Intermediate 29, using Intermediate 50 in place of 3-(2-oxopropyl)benzoic acid. ES/MS calcd. for $C_{51}H_{68}N_7O_5Si^+$ 886.5, found $m/z = 886.5$ ($M+H$)⁺.


10 Intermediate 53: (R)-3-[2-[[2-[4-(Benzylxy)-3-(hydroxymethyl)phenyl]-2-hydroxyethyl]amino]-2-methylpropyl]-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide



The title compound was synthesized in a manner analogous to that described
for Intermediate 3, using Intermediate 51 as a substrate. ES/MS calcd. for
C₄₃H₅₅N₆O₅⁺ 735.2, found $m/z = 735.5$ ($M+H$)⁺.


15

Intermediate 54: (R)-3-[2-[[2-[8-(Benzylxy)-2-oxo-1,2-dihydroquinolin-5-yl]-2-hydroxyethyl]amino]-2-methylpropyl]-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]benzamide




20 The title compound was synthesized in a manner analogous to that described
for Intermediate 3, using Intermediate 52 as a substrate. ES/MS calcd. for
C₄₅H₅₄N₇O₅⁺ 772.4, found $m/z = 772.4$ ($M+H$)⁺.

Intermediate 55: (R)-8-[(2-[(tert-Butyldimethylsilyl)oxy]-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]-N-[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]octanamide




- 5 The title compound was prepared in a manner analogous to that described for
10 Intermediate 37, using Intermediate 33 in place of Intermediate 26. ES/MS
calcd. for $C_{41}H_{64}N_7O_5Si$ 762.5, found $m/z = 762.5(M+H)^+$.

Intermediate 56: (R)-1-(4-(5-((2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pent-1-yn-1-yl)phenyl)-3-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)urea



- The title compound was prepared in a manner analogous to that described for
15 Intermediate 37, using Intermediate 35 in place of Intermediate 26. ES/MS
calcd. for $C_{45}H_{61}N_8O_5Si^+$ 821.5, found $m/z = 821.6 (M+H)^+$.


Intermediate 57: (R)-tert-butyl (2-(2,8-bis((tert-butoxycarbonyl)oxy)quinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(6-(4-(4-(3-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)ureido)phenyl)butoxy)hexyl)carbamate



The title compound was prepared in a manner analogous to that described for Intermediate 35, using Intermediate 42 in place of Intermediate 24. ES/MS calcd. for $C_{65}H_{99}N_8O_{12}Si^+$ 1211.7, found $m/z = 1211.6 (M+H)^+$.

5

Intermediate 58: (R)-1-(4-(5-((2-((tert-butyldimethylsilyl)oxy)-2-(5-hydroxy-3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-8-yl)ethyl)amino)pent-1-yn-1-yl)phenyl)-3-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)urea




10

The title compound was prepared in a manner analogous to that described for Intermediate 56, using (R)-8-(2-amino-1-((tert-butyldimethylsilyl)oxy)ethyl)-5-hydroxy-2H-benzo[b][1,4]oxazin-3(4H)-one in place of Intermediate 2. ES/MS calcd. for $C_{44}H_{61}N_8O_6Si^+$ 825.5, found $m/z = 825.6 (M+H)^+$.

15

Example 1: N-[(1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl]-3-[2-[(R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]propylbenzamide




The title compound was prepared in a manner analogous to that described for Intermediate 43, using Intermediate 4 in place of Intermediate 8.

¹H NMR (400 MHz, DMSO-d6) δ 10.51 (brs, 1H), 9.15 (brs, 1H), 8.86 (brs, 1H), 8.73 (brs, 1H), 8.34 (brs, 1H), 8.15 (dd, 1H, *J* = 4.5, 9.9 Hz), 7.77 (m, 2H), 7.47 5 (m, 2H), 7.19 (dd, 1H, *J* = 6.6, 8.0 Hz), 7.00 (d, 1H, *J* = 8.0 Hz), 6.60 (d, 1H, *J* = 10.0 Hz), 6.23 (brs, 1H), 5.35 (m, 1H), 4.57 (m, 2H), 4.44 (dd, 2H, *J* = 7.1, 14.3 Hz), 4.29 (m, 2H), 3.9-2.64 (m, 11H), 1.95 (m, 2H), 1.65 (m, 2H), 1.37 (t, 3H, *J* = 7.2 Hz), 1.30 (t, 3H, *J* = 6.8 Hz), 1.10 (dd, 3H, *J* = 6.5, 10.7 Hz); ES/MS calcd. for C₃₇H₄₆N₇O₅⁺ 668.4, found *m/z* = 668.3 (M+H)⁺.

10

Example 2: N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[(R)-2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]amino]propyl]benzamide




15 The title compound was synthesized in a manner analogous to that described in Intermediate 3, using Intermediate 43 as substrate.

¹H NMR (400 MHz, DMSO-d6) δ 9.44 (brs, 1H), 9.17 (brs, 1H), 8.80 (brs, 1H), 8.64 (brs, 1H), 8.34 (s, 1H), 7.74 (m, 2H), 7.46 (m, 2H), 7.36 (s, 1H), 7.09 (dd, 1H, *J* = 1.5, 8.9 Hz), 6.77 (d, 1H, *J* = 8.2 Hz), 6.06 (m, 1H), 4.82 (m, 1H), 4.57 20 (d, 2H, *J* = 6.5 Hz), 4.50 (s, 1H), 4.45 (dd, 2H, *J* = 6.2, 13.5 Hz), 4.30 (m, 1H), 3.88 (m, 3H), 3.61 (t, 2H, *J* = 11.5 Hz), 3.48 (m, 2H), 3.30 (m, 1H), 3.09 (m, 4H), 2.70 (m, 1H), 1.95 (m, 2H), 1.66 (m, 2H), 1.37 (t, 3H, *J* = 7.1 Hz), 1.30 (t, 3H, *J* = 7.5 Hz), 1.09 (m, 3H); ES/MS calcd. for C₃₅H₄₇N₆O₅⁺ 631.4, found *m/z* = 631.3 (M+H)⁺.

25


Example 3: (R)-N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[(2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl)amino]-2-methylpropyl]benzamide



The title compound was synthesized in a manner analogous to that described in Intermediate 2, using Intermediate 53 as substrate.

¹H NMR (400 MHz, DMSO-d₆) δ 9.44 (brs, 1H), 9.10 (brs, 1H), 8.66 (brs, 1H), 8.53 (brs, 1H), 8.30 (brs, 1H), 7.80 (d, 1H, *J* = 7.7 Hz), 7.73 (s, 1H), 7.47 (m, 2H), 7.39 (d, 1H, *J* = 1.9 Hz), 7.13 (dd, 1H, *J* = 2.0, 8.1 Hz), 6.79 (d, 1H, *J* = 8.2 Hz), 6.08 (m, 1H), 4.79 (m, 1H), 4.58 (d, 1H, *J* = 6.0 Hz), 4.51 (s, 1H), 4.42 (m, 2H), 4.28 (m, 1H), 3.89 (m, 2H), 3.78-2.78 (m, 11H), 1.95 (m, 2H), 1.63 (m, 2H), 1.37 (t, 3H, *J* = 7.1 Hz), 1.30 (t, 3H, *J* = 7.4 Hz), 1.20 (s, 6H); ES/MS calcd. for C₃₆H₄₉N₆O₅⁺ 645.4, found *m/z* = 645.4 (M+H)⁺.

Example 4: (R)-N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[(2-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]-2-methylpropyl]benzamide




15

The title compound was synthesized in a manner analogous to that described in Intermediate 2, using Intermediate 54 as substrate.


¹H NMR (400 MHz, DMSO-d₆) δ 10.31 (brs, 1H), 8.80 (t, 1H, *J* = 6.1 Hz), 8.19 (d, 1H, *J* = 10.0 Hz), 8.01 (s, 1H), 7.67 (m, 2H), 7.30 (m, 2H), 7.07 (t, 2H, *J* = 7.6 Hz), 6.90 (d, 1H, *J* = 8.1 Hz), 6.50 (d, 1H, *J* = 9.9 Hz), 5.29 (m, 1H), 4.94 (m, 1H), 4.55 (d, 2H, *J* = 6.2 Hz), 4.32 (q, 2H, *J* = 7.2 Hz), 4.11 (m, 1H), 3.85 (m, 2H, *J* = 7.9 Hz), 3.54 (t, 2H, *J* = 11.3 Hz), 3.32 (m, 2H), 3.16 (m, 1H), 2.95 (m, 2H), 2.86-2.54 (m, 3H), 1.91 (d, 2H, *J* = 15.5 Hz), 1.54 (m, 3H), 1.34 (t, 3H, *J* = 7.0 Hz), 1.24 (t, 3H, *J* = 7.3 Hz), 0.92 (m, 6H, *J* = 5.7 Hz); ES/MS calcd. for C₃₈H₄₈N₇O₅⁺ 682.3, found *m/z* = 682.4 (M+H)⁺.

Example 5: (R)-N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]methyl]benzamide



- 5 The title compound was synthesized in a manner analogous to that described for Intermediate 43, using Intermediate 4 in place of Intermediate 8 and Intermediate 30 in place of Intermediate 29.
- 10 ^1H NMR (400 MHz, DMSO-d6) δ 8.88 (t, 1H, J = 5.9 Hz), 8.10 (d, 1H, J = 10.0 Hz), 8.01 (s, 1H), 7.82 (s, 1H), 7.71 (d, 1H, J = 9.1 Hz), 7.47 (d, 1H, J = 7.6 Hz), 7.39 (t, 1H, J = 7.6 Hz), 7.10 (d, 1H, J = 7.7 Hz), 7.04 (d, 1H, J = 8.2 Hz), 6.88 (d, 1H, J = 8.1 Hz), 6.44 (d, 1H, J = 9.9 Hz), 5.32 (brs, 1H), 5.05 (dd, 1H, J = 4.5, 8.2 Hz), 4.55 (d, 2H, J = 6.1 Hz), 4.32 (q, 2H, J = 7.2 Hz), 4.10 (m, 1H), 3.86 (s, 2H), 3.78 (s, 2H), 3.55 (m, 2H), 3.32 (m, 3H), 2.98 (q, 2H, J = 7.4 Hz), 2.65 (m, 2H), 1.91 (m, 2H), 1.54 (m, 2H), 1.34 (t, 3H, J = 7.2 Hz), 1.26 (t, 3H, J = 7.5 Hz); ES/MS calcd. for $\text{C}_{35}\text{H}_{42}\text{N}_7\text{O}_5^+$ 640.3, found m/z = 640.3 ($\text{M}+\text{H}$)⁺.

Example 6: (R)-N-[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]methyl]benzamide




- 20 The title compound was synthesized in a manner analogous to that described for Intermediate 43, using Intermediate 4 in place of Intermediate 8 and Intermediate 31 in place of Intermediate 29.

25 ^1H NMR (400 MHz, DMSO-d6) δ 8.85 (t, 1H, J = 5.9 Hz), 8.11 (d, 1H, J = 10.0 Hz), 8.01 (s, 1H), 7.80 (d, 2H, J = 8.3 Hz), 7.40 (d, 2H, J = 8.4 Hz), 7.12 (d, 1H, J = 7.9 Hz), 7.04 (d, 1H, J = 8.2 Hz), 6.88 (d, 1H, J = 8.1 Hz), 6.44 (d, 1H, J = 9.9 Hz), 5.33 (brs, 1H), 5.05 (dd, 1H, J = 4.1, 8.1 Hz), 4.54 (d, 2H, J = 5.9 Hz),

4.32 (q, 2H, $J = 7.2$ Hz), 4.11 (m, 1H), 3.87 (m, 2H), 3.78 (s, 2H), 3.55 (td, 2H, $J = 2.1, 11.4$ Hz), 3.33 (m, 3H), 2.97 (q, 2H, $J = 7.4$ Hz), 2.65 (m, 2H), 1.91 (m, 2H), 1.53 (m, 2H), 1.34 (t, 3H, $J = 7.2$ Hz), 1.26 (t, 3H, $J = 7.5$ Hz); ES/MS calcd. for $C_{35}H_{42}N_7O_5^+$ 640.3, found $m/z = 640.3$ ($M+H$)⁺.


5

Example 7: (R)-N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[[[2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]amino]methyl]benzamide



- 10 The title compound was synthesized in a manner analogous to that described for Intermediate 3, using HF-Pyridine in place of TBAF and Intermediate 45 as a substrate.
- 15 1H NMR (400 MHz, DMSO-d6) δ 9.24 (brs, 1H), 8.91 (t, 1H, $J = 6.0$), 8.01 (s, 1H), 7.85 (s, 1H), 7.74 (d, 1H, $J = 7.8$), 7.52 (d, 1H, $J = 7.6$), 7.42 (t, 1H, $J = 7.7$), 7.24 (d, 1H, $J = 2.0$), 7.10 (d, 1H, $J = 7.8$), 6.97 (dd, 1H, $J = 2.2, 8.2$), 6.68 (d, 1H, $J = 8.2$), 4.93 (m, 1H), 4.59 (m, 1H), 4.55 (d, 2H, $J = 5.9$), 4.45 (s, 2H), 4.33 (q, 2H, $J = 7.2$), 4.12 (m, 1H), 3.85 (s, 3H), 3.55 (td, 2H, $J = 2.1, 11.4$), 3.34 (brs, 3H), 2.98 (q, 2H, $J = 7.4$), 2.62 (m, 2H), 1.91 (m, 2H), 1.54 (m, 2H), 1.34 (t, 3H, $J = 7.2$), 1.26 (t, 3H, $J = 7.5$); ES/MS calcd. for $C_{33}H_{43}N_6O_5^+$ 603.3, found $m/z = 603.3$ ($M+H$)⁺.
- 20

Example 8: (R)-N-[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[[[2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]amino]methyl]benzamide




25

The title compound was synthesized in a manner analogous to that described for Intermediate 3, using HF·Pyridine in place of TBAF and Intermediate 46 as a substrate.

¹H NMR (400 MHz, DMSO-d₆) δ 9.19 (brs, 1H), 8.87 (t, 1H, J = 5.8), 8.01 (s, 1H), 7.82 (d, 2H, J = 8.4), 7.44 (d, 2H, J = 8.3), 7.24 (d, 1H, J = 2.0), 7.11 (d, 1H, J = 7.7), 6.96 (dd, 1H, J = 2.2, 8.2), 6.68 (d, 1H, J = 8.2), 4.91 (m, 1H), 4.54 (m, 3H, J = 6.0), 4.44 (s, 2H), 4.32 (q, 2H, J = 7.2), 4.11 (m, 1H), 3.84 (s, 3H), 3.56 (td, 2H, J = 1.9, 11.2), 3.34 (m, 3H), 2.97 (q, 2H, J = 7.5), 2.60 (s, 2H), 1.91 (s, 2H), 1.53 (m, 2H), 1.34 (t, 3H, J = 7.2), 1.26 (t, 3H, J = 7.5); ES/MS calcd. for C₃₃H₄₃N₆O₅⁺ 603.3, found m/z = 603.3 (M+H)⁺.

Example 9: (R)-N-[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[5-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]pentyl]benzamide


15



The title compound was synthesized in a manner analogous to that described for Intermediate 3, using Intermediate 44 as substrate.

¹H NMR (400 MHz, DMSO-d₆) δ 10.58-10.39 (m, 2H), 9.19-8.95 (m, 2H), 8.68-8.45 (m, 2H), 8.42-8.25 (s, 1H), 8.14 (d, J = 9.99 Hz, 1H), 7.81 (d, J = 8.28 Hz, 2H), 7.33 (d, J = 8.24 Hz, 2H), 7.14 (d, J = 8.21 Hz, 1H), 6.98 (d, J = 8.17 Hz, 1H), 6.65-6.51 (m, 1H), 6.29-6.04 (s, 1H), 5.41-5.21 (m, 1H), 4.56 (d, J = 6.14 Hz, 2H), 4.49-4.36 (m, 2H), 4.36-4.19 (m, 1H), 3.99-3.82 (m, 2H), 3.72-3.53 (m, 2H), 3.19-2.86 (m, 6H), 2.64 (m, 2H), 2.07-1.86 (m, 2H), 1.79-1.51 (m, 6H), 1.34 (m, J = 30.50, 7.39 Hz, 7H); ES/MS calcd. for C₃₉H₅₀N₇O₅⁺ 696.4, found m/z = 696.4 (M+H)⁺.

Example 10: (R)-N-[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-8-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]octanamide




The title compound was synthesized in a manner analogous to that described in Intermediate 3, using Intermediate 55 as substrate.

- ¹H NMR (400 MHz, DMSO-d₆) δ 10.65 (brs, 2H), 8.65 (brs, 1H), 8.59 (brs, 2H), 8.33 (brs, 1H), 8.15 (d, 1H, *J* = 10 Hz), 7.14 (d, 1H, *J* = 8 Hz), 6.98 (d, 1H, *J* = 8 Hz), 6.20 (brs, 1H), 5.30 (d, 1H, *J* = 8.8 Hz), 4.42 (m, 3H), 4.35 (d, 2H, *J* = 6 Hz), 4.25 (brs, 1H), 3.9 (m, 3H), 3.60 (t, 2H, *J* = 11.2 Hz), 3.18-2.9 (m, 6H), 2.15 (t, 2H, *J* = 7.2 Hz), 1.95 (m, 2H), 1.6-1.5 (m, 6H), 1.35 (t, 3H, *J* = 7.2 Hz), 1.22 (M, 8H); ES/MS calcd. for C₃₅H₅₀N₇O₅⁺ 648.4, found *m/z* = 649.0 (M+H)⁺.

10

Example 11: (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(5-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pent-1-yn-1-yl)phenyl)urea




- 15 A solution of Intermediate 56 (0.16 g, 0.19 mmol) in tetrahydrofuran (2 mL) was treated successively with TBAF solution (1.0 M in THF, 2 mL) and glacial acetic acid (0.60 mL). The mixture was stirred overnight at room temperature and then concentrated to dryness under reduced pressure. The residue was purified by reverse-phase high performance liquid chromatography (RP-HPLC, acetonitrile/water/0.1 % TFA) to provide, after concentration, the trifluoroacetic acid salt of the title compound as a white powder (80 mg).

- 20 ¹H NMR (400 MHz, DMSO-d₆) δ 11.44 (bs, 1H), 10.49 (s, 1H), 9.26 (s, 1H), 8.90 (bs, 1H), 8.67 (s, 1H), 8.35 (s, 1H), 8.17 (d, *J* = 10.0 Hz, 1H), 7.42 (d, *J* = 8.8 Hz, 2H), 7.30 (d, *J* = 8.4 Hz, 2H), 7.16 (d, *J* = 8.4 Hz, 1H), 6.99 (d, *J* = 8.0 Hz, 1H), 6.58 (d, *J* = 10.0 Hz, 1H), 6.65-6.51 (m, 1H), 6.24-6.09 (bs, 1H), 5.31 (m, 1H), 4.41 (m, 2H), 4.37-4.22 (m, 1H), 3.93-3.85 (m, 2H), 3.83-3.75 (m, 2H),

3.68-3.57 (m, 2H), 3.55-3.48 (m, 2H), 3.10-2.96 (m, 6H), 2.07-1.82 (m, 2H), 1.75-1.61 (m, 4H), 1.38 (t, $J = 7.2$ Hz, 3H), 1.30 (t, $J = 7.6$ Hz, 3H); ES/MS calcd. for $C_{39}H_{47}N_8O_5^+$ 707.4, found $m/z = 707.5$ ($M+H$)⁺.


- 5 Example 12: (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(5-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)phenyl)urea



A solution of the trifluoroacetic acid salt of Example 11 (60 mg) in methanol (4 mL) and ethyl acetate (1 mL) was treated with palladium on carbon (10 % w/w, 10 mg) and then stirred for one hour under a balloon of hydrogen gas. The mixture was filtered through a pad of Celite diatomaceous earth, and then the filtrate was concentrated under reduced pressure to provide the trifluoroacetic acid salt of the title compound as a pale golden glass, which was scraped into a powder (27 mg).

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.55-10.48 (m, 2H), 9.83 (s, 1H), 8.82-8.74 (m, 2H), 8.54 (s, 1H), 8.29 (bs, 1H), 8.15 (d, $J = 10.0$ Hz, 1H), 7.32 (d, $J = 8.4$ Hz, 2H), 7.14 (d, $J = 8.0$ Hz, 1H), 7.08 (d, $J = 8.4$ Hz, 2H), 6.98 (d, $J = 8.0$ Hz, 1H), 6.65-6.51 (d, $J = 9.6$ Hz, 1H), 6.19-6.10 (bs, 1H), 5.29 (m, 1H), 4.56 (d, $J = 6.14$ Hz, 2H), 4.48-4.33 (m, 2H), 4.32-4.20 (m, 1H), 3.93-3.83 (m, 2H), 3.65-3.55 (m, 2H), 3.47-3.42 (m, 2H), 3.18-2.86 (m, 6H), 2.64 (m, 2H), 2.05-1.92 (m, 2H), 1.74-1.51 (m, 6H), 1.34 (m, 7H); ES/MS calcd. for $C_{39}H_{51}N_8O_5^+$ 711.4, found $m/z = 711.4$ ($M+H$)⁺.


- 25 Example 13: (R)-3-(((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)amino)-4-((4-(5-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)phenyl)amino)cyclobut-3-ene-1,2-dione



A mixture of Intermediate 38 (0.11 g, 0.19 mmol) and Intermediate 28 (0.15 g, 0.36 mmol) in ethanol (2 mL) was heated for 3 days in a 70 °C oil bath. The mixture was concentrated under reduced pressure and was then purified by
5 automated flash silica gel chromatography, using a 12g Silicycle SiliSep flash column (dichloromethane/methanol) to give the impure intermediate, (R)-tert-butyl 2-(tert-butyldimethylsilyloxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl(5-(4-(2-((1,6-diethyl-4-(tetrahydro-2H-pyran-4-ylamino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methylamino)-3,4-dioxocyclobut-1-enylamino)phenyl)pent-4-ynyl)carbamate. This material was then subjected to reverse-phase high performance liquid chromatography (RP-HPLC, acetonitrile/water/0.1 % TFA) to provide, after concentration, the intermediate
10 (R)-tert-butyl 5-(4-(2-((1,6-diethyl-4-(tetrahydro-2H-pyran-4-ylamino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methylamino)-3,4-dioxocyclobut-1-enylamino)phenyl)pent-4-ynyl(2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)carbamate. This intermediate was dissolved in methanol (5 mL) and treated with palladium on carbon (10 % w/w, wet
15 Degussa-type, catalytic amount). The suspension was stirred under a balloon of hydrogen gas until LC/MS analysis indicated the complete reduction of the triple bond. The mixture was then filtered through a pad of Celite diatomaceous earth, and the filtrate was concentrated to dryness under reduced pressure. The residue was taken up in dichloromethane/methanol (1:1, 5 mL) and treated with a solution of hydrogen chloride in dioxane (4 N, 2 mL). Once LC/MS analysis showed the complete removal of the Boc-protecting
20 group, the mixture was concentrated to dryness. The residue was purified by reverse-phase high performance liquid chromatography (RP-HPLC, acetonitrile/water/0.1 % TFA) to provide, after concentration, the trifluoroacetic acid salt of the title compound as a pale yellow powder (28 mg).

¹H NMR (400 MHz, DMSO-d₆) δ 10.50-10.46 (m, 2H), 9.81 (s, 1H), 8.53 (m, 2H), 8.28 (bs, 1H), 8.15 (d, J = 9.6 Hz, 1H), 7.85 (m, 1H), 7.36 (d, J = 8.4 Hz, 2H), 7.13 – 7.17 (m, 3H), 6.98 (d, J = 8.0 Hz, 1H), 6.58 (d, J = 10.0 Hz, 1H), 6.24-6.04 (bs, 1H), 5.29 (m, 1H), 4.94 (d, J = 5.2 Hz, 2H), 4.42 (m, 2H), 4.30-5 4.18 (m, 1H), 3.92-3.85 (m, 2H), 3.59 (m, 2H), 3.18-2.86 (m, 6H), 2.54 (m, 2H), 2.06-1.88 (m, 2H), 1.79-1.51 (m, 6H), 1.33 (m, 7H); ES/MS calcd. for C₄₂H₅₁N₈O₆⁺ 763.4, found m/z = 763.3 (M+H)⁺.


Example 14: (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(4-((6-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl)oxy)butyl)phenyl)urea



The crude concentrated reaction mixture containing Intermediate 57 was taken up in methanol (approximately 0.50 mL) and treated with a solution of hydrogen chloride in dioxanes (4 N, 2 mL). The mixture was subsequently concentrated to dryness under reduced pressure. The residue was purified by reverse-phase high performance liquid chromatography (RP-HPLC, acetonitrile/water/0.1 % TFA) to provide, after concentration, the trifluoroacetic acid salt of the title compound as a powder (107 mg, 58 % over two steps).

20 ¹H NMR (400 MHz, DMSO-d₆) δ 10.48 (m, 2H), 8.82 (s, 1H), 8.54 (m, 2H), 8.32 (s, 1H), 8.15 (d, J = 10.0 Hz, 1H), 7.31 (d, J = 8.4 Hz, 2H), 7.14 (d, J = 8.4 Hz, 1H), 7.07 (d, J = 8.4 Hz, 2H), 6.98 (d, J = 8.4 Hz, 1H), 6.58 (d, J = 9.9 Hz, 1H), 6.15 (s, 1H), 5.30 (d, J = 8.1 Hz, 1H), 4.44-4.22 (m, 6H), 3.91 (m, 2H), 3.60 (m, 3H), 3.33 (m, 4H), 3.14 – 2.88 (m, 8H), 2.02 – 1.91 (m, 2H), 1.78-1.42 (m, 8H), 25 1.40-1.34 (m, 4H), 1.32-1.24 (m, 8H); ES/MS calcd. for C₄₄H₆₁N₈O₆⁺ 797.5, found m/z = 797.4 (M+H)⁺.

Example 15: (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(5-((2-hydroxy-2-(5-hydroxy-3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-8-yl)ethyl)amino)pent-1-yn-1-yl)phenyl)urea




- 5 The title compound was prepared in a manner analogous to that described for Example 11, using Intermediate 58 as a substrate.

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.05-9.96 (m, 2H), 9.19-8.70 (bs, 2H), 8.63-8.56 (m, 2H), 8.48-8.25 (bs, 1H), 7.41 (d, *J* = 8.4 Hz, 2H), 7.30 (d, *J* = 8.9 Hz, 2H), 6.92 (d, *J* = 8.0 Hz, 1H), 6.56 (d, *J* = 8.8 Hz, 1H), 5.94 (s, 1H), 5.07 (d, *J* = 9.3 Hz, 2H), 5.94 (s, 1H), 5.06 (d, *J* = 9.2 Hz, 1H), 4.58-4.49 (m, 4H), 4.49-4.32 (m, 2H), 4.32-4.19 (m, 1H), 3.91-3.88 (m, 2H), 3.63-3.53 (m, 2H), 3.14-2.86 (m, 6H), 2.68 (m, 2H), 2.07-1.86 (m, 2H), 1.72-1.56 (m, 2H), 1.37 (t, *J* = 7.2 Hz, 3H), 1.29 (t, *J* = 7.6 Hz, 3H); ES/MS calcd. for C₃₈H₄₇N₈O₆⁺ 711.4, found *m/z* = 711.3 (M+H)⁺.


15

Examples 16-20 may be prepared using the general methods described above.


Example 16: (R)-1-Ethyl-N-[4'-[[[2-(3-formamido-4-hydroxyphenyl)-2-hydroxyethyl]amino]methyl]biphenyl-4-yl]-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide




Example 17: (R)-8-[2-[2-[4-[1-Ethyl-4-(tetrahydro-2H-pyran-4-ylamino)-1H-pyrazolo[3,4-b]pyridine-5-carbonyl]piperazin-1-yl]ethylamino]-1-hydroxyethyl]-5-hydroxy-2H-benzo[b][1,4]oxazin-3(4H)-one




Example 18: (R)-N-[[1,6-diethyl-4-(tetrahydro-2H-pyran-4-ylamino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl]-6-[4-[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethylamino]butoxy]nicotinamide
5



Example 19: (R)-N-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-4-(5-((2-hydroxy-2-(4-hydroxy-3-(methylsulfonamido)phenyl)ethyl)amino)pentyl)benzamide
10



Example 20: (R)-6-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl ((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)carbamate
15



BIOLOGICAL EXAMPLES

Inhibition of PDE4B4 Enzyme Activity

The inhibitory activity of test compounds were determined using the PDE Glo Assay (Promega, V1361). The PDE-Glo Reaction Buffer was prepared (1mL
5 5X buffer + 4mL water) and 150 nM solution of test compound dissolved in the Reaction Buffer, followed by 1:3 serial dilutions into Reaction Buffer. DMSO was used for the initial dilution of test compounds and the final concentration used in the assay did not exceed 0.1% DMSO. Human recombinant PDE4B2 (BPS Bioscience) was added to test compound in Reaction Buffer at a
10 concentration of 0.12 nM per reaction. The enzyme and test compound were pre-incubated together at rt for 10 min. 50 nM of the enzyme substrate cAMP was then added to initiate the enzyme reaction, with the reaction terminated after one hour. Termination Buffer (1mL 5X termination buffer + 3.9 mL water + 100 µL of 100 mM IBMX (ICN Chemicals, in DMSO)), was added to each
15 reaction well to terminate the reaction. Detection Buffer (1mL 5X PDE-Glo Detection buffer + 3.96mL water + 40 µL PKA (supplied in kit)) was added to the cAMP-enzyme-test compound mix and this secondary reaction proceeded for 20 minutes at rt. An equal volume of Kinase Glo Reagent was then added to the reaction mixture and after 10 min luminescence was measured using a
20 luminometer (EnVision, 0.1 sec read for luminescence). Luminescence values were directly related to levels of cAMP in the reaction mixture. Data were plotted as relative light units (RLU) versus test compound concentration and the IC₅₀ determined using GraphPad Prism 5.0, using a nonlinear curve fit in a single-site binding model.
25 IC50 determinations for the compounds presented above are seen in the first column of Table 1 (PDE4 IC50).

Beta-2 Adrenoceptor Binding Assay

Test compounds were incubated for 120 min with Chinese hamster oocyte
30 (CHO) cells expressing the human recombinant beta-2 adrenoreceptors. Binding of test compounds to the beta-2 adrenoreceptors was determined by measuring the displacement of the radiolabeled ligand [³H](-)CGP-12177 (0.3 nM) from the receptor measured by a scintillation counting. Concentration

response curves were generated for test compounds and the IC₅₀ (molar concentration of the compound which produced 50% inhibition for the maximal response for that compound) was determined. These assays were performed using a protocol based on the original description of the assay by Joseph *et al.*,

5 (2004) *Naun.-Sch. Arch. Pharm.*; **369**:525-532.

IC50 determinations for the compounds presented above are seen in the second column of Table 1 (Beta 2 IC50).

Functional agonism of human recombinant beta-2 adrenoreceptors expressed

10 in Chinese hamster oocytes (CHO)

Test compounds were incubated for 30 min with CHO cells expressing the human recombinant beta-2 adrenoreceptors. Agonism of the receptor was measured by the elevation of intracellular cyclic-AMP over control levels detected using a homogeneous time resolved fluorescence (HTRF) format.

15 Compounds were determined to be "full" or "partial" agonists based on the maximum level of cAMP accumulated compared to the control compound isoproterenol which is a full agonist in this assay system. Concentration response curves were generated for test compounds and the EC₅₀ (molar concentration of the agonist which produced 50% of the maximal response for that compound) was determined. These assays were performed using a protocol based on the original description of the assay by Baker, J.G.

(2005) *Brit. J. Pharmacol.*; **144**:317-322.

EC50 determinations for the compounds presented above are seen in the third column of Table 1 (Beta 2 EC50).

25

Inhibition of lipoplysaccharide (LPS)-induced tumour necrosis factor alpha (TNF- α) release from human peripheral blood mononuclear cells (PBMC)

Human whole blood was drawn from donors and cell purification initiated within two hours. Peripheral blood mononuclear cells (PBMC) were purified using a standard Ficoll gradient purification technique and aliquoted at a concentration of 100,000 cells/ well in 96 well plates. Beta-2 adrenoreceptor agonists are also anti-inflammatory for some immune cells expressing the receptor and in this assay they can inhibit tumor necrosis factor-alpha (TNF- α) production.

Therefore PBMC were pre-incubated for 30 min in the presence of the beta-2 adrenoreceptor antagonist ICI-118551 (3-(isopropylamino)-1-[(7-methyl-4-indanyl)oxy]butan-2-ol) (10 µM) to determine the PDE4 directed activity of test compounds, and in the absence of ICI-118551 for the combined PDE4 and

5 beta-2 receptor directed activity. Test compounds were dissolved in DMSO and diluted in buffer (final DMSO concentration of 0.1% v/v) and were then added to the cell suspensions and pre-incubated for a further 1 h at 37°C. LPS solution was then added (0.4 ng/mL) and the cells incubated for a further 6 h. At the end of the incubation period the cell supernatants were collected and TNF- α

10 was measured using a Procarta Cytokine Assay Kit (Affymetrix, Santa Clara, CA) according to the manufacturers' instructions. Data were analyzed using MiraiBio Masterplex QT v4.0 software (MasterPlex Version 1.0.1.18 Copyright 2008, Hitachi Software Engineering Co., Ltd) and used to extrapolate Mean Fluorescence Intensity (MFI) into concentrations of TNF- α and mean values

15 from replicate wells were determined. Inhibition curves were plotted using a nonlinear curve fit employing a single-site binding model to determine the IC₅₀ value for each test compound in the presence or absence of ICI-118551 using GraphPad Prism 5 for Windows, version 5.02 (GraphPad Software, San Diego CA).

20 EC50 determinations for the compounds presented above are seen in the fourth column of Table 1 (PBMC TNF EC50).

Inhibition of lipopolysaccharide (LPS)-induced neutrophil recruitment into the lungs of Lewis rats

25 Rats are anesthetized with 4% isoflurane, placed in the supine position at an angle of 30°, the mouth opened and the trachea exposed. A 22-gauge needle with a syringe attached is introduced into the trachea and test compound in suspension (200 µl volume per 400 g) is delivered into the lungs from approximately one centimeter above the carina. The rats are allowed to recover

30 and two hrs later conscious animals are placed into a chamber and exposed to aerosolized LPS (1.0 mg/mL) at a rate of 3.0 L/min for 20 min. Rats are euthanized 4 h post-LPS exposure by an overdose of pentobarbital (90 mg/kg) by intra-peritoneal injection. Bronchoalveolar lavage (BAL) was then

performed with a 14 gauge blunt needle into the exposed trachea. Five, 5 ml washes of PBS are collect from the lungs and placed into Falcon tubes then centrifuged at 1600g for 10 min at 4°C. The supernatant is discarded, the cells are re-suspended in PBS and total cell counts are determined based on a 10 µl sample of re-suspended cells stained with trypan blue and counts performed using a Countess® cell counter (Invitrogen). Differential cell counts to determine the number of neutrophils in the BAL wash are performed on cytospun cells stained with May-Grunwald and Giemsa solution. Manual eye counting is performed to determine the percentage number of cells in the cytospun sample (determined as macrophages, neutrophils, eosinophils, T-lymphocytes and eosinophils) and these values are used to determine the total number of each cell type per sample. Typically experiments contain a minimum of six rats per experimental group and the mean ± SEM number of neutrophils is determined for each group. The level of neutrophil inhibition caused by test compounds dosed directly into the lungs is determined compared to the vehicle-treated and LPS-exposed control rats. Statistical analysis to determine significant differences between groups are performed by one-way analysis of variance (ANOVA) using GraphPad Prism 5 for Windows, version 5.02 (GraphPad Software, San Diego CA).

The compounds of Examples 1 and 9 were tested in this assay and both exhibited a neutrophil inhibition of greater than 40% at a dose of 300 µg/kg.

Inhibition of acetylcholine-induced bronchoconstriction in Dunkin-Hartley guinea pigs

Guinea pigs (Dunkin-Hartley from Charles River Laboratories , male, 500 to 800g) are anesthetized with 4% isoflurane, placed in the supine position at an angle of 30°, the mouth opened and the trachea exposed. A 22-gauge needle with a syringe attached is introduced into the trachea and test compound in suspension (200 µl volume) is delivered into the lungs from approximately one centimeter above the carina. *In vivo* bronchoprotective effects of the test compounds against acetylcholine (ACh)-induced bronchoconstriction are tested in conscious guinea pigs using a whole body plethysmograph system (WBP) (Buxco Research Systems) 4 hrs after intra-tracheal dosing of test compounds.

The lung function is measured in this system and expressed as an enhanced PAUSE (Penh) which has been widely used in scientific research and preclinical drug screening as a surrogate methodology for measuring lung resistance in conscious animals (Chong *et al.*, *J. Pharmacol. Toxicol. Methods* 5 1998;39:163-168. Pennock *et al.*, *J. Appl. Physiol.* 1979;46:399-406). Guinea pigs are placed in the WBP system chambers and exposed to aerosol of either 0.9% saline solution or ACh solution (4mg/mL) for 1 min. Lung function measurements (expressed as Penh and calculated by peak expiratory flow/peak inspiratory flow x pause) are continuously recorded for 20 min 10 immediately after saline or ACh challenge. The results are expressed as area under curves (AUC) of airway response (Penh) over the responding time (20 min). Twenty four hrs before the assessment of test compounds takes place, airway responses of guinea pigs are measured to determine their baseline responses prior to compound treatments. Each animal may therefore act as it's own control for the evaluation of bronchoprotection of test compounds and the efficacy of test compounds are calculated as the percentage inhibition of airway 15 response compared to this value. Assessment of the duration of bronchoprotection could also be determined by re-challenging animals up to 24 hrs post test compound dosing. Typically experiments will contain a minimum of six guinea pigs per experimental group and the mean ± SEM inhibition of PenH is determined for each group. Statistical analysis to determine significant differences between groups are performed by one-way analysis of variance (ANOVA) using GraphPad Prism 5 for Windows, version 5.02 (GraphPad Software, San Diego CA).

25

Table 1

Example #	PDE4 IC50	Beta 2 IC50	Beta 2 EC50	PBMC TNF EC50
1	++	++	++	++
2	++	-	++	++
3	++	+	+	++
4	++	++	++	++
5	++	+	++	++
6	++	-	++	++
7	++	-	-	-
8	++	-	-	+
9	++	++	++	++

10	++	-	++	++
11	++	++	++	++
12	++	++	++	++
13	**	++	++	++
14	**	++	++	++
15	**	++	++	++

< 100 nM = ++
< 300 nM = +
> 300 nM = -
Not measured = **

Dry Powder Formulation

5

A dry powder formulation of one or more compounds of the invention for administration by inhalation may be prepared by as follows:


Particles of a compound of the invention (API) are micronized using conventional processes including but not limited to jet milling, to achieve a distribution with a mass median aerodynamic diameter (MMAD) of about 2 and a GSD < about 2.5. The micronized particles are then blended with a conventional dry powder excipient such as lactose. Specific examples of suitable forms of commercially available lactose include Lactohale LH100 which comprises particles >60 micron and Lactohale LH200 which comprises large (>60 microns) lactose particles mixed with lactose "fines" (<10 microns). A typical formulation will include less than 10% API, with the remainder being the dry powder excipient. This bulk formulation can be filled into a multi-dose DPI, e.g. Valois Prohaler, with a fill weight designed to permit emission of the desired dose.

10
15
20

Claims

That Which Is Claimed Is:


1. A compound of Formula I:



5

or a pharmaceutically acceptable salt thereof,
wherein:

X is selected from:



- 10 R¹ is CH₂OH, CH₂CH₂OH, N(H)C(O)H, N(H)S(O₂)CH₃, and R² is H;
or R¹ and R² together with the phenyl to which they are bound form a bicyclic,
fused heterocyclic ring having 9 or 10 ring atoms wherein 1 or 2 ring
atoms are selected from N, O and S and the remaining ring atoms are C,
wherein said bicyclic fused heterocyclic ring is optionally substituted with
15 one, two or three additional substituents independently selected from
alkyl, oxo and OH;

- R³ is selected from C₄₋₁₂alkylene, C₄₋₁₂alkenylene, C₄₋₁₂alkynylene, R⁴-O-R⁴,
R⁴-N(R⁸)-R⁴, C₃₋₆cycloalkylene, R⁴-C₃₋₆cycloalkylene,
C₃₋₆cycloalkylene-R⁴, R⁴-C₃₋₆cycloalkylene-R⁴, C₆₋₁₀arylene,
20 R⁴-C₆₋₁₀arylene, C₆₋₁₀arylene-R⁴, R⁴-C₆₋₁₀arylene-R⁴,
R⁴-C₆₋₁₀arylene-O-R⁴, R⁴-C₆₋₁₀arylene-N(R⁸)-R⁴,
R⁴-C₆₋₁₀arylene-C₆₋₁₀arylene, Het, R⁴-Het, R⁴-Het-C₆₋₁₀arylene, , R⁴- C₆₋
10arylene -C₆₋₁₀arylene, Het-R⁴, R⁴-Het-R⁴, R⁴-O-Het
25 R⁴-C₆₋₁₀arylene-O-Het, R⁴-C₆₋₁₀arylene-C(O)-Het, and
R⁴-C₆₋₁₀arylene-N(R⁸)-Het;

wherein said alkylene, alkenylene, alkynylene, cycloalkylene, or arylene is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸;

Het is 5-6 membered saturated or unsaturated heterocyclene wherein 1 or 2 ring atoms are selected from N, O and S, and wherein said heterocyclene is optionally substituted with 1, 2 or 3 substituents selected from halo, alkyl, alkoxy, oxo and OH;


R⁴ is C₁₋₁₀alkylene, C₂₋₁₀alkenylene, or C₂₋₁₀alkynylene wherein each R⁴ is optionally substituted with 1, 2 or 3 substituents selected from halo, oxo, and OR⁸; with the proviso that the total number of carbon atoms in the C₁₋₁₀alkylene, C₂₋₁₀alkenylene, or C₂₋₁₀alkynylene chains of two R⁴ groups in any definition of R³ is not greater than 12;

Y is C(O), C(O)N(R⁸)CH₂, N(R⁸)C(O), O-C(O)N(R⁸)CH₂, N(R⁸)C(O)N(R⁸)CH₂, or SO₂N(R⁸)CH₂;

R⁵ is alkyl;

R⁶ is H or alkyl;

R⁷ is selected from unsubstituted C₃₋₆cycloalkyl, substituted C₃₋₆cycloalkyl, and a heterocyclic group selected from formulas (i), (ii), and (iii):



20

wherein Z is O, S, S(O)₂, NH or N-R^{7a}, and

R^{7a} is selected from alkyl, C(O)alkyl, C(O)NH₂, C(O)N(H)alkyl, and

C(O)N(alkyl)₂; and

R⁸ is H or alkyl.

25

2. The compound according to claim 1 wherein R³ is selected from C₄₋₁₂alkylene, C₄₋₁₂alkenylene, C₄₋₁₂alkynylene, R⁴-O-R⁴, R⁴-N(R⁸)-R⁴,

C₃₋₆cycloalkylene, R⁴-C₃₋₆cycloalkylene, C₃₋₆cycloalkylene-R⁴,

R⁴-C₃₋₆cycloalkylene-R⁴, phenylene, R⁴-phenylene, phenylene-R⁴,

30 R⁴-phenylene-R⁴, R⁴-phenylene-O-R⁴, R⁴-phenylene-N(R⁸)-R⁴,

R⁴-phenylene-phenylene, Het, R⁴-Het, R⁴-Het-phenylene, Het-R⁴, R⁴-Het-R⁴,

R^4 -O-Het R^4 -phenylene-O-Het, R^4 -phenylene-C(O)-Het, and
 R^4 -phenylene-N(R^8)-Het;


wherein said alkylene, alkenylene, alkynylene, cycloalkylene, or arylene
is optionally substituted with 1, 2 or 3 substituents selected from
5 halo, oxo, and OR⁸;
or a pharmaceutically acceptable salt thereof.

3. The compound according to claim 1 or a pharmaceutically acceptable
salt thereof wherein R¹ is CH₂OH and R² is H.

10

4. The compound according to claim 1 or a pharmaceutically acceptable
salt thereof wherein R¹ and R² together with the phenyl to which they are
bound form a bicyclic, fused heterocyclic ring having 9 or 10 ring atoms
wherein 1 or 2 ring atoms are selected from N, O and S, and said bicyclic fused
15 heterocyclic ring is optionally substituted with one additional substituent
selected from alkyl, oxo and OH.


5. The compound according to claim 1 or a pharmaceutically acceptable
salt thereof wherein R¹ and R² together with the phenyl to which they are
20 bound form



6. The compound according to any of claims 1-5 or a pharmaceutically
acceptable salt thereof wherein R³ is selected from C₄₋₁₂alkylene, C₄₋₁₂
25 alkenylene, C₄₋₁₂alkynylene, R⁴-O-R⁴, and R⁴-N(R^8)-R⁴, wherein said alkylene,
alkenylene or alkynylene are each optionally substituted with 1, 2 or 3
substituents selected from halo, oxo, and OR⁸.

7. The compound according to any of claims 1-6 or a pharmaceutically acceptable salt thereof wherein R³ is selected from C₅₋₈alkylene, C₅₋₈alkenylene, C₅₋₈alkynylene, R⁴-O-R⁴, and R⁴-N(R⁸)-R⁴, wherein each R⁴ is C₁₋₄alkylene, C₂₋₄alkenylene, or C₂₋₄alkynylene each alkylene, alkenylene and alkynylene
5 optionally substituted with 1 or 2 substituents selected from halo, oxo, and OR⁸.
8. The compound according to any of claims 1-7 or a pharmaceutically acceptable salt thereof wherein R³ is unsubstituted C₅₋₈alkylene.
- 10 9. The compound according to any of claims 1-5 or a pharmaceutically acceptable salt thereof wherein R³ is selected from C₆₋₁₀arylene, R⁴-C₆₋₁₀arylene, R⁴-C₆₋₁₀arylene-R⁴, Het, R⁴-Het, and R⁴-Het-R⁴.
- 15 10. The compound according to any of claims 1-5 or 9 or a pharmaceutically acceptable salt thereof wherein R³ is selected from R⁴-C₆₋₁₀arylene, Het, R⁴-Het, and R⁴-Het-R⁴.
- 20 11. The compound according to any of claims 1-5 or 9-10 or a pharmaceutically acceptable salt thereof wherein R³ is selected from R⁴-phenylene, R⁴-phenylene-R⁴, Het, R⁴-Het, and R⁴-Het-R⁴.
- 25 12. The compound according to any of claims 1-5 or 9-11 or a pharmaceutically acceptable salt thereof wherein R³ is selected from R⁴-phenylene, Het, and R⁴-Het.
13. The compound according to any of claims 1-12 or a pharmaceutically acceptable salt thereof wherein Y is C(O)N(R⁸)CH₂.
- 30 14. The compound according to any of claims 1-13 or a pharmaceutically acceptable salt thereof wherein R⁵ is -CH₂CH₃.
15. The compound according to any of claims 1-14 or a pharmaceutically acceptable salt thereof wherein R⁶ is -CH₂CH₃.

16. The compound according to any of claims 1-15 or a pharmaceutically

acceptable salt thereof wherein R⁷ is .

5 17. A compound selected from

N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[(R)-2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]amino]propyl]benzamide;

10 (R)-N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]amino]-2-methylpropyl]benzamide;

15 (R)-N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[2-[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]-2-methylpropyl]benzamide;

20 (R)-N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]methyl]benzamide;

(R)-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]methyl]benzamide;

25

(R)-N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-3-[[2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]amino]methyl]benzamide;

(R)-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[[[2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]amino]methyl]benzamide;

5 (R)-N-[[1,6-diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-4-[5-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]pentyl]benzamide;

10 (R)-N-[[1,6-Diethyl-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-8-[[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl]amino]octanamide;

15 (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(5-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pent-1-yn-1-yl)phenyl)urea;

20 (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(5-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)phenyl)urea;

(R)-3-(((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)amino)-4-((4-(5-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)phenyl)amino)cyclobut-3-ene-1,2-dione;

25 (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(4-((6-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl)oxy)butyl)phenyl)urea;

30 (R)-1-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-3-(4-(5-((2-hydroxy-2-(5-hydroxy-3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-8-yl)ethyl)amino)pent-1-yn-1-yl)phenyl)urea;

(R)-1-Ethyl-N-[4'-[[[2-(3-formamido-4-hydroxyphenyl)-2-hydroxyethyl]amino]methyl]biphenyl-4-yl]-4-[(tetrahydro-2H-pyran-4-yl)amino]-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

5 (R)-8-[2-[2-[4-[1-Ethyl-4-(tetrahydro-2H-pyran-4-ylamino)-1H-pyrazolo[3,4-b]pyridine-5-carbonyl]piperazin-1-yl]ethylamino]-1-hydroxyethyl]-5-hydroxy-2H-benzo[b][1,4]oxazin-3(4H)-one;

10 (R)-N-[[1,6-diethyl-4-(tetrahydro-2H-pyran-4-ylamino)-1H-pyrazolo[3,4-b]pyridin-5-yl]methyl]-6-[4-[2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethylamino]butoxy]nicotinamide;

15 (R)-N-((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)-4-(5-((2-hydroxy-2-(4-hydroxy-3-(methylsulfonamido)phenyl)ethyl)amino)pentyl)benzamide;

(R)-6-((2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl ((1,6-diethyl-4-((tetrahydro-2H-pyran-4-yl)amino)-1H-pyrazolo[3,4-b]pyridin-5-yl)methyl)carbamate;

20 or a pharmaceutically acceptable salt thereof.

18. A composition comprising a compound of Claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent or excipient.

19. The composition according to Claim 18, wherein said composition is suitable for inhalation.

30 20. The composition according to Claim 18 further comprising a therapeutically active agent selected from anti-inflammatory agents, anticholinergic agents, peroxisome proliferator-activated receptor agonists,

epithelial sodium channel blockers, kinase inhibitors, protease inhibitors, antiinfective agents and antihistamines.

21. A method comprising administering to a human, an effective amount of a compound according to any of Claims 1-17 or a pharmaceutically acceptable salt thereof.
22. A method for treating pulmonary inflammation or bronchoconstriction in a human in need thereof, comprising administering to said human an effective amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof.
23. A method for treating a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associated bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human in need thereof, said method comprising administering to said human an effective amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof.
24. A method for treating chronic obstructive pulmonary disease (COPD) in a human in need thereof, said method comprising administering to said human an effective amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof.
25. A method for treating asthma in a human in need thereof, said method comprising administering to said human an effective amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof.
26. A compound according to any of claims 1-17 or a pharmaceutically acceptable salt thereof for use as a medicament.

27. A compound according to any of claims 1-17 or a pharmaceutically acceptable salt thereof for use in a method of treating of pulmonary inflammation or bronchoconstriction in a human.

5

28. A compound according to any of claims 1-17 or a pharmaceutically acceptable salt thereof for use in a method of treating a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associate bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human.

15

29. A compound according to any of claims 1-17 or a pharmaceutically acceptable salt thereof for use in a method of treating a disease associated with chronic obstructive pulmonary disease (COPD) or asthma, in a human.

20 30. The use of a compound according to any one of claims 1-17 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of pulmonary inflammation or bronchoconstriction in a human.

25 31. The use of a compound according to any one of claims 1-17 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD), asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associate bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human.

32. A composition comprising a compound according to any one of claims 1-17 or a pharmaceutically acceptable salt thereof for use in the preparation of a medicament for the treatment of pulmonary inflammation or bronchoconstriction in a human.

5

33. A composition comprising a compound according to any one of claims 1-17 or a pharmaceutically acceptable salt thereof for use in the preparation of a medicament for the treatment of a disease associated with reversible or irreversible airway obstruction, chronic obstructive pulmonary disease (COPD),

10 asthma, bronchiectasis (including bronchiectasis due to conditions other than cystic fibrosis), acute bronchitis, chronic bronchitis, post-viral cough, cystic fibrosis, emphysema, pneumonia, panbronchiolitis, transplant-associate bronchiolitis, sinusitis, and ventilator-associated tracheobronchitis or preventing ventilator-associated pneumonia, or treating sinusitis in a human.

15

INTERNATIONAL SEARCH REPORT

International application No PCT/US2011/035741

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D471/04 A61K31/437 A61P29/00 A61P37/08 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2009/100169 A1 (GLAXO GROUP LTD [GB]; CALLAHAN JAMES FRANCIS [US]; LI TINDY [US]; WAN) 13 August 2009 (2009-08-13) cited in the application the whole document -----	1-33
A	WO 2009/100170 A1 (GLAXO GROUP LTD [GB]; CALLAHAN JAMES FRANCIS [US]; WAN ZEHONG [CN]; YA) 13 August 2009 (2009-08-13) the whole document -----	1-33
A	WO 2004/024728 A2 (GLAXO GROUP LTD [GB]; ALLEN DAVID GEORGE [GB]; COE DIANE MARY [GB]; CO) 25 March 2004 (2004-03-25) the whole document -----	1-33

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
23 August 2011	05/09/2011
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Bissmire, Stewart

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/US2011/035741

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 2009100169	A1	13-08-2009	EP 2249646 A1	17-11-2010

WO 2009100170	A1	13-08-2009	EP 2249830 A1	17-11-2010

WO 2004024728	A2	25-03-2004	AT 442367 T AU 2003285300 A1 AU 2010201417 A1 BR 0314392 A CA 2497550 A1 CN 1694882 A DK 1539753 T3 EP 1539753 A2 ES 2331119 T3 IL 166898 A IS 7709 A JP 2006503108 A KR 20050052501 A MA 27475 A1 MX PA05002887 A NZ 538515 A PT 1539753 E SI 1539753 T1 US 2008175914 A1 US 2006089375 A1	15-09-2009 30-04-2004 29-04-2010 12-07-2005 25-03-2004 09-11-2005 04-01-2010 15-06-2005 22-12-2009 30-06-2010 24-02-2005 26-01-2006 02-06-2005 01-08-2005 27-05-2005 27-04-2007 10-12-2009 31-12-2009 24-07-2008 27-04-2006
