



US 20110190248A1

(19) **United States**(12) **Patent Application Publication****Himmelsbach et al.**(10) **Pub. No.: US 2011/0190248 A1**(43) **Pub. Date: Aug. 4, 2011**

(54) **CYCLOHEXYLOXY SUBSTITUTED  
HETEROCYCLES, PHARMACEUTICAL  
COMPOSITIONS CONTAINING THESE  
COMPOUNDS AND PROCESSES FOR  
PREPARING THEM**

*A61P 29/00* (2006.01)  
*A61P 37/08* (2006.01)  
*A61P 11/00* (2006.01)  
*A61P 11/06* (2006.01)  
*A61P 11/14* (2006.01)  
*A61P 35/00* (2006.01)

(75) Inventors: **Frank Himmelsbach**,  
Mittelbiberach (DE); **Birgit Jung**,  
Laupheim (DE); **Ralf Lotz**,  
Schemmerhofen (DE)

(52) **U.S. Cl. .... 514/171; 514/252.17; 544/293**

(73) Assignee: **Boehringer Ingelheim  
International GmbH**, Ingelheim  
am Rhein (DE)

(57) **ABSTRACT**


The present invention relates to cyclohexyloxy-substituted heterocycles of general formula (I)

(21) Appl. No.: **13/057,872**(22) PCT Filed: **Jul. 23, 2009**(86) PCT No.: **PCT/EP2009/059510**

§ 371 (c)(1),  
(2), (4) Date:

**Mar. 21, 2011**(30) **Foreign Application Priority Data**


Aug. 8, 2008 (EP) ..... 08104995.9

**Publication Classification**(51) **Int. Cl.***A61K 31/517* (2006.01)*A61K 31/56* (2006.01)*C07D 403/12* (2006.01)

the tautomers, the stereoisomers, the mixtures thereof and the salts thereof, particularly the physiologically acceptable salts thereof with inorganic or organic acids, which have valuable pharmacological properties, particularly an inhibitory effect on signal transduction mediated by tyrosine kinases, the use thereof for the treatment of diseases, particularly tumoral diseases as well as benign prostatic hyperplasia (BPH), diseases of the lungs and airways and the preparation thereof.

**CYCLOHEXYLOXY SUBSTITUTED  
HETEROCYCLES, PHARMACEUTICAL  
COMPOSITIONS CONTAINING THESE  
COMPOUNDS AND PROCESSES FOR  
PREPARING THEM**

**[0001]** The present invention relates to cyclohexyloxy-substituted heterocycles of general formula




the tautomers, the stereoisomers, the mixtures thereof and the salts thereof, particularly the physiologically acceptable salts thereof with inorganic or organic acids, which have valuable pharmacological properties, particularly an inhibitory effect on signal transduction mediated by tyrosine kinases, the use thereof for the treatment of diseases, particularly tumoral diseases as well as benign prostatic hyperplasia (BPH), diseases of the lungs and airways and the preparation thereof.

**[0002]** The problem of the present invention is to prepare new compounds which on the basis of their pharmaceutical effectiveness as tyrosine-kinase inhibitors, may be used therapeutically, i.e. for the treatment of pathophysiological processes caused by hyperfunction of tyrosine kinases.

**DETAILED DESCRIPTION OF THE INVENTION**

**[0003]** It has surprisingly been found that the problem mentioned above is solved by compounds of formula (I), wherein the groups  $R^a$  to  $R^d$  and A have the meanings given hereinafter. The present invention therefore relates to compounds of general formula (I),



wherein

**[0004]**  $R^a$  denotes a phenyl or 1-phenylethyl group, wherein the phenyl nucleus is substituted in each case by the groups  $R^1$  to  $R^3$ , where

**[0005]**  $R^1$  and  $R^2$  which may be identical or different, denote hydrogen or a group selected from among

**[0006]** F, Cl, Br, I,  $\text{OCH}_2\text{F}$ ,  $\text{OCHF}_2$ ,  $\text{OCF}_3$ ,  $\text{CH}_2\text{F}$ ,  $\text{CHF}_2$ ,  $\text{CF}_3$ , CN,  $\text{NO}_2$ ,  $\text{NH}_2$  and OH,

**[0007]** or

**[0008]** a group selected from among

**[0009]**  $\text{C}_{1-4}$ -alkyl,  $\text{C}_{1-4}$ -alkyl-O,  $\text{C}_{2-3}$ -alkenyl,  $\text{C}_{2-3}$ -alkynyl, phenyl, phenyl-O, phenyl- $\text{C}_{1-3}$ -alkyl- and phenyl- $\text{C}_{1-3}$ -alkyl-O, heteroaryl, heteroaryl-O, heteroaryl- $\text{C}_{1-3}$ -alkyl and heteroaryl- $\text{C}_{1-3}$ -alkyl-O, while the above-mentioned phenyl groups are mono- or disubstituted by  $R^5$  groups,

**[0010]** and

**[0011]**  $R^3$  denotes hydrogen,

**[0012]** or

**[0013]** a group selected from among

**[0014]** F, Cl, Br and  $\text{CH}_3$ ,

**[0015]**  $R^b$  denotes hydrogen, or an optionally substituted group selected from among  $\text{C}_{1-6}$ -alkyl,  $\text{C}_{3-6}$ -cycloalkyl and  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl,

**[0016]**  $R^c$  denotes hydrogen, or an optionally substituted group selected from among  $\text{C}_{1-6}$ -alkyl,  $\text{C}_{3-6}$ -cycloalkyl,  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl,  $\text{C}_{1-6}$ -alkyl-CO,  $\text{C}_{3-6}$ -cycloalkyl-CO,  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl-CO,  $\text{C}_{1-6}$ -alkyl- $\text{SO}_2$ ,  $\text{C}_{3-6}$ -cycloalkyl- $\text{SO}_2$ ,  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl- $\text{SO}_2$ , phenyl-CO— and phenyl- $\text{SO}_2$ ,

**[0017]**  $R^d$  denotes hydrogen or

**[0018]** a group selected from among

**[0019]** F, Cl, Br, I, OH,  $\text{C}_{1-4}$ -alkyl,  $\text{C}_{1-4}$ -alkyl-O,  $\text{C}_{1-2}$ -alkyl-O substituted by 1 to 3 fluorine atoms,  $\text{C}_{3-7}$ -cycloalkyl-O,  $\text{C}_{3-7}$ -cycloalkyl- $\text{C}_{1-4}$ -alkyl-O, tetrahydrofuran-3-yl-O, tetrahydropyran-3-yl-O, tetrahydro-pyran-4-yl-O, tetrahydrofuranyl- $\text{C}_{1-4}$ -alkyl-O and tetrahydropyranyl- $\text{C}_{1-4}$ -alkyl-O,

**[0020]** or

**[0021]**  $R^4$ — $\text{C}_{1-4}$ -alkyl, wherein the linking of the groups  $R^4$  may take place via each C atom of the alkyl group,

**[0022]** or

**[0023]**  $R^4$ — $\text{C}_{2-4}$ -alkyl-O, wherein the group  $R^4$  is separated from the oxygen atom by at least 2 C atoms,

**[0024]** or

**[0025]** a group selected from among

**[0026]** pyrrolidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, pyrrolidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, piperidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, piperidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, piperidin-4-yl- $\text{C}_{1-4}$ -alkyl-O, azepan-2-yl- $\text{C}_{1-4}$ -alkyl-O, azepan-3-yl- $\text{C}_{1-4}$ -alkyl-O, azepan-4-yl- $\text{C}_{1-4}$ -alkyl-O, morpholin-2-yl- $\text{C}_{1-4}$ -alkyl-O, morpholin-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-pyrrolidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-pyrrolidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-piperidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-piperidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-piperidin-4-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-azepan-2-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-azepan-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-azepan-4-yl- $\text{C}_{1-4}$ -alkyl-O, 4-( $\text{C}_{1-3}$ -alkyl)-morpholin-2-yl- $\text{C}_{1-4}$ -alkyl-O and 4-( $\text{C}_{1-3}$ -alkyl)-morpholin-3-yl- $\text{C}_{1-4}$ -alkyl-O,

**[0027]** wherein

**[0028]**  $R^4$  denotes a group, which may be identical or different, selected from among

**[0029]** OH,  $\text{C}_{1-3}$ -alkyl-O,  $\text{C}_{3-6}$ -cycloalkyl-O,  $\text{NH}_2$ ,  $\text{C}_{1-3}$ -alkyl-NH, ( $\text{C}_{1-3}$ -alkyl) $_2\text{N}$ , (2-methoxyethyl) $_2\text{N}$ , pyrrolidin-1-yl, piperidin-1-yl, azepan-1-yl, morpholin-4-yl, 1,4-oxazepan-4-yl, 2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl, 3-oxa-8-aza-bicyclo[3.2.1]oct-8-yl, 8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl, piperazin-1-yl, 4-( $\text{C}_{1-3}$ -alkyl)-piperazin-1-yl, 1,4-diazepan-1-yl, 4-( $\text{C}_{1-3}$ -alkyl)-1,4-diazepan-1-yl,

HCO—NH, C<sub>1-4</sub>-alkyl-CO—NH, C<sub>1-3</sub>-alkyl-O—C<sub>1-3</sub>-alkyl-CO—NH, C<sub>1-4</sub>-alkyl-O—CO—NH, H<sub>2</sub>NCONH, C<sub>1-3</sub>-alkyl-NH—CO—NH, (C<sub>1-3</sub>-alkyl)<sub>2</sub>N—CONN, pyrrolidin-1-yl-CO—NH, piperidin-1-yl-CO—NH, piperazin-1-yl-CO—NH, 4-(C<sub>1-3</sub>-alkyl)-piperazin-1-yl-CO—NH, morpholin-4-yl-CO—NH and C<sub>1-4</sub>-alkyl-SO<sub>2</sub>—NH,

while the pyrrolidinyl, piperidinyl, azepan-1-yl, piperazinyl, 1,4-diazepan-1-yl, morpholinyl and 1,4-oxazepan-4-yl groups mentioned above in the definition of the group R<sup>d</sup> may each additionally be substituted by one or two C<sub>1-3</sub>-alkyl groups,

[0030] and

wherein the above-mentioned phenyl groups are mono- or disubstituted by groups R<sup>5</sup>, wherein

[0031] R<sup>5</sup> denotes hydrogen, or

[0032] a group, which may be identical or different, selected from among F, Cl, Br, I, OH, CN, C<sub>1-3</sub>-alkyl, C<sub>1-3</sub>-alkyl-O, CHF<sub>2</sub>, CF<sub>3</sub>, —O—CHF<sub>2</sub> and —O—CF<sub>3</sub>,

[0033] and

unless stated otherwise, the above-mentioned alkyl groups may be straight-chain or branched,

[0034] A denotes —CO or —C<sub>1-3</sub>-alkylene,

[0035] wherein the —C<sub>1-3</sub>-alkylene group may be 1-, 2-, 3- or 4-substituted by a group R<sup>6</sup>,

[0036] and

[0037] R<sup>6</sup> which may be identical or different, denotes hydrogen, or

[0038] a group selected from among OH, C<sub>1</sub>-C<sub>4</sub>-alkyl and —O—C<sub>1</sub>-C<sub>4</sub>-alkyl

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof, and the solvates and hydrates thereof.

[0039] Preferred compounds of formula (I) are those wherein

[0040] R<sup>a</sup> denotes a group selected from among 3-chloro-2-fluoro-phenyl, 3-chloro-4-fluoro-phenyl, 5-chloro-2-fluoro-phenyl, 2-fluoro-3-methyl-phenyl, 2-fluoro-5-methyl-phenyl, 4-fluoro-3-methyl-phenyl and 3-chloro-2-methyl-phenyl,

[0041] R<sup>b</sup> and R<sup>c</sup> which may be identical or different,

[0042] denote hydrogen or C<sub>1-3</sub>-alkyl,

[0043] R<sup>d</sup> denotes C<sub>1-3</sub>-alkyl-O,

unless stated otherwise, the above-mentioned alkyl groups may be straight-chain or branched,

[0044] A denotes —CH<sub>2</sub>CH<sub>2</sub>, wherein the —CH<sub>2</sub>CH<sub>2</sub>— group may be substituted by 1 or 2 methyl groups,

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof, and the solvates and hydrates thereof.

[0045] The invention further relates to compounds of formula (I) for use as medicaments. Preferably the compounds of formula (I) are used in cases of inflammatory or allergic diseases of the airways.

[0046] The compounds of formula (I) are particularly preferably used in cases of a disease selected from among chronic bronchitis, acute bronchitis, bronchitis caused by bacterial or viral infection or fungi or helminths, allergic bronchitis, toxic bronchitis, chronic obstructive bronchitis (COPD), asthma (intrinsic or allergic), paediatric asthma, bronchiectasis, allergic alveolitis, allergic or non-allergic rhinitis, chronic sinusi-

tis, cystic fibrosis or mucoviscidosis, alpha-1-antitrypsin deficiency, cough, pulmonary emphysema, interstitial lung diseases, alveolitis, hyperreactive airways, nasal polyps, pulmonary oedema, pneumonitis of different origins, e.g. radiation-induced or caused by aspiration or infectious pneumonitis, collagenoses such as lupus erythematoses, systemic sclerodermy, sarcoidosis and Boeck's disease.

[0047] It is also particularly preferred to use the compounds of formula (I) in cases of inflammatory or allergic complaints in which autoimmune reactions are involved. It is also particularly preferred to use the compounds of formula (I) in cases of a disease in the form of benign or malignant tumours.

[0048] The invention further relates to a pharmaceutical formulation containing a compound of formula (I).

[0049] Preferably an orally administered pharmaceutical formulation containing a compound of formula (I) is used.

[0050] The invention further relates to medicament combinations which contain, besides one or more compounds of formula (I), as further active substances, one or more compounds selected from among the categories of betamimetics, anticholinergics, corticosteroids, further PDE4-inhibitors, LTD4-antagonists, EGFR-inhibitors, dopamine agonists, H1-antihistamines, PAF-antagonists and PI3-kinase inhibitors or double or triple combinations thereof.

[0051] Suitable betamimetics used are preferably compounds selected from among arformoterol, carmoterol, formoterol, indacaterol, salmeterol, albuterol, bambuterol, bitolterol, broxaterol, carbutoleol, clenbuterol, fenoterol, hexoprenaline, ibuterol, isoetharine, isoprenaline, levosalbutamol, mabuterol, meluadrine, metaproterenol, milveterol, orciprenaline, pirbuterol, procaterol, reproterol, rimiterol, ritodrine, salmefamol, soterenol, sulphonterol, terbutaline, tiaramide, tolubuterol, zinterol and

[0052] 6-hydroxy-8-{1-hydroxy-2-[2-(4-methoxy-phenyl)-1,1-dimethyl-ethylamino]-ethyl}-4H-benzo[1,4]oxazin-3-one

[0053] 8-{2-[2-(2,4-difluoro-phenyl)-1,1-dimethyl-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one

[0054] 8-{2-[2-(3,5-difluoro-phenyl)-1,1-dimethyl-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one

[0055] 8-{2-[2-(4-ethoxy-phenyl)-1,1-dimethyl-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one

[0056] 8-{2-[2-(4-fluoro-phenyl)-1,1-dimethyl-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one

[0057] N-(5-{2-[3-(4,4-diethyl-2-oxo-4H-benzo[d][1,3]oxazin-1-yl)-1,1-dimethyl-propylamino]-1-hydroxy-ethyl}-2-hydroxy-phenyl)-methanesulphonamide

[0058] N-(5-{2-[3-(4,4-diethyl-6-fluoro-2-oxo-4H-benzo[d][1,3]oxazin-1-yl)-1,1-dimethyl-propylamino]-1-hydroxy-ethyl}-2-hydroxy-phenyl)-methanesulphonamide

[0059] N-(5-{2-[3-(4,4-diethyl-6-methoxy-2-oxo-4H-benzo[d][1,3]oxazin-1-yl)-1,1-dimethyl-propylamino]-1-hydroxy-ethyl}-2-hydroxy-phenyl)-methanesulphonamide

[0060] N-(5-{2-[1,1-dimethyl-3-(2-oxo-4,4-dipropyl-4H-benzo[d][1,3]oxazin-1-yl)-propylamino]-1-hydroxy-ethyl}-2-hydroxy-phenyl)-methanesulphonamide

[0061] 8-{2-[1,1-dimethyl-3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-propylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one

- [0062] 8-{2-[1,1-dimethyl-3-(6-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one
- [0063] 8-{2-[1,1-dimethyl-3-(2-oxo-5-trifluoromethyl-2,3-dihydro-benzimidazol-1-yl)-propylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one
- [0064] 8-{2-[1,1-dimethyl-3-(3-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one
- [0065] N-[2-hydroxy-5-((1R)-1-hydroxy-2-{2-[4-(2-hydroxy-2-phenyl-ethylamino)-phenyl]-ethylamino}-ethyl)-phenyl]-formamide
- [0066] 8-hydroxy-5-((1R)-1-hydroxy-2-{2-[4-(6-methoxy-biphenyl-3-ylamino)-phenyl]-ethylamino}-ethyl)-1H-quinolin-2-one
- [0067] 8-hydroxy-5-[(1R)-1-hydroxy-2-(6-phenethylamino-hexylamino)-ethyl]-1H-quinolin-2-one
- [0068] 5-[(1R)-2-(2-{4-[4-(2-Amino-2-methyl-propoxy)-phenylamino]-phenyl]-ethylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one
- [0069] [3-(4-{6-[(2R)-2-hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]-hexyloxy}-butyl)-5-methyl-phenyl]-urea
- [0070] 4-((1R)-2-{6-[2-(2,6-dichloro-benzyloxy)-ethoxy]-hexylamino}-1-hydroxy-ethyl)-2-hydroxymethyl-phenol
- [0071] 3-(4-{6-[(2R)-2-hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]-hexyloxy}-butyl)-benzenesulphonamide
- [0072] 3-(3-{7-[(2R)-2-hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]-heptyloxy}-propyl)-benzenesulphonamide
- [0073] 4-((1R)-2-{6-[4-(3-cyclopentanesulphonyl-phenyl)-butoxy]-hexylamino}-1-hydroxy-ethyl)-2-hydroxymethyl-phenol
- [0074] N-1-adamantanyl-2-{3-[(2R)-2-((2R)-2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl)amino]propyl}phenyl]acetamide
- [0075] (1R)-5-{2-[6-(2,2-difluoro-2-phenyl-ethoxy)-hexylamino]-1-hydroxy-ethyl}-8-hydroxy-1H-quinolin-2-one
- [0076] (R,S)-4-(2-{[6-(2,2-difluoro-4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol
- [0077] (R,S)-4-(2-{[6-(2,2-difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol
- [0078] (R,S)-4-(2-{[4,4-difluoro-6-(4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol
- [0079] (R,S)-4-(2-{[6-(4,4-difluoro-4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol
- [0080] (R,S)-5-(2-{[6-(2,2-difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxy-ethyl)-8-hydroxyquinolin-2(1H)-one
- [0081] (R,S)-[2-({[6-[2,2-difluoro-2-(3-methylphenyl)ethoxy]hexyl]amino)-1-hydroxyethyl]-2-(hydroxymethyl)phenol
- [0082] 4-(1R)-2-{[6-(2,2-difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxyethyl)-2-(hydroxymethyl)phenol
- [0083] (R,S)-2-(hydroxymethyl)-4-(1-hydroxy-2-{[4,4,5,5-tetrafluoro-6-(3-phenylpropoxy)-hexyl]amino}ethyl)phenol
- [0084] (R,S)-[5-(2-{[6-(2,2-difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxyphenyl)]formamide
- [0085] (R,S)-4-[2-({[6-[2-(3-bromophenyl)-2,2-difluoroethoxy]hexyl]amino)-1-hydroxyethyl]-2-(hydroxymethyl)phenol
- [0086] (R,S)-N-[3-(1,1-difluoro-2-{[6-({2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]-ethyl]amino)hexyl]oxy}ethyl)phenyl]-urea
- [0087] 3-[3-(1,1-difluoro-2-{[6-({2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl]-amino)hexyl]oxy}ethyl)phenyl]imidazolidine-2,4-dione
- [0088] (R,S)-4-[2-({[6-[2,2-difluoro-2-(3-methoxyphenyl)ethoxy]hexyl]amino)-1-hydroxyethyl]-2-(hydroxymethyl)phenol
- [0089] 5-((1R)-2-{[6-(2,2-difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxyethyl)-8-hydroxyquinolin-2(1H)-one
- [0090] 4-((1R)-2-{[4,4-difluoro-6-(4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol
- [0091] (R,S)-4-(2-{[6-(3,3-difluoro-3-phenylpropoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol
- [0092] (R,S)-(2-{[6-(2,2-difluoro-2-phenylethoxy)-4,4-difluoro]hexyl]amino}-1-hydroxyethyl)-2-(hydroxymethyl)phenol
- [0093] (R,S)-4-(2-{[6-(2,2-difluoro-3-phenylpropoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol
- [0094] 3-[2-(3-chloro-phenyl)-ethoxy]-N-(2-diethylamino-ethyl)-N-{2-[2-(4-hydroxy-2-oxo-2,3-dihydro-benzothiazol-7-yl)-ethylamino]-ethyl}-propionamide
- [0095] N-(2-diethylamino-ethyl)-N-{2-[2-(4-hydroxy-2-oxo-2,3-dihydro-benzothiazol-7-yl)-ethylamino]-ethyl}-3-(2-naphthalen-1-yl-ethoxy)-propionamide
- [0096] 7-[2-(2-{3-[2-(2-chloro-phenyl)-ethylamino]-propylsulphonyl]-ethylamino)-1-hydroxy-ethyl]-4-hydroxy-3H-benzothiazol-2-one

optionally in the form of the racemates, enantiomers, diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts, solvates or hydrates thereof. According to the invention the acid addition salts of the betamimetics are preferably selected from among the hydrochloride, hydrobromide, hydriodide, hydrosulphate, hydrophosphate, hydromethanesulphonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulphonate.

[0097] The anticholinergics used are preferably compounds selected from among the tiotropium salts, preferably the bromide salt, oxitropium salts, preferably the bromide salt, flutropium salts, preferably the bromide salt, ipratropium salts, preferably the bromide salt, aclidinium salts, preferably the bromide salt, glycopyrronium salts, preferably the bromide salt, trospium salts, preferably the chloride salt, tolterodine, (3R)-1-phenethyl-3-(9H-xanthen-9-carbonyl)-1-azoniabicyclo[2.2.2]octane salts. In the above-mentioned salts the cations are the pharmacologically active constituents. As anions the above-mentioned salts may preferably contain the chloride, bromide, iodide, sulphate, phosphate, methanesulphonate, nitrate, maleate, acetate, citrate, fumarate, tartrate, oxalate, succinate, benzoate or p-toluenesulphonate, while chloride, bromide, iodide, sulphate, methanesulpho-

nate or p-toluenesulphonate are preferred as counter-ions. Of all the salts the chlorides, bromides, iodides and methane-sulphonates are particularly preferred.

[0098] Other specified compounds are:

- [0099] tropenol 2,2-diphenylpropionate methobromide,
- [0100] scopine 2,2-diphenylpropionate methobromide,
- [0101] scopine 2-fluoro-2,2-diphenylacetate methobromide,
- [0102] tropenol 2-fluoro-2,2-diphenylacetate methobromide;
- [0103] tropenol 3,3',4,4'-tetrafluorobenzilate methobromide,
- [0104] scopine 3,3',4,4'-tetrafluorobenzilate methobromide,
- [0105] tropenol 4,4'-difluorobenzilate methobromide,
- [0106] scopine 4,4'-difluorobenzilate methobromide,
- [0107] tropenol 3,3'-difluorobenzilate methobromide,
- [0108] scopine 3,3'-difluorobenzilate methobromide;
- [0109] tropenol 9-hydroxy-fluorene-9-carboxylate methobromide;
- [0110] tropenol 9-fluoro-fluorene-9-carboxylate methobromide;
- [0111] scopine 9-hydroxy-fluorene-9-carboxylate methobromide;
- [0112] scopine 9-fluoro-fluorene-9-carboxylate methobromide;
- [0113] tropenol 9-methyl-fluorene-9-carboxylate methobromide;
- [0114] scopine 9-methyl-fluorene-9-carboxylate methobromide;
- [0115] cyclopropyltropine benzilate methobromide;
- [0116] cyclopropyltropine 2,2-diphenylpropionate methobromide;
- [0117] cyclopropyltropine 9-hydroxy-xanthene-9-carboxylate methobromide;
- [0118] cyclopropyltropine 9-methyl-fluorene-9-carboxylate methobromide;
- [0119] cyclopropyltropine 9-methyl-xanthene-9-carboxylate methobromide;
- [0120] cyclopropyltropine 9-hydroxy-fluorene-9-carboxylate methobromide;
- [0121] cyclopropyltropine methyl 4,4'-difluorobenzilate methobromide.
- [0122] tropenol 9-hydroxy-xanthene-9-carboxylate methobromide;
- [0123] scopine 9-hydroxy-xanthene-9-carboxylate methobromide;
- [0124] tropenol 9-methyl-xanthene-9-carboxylate methobromide;
- [0125] scopine 9-methyl-xanthene-9-carboxylate methobromide;
- [0126] tropenol 9-ethyl-xanthene-9-carboxylate methobromide;
- [0127] tropenol 9-difluoromethyl-xanthene-9-carboxylate methobromide;
- [0128] scopine 9-hydroxymethyl-xanthene-9-carboxylate methobromide,
- [0129] The above-mentioned compounds may also be used as salts within the scope of the present invention, wherein instead of the methobromide the salts metho-X are used, wherein X may have the meanings given hereinbefore for X<sup>-</sup>.
- [0130] As corticosteroids it is preferable to use compounds selected from among beclomethasone, betamethasone, budesonide, butixocort, ciclesonide, deflazacort, dexametha-

sone, etiprednol, flunisolide, fluticasone, loteprednol, mometasone, prednisolone, prednisone, rofleponide, triamcinolone, tipredane and

- [0131] pregna-1,4-diene-3,20-dione, 6-fluoro-11-hydroxy-16,17-[(1-methylethylidene) bis(oxy)]-21-[[4-[(nitrooxy)methyl]benzoyl]oxy]-, (6- $\alpha$ ,11- $\beta$ ,16- $\alpha$ -ph)-(9CI) (NCX-1024)
  - [0132] 16,17-butylenedioxy-6,9-difluoro-11-hydroxy-17-(methylthio)androst-4-en-3-one (RPR-106541),
  - [0133] (S)-fluoromethyl 6,9-difluoro-17-[(2-furanylcarbonyl)oxy]-11-hydroxy-16-methyl-3-oxo-androsta-1,4-diene-17-carbothionate
  - [0134] (S)-(2-oxo-tetrahydro-furan-3S-yl) 6,9-difluoro-11-hydroxy-16-methyl-3-oxo-17-propionyloxy-androsta-1,4-diene-17-carbothionate,
  - [0135] cyanomethyl 6- $\alpha$ ,9- $\alpha$ -difluoro-11- $\beta$ -hydroxy-16 $\alpha$ -methyl-3-oxo-17 $\alpha$ -(2,2,3,3-tetramethylcyclopropylcarbonyl)oxy-androsta-1,4-diene-17 $\beta$ -carboxylate,
- optionally in the form of the racemates, enantiomers or diastereomers thereof and optionally in the form of the salts and derivatives thereof, the solvates and/or hydrates thereof. Any reference to steroids includes a reference to any salts or derivatives, hydrates or solvates thereof which may exist. Examples of possible salts and derivatives of the steroids may be: alkali metal salts, such as for example sodium or potassium salts, sulphobenzoates, phosphates, isonicotinates, acetates, dichloroacetates, propionates, dihydrogen phosphates, palmitates, pivalates or furoates.
- [0136] PDE4-inhibitors which may be used are preferably compounds selected from among enprofyllin, theophyllin, roflumilast, ariflo (cilomilast), tofomilast, pumafentrin, lir- imilast, apremilast, arofyllin, atizoram, oglemilast, tetomilas- tnd
  - [0137] 5-[(N-(2,5-dichloro-3-pyridinyl)-carboxamid]-8-methoxy-quinoline (D-4418),
  - [0138] N-(3,5-dichloro-1-oxido-4-pyridinyl)-carboxa- mid]-8-methoxy-2-(trifluoromethyl)-quinoline (D-4396 (Sch-351591)), N-(3,5-dichloropyrid-4-yl)-[1-(4-fluo- robenzyl)-5-hydroxy-indol-3-yl]glyoxylic acid amide (AWD-12-281 (GW-842470)), 9-[(2-fluorophenyl)methyl]-N-methyl-2-(trifluoromethyl)-9H-purin-6-amine (NCS-613),
  - [0139] 4-[(2R)-2-[3-(cyclopentylloxy)-4-methoxyphenyl]- 2-phenylethyl]-pyridine (CDP-840),
  - [0140] N-[(3R)-3,4,6,7-tetrahydro-9-methyl-4-oxo-1-phe- nylpyrrolo[3,2,1-jk][1,4]benzodiazepin-3-yl]-4-pyridin- ecarboxamide (PD-168787),
  - [0141] 4-[6,7-diethoxy-2,3-bis(hydroxymethyl)-1-naph- thalenyl]-1-(2-methoxyethyl)-2(1H)-pyridinone (T-440),
  - [0142] 2-[4-[6,7-diethoxy-2,3-bis(hydroxymethyl)-1- naphthalenyl]-2-pyridinyl]-4-(3-pyridinyl)-1(2H)-ph- thalazinone (T-2585),
  - [0143] (3-(3-cyclopentylloxy-4-methoxybenzyl)-6-ethy- lamino-8-isopropyl-3H-purine (V-11294A),
  - [0144] beta-[3-(cyclopentylloxy)-4-methoxyphenyl]-1,3- dihydro-1,3-dioxo-2H-Isoindole-2-propanamide (CDC- 801),
  - [0145] ilmidazo[1,5-a]pyrido[3,2-e]pyrazin-6(5H)-one, 9-ethyl-2-methoxy-7-methyl-5-propyl-(D-22888)
  - [0146] 5-[3-(cyclopentylloxy)-4-methoxyphenyl]-3-[(3- methylphenyl)methyl]-, (3S,5S)-2-piperidinone (HT- 0712),

- [0147] 4-[1-[3,4-bis(difluoromethoxy)phenyl]-2-(3-methyl-1-oxido-4-pyridinyl)ethyl]-alpha,alpha-bis(trifluoromethyl)-benzenemethanol (L-826141)
- [0148] N-(3,5-dichloro-1-oxo-pyridin-4-yl)-4-difluoromethoxy-3-cyclopropylmethoxybenzamide
- [0149] (-)-p-[(4aR\*,10bS\*)-9-ethoxy-1,2,3,4,4a,10b-hexahydro-8-methoxy-2-methylbenzo[s][1,6]naphthyridin-6-yl]-N,N-diisopropylbenzamide
- [0150] (R)-(+)-1-(4-bromobenzyl)-4-[(3-cyclopentyl-4-methoxyphenyl)-2-pyrrolidone
- [0151] 3-(cyclopentyl-4-methoxyphenyl)-1-(4-N'-[N-2-cyano-S-methyl-isothioureido]benzyl)-2-pyrrolidone
- [0152] cis[4-cyano-4-(3-cyclopentyl-4-methoxyphenyl)cyclohexan-1-carboxylic acid]
- [0153] 2-carbomethoxy-4-cyano-4-(3-cyclopropyl-methoxy-4-difluoromethoxyphenyl)cyclohexan-1-one
- [0154] cis[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol]
- [0155] (R)-(+)-ethyl[4-(3-cyclopentyl-4-methoxyphenyl)pyrrolidin-2-ylidene]acetate
- [0156] (S)-(-)-ethyl[4-(3-cyclopentyl-4-methoxyphenyl)pyrrolidin-2-ylidene]acetate
- [0157] 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(2-thienyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine
- [0158] 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(tert-butyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine, optionally in the form of the racemates, enantiomers or diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts thereof, the solvates and/or hydrates thereof. According to the invention the acid addition salts are preferably selected from among the hydrochloride, hydrobromide, hydriodide, hydrosulphate, hydrophosphate, hydromethanesulphonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulphonate.
- [0159] Preferred EGFR-inhibitors used are preferably compounds selected from among cetuximab, trastuzumab, panitumumab (=ABX-EGF), Mab ICR-62, gefitinib, caneritinib, erlotinib, and
- [0160] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-cyclopropylmethoxy-quinazoline,
- [0161] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-cyclopropylmethoxy-quinazoline,
- [0162] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-cyclopropylmethoxy-quinazoline,
- [0163] 4-[(R)-(1-phenyl-ethyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-cyclopentyl-oxo-quinazoline,
- [0164] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[[4-((R)-6-methyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-cyclopropylmethoxy-quinazoline
- [0165] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[[4-((R)-6-methyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-[(S)-(tetrahydrofuran-3-yl)oxy]-quinazoline
- [0166] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[[4-((R)-2-methoxymethyl-6-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-cyclopropylmethoxy-quinazoline
- [0167] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[2-((S)-6-methyl-2-oxo-morpholin-4-yl)-ethoxy]-7-methoxy-quinazoline 4-[(3-chloro-4-fluorophenyl)amino]-6-([4-
- [N-(2-methoxy-ethyl)-N-methyl-amino]-1-oxo-2-buten-1-yl]-amino)-7-cyclopropylmethoxy-quinazoline,
- [0168] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-cyclopentyl-oxo-quinazoline,
- [0169] 4-[(R)-(1-phenyl-ethyl)amino]-6-[[4-(N,N-bis-(2-methoxy-ethyl)-amino)-1-oxo-2-buten-1-yl]-amino]-7-cyclopropylmethoxy-quinazoline,
- [0170] 4-[(R)-(1-phenyl-ethyl)amino]-6-([4-N-(2-methoxy-ethyl)-N-ethyl-amino]-1-oxo-2-buten-1-yl)-amino)-7-cyclopropylmethoxy-quinazoline,
- [0171] 4-[(R)-(1-phenyl-ethyl)amino]-6-([4-N-(2-methoxy-ethyl)-N-methyl-amino]-1-oxo-2-buten-1-yl)-amino)-7-cyclopropylmethoxy-quinazoline,
- [0172] 4-[(R)-(1-phenyl-ethyl)amino]-6-([4-N-(tetrahydrofuran-4-yl)-N-methyl-amino]-1-oxo-2-buten-1-yl)-amino)-7-cyclopropylmethoxy-quinazoline,
- [0173] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-((R)-tetrahydrofuran-3-yloxy)-quinazoline,
- [0174] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline
- [0175] 4-[(3-chloro-4-fluorophenyl)amino]-6-([4-N-(2-methoxy-ethyl)-N-methyl-amino]-1-oxo-2-buten-1-yl)-amino)-7-cyclopentyl-oxo-quinazoline,
- [0176] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N-cyclopropyl-N-methyl-amino)-1-oxo-2-buten-1-yl]-amino]-7-cyclopentyl-oxo-quinazoline,
- [0177] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-[(R)-(tetrahydrofuran-2-yl)methoxy]-quinazoline,
- [0178] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]-quinazoline,
- [0179] 4-[(3-ethynyl-phenyl)amino]-6,7-bis-(2-methoxy-ethoxy)-quinazoline,
- [0180] 4-[(3-chloro-4-fluorophenyl)amino]-7-[3-(morpholin-4-yl)-propyloxy]-6-Rvinylcarbonyl amin\*quinazoline,
- [0181] 4-[(R)-(1-phenyl-ethyl)amino]-6-(4-hydroxy-phenyl)-7H-pyrrolo[2,3-d]pyrimidin,
- [0182] 3-cyano-4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]-amino]-7-ethoxy-quinoline,
- [0183] 4-[[3-chloro-4-(3-fluoro-benzyloxy)-phenyl]amino]-6-(5-[[2-methanesulphonyl-ethyl]amino]methyl)-furan-2-yl)quinazoline,
- [0184] 4-[(R)-(1-phenyl-ethyl)amino]-6-[[4-((R)-6-methyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-methoxy-quinazoline,
- [0185] 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-[(tetrahydrofuran-2-yl)methoxy]-quinazoline,
- [0186] 4-[(3-chloro-4-fluorophenyl)amino]-6-([4-N,N-bis-(2-methoxy-ethyl)-amino]-1-oxo-2-buten-1-yl)-amino)-7-[(tetrahydrofuran-2-yl)methoxy]-quinazoline,
- [0187] 4-[(3-ethynyl-phenyl)amino]-6-[[4-(5,5-dimethyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]-amino]-7-quinazoline,
- [0188] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-7-methoxy-quinazoline,

- [0189] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-7-[(R)-(tetrahydrofuran-2-ylmethoxy)-quinazoline,
- [0190] 4-[(3-chloro-4-fluoro-phenyl)amino]-7-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-6-[(S)-(tetrahydrofuran-2-yl)methoxy]-quinazoline,
- [0191] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[2-[4-(2-oxo-morpholin-4-yl)-piperidin-1-yl]-ethoxy]-7-methoxy-quinazoline,
- [0192] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-(tert.-butyloxycarbonyl)-piperidin-4-yloxy]-7-methoxy-quinazoline,
- [0193] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(trans-4-amino-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0194] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(trans-4-methanesulphonylamino-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0195] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(tetrahydropyran-3-yloxy)-7-methoxy-quinazoline,
- [0196] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-methyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0197] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy]-7-methoxy-quinazoline,
- [0198] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-[(methoxymethyl)carbonyl]-piperidin-4-yl-oxy]-7-methoxy-quinazoline,
- [0199] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(piperidin-3-yloxy)-7-methoxy-quinazoline,
- [0200] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-(2-acetylamino-ethyl)-piperidin-4-yloxy]-7-methoxy-quinazoline,
- [0201] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-ethoxy-quinazoline,
- [0202] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[(S)-tetrahydrofuran-3-yloxy]-7-hydroxy-quinazoline,
- [0203] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methoxy-ethoxy)-quinazoline,
- [0204] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[trans-4-[(dimethylamino)sulphonylamino]-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0205] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[trans-4-[(morpholin-4-yl)carbonylamino]-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0206] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[trans-4-[(morpholin-4-yl)sulfonylamino]-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0207] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-acetylamino-ethoxy)-quinazoline,
- [0208] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methanesulphonylamino-ethoxy)-quinazoline,
- [0209] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-[(piperidin-1-yl)carbonyl]-piperidin-4-yloxy]-7-methoxy-quinazoline,
- [0210] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-aminocarbonylmethyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0211] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(cis-4-[N-[(tetrahydropyran-4-yl)carbonyl]-N-methyl-amino]-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0212] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(cis-4-[N-[(morpholin-4-yl)carbonyl]-N-methyl-amino]-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0213] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(cis-4-[N-[(morpholin-4-yl)sulfonyl]-N-methyl-amino]-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0214] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(trans-4-ethanesulphonylamino-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0215] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-methanesulphonyl-piperidin-4-yloxy)-7-ethoxy-quinazoline,
- [0216] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-methanesulphonyl-piperidin-4-yloxy)-7-(2-methoxy-ethoxy)-quinazoline,
- [0217] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-(2-methoxy-acetyl)-piperidin-4-yloxy]-7-(2-methoxy-ethoxy)-quinazoline,
- [0218] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(cis-4-acetylamino-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0219] 4-[(3-ethynyl-phenyl)amino]-6-[1-(tert.-butyloxycarbonyl)-piperidin-4-yloxy]-7-methoxy-quinazoline,
- [0220] 4-[(3-ethynyl-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-methoxy-quinazoline,
- [0221] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(cis-4-[N-[(piperidin-1-yl)carbonyl]-N-methyl-amino]-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0222] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(cis-4-[N-[(4-methyl-piperazin-1-yl)carbonyl]-N-methyl-amino]-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0223] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[cis-4-[(morpholin-4-yl)carbonylamino]-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0224] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-[2-(2-oxopyrrolidin-1-yl)ethyl]-piperidin-4-yloxy]-7-methoxy-quinazoline,
- [0225] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-[(morpholin-4-yl)carbonyl]-piperidin-4-yloxy]-7-(2-methoxy-ethoxy)-quinazoline,
- [0226] 4-[(3-ethynyl-phenyl)amino]-6-(1-acetyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0227] 4-[(3-ethynyl-phenyl)amino]-6-(1-methyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0228] 4-[(3-ethynyl-phenyl)amino]-6-(1-methanesulphonyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0229] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-methyl-piperidin-4-yloxy)-7-(2-methoxy-ethoxy)-quinazoline,
- [0230] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-isopropylloxycarbonyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0231] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(cis-4-methylamino-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0232] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[cis-4-[N-(2-methoxy-acetyl)-N-methyl-amino]-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0233] 4-[(3-ethynyl-phenyl)amino]-6-(piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0234] 4-[(3-ethynyl-phenyl)amino]-6-[1-(2-methoxy-acetyl)-piperidin-4-yloxy]-7-methoxy-quinazoline,
- [0235] 4-[(3-ethynyl-phenyl)amino]-6-[1-[(morpholin-4-yl)carbonyl]-piperidin-4-yloxy]-7-methoxy-quinazoline,
- [0236] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[1-[(cis-2,6-dimethyl-morpholin-4-yl)carbonyl]-piperidin-4-yloxy]-7-methoxy-quinazoline,

- [0237] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-{1-[(2-methyl-morpholin-4-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-quinazoline,
- [0238] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-{1-[(S,S)-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-quinazoline,
- [0239] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-{1-[(N-methyl-N-2-methoxyethyl-amino)carbonyl]-piperidin-4-yloxy}-7-methoxy-quinazoline,
- [0240] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-ethyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0241] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-{1-[(2-methoxyethyl)carbonyl]-piperidin-4-yloxy}-7-methoxy-quinazoline,
- [0242] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-{1-[(3-methoxypropyl-amino)-carbonyl]-piperidin-4-yloxy}-7-methoxy-quinazoline,
- [0243] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[cis-4-(N-methanesulphonyl-N-methyl-amino)-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0244] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[cis-4-(N-acetyl-N-methyl-amino)-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0245] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(trans-4-methylamino-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0246] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[trans-4-(N-methanesulphonyl-N-methyl-amino)-cyclohexan-1-yloxy]-7-methoxy-quinazoline,
- [0247] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(trans-4-dimethylamino-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0248] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(trans-4-{N-[(morpholin-4-yl)carbonyl]-N-methyl-amino}-cyclohexan-1-yloxy)-7-methoxy-quinazoline,
- [0249] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]-quinazoline,
- [0250] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-methanesulphonyl-piperidin-4-yloxy)-7-methoxy-quinazoline,
- [0251] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-(1-cyanopiperidin-4-yloxy)-7-methoxy-quinazoline,
- [0252] 3-Cyano-4-[(3-chloro-4-fluorophenyl)amino]-6-{4-[(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino}-7-ethoxy-quinoline;
- [0253] 4-[(3-chloro-4-fluoro-phenyl)amino]-6-{[4-(homomorpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-[(S)-(tetrahydrofuran-3-yl)oxy]-quinazoline,
- [0254] 4-[(3-chloro-4-fluoro-phenyl)amino]-7-2-{4-[(S)-(2-oxo-tetrahydrofuran-5-yl)-carbonyl]-piperazin-1-yl}-ethoxy-6-[(vinylcarbonyl)amino]-quinazoline,
- [0255] 4-[(3-chloro-4-fluoro-phenyl)amino]-7-2-[(S)-6-methyl-2-oxo-morpholin-4-yl)-ethoxy]-6-[(vinylcarbonyl)amino]-quinazoline,
- [0256] 4-[(3-chloro-4-fluoro-phenyl)amino]-7-[4-((R)-6-methyl-2-oxo-morpholin-4-yl)-butyloxy]-6-[(vinylcarbonyl)amino]-quinazoline,
- [0257] 4-[(3-chloro-4-fluoro-phenyl)amino]-7-[4-((S)-6-methyl-2-oxo-morpholin-4-yl)-butyloxy]-6-[(vinylcarbonyl)amino]-quinazoline, and
- [0258] 4-[(3-chloro-4-fluorophenyl)amino]-6-[(4-{N-[2-(ethoxycarbonyl)-ethyl]-N-[(ethoxycarbonyl)methyl]amino}-1-oxo-2-buten-1-yl)amino]-7-cyclopropyl-methoxy-quinazoline,

optionally in the form of the racemates, enantiomers or diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts thereof, the solvates and/or hydrates thereof. Preferably, according to the invention, acid addition salts are selected from among the hydrochloride, hydrobromide, hydriodide, hydrosulphate, hydrophosphate, hydromethanesulphonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulphonate.

[0259] The LTD4-receptor antagonists used are preferably compounds selected from among montelukast, pranlukast, zafirlukast, and (E)-8-[2-[4-[4-(4-fluorophenyl)butoxy]phenyl]ethenyl]-2-(1H-tetrazol-5-yl)-4H-1-benzopyran-4-one (MEN-91507)

[0260] 4-[6-Acetyl-3-[3-(4-acetyl-3-hydroxy-2-propylphenylthio)propoxy]-2-propylphenoxy]-butyric acid (MN-001)

[0261] 1-(((R)-3-(2-(6,7-difluoro-2-quinolinyl)ethenyl)phenyl)-3-(2-(2-hydroxy-2-propyl)phenyl)thio)methylcyclopropane-acetic acid,

[0262] 1-(((1(R)-3(3-(2-(2,3-dichlorothieno[3,2-b]pyridin-5-yl)-(E)-ethenyl)phenyl)-3-(2-(1-hydroxy-1-methyl-ethyl)phenyl)propyl)thio)methyl)cyclopropaneacetic acid

[0263] [2-[[2-(4-tert-butyl-2-thiazolyl)-5-benzofuranyl]oxymethyl]phenyl]acetic acid

optionally in the form of the racemates, enantiomers or diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts thereof, the solvates and/or hydrates thereof. Preferably, according to the invention, acid addition salts are selected from among the hydrochloride, hydrobromide, hydriodide, hydrosulphate, hydrophosphate, hydromethanesulphonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulphonate. By salts or derivatives which the LTD4-receptor antagonists may optionally be capable of forming are meant, for example: alkali metal salts, such as for example sodium or potassium salts, alkaline earth metal salts, sulphobenzoates, phosphates, isonicotinates, acetates, propionates, dihydrogen phosphates, palmitates, pivalates or furoates.

[0264] The histamine H1 receptor antagonists used are preferably compounds selected from among epinastine, cetirizine, azelastine, fexofenadine, levocabastine, loratadine, mizolastine, ketotifen, emedastine, dimetindene, clemastine, bamipine, cexchlorpheniramine, pheniramine, doxylamine, chlorophenoxamine, dimenhydrinate, diphenhydramine, promethazine, ebastine, olopatadine, desloratadine and meclozine, optionally in the form of the racemates, enantiomers, diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts, solvates or hydrates thereof. According to the invention these acid addition salts are preferably selected from among the hydrochloride, hydrobromide, hydriodide, hydrosulphate, hydrophosphate, hydromethanesulphonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulphonate.

[0265] The PAF antagonists used are preferably compounds selected from among lexipafant and 4-(2-chlorophenyl)-9-methyl-2-[3(4-morpholinyl)-3-propanon-1-yl]-6H-thieno-[3,2,4]-[1,2,4]triazolo[4,3-a][1,4]diazepine



**[0266]** 6-(2-chlorophenyl)-8,9-dihydro-1-methyl-8-[(4-morpholinyl)carbonyl]-4H,7H-cyclo-penta-[4,5]thieno-[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine,

optionally in the form of the racemates, enantiomers, diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts, solvates or hydrates thereof. According to the invention these acid addition salts are preferably selected from among the hydrochloride, hydrobromide, hydriodide, hydrosulphate, hydrophosphate, hydromethanesulphonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulphonate.

**[0267]** The dopamine receptor agonists used are preferably compounds selected from among bromocriptin, cabergoline, alpha-dihydroergocryptine, lisuride, pergolide, pramipexol, roxindol, ropinirol, talipexol, tergurid and viozan, optionally in the form of the racemates, enantiomers, diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts, solvates or hydrates thereof. According to the invention these acid addition salts are preferably selected from among the hydrochloride, hydrobromide, hydriodide, hydrosulphate, hydrophosphate, hydromethanesulphonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydroxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulphonate.

**[0268]** Substances of preferred PI3 kinase antagonists that may be used are preferably compounds selected from among

**[0269]** 5-(Quinoxalin-6-ylmethylene)thiazolidine-2,4-dione (AS-605240),

**[0270]** 2-[(6-amino-9H-purin-9-yl)methyl]-5-methyl-3-(2-methylphenyl)-4(3H)-quinazolinone (C-87114),

**[0271]** 2-Methyl-2-[4-[3-methyl-2-oxo-8-(quinoline-3-yl)-2,3-dihydroimidazo[4,5-c]quinolin-1-yl]phenyl]propionitrile (BEZ-235),

optionally in the form of the racemates, enantiomers, diastereomers thereof and optionally in the form of the pharmacologically acceptable acid addition salts, prodrugs, solvates or hydrates thereof.

#### Terms and Definitions Used

**[0272]** By the term “optionally substituted” is meant within the scope of the invention the above-mentioned group, optionally substituted by a lower-molecular group. Examples of lower-molecular groups regarded as chemically meaningful are groups consisting of 1-25 atoms. Preferably such groups have no negative effect on the pharmacological efficacy of the compounds. For example the groups may comprise:

**[0273]** Straight-chain or branched carbon chains, optionally interrupted by heteroatoms, optionally substituted by rings, heteroatoms or other common functional groups.

**[0274]** Aromatic or non-aromatic ring systems consisting of carbon atoms and optionally heteroatoms, which may in turn be substituted by functional groups.

**[0275]** A number of aromatic or non-aromatic ring systems consisting of carbon atoms and optionally heteroatoms which may be linked by one or more carbon chains, optionally interrupted by heteroatoms, optionally substituted by heteroatoms or other common functional groups.

**[0276]** Also included in the subject-matter of this invention are the compounds according to the invention, including the salts thereof, wherein one or more hydrogen atoms, for example one, two, three, four or five hydrogen atoms, are replaced by deuterium.

**[0277]** Where a hyphen open on one side “-” is used in the structural formula of a substituent, this hyphen is to be understood as the linkage point to the remainder of the molecule. The substituent replaces the corresponding groups  $R^a$ ,  $R^b$ , etc. If no hyphen open on one side is used in the structural formula of a substituent, the linkage point to the remainder of the molecule is clear from the structural formula itself.

**[0278]** Compounds of general formula (I) may contain acid groups, primarily carboxyl groups, and/or basic groups such as e.g. amino functions. Compounds of general formula (I) may therefore be present as internal salts, as salts with pharmaceutically useable inorganic acids such as hydrochloric acid, sulphuric acid, phosphoric acid, sulphonic acid or organic acids (such as for example maleic acid, fumaric acid, citric acid, tartaric acid or acetic acid) or as salts with pharmaceutically useable bases such as alkali metal or alkaline earth metal hydroxides or carbonates, zinc or ammonium hydroxides or organic amines such as e.g. diethylamine, triethylamine, triethanolamine, inter alia. For preparing the alkali metal and alkaline earth metal salts of the compound of formula (I), it is preferable to use the alkali metal and alkaline earth metal hydroxides and hydrides, while the hydroxides and hydrides of the alkali metals, particularly sodium and potassium are preferred, and sodium and potassium hydroxide are particularly preferred. (See also *Pharmaceutical Salts*, S. M. Birge et al., *J. Pharm. Sci.* (1977), 66, 1-19)

**[0279]** As already mentioned, the compounds of general formula (I) may be converted into the salts thereof, particularly for pharmaceutical use, into the pharmacologically acceptable acid addition salts thereof with an inorganic or organic acid. Suitable acids for this purpose include for example succinic acid, hydrobromic acid, acetic acid, fumaric acid, maleic acid, methanesulphonic acid, lactic acid, phosphoric acid, hydrochloric acid, sulphuric acid, tartaric acid or citric acid. In addition, mixtures of the above-mentioned acids may be used.

**[0280]** The present invention relates to the respective compounds, optionally in the form of the individual diastereomers, mixtures of the individual diastereomers and/or individual enantiomers, mixtures of the individual enantiomers or racemates thereof, in the form of the tautomers as well as in the form of the free bases or the corresponding acid addition salts with pharmacologically acceptable acids—such as for example acid addition salts with hydrohalic acids—for example hydrochloric or hydrobromic acid or organic acids—such as for example tartaric acid, fumaric acid, citric acid or methanesulphonic acid.

**[0281]** “Protective groups” for the purposes of the present invention is a collective term for organic groups with which certain functional groups of a molecule containing a number of active centres can temporarily be protected from attack by reagents so that reactions take place only at the desired (unprotected) sites. The protective groups should be introduced selectively under mild conditions. They must be stable for the duration of the protection under all the conditions of the reactions and purifying procedures which are to be carried out; racemisations and epimerisations must be suppressed. Protective groups should be capable of being cleaved again under mild conditions selectively and ideally in high yields.

The choice of a suitable protective group, the reaction conditions (solvent, temperature, duration, etc.), and also the options for removing a protective group are known in the art (e.g. Philip Kocienski, *Protecting Groups*, 3rd ed. 2004, THIEME, Stuttgart, ISBN: 3131370033).

**[0282]** By an “organic solvent” is meant, within the scope of the invention, an organic, low-molecular substance which can dissolve other organic substances by a physical method. To be suitable the prerequisite for the solvent is that neither the dissolving substance nor the dissolved substance should be chemically altered during the dissolving process, i.e. the components of the solution should be recoverable in their original form by physical separation processes such as distillation, crystallisation, sublimation, evaporation or adsorption. For various reasons, not only the pure solvents but also mixtures that combine the dissolving properties may be used. Examples include:

- [0283]** alcohols, preferably methanol, ethanol, propanol, butanol, octanol, cyclohexanol;
- [0284]** glycols, preferably ethyleneglycol, diethyleneglycol;
- [0285]** ethers/glycolethers, preferably diethyl ether, tert-butyl-methylether, dibutylether, anisol, dioxane, tetrahydrofuran, mono-, di-, tri-, polyethyleneglycol ethers;
- [0286]** ketones, preferably acetone, butanone, cyclohexanone;
- [0287]** esters, preferably acetic acid esters, glycolesters;
- [0288]** amides and other nitrogen compounds, preferably dimethylformamide, pyridine, N-methylpyrrolidone, acetonitrile;
- [0289]** sulphur compounds, preferably carbon disulphide, dimethylsulphoxide, sulpholane;
- [0290]** nitro compounds, preferably nitrobenzene;
- [0291]** halogenated hydrocarbons, preferably dichloromethane, chloroform, tetrachloromethane, tri- and tetrachloroethane, 1,2-dichloroethane, chlorofluorocarbons;
- [0292]** aliphatic or alicyclic hydrocarbons, preferably benzines, petroleum ether, cyclohexane, methylcyclohexane, decaline, terpene-L.; or
- [0293]** aromatic hydrocarbons, preferably benzene, toluene, o-xylene, m-xylene, p-xylene;

or corresponding mixtures thereof.

**[0294]** The term diastereomerically pure describes within the scope of the present invention compounds of formula (I), which are present in a diastereomeric purity of at least 85% de, preferably at least 90% de, particularly preferably >95% de. The term de (diastereomeric excess) is known in the art and describes the optical purity of diastereomeric compounds.

**[0295]** The term enantiomerically pure describes within the scope of the present invention compounds of formula (I), which are present in an enantiomeric purity of at least 85% ee, preferably at least 90% ee, particularly preferably >95% ee. The term ee (enantiomeric excess) is known in the art and describes the optical purity of chiral compounds.

**[0296]** By the term “C<sub>1-6</sub>-alkyl” (including those which are part of other groups) are meant branched and unbranched alkyl groups with 1 to 6 carbon atoms and by the term “C<sub>1-4</sub>-alkyl” are meant branched and unbranched alkyl groups with 1 to 4 carbon atoms. Preferred are alkyl groups with 1 to 4 carbon atoms, particularly preferably alkyl groups with 1 to 2 carbon atoms. Examples include: methyl, ethyl, n-propyl,

iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, iso-pentyl, neo-pentyl or hexyl. The abbreviations Me, Et, n-Pr, i-Pr, n-Bu, i-Bu, t-Bu, etc. may optionally also be used for the above-mentioned groups. Unless stated otherwise, the definitions propyl, butyl, pentyl and hexyl include all the possible isomeric forms of the groups in question. Thus, for example, propyl includes n-propyl and iso-propyl, butyl includes iso-butyl, sec-butyl and tert-butyl etc.

**[0297]** By the term “C<sub>1-3</sub>-alkylene” (including those which are part of other groups) are meant branched and unbranched alkylene groups with 1 to 3 carbon atoms. Preferred are alkylene groups with 1 to 2 carbon atoms. Examples include: methylene, ethylene, propylene and 1-methylethylene. Unless stated otherwise, the definition propylene includes all the possible isomeric forms of the groups in question with the same number of carbons. Thus, for example, propylene also includes 1-methylethylene.

**[0298]** By the term “C<sub>2-3</sub>-alkenyl” (including those which are part of other groups) are meant alkenyl groups with 2 to 3 carbon atoms, if they have at least one double bond. Examples include: ethenyl or vinyl. Unless stated otherwise, the definition “C<sub>2-3</sub>-alkenyl” includes all the possible isomeric forms of the groups in question with the same number of carbons. Thus, for example, propenyl includes 1-propen-1-yl, 1-propen-2-yl and 2-propen-1-yl.

**[0299]** By the term “C<sub>2-3</sub>-alkynyl” (including those which are part of other groups) are meant alkynyl groups with 2 to 3 carbon atoms, if they have at least one triple bond. Examples include: ethynyl or propynyl. Unless stated otherwise, the definition “C<sub>2-3</sub>-alkynyl” includes all the possible isomeric forms of the groups in question with the same number of carbons. Thus, for example, propynyl includes propyn-1-yl and 2-propyn-1-yl.

**[0300]** By the term “C<sub>3-7</sub>-cycloalkyl” (including those which are part of other groups) are meant cyclic alkyl groups with 3 to 7 carbon atoms. Examples include: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl. Unless otherwise stated, the cyclic alkyl groups may be substituted by one or more groups selected from among methyl, ethyl, iso-propyl, tert-butyl, hydroxy and fluorine.


**[0301]** By the term “aryl” (including those which are part of other groups) are meant aromatic ring systems with 6, 10 or 14 carbon atoms. Examples include: phenyl, naphthyl, anthracenyl or phenanthrenyl, the preferred aryl group being phenyl. Unless otherwise stated, the aromatic groups may be substituted by one or more groups R<sup>5</sup>. Particularly preferably the term “aryl” in each case denotes a phenyl group which is mono- or disubstituted by R<sup>5</sup>, wherein the substituents R<sup>5</sup> may be identical or different and

**[0302]** R<sup>5</sup> denotes hydrogen, or

**[0303]** a group, which may be identical or different, selected from among F, Cl, Br, I, OH, CN, C<sub>1-3</sub>-alkyl, C<sub>1-3</sub>-alkyl-O, CHF<sub>2</sub>, CF<sub>3</sub>, —O—CHF<sub>2</sub> and —O—CF<sub>3</sub>.

**[0304]** By the term “heteroaryl” are meant 5-10-membered mono- or bicyclic heteroaryl rings, wherein up to three C atoms may be replaced by one or more heteroatoms selected from among oxygen, nitrogen or sulphur, these rings containing sufficient conjugated double bonds to form an aromatic system. Each of the above-mentioned heterocycles may optionally also be fused to a benzene ring. The heteroaryl rings may, unless otherwise described, carry one or more substituents, for example. The ring may be linked to the

molecule via a carbon atom or, if available, via a nitrogen atom. The following are examples of five- or six-membered heterocyclic aromatic groups:



**[0305]** Examples of 5-10-membered bicyclic heteroaryl rings include pyrrolizine, indole, indolizine, isoindole, indazole, purine, quinoline, isoquinoline, quinoxaline, benzimidazole, benzofuran, benzothiophene, benzothiazole, benzisothiazole, pyridopyrimidine, pteridine, pyrimidopyrimidine.

**[0306]** Particularly preferably, the term “heteroaryl” denotes a pyridinyl, pyridazinyl, pyrimidinyl or pyrazinyl group, which is mono- or disubstituted in each case by the group  $R^5$ , wherein the substituents  $R^5$  may be identical or different and  $R^5$  is as hereinbefore defined.

**[0307]** “Halogen” within the scope of the present invention denotes fluorine, chlorine, bromine or iodine. Unless stated to the contrary, fluorine, chlorine and bromine are regarded as preferred halogens.

**[0308]** The substituent  $R^a$  may represent a phenyl or 1-phenylethyl group, preferably a phenyl group, wherein the phenyl nucleus is substituted in each case by the groups  $R^1$  to  $R^3$ .

**[0309]** Particularly preferably the substituent  $R^a$  denotes a group, selected from among 3-chloro-2-fluoro-phenyl, 3-chloro-4-fluoro-phenyl, 5-chloro-2-fluoro-phenyl, 2-fluoro-3-methyl-phenyl, 2-fluoro-5-methyl-phenyl, 4-fluoro-3-methyl-phenyl- and 3-chloro-2-methyl-phenyl. Most particularly preferably the substituent  $R^a$  denotes a 3-chloro-2-fluoro-phenyl group.

**[0310]** The substituent  $R^b$  may represent hydrogen, or an optionally substituted group selected from among  $C_{1-6}$ -alkyl,  $C_{3-6}$ -cycloalkyl- and  $C_{3-6}$ -cycloalkyl- $C_{1-3}$ -alkyl, preferably hydrogen and  $C_{1-3}$ -alkyl, particularly preferably hydrogen and methyl.

**[0311]** The substituent  $R^c$  may represent hydrogen, or an optionally substituted group selected from among  $C_{1-6}$ -alkyl,  $C_{3-6}$ -cycloalkyl- and  $C_{3-6}$ -cycloalkyl- $C_{1-3}$ -alkyl,  $C_{1-6}$ -alkyl-CO,  $C_{3-6}$ -cycloalkyl-CO— and  $C_{3-6}$ -cycloalkyl- $C_{1-3}$ -alkyl-CO,  $C_{1-6}$ -alkyl-SO<sub>2</sub>,  $C_{3-6}$ -cycloalkyl-SO<sub>2</sub>— and  $C_{3-6}$ -cycloalkyl- $C_{1-3}$ -alkyl-SO<sub>2</sub>, phenyl-CO— and phenyl-SO<sub>2</sub>, preferably hydrogen and  $C_{1-3}$ -alkyl, particularly preferably hydrogen and methyl.

**[0312]** The substituent  $R^d$  may denote hydrogen or

**[0313]** a group selected from among

**[0314]** F, Cl, Br, I, OH,  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkyl-O,  $C_{1-2}$ -alkyl-O substituted by 1 to 3 fluorine atoms,  $C_{3-7}$ -cycloalkyl-O,  $C_{3-7}$ -cycloalkyl- $C_{1-4}$ -alkyl-O, tetrahydrofuran-3-yl-O, tetrahydropyran-3-yl-O, tetrahydro-pyran-

4-yl-O, tetrahydrofuran-yl- $C_{1-4}$ -alkyl-O— and tetrahydropyran-yl- $C_{1-4}$ -alkyl-O,

or

**[0315]**  $R^4$ - $C_{1-4}$ -alkyl, wherein the linking of the groups  $R^4$  may take place via each C atom of the alkyl group,

or

**[0316]**  $R^4$ - $C_{2-4}$ -alkyl-O, wherein the group  $R^4$  is separated from the oxygen atom by at least 2 C atoms,

or

**[0317]** a group selected from among

**[0318]** pyrrolidin-2-yl- $C_{1-4}$ -alkyl-O, pyrrolidin-3-yl- $C_{1-4}$ -alkyl-O, piperidin-2-yl- $C_{1-4}$ -alkyl-O, piperidin-3-yl- $C_{1-4}$ -alkyl-O, piperidin-4-yl- $C_{1-4}$ -alkyl-O, azepan-2-yl- $C_{1-4}$ -alkyl-O, azepan-3-yl- $C_{1-4}$ -alkyl-O, azepan-4-yl- $C_{1-4}$ -alkyl-O, morpholin-2-yl- $C_{1-4}$ -alkyl-O, morpholin-3-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-pyrrolidin-2-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-pyrrolidin-3-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-piperidin-2-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-piperidin-3-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-piperidin-4-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-azepan-2-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-azepan-3-yl- $C_{1-4}$ -alkyl-O, 1-( $C_{1-3}$ -alkyl)-azepan-4-yl- $C_{1-4}$ -alkyl-O, 4-( $C_{1-3}$ -alkyl)-morpholin-2-yl- $C_{1-4}$ -alkyl-O and 4-( $C_{1-3}$ -alkyl)-morpholin-3-yl- $C_{1-4}$ -alkyl-O—,

preferably  $C_{1-3}$ -alkyl-O—, particularly preferably  $CH_3$ -O—,

wherein the pyrrolidinyl, piperidinyl, azepan-1-yl, piperazinyl, 1,4-diazepan-1-yl, morpholinyl and 1,4-oxazepan-4-yl groups mentioned above in the definition of the group  $R^d$  may each additionally be substituted by one or two  $C_{1-3}$ -alkyl groups,

and

wherein the above-mentioned phenyl groups are mono- or disubstituted by groups  $R^5$ .

**[0319]** The substituent  $R^1$  may denote hydrogen or

**[0320]** a group selected from among

**[0321]** F, Cl, Br, I,  $OCH_2F$ ,  $OCHF_2$ ,  $OCF_3$ ,  $CH_2F$ ,  $CHF_2$ ,  $CF_3$ , CN,  $NO_2$ ,  $NH_2$  and OH, or

**[0322]** a group selected from among

**[0323]**  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkyl-O,  $C_{2-3}$ -alkenyl,  $C_{2-3}$ -alkynyl,

**[0324]** phenyl, phenyl-O, phenyl- $C_{1-3}$ -alkyl- and phenyl- $C_{1-3}$ -alkyl-O, heteroaryl, heteroaryl-O, heteroaryl- $C_{1-3}$ -alkyl and heteroaryl- $C_{1-3}$ -alkyl-O,

**[0325]** wherein the above-mentioned phenyl groups are mono- or disubstituted by groups  $R^5$ ,

preferably hydrogen, fluorine, chlorine, bromine or methyl, particularly preferably hydrogen, fluorine, chlorine or methyl.

**[0326]** The substituent  $R^2$  may represent hydrogen or

**[0327]** a group selected from among

**[0328]** F, Cl, Br, I,  $OCH_2F$ ,  $OCHF_2$ ,  $OCF_3$ ,  $CH_2F$ ,  $CHF_2$ ,  $CF_3$ , CN,  $NO_2$ ,  $NH_2$  and OH, or

**[0329]** a group selected from among

**[0330]**  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkyl-O,  $C_{2-3}$ -alkenyl,  $C_{2-3}$ -alkynyl, phenyl, phenyl-O, phenyl- $C_{1-3}$ -alkyl and phenyl- $C_{1-3}$ -alkyl-O, heteroaryl, heteroaryl-O, heteroaryl- $C_{1-3}$ -alkyl and heteroaryl- $C_{1-3}$ -alkyl-O, wherein the above-mentioned phenyl groups are mono- or disubstituted by groups  $R^5$ ,

preferably hydrogen, fluorine, chlorine or methyl, particularly preferably hydrogen, fluorine or chlorine.

[0331] The substituent  $R^3$  may represent hydrogen, or

[0332] a group selected from among F, Cl, Br and  $CH_3$ , preferably hydrogen.

[0333] The substituent  $R^4$  may represent a group, which may be identical or different, selected from among

[0334] OH,  $C_{1-3}$ -alkyl-O,  $C_{3-6}$ -cycloalkyl-O,  $NH_2$ ,  $C_{1-3}$ -alkyl-NH,  $(C_{1-3}$ -alkyl) $_2N$ , (2-methoxyethyl) $_2N$ , pyrrolidin-1-yl, piperidin-1-yl, azepan-1-yl, morpholin-4-yl, 1,4-oxazepan-4-yl, 2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl, 3-oxa-8-aza-bicyclo[3.2.1]oct-8-yl, 8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl, piperazin-1-yl, 4-( $C_{1-3}$ -alkyl)-piperazin-1-yl, 1,4-diazepan-1-yl, 4-( $C_{1-3}$ -alkyl)-1,4-diazepan-1-yl,  $HCO-NH$ ,  $C_{1-4}$ -alkyl-CO-NH,  $C_{1-3}$ -alkyl-O- $C_{1-3}$ -alkyl-CO-NH,  $C_{1-4}$ -alkyl-O-CO-NH,  $H_2NCONH$ ,  $C_{1-3}$ -alkyl-NH-CO-NH,  $(C_{1-3}$ -alkyl) $_2N-CONH$ , pyrrolidin-1-yl-CO-NH, piperidin-1-yl-CO-NH, piperazin-1-yl-CO-NH, 4-( $C_{1-3}$ -alkyl)-piperazin-1-yl-CO-NH and morpholin-4-yl-CO-NH and  $C_{1-4}$ -alkyl-SO $_2$ -NH—.

[0335] The substituent  $R^5$  may represent hydrogen, or

[0336] a group, which may be identical or different, selected from among

[0337] F, Cl, Br, I, OH, CN,  $C_{1-3}$ -alkyl,  $C_{1-3}$ -alkyl-O,  $CHF_2$ ,  $CF_3$ ,  $-O-CHF_2$  and  $-O-CF_3$ .

[0338] A may denote  $-CO-$  or  $-C_1-C_3$ -alkylene, preferably  $-CH_2CH_2-$ ,


[0339] wherein the  $-C_1-C_3$ -alkylene group may be 1-, 2-, 3- or 4-, preferably 1- or 2-substituted by a group  $R^6$ ,

[0340] The substituent  $R^6$ , which may be identical or different, may represent hydrogen, or a group selected from among OH,  $C_1-C_4$ -alkyl and  $-O-C_1-C_4$ -alkyl, preferably methyl. A particularly preferred definition of A is  $-CH_2CH_2-$ .

#### Methods of Preparation


[0341] The following methods are suitable, for example, for preparing compounds of general formula (I):

a) reacting a compound of general formula



wherein

$R^a$  and  $R^d$  are as hereinbefore defined, with a compound of general formula




wherein

$R^b$ ,  $R^c$  and A are as hereinbefore defined and  $Z^1$  denotes a leaving group such as a halogen atom, e.g. a chlorine or bromine atom, a sulphonyloxy group such as a methane-sulphonyloxy or p-toluenesulphonyloxy group or a hydroxy group.

[0342] With a compound of general formula (III), wherein  $Z^1$  denotes a halogen atom or a sulphonyloxy group, the reaction is expediently carried out in a solvent such as ethanol, isopropanol, acetonitrile, toluene, tetrahydrofuran, dioxane, dimethylformamide, dimethylsulphoxide or N-methylpyrrolidinone, preferably in the presence of a base such as potassium carbonate, potassium-tert-butoxide, sodium hydride or N-ethyl-diisopropylamine, at temperatures in the range from 20° C. to 160° C., for example at temperatures in the range from 60° C. to 140° C.

[0343] With a compound of general formula III wherein  $Z^1$  denotes a hydroxy group, the reaction is carried out in the presence of a dehydrating agent, preferably in the presence of a phosphine and an azodicarboxylic acid derivative such as e.g. triphenylphosphine/diethyl azodicarboxylate, conveniently in a solvent such as methylene chloride, acetonitrile, tetrahydrofuran, dioxane, toluene or ethyleneglycol diethylether at temperatures between  $-50$  and  $150^\circ\text{C}$ ., but preferably at temperatures between  $-20$  and  $80^\circ\text{C}$ .

[0344] b) In order to prepare compounds of general formula I wherein  $R^b$  and  $R^c$  each denote a hydrogen atom and A represents a  $-CO-$  group, reacting a compound of general formula




wherein

$R^a$  and  $R^d$  are as hereinbefore defined, with an alkali metal cyanide and ammonium carbonate.

[0345] The reaction is carried out for example in a solvent or mixture of solvents such as methanol, ethanol, ethanol/water or isopropanol at temperatures between ambient temperature and  $120^\circ\text{C}$ . Further references to the synthesis of hydantoins can be found for example in the following publication:


[0346] Meusel, M.; Guetschow, M., Organic Preparations and Procedures International (2004), 36(5), 391-443.

c) reacting a compound of general formula (V)



wherein  $R^b$ ,  $R^c$ ,  $R^d$  and A are as hereinbefore defined, with a halogenating agent, for example an acid halide such as thionyl chloride, thionyl bromide, phosphorus trichloride, phosphorus pentachloride, phosphorus oxychloride, or triph-

enylphosphine/carbon tetrachloride or triphenylphosphine/N-chlorosuccinimide) to obtain an intermediate compound of general formula (VI),



wherein  $R^b$ ,  $R^c$ ,  $R^d$  and A are as hereinbefore defined and  $Z^2$  denotes a halogen atom such as a chlorine or bromine atom, and subsequent reaction with a compound of general formula (VII),



wherein  $R^a$  is as hereinbefore defined, or the salts thereof.


**[0347]** The reaction with the halogenating agent is optionally carried out in a solvent such as methylene chloride, chloroform, acetonitrile or toluene and optionally in the presence of a base such as N,N-diethylaniline, triethylamine or N-ethyl-diisopropylamine at temperatures in the range from 20° C. to 160° C., preferably from 40° C. to 120° C. Preferably, however, the reaction is carried out with thionyl chloride and catalytic amounts of dimethylformamide at the boiling temperature of the reaction mixture or with phosphorus oxychloride in acetonitrile in the presence of triethylamine at the boiling temperature of the reaction mixture or with triphenylphosphine/carbon tetrachloride or with triphenylphosphine/N-chlorosuccinimide in acetonitrile.

**[0348]** The reaction of the compound of general formula (VI) with the compound of general formula (VII) or the salts thereof is conveniently carried out in a solvent such as ethanol, isopropanol, acetonitrile, dioxane or dimethylformamide, optionally in the presence of a base such as potassium carbonate, triethylamine or N-ethyl-diisopropylamine, at temperatures in the range from 20° C. and 160° C., preferably from 60° C. to 120° C. However, the reaction is preferably carried out in isopropanol at the boiling temperature of the reaction mixture.

**[0349]** The reaction of a compound of general formula (V) to obtain a compound of general formula (I) may also be carried out as a one-pot reaction, for example in acetonitrile in the presence of triethylamine.

d) In order to prepare compounds of general formula I wherein  $R^d$  denotes one of the optionally substituted alkyloxy groups mentioned hereinbefore:

reacting a compound of general formula



wherein  $R^a$ ,  $R^b$ ,  $R^c$  and A are as hereinbefore defined, with a compound of general formula



wherein  $R^d$  denotes a group selected from among  $C_{1-4}$ -alkyl,  $C_{1-2}$ -alkyl substituted by 1 to 3 fluorine atoms,  $C_{3-7}$ -cycloalkyl,  $C_{3-7}$ -cycloalkyl- $C_{1-4}$ -alkyl, tetrahydrofuran-3-yl, tetrahydropyran-3-yl, tetrahydro-pyran-4-yl, tetrahydrofuran-yl- $C_{1-4}$ -alkyl and tetrahydropyranyl- $C_{1-4}$ -alkyl,

or

**[0350]**  $R^4$ - $C_{2-4}$ -alkyl, wherein the group  $R^4$  is separated from  $Z^3$  by at least 2 C atoms,

or

**[0351]** a group selected from among


**[0352]** pyrrolidin-2-yl- $C_{1-4}$ -alkyl, pyrrolidin-3-yl- $C_{1-4}$ -alkyl, piperidin-2-yl- $C_{1-4}$ -alkyl, piperidin-3-yl- $C_{1-4}$ -alkyl, piperidin-4-yl- $C_{1-4}$ -alkyl, azepan-2-yl- $C_{1-4}$ -alkyl, azepan-3-yl- $C_{1-4}$ -alkyl, azepan-4-yl- $C_{1-4}$ -alkyl, morpholin-2-yl- $C_{1-4}$ -alkyl, morpholin-3-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-pyrrolidin-2-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-pyrrolidin-3-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-piperidin-2-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-piperidin-3-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-piperidin-4-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-azepan-2-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-azepan-3-yl- $C_{1-4}$ -alkyl, 1-( $C_{1-3}$ -alkyl)-azepan-4-yl- $C_{1-4}$ -alkyl, 4-( $C_{1-3}$ -alkyl)-morpholin-2-yl- $C_{1-4}$ -alkyl, 4-( $C_{1-3}$ -alkyl)-morpholin-3-yl- $C_{1-4}$ -alkyl,

and

$Z^3$  denotes a leaving group such as a halogen atom, an alkylsulphonyloxy, arylsulphonyloxy or a hydroxy group.

**[0353]** If the leaving group is a halogen atom such as a chlorine, bromine or iodine atom or an alkylsulphonyloxy or arylsulphonyloxy group such as the methanesulphonyloxy or p-toluenesulphonyloxy group, the reaction is preferably carried out in the presence of an organic or inorganic base such as potassium carbonate, sodium hydride or N-ethyl-diisopropylamine. If the leaving group is a hydroxy group, the reaction is carried out in the presence of a dehydrating agent, preferably in the presence of a phosphine and an azodicarboxylic acid derivative such as e.g. triphenylphosphine/diethyl azodicarboxylate.

e) In order to prepare compounds of general formula I wherein  $R^d$  denotes a  $R^{4'}$ - $C_{2-4}$ -alkyl-O- group, wherein the group  $R^{4'}$  is separated from the oxygen atom by at least 2 C atoms, and  $R^{4'}$  denotes a group selected from among  $NH_2$ ,  $C_{1-3}$ -alkyl-NH, ( $C_{1-3}$ -alkyl) $_2N$ , (2-methoxyethyl) $_2N$ , pyrrolidin-1-yl, piperidin-1-yl, azepan-1-yl, morpholin-4-yl, 1,4-oxazepan-4-yl, 2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl, 3-oxa-8-aza-bicyclo[3.2.1]oct-8-yl, 8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl, piperazin-1-yl, 4-( $C_{1-3}$ -alkyl)piperazin-1-yl, 1,4-diazepan-1-yl, 4-( $C_{1-3}$ -alkyl)-1,4-diazepan-1-yl: reacting a compound of general formula




wherein  $R^a$ ,  $R^b$ ,  $R^c$  and A are as hereinbefore defined and  $Z^4$  denotes a leaving group such as a halogen atom, e.g. a chlorine or bromine atom or a sulphonyloxy group such as a methanesulphonyloxy or p-toluenesulphonyloxy group, with



wherein  $R^{4'}$  is as hereinbefore defined.

f) In order to prepare compounds of general formula I wherein  $R^b$  denotes a hydrogen atom:

cleaving a protective group from a compound of general formula



wherein  $R^a$ ,  $R^c$ ,  $R^d$  and A are as hereinbefore defined and  $R^b$  denotes a protective group, for example an optionally substituted benzyl group, a tert.-butyl group or a 2-(trimethylsilyl)ethyl group.

**[0354]** An optionally substituted benzyl group is for example cleaved hydrogenolytically, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a suitable solvent such as methanol, ethanol, ethyl acetate or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid, at temperatures between 0 and 100° C., but preferably at ambient temperatures between 20 and 60° C., and under a hydrogen pressure of 1 to 7 bar, but preferably from 3 to 5 bar. However, a 2,4-dimethoxybenzyl group is preferably cleaved in trifluoroacetic acid in the presence of anisole, thioanisole, pentamethylbenzene or triethylsilane.


**[0355]** An optionally substituted benzyl group or a tert.-butyl group may for example also be cleaved by treating with an acid such as trifluoroacetic acid, hydrochloric acid or hydrobromic acid, optionally using a solvent such as methylene chloride or toluene, optionally in the presence of anisole, thioanisole, pentamethylbenzene or triethylsilane.

**[0356]** A 2-(trimethylsilyl)ethyl group is cleaved for example by treatment with fluorides such as tetrabutylammonium fluoride, optionally using a solvent such as tetrahydrofuran or dioxane.

**[0357]** Other suitable protective groups and possible ways of introducing and cleaving them are described for example in "Protective Groups in Organic Synthesis" by Theodora W. Greene and Peter G. M. Wuts, Wiley-VCH, or Philip Kocienski, Protecting Groups, 3rd ed. 2004, THIEME.

g) In order to prepare compounds of general formula I wherein  $R^c$  denotes a hydrogen atom:

cleaving a protective group from a compound of general formula



wherein  $R^a$ ,  $R^b$ ,  $R^d$  and A are as hereinbefore defined and  $R^c$  denotes a protective group, for example an optionally substituted benzyl group or a formyl, acetyl, trifluoroacetyl, methoxycarbonyl, ethoxycarbonyl, tert.-butoxycarbonyl or benzyloxycarbonyl group.

**[0358]** The protective group is cleaved, for example, hydrolytically in an aqueous solvent, e.g. in water, isopropanol/water, acetic acid/water, tetrahydrofuran/water or dioxane/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali metal base such as sodium hydroxide or potassium hydroxide or aprotically, e.g. in the presence of iodotrimethylsilane, at temperatures between 0 and 120° C., preferably at temperatures between 10 and 100° C.


**[0359]** An optionally substituted benzyl group, or a benzyloxycarbonylbenzyl group is cleaved, for example, hydrogenolytically, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a suitable solvent such as methanol, ethanol, ethyl acetate or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid at temperatures between 0 and 100° C., but preferably at ambient temperatures between 20 and 60° C., and under a hydrogen pressure of 1 to 7 bar, but preferably from 3 to 5 bar.

**[0360]** A tert.-butoxycarbonyl group is preferably cleaved by treatment with an acid such as trifluoroacetic acid or hydrochloric acid, optionally using a solvent such as methylene chloride, dioxane, methanol or diethyl ether.

**[0361]** A trifluoroacetyl group is preferably cleaved by treatment with an acid such as hydrochloric acid, optionally in the presence of a solvent such as acetic acid at temperatures between 50 and 120° C. or by treatment with sodium hydroxide solution, optionally in the presence of a solvent such as tetrahydrofuran at temperatures between 0 and 50° C.

**[0362]** Other suitable protective groups and possible ways of introducing and cleaving them are described for example in "Protective Groups in Organic Synthesis" by Theodora W. Greene and Peter G. M. Wuts, Wiley-VCH, or Philip Kocienski, Protecting Groups, 3rd ed. 2004, THIEME.

h) In order to prepare compounds of general formula I wherein A denotes a  $-C_2-C_3$ -alkylene group: cyclising a compound of general formula



wherein  $R^a$ ,  $R^b$ ,  $R^c$  and  $R^d$  are as hereinbefore defined, A' denotes a  $-C_2-C_3$ -alkylene group and  $Z^5$  denotes a leaving group such as a halogen atom, a hydroxy or alkyloxy group.

**[0363]** If the leaving group is a hydroxy group, the reaction is carried out in the presence of a dehydrating agent such as N,N'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium-tetrafluoroborate (TBTU) or O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium-hexafluorophosphate (HATU), conveniently in a solvent such as methylene chloride, N,N-

dimethylformamide, acetonitrile, tetrahydrofuran, dioxane or ethyleneglycol diethyl ether at temperatures between  $-50^{\circ}\text{C}$ . and  $100^{\circ}\text{C}$ ., but preferably at temperatures between  $-20^{\circ}\text{C}$ . and  $60^{\circ}\text{C}$ .

**[0364]** If the leaving group is a halogen atom, the reaction is preferably carried out in the presence of a base such as triethylamine or N-ethyl-diisopropylamine, conveniently in a solvent such as methylene chloride, N,N-dimethylformamide, acetonitrile, tetrahydrofuran, dioxane or ethyleneglycol diethyl ether at temperatures between  $-50^{\circ}\text{C}$ . and  $100^{\circ}\text{C}$ ., but preferably at temperatures between  $-20^{\circ}\text{C}$ . and  $60^{\circ}\text{C}$ .

**[0365]** If the leaving group is an alkoxy group, the reaction is optionally carried out in the presence of a base such as potassium carbonate, sodium hydroxide, triethylamine or N-ethyl-diisopropylamine, conveniently in a solvent such as methanol, ethanol, isopropanol, methylene chloride, N,N-dimethylformamide, acetonitrile, tetrahydrofuran, dioxane or ethyleneglycol diethyl ether at temperatures between  $-50^{\circ}\text{C}$ . and  $120^{\circ}\text{C}$ ., but preferably at temperatures between  $0^{\circ}\text{C}$ . and  $80^{\circ}\text{C}$ .

**[0366]** If according to the invention a compound of general formula I is obtained which contains an amino, alkylamino or imino group, this may be converted by acylation or sulphonylation into a corresponding acyl or sulphonyl compound of general formula I, wherein the acylating agents used may be for example carboxylic acid halides, carboxylic acid anhydrides and carboxylic acids with activating agents such as N,N'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide or O-(benzotriazol-1-yl)-N,N,N'-tetramethyluronium-tetrafluoroborate and sulphonyl halides as sulphonylating agents, and/or

**[0367]** if a compound of general formula I is obtained which contains an amino, alkylamino or imino group, it may be converted by alkylation or reductive alkylation into a corresponding alkyl compound of general formula I and/or

**[0368]** if a compound of general formula I is obtained which contains an alkoxycarbonyl group, it may be converted by ester cleaving into a carboxylic acid, and/or

**[0369]** if a compound of general formula I is obtained which contains an alkoxycarbonyl group, it may be converted by reaction with an amine into a carboxylic acid amide derivative and/or

**[0370]** if a compound of general formula I is obtained which contains a carboxy group, it may be converted by reaction with an amine into a carboxylic acid amide derivative.

**[0371]** In the reactions described hereinbefore any reactive groups present such as hydroxy, amino, alkylamino or imino groups may be protected during the reaction by conventional protective groups which are cleaved again after the reaction.

**[0372]** For example a protecting group for a hydroxy group might be the trimethylsilyl, acetyl, trityl, benzyl or tetrahydropyranyl group.

**[0373]** Protecting groups for an amino, alkylamino or imino group might be, for example, the formyl, acetyl, trifluoroacetyl, ethoxycarbonyl, tert.-butoxycarbonyl, benzyloxycarbonyl, benzyl, methoxybenzyl or 2,4-dimethoxybenzyl group.

**[0374]** Any protective group used is optionally subsequently cleaved for example by hydrolysis in an aqueous solvent, e.g. in water, isopropanol/water, acetic acid/water, tetrahydrofuran/water or dioxane/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali metal base such as

sodium hydroxide or potassium hydroxide or aprotically, e.g. in the presence of iodotrimethylsilane, at temperatures between  $0$  and  $120^{\circ}\text{C}$ ., preferably at temperatures between  $10$  and  $100^{\circ}\text{C}$ .

**[0375]** A benzyl, methoxybenzyl or benzyloxycarbonyl group, however, is cleaved by hydrogenolysis, for example, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a suitable solvent such as methanol, ethanol, ethyl acetate or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid at temperatures between  $0$  and  $100^{\circ}\text{C}$ ., but preferably at ambient temperatures between  $20$  and  $60^{\circ}\text{C}$ ., and under a hydrogen pressure of  $1$  to  $7$  bar, but preferably from  $3$  to  $5$  bar. A 2,4-dimethoxybenzyl group, however, is preferably cleaved in trifluoroacetic acid in the presence of anisole, thioanisole, pentamethylbenzene or triethylsilane.

**[0376]** A tert.-butyl or tert.-butoxycarbonyl group is preferably cleaved by treatment with an acid such as trifluoroacetic acid or hydrochloric acid or by treating with iodotrimethylsilane, optionally using a solvent such as methylene chloride, dioxane, methanol or diethyl ether.

**[0377]** A trifluoroacetyl group is preferably cleaved by treatment with an acid such as hydrochloric acid, optionally in the presence of a solvent such as acetic acid at temperatures between  $50$  and  $120^{\circ}\text{C}$ . or by treatment with sodium hydroxide solution, optionally in the presence of a solvent such as tetrahydrofuran at temperatures between  $0$  and  $50^{\circ}\text{C}$ .

**[0378]** Other suitable protective groups and possible ways of introducing and cleaving them are described for example in "Protective Groups in Organic Synthesis" by Theodora W. Greene and Peter G. M. Wuts, Wiley-VCH, or Philip Kocienski, Protecting Groups, 3rd ed. 2004, THIEME.

**[0379]** Moreover, the compounds of general formula I obtained may be resolved into their enantiomers and/or diastereomers, as mentioned hereinbefore. Thus, for example, cis/trans mixtures may be resolved into their cis and trans isomers, and compounds with at least one optically active carbon atom may be separated into their enantiomers.

**[0380]** Thus, for example, the cis/trans mixtures obtained may be resolved by chromatography into the cis and trans isomers thereof, the compounds of general formula I obtained which occur as racemates may be separated by methods known per se (cf. Allinger N. L. and Eliel E. L. In "Topics in Stereochemistry", Vol. 6, Wiley Interscience, 1971) into their optical antipodes and compounds of general formula I with at least 2 asymmetric carbon atoms may be resolved into their diastereomers on the basis of their physical-chemical differences using methods known per se, e.g. by chromatography and/or fractional crystallisation, and, if these compounds are obtained in racemic form, they may subsequently be resolved into the enantiomers as mentioned above.

**[0381]** The enantiomers are preferably separated by column separation on chiral phases or by recrystallisation from an optically active solvent or by reacting with an optically active substance which forms salts or derivatives such as e.g. esters or amides with the racemic compound, particularly acids and the activated derivatives or alcohols thereof, and separating the diastereomeric mixture of salts or derivatives thus obtained, e.g. on the basis of their differences in solubility, whilst the free antipodes may be released from the pure diastereomeric salts or derivatives by the action of suitable agents. Optically active acids in common use are e.g. the D- and L-forms of tartaric acid or dibenzoyltartaric acid, di-o-tolyltartaric acid, malic acid, mandelic acid, camphorsul-



phonic acid, glutamic acid, aspartic acid or quinic acid. An optically active alcohol may be for example (+) or (–)-menthol and an optically active acyl group in amides, for example, may be a (+)- or (–)-menthyloxycarbonyl.

[0382] Furthermore, the compounds of formula I obtained may be converted into the salts thereof, particularly for pharmaceutical use into the physiologically acceptable salts with inorganic or organic acids or bases. Acids which may be used for this purpose include for example hydrochloric acid, hydrobromic acid, sulphuric acid, methanesulphonic acid, ethanesulphonic acid, benzenesulphonic acid, p-toluenesulphonic acid, phosphoric acid, fumaric acid, succinic acid, benzoic acid, salicylic acid, mandelic acid, lactic acid, malonic acid, citric acid, L-malic acid, L-tartaric acid or maleic acid. Suitable bases for this purpose include for example sodium hydroxide solution, potassium hydroxide solution, calcium hydroxide, diethanolamine or N-methyl-D-glucamine.

[0383] The compounds of general formulae II to XXIII used as starting materials are known from the literature to some extent or may be obtained by methods known from the literature (cf. Examples I to V), optionally with the additional introduction of protecting groups.

[0384] Standard processes for preparing the starting materials are described for example in "March's Advanced Organic Chemistry" by Michael B. Smith and Jerry March, Wiley-VCH or in "Science of Synthesis/Houben-Weyl" published by Thieme.

[0385] One possible method of obtaining compounds of general formula (V) and (VI) is as follows:




[0386] Starting from a compound of general formula (XV), wherein PG denotes a protective group such as for example benzyl, 4-methoxybenzyl or 2,4-dimethoxybenzyl, the reaction is carried out with a compound of general formula (III) analogously to the previously described process a) to obtain a compound of general formula (XVI). The compounds of general formula (XV) are known from the literature (cf. E.g. WO 2004/108664 or WO 2007/003486) or may be obtained by methods known from the literature.


[0387] The cleaving of the protective group from a compound of general formula (XVI) to obtain a compound of general formula (V) is carried out, if PG denotes benzyl, with hydrogen, for example, in the presence of a catalyst such as palladium/charcoal. The cleaving of the protective group if PG denotes 4-methoxybenzyl or 2,4-dimethoxybenzyl may also be carried out oxidatively (e.g. with cerium (IV)-ammonium nitrate or with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone) or with acids (e.g. with trifluoroacetic acid in the presence of anisole, thioanisole, pentamethylbenzene or triethylsilane).

[0388] A compound of general formula (V) may then be converted into a compound of general formula (VI), as described in the previous process c). The meanings for  $R^b$ ,  $R^c$ ,  $R^d$ , A,  $Z^1$  and  $Z^2$  in the compounds of Scheme 1 are defined as mentioned hereinbefore.

[0389] Another possible way of obtaining compounds of general formula (XVI) is as follows:








**[0390]** Starting from a compound of general formula (XV), wherein PG denotes a protective group such as for example benzyl, 4-methoxybenzyl or 2,4-dimethoxybenzyl, the reaction is carried out with a compound of general formula (XVII) analogously to the previously described process a) to form a compound of general formula (XVIII). After cleaving of the ketal to form the ketone, the reaction is carried out to obtain a compound of general formula (XX) (analogously to e.g. WO 2008/079735). After reduction of the double bond (analogously to e.g. WO 2008/079735) the protective group PG' is cleaved to form a compound of general formula (XXII). The other steps leading to a compound of general formula (XVI) are carried out for example analogously to Examples III, II, I, and 1. The meanings of R<sup>b</sup>, R<sup>c</sup>, R<sup>d</sup>, A and Z<sup>1</sup> in the compounds of Scheme 2 are defined as hereinbefore. PG' denotes a protective group, for example formyl, acetyl, trifluoroacetyl, ethoxycarbonyl, tert.-butoxycarbonyl or benzyloxycarbonyl and R' denotes a C<sub>1-4</sub>-alkyl group, for example methyl or ethyl.

**[0391]** Another possible way of obtaining compounds of general formula (XXI) is as follows:



**[0392]** Starting from a compound of general formula (XV), wherein PG denotes a protective group such as for example benzyl, 4-methoxybenzyl or 2,4-dimethoxybenzyl, the reaction is carried out with a compound of general formula (XXIII) analogously to the previously described process a) to form a compound of general formula (XXI). The meanings of R<sup>d</sup>, PG, PG', R' and Z<sup>1</sup> in the compounds of Scheme 3 are as hereinbefore defined.

**[0393]** As already mentioned hereinbefore, the compounds of general formula (I) according to the invention and the physiologically acceptable salts thereof have valuable pharmacological properties, particularly an inhibiting effect on signal transduction mediated by the Epidermal Growth Factor receptor (EGF-R), whilst this may be achieved for example by inhibiting ligand bonding, receptor dimerisation or tyrosine kinase itself. It is also possible to block the transmission of signals to components located further downstream.


**[0394]** The following Examples are intended to illustrate the present invention without restricting it:

#### Preparation of the Starting Compounds

##### Example I

Methyl cis/trans-(S)-[(2-amino-ethyl)-methyl-amino]-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

##### [0395]



**[0396]** A mixture of 970 mg methyl cis/trans-(S)-[(2-tert.-butoxycarbonylamino-ethyl)-methyl-amino]-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-

xyloxy]-cyclohexyl}-acetate, 2 ml trifluoroacetic acid and 8 ml methylene chloride is stirred overnight at ambient temperature. Then the reaction mixture is evaporated down in vacuo, leaving a brownish solid, which is reacted further without any further purification.


[0397] Yield: 700 mg (85% of theory)

[0398] Mass spectrum (ESI<sup>+</sup>): m/z=546, 548 [M+H]<sup>+</sup>

[0399] The following compounds are obtained analogously to Example I:

(1) methyl cis/trans-(S)-amino-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

[0400]




[0401] R<sub>f</sub> value: 0.42 (silica gel, methylene chloride/methanol/conc. Ammonia=90:10:1)

[0402] Mass spectrum (ESI<sup>+</sup>): m/z=489, 491 [M+H]<sup>+</sup>

(2) methyl cis/trans-(S)-(2-amino-ethylamino)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate


[0403]



[0404] Mass spectrum (ESI<sup>+</sup>): m/z=532, 534 [M+H]<sup>+</sup>

(3) methyl cis/trans-(S)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-(2-methylamino-ethylamino)-acetate


[0405]



[0406] Mass spectrum (ESI<sup>+</sup>): m/z=546, 548 [M+H]<sup>+</sup>

(4) methyl cis/trans-(S)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-[methyl-(2-methylamino-ethyl)-amino]-acetate


[0407]



[0408] Mass spectrum (ESI<sup>+</sup>): m/z=560, 562 [M+H]<sup>+</sup>

(5) methyl cis-(R)-(2-amino-ethylamino)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate


[0409]



[0410] Mass spectrum (ESI<sup>+</sup>): m/z=530, 532 [M+H]<sup>+</sup>

(6) methyl cis-(R)-amino-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate


[0411]



[0412] Mass spectrum (ESI<sup>+</sup>): m/z=489, 491 [M+H]<sup>+</sup>

(7) methyl trans-(R)-(2-amino-ethylamino)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate


[0413]



[0414] Mass spectrum (ESI<sup>+</sup>): m/z=532, 534 [M+H]<sup>+</sup>

(8) methyl trans-(R)-amino-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

[0415]




[0416] Mass spectrum (ESI<sup>+</sup>): m/z=489, 491 [M+H]<sup>+</sup>

Example II

Methyl cis/trans-(S)-[(2-tert.-butoxycarbonylamino-ethyl)-methyl-amino]-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

[0417]



[0418] 240 µl of a 37% aqueous formaldehyde solution, 100 µl glacial acetic acid and 500 mg sodium triacetoxyborohydride are added to 1.00 g methyl cis/trans-(S)-[(2-tert.-butoxycarbonylamino-ethylamino)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate in 30 ml of tetrahydrofuran. The reaction mixture is stirred overnight at ambient temperature, diluted with ethyl acetate, combined with 10% potassium carbonate solution and vigorously stirred. The organic phase is separated off, washed with water and saturated sodium chloride solution, dried on magnesium sulphate and evaporated down.

[0419] Yield: 970 mg (95% of theory)


[0420] Mass spectrum (ESI<sup>+</sup>): m/z=646, 648 [M+H]<sup>+</sup>

**[0421]** The following compound is obtained analogously to Example II:

(1) methyl cis/trans-(S)-{[2-(tert.-butoxycarbonyl-methyl-amino)-ethyl]-methyl-amino}-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-chazolin-6-yloxy]-cyclohexyl}-acetate

**[0422]**

**[0425]** 490 mg tert. Butyl (2-oxo-ethyl)-carbamate followed by 180  $\mu$ l glacial acetic acid and 150 mg sodium triacetoxyborohydride are added to 1.50 g methyl cis/trans-(S)-amino-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate in 40 ml of tetrahydrofuran. The reaction mixture is stirred overnight at ambient temperature, then another 70 mg tert. Butyl (2-oxo-ethyl)-carbamate and 150 mg sodium triacetoxyborohydride are added and the mixture is stirred for a further four hours at ambient temperature. For working up the reaction mixture is



**[0423]** Mass spectrum (ESI<sup>+</sup>): m/z=660, 662 [M+H]<sup>+</sup>


#### Example III

Methyl cis/trans-(S)-(2-tert.-butoxycarbonylamino-ethylamino)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

**[0424]**

diluted with 20 ml of ethyl acetate, combined with 10 ml 10% potassium carbonate solution and stirred thoroughly. The aqueous phase is separated off and extracted with ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried on magnesium sulphate and evaporated down. The crude product is reacted further without any further purification.


**[0426]** Mass spectrum (ESI<sup>+</sup>): m/z=632, 634 [M+H]<sup>+</sup>



[0427] The following compounds are obtained analogously to Example III:

- (1) methyl cis/trans-(S)-[2-(tert.-butoxycarbonylmethyl-amino)-ethylamino]-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

[0428]




[0429]  $R_f$  value: 0.16 (silica gel, methylene chloride/methanol/conc. Ammonia=95:5:0.1)

[0430] Mass spectrum (ESI<sup>+</sup>):  $m/z$ =646, 648 [M+H]<sup>+</sup>

- (2) methyl cis-(R)-[2-(tert.-butoxycarbonylamino-ethylamino)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate


[0431]



[0432] Mass spectrum (ESI<sup>+</sup>):  $m/z$ =632, 634 [M+H]<sup>+</sup>

- (3) methyl trans-(R)-[2-(tert.-butoxycarbonylamino-ethylamino)-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

[0433]




[0434] Mass spectrum (ESI<sup>+</sup>):  $m/z$ =632, 634 [M+H]<sup>+</sup>

## Example IV

methyl cis/trans-(S)-tert.-butoxycarbonylamino-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

[0435]



[0436] 10.00 g 4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-ol in 70 ml N,N-dimethylformamide are heated to 50° C., combined with 7.00 g potassium carbonate as well as 14.00 g methyl cis/trans-(S)-tert.-butoxycarbonylamino-(4-methanesulphonyloxy-cyclohexyl)-acetate and heated to 80° C. The reaction mixture is stirred overnight at 80° C., then another 3.50 g methyl cis/trans-(S)-tert.-butoxycarbonylamino-(4-methanesulphonyloxy-cyclohexyl)-acetate and 2.60 g potassium carbonate are added.

[0437] After a further four hours at 80° C. the reaction mixture is cooled and diluted with ethyl acetate and water. The aqueous phase is separated off and extracted with ethyl acetate. The combined organic phases are dried on magnesium sulphate and evaporated down. The crude product is purified by chromatography through a silica gel column with methylene chloride/methanol/conc. Ammonia (99/1/0.1 auf 8/2/0.1) as eluant.


[0438] Yield: 5.90 g (32% of theory)

[0439] Mass spectrum (ESI<sup>+</sup>): m/z=589, 591 [M+H]<sup>+</sup>

[0440] The following compounds are obtained analogously to Example IV:

(1) methyl cis-(R)-tert.-butoxycarbonylamino-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate


[0441]



[0442] Mass spectrum (ESI<sup>+</sup>): m/z=589, 591 [M+H]<sup>+</sup>

(2) methyl trans-(R)-tert.-butoxycarbonylamino-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate

[0443]




[0444] Mass spectrum (ESI<sup>+</sup>): m/z=589, 591 [M+H]<sup>+</sup>

## Example V

Methyl cis/trans-(S)-tert.-butoxycarbonylamino-(4-methanesulphonyloxy-cyclohexyl)-acetate

[0445]



[0446] 5.20 ml methanesulphonic acid chloride are slowly added dropwise to 17.55 g methyl cis/trans-(S)-tert.-butoxycarbonylamino-(4-hydroxy-cyclohexyl)-acetate (see WO2004/110436) and 10.50 ml triethylamine in 170 ml methylene chloride while cooling with an ice bath, the temperature being kept below 10° C. Then the reaction mixture is heated to ambient temperature and stirred overnight.

[0447] For working up 50 ml saturated sodium hydrogen carbonate solution are added. The aqueous phase is separated off and extracted with methylene chloride. The combined organic phases are washed with saturated sodium chloride solution, dried on magnesium sulphate and evaporated down. A viscous oil remains which is further reacted without any further purification.


[0448] Yield: 21.70 g (97% of theory)

[0449] Mass spectrum (ESI<sup>+</sup>): m/z=383 [M<sub>+</sub>NH<sub>4</sub>]<sup>+</sup>

[0450] The following compounds are obtained analogously to Example V:

- (1) methyl cis-(R)-tert.-butoxycarbonylamino-(4-methanesulphonyloxy-cyclohexyl)-acetate

[0451]




[0452] The starting material, methyl cis-(R)-tert.-butoxycarbonylamino-(4-hydroxy-cyclohexyl)-acetate, is obtained starting from (R)-4-hydroxy-phenylglycine analogously to WO2004/110436.

[0453] Mass spectrum (ESI<sup>+</sup>): m/z=366 [M+H]<sup>+</sup>

- (2) methyl trans-(R)-tert.-butoxycarbonylamino-(4-methanesulphonyloxy-cyclohexyl)-acetate

[0454]



[0455] The starting material, methyl trans-(R)-tert.-butoxycarbonylamino-(4-hydroxy-cyclohexyl)-acetate, is obtained starting from (R)-4-hydroxy-phenylglycine analogously to WO2004/110436.


[0456] Mass spectrum (ESI<sup>+</sup>): m/z=366 [M+H]<sup>+</sup>

Preparation of the End Compounds

#### Example 1

cis/trans-(S)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-4-methyl-piperazin-2-one

[0457]



[0458] A mixture of 700 mg methyl [(2-amino-ethyl)-methyl-amino]-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-acetate, 7 ml of methanol and 0.65 ml 4N sodium hydroxide solution is stirred for three hours at ambient temperature. Then the reaction mixture is evaporated down and extracted with ethyl acetate. The organic phase is dried on magnesium sulphate and evaporated down. The flask residue is purified by chromatography through a silica gel column with methylene chloride/methanol/conc. ammonia (98/2/0.1 auf 8/2/0.1) as eluant. The crude product is stirred with diisopropylether, suction filtered and dried.


[0459] Yield: 300 mg (46% of theory)

[0460] Mass spectrum (ESI<sup>+</sup>): m/z=514, 516 [M+H]<sup>+</sup>

[0461] The following compounds are obtained analogously to Example 1:

- (1) cis/trans-(S)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-piperazin-2-one

[0462]



[0463] R<sub>f</sub>-value: 0.20 (silica gel, methylene chloride/methanol/conc. ammonia=90:10:1)

[0464] Mass spectrum (ESI<sup>+</sup>): m/z=500, 502 [M+H]<sup>+</sup>

[0465] The cis/trans-mixture is separated by supercritical liquid chromatography under the following conditions:

column: Daicel ASH 20x250 mm

eluant: CO<sub>2</sub>/methanol (+0.2% diethylamine) (60:40)

flow: 70 ml/min


injection volume: 450 µl

sample concentration: 100 mg/l

[0466] The assignment of the isomers is carried out by 1H-NMR spectroscopy (400 MHz, dimethylsulphoxide-d<sub>6</sub>)

- a) cis-(S)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-piperazin-2-one


[0467]



characteristic signal at 4.75 (1H, m)

b) trans-(S)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-piperazin-2-one


[0468]



characteristic signal at 4.42 (1H, m)

(2) trans-(S)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-1-methyl-piperazin-2-one

[0469]




[0470]  $R_f$  value: 0.31 (silica gel, methylene chloride/methanol/conc. Ammonia=90:10:1)

[0471] Mass spectrum (ESI<sup>+</sup>):  $m/z$ =514, 516 [M+H]<sup>+</sup>

(3) cis/trans-(S)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-1,4-dimethyl-piperazin-2-one

[0472]




[0473]  $R_f$  value: 0.35 (silica gel, methylene chloride/methanol/conc. Ammonia=90:10:1)

[0474] Mass spectrum (ESI<sup>+</sup>):  $m/z$ =528, 530 [M+H]<sup>+</sup>

(4) cis-(R)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-piperazin-2-one

[0475]




[0476] The cyclisation is carried out in methanol in the presence of triethylamine at ambient temperature.

[0477] Mass spectrum (ESI<sup>+</sup>):  $m/z$ =500, 502 [M+H]<sup>+</sup>

(5) trans-(R)-3-{4-[4-(3-chloro-2-fluoro-phenylamino)-7-methoxy-quinazolin-6-yloxy]-cyclohexyl}-piperazin-2-one

[0478]



[0479] The cyclisation is carried out in methanol in the presence of triethylamine at ambient temperature.

[0480] Mass spectrum (ESI<sup>+</sup>):  $m/z$ =500, 502 [M+H]<sup>+</sup>

#### Biological Test

[0481] The biological properties of the new compounds are investigated as follows, for example:

[0482] The inhibition of the EGF-R-mediated signal transduction can be demonstrated e.g. with cells which express human EGF-R and whose survival and proliferation depend on stimulation by EGF or TGF- $\alpha$ . A murine haematopoietic cell line is genetically modified so as to express functional human EGF-R. The proliferation of this cell line can therefore be stimulated by EGF.

[0483] The test is carried out as follows:

[0484] The cells are cultivated in RPMI/1640 medium. The proliferation is stimulated with 20 ng/ml of human EGF (Promega). To investigate the inhibitory activity of the compounds according to the invention these compounds are dissolved in 100% dimethylsulphoxide (DMSO) and added to



the cultures in various dilutions, the maximum DMSO concentration being 1%. The cultures are incubated for 48 hours at 37° C.

**[0485]** In order to determine the inhibitory activity of the compounds according to the invention the relative cell number is measured in O.D. Units using the Cell Titer 96™ Aqueous Non-Radioactive Cell Proliferation Assay (Promega). The relative cell number is calculated as a percentage of the control and the concentration of active substance which inhibits the proliferation of the cells by 50% (IC<sub>50</sub>) is derived therefrom.

**[0486]** The compounds of general formula (I) according to the invention exhibit IC<sub>50</sub> values of <10 micromolar, preferably <1 micromolar, for example.

Compound (Example No.)	Inhibition of the EGFR- dependent proliferation IC <sub>50</sub> [nM]
1	1.0
1(1a)	10
1(1b)	1.4
1(2)	1.0
1(3)	1.0
1(4)	12.5
1(5)	1.6

#### Indications

**[0487]** As has been found, the compounds of formula (I) are characterised in that by their versatility in the therapeutic field. Particular mention should be made of the possible applications for which the compounds of formula (I) according to the invention are preferably used on the basis of their pharmaceutical efficacy as tyrosine inhibitors.

**[0488]** The compounds of general formula (I) according to the invention thus inhibit signal transduction by tyrosine kinases, as demonstrated by the example of the human EGF receptor, and are therefore useful for treating pathophysiological processes caused by hyperfunction of tyrosine kinases. These are e.g. benign or malignant tumours, particularly tumours of epithelial and neuroepithelial origin, metastatisation and the abnormal proliferation of vascular endothelial cells (neovascularisation).

**[0489]** The compounds according to the invention are also useful for preventing and treating diseases of the airways and lungs which are accompanied by increased or altered production of mucus caused by stimulation of tyrosine kinases, e.g. in inflammatory diseases of the airways such as chronic bronchitis, chronic obstructive bronchitis, asthma, bronchiectasis, allergic or non-allergic rhinitis or sinusitis, cystic fibrosis,  $\alpha$ 1-antitrypsin deficiency, or coughs, pulmonary emphysema, pulmonary fibrosis and hyperreactive airways.

**[0490]** The compounds are also suitable for treating diseases of the gastrointestinal tract and bile duct and gall bladder which are associated with disrupted activity of the tyrosine kinases, such as may be found e.g. in chronic inflammatory changes such as cholecystitis, Crohn's disease, ulcerative colitis, and ulcers in the gastrointestinal tract or such as may occur in diseases of the gastrointestinal tract which are associated with increased secretions, such as Menetrier's disease, secreting adenomas and protein loss syndrome.

**[0491]** In addition, the compounds of general formula I and the physiologically acceptable salts thereof may be used to

treat other diseases caused by abnormal function of tyrosine kinases, such as e.g. epidermal hyperproliferation (psoriasis), benign prostatic hyperplasia (BPH), inflammatory processes, diseases of the immune system, hyperproliferation of haematopoietic cells, the treatment of nasal polyps, etc.

**[0492]** By reason of their biological properties the compounds according to the invention may be used on their own or in conjunction with other pharmacologically active compounds, for example in tumour therapy, in monotherapy or in conjunction with other anti-tumour therapeutic agents, for example in combination with topoisomerase inhibitors (e.g. etoposide), mitosis inhibitors (e.g. vinblastine), compounds which interact with nucleic acids (e.g. cisplatin, cyclophosphamide, adriamycin), hormone antagonists (e.g. tamoxifen), inhibitors of metabolic processes (e.g. 5-FU etc.), cytokines (e.g. interferons), antibodies, etc. For treating respiratory tract diseases, these compounds may be used on their own or in conjunction with other therapeutic agents for the airways, such as substances with a secretolytic (e.g. ambroxol, N-acetylcysteine), broncholytic (e.g. tiotropium or ipratropium or fenoterol, salmeterol, salbutamol) and/or anti-inflammatory activity (e.g. theophylline or glucocorticoids).

**[0493]** For treating diseases in the region of the gastrointestinal tract, these compounds may also be administered on their own or in conjunction with substances having an effect on motility or secretion. These combinations may be administered either simultaneously or sequentially.

#### Formulations

**[0494]** The compounds according to the invention may be administered by oral, transdermal, inhalative, parenteral or sublingual route. The compounds according to the invention are present as active ingredients in conventional preparations, for example in compositions consisting essentially of an inert pharmaceutical carrier and an effective dose of the active substance, such as for example tablets, coated tablets, capsules, lozenges, powders, solutions, suspensions, emulsions, syrups, suppositories, transdermal systems etc. An effective dose of the compounds according to the invention is between 0.1 and 5000, preferably between 1 and 500, more preferably between 5-300 mg/dose for oral administration, and between 0.001 and 50, preferably between 0.1 and 10 mg/dose for intravenous, subcutaneous or intramuscular administration. For inhalation, according to the invention, solutions containing 0.01 to 1.0, preferably 0.1 to 0.5% active substance are suitable. For administration by inhalation the use of powders, ethanolic or aqueous solutions is preferred. It is also possible to use the compounds according to the invention as a solution for infusion, preferably in a physiological saline or nutrient saline solution.

**[0495]** The compounds according to the invention may be used on their own or in conjunction with other active substances according to the invention, optionally also in conjunction with other pharmacologically active substances. Suitable formulations include, for example, tablets, capsules, suppositories, solutions, syrups, emulsions or dispersible powders. Corresponding tablets may be obtained for example by mixing the active substance(s) with known excipients, for example inert diluents, such as calcium carbonate, calcium phosphate or lactose, disintegrants such as maize starch or alginic acid, binders such as starch or gelatine, lubricants such as magnesium stearate or talc and/or agents for delaying

release, such as carboxymethyl cellulose, cellulose acetate phthalate, or polyvinyl acetate. The tablets may also comprise several layers.

**[0496]** Coated tablets may be prepared accordingly by coating cores produced analogously to the tablets with substances normally used for tablet coatings, for example colli-done or shellac, gum arabic, talc, titanium dioxide or sugar. To achieve delayed release or prevent incompatibilities the core may also consist of a number of layers. Similarly the tablet coating may consist of a number of layers to achieve delayed release, possibly using the excipients mentioned above for the tablets.

**[0497]** Syrups containing the active substances or combinations thereof according to the invention may additionally contain a sweetener such as saccharine, cyclamate, glycerol or sugar and a flavour enhancer, e.g. a flavouring such as vanillin or orange extract. They may also contain suspension adjuvants or thickeners such as sodium carboxymethyl cellulose, wetting agents such as, for example, condensation products of fatty alcohols with ethylene oxide, or preservatives such as p-hydroxybenzoates.

**[0498]** Solutions for injection are prepared in the usual way, e.g. with the addition of preservatives such as p-hydroxybenzoates, or stabilisers such as alkali metal salts of ethylenediamine tetraacetic acid, and transferred into injection vials or ampoules.

**[0499]** Capsules containing one or more active substances or combinations of active substances may for example be prepared by mixing the active substances with inert carriers such as lactose or sorbitol and packing them into gelatine capsules.

**[0500]** Suitable suppositories may be made for example by mixing with carriers provided for this purpose, such as neutral fats or polyethyleneglycol or the derivatives thereof.

**[0501]** For pharmaceutical use the compounds according to the invention are generally used for warm-blooded vertebrates, particularly humans, in doses of 0.01-100 mg/kg of body weight, preferably 0.1-15 mg/kg. For administration they are formulated with one or more conventional inert carriers and/or diluents, e.g. with corn starch, lactose, glucose, microcrystalline cellulose, magnesium stearate, polyvinylpyrrolidone, citric acid, tartaric acid, water, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol, stearyl alcohol, carboxymethylcellulose or fatty substances such as hard fat or suitable mixtures thereof to produce conventional galenic preparations such as plain or coated tablets, capsules, powders, suspensions, solutions, sprays or suppositories.

**[0502]** The Examples which follow illustrate the present invention without restricting its scope:

#### Examples of Pharmaceutical Formulations

##### A) Coated Tablets Containing 75 mg of Active Substance

###### Composition:

**[0503]** 1 tablet core contains:

active substance	75.0 mg
calcium phosphate	93.0 mg
corn starch	35.5 mg

###### -continued

polyvinylpyrrolidone	10.0 mg
hydroxypropylmethylcellulose	15.0 mg
magnesium stearate	1.5 mg
	230.0 mg

###### Preparation:

**[0504]** The active substance is mixed with calcium phosphate, corn starch, polyvinyl-pyrrolidone, hydroxypropylmethylcellulose and half the specified amount of magnesium stearate. Blanks 13 mm in diameter are produced in a tablet-making machine and these are then rubbed through a screen with a mesh size of 1.5 mm using a suitable machine and mixed with the rest of the magnesium stearate. This granulate is compressed in a tablet-making machine to form tablets of the desired shape.

**[0505]** Weight of core: 230 mg

**[0506]** die: 9 mm, convex

**[0507]** The tablet cores thus produced are coated with a film consisting essentially of hydroxypropylmethylcellulose. The finished film-coated tablets are polished with beeswax.

**[0508]** Weight of coated tablet: 245 mg.

##### B) Tablets Containing 100 mg of Active Substance

###### Composition:

**[0509]** 1 tablet contains:

active substance	100.0 mg
lactose	80.0 mg
corn starch	34.0 mg
polyvinylpyrrolidone	4.0 mg
magnesium stearate	2.0 mg
	220.0 mg

###### Method of Preparation:

**[0510]** The active substance, lactose and starch are mixed together and uniformly moistened with an aqueous solution of the polyvinylpyrrolidone. After the moist composition has been screened (2.0 mm mesh size) and dried in a rack-type drier at 50° C. it is screened again (1.5 mm mesh size) and the lubricant is added. The finished mixture compressed to form tablets.

**[0511]** Weight of tablet: 220 mg

**[0512]** Diameter: 10 mm, biplanar, faceted on both sides and notched on one side.

##### C) Tablets Containing 150 mg of Active Substance

###### Composition:

**[0513]** 1 tablet contains:

active substance	150.0 mg
powdered lactose	89.0 mg
corn starch	40.0 mg

-continued

colloidal silica	10.0 mg
polyvinylpyrrolidone	10.0 mg
magnesium stearate	1.0 mg
	300.0 mg

Preparation:

**[0514]** The active substance mixed with lactose, corn starch and silica is moistened with a 20% aqueous polyvinylpyrrolidone solution and passed through a screen with a mesh size of 1.5 mm. The granules, dried at 45° C., are passed through the same screen again and mixed with the specified amount of magnesium stearate. Tablets are pressed from the mixture.

**[0515]** Weight of tablet: 300 mg

**[0516]** die: 10 mm, flat

D) Hard Gelatine Capsules Containing 150 mg of Active Substance

Composition:

**[0517]** 1 capsule contains:

active substance	150.0 mg
corn starch (dried)	approx. 180.0 mg
lactose (powdered)	approx. 87.0 mg
magnesium stearate	3.0 mg
	approx. 420.0 mg

Preparation:

**[0518]** The active substance is mixed with the excipients, passed through a screen with a mesh size of 0.75 mm and homogeneously mixed using a suitable apparatus. The finished mixture is packed into size 1 hard gelatine capsules.

**[0519]** Capsule filling: approx. 320 mg

**[0520]** Capsule shell: size 1 hard gelatine capsule.

E) Suppositories Containing 150 mg of Active Substance

Composition:

**[0521]** 1 suppository contains:

active substance	150.0 mg
polyethyleneglycol 1500	550.0 mg
polyethyleneglycol 6000	460.0 mg
polyoxyethylene sorbitan monostearate	840.0 mg
	2,000.0 mg

Preparation:

**[0522]** After the suppository mass has been melted the active substance is homogeneously distributed therein and the melt is poured into chilled moulds.

F) Suspension Containing 50 mg of Active Substance

Composition:

**[0523]** 100 ml of suspension contain:

active substance	1.00 g
carboxymethylcellulose-Na-salt	0.10 g
methyl p-hydroxybenzoate	0.05 g
propyl p-hydroxybenzoate	0.01 g
glucose	10.00 g
glycerol	5.00 g
70% sorbitol solution	20.00 g
flavouring	0.30 g
dist. water	ad 100 ml

Preparation:

**[0524]** The distilled water is heated to 70° C. The methyl and propyl p-hydroxybenzoates together with the glycerol and sodium salt of carboxymethylcellulose are dissolved therein with stirring. The solution is cooled to ambient temperature and the active substance is added and homogeneously dispersed therein with stirring. After the sugar, the sorbitol solution and the flavouring have been added and dissolved, the suspension is evacuated with stirring to eliminate air.

**[0525]** 5 ml of suspension contain 50 mg of active substance.

G) Ampoules Containing 10 mg Active Substance

Composition:

**[0526]**

active substance	10.0 mg
0.01N hydrochloric acid	q.s.
double-distilled water	ad 2.0 ml

Preparation:

**[0527]** The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with common salt, filtered sterile and transferred into 2 ml ampoules.

H) Ampoules Containing 50 mg of Active Substance

Composition:

**[0528]**

active substance	50.0 mg
0.01N hydrochloric acid	q.s.
double-distilled water	ad 10.0 ml

Preparation:

**[0529]** The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with common salt, filtered sterile and transferred into 10 ml ampoules.

## I) Capsules for Powder Inhalation Containing 5 Mg of Active Substance

[0530] 1 capsule contains:

active substance	5.0 mg
lactose for inhalation	15.0 mg
	20.0 mg

## Preparation:

[0531] The active substance is mixed with lactose for inhalation. The mixture is packed into capsules in a capsule-making machine (weight of the empty capsule approx. 50 mg).

weight of capsule:	70.0 mg
size of capsule =	3

## J) Solution for Inhalation for Hand-Held Nebulisers Containing 2.5 mg Active Substance

[0532] 1 spray contains:


active substance	2,500 mg
benzalkonium chloride	0.001 mg
1N hydrochloric acid	q.s.
ethanol/water (50/50)	ad 15,000 mg

## Preparation:

[0533] The active substance and benzalkonium chloride are dissolved in ethanol/water (50/50). The pH of the solution is adjusted with 1N hydrochloric acid. The resulting solution is filtered and transferred into suitable containers for use in hand-held nebulisers (cartridges).

[0534] Contents of the container: 4.5 g

1. A compound of formula (I)



wherein:

$R^a$  denotes a phenyl or 1-phenylethyl group, wherein the phenyl nucleus is substituted in each case by the groups  $R^1$  to  $R^3$ , wherein

$R^1$  and  $R^2$  which may be identical or different, denote hydrogen or

a group selected from among

F, Cl, Br, I,  $\text{OCH}_2\text{F}$ ,  $\text{OCHF}_2$ ,  $\text{OCF}_3$ ,  $\text{CH}_2\text{F}$ ,  $\text{CHF}_2$ ,  $\text{CF}_3$ , CN,  $\text{NO}_2$ ,  $\text{NH}_2$  and OH,

or

a group selected from among

$\text{C}_{1-4}$ -alkyl,  $\text{C}_{1-4}$ -alkyl-O,  $\text{C}_{2-3}$ -alkenyl,  $\text{C}_{2-3}$ -alkynyl, phenyl, phenyl-O, phenyl- $\text{C}_{1-3}$ -alkyl, phenyl- $\text{C}_{1-3}$ -alkyl-O, heteroaryl, heteroaryl-O, heteroaryl- $\text{C}_{1-3}$ -alkyl, heteroaryl- $\text{C}_{1-3}$ -alkyl-O, wherein the above-mentioned phenyl groups are mono- or disubstituted by  $R^5$  groups,

and

 $R^3$  denotes hydrogen,

or

a group selected from among

F, Cl, Br and  $\text{CH}_3$ ,

$R^b$  denotes hydrogen, or an optionally substituted group selected from among  $\text{C}_{1-6}$ -alkyl,  $\text{C}_{3-6}$ -cycloalkyl and  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl,

$R^c$  denotes hydrogen, or an optionally substituted group selected from among  $\text{C}_{1-6}$ -alkyl,  $\text{C}_{3-6}$ -cycloalkyl,  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl,  $\text{C}_{1-6}$ -alkyl-CO,  $\text{C}_{3-6}$ -cycloalkyl-CO,  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl-CO,  $\text{C}_{1-6}$ -alkyl- $\text{SO}_2$ ,  $\text{C}_{3-6}$ -cycloalkyl- $\text{SO}_2$ ,  $\text{C}_{3-6}$ -cycloalkyl- $\text{C}_{1-3}$ -alkyl- $\text{SO}_2$ , phenyl-CO— and phenyl- $\text{SO}_2$ ,

 $R^d$  denotes hydrogen or

a group selected from among

F, Cl, Br, I, OH,  $\text{C}_{1-4}$ -alkyl,  $\text{C}_{1-4}$ -alkyl-O,  $\text{C}_{1-2}$ -alkyl-O substituted by 1 to 3 fluorine atoms,  $\text{C}_{3-7}$ -cycloalkyl-O,  $\text{C}_{3-7}$ -cycloalkyl- $\text{C}_{1-4}$ -alkyl-O, tetrahydrofuran-3-yl-O, tetrahydropyran-3-yl-O, tetrahydro-pyran-4-yl-O, tetrahydrofuran-yl- $\text{C}_{1-4}$ -alkyl-O— and tetrahydropyran-yl- $\text{C}_{1-4}$ -alkyl-O,

or

$R^4$ — $\text{C}_{1-4}$ -alkyl, wherein the linking of the groups  $R^4$  may take place via each C atom of the alkyl group,

or

$R^4$ — $\text{C}_{2-4}$ -alkyl-O, wherein the group  $R^4$  is separated from the oxygen atom by at least 2 C atoms,

or

a group selected from among

pyrrolidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, pyrrolidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, piperidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, piperidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, piperidin-4-yl- $\text{C}_{1-4}$ -alkyl-O, azepan-2-yl- $\text{C}_{1-4}$ -alkyl-O, azepan-3-yl- $\text{C}_{1-4}$ -alkyl-O, azepan-4-yl- $\text{C}_{1-4}$ -alkyl-O, morpholin-2-yl- $\text{C}_{1-4}$ -alkyl-O, morpholin-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-pyrrolidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-pyrrolidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-piperidin-2-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-piperidin-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-piperidin-4-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-azepan-2-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-azepan-3-yl- $\text{C}_{1-4}$ -alkyl-O, 1-( $\text{C}_{1-3}$ -alkyl)-azepan-4-yl- $\text{C}_{1-4}$ -alkyl-O, 4-( $\text{C}_{1-3}$ -alkyl)-morpholin-2-yl- $\text{C}_{1-4}$ -alkyl-O and 4-( $\text{C}_{1-3}$ -alkyl)-morpholin-3-yl- $\text{C}_{1-4}$ -alkyl-O,

wherein

$R^4$  denotes a group, which may be identical or different, selected from among OH,  $\text{C}_{1-3}$ -alkyl-O,  $\text{C}_{3-6}$ -cycloalkyl-O,  $\text{NH}_2$ ,  $\text{C}_{1-3}$ -alkyl-NH, ( $\text{C}_{1-3}$ -alkyl) $_2$ N, (2-methoxyethyl) $_2$ N, pyrrolidin-1-yl, piperidin-1-yl, azepan-1-yl, morpholin-4-yl, 1,4-oxazepan-4-yl, 2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl, 3-oxa-8-aza-bicyclo[3.2.1]oct-8-yl, 8-oxa-3-aza-bicyclo[3.2.1]oct-3-yl, piperazin-1-yl, 4-( $\text{C}_{1-3}$ -alkyl)-piperazin-1-yl, 1,4-diazepan-

1-yl, 4-(C<sub>1-3</sub>-alkyl)-1,4-diazepan-1-yl, HCO—NH, C<sub>1-4</sub>-alkyl-CO—NH, C<sub>1-3</sub>-alkyl-O—C<sub>1-3</sub>-alkyl-CO—NH, C<sub>1-4</sub>-alkyl-O—CO—NH, H<sub>2</sub>NCONH, C<sub>1-3</sub>-alkyl-NH—CO—NH, (C<sub>1-3</sub>-alkyl)<sub>2</sub>N—CONH, pyrrolidin-1-yl-CO—NH, piperidin-1-yl-CO—NH, piperazin-1-yl-CO—NH, 4-(C<sub>1-3</sub>-alkyl)-piperazin-1-yl-CO—NH, morpholin-4-yl-CO—NH and C<sub>1-4</sub>-alkyl-SO<sub>2</sub>—NH, wherein the pyrrolidinyl, piperidinyl, azepan-1-yl, piperazinyl, 1,4-diazepan-1-yl, morpholinyl and 1,4-oxazepan-4-yl groups mentioned above in the definition of the group R<sup>d</sup> may each additionally be substituted by one or two C<sub>1-3</sub>-alkyl groups,

and

wherein the above-mentioned phenyl groups is mono- or disubstituted by R<sup>5</sup> groups, wherein

R<sup>5</sup> denotes hydrogen, or

a group, which may be identical or different, selected from among

F, Cl, Br, I, OH, CN, C<sub>1-3</sub>-alkyl, C<sub>1-3</sub>-alkyl-O, CHF<sub>2</sub>, CF<sub>3</sub>, —O—CHF<sub>2</sub> and —O—CF<sub>3</sub>,

and

unless stated otherwise, the above-mentioned alkyl groups may be straight-chain or branched,

A denotes —CO or —C<sub>1-3</sub>-alkylene,

wherein the —C<sub>1-3</sub>-alkylene group may be 1-, 2-, 3- or 4-substituted by a group R<sup>6</sup>,

and

R<sup>6</sup> which may be identical or different, denotes hydrogen, or

a group selected from among OH, C<sub>1-4</sub>-alkyl and —O—C<sub>1-4</sub>-alkyl,

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

2. The compounds according to claim 1,

wherein:

R<sup>a</sup> denotes a group selected from among 3-chloro-2-fluoro-phenyl, 3-chloro-4-fluoro-phenyl, 5-chloro-2-fluoro-phenyl, 2-fluoro-3-methyl-phenyl, 2-fluoro-5-methyl-phenyl, 4-fluoro-3-methyl-phenyl and 3-chloro-2-methyl-phenyl group,

R<sup>b</sup> and R<sup>c</sup> which may be identical or different,

denote hydrogen or C<sub>1-3</sub>-alkyl,

R<sup>d</sup> denotes C<sub>1-3</sub>-alkyl-O,

unless stated otherwise, the above-mentioned alkyl groups may be straight-chain or branched,

A denotes —CH<sub>2</sub>CH<sub>2</sub>, wherein the —CH<sub>2</sub>CH<sub>2</sub>— group may be substituted by 1 or 2 methyl groups,

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

3. (canceled)

4. A method for the treatment of inflammatory or allergic diseases of the airways comprising administering to a patient in need thereof a therapeutically effective amount of a compound according to claim 1 or a pharmacologically acceptable acid addition salt thereof.

5. The method according to claim 4 wherein the inflammatory or allergic disease of the airways is selected from among chronic bronchitis, acute bronchitis, bronchitis caused by bacterial or viral infection or fungi or helminths, allergic bronchitis, toxic bronchitis, chronic obstructive bronchitis (COPD), asthma (intrinsic or allergic), paediatric asthma, bronchiectasis, allergic alveolitis, allergic or non-allergic rhinitis, chronic sinusitis, cystic fibrosis or mucoviscidosis, alpha-1-antitrypsin deficiency, cough, pulmonary emphysema, interstitial lung diseases, alveolitis, hyperreactive airways, nasal polyps, pulmonary oedema, pneumonitis of different origins, e.g. radiation-induced or caused by aspiration or infectious pneumonitis, collagenoses such as lupus erythematoses, systemic sclerodermy, sarcoidosis and Boeck's disease.

6. The method according to claim 4, characterised in that it relates to inflammatory or allergic conditions in which autoimmune reactions are involved.

7. A method for the treatment of benign or malignant tumours comprising administering to a patient in need thereof a therapeutically effective amount of a compound according to claim 1 or a pharmacologically acceptable acid addition salt thereof.

8. A pharmaceutical formulation containing a compound of formula (I) according to claim 1.

9. An orally administered pharmaceutical formulation according to claim 8.

10. A medicament combinations which contains, besides one or more compounds of formula (I) according to claim 1, as further active substances, one or more compounds selected from among the categories of betamimetics, anticholinergics, corticosteroids, further PDE4-inhibitors, LTD4-antagonists, EGFR-inhibitors, dopamine agonists, H1-antihistamines, PAF-antagonists and PI3-kinase inhibitors or double or triple combinations thereof.

\* \* \* \* \*