

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
24 September 2009 (24.09.2009)(10) International Publication Number
WO 2009/117540 A1

(51) International Patent Classification:

<i>C07D 498/18</i> (2006.01)	<i>A61K 31/55</i> (2006.01)
<i>A61K 31/5386</i> (2006.01)	<i>C07D 471/08</i> (2006.01)
<i>C07D 471/18</i> (2006.01)	<i>A61P 31/18</i> (2006.01)
<i>A61K 31/529</i> (2006.01)	<i>C07D 493/18</i> (2006.01)
<i>C07D 487/08</i> (2006.01)	

(21) International Application Number:

PCT/US2009/037596

(22) International Filing Date:

19 March 2009 (19.03.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/037,729	19 March 2008 (19.03.2008)	US
12/406,268	18 March 2009 (18.03.2009)	US

(71) Applicant (for all designated States except US): **BRISTOL-MYERS SQUIBB COMPANY** [US/US]; Route 206 And Province Line Road, Princeton, NJ 08543-4000 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BEAULIEU, Francis** [CA/CA]; 370 Leotable Dubuc, La Prairie, Quebec, J5R 5M4 (CA). **OUELLET, Carl** [CA/CA]; 555 Rue De La Verendrye, Boucherville, Quebec, J4B 2Y3 (CA). **NAIDU, B., Narasimhulu** [IN/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **PATEL, Manoj** [CA/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **UEDA, Yasutsugu** [JP/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **CONNOLLY, Timothy, P.**

[CA/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **WEISS, Jonathan, R.** [US/US]; 700 Huron Avenue, Apt. 13d, Cambridge, MA 02138 (US). **WALKER, Michael, A.** [US/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **MEANWELL, Nicholas, A.** [US/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **PEESE, Kevin, M.** [US/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **SORENSEN, Margaret, E.** [US/US]; C/o Bristol-myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US).


(74) Agents: **EPPERSON, James** et al.; Bristol-myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

[Continued on next page]

(54) Title: BRIDGED HETEROCYCLES AS HIV INTEGRASE INHIBITORS



(57) Abstract: The disclosure generally relates to the novel compounds of formula I, including their salts, which inhibit HIV integrase and prevent viral integration into human DNA. This action makes the compounds useful for treating HIV infection and AIDS. The invention also encompasses pharmaceutical compositions and methods for treating those infected with HIV. Formula I.



MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR),
OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

BRIDGE HETEROCYCLES AS HIV INTEGRASE INHIBITORS.

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of U.S. non-provisional application serial number 12/406,268 filed March 18, 2009 and U.S. provisional application serial number 61/037,729 filed March 19, 2008.

BACKGROUND OF THE INVENTION

10

The disclosure generally relates to the novel compounds of formula I, including their salts, which inhibit HIV integrase and prevent viral integration into human DNA. This action makes the compounds useful for treating HIV infection and AIDS. The invention also encompasses pharmaceutical compositions and 15 methods for treating those infected with HIV.

Human immunodeficiency virus (HIV) has been identified as the etiological agent responsible for acquired immune deficiency syndrome (AIDS), a fatal disease characterized by destruction of the immune system and the inability to fight off life 20 threatening opportunistic infections. Recent statistics (UNAIDS: Report on the Global HIV/AIDS Epidemic, December 1998), indicate that as many as 33 million people worldwide are infected with the virus. In addition to the large number of individuals already infected, the virus continues to spread. Estimates from 1998 point to close to 6 million new infections in that year alone. In the same year there 25 were approximately 2.5 million deaths associated with HIV and AIDS.

There are currently a number of antiviral drugs available to combat the infection. These drugs can be divided into four classes based on the viral protein they target and their mode of action. In particular, saquinavir, indinavir, ritonavir, 30 nelfinavir atazanavir darunavir, amprenavir, fosamprenavir, lopinavir and tipranavir are competitive inhibitors of the aspartyl protease expressed by HIV. Zidovudine, didanosine, stavudine, lamivudine, zalcitabine, emtricitabine, tenofovir and abacavir are nucleoside reverse transcriptase inhibitors that behave as substrate mimics to halt viral cDNA synthesis. The non-nucleoside reverse transcriptase inhibitors,

nevirapine, delavirdine, efavirenz and etravirine inhibit the synthesis of viral cDNA via a non-competitive (or uncompetitive) mechanism. Enfuvirtide and maraviroc inhibit the entry of the virus into the host cell. Used alone these drugs are effective in reducing viral replication. The effect is only temporary as the virus readily develops 5 resistance to all known agents. However, combination therapy has proven very effective at both reducing virus and suppressing the emergence of resistance in a number of patients. In the US, where combination therapy is widely available, the number of HIV-related deaths has declined (Palella, F. J.; Delany, K. M.; Moorman, A. C.; Loveless, M. O.; Furher, J.; Satten, G. A.; Aschman, D. J.; Holmberg, S. D. N. 10 *Engl. J. Med.* 1998, 338, 853-860).

Unfortunately, not all patients are responsive and a large number fail this therapy. In fact, approximately 30-50% of patients ultimately fail combination therapy. Treatment failure in most cases is caused by the emergence of viral 15 resistance. Viral resistance in turn is caused by the rapid turnover of HIV-1 during the course of infection combined with a high viral mutation rate. Under these circumstances incomplete viral suppression caused by insufficient drug potency, poor compliance to the complicated drug regimen as well as intrinsic pharmacological barriers to exposure provides fertile ground for resistance to emerge. More 20 disturbing are recent findings which suggest that low-level replication continues even when viral plasma levels have dropped below detectable levels (< 50 copies/ml) (Carpenter, C. C.; Cooper, D. A.; Fischl, M. A.; Gatell, J. M.; Gazzard, B. G.; Hammer, S. M.; Hirsch, M. S.; Jacobsen, D. M.; Katzenstein, D. A.; Montaner, J. S.; Richman, D. D.; Saag, M. S.; Schechter, M.; Schooley, R. T.; Thompson, M. A.; 25 Vella, S.; Yeni, P. G.; Volberding, P. A. *JAMA* 2000, 283, 381-390). Clearly, there is a need for new antiviral agents, preferably targeting other viral enzymes to reduce the rate of resistance and suppress viral replication even further.

HIV expresses three enzymes, reverse transcriptase, an aspartyl protease, and 30 integrase. All three are targets for treating AIDS and HIV infection. HIV integrase is a component of the pre-integration complex of the virus that is assembled in the cell shortly after infection (Chiu, T.K.; Davies, D.R. *Curr. Top. Med. Chem.* 2004, 4, 965-

977). This enzyme catalyzes the integration of proviral DNA into the host genome and is absolutely required for viral infectivity. Early experiments showed that mutating the active site of integrase within a proviral clone produces virus unable to replicate due to its inability to insert into the host chromosome (Englund, G.;
5 Theodore, T.S.; Freed, E.O.; Engleman, A.; Martin, M.A. *J. Virol.* 1995, 69, 3216-3219). Selective HIV integrase inhibitors have been shown to possess effective anti-HIV activity in cell culture (Hazuda, D.J.; Felock, P.; Witmer, M.; Wolfe, A; Stillmock, K.; Grobler, J.A.; Espeseth, A.; Gabryelski, L.; Schleif, W.; Blau, C.; Miller, M.D. *Science*, 2000, 287, 646-650), and it is clear that this class of inhibitors
10 is very effective as part of a combination regimen containing HIV inhibitors of different classes. An HIV integrase inhibitor, raltegravir (ISENTRESS®), has been approved for use in treatment experienced patients based upon 48 week trial results (Cooper, D.A.; Gatell, J.; Rockstroh, J.; Katlama, C.; Yeni, P.; Lazzarin, A.; Xu, X.; Isaacs, R.; Teppler, H.; Nguyen, B.Y. *15th Conference on Retroviruses and*
15 *Opportunistic Infections*, Boston, MA., Feb 3-6, 2008 Abst. 105LB: Evering, T.H.; Markowitz, M. *Drugs Today*, 2007, 43, 865-877). In addition, a second integrase inhibitor, elvitegravir (GS-9137), completed a successful Phase II trial in combination with ritonavir boosting in naive and treatment experienced patients (Zolopa, A. *14th Conference on Retroviruses and Opportunistic Infections*, Los Angeles, CA. Feb 25-28, 2007 Abst. 143LB). Thus, HIV-1 integrase is a promising
20 target for novel anti-HIV-1 therapeutics.


The invention provides technical advantages, for example, the compounds are novel and inhibit HIV integrase. Additionally, the compounds provide advantages for pharmaceutical uses, for example, with regard to one or more of their mechanism
25 of action, binding, inhibition efficacy, target selectivity, solubility, safety profiles, or bioavailability.

HIV integrase inhibitors have been disclosed in U.S. patents 7,176,196 and 7,173,022 and PCT publications WO 2004/58756 and WO 2004/58757.

DESCRIPTION OF THE INVENTION

The invention encompasses compounds of Formula I, including pharmaceutically acceptable salts, their pharmaceutical compositions, and their use in 5 inhibiting HIV integrase and treating those infected with HIV or AIDS.

One aspect of the invention are compounds of Formula I



10

I

wherein:

R¹ is (Ar¹)alkyl, alkyl, (cycloalkyl)alkyl, diphenylalkyl, phenoxyalkyl, (PhNH)alkyl, (methylpyrrolidinyl)alkyl, (imidazolyl)alkyl (valerolactamyl)alkyl, 15 (tetrahydrofuranyl)alkyl, ((fluoro)(methyl)pyridinyl)methyl, phenylcyclopropyl, or benzylpyrrolidinyl;

R² is hydrogen, alkyl, hydroxy or alkoxy;

20 R³ is hydrogen, halo, hydroxy, alkoxy, cyano, alkyl, cycloalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, alkoxy, haloalkoxy, N(R⁸)(R⁹), N(R⁶)COR⁷, N(R⁶)SO₂R⁷, N(R⁶)CO₂R⁷, N(R⁶)SO₂N(R⁸)(R⁹), CO₂R⁶, CON(R⁸)(R⁹), SOR⁷, SO₂R⁷, SO₂N(R⁸)(R⁹), PO(OR⁶)₂, R¹², or Ar²;

25 R⁴ is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or N(R⁶)(R⁶);

R⁵ is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or N(R⁶)(R⁶);

30

R^6 is hydrogen, alkyl, haloalkyl, or cycloalkyl;

R^7 is alkyl, haloalkyl, or cycloalkyl;

5 R^8 is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl;

R^9 is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl; or

N(R^8)(R^9) taken together is azetidinyl, pyrrolidinyl, (R^{10})-piperidinyl,

10 N-(R^{11})-piperazinyl, morpholinyl, thiomorpholinyl, or dioxothiazinyl;

R^{10} is hydrogen, alkyl, hydroxy, or hydroxyalkyl;


R^{11} is hydrogen, alkyl, cyclolkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl,

15 phenyl, pyridinyl, (methylimidazolyl)methyl, COR⁶, CO₂R⁶, (hydroxyalkyl)CO,

(alkoxyalkyl)CO, (tetrahydrofuranyl)CO, (methylisoxazolyl)CO, (thienyl)CO,

(furanyl)CO, (pyridinyl)CO, CON(R^8)(R^9), SO₂R⁷, SO₂N(R^8)(R^9),

(dimethylisoxazolyl)SO₂, ((carboethoxy)thienyl)SO₂, (methylimidazolyl)SO₂,



20

R^{12} is azetidinonyl, pyrrolidinonyl, valerolactamyl, caprolactamyl, maleimido, oxazolidinonyl, imidazolidinonyl, triazolonyl, N-(R^6)-dioxothiazolidinyl or dioxothiazinyl, and is substituted with 0-2 substituents selected from the group consisting of alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, and aminoalkyl;

25

R^{13} is hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, benzyloxycarbonyl, or alkylsulfonyl;


R^{14} is N(R^{15})(R^{15}), N(R^{15})(benzyloxycarbonyl), N(R^{15})(alkyloxycarbonyl),


30 N(R^{15})((hydroxyalkyl)oxycarbonyl), N(R^{15})((alkyloxy)alkyloxycarbonyl),

$N(R^{15})CO(R^{16})$, $N(R^{15})((CO(N(R^{15})_2)alkyl)$, $N(R^{15})((CO_2CO(N(R^{15})_2)alkyl)$,
 $N(R^{15})CO((N(R^{15})_2)alkyl)$, $N(R^{15})CO((CON(R^{15})(R^{15})alkyl)$, $N(R^{15})COCO_2(R^{15})$,
 $N(R^{15})CO((CO_2(R^{15})alkyl)$, $N(R^{15})COCON(R^{15})(R^{15})$,
 $N(R^{15})COCO((N(R^{15})(R^{15}))alkyl)$, $N(R^{15})CO(N(R^{15})_2)$, $N(R^{15})SO_2R^{16}$,
5 $N(R^{15})SO_2(N(R^{15})_2)$, or $N(R^{15})COAr^3$;

or R¹⁴ is hydroxy, alkoxy, (hydroxy)alkoxy, (alkoxy)alkoxy, (R¹⁶CO₂)alkoxy, or (PhCO₂)alkoxy;

10 or R^{14} is $N(R^{15})COCON(R^{15})(alkoxyalkyl)$, $N(R^{15})COCON(R^{15})(hydroxyalkyl)$,
 $N(R^{15})COCON(R^{15})((tetrahydrofuran-2-yl)methyl)$, $N(R^{15})COCON(R^{15})((N,N-$
 $dimethylpyrazol-1-yl)methyl)$, or $N(R^{15})CO((Ar^4)alkyl)$;







5 R^{15} is hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, haloalkyl, or cycloalkyl;

or $N(R^{15})(R^{15})$ taken together is azetidinyl, pyrrolidinyl, (R^{10}) -piperidinyl, $N(R^{11})$ -piperazinyl, morpholinyl, or thiomorpholinyl, and is substituted with 0-2 substituents selected from halo, hydroxy, alkoxy, alkyl, hydroxyalkyl, alkoxyalkyl, 10 hydroxyalkyloxyalkyl, acetamido, CO_2R^6 , and $CON(R^8)(R^9)$;

R^{16} is alkyl, haloalkyl, cycloalkyl, (cycloalkyl)alkyl, hydroxyalkyl, alkoxyalkyl, tetrahydropyranyl, or Ar^3 ;

15 R^{17} is hydrogen, alkyl, alkylCO, cycloalkylCO, alkyloxyCO, $CON(R^{15})(R^{15})$, $COCON(R^{15})(R^{15})$, $COAr_3$, alkylSO₂, cycloalkylSO₂, furanylSO₂, triazolylSO₂, or N-methylpyrrolylSO₂;





Ar² is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, 5 thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyrimidinyl, pyrazinyl, pyridinyl, or hydroxypyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, cyano, benzyl, alkyl, alkoxy, N(R⁸)(R⁹), CO₂R⁶, and CON(R⁸)(R⁹);


10 Ar³ is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, imidazopyridinyl, quinolinyl, isoquinolinyl, or phenyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, alkoxy, haloalkoxy, cyano, benzyl, alkyl, haloalkyl, N(R⁸)(R⁹), and N(R¹⁵)CO(R¹⁶);

15

Ar⁴ is pyrazolyl or triazolyl, and is substituted with 0-1 substituents selected from CO₂R⁶ and CON(R⁸)(R⁹);

X-Y-Z is

20



5 or a pharmaceutically acceptable salt thereof.

Another aspect of the invention is a compound of formula I wherein:

R^1 is $(Ar^1)alkyl$;

10

R^2 is hydrogen, alkyl, hydroxy or alkoxy;

R^3 is hydrogen, halo, hydroxy, cyano, alkyl, haloalkyl, alkoxy, haloalkoxy,
 $N(R^8)(R^9)$, $N(R^6)COR^7$, $N(R^6)SO_2R^7$, $N(R^6)CO_2R^7$, $N(R^6)SO_2N(R^8)(R^9)$, CO_2R^6 ,
15 $CON(R^8)(R^9)$, SOR^7 , SO_2R^7 , $SO_2N(R^8)(R^9)$, $PO(OR^6)_2$, R^{12} , or Ar^2 ;

R^4 is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or
 $N(R^6)(R^6)$;

20 R^5 is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or
 $N(R^6)(R^6)$;

R^6 is hydrogen, alkyl, haloalkyl, or cycloalkyl;

25 R^7 is alkyl, haloalkyl, or cycloalkyl;

R^8 is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl;

R^9 is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl; or

5 $N(R^8)(R^9)$ taken together is azetidinyl, pyrrolidinyl, (R^{10})-piperidinyl,
 N-(R^{11})-piperazinyl, morpholinyl, thiomorpholinyl, or dioxothiazinyl;

R^{10} is hydrogen, alkyl, hydroxy, or hydroxyalkyl;


10 R^{11} is hydrogen, alkyl, cycloalkyl, COR^6 , or CO_2R^6 ;

R^{12} is azetidinonyl, pyrrolidinonyl, valerolactamyl, caprolactamyl, maleimido,
 oxazolidinonyl, imidazolidinonyl, triazolonyl, $N-(R^6)$ -dioxothiazolidinyl or
 dioxothiazinyl, and is substituted with 0-2 substituents selected from the group
 15 consisting of alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, and aminoalkyl;

R^{13} is hydrogen, alkyl, alkylcarbonyl, alkoxycarbonyl, benzyloxycarbonyl, or
 alkylsulfonyl;

20 R^{14} is $N(R^{15})(R^{15})$, $N(R^{15})(benzyloxycarbonyl)$, $N(R^{15})(alkyloxycarbonyl)$,
 $N(R^{15})CO(R^{16})$, $N(R^{15})CO(alkyl(N(R^{15})_2))$, $N(R^{15})COCO_2(R^{15})$,
 $N(R^{15})COCON(R^{15})(R^{15})$, $N(R^{15})CO(N(R^{15})_2)$, $N(R^{15})SO_2R^{16}$, $N(R^{15})SO_2(N(R^{15})_2)$,
 or $N(R^{15})COAr^3$;

25 or R^{14} is hydroxy, alkoxy, (hydroxy)alkoxy, (alkoxy)alkoxy, $(R^{16}CO_2)$ alkoxy, or
 $(PhCO_2)$ alkoxy;




30 R^{15} is hydrogen, alkyl, haloalkyl, or cycloalkyl;

or N(R¹⁵)(R¹⁵) taken together is azetidinyl, pyrrolidinyl, (R¹⁰)-piperidinyl, N-(R¹¹)-piperazinyl, morpholinyl, or thiomorpholinyl;

R¹⁶ is alkyl, haloalkyl, or cycloalkyl;

5




10 Ar² is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyrimidinyl, pyrazinyl, pyridinyl, or hydroxypyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, cyano, benzyl, alkyl, alkoxy, N(R⁸)(R⁹), CO₂R⁶, and CON(R⁸)(R⁹);

15

Ar³ is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, or pyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, alkoxy, cyano, benzyl, and alkyl; and

20

X-Y-Z is



or a pharmaceutically acceptable salt thereof.

5

Another aspect of the invention is a compound of formula I where

- R¹ is (Ar¹)alkyl, alkyl, (cycloalkyl)alkyl, diphenylalkyl, phenoxyalkyl, (PhNH)alkyl, (methylpyrrolidinyl)alkyl, (imidazolyl)alkyl (valerolactamyl)alkyl,
- 10 (tetrahydrofuranyl)alkyl, ((fluoro)(methyl)pyridinyl)methyl, phenylcyclopropyl, or benzylpyrrolidinyl;
- R² is hydrogen;
- 15 R³ is hydrogen, halo, hydroxy, alkoxy, cyano, alkyl, cycloalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, alkoxy, haloalkoxy, N(R⁸)(R⁹), N(R⁶)COR⁷, N(R⁶)SO₂R⁷, N(R⁶)CO₂R⁷, N(R⁶)SO₂N(R⁸)(R⁹), CO₂R⁶, CON(R⁸)(R⁹), SO₂R⁷, SO₂N(R⁸)(R⁹), or Ar²;
- 20 R⁴ is hydrogen, halo, or alkyl;
- R⁵ is hydrogen, halo, or alkyl;
- R⁶ is hydrogen, alkyl, haloalkyl, or cycloalkyl;
- 25 R⁷ is alkyl, haloalkyl, or cycloalkyl;
- R⁸ is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl;

R^9 is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl; or

$N(R^8)(R^9)$ taken together is azetidinyl, pyrrolidinyl, (R^{10}) -piperidinyl,

$N-(R^{11})$ -piperazinyl, morpholinyl, thiomorpholinyl, or dioxothiazinyl;

5


R^{10} is hydrogen, alkyl, hydroxy, or hydroxyalkyl;

R^{11} is hydrogen, alkyl, cycloalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, phenyl, pyridinyl, (methylimidazolyl)methyl, COR^6 , CO_2R^6 , $(hydroxyalkyl)CO$,

10 $(alkoxyalkyl)CO$, $(tetrahydrofuran-2-yl)CO$, $(methylisoxazolyl)CO$, $(thienyl)CO$,

$(furanyl)CO$, $(pyridinyl)CO$, $CON(R^8)(R^9)$, SO_2R^7 , $SO_2N(R^8)(R^9)$,

$(dimethylisoxazolyl)SO_2$, $((carboethoxy)thienyl)SO_2$, $(methylimidazolyl)SO_2$,



15 R^{13} is hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, benzyloxycarbonyl, or alkylsulfonyl;

R^{14} is $N(R^{15})(R^{15})$, $N(R^{15})(benzyloxycarbonyl)$, $N(R^{15})(alkyloxycarbonyl)$,

$N(R^{15})((hydroxyalkyl)oxycarbonyl)$, $N(R^{15})((alkyloxy)alkyloxycarbonyl)$,

20 $N(R^{15})CO(R^{16})$, $N(R^{15})((CO(N(R^{15})_2)alkyl)$, $N(R^{15})((CO_2CO(N(R^{15})_2)alkyl)$,

$N(R^{15})CO((N(R^{15})_2)alkyl)$, $N(R^{15})CO((CON(R^{15})(R^{15})alkyl)$, $N(R^{15})COCO_2(R^{15})$,

$N(R^{15})CO((CO_2(R^{15})alkyl)$, $N(R^{15})COCON(R^{15})(R^{15})$,


$N(R^{15})COCO((N(R^{15})(R^{15}))alkyl)$, $N(R^{15})CO(N(R^{15})_2)$, $N(R^{15})SO_2R^{16}$,

$N(R^{15})SO_2(N(R^{15})_2)$, or $N(R^{15})COAr^3$;

25

or R^{14} is hydroxy, alkoxy, (hydroxy)alkoxy, (alkoxy)alkoxy, $(R^{16}CO_2)alkoxy$, or $(PhCO_2)alkoxy$;

or R¹⁴ is N(R¹⁵)COCON(R¹⁵)(alkoxyalkyl), N(R¹⁵)COCON(R¹⁵)(hydroxyalkyl), N(R¹⁵)COCON(R¹⁵)((tetrahydrofuran-2-yl)methyl), N(R¹⁵)COCON(R¹⁵)((N,N-dimethylpyrazol-1-yl)methyl), or N(R¹⁵)CO((Ar⁴)alkyl);




R^{15} is hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, haloalkyl, or cycloalkyl;

or $N(R^{15})(R^{15})$ taken together is azetidinyl, pyrrolidinyl, (R^{10})-piperidinyl, $N-(R^{11})$ -piperazinyl, morpholinyl, or thiomorpholinyl, and is substituted with 0-2 substituents selected from halo, hydroxy, alkoxy, alkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkyloxyalkyl, acetamido, CO_2R^6 , and $CON(R^8)(R^9)$;

R^{16} is alkyl, haloalkyl, cycloalkyl, (cycloalkyl)alkyl, hydroxyalkyl, alkoxyalkyl, tetrahydropyranyl, or Ar^3 ;

10

R^{17} is hydrogen, alkyl, $alkylCO$, $cycloalkylCO$, $alkyloxyCO$, $CON(R^{15})(R^{15})$, $COCON(R^{15})(R^{15})$, $COAr_3$, $alkylSO_2$, $cycloalkylSO_2$, $furanylSO_2$, $triazolylSO_2$, or N -methylpyrrolyl SO_2 ;




20 Ar^2 is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyrimidinyl, pyrazinyl, pyridinyl, or hydroxypyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, cyano, benzyl, alkyl, alkoxy, $N(R^8)(R^9)$, CO_2R^6 , and $CON(R^8)(R^9)$;

25 Ar^3 is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, imidazopyridinyl, quinolinyl, isoquinolinyl, or phenyl, and is substituted

with 0-3 substituents selected from the group consisting of oxo, halo, alkoxy, haloalkoxy, cyano, benzyl, alkyl, haloalkyl, $N(R^8)(R^9)$, and $N(R^{15})CO(R^{16})$;

Ar⁴ is pyrazolyl or triazolyl, and is substituted with 0-1 substituents selected from
 5 CO_2R^6 and $CON(R^8)(R^9)$; and

X-Y-Z is




or a pharmaceutically acceptable salt thereof.


15 Another aspect of the invention is a compound of Formula I where R¹ is (Ar¹)alkyl.

Another aspect of the invention is a compound of Formula I where R¹ is (Ar¹)alkyl and R₂ is hydrogen.

20 Another aspect of the invention is a compound of Formula I where R¹ is



Another aspect of the invention is a compound of Formula I where R¹ is



and R³ is other than hydrogen or halo.

Another aspect of the invention is a compound of Formula I where R² is hydrogen.

5

Another aspect of the invention is a compound of Formula I where R³ is R¹².

Another aspect of the invention is a compound of Formula I where R³ is Ar².

- 10 Another aspect of the invention is a compound of Formula I where R³ is CON(R⁸)(R⁹), SO₂N(R⁸)(R⁹), azetidinonyl, pyrrolidinonyl, valerolactamyl, caprolactamyl, oxazolidinonyl, imidazolidinonyl, dioxothiazinyl, N-(R⁶)-dioxothiazolidinyl, or triazolyl where triazolyl is substituted with 0-2 alkyl or oxo substituents.

15

Another aspect of the invention is a compound of Formula I where R³ is halo, CON(R⁸)(R⁹), oxazolidinonyl, or triazolyl where triazolyl is substituted with 0-2 alkyl or oxo substituents.

- 20 Another aspect of the invention is a compound of Formula I where R¹⁴ is N(R¹⁵)(R¹⁵), N(R¹⁵)(benzyloxycarbonyl), N(R¹⁵)(alkyloxycarbonyl), N(R¹⁵)((hydroxyalkyl)oxycarbonyl), N(R¹⁵)((alkyloxy)alkyloxycarbonyl), N(R¹⁵)CO(R¹⁶), N(R¹⁵)((CO(N(R¹⁵)₂)alkyl), N(R¹⁵)((CO₂CO(N(R¹⁵)₂)alkyl), N(R¹⁵)CO((N(R¹⁵)₂)alkyl), N(R¹⁵)CO((CON(R¹⁵)(R¹⁵)alkyl), N(R¹⁵)COCO₂(R¹⁵),

- 25 N(R¹⁵)CO((CO₂(R¹⁵)alkyl), N(R¹⁵)COCON(R¹⁵)(R¹⁵), N(R¹⁵)COCO((N(R¹⁵)(R¹⁵))alkyl), N(R¹⁵)CO(N(R¹⁵)₂), N(R¹⁵)SO₂R¹⁶, N(R¹⁵)SO₂(N(R¹⁵)₂), or N(R¹⁵)COAr³.

Another aspect of the invention is a compound of Formula I where R¹⁴ is

- 30 N(R¹⁵)COCON(R¹⁵)(R¹⁵).


Another aspect of the invention is a compound of Formula I where Ar² is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, or pyrrolyl, and is substituted with 0-2 substituents selected from the group consisting of halo and alkyl.

5

Another aspect of the invention is a compound of Formula I where Ar³ is oxadiazolyl, pyrazolyl, or isoxazolyl.

10



Another aspect of the invention is a compound of Formula I where X-Y-Z is




For a compound of Formula I, the scope of any instance of a variable substituent, including R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, Ar¹, 15 Ar², Ar³, Ar⁴, and X-Y-Z, can be used independently with the scope of any other instance of a variable substituent. As such, the invention includes combinations of the different aspects.

Unless specified otherwise, these terms have the following meanings. “Alkyl” 20 means a straight or branched alkyl group composed of 1 to 6 carbons. “Alkenyl” means a straight or branched alkyl group composed of 2 to 6 carbons with at least one double bond. “Alkynyl” means a straight or branched alkyl group composed of 2 to 6 carbons with at least one triple bond. “Cycloalkyl” means a monocyclic ring system composed of 3 to 7 carbons. “Haloalkyl” and “haloalkoxy” include all 25 halogenated isomers from monohalo to perhalo. Terms with a hydrocarbon moiety (e.g. alkoxy) include straight and branched isomers for the hydrocarbon portion. Parenthetic and multiparenthetical terms are intended to clarify bonding relationships to those skilled in the art. For example, a term such as ((R)alkyl) means an alkyl substituent further substituted with the substituent R.


30



5 “Dioxothiazinyl” means .


The invention includes all pharmaceutically acceptable salt forms of the compounds. Pharmaceutically acceptable salts are those in which the counter ions do not contribute significantly to the physiological activity or toxicity of the compounds and as such function as pharmacological equivalents. These salts can be made according to common organic techniques employing commercially available reagents. Some anionic salt forms include acetate, acistrate, besylate, bromide, chloride, citrate, fumarate, glucouronate, hydrobromide, hydrochloride, hydroiodide, iodide, lactate, maleate, mesylate, nitrate, pamoate, phosphate, succinate, sulfate, tartrate, tosylate, and xinofoate. Some cationic salt forms include ammonium, aluminum, benzathine, bismuth, calcium, choline, diethylamine, diethanolamine, lithium, magnesium, meglumine, 4-phenylcyclohexylamine, piperazine, potassium, sodium, tromethamine, and zinc.

20 Some of the compounds of the invention exist in stereoisomeric forms. The invention includes all stereoisomeric forms of the compounds including enantiomers and diastereromers. Methods of making and separating stereoisomers are known in the art. One example of a stereoisomeric form is shown below.



The invention includes all tautomeric forms of the compounds. An example of a tautomeric pair is shown below.

5




Synthetic Methods

- 10 The compounds may be made by methods known in the art including those described below and including variations within the skill of the art. Some reagents and intermediates are known in the art. Other reagents and intermediates can be made by methods known in the art using readily available materials. The following methods are for illustrative purposes and are not intended to limit the scope of the invention. It will be appreciated by those skilled in the art that there are a number of methods available for the synthesis of an appropriately substituted pyrimidinone heterocycles useful for making the compounds of the current invention and that their synthesis is not limited to the methods provided in the following examples. For example, one such method is shown in Scheme I and is similar to that reported in
- 15 Journal of Heterocyclic Chemistry (1979), 16(7), 1423-4. In this scheme an alkyl- or arylnitrile 1-1 is combined with hydroxylamine to form intermediate 1-2 which in turn is reacted with dialkyl-butynoate 1-3 to form intermediate 1-4. At high temperatures, such as that of refluxing xylenes, intermediate 1-4 undergoes a rearrangement reaction to yield the 5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-
- 20 carboxylate heterocycle 1-5 which can serve as an advanced intermediate for synthesizing some of the compounds of this invention. The variables (e.g. numbered
- 25

“R” substituents) used to describe the synthesis of the compounds are intended only to illustrate how to make the compounds and are not to be confused with variables used in the claims or in other sections of the specification.

- 5 Abbreviations used in the schemes generally follow conventions used in the art. Chemical abbreviations used in the specification and examples are defined as follows: "NaHMDS" for sodium bis(trimethylsilyl)amide; "DMF" for N,N-dimethylformamide; "MeOH" for methanol; "NBS" for N-bromosuccinimide; "Ar" for aryl; "TFA" for trifluoroacetic acid; "LAH" for lithium aluminum hydride; 10 "BOC", "DMSO" for dimethylsulfoxide; "h" for hours; "rt" for room temperature or retention time (context will dictate); "min" for minutes; "EtOAc" for ethyl acetate; "THF" for tetrahydrofuran; "EDTA" for ethylenediaminetetraacetic acid; "Et₂O" for diethyl ether; "DMAP" for 4-dimethylaminopyridine; "DCE" for 1,2-dichloroethane; "ACN" for acetonitrile; "DME" for 1,2-dimethoxyethane; "HOBr" for 1-hydroxybenzotriazole hydrate; "DIEA" for diisopropylethylamine, "Nf" for CF₃(CF₂)₃SO₂-; and "TMOF" for trimethylorthoformate.
- 15

Scheme I




20

- Scheme II demonstrates an alternative method for providing the 5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate heterocyclic system which is adapted from the procedure in Inorganic Chemistry (2001), 40(26), 6746-6756. In this scheme, a substituted derivative of glycolic acid, 2-1 is reacted with a diester of oxalic acid 25 under Claisen condensation conditions (Org. React. 1, 266-322 (1942); Modern Synthetic Reactions (W. A. Benjamin, Menlo Park, California, 2nd ed., 1972) pp 734-746, J. Chem. Educ. 56, 721 (1979); J. Am. Chem. Soc. 103, 1338 (1981); Comp. Org. Syn. 2, 795-805 (1991)) to provide intermediate 2-2. This intermediate

can be combined with an amidine, 2-3, under heating to yield the 5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate heterocycle, 2-4.

Scheme II




5

In Scheme III, peracylated glycoside 3-1 can be converted to intermediate 3-2 using synthetic procedures known to those skilled in the art, for example the method described in Carbohydrate Research (1982), 108(2), 229-35. Following this, a 10 method similar to that described in Journal of Organic Chemistry (1988), 53(16), 3723-31 can be used to eliminate the acetoxy group at C-4 yielding a mixture of compounds, 3-3 and 3-4, which upon hydrogenation produce intermediate 3-5. The acetyl group of 3-5 can be removed using common conditions familiar to those skilled in the art. The synthesis of the 5-hydroxy-6-oxo-1,6-dihdropyrimidine-4- 15 carboxylate ring of 3-7 can be assembled as described in Scheme I. Intermediate 3-7 is then treated with \geq 3 equivalents of methanesulfonyl chloride and diisopropyl-ethylamine to yield 3-8 which when treated with a base such as K_2CO_3 produces the bridged tricyclic-fused, 5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate template, 3-8. This intermediate can be converted to the corresponding amide, 3-10, 20 using standard amide bond forming procedures familiar to those skilled in the art. For example, in one method intermediate 3-9 can be combined with the amine, 3-11 and a base such as triethylamine and the mixture heated to effect aminolysis of the ester bond. In another method the hydroxyl group of 3-9 is modified with a protecting group such as benzyl, or a group such as those mentioned in Greene's Protective 25 Groups in Organic Synthesis, 4th Edition, Peter G. M. Wuts, Theodora W. Greene, John Wiley & Sons, Inc. 2007, Hoboken New Jersey. Following this the ester is hydrolyzed to the corresponding carboxylic acid and this is coupled with the amine 3-


11 using an amide bond forming reagent such as N,N'-dicyclohexylcarbodiimide, O-benzotriazol-1-yl-N,N,N',N'-bis(pentamethylene)uronium hexafluorophosphate, 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide methiodide, bromotripyrrolidinophosphonium hexafluorophosphate or other amide bond forming
5 reagents familiar to those skilled in the art. Following this the protecting is removed to produce 3-10.

Scheme III



In Scheme IV, intermediate 4-1 (Tetrahedron Letters (1994), 35(26), 4515-18 and references cited therein) is reacted with benzyl bromide, in the presence of a base such as K₂CO₃, to form the N-benzyl derivative 4-2 (Synth. Commun. 1997, 27 (1), 69-71). It will be appreciated by those skilled in the art that the benzyl group attached to 4-2 serves as a protecting group and that other nitrogen protecting groups could be used in its place. The diester, 4-2 can be treated with lithium aluminum hydride to form the bis-alcohol 4-3. Upon treatment of this intermediate with 1 molar equivalent of tert-butyldiphenylsilyl chloride and sodium hydride the protected intermediate 4-3 is formed. Treatment of this intermediate using Swern reaction conditions (Tetrahedron 34, 1651 (1978)), affords intermediate 4-5. It will be appreciated by those skilled in the art that a number of procedures are available for the conversion of an alcohol to an aldehyde, such as those described in Comprehensive Organic Transformations: A Guide to Functional Group Preparations, Second Edition, Richard C. Larock, John Wiley and Sons, 1999, New York, NY. The aldehyde 4-5 can be condensed with hydroxylamide to form intermediate 4-6, which is then transformed to the nitrile 4-7. One skilled in the art will appreciate that there are a number of methods available for transforming intermediates such as 4-6 into 4-7. For example Comprehensive Organic Functional Group Transformations II, Alan R. Katritzky, Otho Meth-Cohn, and Charles W. Rees, Elsevier, 1995. Upon treatment of this intermediate with hydroxylamine followed by dialkyl-butynoate, intermediate 4-9 is produced. Formation of the 5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate heterocycle and cyclization to form the bridged tricyclic fused ring system can be carried out as described in the previous schemes to afford intermediate 4-11. The ethylene-dicarboxylate group attached to the nitrogen atom of 4-11 can be removed under acidic conditions. The nitrogen is next protected as the corresponding benzylcarbamate (Cbz) using benzylchloroformate and diisopropylidethylamine to yield 4-13. Amidation to form 4-14 can be accomplished as described previously and following this, removal of the Cbz-protecting group produces 4-15.


Scheme IV



In Scheme V, diol 5-1 (*Tetrahedron Letters* (1971), (39), 3587-9; *Journal of Organic Chemistry* (1975), 40(17), 2488-95) is treated with tert-butyl dimethylsilylchloride in the presence of sodium hydride to deliver intermediate 5-2. Oxidation of this compound under Swern conditions, similar to that described above, yields aldehyde 5-3. Intermediate 5-4 can be synthesized by reacting 5-3 with


methylamine and potassium cyanide in methanol. Following protection of the nitrogen group as the corresponding Cbz-derivative, the tert-butyldisilyl protecting group can be removed to provide intermediate 5-6. Formation of the 5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate heterocycle and cyclization to form the bridged tricyclic fused ring system, as described in the previous schemes, affords intermediate 5-9.

Scheme V



Scheme VI describes yet another method for the synthesis of compounds of this invention. In this method, intermediate 6-1 is converted to the corresponding diester, 6-3, via the anhydride 6-2. Reduction of the ester groups to provide intermediate 6-4, can be accomplished using procedures familiar to those skilled in the art. The subsequent synthetic steps from intermediate 6-4 to the final compound 6-10 can be accomplished using methods similar to those described in the schemes above.


Scheme VI



10

In Scheme VII, intermediate 5-3 is reacted with hydroxylamine to form, 7-1 and converted into the corresponding nitrile 7-2, as described above in Scheme IV for the synthesis of intermediate 4-7. The synthesis of compound 7-8 can be 15 accomplished using methods similar to those described in the schemes above.

Scheme VII




Treatment of intermediate 8-1, in Scheme VIII, under Strecker reaction

5 conditions (Greenstein, M. Winitz, Chemistry of the Amino Acids vol. 3 (New York, 1961) pp 698-700, G. C. Barrett, Chemistry and Biochemistry of the Amino Acids (Chapman and Hall, New York, 1985) pp 251, 261) provides the amino nitrile 8-2. After protection of the amino-group as the corresponding Cbz-derivative, the 5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate heterocycle is formed by
10 sequentially treating the nitrile with, hydroxylamine and diethyl but-2-ynedioate, then heating in xylenes. The cyclic ketal protecting group can be hydrolyzed under acidic conditions to furnish the ketone 8-5 which is then reduced with NaBH₄ to provide intermediate 8-6. Cyclization to form intermediate 8-7, can be achieved by first treating this intermediate with methane sulfonyl chloride/triethylamine, to effect
15 mesylation of the hydroxyl group followed by treatment with cesium carbonate. The Cbz-group is hydrolytically removed under standard conditions resulting in intermediate 8-8. Formation of the corresponding 4-fluorobenzylamide can be accomplished using the methods described above. Intermediate 8-9 is then treated


with an activated carboxylic acid, for example and acid chloride, or a carboxylic acid in the presence of an amide bond forming reagent to yield compound 8-10.

Scheme VIII



Scheme IX illustrates a method for the synthesis of intermediate 9-10. In the first step of the sequence compound 9-1 is treated with ethane 1,2-diol and toluene sulfonic acid which yields intermediate 9-2. The carboxylic ester group of 9-2 is reduced using lithium aluminum hydride affording 9-3. Treatment of this 5 intermediate with SOCl_2 produces the chloride derivative 9-4. It will be appreciated by those skilled in the art that numerous methods exist for the substitution of an alcohol group with chloride some of which can be found in Comprehensive Organic Transformations: A Guide to Functional Group Preparations, Second Edition, Richard C. Larock, John Wiley and Sons, 1999, New York, NY. Hydrolysis of the 10 cyclic ketal group of intermediate 9-4 provides a ketone, 9-5, which can be taken on as described in the previous schemes, to the 5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate derivative 9-9. Under basic conditions the tricyclic ring system is formed via nucleophilic displacement of the chloride group. Compound 9-10 can serve as an advanced intermediate in the synthesis of compounds of this invention.

Scheme IX




Another method for the synthesis of intermediate 9-10 is shown in Scheme X.

- 5 In Scheme X, compound 10-1 can be prepared according to general procedure described in Journal of Organic Chemistry 1997, 62, 5284-5292 from commercially available ethyl 4-oxocyclohexanecarboxylate. A Strecker reaction, like that described in the schemes above produces the amino-nitrile intermediate 10-2 which is carried on to the N-protected intermediate 10-3. Treatment of this compound with

hydroxylamine, followed by diethyl but-2-ynedioate and heating in xylenes affords 10-4. The final cyclization reaction can be carried out as before, via activation of the alcohol with methane sulfonyl chloride followed by ring closing nucleophilic attack by the pyrimidinone nitrogen-group.


5

Scheme X




- In Scheme XI intermediate 9-10 is treated with a substituted benzylic amine
- 10 in the presence of triethylamine, with heating, to form the corresponding benzylamide 11-1. The Cbz protecting of 11-1 can be removed using standard hydrogenation conditions (e.g. H₂ and Pd/C) or other methods commonly known to one skilled in the art. In the final step amide, urea, carbamate, sulfonamide and related analogues can be synthesized directly from 11-3 under appropriate conditions.
- 15 For example amides can be synthesized by treating 11-3 with a carboxylic acid chloride or by reacting it with a carboxylic acid in the presence of an amide bond forming reagent such as HATU. Other analogues are available by treating 11-3 with isocyantes, chlorosulfonates or chloroformates, to name a few. The analogues are treated with MeNH₂ in the event that the C3-hydroxyl group of the pyrimidinone ring
- 20 has been modified.

Scheme XI



Compounds of this invention can also be synthesized according to the method illustrated in Scheme XII. In this Scheme 11-2 can be synthesized as before. This intermediate is then reacted with 2-chloro-2-oxoacetate in the presence of triethylamine to deliver 12-1. Following this 12-1 is heated with an amine to affect the formation of 12-2. In certain cases intermediate 12-1 need not be isolated but can be carried on to the 12-2 forming step immediately after work-up.


Scheme XII



In Scheme XIII, intermediate 13-1 can be coupled with intermediate 13-2 under standard amide-bond forming conditions, for example using the HATU as the coupling reagent in the presence of diisopropyl ethylamine and DMAP. Intermediate 13-3 is then treated with a benzylamine, with heating, as described in the above schemes to produce 13-4.


10

Scheme XIII



In the method shown in Scheme XIV, intermediate 9-10 is benzoylated by treating it with benzoic acid anhydride and pyridine to form intermediate 14-1. The Cbz protecting group can then be removed by hydrogenolysis to yield intermediate 14-2. Intermediate 14-2 can be coupled with acid 14-3 and the resulting product, 14-4 reacted with a substituted benzyl or substituted phenylpropylamine to provide 14-5.


Scheme XIV



10

Scheme XV illustrates another method for the synthesis of compounds of this invention. Intermediate 15-1, prepared using methods similar to those described previously, is heated with a substituted piperazine to provide 15-2.


Scheme XV



As shown in Scheme XVI, when 15-1 is reacted with piperazine as described
5 in the previous scheme, intermediate 16-1 is produced. The unsubstituted nitrogen of
the piperazine ring can be modified by the addition of R to form 16-2. R can be a
substituted alkyl group, an acyl group, a sulfonyl group, a urea or related
functionality.

10


Scheme XVI



As shown in Scheme XVII, the bridgehead nitrogen of intermediate 17-1 can be transformed into the corresponding dimethylated analogue, 17-2 via reductive amination, using for example formaldehyde and sodium triacetoxy borohydride. In a similar fashion intermediate 17-3 can be synthesized by sequential introduction of the 5 benzyl group via reductive amination (benzaldehyde, NaCNBH₃) followed by introducing of the methyl group in the same way. The benzyl group is then cleaved under reductive conditions, H₂ and Pd/C, to provide 17-4. The nitrogen atom 17-4 can be modified using methods similar to those described above.


10

Scheme XVII




Additional methods for the synthesis of compounds of this invention are illustrated in Schemes XVIII and XIX.

Scheme XVIII




Scheme XIX




The method described in Scheme XX for the synthesis of intermediate 20-2 is
5 similar to that described for intermediate 18-1 in Scheme XVIII except acetraldehyde
is used in place of formaldehyde. The subsequent transformations rely on methods
described above.

Scheme XX




Schemes XXI through XXV illustrate additional methods for synthesizing
5 compounds of this invention.

Scheme XXI




Scheme XXII




5


Scheme XXIII



Scheme XXIV




Scheme XXV



Scheme XXVI illustrates another method for the synthesis of compounds of this invention. In this method 3-oxocyclopentane carboxylic acid (26-1) is converted into the ketal-ester derived intermediate 26-2 (Tetrahedron (1977), 33, 1113) by stirring in methanol and acetyl chloride. Following this the ester group can be reduced, using LAH, to the corresponding alcohol, 26-3. The ketal group of intermediate 36-3 is removed under acidic conditions (p-toluenesulfonic acid) and


the 5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate derivative 26-9 synthesized using the same reaction sequence used in the previous schemes. Formation of the final ring can be carried out by first treating 26-9 with methane sulfonyl chloride and triethylamine followed by a second step involving treatment 5 with potassium carbonate. The product of this sequence, 26-10 can be esterified to form 26-11 using benzoic acid anhydride. Further transformation of this intermediate using the methods described previously provides, 26-12, 26-13 and 26-14.

Scheme XXVI



A method for the synthesis of advanced intermediate 27-11 is shown in Scheme XXVII. Cyclopentenone, 27-1 can be converted to 27-2 (*Tetrahedron Lett.* 2005, 46, 6875-6878) by the nucleophilic addition of dimethylmalonate using 2,3,4,6,7,8-hexahydro-1H-pyrimido[1,2-a]pyrimidine as a base. Those skilled in the art will recognize this synthetic transformation as a Michael reaction which can be carried out under a number of different conditions in addition to those shown here as illustrated in; E. D. Bergmann et al., *Org. React.* 10, 179-555 (1959); H. O. House, *Modern Synthetic Reactions* (W. A. Benjamin, Menlo Park, California, 2nd ed., 1972) pp 595-623; M. E. Jung, *Comp. Org. Syn.* 4, 1-67 (1991). In the next step, a methylcarboxylate group of 27-2 is removed at elevated temperature in the presence of dimethyl 2-(3-oxocyclopentyl)malonate to yield intermediate 27-3. This intermediate is then converted to the corresponding dimethyl ketal and the methylcarboxylate group reduced to the corresponding alcohol using LiBH₄ to provide 27-5 (*J. Org. Chem.*, 1986, 51 (21), 4000-4005). The ketal group is then hydrolyzed, yielding 27-6, then the pyrimidinone derivative, 27-10, synthesized in the same manner as that described in the above schemes wherein after formation of the amino nitrile, under Strecker reaction conditions, the amino group is protected with Cbz and the nitrile treated with hydroxylamine then diethyl but-2-ynedioate. The bridged bicyclic portion of 27-11 can be formed by treating intermediate 27-10 with methane sulfonyl chloride/triethylamine followed by treatment with a base such as K₂CO₃. Compound 27-11 can serve as an advanced intermediate for the synthesis of compounds of this invention as illustrated in Scheme XVIII.


Scheme XXVII




Compounds of this invention can also be synthesized according to Scheme

5 XVIII through XXXIV.


Scheme XXVIII




Scheme XXIX




Scheme XXX




Scheme XXXI




Scheme XXXII



Scheme XXXIII



Scheme XXXIV



Biological Methods

5

HIV-Integrase Inhibition Activity. To evaluate *in-vitro* activity against HIV-integrase, 5 pmole of biotin labeled substrate DNA was bound to 100 µg of Streptavidin coated PVT SPA beads (Amersham Pharmacia Biotech). Recombinant integrase (0.26 ng) was incubated with the beads for 90 min at 37 °C. Unbound enzyme was removed by washing the complex followed by addition of inhibitors and 0.1 fmol of P33 labeled target DNA. The reaction was stopped by adding EDTA to a

final concentration of 10 mM. Samples were counted in TopCountNXT (Packard) and the CPM was used as a measure of integration. The reaction condition was as described in A. Engelman and R. Craigie , *J. Virol.* 69, 5908-5911 (1995). The sequences of substrate and target DNA were described in *Nucleic Acid Research* 22,1121-1122 (1994). Results are shown in the Table 1. Activity equal to A refers to a compound having $IC_{50} = < 10$ nM while B and C denote compounds having $IC_{50} = < 10$ nM and $IC_{50} < 100$ nM and $IC_{50} > 100$ nM respectively.

Table 2.

Compound	in vitro activity
Example 6	B
Example 7	B
Example 8	A
Example 9	C
Example 11	B
Example 12	A
Example 13	A
Example 14	B
Example 15	A
Example 16	A
Example 17	A
Example 18	A
Example 19	C
Example 20	A
Example 21	A
Example 22	B
Example 23	A
Example 24	A
Example 25	A
Example 26	A
Example 27	B

Compound	in vitro activity
Example 28	A
Example 29	B
Example 30	B
Example 32	A
Example 33	A
Example 34	A
Example 35	A
Example 36	A
Example 37	B
Example 38	A
Example 39	A
Example 40	A
Example 41	A
Example 42	A
Example 43	A
Example 45	A
Example 46	A
Example 47	A
Example 48	A
Example 49	A
Example 50	A
Example 51	A
Example 52	A
Example 53	A
Example 54	A
Example 56	A
Example 57	A
Example 58	A
Example 59	A
Example 60	A

Compound	in vitro activity
Example 61	A
Example 62	A
Example 63	A
Example 64	A
Example 65	A
Example 66	A
Example 67	A
Example 68	A
Example 69	A
Example 70	A
Example 71	A
Example 72	A
Example 73	A
Example 74	A
Example 75	A
Example 76	A
Example 77	A
Example 78	A
Example 79	A
Example 80	A
Example 81	A
Example 82	A
Example 83	A
Example 84	A
Example 85	A
Example 86	A
Example 87	A
Example 88	A
Example 89	A
Example 90	A

Compound	in vitro activity
Example 91	A
Example 92	A
Example 93	A
Example 94	A
Example 95	A
Example 96	A
Example 97	A
Example 98	A
Example 99	A
Example 100	A
Example 101	A
Example 102	A
Example 103	A
Example 104	A
Example 105	A
Example 106	B
Example 107	B
Example 108	B
Example 109	B
Example 111	A
Example 112	A
Example 113	A
Example 114	A
Example 115	A
Example 116	A
Example 117	B
Example 118	A
Example 119	A
Example 120	A
Example 121	A

Compound	in vitro activity
Example 122	A
Example 123	B
Example 124	A
Example 125	A
Example 126	A
Example 127	B
Example 128	A
Example 129	A
Example 130	A
Example 131	B
Example 132	A
Example 133	A
Example 134	A
Example 135	A
Example 136	A
Example 137	A
Example 139	A
Example 140	A
Example 141	A
Example 142	A
Example 143	A
Example 144	A
Example 145	A
Example 146	A
Example 147	A
Example 148	A
Example 149	A
Example 150	C
Example 153	A
Example 154	A

Compound	in vitro activity
Example 155	A
Example 156	A
Example 157	A
Example 158	A
Example 159	A
Example 160	A
Example 161	A
Example 162	A
Example 163	A
Example 164	A
Example 165	A
Example 166	A
Example 167	A
Example 168	A
Example 169	A
Example 170	A
Example 171	A
Example 172	A
Example 173	A
Example 175	A
Example 176	B
Example 180	A
Example 181	B
Example 182	C
Example 183	B
Example 184	B
Example 185	C
Example 186	A
Example 187	A
Example 188	A

Compound	in vitro activity
Example 189	A
Example 190	B
Example 191	A
Example 192	A
Example 193	A
Example 194	A
Example 195	B
Example 196	A
Example 199	A
Example 200	A
Example 201	A
Example 202	A
Example 203	A
Example 204	A
Example 205	A
Example 206	A
Example 207	A
Example 208	A
Example 209	A
Example 210	A
Example 211	A
Example 212	A
Example 213	A
Example 214	A
Example 215	A
Example 216	A
Example 217	A
Example 218	A
Example 219	B
Example 220	B

Compound	in vitro activity
Example 221	A
Example 222	A
Example 225	A
Example 226	B
Example 234	A
Example 235	A
Example 236	B
Example 237	A
Example 238	A
Example 239	A
Example 240	A
Example 241	A
Example 242	A
Example 243	B
Example 244	B
Example 245	A
Example 246	A
Example 247	A
Example 248	A
Example 249	A
Example 250	A
Example 251	A
Example 252	B
Example 253	B
Example 254	A
Example 255	A
Example 256	A
Example 257	B
Example 258	B
Example 259	A

Compound	in vitro activity
Example 260	A
Example 261	A
Example 262	A
Example 263	A
Example 264	A
Example 265	A
Example 266	A
Example 267	A
Example 268	A
Example 269	A
Example 270	A
Example 271	A
Example 272	A
Example 273	A
Example 274	A
Example 275	A
Example 276	A
Example 277	B
Example 278	A
Example 279	A
Example 280	A
Example 281	A
Example 282	A
Example 283	A
Example 284	A
Example 285	A
Example 286	A
Example 287	A
Example 288	A
Example 289	A

Compound	in vitro activity
Example 290	A
Example 291	A
Example 292	A
Example 293	A
Example 294	A
Example 295	A
Example 296	A
Example 297	A
Example 298	A
Example 299	A
Example 300	A
Example 301	B
Example 302	A
Example 303	C
Example 304	A
Example 305	A
Example 306	C
Example 307	A
Example 308	C
Example 309	A
Example 310	A
Example 311	A
Example 312	A
Example 313	A
Example 314	A
Example 315	A
Example 316	A
Example 317	A
Example 318	A
Example 319	A

Compound	in vitro activity
Example 320	A
Example 321	A
Example 322	A
Example 323	A
Example 324	A
Example 325	A
Example 326	A
Example 327	A
Example 328	B
Example 329	A
Example 330	A
Example 331	A
Example 332	B
Example 333	A
Example 334	A
Example 335	A
Example 336	C
Example 337	C
Example 338	C
Example 339	B
Example 340	C
Example 341	C
Example 342	B
Example 343	B
Example 344	B
Example 345	B
Example 346	C
Example 347	C
Example 348	C
Example 349	A

Compound	in vitro activity
Example 350	C
Example 351	C
Example 352	C
Example 353	A
Example 354	B
Example 355	A
Example 356	C
Example 357	A
Example 358	B
Example 359	A
Example 360	A
Example 361	A
Example 362	A
Example 363	B
Example 364	B
Example 365	C
Example 366	A
Example 367	A
Example 368	A
Example 369	A
Example 370	A
Example 371	A
Example 372	A
Example 373	A
Example 374	A
Example 375	A
Example 376	A
Example 377	B
Example 378	B
Example 379	A

Compound	in vitro activity
Example 380	A
Example 381	A
Example 382	B
Example 383	A
Example 384	A
Example 385	B
Example 386	A
Example 387	A
Example 388	A
Example 389	B
Example 390	B
Example 391	A
Example 392	C
Example 393	A
Example 395	A
Example 396	C
Example 397	A
Example 398	A
Example 399	A
Example 400	A
Example 401	A
Example 402	A
Example 403	A
Example 404	A
Example 405	A
Example 406	A
Example 407	A
Example 408	A
Example 409	A
Example 410	A

Compound	in vitro activity
Example 411	A
Example 412	A
Example 413	A
Example 414	A
Example 415	A
Example 416	A
Example 417	A
Example 418	A

Inhibition of HIV replication. A recombinant NL-Rluc virus was constructed in which a section of the nef gene from NL4-3 was replaced with the *Renilla Luciferase* gene. The NL-RLuc virus was prepared by co-transfection of two plasmids, pNLRLuc and pVSVenv. The pNLRLuc contains the NL-Rluc DNA cloned into pUC18 at the *Pvu*II site, while the pVSVenv contains the gene for VSV G protein linked to an LTR promoter. Transfections were performed at a 1:3 ratio of pNLRLuc to pVSVenv on 293T cells using the LipofectAMINE PLUS kit from Invitrogen (Carlsbad, CA) according to manufactures instruction, and the pseudotype virus generated was titered in MT-2 cells.

Susceptibility of viruses to compounds was determined by incubation in the presence of serial dilutions of the compound. The 50% effective concentration (EC_{50}) was calculated by using the exponential form of the median effect equation where $(Fa) = 1/[1 + (ED_{50}/\text{drug conc.})^m]$ (Johnson VA, Byington RT. Infectivity Assay. In *Techniques in HIV Research*. ed. Aldovini A, Walker BD. 71-76. New York: Stockton Press.1990). The anti-viral activity of compounds was evaluated under three serum conditions, 10% FBS, 15mg/ml human serum albumin/10% FBS or 40% human serum/5% FBS, and the results from at least 2 experiments were used to calculate the EC_{50} values. Results are shown in the Table 2. Activity equal to A refers to a compound having $IC_{50} = < 10$ nM while B and C denote compounds having $IC_{50} = < 10$ nM and $IC_{50} < 100$ nM and $IC_{50} > 100$ nM respectively.

Table 2.

Example	Antiviral activity
Example 1	B
Example 2	B
Example 3	B
Example 4	C
Example 5	B
Example 6	C
Example 7	C
Example 8	B
Example 9	C
Example 11	C
Example 12	B
Example 13	B
Example 14	B
Example 15	B
Example 16	A
Example 17	A
Example 18	A
Example 19	C
Example 20	A
Example 21	A
Example 22	A
Example 23	A
Example 24	A
Example 25	A
Example 26	A
Example 27	B
Example 28	A
Example 29	B
Example 30	A

Example	Antiviral activity
Example 31	A
Example 32	A
Example 33	A
Example 34	A
Example 35	A
Example 36	A
Example 38	B
Example 39	B
Example 40	A
Example 41	A
Example 42	A
Example 43	A
Example 44	A
Example 45	A
Example 47	B
Example 48	A
Example 50	A
Example 51	A
Example 52	A
Example 53	A
Example 54	A
Example 55	A
Example 56	A
Example 57	A
Example 58	A
Example 59	A
Example 60	A
Example 61	A
Example 62	A
Example 63	A

Example	Antiviral activity
Example 64	A
Example 65	A
Example 66	A
Example 67	A
Example 68	B
Example 69	A
Example 70	A
Example 71	A
Example 72	A
Example 73	A
Example 74	A
Example 75	A
Example 76	A
Example 77	A
Example 78	A
Example 79	A
Example 80	A
Example 81	A
Example 82	A
Example 83	B
Example 84	C
Example 85	A
Example 86	A
Example 87	A
Example 88	A
Example 89	A
Example 90	A
Example 91	A
Example 92	A
Example 93	A

Example	Antiviral activity
Example 94	A
Example 95	A
Example 96	A
Example 97	A
Example 98	A
Example 99	A
Example 100	A
Example 101	A
Example 102	A
Example 103	A
Example 104	A
Example 105	A
Example 106	B
Example 107	B
Example 108	C
Example 109	C
Example 110	B
Example 111	A
Example 112	A
Example 113	A
Example 114	A
Example 115	A
Example 116	A
Example 117	B
Example 118	A
Example 119	A
Example 120	A
Example 121	A
Example 122	A
Example 123	B

Example	Antiviral activity
Example 124	A
Example 125	B
Example 126	A
Example 127	B
Example 128	A
Example 129	A
Example 130	A
Example 131	B
Example 132	A
Example 133	A
Example 134	A
Example 135	A
Example 136	B
Example 137	A
Example 138	A
Example 139	A
Example 140	A
Example 141	A
Example 142	A
Example 143	A
Example 144	B
Example 145	A
Example 146	A
Example 147	A
Example 148	A
Example 149	B
Example 150	C
Example 153	A
Example 154	A
Example 155	A

Example	Antiviral activity
Example 156	A
Example 157	A
Example 158	A
Example 159	A
Example 160	A
Example 161	A
Example 162	A
Example 163	A
Example 164	A
Example 165	A
Example 166	A
Example 167	A
Example 168	A
Example 169	A
Example 170	A
Example 171	A
Example 172	A
Example 173	A
Example 180	A
Example 186	A
Example 187	A
Example 188	A
Example 189	A
Example 190	C
Example 191	A
Example 192	A
Example 193	A
Example 194	B
Example 195	B
Example 196	A

Example	Antiviral activity
Example 197	B
Example 198	C
Example 199	A
Example 200	A
Example 201	A
Example 202	A
Example 203	A
Example 204	A
Example 205	A
Example 206	A
Example 207	A
Example 208	A
Example 209	A
Example 210	C
Example 211	A
Example 212	A
Example 213	A
Example 214	A
Example 215	A
Example 216	A
Example 218	A
Example 219	A
Example 220	C
Example 221	A
Example 222	A
Example 223	B
Example 224	A
Example 225	A
Example 227	C
Example 228	C

Example	Antiviral activity
Example 229	C
Example 230	C
Example 231	C
Example 232	C
Example 233	C
Example 234	A
Example 235	A
Example 236	B
Example 237	A
Example 238	A
Example 239	B
Example 240	B
Example 241	A
Example 242	A
Example 243	B
Example 245	A
Example 246	A
Example 247	A
Example 248	A
Example 249	A
Example 250	A
Example 251	A
Example 253	A
Example 254	A
Example 255	A
Example 256	A
Example 259	A
Example 262	A
Example 263	A
Example 264	A

Example	Antiviral activity
Example 265	A
Example 266	A
Example 267	A
Example 268	A
Example 270	A
Example 271	A
Example 272	A
Example 273	A
Example 274	A
Example 275	A
Example 276	A
Example 280	A
Example 281	A
Example 282	B
Example 283	B
Example 284	A
Example 285	C
Example 286	A
Example 287	A
Example 288	A
Example 289	A
Example 290	A
Example 291	A
Example 292	A
Example 293	A
Example 294	A
Example 295	A
Example 296	A
Example 297	B
Example 298	A

Example	Antiviral activity
Example 299	B
Example 300	A
Example 302	A
Example 303	C
Example 304	B
Example 305	A
Example 306	C
Example 309	A
Example 310	A
Example 312	A
Example 313	A
Example 314	A
Example 315	B
Example 316	A
Example 317	A
Example 318	A
Example 319	A
Example 320	A
Example 321	B
Example 323	A
Example 324	A
Example 325	A
Example 326	B
Example 327	A
Example 328	B
Example 329	A
Example 330	A
Example 334	A
Example 339	B
Example 342	A

Example	Antiviral activity
Example 343	A
Example 344	A
Example 345	B
Example 349	A
Example 353	A
Example 354	B
Example 355	A
Example 357	A
Example 358	A
Example 359	A
Example 360	A
Example 362	A
Example 371	A
Example 372	A
Example 373	A
Example 374	A
Example 375	A
Example 380	A
Example 382	B
Example 383	A
Example 384	A
Example 386	A
Example 387	A
Example 388	A
Example 391	A
Example 392	C
Example 393	B
Example 395	A
Example 396	C
Example 397	A

Example	Antiviral activity
Example 398	A
Example 399	A
Example 400	A
Example 401	A
Example 402	A
Example 403	A
Example 404	A
Example 406	A
Example 407	A
Example 408	A
Example 409	B
Example 410	A
Example 411	A
Example 412	A
Example 413	A
Example 414	A
Example 415	A
Example 416	A
Example 417	A
Example 418	A

Table 3 shows the activity of a set representative compounds.

Table 3.

Example	K _i μM	EC ₅₀ μM
Example 6	0.012	0.141
Example 19	0.126	0.302
Example 33	0.001	0.002
Example 38	0.009	0.037
Example 41	0.001	<0.001

Example	K _i μM	EC ₅₀ μM
Example 42	0.001	0.001
Example 48	0.001	0.004
Example 219	0.012	0.037
Example 303	0.180	0.192
Example 306	0.860	0.385
Example 314	<0.001	<0.001
Example 357	0.009	0.006
Example 382	0.011	0.016
Example 392	0.467	0.618
Example 396	0.294	0.455

Pharmaceutical Composition and Methods of Use

The compounds of this invention inhibit HIV integrase. HIV integrase inhibitors belonging to a class of diketo acid compounds prevented viral integration and inhibited HIV-1 replication in cells (Hazuda et al. *Science* **2000**, 287, 646). Recently raltegravir, an HIV integrase inhibitor, has been approved by the FDA for treating AIDS and HIV infection.

Accordingly, another aspect of the invention is a method for treating HIV infection in a human patient comprising administering a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, with a pharmaceutically acceptable carrier.

Another aspect of the invention is the use of a compound of formula I in the manufacture of a medicament for the treatment of AIDS or HIV infection.

Another aspect of the invention is a method for treating HIV infection in a human patient comprising the administration of a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, with a therapeutically effective amount of at least one other agent used for treatment of

AIDS or HIV infection selected from the group consisting of nucleoside HIV reverse transcriptase inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, HIV protease inhibitors, HIV fusion inhibitors, HIV attachment inhibitors, CCR5 inhibitors, CXCR4 inhibitors, HIV budding or maturation inhibitors, and HIV integrase inhibitors.

Another aspect of the invention is a method wherein the agent is a nucleoside HIV reverse transcriptase inhibitor.

10 Another aspect of the invention is a method wherein the nucleoside HIV reverse transcriptase inhibitor is selected from the group consisting of abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine, and zidovudine, or a pharmaceutically acceptable salt thereof.

15 Another aspect of the invention is a method wherein the agent is a non-nucleoside HIV reverse transcriptase inhibitor.

Another aspect of the invention is a method wherein the non-nucleoside HIV reverse transcriptase inhibitor is selected from the group consisting of delavirdine, 20 efavirenz, and nevirapine, or a pharmaceutically acceptable thereof.

Another aspect of the invention is a method wherein the agent is an HIV protease inhibitor.

25 Another aspect of the invention is a method wherein the HIV protease inhibitor is selected from the group consisting of amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir and fosamprenavir, or a pharmaceutically acceptable salt thereof.

30 Another aspect of the invention is a method wherein the agent is an HIV fusion inhibitor.

Another aspect of the invention is a method wherein the HIV fusion inhibitor is enfuvirtide or T-1249, or a pharmaceutically acceptable salt thereof.

5 Another aspect of the invention is a method wherein the agent is an HIV attachment inhibitor.

Another aspect of the invention is a method wherein the agent is a CCR5 inhibitor.

10 Another aspect of the invention is a method wherein the CCR5 inhibitor is selected from the group consisting of Sch-C, Sch-D, TAK-220, PRO-140, and UK-427,857, or a pharmaceutically acceptable salt thereof.

15 Another aspect of the invention is a method wherein the agent is a CXCR4 inhibitor.

Another aspect of the invention is a method wherein the CXCR4 inhibitor is AMD-3100, or a pharmaceutically acceptable salt thereof.

20 Another aspect of the invention is a method wherein the agent is an HIV budding or maturation inhibitor.

Another aspect of the invention is a method wherein the budding or maturation inhibitor is PA-457, or a pharmaceutically acceptable salt thereof.

25

Another aspect of the invention is a method wherein the agent is an HIV integrase inhibitor.

30 Another aspect of the invention is a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, with at least one other agent used for treatment of AIDS or HIV infection selected from the group consisting of nucleoside HIV reverse transcriptase inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, HIV

protease inhibitors, HIV fusion inhibitors, HIV attachment inhibitors, CCR5 inhibitors, CXCR4 inhibitors, HIV budding or maturation inhibitors, and HIV integrase inhibitors, and a pharmaceutically acceptable carrier.

5 Another aspect of the invention is the composition wherein the agent is a nucleoside HIV reverse transcriptase inhibitor.

Another aspect of the invention is the composition wherein the nucleoside HIV transcriptase inhibitor is selected from the group consisting of abacavir, 10 didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine, and zidovudine, or a pharmaceutically acceptable salt thereof.

Another aspect of the invention is the composition wherein the agent is a non-nucleoside HIV reverse transcriptase inhibitor.

15 Another aspect of the invention is the composition wherein the non-nucleoside HIV reverse transcriptase inhibitor is selected from the group consisting of delavirdine, efavirenz, and nevirapine, or a pharmaceutically acceptable salt thereof.

20 Another aspect of the invention is the composition wherein the agent is an HIV protease inhibitor.

Another aspect of the invention is the composition wherein the HIV protease 25 inhibitor is selected from the group consisting of amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir and fosamprenavir, or a pharmaceutically acceptable salt thereof.

30 Another aspect of the invention is the composition wherein the agent is an HIV fusion inhibitor.

Another aspect of the invention is the composition method wherein the HIV fusion inhibitor is enfuvirtide or T-1249, or a pharmaceutically acceptable salt thereof.

5 Another aspect of the invention is the composition wherein the agent is an HIV attachment inhibitor.

Another aspect of the invention is the composition wherein the agent is a CCR5 inhibitor.

10 Another aspect of the invention is the composition wherein the CCR5 inhibitor is selected from the group consisting of Sch-C, Sch-D, TAK-220, PRO-140, and UK-427,857, or a pharmaceutically acceptable salt thereof.

15 Another aspect of the invention is a method wherein the agent is a CXCR4 inhibitor.

Another aspect of the invention is a method wherein the CXCR4 inhibitor is AMD-3100 or a pharmaceutically acceptable salt thereof.

20 Another aspect of the invention is the composition wherein the agent is an HIV budding or maturation inhibitor.

25 Another aspect of the invention is the composition wherein the budding or maturation inhibitor is PA-457, or a pharmaceutically acceptable salt thereof.

Another aspect of the invention is the composition wherein the agent is an HIV integrase inhibitor.

30 “Combination,” “coadministration,” “concurrent,” and similar terms referring to the administration of a compound of Formula I with at least one anti-HIV agent mean that the components are part of a combination antiretroviral therapy or highly

active antiretroviral therapy (HAART) as understood by practitioners in the field of AIDS and HIV infection.

“Therapeutically effective” means the amount of agent required to provide a meaningful patient benefit as understood by practitioners in the field of AIDS and HIV infection. In general, the goals of treatment are suppression of viral load, restoration and preservation of immunologic function, improved quality of life, and reduction of HIV-related morbidity and mortality.

10 “Patient” means a person infected with the HIV virus and suitable for therapy as understood by practitioners in the field of AIDS and HIV infection.

15 “Treatment,” “therapy,” “regimen,” “HIV infection,” “ARC,” “AIDS” and related terms are used as understood by practitioners in the field of AIDS and HIV infection.

20 The compounds of this invention are generally given as pharmaceutical compositions comprised of a therapeutically effective amount of a compound of Formula I or its pharmaceutically acceptable salt and a pharmaceutically acceptable carrier and may contain conventional excipients. A therapeutically effective amount is that which is needed to provide a meaningful patient benefit. Pharmaceutically acceptable carriers are those conventionally known carriers having acceptable safety profiles. Compositions encompass all common solid and liquid forms including capsules, tablets, losenges, and powders as well as liquid suspensions, syrups, elixers, 25 and solutions. Compositions are made using common formulation techniques, and conventional excipients (such as binding and wetting agents) and vehicles (such as water and alcohols) are generally used for compositions.

30 Solid compositions are normally formulated in dosage units and compositions providing from about 1 to 1000 mg of the active ingredient per dose are preferred. Some examples of dosages are 1 mg, 10 mg, 100 mg, 250 mg, 500 mg, and 1000 mg. Generally, other antiretroviral agents will be present in a unit range similar to agents of that class used clinically. Typically, this is 0.25-1000 mg/unit.

Liquid compositions are usually in dosage unit ranges. Generally, the liquid composition will be in a unit dosage range of 1-100 mg/mL. Some examples of dosages are 1 mg/mL, 10 mg/mL, 25 mg/mL, 50 mg/mL, and 100 mg/mL. Generally, other antiretroviral agents will be present in a unit range similar to agents of that class used clinically. Typically, this is 1-100 mg/mL.

The invention encompasses all conventional modes of administration; oral and parenteral methods are preferred. Generally, the dosing regimen will be similar to other antiretroviral agents used clinically. Typically, the daily dose will be 1-100 mg/kg body weight daily. Generally, more compound is required orally and less parenterally. The specific dosing regime, however, will be determined by a physician using sound medical judgement.

The invention also encompasses methods where the compound is given in combination therapy. That is, the compound can be used in conjunction with, but separately from, other agents useful in treating AIDS and HIV infection. Some of these agents include HIV attachment inhibitors, CCR5 inhibitors, CXCR4 inhibitors, HIV cell fusion inhibitors, HIV integrase inhibitors, HIV nucleoside reverse transcriptase inhibitors, HIV non-nucleoside reverse transcriptase inhibitors, HIV protease inhibitors, budding and maturation inhibitors, immunomodulators, and anti-infectives. In these combination methods, the compound of Formula I will generally be given in a daily dose of 1-100 mg/kg body weight daily in conjunction with other agents. The other agents generally will be given in the amounts used therapeutically. The specific dosing regime, however, will be determined by a physician using sound medical judgement.

Table 4 lists some agents useful in treating AIDS and HIV infection which are suitable for this invention.

Table 4. ANTIVIRALS

DRUG NAME	MANUFACTURER	INDICATION
097 (non-nucleoside reverse transcriptase inhibitor)	Hoechst/Bayer	HIV infection, AIDS, ARC
Amprenavir 141 W94 GW 141 (protease inhibitor)	Glaxo Wellcome	HIV infection, AIDS, ARC
Abacavir (1592U89) GW 1592 (RT inhibitor)	Glaxo Wellcome	HIV infection, AIDS, ARC
Acemannan	Carrington Labs (Irving, TX)	ARC
Acyclovir	Burroughs Wellcome	HIV infection, AIDS, ARC, in combination with AZT
AD-439	Tanox Biosystems	HIV infection, AIDS, ARC
AD-519	Tanox Biosystems	HIV infection, AIDS, ARC
Adefovir dipivoxil AL-721	Gilead Sciences Ethigen (Los Angeles, CA)	HIV infection, ARC, PGL HIV positive, AIDS
Alpha Interferon HIV in combination w/Retrovir	Glaxo Wellcome	Kaposi's sarcoma
Ansamycin LM 427	Adria Laboratories (Dublin, OH) Erbamont (Stamford, CT)	ARC

DRUG NAME	MANUFACTURER	INDICATION
Antibody which Neutralizes pH Labile alpha aberrant Interferon	Advanced Biotherapy Concepts (Rockville, MD)	AIDS, ARC
AR177	Aronex Pharm	HIV infection, AIDS, ARC
Beta-fluoro-ddA	Nat'l Cancer Institute	AIDS-associated diseases
BMS-232623 (CGP-73547) (protease inhibitor)	Bristol-Myers Squibb/ Novartis	HIV infection, AIDS, ARC
BMS-234475 (CGP-61755) (protease inhibitor)	Bristol-Myers Squibb/ Novartis	HIV infection, AIDS, ARC
CI-1012	Warner-Lambert	HIV-1 infection
Cidofovir	Gilead Science	CMV retinitis, herpes, papillomavirus
Curdian sulfate	AJI Pharma USA	HIV infection
Cytomegalovirus Immune globin	MedImmune	CMV retinitis
Cytovene	Syntex	Sight threatening
Ganciclovir		CMV peripheral, CMV retinitis
Delavirdine (RT inhibitor)	Pharmacia-Upjohn	HIV infection, AIDS, ARC
Dextran Sulfate	Ueno Fine Chem. Ind. Ltd. (Osaka, Japan)	AIDS, ARC, HIV positive asymptomatic
ddC Dideoxycytidine	Hoffman-La Roche	HIV infection, AIDS, ARC

DRUG NAME	MANUFACTURER	INDICATION
ddI Dideoxyinosine	Bristol-Myers Squibb	HIV infection, AIDS, ARC; combinationwith AZT/d4T
DMP-450 (protease inhibitor)	AVID (Camden, NJ)	HIV infection, AIDS, ARC
Efavirenz (DMP 266) (-)-6-Chloro-4-(S)- cyclopropylethynyl- 4(S)-trifluoro- methyl-1,4-dihydro- 2H-3,1-benzoxazin- 2-one, STOCRINE (non-nucleoside RT inhibitor)	DuPont Merck	HIV infection, AIDS, ARC
EL10	Elan Corp, PLC (Gainesville, GA)	HIV infection
Famciclovir	Smith Kline	herpes zoster, herpes simplex
FTC (reverse transcriptase inhibitor)	Emory University	HIV infection, AIDS, ARC
GS 840 (reverse transcriptase inhibitor)	Gilead	HIV infection, AIDS, ARC
HBY097 (non-nucleoside reverse transcriptaseinhibitor)	Hoechst Marion Roussel	HIV infection, AIDS, ARC
Hypericin	VIMRx Pharm.	HIV infection, AIDS, ARC

DRUG NAME	MANUFACTURER	INDICATION
Recombinant Human Interferon Beta	Triton Biosciences (Almeda, CA)	AIDS, Kaposi's sarcoma, ARC
Interferon alfa-n3	Interferon Sciences	ARC, AIDS
Indinavir	Merck	HIV infection, AIDS, ARC, asymptomatic HIV positive, also in combination with AZT/ddI/ddC
ISIS 2922	ISIS Pharmaceuticals	CMV retinitis
KNI-272	Nat'l Cancer Institute	HIV-associated diseases
Lamivudine, 3TC (reverse transcriptase inhibitor)	Glaxo Wellcome	HIV infection, AIDS, ARC, also with AZT
Lobucavir	Bristol-Myers Squibb	CMV infection
Nelfinavir (protease inhibitor)	Agouron Pharmaceuticals	HIV infection, AIDS, ARC
Nevirapine (RT inhibitor)	Boehringer Ingleheim	HIV infection, AIDS, ARC
Novapren	Novaferon Labs, Inc. (Akron, OH)	HIV inhibitor
Peptide T Octapeptide Sequence	Peninsula Labs (Belmont, CA)	AIDS
Trisodium Phosphonoformate	Astra Pharm. Products, Inc.	CMV retinitis, HIV infection, other CMV infections
PNU-140690 (protease inhibitor)	Pharmacia Upjohn	HIV infection, AIDS, ARC
Probucol	Vyrex	HIV infection, AIDS
RBC-CD4	Sheffield Med. Tech (Houston, TX)	HIV infection, AIDS, ARC

DRUG NAME	MANUFACTURER	INDICATION
Ritonavir (protease inhibitor)	Abbott	HIV infection, AIDS, ARC
Saquinavir (protease inhibitor)	Hoffmann-LaRoche	HIV infection, AIDS, ARC
Stavudine; d4T Didehydrodeoxy- thymidine	Bristol-Myers Squibb	HIV infection, AIDS, ARC
Valaciclovir	Glaxo Wellcome	Genital HSV & CMV infections
Virazole Ribavirin	Viratek/ICN (Costa Mesa, CA)	asymptomatic HIV- positive, LAS, ARC
VX-478	Vertex	HIV infection, AIDS, ARC
Zalcitabine	Hoffmann-LaRoche	HIV infection, AIDS, ARC, with AZT
Zidovudine; AZT	Glaxo Wellcome	HIV infection, AIDS, ARC, Kaposi's sarcoma, in combination with other therapies
Tenofovir disoproxil, fumarate salt (VIREAD®) (reverse transcriptase inhibitor)	Gilead	HIV infection, AIDS
COMBIVIR® (reverse transcriptase inhibitor)	GSK	HIV infection, AIDS
abacavir succinate (or ZIAGEN®) (reverse transcriptase inhibitor)	GSK	HIV infection, AIDS

DRUG NAME	MANUFACTURER	INDICATION
REYATAZ® (atazanavir)	Bristol-Myers Squibb	HIV infection, AIDS
Fuzeon (Enfuvirtide, T-20)	Roche/Trimeris	HIV infection, AIDS, viral fusion inhibitor
TRIZIVIR®		HIV infection, AIDS
KALETRA®	Abbott	HIV infection, AIDS, ARC

IMMUNOMODULATORS

DRUG NAME	MANUFACTURER	INDICATION
AS-101	Wyeth-Ayerst	AIDS
Propirimine	Pharmacia Upjohn	Advanced AIDS
Acemannan	Carrington Labs, Inc. (Irving, TX)	AIDS, ARC
CL246,738	American Cyanamid Lederle Labs	AIDS, Kaposi's sarcoma
EL10	Elan Corp, PLC (Gainesville, GA)	HIV infection
FP-21399	Fuki ImmunoPharm	Blocks HIV fusion with CD4+ cells
Gamma Interferon	Genentech	ARC, in combination w/TNF (tumor necrosis factor)
Granulocyte Macrophage Colony Stimulating Factor	Genetics Institute Sandoz	AIDS
Granulocyte Macrophage Colony Stimulating Factor	Hoechst-Roussel Immunex	AIDS

DRUG NAME	MANUFACTURER	INDICATION
Granulocyte Macrophage Colony Stimulating Factor	Schering-Plough	AIDS, combination w/AZT
HIV Core Particle Immunostimulant	Rorer	Seropositive HIV
IL-2 Interleukin-2	Cetus	AIDS, in combination w/AZT
IL-2 Interleukin-2	Hoffman-LaRoche Immunex	AIDS, ARC, HIV, in combination w/AZT
IL-2 Interleukin-2 (aldeslukin)	Chiron	AIDS, increase in CD4 cell counts
Immune Globulin Intravenous (human)	Cutter Biological (Berkeley, CA)	Pediatric AIDS, in combination w/AZT
IMREG-1	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
IMREG-2	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
Imuthiol Diethyl Dithio Carbamate	Merieux Institute	AIDS, ARC
Alpha-2 Interferon	Schering Plough	Kaposi's sarcoma w/AZT, AIDS
Methionine- Enkephalin	TNI Pharmaceutical (Chicago, IL)	AIDS, ARC
MTP-PE Muramyl-Tripeptide Granulocyte Colony Stimulating Factor	Ciba-Geigy Corp. Amgen	Kaposi's sarcoma AIDS, in combination w/AZT

DRUG NAME	MANUFACTURER	INDICATION
Remune	Immune Response Corp.	Immunotherapeutic
rCD4 Recombinant Soluble Human CD4	Genentech	AIDS, ARC
rCD4-IgG hybrids		AIDS, ARC
Recombinant Soluble Human CD4	Biogen	AIDS, ARC
Interferon Alfa 2a	Hoffman-La Roche in combination w/AZT	Kaposi's sarcoma, AIDS, ARC
SK&F106528 Soluble T4	Smith Kline	HIV infection
Thymopentin	Immunobiology Research Institute (Annandale, NJ)	HIV infection
Tumor Necrosis Factor; TNF	Genentech	ARC, in combination w/gamma Interferon

ANTI-INFECTIVES

DRUG NAME	MANUFACTURER	INDICATION
Clindamycin with Primaquine	Pharmacia Upjohn	PCP
Fluconazole	Pfizer	Cryptococcal meningitis, candidiasis
Pastille Nystatin Pastille	Squibb Corp.	Prevention of oral candidiasis
Ornidyl Eflornithine	Merrell Dow	PCP
Pentamidine Isethionate (IM & IV)	LyphoMed (Rosemont, IL)	PCP treatment

DRUG NAME	MANUFACTURER	INDICATION
Trimethoprim		Antibacterial
Trimethoprim/sulfa		Antibacterial
Piritrexim	Burroughs Wellcome	PCP treatment
Pentamidine Isethionate for Inhalation	Fisons Corporation	PCP prophylaxis
Spiramycin	Rhone-Poulenc diarrhea	Cryptosporidial
Intraconazole- R51211	Janssen-Pharm.	Histoplasmosis; cryptococcal meningitis
Trimetrexate	Warner-Lambert	PCP
Daunorubicin	NeXstar, Sequus	Kaposi's sarcoma
Recombinant Human Erythropoietin	Ortho Pharm. Corp.	Severe anemia assoc. with AZT therapy
Recombinant Human Growth Hormone	Serono	AIDS-related wasting, cachexia
Megestrol Acetate	Bristol-Myers Squibb	Treatment of anorexia assoc. W/AIDS
Testosterone	Alza, Smith Kline	AIDS-related wasting
Total Enteral Nutrition	Norwich Eaton Pharmaceuticals	Diarrhea and malabsorption related to AIDS

DESCRIPTION OF SPECIFIC EMBODIMENTS


Abbreviations used in the schemes generally follow conventions used in the art. Chemical abbreviations used in the specification and Examples are defined as follows: "NaHMDS" for sodium bis(trimethylsilyl)amide; "DMF" for N,N-dimethylformamide; "MeOH" for methanol; "NBS" for N-bromosuccinimide; "Ar" for aryl; "TFA" for trifluoroacetic acid; "LAH" for lithium aluminum hydride; "BOC", "DMSO" for dimethylsulfoxide; "h" for hours; "rt" for room temperature or

retention time (context will dictate); "min" for minutes; "EtOAc" for ethyl acetate; "THF" for tetrahydrofuran; "EDTA" for ethylenediaminetetraacetic acid; "Et₂O" for diethyl ether; "DMAP" for 4-dimethylaminopyridine; "DCE" for 1,2-dichloroethane; "ACN" for acetonitrile; "DME" for 1,2-dimethoxyethane; "HOBr" for 1-hydroxybenzotriazole hydrate; "DIEA" for diisopropylethylamine, "Nf" for CF₃(CF₂)₃SO₂⁻; and "TMOF" for trimethylorthoformate.

Abbreviations as used herein, are defined as follows: "1 x" for once, "2 x" for twice, "3 x" for thrice, "°C" for degrees Celsius, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams, "L" for liter or liters, "mL" for milliliter or milliliters, "μL" for microliter or microliters, "N" for normal, "M" for molar, "mmol" for millimole or millimoles, "min" for minute or minutes, "h" for hour or hours, "rt" for room temperature, "RT" for retention time, "atm" for atmosphere, "psi" for pounds per square inch, "conc." for concentrate, "sat" or "sat'd" for saturated, "MW" for molecular weight, "mp" for melting point, "ee" for enantiomeric excess, "MS" or "Mass Spec" for mass spectrometry, "ESI" for electrospray ionization mass spectroscopy, "HR" for high resolution, "HRMS" for high resolution mass spectrometry, "LCMS" for liquid chromatography mass spectrometry, "HPLC" for high pressure liquid chromatography, "RP HPLC" for reverse phase HPLC, "TLC" or "tlc" for thin layer chromatography, "NMR" for nuclear magnetic resonance spectroscopy, "¹H" for proton, "δ" for delta, "s" for singlet, "d" for doublet, "t" for triplet, "q" for quartet, "m" for multiplet, "br" for broad, "Hz" for hertz, and "α", "β", "R", "S", "E", and "Z" are stereochemical designations familiar to one skilled in the art.

25

Intermediate 1




30 *3-(Dimethylamino)-3-oxopropanoic acid.* To a stirred solution of 3-tert-butoxy-3-oxopropanoic acid (680 mg, 4.25 mmol) and 2M dimethylamine in THF

(3.18 mL, 6.37 mmol) in DMF (10 mL) was added diisopropyl-ethylamine (2.224 mL, 12.74 mmol), 4-(dimethylamino)pyridine (DMAP) (104 mg, 0.849 mmol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (2421 mg, 6.37 mmol) and the resulting mixture was stirred at room 5 temperature for 16 h. Water was then added and the resulting mixture was extracted with ethyl acetate (2 x 50 mL), washed with 1N HCl (20 mL), dried (Na_2SO_4), filtered and concentrated to give crude product which was dissolved in CH_2Cl_2 (5.00 mL), treated with trifluoroacetic acid (4.91 mL, 63.7 mmol) and stirred at room 10 temperature for 16h. The solvent was then removed under reduced pressure to afford the title compound (230 mg, 41% yield) as a light-yellow solid, which was used in the next step without further purification. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ : 12.1 (1 H, br s), 3.4 (2 H, s), 2.9 (3 H, s), 2.8 (3 H, s). LCMS ($\text{M}+\text{H}$) = 132.33.

Intermediate 2


15



2-(Methyl(2,2,2-trifluoroethyl)amino)-2-oxoacetic acid. To a solution of methyl 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoacetate (1.2 g, 6.03 mmol) in 20 methanol (15 mL) and water (15.00 mL) was added LiOH (0.506 g, 12.05 mmol) and the mixture stirred for 2h. The solution was then acidified with conc. HCl and concentrated. The solid was washed with CH_2Cl_2 (2 x 100 mL), filtered and the filtrate concentrated to afford the title compound (600 mg, 53.8% yield) (mixture of rotational isomers at room temperature) as an off-white solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ : 4.32 (1 H, q, $J=9.36$ Hz), 4.23 (1 H, q, $J=9.46$ Hz), 3.07 (2 H, s), 2.97 (1 H, s).

Intermediate 3

30



tert-Butyl 2-fluoroethylcarbamate. To a solution of 2-fluoroethanamine hydrochloride salt (2.5 g, 25.1 mmol) in THF (40 mL)/water (40 mL) at 0 °C was added sodium bicarbonate (8.44 g, 100 mmol), portion-wise. After 5 min, di-*tert*-butyl dicarbonate (7.00 mL, 30.1 mmol) was slowly added. The reaction was stirred under N₂ and allowed to warm to room temperature. The solution was concentrated, diluted with water (60 mL) and extracted with CH₂Cl₂ (3 x 75 mL). The organic layer was dried (MgSO₄) and concentrated to a give a crude oil which was purified by flash chromatography (Biotage flash chromatography system; 300g SiO₂; 0-70% ethyl acetate:hexane over 2000 mL) to give the title compound as a clear oil (4.1 g, 100% yield). ¹H NMR (400 MHz, CDCl₃) δ: 4.86 (1 H, brs), 4.47 (2 H, dt, *J* = 47.43, 4.77 Hz), 3.43 (2 H, dq, *J* = 27.92, 5.08 Hz), 1.45 (9 H, s).

Intermediate 4



tert-Butyl 2-fluoroethyl(methyl)carbamate. To a solution of *tert*-butyl 2-fluoroethylcarbamate (4.1 g, 25.1 mmol) in THF (100 mL) at 0 °C was added sodium hydride, 60 wt % in mineral oil (1.507 g, 37.7 mmol) portion-wise. The reaction was stirred under N₂ for 15 minutes after which gas evolution ceased. Iodomethane (2.356 mL, 37.7 mmol) was added drop-wise and the reaction stirred under N₂. After 6 h, the reaction was quenched with water (100 mL) and concentrated. The aqueous solution was washed with CH₂Cl₂ (3 x 50 mL). The CH₂Cl₂ extracts were combined, washed with water (2 x 100mL), dried (MgSO₄) and concentrated to afford the title compound (4.0 g, 90% yield) as a clear oil. ¹H NMR (400 MHz, CDCl₃) δ: 4.58 (1 H, brs), 4.47 (1 H, brs), 3.53 (1 H, brs), 3.47 (1 H, brs), 2.94 (3 H, s), 1.45 (9 H, s).


Intermediate 5



2-Fluoro-N-methylethanamine hydrochloride. To a solution of tert-butyl 2-fluoroethyl(methyl)carbamate (4.0 g, 22.57 mmol) in CH₂Cl₂ (10 mL) was added trifluoroacetic acid (10 mL, 130 mmol) and the reaction stirred under N₂ for 2 h. The solution was concentrated, dissolved in 1N HCl (30 mL), washed with Et₂O (2 x 30 mL) and the water removed to give the title compound HCl salt as a white solid. ¹H NMR (500 MHz, CDCl₃) δ: 4.76 (2 H, dt, *J* = 46.69, 4.27 Hz), 3.36 (2 H, dt, *J* = 25.79, 3.97, 3.81 Hz), 2.84 (3 H, s).

Intermediate 6


10



Methyl 2-((2-fluoroethyl)(methyl)amino)-2-oxoacetate. To a solution of 2-fluoro-N-methylethanamine hydrochloride salt (1 g, 8.81 mmol) in CH₂Cl₂ (40.0 mL) was added drop-wise diisopropyl-ethylamine (5.59 mL, 32.0 mmol) followed by the drop-wise addition of methyl 2-chloro-2-oxoacetate (0.981 g, 8.01 mmol) and the resulting mixture stirred for 3 h under N₂. The solution was concentrated and the residue was taken up in ethyl acetate (20 mL), washed with 1N HCl (2 x 15 mL) followed by saturated aqueous sodium bicarbonate (15 mL). The organic phase was dried (MgSO₄), filtered, and concentrated to give the title compound as a clear oil (1.2 g, 92% yield). ¹H NMR (500 MHz, CDCl₃) δ: 4.46 – 4.73 (2 H, m) 3.82 – 3.93 (3 H, m) 3.60 – 3.77 (2 H, m) 3.01 – 3.20 (3 H, m). LCMS (M+H) calcd for C₆H₁₁FNO₃: 164.07; found: 164.11.

25


Intermediate 7



2-((2-Fluoroethyl)(methyl)amino)-2-oxoacetic acid. To a solution of ethyl 2-((2-fluoroethyl)(methyl)amino)-2-oxoacetate (1.2 g, 7.36 mmol) in methanol/ water (30 mL, 1:1) was added LiOH (0.352 g, 14.71 mmol). The solution was stirred for 2 h. The methanol was removed by concentration and concentrated HCl (2 mL) was added. The solvent was removed and the residue dissolved in CH₂Cl₂ and filtered. The filtrate was concentrated to give the title compound as an off-white crystalline powder (0.615 g, 56% yield). ¹H NMR (400 MHz, CDCl₃) δ: 4.47 - 4.89 (2 H, m) 3.67 - 4.39 (2 H, m) 3.07 - 3.60 (3 H, m).


10

Intermediate 8



2-((4-Methoxybenzylamino)-2-methylpropan-1-ol. To a solution of 2-amino-2-methyl-1-propanol (8.4, 94 mmol) in 1,2-dichloroethane (150 mL) and acetic acid (4.3 mL, 77 mmol) was added 4-methoxy benzaldehyde (11.5 mL, 94 mmol). After stirring for 10 min, sodium triacetoxyborohydride (49.9g, 235.5 mmol) was added portion-wise over 20 min. The resulting mixture was stirred at 60 °C for 2 h then cooled to room temperature and quenched with saturated NaHCO₃. The aqueous layer was washed with dichloromethane and made basic by the addition of 10N NaOH. The product was extracted with ethyl acetate, concentrated and crystallized from Et₂O/hexane to give the title compound as a white solid (11.95 g, 61% yield). ¹H NMR (300 MHz, CDCl₃) δ: 7.32-7.20 (2H, m), 6.86 - 6.80 (2H, m), 3.76 (3H, s), 3.58 (2H, s), 3.30 (2H, s), 1.11 (6H, s). LCMS (M+H) calcd for C₁₂H₂₀NO₂: 210.14; found: 210.20.


Intermediate 9



5 *Methyl 5-(4-methoxybenzyl)-4,4-dimethyl-1,2,5-thiadiazolidine-2-carboxylate 1,1-dioxide.* To a solution of 2-(4-methoxybenzylamino)-2-methylpropan-1-ol (12 g, 57.3 mmol) in THF (130 mL) was added (methoxycarbonylsulfamoyl)triethylammonium hydroxide(Burgess reagent) (44 g, 172 mmol) and the resulting mixture stirred at room temperature for 2 days. The
10 mixture was diluted with ethyl acetate and washed with 1N HCl followed by brine. The organic phase was dried (Na_2SO_4), filtered and concentrated. The residue was purified by flash chromatography (Biotage flash chromatography system; 0% -
70%ethyl acetate/hexane) to give a pale yellow oil that was further purified by flash chromatography eluting with 25% then 50% ethyl acetate/hexane to afford the title
15 compound as a white solid (17.56 g, 93% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.32 - 7.31 (2H, m), 6.86 - 6.84 (2H, m), 4.19 (2H, s), 3.89 (3H, s), 3.78 (3H, s), 3.64 (2H, s), 1.28 (6H, s).

Intermediate 10


20



25 *2-(4-Methoxybenzyl)-3,3-dimethyl-1,2,5-thiadiazolidine 1,1-dioxide.* To a suspension of methyl 5-(4-methoxybenzyl)-4,4-dimethyl-1,2,5-thiadiazolidine-2- carboxylate 1,1-dioxide (17.56 g, 53.5 mmol) in methanol/water (300 mL, 2:1) was added 10 N NaOH (10.7 mL, 107 mmol) and the mixture stirred at room temperature

for 4 h. The mixture was made acidic with 10 N HCl and washed with ethyl acetate. The combined organic layers were dried (Na_2SO_4), filtered and concentrated to give the title compound as a white solid (12.91 g, 89% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.34 - 7.32 (2H, m), 6.86 - 6.84 (2H, m), 4.74 (1H, brs), 4.18 (2H, s), 3.78 (3H, s), 3.24 (2H, s), 1.21 (9H, s).


Intermediate 11



10

4-Fluoro-2-(5-(4-methoxybenzyl)-4,4-dimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)benzonitrile. 2-Bromo-4-fluorobenzonitrile, tris(dibenzylideneacetone)dipalladium(0) (0.238 g, 0.26 mmol), 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene (0.231 g, 0.4 mmol) and potassium carbonate (5.72 g, 41.4 mmol) were stirred together. Added to this was a solution of 2-(4-methoxybenzyl)-3,3-dimethyl-1,2,5-thiadiazolidine 1,1-dioxid (8.00 g, 29.6 mmol) in dioxane (50 mL) and the resulting mixture stirred at 100 °C for 18 h. The cooled mixture was diluted with CH_2Cl_2 and the solids were removed by filtration. The solvent was evaporated leaving a yellow oil that was purified by flash chromatography (Biotage flash chromatography system, 5% - 100% ethyl acetate/hexane) to give a cream colored solid that was triturated with ethyl acetate/hexane (1:4) to give the title compound as a white powder (4.6048 g, 42% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.71 - 7.65 (2H, m), 7.37 (2H, d, J = 8.5 Hz), 7.12 - 7.08 (1H, m), 6.88 (2H, d, J = 8.5 Hz), 4.28 (2H, s), 3.80 (3H, s), 3.78 (2H, s), 1.42 (6H, s).


Intermediate 12



5 *2-(4,4-Dimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)-4-fluorobenzonitrile.*

To a solution of 4-fluoro-2-(5-(4-methoxybenzyl)-4,4-dimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)benzonitrile (4.6 g, 11.81 mmol) in dichloromethane (30 mL) was added trifluoroacetic acid (2.73 mL, 35.4 mmol) and the mixture stirred at room temperature for 18 h. The mixture was diluted with methanol and the white solid collected by filtration and discarded. The cloudy solution was concentrated and the resulting solid was triturated with methanol and then filtered to give the title compound as a white solid (3.18 g, 100% yield). LCMS (M+H) calcd for C₁₁H₁₃FN₃O₂S: 270.07; found: 270.00. ¹H NMR (300 MHz, CDCl₃) δ: 7.66 (.2H, dd, *J* = 8.8, 5.8 Hz), 7.58 (1H, dd, *J* = 9.5, 2.6 Hz), 7.10 - 7.04 (1H, m), 4.48 (1H, brs), 3.85 (2H, s), 1.55 (6H, s).

Intermediate 13




20

4-Fluoro-2-(4,4,5-trimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)benzonitrile. To a solution of 2-(4,4-dimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)-4-fluorobenzonitrile (3.18 g, 11.81 mmol) in DMF (40 mL) was added NaH (0.312 g, 12.99 mmol) and the mixture stirred for 5 min. Iodomethane (1.477 mL, 23.62 mmol) was added and the mixture stirred at 90 °C for 2 h then at room

temperature for 16 h. The mixture was diluted with H₂O and extracted with ethyl acetate. The organic phase was washed two times with H₂O followed by brine, dried (Na₂SO₄), filtered and concentrated to give the title compound as a white solid (3.35 g, 100% yield). LCMS (M+H) calcd for C₁₂H₁₅FN₃O₂S: 284.08; found: 284.14. ¹H NMR (300 MHz, CDCl₃) δ: 7.70 - 7.61 (2H, m), 7.11 - 7.05 (1H, m), 3.80 (2H, s), 2.73 (3H, s), 1.42 (6H, s).


Intermediate 14



10

I-(4-Fluoro-2-(4,4,5-trimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)phenyl)methanamine. To a solution of 4-fluoro-2-(4,4,5-trimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)benzonitrile (3.34 g, 11.8 mmol) in CH₂Cl₂ (30 mL), ethyl acetate (30.0 mL) and ethanol (30.0 mL) was added 1N HCl (12 mL) followed by 10% palladium on carbon (0.377 g, 3.54 mmol). The mixture was shaken under H₂ at 15 50 psi for 22 h then filtered over CELITE® and concentrated. The solids were triturated with H₂O and removed by filtration. The aqueous solution was washed with ethyl acetate and lyophilized to give the title compound, as the HCl salt, as a white powder (3.44 g, 90% yield). LCMS (M+H) calcd for C₁₂H₁₉FN₃O₂S: 288.11; found: 288.19. ¹H NMR (300 MHz, DMSO) δ: 8.35 (2H, brs), 7.71 (1H, dd, *J* = 8.6, 6.4 Hz), 7.54 (1H, dd, *J* = 9.9, 2.6 Hz), 7.43 (1H, td, *J* = 8.4, 2.6 Hz), 4.19 (2H, s), 3.75 (2H, s), 2.63 (3H, s), 1.35 (6H, s).


Intermediate 15



5 *2,6-Dibromo-4-fluoro-3-methylaniline.* To a stirred solution of 4-fluoro-3-methylaniline (12.76 g, 102 mmol) in 1:1 CH₂Cl₂/methanol (125 ml) was added, dropwise, a solution of bromine (13.14 ml, 255 mmol) in 1:1 CH₂Cl₂/methanol (125 ml) at room temperature over 1.5 h using an addition funnel. After 4 h, the reaction mixture was concentrated and the resulting residue was stirred with 1N Na₂S₂O₃ (100 mL) and ethyl acetate (150 mL) for 10 min then carefully basified with 1N Na₂CO₃ (150 mL). The organic layer was separated and the aqueous layer extracted with ethyl acetate (2 x 100 mL). The combined organic layers were washed with 1N Na₂S₂O₃ (100 mL) followed by brine (50 mL), then dried (Na₂SO₄/carbon), filtered and concentrated to give 2,6-dibromo-4-fluoro-3-methylaniline (28.6 g, 101 mmol, 10 99% yield) as a tan color solid. ¹H NMR (400 MHz, CDCl₃) δ: 7.19 (1 H, d, *J* = 8.5 Hz), 4.42 (2 H, brs), 2.31 (3 H, d, *J* = 2.5 Hz). LCMS (M+H) calcd for C₇H₇FN: 283.89; found: 283.83.

Intermediate 16

20



25 *2,6-Dibromo-4-fluoro-3-methylbenzonitrile.* To a mixture of 2,6-dibromo-4-fluoro-3-methylaniline (2.83 g, 10 mmol), acetic acid (7 mL) and water (4 mL) was added H₂SO₄ (1.492 mL, 28 mmol) and the resulting mixture stirred at 70 °C until it turned into a homogeneous solution. The reaction mixture was then cooled in ice-


water bath and treated with 1N sodium nitrite (0.759 g, 11 mmol) over 15 min., to produce the corresponding diazonium salt.

In a separate flask, KCN (3.26 g, 50 mmol) was added in small portions over 5 15 min., to a solution of copper (II) sulfate (1.915 g, 12 mmol) in water (25 mL) at ice bath temperature. The ice bath was then removed after which sodium bicarbonate (6.72 g, 80 mmol) and benzene (20 mL) was added and the mixture warmed to 55 °C. To this vigorously stirred mixture was added, dropwise over 15 min., the diazonium salt synthesized above. After 1 h, the mixture was carefully neutralized with 10 saturated Na₂CO₃, then cooled to room temperature and extracted with Et₂O (3 x 50 mL). The combined organic layers were washed with water (25 mL) followed by brine (25 mL), then dried (Na₂SO₄), filtered and concentrated to give the crude product which was purified by flash column chromatography on a silica gel column using 5%ethyl acetate/hexane as eluent to afford 2,6-dibromo-4-fluoro-3-methylbenzonitrile (1.1474 g, 3.92 mmol, 39% yield) as an off-white solid. ¹H NMR (500 MHz, CDCl₃) δ: 7.41 (1 H, d, *J* = 8.5 Hz), 2.38 (3 H, d, *J* = 2.4 Hz).

15

20


Intermediate 17



(2,6-Dibromo-4-fluoro-3-methylphenyl)methanamine. To a stirred solution of 2,6-dibromo-4-fluoro-3-methylbenzonitrile (1.145 g, 3.91 mmol) in THF (7 mL) was added 2M borane-methylsulfide complex (8 mL, 16.00 mmol) over 5 min at room 25 temperature. After 30 min, the reaction mixture was warmed to 60 °C and stirred for an additional 1.6 h. The mixture was cooled then quenched by the careful addition of methanol (10 mL) and 4M HCl/dioxane (2 mL). After 30 min, the reaction mixture was heated to 50 °C for 45 min and concentrated. The residue was taken up in sat. Na₂CO₃ (20 mL) and extracted with CH₂Cl₂ (3 x 25 mL). The combined CH₂Cl₂ 30 extracts were washed with water (20 mL) followed by brine (20 mL), then dried

(Na₂SO₄), filtered and concentrated to afford (2,6-dibromo-4-fluoro-3-methylphenyl)methanamine (1.16 g, 3.91 mmol, 100% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ: 7.31 (1 H, d, *J* = 8.9 Hz), 4.19 (2 H, s), 2.34 (3 H, d, *J* = 2.4 Hz), 1.59 (2 H, brs). LCMS (M+H) calcd for C₁₅H₂₁N₂O₃: 297.91; found: 5 297.85.

Intermediate 18




10

5-(Aminomethyl)-2-fluorobenzonitrile. A solution of 5-(bromomethyl)-2-fluorobenzonitrile (300 mg, 1.402 mmol) in 7M NH₃/methanol (5 mL, 35.0 mmol) was stirred at room temp for 18h. After the removing the solvent the resulting residue was stirred with Et₂O/sat. Na₂CO₃. The aqueous layer was then separated 15 and the organic layer dried (Na₂SO₄), filtered and concentrated to give a 3:1 mixture of the title compound, amine 5-(aminomethyl)-2-fluorobenzonitrile contaminated and 5,5'-azanediylbis(methylene)bis(2-fluorobenzonitrile). The mixture was used without further purification. ¹H NMR (500 MHz, CDCl₃) δ: 7.48 - 7.66 (2 H, m), 7.07 - 7.21 (1 H, m), 3.89 (2 H, s).

20

Intermediate 19




25 *tert-Butyl 3-bromo-4-fluorobenzylcarbamate.* To a solution of (3-bromo-4-fluorophenyl)methanamine, HCl (1 g, 4.16 mmol) in THF (10 mL)/water (10 mL) at 0 °C was added, portion-wise, sodium bicarbonate (1.397 g, 16.63 mmol). After 5 min, di-*tert*-butyl dicarbonate (1.062 mL, 4.57 mmol) was slowly added and the reaction stirred while gradually warming to room temperature. The solution was

diluted with Et₂O (20 mL) and washed with 1N HCl (2 x 30 mL). The organic layer was dried (MgSO₄) and concentrated. The residue was purified by flash chromatography (Biotage flash chromatography system; 40g SiO₂; 0-20% ethyl acetate:hexane) to give the title compound as a clear oil (1.26 g, 100% yield).

5 ¹H NMR (500 MHz, CDCl₃) δ: 7.49 (1 H, dd, *J* = 6.53, 2.01 Hz), 7.17 - 7.25 (1 H, m), 7.08 (1 H, t, *J* = 8.28 Hz), 4.87 (1 H, brs), 4.27 (2 H, d, *J* = 5.77 Hz), 1.48 (9 H, s).

Intermediate 20

10




tert-Butyl 3-cyclopropyl-4-fluorobenzylcarbamate. Under N₂, a toluene (4.70 mL)/ water (0.235 mL) solution of *tert*-butyl 3-bromo-4-fluorobenzylcarbamate (0.300 g, 0.986 mmol) was treated with potassium phosphate tribasic (0.733 g, 3.45 mmol), cyclopropylboronic acid (0.110 g, 1.282 mmol), and tricyclohexenylphosphene (0.028 g, 0.099 mmol) in was added palladium(II) acetate (0.011 g, 0.049 mmol). The reaction was stirred at 100 °C for 3 h. The solution was diluted with water (15 mL) then extracted with ethyl acetate (2 x 15 mL). The organic extracts were combined, washed with brine (20 mL), then dried (MgSO₄), filtered and concentrated. The residue was purified by flash chromatography (Biotage flash chromatography system; 60 g SiO₂; 0-20% ethyl acetate:hexane over 900 mL) to give the title compound as a white crystalline solid (0.230 g, 88% yield).

15 ¹H NMR (500 MHz, CDCl₃) δ: 6.98 - 7.05 (1 H, m), 6.90 - 6.97 (1 H, m), 6.79 (1 H, dd, *J* = 7.02, 1.53 Hz), 4.75 (1 H, brs), 4.21 (2 H, d, *J* = 5.49 Hz), 2.02 - 2.09 (1 H, m), 1.45 (9 H, s), 0.93 - 1.01 (2 H, m), 0.66 - 0.75 (2 H, m).

20


25

Intermediate 21



5 *(3-Cyclopropyl-4-fluorophenyl)methanamine hydrochloride salt.* To a solution of tert-butyl 3-cyclopropyl-4-fluorobenzylcarbamate (0.220 g, 0.829 mmol) in CH₂Cl₂ (5 mL) was added trifluoroacetic acid (5 mL, 64.9 mmol). The solution was stirred at room temperature for 1h. To the solution was added 2M HCl in Et₂O (5 mL). The solids were collected via filtration to give the title compound as a white solid (0.154 g, 92% yield). ¹H NMR (400 MHz, MeOD) δ: 7.21 - 7.33 (1 H, m), 7.03 - 7.16 (2 H, m), 4.06 (2 H, s), 2.06 - 2.18 (1 H, m), 0.99 - 1.09 (2 H, m), 0.74 - 0.86 (2 H, m). ¹³C NMR (500 MHz, MeOD) δ: 162.09 (1 C, d, *J* = 245.84 Hz), 131.64 (1 C, d, *J* = 14.64 Hz), 128.98 (1 C, d, *J* = 3.08 Hz), 127.33 (1 C, d, *J* = 8.48 Hz), 126.78 (1 C, d, *J* = 4.62 Hz), 115.15 (1 C, d, *J* = 23.12 Hz), 42.43 (1 C, s), 8.03 (1 C, d, *J* = 5.39 Hz), 7.04 (1 C, s).

Intermediate 22




20 *((2R,3S,6R)-3-Acetoxy-6-cyano-3,6-dihydro-2H-pyran-2-yl)methyl acetate.* To (2R,3S,4R)-2-(acetoxymethyl)-3,4-dihydro-2H-pyran-3,4-diyl-diacetate (1.0 g, 3.67 mmol) in anhydrous CH₂Cl₂ (15 mL) was added dropwise, TMSCN (0.744g, 7.51 mmol). The reaction mixture was stirred at 23 °C for 15 min. then BF₃-etherate (4 drops) added. The reaction mixture was stirred at 23 °C for 30 min. The solvent and excess TMSCN were removed *in vacuo* and the crude material purified by silica gel column chromatography (Biotage flash chromatography system) using ethyl acetate : hexane (20:80) to (1:1) resulting a complete separation of the C6-epimeric

diastereomers. The first eluting compound (400 mg, 46% yield) was confirmed by NMR to be ((2R,3S,6S)-3-acetoxy-6-cyano-3,6-dihydro-2H-pyran-2-yl)methyl acetate while the slower eluting compound (260 mg, 30% yield) confirmed by NMR to be ((2R,3S,6R)-3-acetoxy-6-cyano-3,6-dihydro-2H-pyran-2-yl)methyl acetate. ¹H NMR (400 MHz, CDCl₃) δ ppm 6.07 (1 H, dt, *J*=10.1, 2.5 Hz), 5.95 (1 H, ddd, *J*=10.2, 2.0, 1.9 Hz), 5.28 - 5.35 (1 H, m, *J*=8.1, 2.4, 2.2, 2.2 Hz), 5.14 (1 H, q, *J*=2.5 Hz), 4.27 - 4.33 (1 H, m), 4.19 - 4.25 (1 H, m), 3.84 (1 H, ddd, *J*=8.4, 5.9, 2.9 Hz), 2.11 (3 H, s), 2.08 (3H, s).

10


Intermediate 23



((2S,6R)-6-Cyanotetrahydro-2H-pyran-2-yl)methyl acetate. To a dry THF (200 mL) solution containing ((2R,3S,6R)-3-acetoxy-6-cyano-3,6-dihydro-2H-pyran-2-yl)methyl acetate (5.5g, 23 mmol), diphenylsilane (8.47, 46 mmol) and zinc chloride(7.52g, 55.2 mmol) was added after N₂ bubbling, Pd(PPh₃)₄(2.66g, 2.30mmol). The reaction mixture was stirred at 23 °C for 4 h. (monitored by TLC). The reaction mixture was filtered through a short silica gel column using CH₂Cl₂ as solvent. The solvent was removed under reduced pressure and the crude material was purified by a silica gel column (Biotage flash chromatography system) using ethyl acetate : hexane (1:5) to (1:2) to afford 2.8g (67% yield) mixture of ((2S,6R)-6-cyano-5,6-dihydro-2H-pyran-2-yl)methyl acetate and((2S,6R)-6-cyano-3,6-dihydro-2H-pyran-2-yl)methyl acetate. ¹H NMR (400 MHz, CDCl₃) δ ppm 6.06 - 6.15 (0.5 H, m, *J*=7.6, 5.3, 2.8, 2.8 Hz), 5.93 - 6.02 (0.5 H, m, *J*=10.4, 5.2, 2.6, 2.6 Hz), 5.69 - 5.79 (1 H, m), 4.54 (0.5 H, dd, *J*=9.9, 3.8 Hz), 4.48 (0.5 H, m), 4.11 - 4.23 (2 H, m), 3.89 (1 H, m), 2.59 - 2.69 (1 H, m), 2.33 - 2.42 (1 H, m,), 2.16 - 2.22 (1 H, m), 2.14 (3 H, s), 2.12 (3H, s).To the mixture of ((2S,6R)-6-cyano-5,6-dihydro-2H-pyran-2-yl)methyl acetate and ((2S,6R)-6-cyano-3,6-dihydro-2H-pyran-2-yl)methyl acetate. 30 (2.6 g, 14.4 mmol) in ethyl acetate (200 mL) was added Pd/C (5%) (0.200g). The reaction mixture was stirred at 23 °C for 6 h. under a hydrogen atmosphere (balloon).

The Pd/C was then removed by filtration and the solvent evaporated *in vacuo* to afford 2.20g (84% yield) of the target compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 4.28 (1 H, dd, *J*=11.7, 2.7 Hz), 4.03 - 4.14 (2 H, m), 3.59 - 3.67 (1 H, m), 2.11 (3 H, s), 1.94 - 2.04 (2 H, m), 1.80 - 1.91 (1 H, m), 1.54 - 1.65 (2 H, m), 1.33 - 1.45 (1 H, m).


Intermediate 24



10

(2*R*,6*S*)-6-(Hydroxymethyl)tetrahydro-2*H*-pyran-2-carbonitrile. To ((2*S*,6*R*)-6-cyanotetrahydro-2*H*-pyran-2-yl)methyl acetate (3.90g, 21.29 mmol) in methanol (300 mL) was added Et₃N (30 mL). The reaction mixture was stirred at 23 °C for 18 h. The solvents were evaporated *in vacuo*. The residue was redissolved in methanol 15 the solvent removed *in vacuo* (2 x 20 mL). The crude material was then purified on a silica gel column (Biotage flash chromatography system) eluting with a gradient mixture ethyl acetate (100%) to ethyl acetate :methanol (8:2) to afford 2.20g (73% yield) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 4.31 (1 H, dd, *J*=11.6, 2.5 Hz), 3.58 - 3.63 (2 H, m), 3.51 - 3.57 (1 H, m), 2.06 (1 H, s), 1.94 - 2.03 (2 H, m), 1.79 - 1.90 (1 H, m), 1.53 - 1.65 (2 H, m), 1.36 - 1.47 (1 H, m).

Intermediate 25




25

((2*R*,3*S*,6*R*)-3-Acetoxy-6-(N-hydroxycarbamimidoyl)-3,6-dihydro-2*H*-pyran-2-yl)methyl acetate. To (2*R*,6*S*)-6-(hydroxymethyl)tetrahydro-2*H*-pyran-2-carbonitrile (1.90g, 13.46 mmol) in EtOH (11.0 mL) and H₂O (11.0 mL) was added

hydroxylamine (1.24 mL, 20.19 mmol). The reaction mixture was stirred at 80 °C for 3h. The solvent was evaporated *in vacuo* and residual volatile impurities removed by azeotropic distillation with EtOH (2 x 20 mL) to afford 2.10g (90% yield) of the title compound. LCMS (⁺ESI, M+H⁺) m/z 175.

5

Intermediate 26




10 *Diethyl 2-((2R,6S)-6-(hydroxymethyl)tetrahydro-2H-pyran-2-carboximidoxy)maleate.* To ((2R,3S,6R)-3-acetoxy-6-(N-hydroxycarbamimidoyl)-3,6-dihydro-2H-pyran-2-yl)methyl acetate (2.33g, 13.4 mmol) in EtOH (15 mL) was added diethylacetylenedicarboxylate (3.20 mL, 20.1 mmol) and the reaction mixture was stirred at 60 °C for 3h. The solvent was removed *in vacuo* to afford 2.40g (52% yield) of the title compound as an amber oil. LCMS (⁺ESI, M+H⁺) m/z 345.

15 Intermediate 27

20 *Ethyl 5-hydroxy-2-((2S,6S)-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl)-6-oxo-1,6-dihdropyrimidine-4-carboxylate.* Diethyl 2-((2R,6S)-6-(hydroxymethyl)tetrahydro-2H-pyran-2-carboximidoxy) maleate (2.20 g, 6.39 mmol) in trimethylbenzene (25 mL) was heated in a sealed tube at 180 °C for 18 h. The solvent was removed *in vacuo* and the crude material was purified by preparative

HPLC using a C18 column and trifluoroacetic acid in the mobile phase to afford 1.90 g (99% yield) of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 12.52 (1 H, br s), 10.5-11.0 (1 H, br s), 4.46 - 4.57 (3 H, m), 3.70 - 3.82 (3 H, m), 3.56 - 3.64 (1 H, m), 2.24 - 2.33 (1 H, m), 1.96 - 2.07 (1 H, m), 1.58 - 1.76 (4H, m), 1.46 - 1.64 (3 H, m), $J=7.4$ Hz).

Intermediate 28




10

Ethyl (7S,11R)-3-hydroxy-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,11-epoxypyrimido[1,2-a]azocine-2-carboxylate. To ethyl 5-hydroxy-2-((2S,6S)-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl)-6-oxo-1,6-dihydropyrimidine-4-carboxylate (1.90 g, 6.37 mmol) in dry THF (60mL) was added methanesulfonyl chloride (1.49 mL, 19.1 mmol) and di-isopropyl-ethylamine (3.56mL, 20.0 mmol) at 0°C. The reaction mixture was stirred at 23 °C for 3h. The THF was removed *in vacuo*, the crude material was dissolved in ethyl acetate (50 mL) and washed with 1N NaHCO_3 (50 mL). The aqueous solution was then washed with ethyl acetate (3 x 20mL) and the combined organic fractions were dried (MgSO_4) filtered and concentrated *in vacuo* to give a brown oil. The crude oil was dissolved in EtOH:THF (20mL:20mL) and to this was added K_2CO_3 (0.881g, 6.37 mmol) and H_2O (1.0 mL). The reaction mixture was stirred at 65 °C for 17 h. The solvent was removed *in vacuo*, the crude material was dissolved in ethyl acetate (50 mL) and washed with H_2O (50 mL) and HCl aq. 1N (50mL). The acidic aqueous phase was extracted with ethyl acetate (2 x 50 mL). The combined organic fractions were dried (MgSO_4) and concentrated *in vacuo*. The crude material was purified by preparative HPLC using a C18 column and trifluoroacetic acid in the mobile phase to afford the compound 1.16g (65% yield) of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 10.75

(1 H, s), 4.94 (1 H, d, $J=2.3$ Hz), 4.49 - 4.58 (3 H, m), 4.01 - 4.13 (2 H, m), 2.05 - 2.17 (2 H, m,), 1.96 (1 H, dd, $J=13.6, 2.0$ Hz), 1.76 (2 H, m), 1.39 - 1.49 (4 H, m). LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z 281. HRMS (ESI $^+$) calculated for $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}_5$ [$\text{M}+\text{H}^+$] :281.1137; found : 281.1147.

5


Intermediate 29



10 *Dimethyl pyridine-3,5-dicarboxylate hydrochloride.* To pyridine-3,5-dicarboxylic acid (19.9g, 119.1 mmol) in methanol (400 mL) was added of 2,2-dimethoxypropane (160 mL, 1.3 mol) and concentrated HCl (15 mL). The reaction mixture was refluxed for 18h under an atmosphere of Ar. The solvent was evaporated then ether (100 mL) was added to the crude material and the resulting mixture heated to reflux. The mixture was allowed to remain at reflux for 2 hours then s allowed to cool to 23 °C and the precipitate filtered. The precipitate was dried to afford 27.5 g (86% yield) of the title compound as described in J.Org. Chem. Vol. 16, No 10, 1996. LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z : 196.

15

Intermediate 30




20 *Dimethyl piperidine-3,5-dicarboxylate.* To dimethyl pyridine-3,5-dicarboxylate (15.0g, 76.87 mmol) in acetic acid (150 mL) was added platinum oxide (0,250 mg, 1.1 mmol). The reaction mixture was stirred at 23 °C for 36 h under 45 psi of H_2 . PtO_2 was removed by filtration on CELITE®, washed with acetic acid (20 mL) and the solvent removed *in vacuo*. A solution of NaHCO_3 (1M, 100 mL)) was

then added and the organic material was extracted with CH₂Cl₂ (3 x 100 mL). The combined extracts were dried (MgSO₄) and concentrated to afford 13.7g (88.6% yield) of the title compound. LCMS (ESI, M+H⁺) m/z : 202.

5


Intermediate 31



Dimethyl 1-benzylpiperidine-3,5-dicarboxylate. To dimethyl piperidine-3,5-dicarboxylate (7.5 g, 37.3 mmol) in acetonitrile (100 mL) was added K₂CO₃ (20.6 g, 149.1) and benzylbromide (4.43 mL, 37.3 mmol). The reaction mixture was stirred at 23°C for 4 hours. The K₂CO₃ was removed by filtration and the solvent evaporated *in vacuo*. A solution of NaHCO₃ (1M, 150 mL) was added and the organic material extracted with ethyl acetate (3 x 50 mL), then dried (MgSO₄) and concentrated *in vacuo*. The crude material was purified on a Silica gel column (Biotage flash chromatography system), eluting with a gradient of ethyl acetate/hexane (10:90) to (35:65) to afford the title compound as the cis diastereomer 1.6 g which eluted after the corresponding trans diastereomer. (based on the NMR reported in Synthetic Communications 1997, 27 (1), 69-71.). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.26 - 7.36 (5 H, m), 3.68 (6 H, s), 3.59 (2 H, s), 3.15 (2 H, m), 2.66 (2 H, t, J=12.0 Hz), 2.36 (1 H, d, J=13.1 Hz), 2.05 (2 H, t, J=11.4 Hz), 1.56 (1 H, q, J=12.9 Hz). LCMS (ESI, M+H⁺) m/z : 292.

Intermediate 32


25



(1-Benzylpiperidine-3,5-diyl)dimethanol. To dimethyl 1-benzylpiperidine-3,5-dicarboxylate (1.5 g, 5.15 mmol) in dry THF (30 mL) was added a solution of lithium aluminum hydride in THF (2M, 3.87mL, 7.73 mmol) at 0° C. The reaction mixture was stirred at 0 °C for 30 min. then warmed to 23 °C and stirred for 1.5 hour.

- 5 A solution of methanol (0.5 mL) and (0.25 mL) was added followed by a 15% solution of NaOH (0.250 mL). The mixture was stirred for 1 hour at 23°C then filtered. The organic material was extracted with ethyl acetate (3 x 20 mL) and the combined extracts were dried (MgSO_4) and concentrated *in vacuo* to afford 0.91g (75% yield) of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 7.26 - 7.35 (5H, m), 3.59 (2H, s), 3.44 - 3.55 (4 H, m), 3.06 (2 H, dt, $J=10.6, 1.8$ Hz), 1.88 - 1.98 (2 H, m), 1.79 - 1.86 (1 H, m), 1.68 (3 H, m). LCMS ($^+\text{ESI}, \text{M}+\text{H}^+$) m/z : 236.
- 10


Intermediate 33



15


- (1-Benzyl-5-((tert-butyldimethylsilyloxy)methyl)piperidin-3-yl)methanol.* To 1-benzylpiperidine-3,5-diyl)dimethanol (0.5g, 2.125 mmol) in dry THF (20 mL) was added NaH (60 % in mineral oil)(90 mg, 2.23 mmol). The reaction mixture was stirred at 23°C for 45 min. then, t-butyldimethylchlorosilane (0.352g, 2.37 mmol) added and the reaction mixture stirred at 23°C for 2 hours. The mixture was poured into ether (75 mL), washed with 10% aqueous K_2CO_3 (30 mL) then dried (MgSO_4) and concentrated *in vacuo*. The crude material was purified on a Silica gel column (Biotage flash chromatography system), eluting with a gradient of ethyl acetate : hexane (9:1) to ethyl acetate 100% to afford 0.340g (43% yield) of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 7.25 - 7.36 (5 H, m), 3.59 (2 H, s), 3.45 - 3.55 (4 H, m), 2.99 - 3.09 (2 H, m), 1.93-1.55 (5 H, m), 1.24 - 1.34 (1 H, m), 0.87(9 H, s), 0.62 - 0.74 (1 H, m), 0.02 (6 H, s). LCMS ($^+\text{ESI}, \text{M}+\text{H}^+$) m/z : 350.
- 20
- 25

Intermediate 34

5 *1-Benzyl-5-((tert-butyldimethylsilyloxy)methyl)piperidine-3-carbaldehyde.*

To a solution of oxalyl chloride (1.40 mL, 16.02 mmol) in dichloromethane (100mL) at -78°C was added, dropwise, dry DMSO (1.78 mL, 25.03 mmol). After 10 min at -78°C, a solution of 1-benzyl-5-((tert-butyldimethylsilyloxy)methyl)piperidin-3-yl)methanol (3.50g, 19.01 mmol) in CH₂Cl₂ (100mL) was slowly added and the reaction stirred at -78°C. After 1h, triethylamine (6.98 mL, 50.1 mmol) was added and stirring continued at -78°C for 10 min after which the reaction was allowed to warm to 23°C. The resulting solution was neutralized with 1N HCl (pH~6)and the phases separated. The aqueous solution was washed with CH₂Cl₂ (3 x 50 mL) and the combined organic fractions dried over MgSO₄, filtered and concentrated *in vacuo* to afford the title compound 3.39g (76% yield) as a light yellow oil. ¹H NMR (400 MHz, CDCl₃) δ ppm 9.61 (1 H, s), 7.59 - 7.67 (2 H, m), 7.42 - 7.49 (3 H, m), 4.22 - 4.30 (1 H, m), 4.10 - 4.18 (1 H, m), 3.70 - 3.81 (1 H, m), 3.52 (1 H, dt, *J*=10.4, 5.2 Hz), 3.35 (1 H, dd, *J*=11.9, 1.8 Hz), 3.14 (1H, m), 2.76 (1 H, tt, *J*=12.1, 3.9 Hz), 2.52 - 2.58 (1 H, m), 2.39 - 2.48 (1 H, m), 2.17 - 2.28 (1 H, m), 1.43 (1 H, t, *J*=7.3 Hz), 1.18 - 1.27 (1 H, m), 0.83 (9 H, m), 0.01 (6 H, m).


Intermediate 35



(Z)-1-Benzyl-5-(hydroxymethyl)piperidine-3-carbaldehyde oxime. To a mixture of 1-benzyl-5-((tert-butyldimethylsilyloxy)methyl)piperidine-3-carbaldehyde (3.39g, 9.75 mmol) in EtOH (50 mL) was added an aqueous solution of NH₂OH-HCl (3.75M, 3.12 mL, 11.7 mmol) followed by an aqueous solution of NaOAc (1.5M, 5.2 mL, 7.8 mmol). The reaction mixture was stirred at room temperature for 18 h. H₂O (30 mL) was then added and the EtOH was partially removed *in vacuo* (~40 mL). The reaction mixture was diluted with ethyl acetate and washed with saturated NaHCO₃. The combined aqueous fractions were extracted with ethyl acetate (3 x 50 mL) and the combined organic fractions were washed with brine and dried (MgSO₄) and concentrated in *vacuo* to afford 3.31g (94% yield) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.25 - 7.35 (6 H, m), 3.58 (2 H, d, *J*=4.5 Hz), 3.43 - 3.54 (2 H, m), 3.03 (2 H, d, *J*=4.5 Hz), 2.57 - 2.68 (1 H, m) 1.90 - 2.01 (2 H, m), 1.80 - 1.89 (1 H, m), 1.65 - 1.77 (2 H, m), 0.93 - 1.01 (1 H, m). LCMS (ESI, M+H⁺) m/z : 249.

15

Intermediate 36




1-Benzyl-5-(hydroxymethyl)piperidine-3-carbonitrile. To 1-benzyl-5-(hydroxymethyl)piperidine-3-carbaldehyde oxime (3.3 g, 13.30 mmol) in CH₂Cl₂ (15 mL) at 23 °C was added 2-chloro-1-methylpyridine iodide (3.73g, 14.62 mmol) in one portion and the resulting mixture stirred at 23 °C for 10 min. Et₃N was then added and the reaction mixture stirred for 3 hours (color changed from light yellow to light brown). The solution was neutralized with HCl (5%), to approximately pH 6 and the organic material was extracted with CH₂Cl₂ (3 x 25 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo* to afford 3.06g (95% yield) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.28 - 7.37 (5

H, m), 3.51 - 3.60 (4 H, m), 3.11 (1 H, ddd, $J=11.0, 1.9, 1.8$ Hz), 2.98 (1 H, d, $J=7.6$ Hz), 2.73 - 2.82 (1 H, m), 2.12 - 2.22 (2 H, m), 1.80 - 1.92 (2 H, m), 1.45 (1 H, br s), 1.27 - 1.38 (1 H, m). LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z : 231.


5

Intermediate 37



Diethyl 2-(1-((E)-1,4-diethoxy-1,4-dioxobut-2-en-2-yl)-5-(hydroxymethyl)piperidine-3-carboximidooxy)maleate. To a solution of 1-
 10 benzyl-5-(hydroxymethyl)piperidine-3-carbonitrile (2.93 g, 11.14 mmol) in EtOH (45.0 mL) was added diethyl acetylenedicarboxylate (2.67 mL, 16.71 mmol). The mixture was allowed to stir at 60 °C for 2 h. The solvent was removed *in vacuo*. The crude product was purified using a silica gel column (Biotage flash chromatography
 15 system) using ethyl acetate and hexane as eluent in a ratio 60:40 to followed by ethyl acetate : methanol (8:2) to afford 1.21g (21% yield) of the title compound. ^1H NMR (400 MHz, CDCl₃) δ ppm 5.77 (1 H, d, $J=15.4$ Hz), 5.26 (1 H, s), 4.98 (1 H, s), 4.82 (1 H, d, $J=7.3$ Hz), 4.31 - 4.44 (4 H, m), 4.08 - 4.21 (4 H, m), 3.42- 3.66 (2 H, m), 2.92 - 3.03 (1 H, m), 2.62 - 2.73 (1 H, m), 2.44 - 2.55 (1 H, m), 2.04 - 2.14 (2 H, m), 1.86 - 1.96 (2 H, m), 1.23 - 1.41 (12 H, m). LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z : 514.
 20


Intermediate 38



5 *Diethyl 2-(3-(4-(ethoxycarbonyl)-5-hydroxy-6-oxo-1,6-dihydropyrimidin-2-yl)-5-(hydroxymethyl)piperidin-1-yl)maleate.* To diethyl 2-(1-((E)-1,4-diethoxy-1,4-dioxobut-2-en-2-yl)-5-(hydroxymethyl)piperidine-3-carboximidamidoxy)maleate (1.20g, 2.34 mmol) was added xylene (15 mL) and the reaction mixture was stirred at 160 °C for 6 hours in a sealed tube. The solvent was removed *in vacuo* to afford 1.05g (99% yield) of the title compound. LCMS (⁺ESI, M+H⁺) m/z : 468.

10

Intermediate 39




15 *Diethyl 2-(2-(ethoxycarbonyl)-3-hydroxy-4-oxo-7,8,10,11-tetrahydro-4H-7,11-methanopyrimido[1,2-a][1,5]diazocin-9(6H)-yl)-2-butenedioate.* To diethyl 2-(3-(4-(ethoxycarbonyl)-5-hydroxy-6-oxo-1,6-dihydropyrimidin-2-yl)-5-(hydroxymethyl)piperidin-1-yl)maleate (1.05g, 2.32mmol) in THF at 0 °C was added triethylamine (0.975 mL, 6.99 mmol)and methanesulfonyl chloride (0.541 mL, 6.99

mmol). The reaction mixture was stirred at 23 °C for 3 hours. H₂O (50 mL) was then added and the mixture was extracted with ethyl acetate (3 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. To the crude material dissolved in EtOH (40 mL) were added K₂CO₃ (0.643 g, 4.66 mmol) and 5 H₂O (1.7 mL). The reaction mixture was stirred at 23 °C for 36 hours. The solvent was removed *in vacuo* and the crude material was purified by silica gel column chromatography (Biotage flash chromatography system) using a gradient of ethyl acetate:methanol:AcOH (95:4:1) to (40:40:10) to afford 1.05g (33% yield) of the title compound. ¹H NMR (400 MHz, DMSO-D6) δ ppm 3.83-4.26 (7 H, m), 2.68-3.40 (6 H, m), 2.02 (1H, m), 1.79-1.89 (2 H, m), 1.07 - 1.31 (10 H, m). LCMS (⁺ESI, M+H⁺) m/z : 450.


Intermediate 40

15



Ethyl 3-hydroxy-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,11-methanopyrimido[1,2-a][1,5]diazocine-2-carboxylate. To diethyl-2-(2-(ethoxycarbonyl)-3-hydroxy-4-oxo-7,8,10,11-tetrahydro-4H-7,11-methanopyrimido[1,2-a][1,5]diazocin-9(6H)-yl)-2-butenedioate (0.300 g, 0.673 mmol) in acetonitrile (15 mL) was added trifluoroacetic acid (1 mL) at 23 °C. The reaction mixture was stirred at 23 °C for 6 hours. CH₃CN and trifluoroacetic acid were then removed *in vacuo* and the crude material was redissolved in toluene and dried again (2 x 30 mL) to afford 234 mg (90% yield) of the title trifluoroacetic salt.
20 LCMS (⁺ESI, M+H⁺) m/z : 280.


Intermediate 41



5 *9-Benzyl 2-ethyl 3-hydroxy-4-oxo-7,8,10,11-tetrahydro-4H-7,11-*
methanopyrimido[1,2-a][1,5]diazocine-2,9(6H)-dicarboxylate. To ethyl 3-hydroxy-
4-oxo-6,7,8,9,10,11-hexahydro-4H-7,11-methanopyrimido[1,2-a][1,5]diazocine-2-
carboxylate (0.26 g, 0.661 mmol) in CH₂Cl₂ (5 mL) was added
diisopropylethylamine (0.576 mL, 3.31 mmol) and benzylchloroformate (0.102 mL,
10 0.727 mmol). The reaction mixture was stirred at 23 °C for 16 hours. HCl (1N, 15
mL) was then added to the reaction mixture and the organic material extracted with
CH₂Cl₂ (3x 15 mL) and ethyl acetate (3 x 15 mL). The combined organic fractions
were dried (MgSO₄) and concentrated in vacuo. The organic material was purified
by Silica gel column chromatography (Biotage flash chromatography system) eluting
15 with gradient of ethyl acetate 100% to ethyl acetate methanol (1:9) to afford 0.036 g
(13% yield) of the title compound. ¹H NMR (400 MHz, MeOD) δ ppm 7.36 (5 H, s),
4.95-5.10 (2 H, m) 3.95 - 4.40 (2H, br m), 2.75 - 3.30 (5 H, br m), 1.75-2.50 (4 H,
m), 0.95- 1.45 (4 H, m). LCMS (ESI, M+H⁺) m/z : 414.

20


Intermediate 42



(3-((Tert-butyldimethylsilyloxy)methyl)cyclopentyl)methanol. To a solution of NaH (60 % in oil) (3.53g, 88.3 mmol) in dry THF(400 mL) was added cyclopentane-1,3-diyldimethanol (10.0g, 76.8 mmol) in dry THF (100 mL). The reaction mixture was stirred at 23°C for 45 min. A solution of t-butyldimethylchlorosilane (13.89g, 92.2 mmol) in dry THF (30 mL) was then added and the reaction mixture stirred at 23°C for 4 hours. The reaction mixture was poured into ether (150 mL), washed with 10% aqueous K₂CO₃ (30 mL), dried (MgSO₄) and concentrated *in vacuo*. The crude material was purified on a Silica gel column (Biotage flash chromatography system), eluted with ethyl acetate : hexane (1:9) to ethyl acetate (4:6) to afford 12.3 g (66 % yield) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 3.49 - 3.58 (4 H, m), 2.09 - 2.19 (2 H, m), 1.94 (1 H, dt, *J*=12.6, 7.5 Hz), 1.69 - 1.79 (2 H, m), 1.56 (1 H, br s), , 1.32 - 1.44 (2 H, m), 0.91 (9 H, m), 0.06 (6 H, m).


Intermediate 43

15




3-((Tert-butyldimethylsilyloxy)methyl)cyclopentanecarbaldehyde. To a solution of oxalyl chloride (3.44 mL, 39.3 mmol) in dichloromethane (200mL) at -78°C was added, dropwise, dry DMSO (4.36 mL, 61.4 mmol). After 10 min at -78°C, a solution of (3-((tert-butyldimethylsilyloxy)methyl)cyclopentyl)methanol (6.0 g, 24.56 mmol) in CH₂Cl₂ (180mL) was slowly added and the mixture stirred at -78°C. After 1h, triethylamine (17.11 mL, 122.8 mmol) was added and stirring continued at -78°C for 10 min after which the mixture was allowed to warm to 23°C. The reaction mixture was diluted with HCl 1N (pH~6) and the phases separated. The organic fraction was dried (MgSO₄), filtered and concentrated *in vacuo* to give 5.81g (97% yield) of the title compound as a light yellow oil. ¹H NMR (400 MHz, CDCl₃) δ ppm 9.62 (1 H, d, *J*=2.5 Hz), 3.54 (2 H, ddd, *J*=12.0, 10.0, 6.3 Hz), 2.73 - 2.82 (1 H, m), 2.14 - 2.24 (1 H, m), 1.89 - 2.00 (2 H, m), 1.74 - 1.85 (2 H, m), 1.57 (1 H, dt, *J*=13.1, 8.6 Hz), 1.33 - 1.44 (1 H, m), 0.91 (9 H, s), 0.06 (6 H, s).

Intermediate 44



5 *2-((3-((tert-butyldimethylsilyloxy)methyl)cyclopentyl)-2-(methylamino)acetonitrile.* To 3-((tert-butyldimethylsilyloxy)methyl)cyclopentanecarbaldehyde (5.8 g, 23.94 mmol) in H₂O (8 mL) and methanol (10 mL) was added methylamine (.0755g, 23.94 mmol) at 0 °C. This mixture was then added at 0 °C to KCN (1.56g, 23.94 mmol) in H₂O (3 mL). The resulting reaction mixture
10 was stirred at 23 °C for 18 hours. The organic material was extracted with ethyl acetate (3 x 100 mL) and the combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. ¹H NMR (400 MHz, CDCl₃) δ ppm 4.27 - 4.38 (2 H, m), 3.49 - 3.60 (1 H, m), 2.55 (3 H, s), 2.00 - 2.35 (4 H, m), 1.65 - 1.96 (2 H, m), 1.42 - 1.54 (2 H, m), 1.22 (1 H, ddd, *J*=13.4, 9.3, 3.8 Hz), 0.89 (9 H, s), 0.05 (6 H, m). LCMS
15 (⁺ESI, M+H⁺) m/z 283.

Intermediate 45




20
Benzyl (3-((tert-butyldimethylsilyloxy)methyl)cyclopentyl)(cyano)methyl(methyl)-carbamate. To 2-((3-((tert-butyldimethylsilyloxy)methyl)cyclopentyl)-2-(methylamino)acetonitrile (6.0g, 21.24 mmol) in THF (33mL), H₂O (15 mL) was added, at 0, °C Na₂CO₃ (4.50 g, 42.48 mmol) followed by the dropwise addition of benzyl chloroformate (3.89 mL,
25

27.6 mmol) over 10 min. The reaction mixture was stirred at 0 °C for 30 min. then at 23 °C for 18 hours. H₂O (150 mL) was then added and the organic material was extracted with ethyl acetate (3 x 100 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The crude material was purified on a 5 Silica gel column (Biotage flash chromatography system), eluted with a gradient of ethyl acetate:hexane (1:4) to (1:1) to afford 3.90g (44% yield) of the title compound.
¹H NMR (400 MHz, CDCl₃) δ ppm 7.34 - 7.45 (5 H, m), 5.33 (2 H, s), 5.18 - 5.25 (1 H, m), 3.48 - 3.59 (2H, m), 3.00 (3H, s), 2.35 - 2.47 (2H, m), 1.55- 2.25 (6 H, m), 0.94 (9 H, s), 0.05 (6 H, s). LCMS (ESI, M+H⁺) m/z 417.


10

Intermediate 46



15 *Benzyl cyano(3-(hydroxymethyl)cyclopentyl)methyl(methyl)carbamate*. To benzyl (3-((tert-butyldimethylsilyloxy)methyl)cyclopentyl)(cyano)methyl(methyl)-carbamate (2.0 g, 4.80 mmol) in dry THF (40 mL) cooled to 0 °C was added tetrabutyl-ammoniumfluoride (1M in THF 17.28 mL). The reaction mixture was stirred at 23 °C for 5 hours. H₂O (20 mL) was added to the reaction mixture and the 20 organic material extracted with ethyl acetate (3 x 30 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The crude material was purified on a Silica gel column (Biotage flash chromatography system), eluted with a gradient of ethyl acetate:hexane (3:7) to ethyl acetate 100% to afford 0.550g (38% yield) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.39 (5 H, m), 5.13 - 5.25 (2 H, m), 5.07 (1 H, br s), 3.50-3.60 (2 H, m), 3.01 (3 H, s), 1.30- 2.52 (8 H, m). LCMS (ESI, M+H⁺) m/z 303.


Intermediate 47



5 *Benzyl 2-amino-2-(hydroxyimino)-1-(3-(hydroxymethyl)cyclopentyl)ethyl(methyl)carbamate.* To a solution of benzyl cyano(3-(hydroxymethyl)cyclopentyl)methyl(methyl)carbamate (0.540g, 1.79 mmol) in ethanol (20 mL) was added 50% aqueous solution of hydroxylamine (0.07mL, 2.15 mmol). The reaction mixture was stirred at 60 °C for 18 hours. The
10 solvent was removed *in vacuo* to afford 0.610g (76% yield) of title compound ¹H NMR (400 MHz, CDCl₃) δ ppm 7.33 - 7.43 (5 H, m), 7.14 - 7.23 (2 H, m), 5.18 (2 H, br s), 4.87-4.93 (1 H, m), 4.24 (1 H, m), 3.49 - 3.60 (2 H, m), 2.83 - 2.94 (3 H, br s), 2.60 - 2.71 (1 H, m), 2.11 - 2.49 (2 H, m), 1.60 - 2.11 (3 H, m), 1.25 - 1.55 (2 H, m).
(+ESI, M+H⁺) 2 peaks m/z 336.

15

Intermediate 48




20

Diethyl 2-(1-amino-2-((benzyloxycarbonyl)(methyl)amino)-2-(3-(hydroxymethyl)cyclopentyl)ethylideneaminoxy)maleate. To benzyl 2-amino-2-(hydroxyimino)-1-(3-(hydroxymethyl)cyclopentyl)ethyl(methyl)carbamate (0.800g, 1.78 mmol) in EtOH (40 mL) was added diethylacetylenedicarboxylate (0.428 ml, 2.67 mmol). The reaction mixture was stirred at 23 °C for 2 days. The solvent was removed *in vacuo* and the crude material purified on a Silica gel column (Biotage flash chromatography system), eluted with a gradient of ethyl acetate : hexane (1:5) to (1:1) to afford 0.570 g (52% yield) of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 7.33 - 7.42 (5 H, m), 5.17 - 5.25 (3 H, m), 4.12 - 4.40 (4 H, m), 3.55 (2 H, m), 2.83 - 2.94 (4 H, m), 2.70-2.75 (1H, m), 2.17 - 2.09 (2 H, m), 1.82 - 2.00 (3 H, m), 1.21 - 1.342 (7 H, m). LCMS ($^{\dagger}\text{ESI}, \text{M}+\text{H}^+$) m/z 506.


Intermediate 49

15



Ethyl 2-(((benzyloxycarbonyl)(methyl)amino)(3-(hydroxymethyl)cyclopentyl)methyl)-5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate. To the diethyl 2-(1-amino-2-((benzyloxycarbonyl)(methyl)amino)-2-(3-(hydroxymethyl)cyclopentyl)ethylideneaminoxy)maleate (0.710 g, 1.40 mmol) was added xylene (30 mL) and the reaction mixture was stirred at 120°C for 36 hours. Xylene was removed *in vacuo* to afford 0.53g (82% yield) of the title compound. LCMS ($^{\dagger}\text{ESI}, \text{M}+\text{H}^+$) m/z 460.


Intermediate 50



5 *Ethyl 11-((benzyloxy)carbonyl)(methyl)amino)-3-hydroxy-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocine-2-carboxylate.*

To ethyl 2-((benzyloxycarbonyl)(methyl)amino)(3-(hydroxymethyl)cyclopentyl)methyl-5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate (0.235 g, 0.511 mmol)in THF (30 mL) at 0 °C was added triethylamine (0.214 mL, 1.53 mmol) and methanesulfonyl chloride (0.119 mL, 1.53 mmol). The reaction mixture was stirrer at 23 °C for 3 hours. H₂O (50 mL) was then added and the reaction mixture extracted with ethyl acetate (3 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated in vacuo. LCMS (⁺ESI, M+H⁺) m/z 694. To the crude ethyl 2-((benzyloxycarbonyl)(methyl)amino)(3-((methylsulfonyloxy)methyl)cyclopentyl)methyl-1-(methylsulfonyl)-5-(methylsulfonyloxy)-6-oxo-1,6-dihdropyrimidine-4-carboxylate (0.252 g, 0.363 mmol) dissolved in DMF(40 mL) was added Cs₂CO₃ (0.590g, 1.816 mmol). The reaction mixture was stirred at 80 °C for 12 hours. HCl (1N, 40 ml) was added and the organic material was extracted with ethyl acetate (3 x 30 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo* to afford the title compound. LCMS (⁺ESI, M+H⁺) m/z 442.


Intermediate 51



5 *Ethyl 3-(benzoyloxy)-11-(((benzyloxy)carbonyl)(methyl)amino)-4-oxo-
6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocine-2-carboxylate.* To
Ethyl 11-((benzyloxy)carbonyl)(methyl)amino-3-hydroxy-4-oxo-6,7,8,9,10,11-
hexahydro-4H-7,10-methanopyrimido[1,2-a]azocine-2-carboxylate in pyridine (5
mL) was added benzoic anhydride (0.136 g, 0.600 mmol) and the reaction mixture
10 was stirred at 23 °C for 2 days. Pyridine was then removed under vacuum. H₂O (25
mL) was added and the organic material was extracted with ethyl acetate (3 x 25
mL). The combined organic fractions were dried (MgSO₄) and concentrated *in
vacuo*. The crude material was purified using preparative HPLC on a LUNA C18
column to afford 0.033g (12 % yield) of the title compound. ¹H NMR (400 MHz,
15 CDCl₃) δ ppm 8.20 (2 H, d, *J*=7.8 Hz), 7.67 (1 H, t, *J*=6.9 Hz), 7.52 (2 H, t, *J*=7.7
Hz), 7.28 - 7.41 (4H, m), 5.15 - 5.27 (2 H, m), 4.22 - 4.33 (2 H, m), 3.47 (1H, m),
3.15 (3 H, m), 2.65 (2 H, s), 2.11 (1 H, m), 2.03 (1 H, d, *J*=4.8 Hz), 1.60-1.89 (3 H,
m), 1.25-1.55 (3 H, m), 1.16 (3 H, m). LCMS (+ ESI, M+H⁺) m/z 546. HRMS (ESI⁺)
calculated for C₃₀H₃₁N₃O₇ [M+H⁺]: 546.2240 found: 546.2240.

20


Intermediate 52



8-Amino-1,4-dioxaspiro[4.5]decane-8-carbonitrile. To a stirred solution of NH₄OH (250 mL) and methanol (250 mL) was added 1,4-cyclohexanedione monoethylene ketal (29.25 g, 186.64 mmol), NH₄Cl (19.97 g, 373 mmol) and NaCN (18.28 g, 373 mmol). After 16 h at room temperature, the reaction mixture was extracted with CH₂Cl₂ (250 mL x 2), and the combined organic fractions were washed with water (500 mL), dried (Na₂SO₄), filtered and concentrated to afforded the title compound as a colorless liquid (29 g, 85% yield). ¹H NMR (500 MHz, CDCl₃) δ: 3.90-3.96 (m, 4H), 2.01-2.04 (m, 2H), 1.80-1.97 (m, 6H), 1.75 (s, 2H). LCMS (M+H) = 183.23.


10

Intermediate 53



15 Benzyl 8-cyano-1,4-dioxaspiro[4.5]decan-8-ylcarbamate. Benzyl chloroformate (24.18 mL, 169.4 mmol) was added dropwise to a biphasic system of water (300 mL) and CH₂Cl₂ (150 mL) containing 8-amino-1,4-dioxaspiro[4.5]decane-8-carbonitrile (28 g, 154 mmol) and Na₂CO₃•H₂O (22.9 g, 185 mmol) at 0 °C and stirred for 2 h. The mixture was then allowed to warm to room 20 temperature and stirred for 18 h. The mixture was then diluted with CH₂Cl₂ (250 mL), washed with sat. NaHCO₃ (200 mL), water (200 mL), followed by brine (100 mL) then dried (Na₂SO₄), filtered and concentrated to give a brown oil. Flash chromatography on silica gel (using a 5% to 40% ethyl acetate-hexane gradient) afforded the title compound as a white solid (45 g, 92% yield). ¹H NMR (500 MHz, CDCl₃) δ: 7.30-7.35 (m, 5H), 5.11 (s, 2H), 4.90 (brs, 1H), 3.86-3.96 (m, 4H), 2.36-2.41 (m, 2H), 1.84-2.01 (m, 4H), 1.73-1.76 (m, 2H). LCMS (M+H) = 317.30.


Intermediate 54



5 *Ethyl 3-(8-(benzyloxycarbonylamino)-1,4-dioxaspiro[4.5]decan-8-yl)-5-(2-ethoxy-2-oxoethyl)-2,5-dihydro-1,2,4-oxadiazole-5-carboxylate.* A mixture of benzyl 8-cyano-1,4-dioxaspiro[4.5]decan-8-ylcarbamate (37.4 g, 118.33 mmol), hydroxylamine (7.98 mL, 130 mmol, 50 wt% in water) in EtOH (750 mL) was heated at 70 °C for 48 h. The mixture was then cooled to room temperature and treated with 10 diethyl acetylenedicarboxylate (22.73 mL, 142 mmol). After 48 h at room temperature, the reaction mixture was concentrated and the resulting yellow residue was taken up in ethyl acetate (500 mL), washed with water (2 x 100 mL) followed by brine (100 mL), then dried (Na_2SO_4), filtered and concentrated to give a yellow oil. Flash chromatography on silica gel (using 5% to 50% ethyl acetate-hexane gradient) 15 afforded the title compound as a yellow oil (50 g, 82% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.33-7.37 (m, 5H), 5.80 (s, 1H), 5.07 (s, 2H), 4.84 (d, 1H, J = 23.8 Hz), 4.33 (q, 1H, J = 7.02 Hz), 4.27 (q, 1H, J = 7.22 Hz), 4.09-4.20 (m, 3H), 3.93 (s, 2H), 3.90-3.92 (m, 2H), 2.16-2.23 (m, 4H), 2.03 (s, 1H), 1.62-1.74 (m, 4H), 1.24-1.37 (m, 6H). LCMS ($\text{M}+\text{H}$) = 336.14.

20

Intermediate 55




Ethyl 2-(8-(benzyloxycarbonylamino)-1,4-dioxaspiro[4.5]decan-8-yl)-5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate. A solution of ethyl 3-(8-(benzyloxycarbonylamino)-1,4-dioxaspiro[4.5]decan-8-yl)-5-(2-ethoxy-2-oxoethyl)-2,5-dihydro-1,2,4-oxadiazole-5-carboxylate (6.2 g, 11.9 mmol) in xylenes (300 mL)

5 was placed in a pre-heated oil bath and stirred for 16 h at 140 °C. The reaction mixture was then cooled and extracted with 0.2 M aq. NaOH (3 x 100 mL) and the combined aqueous fractions were extracted with ethyl acetate (250 mL). The aqueous layer was then acidified with conc. HCl and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ fractions were dried (Na₂SO₄), filtered and concentrated to afford

10 the title compound as a light yellow solid (4.1 g, 70% yield). ¹H NMR (500 MHz, DMSO-D₆) δ: 12.55 (s, 1H), 10.25 (s, 1H), 7.28-7.36 (m, 6H), 4.99 (s, 2H), 4.29 (q, 2H, J = 7.5 Hz), 3.85 (brs, 4H), 2.03-2.08 (m, 4H), 1.71-1.76 (m, 2H), 1.29 (t, 3H, J = 7.5 Hz). LCMS (M+H) = 474.48.

15

Intermediate 56



Ethyl 2-(1-(benzyloxycarbonylamino)-4-oxocyclohexyl)-5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate. A solution of ethyl 2-(8-(benzyloxycarbonylamino)-1,4-dioxaspiro[4.5]decan-8-yl)-5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate (1.8 g, 3.8 mmol) in THF (200 mL) was added 2N HCl (12 ml, 24 mmol) and the resulting mixture was heated for 2 h at 60 °C. The reaction mixture was then cooled and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ fractions were dried (Na₂SO₄), filtered and concentrated to afford the title compound as a light yellow solid (1.5 g, 92% yield). ¹H NMR (500 MHz, CDCl₃) δ:


20 12.11 (s, 1H), 11.09 (s, 1H), 7.17-7.29 (m, 5H), 6.36 (s, 1H), 5.00 (s, 2H), 4.44 (q,

25 1.88 (s, 2H), 1.65 (s, 2H), 1.55 (s, 2H), 1.45 (s, 2H), 1.35 (s, 2H), 1.25 (s, 2H), 1.15 (s, 2H), 1.05 (s, 2H), 0.95 (s, 2H), 0.85 (s, 2H), 0.75 (s, 2H), 0.65 (s, 2H), 0.55 (s, 2H), 0.45 (s, 2H), 0.35 (s, 2H), 0.25 (s, 2H), 0.15 (s, 2H), 0.05 (s, 2H).

2H, J = 7.02 Hz), 2.49-2.57 (m, 8H), 1.42 (q, 3H, J = 7.05 Hz). LCMS (M+H) = 430.38.


Intermediate 57

5




Ethyl 2-(1-(benzyloxycarbonylamino)-4-hydroxycyclohexyl)-5-hydroxy-1,6-dihdropyrimidine-4-carboxylate. To a solution of ethyl 2-(1-(benzyloxycarbonylamino)-4-oxocyclohexyl)-5-hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate (1.5 g, 3.5 mmol) in methanol (35 mL) at room temperature was added NaBH₄ (138 mg, 3.66 mol) and the resulting mixture was stirred for 1 h. Methanol was then removed under reduced pressure and the mixture was diluted with CH₂Cl₂ (200 mL), washed with 1N HCl (100 mL) followed by brine (100 mL). The organic fraction was dried (Na₂SO₄), filtered and concentrated to afford the title compound as a light yellow solid (1.3 g, 86% yield). ¹H NMR (500 MHz, DMSO-D₆) δ: 12.46 (s, 1H), 10.24 (s, 1H), 7.31-7.35 (m, 5H), 7.11 (brs, 1H), 4.28-4.32 (m, 2H), 4.29 (q, 2H, J = 7.22 Hz), 3.43-3.46 (m, 1H), 2.31-2.33 (m, 1H), 2.15-2.22 (m, 2H), 1.77-1.83 (m, 2H), 1.65-1.67 (m, 2H), 1.44-1.49 (m, 2H), 1.28 (t, 20 3H, J = 7.22 Hz). LCMS (M+H) = 432.37.

Intermediate 58



5 *Ethyl 2-(1-(benzyloxycarbonylamino)-4-(methylsulfonyloxy)cyclohexyl)-5-(methylsulfonyloxy)-1,6-dihdropyrimidine-4-carboxylate.* To a stirred solution of ethyl 2-(1-(benzyloxycarbonylamino)-4-hydroxycyclohexyl)-5-hydroxy-1,6-dihdropyrimidine-4-carboxylate (1.75 g, 4.05 mmol) in THF (25 mL) was added methanesulfonyl chloride (0.94 mmol, 12.17 mmol) and triethylamine (1.7 mL, 12.75 mmol) at 5 °C and the mixture stirred at room temperature for 18 h. The resulting white precipitate was filtered off and washed with THF. The filtrate and washings were concentrated to afford an amber oil. A solution of this oil in methanol (20 mL) was treated with K₂CO₃ (560 mg, 4.05 mmol) at room temperature for 3 h. The mixture was then concentrated and the residue dissolved in ethyl acetate (200 mL), washed with 1N HCl (100 mL) followed by brine (100 mL) and then dried (Na₂SO₄), filtered and concentrated to give 2.3 g of the title compound as an amber oil, which was used in the next step without further purification. LCMS (M+H) = 588.41.


Intermediate 59



- 5 *6,9-Ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxylic acid, 6,7,8,9-tetrahydro-3-hydroxy-4-oxo-9-[(phenylmethoxy)carbonyl]amino]-, ethyl ester.* A mixture of ethyl 2-(1-(benzyloxycarbonylamino)-4-(methylsulfonyloxy)cyclohexyl)-5-(methylsulfonyloxy)-1,6-dihdropyrimidine-4-carboxylate (2.35 g, 4 mmol) and cesium carbonate (1.3 g, 4 mmol) in DMF (25 mL) was heated at 60 °C for 18 h.
- 10 The reaction mixture was then cooled and concentrated under vacuum. The residue was taken up in to ethyl acetate (200 mL) and extracted with 0.2 M aq. NaOH (3 x 100 mL). The combined aqueous fractions were extracted with ethyl acetate. The aqueous layer was then acidified with conc. HCl and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ fractions were dried (Na₂SO₄), filtered and concentrated to afford 15 the title compound as a brown solid (800 mg). ¹H NMR (500 MHz, CDCl₃) δ: 10.81 (s, 1H), 7.30-7.40 (m, 6H), 6.74 (s, 1H), 5.30 (s, 1H), 5.13 (s, 2H), 4.46 (q, 2H, J = 7.17 Hz), 3.05-3.10 (m, 2H), 2.15-2.22 (m, 2H), 1.99-2.04 (m, 2H), 1.52-1.55 (m, 2H), 1.43 (t, 3H, J = 7.17). LCMS (M+H) = 414.07.

20

Intermediate 60




- 25 *Ethyl 1,4-dioxaspiro[4.5]decane-8-carboxylate.* A mixture of ethyl 4-oxocyclohexanecarboxylate (500 g, 2.94 mol), ethane-1,2-diol (656 mL, 11.76 mol)

and p-toluene sulfonic acid (10.06 g, 0.053 mol) was stirred at room temperature under nitrogen atmosphere for 3 days. After pouring in to diethyl ether (1 L), the mixture was washed with water (500 ml), dilute NaHCO₃ solution (300 mL) followed by brine (500 mL). The organic layer was then dried (Na₂SO₄) filtered and concentrated to afford (590 g, 94% yield) the title compound as a colorless oil. Ref. *J.Org. Chem.* **1997**, *62*, 5284-5292.

Intermediate 61

10



1,4-Dioxaspiro[4.5]decan-8-ylmethanol. A 12-L, four-necked flask equipped with a mechanical stirrer, dropping funnel, thermometer and a nitrogen-inlet tube was flushed with nitrogen and then charged with a 1.0 M LAH (2754 mL, 2754 mmol).


15 The mixture was cooled (0-4 °C internal temperature, 2-propanol/dry ice bath) and a solution of ethyl 1,4-dioxaspiro[4.5]decane-8-carboxylate (590 g, 2754 mmol) was added over 3.5 h via dropping funnel. The reaction mixture was then stirred for 3h while allowing it to warm to room temperature then stirred for an additional 16 h at room temp. The reaction mixture was then cooled (0 °C) bath and diluted with ethyl ether (3 L). The reaction mixture was then carefully quenched by adding water (100 mL, over 1h), followed by 15% aqueous NaOH (100 mL, over 40 min) and water (300 ml, over 30 min). After the mixture was stirred for 1h, sodium sulfate was added and stirring continued for 10 min. The white precipitate was the filtered off and the cake was washed with ether and the filtrate concentrated to afford the title

20 compound (440g, 92% yield) as a light yellow liquid. Ref. *J.Org. Chem.* **1997**, *62*, 5284-5292.

25

Intermediate 62


30



8-(Chloromethyl)-1,4-dioxaspiro[4.5]decane. SOCl_2 (37.1 mL, 509 mmol, 1.3 equiv) was slowly added to a stirred solution of 1,4-dioxaspiro[4.5]decan-8-ylmethanol (67.3 g, 391 mmol) and pyridine (94.8 mL, 1175 mmol, 3 equiv.) in CHCl_3 (400 mL). Caution: The reaction is exothermic. The resulting yellow mixture was heated at reflux for 1.5 h and concentrated. The residue was partitioned between water (500 mL) and ether (500 mL). The layers were separated and the aqueous layer was re-extracted with ether (200 mL). The combined ether layers were washed with 1N HCl (2 x 100 mL), water (200 mL) followed by brine (200 mL), then dried (MgSO_4), filtered and concentrated to give the title compound as a dark amber oil (65.7 g, 88% yield). ^1H NMR (500 MHz, CDCl_3) δ : 3.94 (4 H, t, $J = 2.9$ Hz), 3.41 (2 H, d, $J = 6.4$ Hz), 1.83 - 1.88 (2 H, m), 1.75 - 1.79 (2 H, m), 1.63 - 1.73 (1 H, m), 1.55 (2 H, td, $J = 4.3, 13.2$ Hz), 1.29 - 1.39 (2 H, m).

Intermediate 63

15




Benzyl 4-(chloromethyl)-1-cyanocyclohexylcarbamate. To a stirred solution of 8-(chloromethyl)-1,4-dioxaspiro[4.5]decane (78 g, 409 mmol) in THF (200 mL) was added 1N aq. HCl (500 mL, 500 mmol) at room temperature. After 24 h, 30% NH_4OH (106 mL, 818 mmol) followed by NaCN (22.05 g, 450 mmol) were added and stirring continued for an additional 72 h at rt. The reaction mixture was then extracted with CH_2Cl_2 (5 x 100 mL). The combined CH_2Cl_2 extracts were treated with Na_2CO_3 (50.7 g, 409 mmol) in water (200 mL) followed by 25 benzylchloroformate (58.4 mL, 409 mmol) at 0 °C and the resulting mixture allowed to warm to room temp and stirred overnight (16 h). After this, the reaction mixture was washed with sat. NaHCO_3 (200 mL), then dried (Na_2SO_4), filtered and concentrated to afford a yellow residue. To this residue was added 20% ethyl acetate/hexane (150 mL) and the precipitated product was triturated with hexane

(500 mL) and filtered. The filter cake was washed with hexane to afford benzyl 4-(chloromethyl)-1-cyanocyclohexylcarbamate (77 g, 251 mmol, 61% yield) as a pale yellow solid. ¹H NMR (500 MHz, CDCl₃) δ: 7.31 - 7.40 (5 H, m), 5.14 (2 H, s), 4.92 (0.8 H, br.s.), 4.78 (0.2 H, br.s.), 3.42 (2 H, dd, *J* = 6.6, 2.3 Hz), 2.55 - 2.61 (2 H, m), 5 2.21 - 2.28 (1 H, m), 2.04 (2 H, s), 1.30 - 1.83 (4 H, m). LCMS (M+H) calcd for C₁₆H₂₀ClN₂O₂: 307.12; found: 307.07.

Intermediate 64


10



Ethyl 10-((benzyloxy)carbonyl)amino-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate. A solution of benzyl 4-(chloromethyl)-1-cyanocyclohexylcarbamate (10 g, 32.6 mmol) and 50% aq. 15 hydroxylamine (19.97 mL, 326 mmol) in THF (120 mL) was heated at reflux for 5 h. It was then cooled, diluted with ethyl acetate (300 mL), washed with water (2 x 50 mL) followed by brine (100 mL). The organic layer was then concentrated to half its original volume. Water (2 mL) and diethyl but-2-yne dioate (5.22 mL, 32.6 mmol) were then added and the mixture stirred at room temperature 24 h. The reaction 20 mixture was diluted with Et₂O (150 mL), washed with water (2 x 50 mL) followed by brine (75 mL), then dried (Na₂SO₄), filtered and concentrated to give a yellow oil (17.2 g). A solution of the oil in xylene (250 mL) was heated at 125 °C (internal 25 temperature) until reaction was complete (5-12 h). It was then cooled, concentrated and the resulting yellow residue stirred with K₂CO₃ (9.21 g, 66.7 mmol) in DMF (300 mL) at 90 °C for 3 h. After cooling it was diluted with Et₂O (300 mL), extracted with water (100 mL) and 0.1 N NaOH (100 mL). The organic layer was discarded and the aqueous layer carefully acidified with conc. HCl and extracted with Et₂O (3 x 300 mL). The combined ether layers were dried (Na₂SO₄), filtered and concentrated. The crude material was then crystallized or precipitated with ethyl

acetate to give the title compound as a light yellow solid (3.4g, 24% yield). ^1H NMR (500 MHz, CDCl_3) δ : 10.81 (s, 1H), 7.29-7.36 (m, 6H), 5.29 (s, 1H), 5.10 (s, 2H), 4.41 (q, 2H, J = 7.27 Hz), 4.13 (d, 2H, J = 3.66 Hz), 2.43-2.47 (m, 2H), 1.94-2.02 (m, 2H), 1.79-1.85 (m, 2H), 1.65-1.72 (m, 2H), 1.40 (t, 3H, J = 7.17). LCMS ($M+\text{H}$) = 5 428.26.

Intermediate 65




10

4-(Hydroxymethyl)cyclohexanone. Prepared according to general procedure described in Journal of Organic Chemistry 1997, 62, 5284-5292 from commercially available ethyl 4-oxocyclohexanecarboxylate.

15


Intermediate 66



1-Amino-4-(hydroxymethyl)cyclohexanecarbonitrile. To a stirred solution of 20 NH_4OH (125 mL) and methanol (50mL) was added 4-(hydroxymethyl)cyclohexanone (10 g, 78 mmol), NH_4Cl (8.35 g, 156 mmol) and NaCN (7.65 g, 156 mmol). After 16 h at room temperature, the reaction mixture was extracted with CH_2Cl_2 (250 mL x 2), and 4:1 chloroform:2-propanol, and the combined organic fractions washed with water (500 mL), dried (Na_2SO_4), filtered and concentrated to afforded the title compound as a colorless liquid (8 g, 67% yield).


25 ^1H NMR (500 MHz, CDCl_3) δ : 3.51 (d, 2H, J = 4.58 Hz), 2.07-2.09 (m, 2H), 1.86-1.91 (m, 4H), 1.68 (s, 2H), 1.47-1.53 (m, 3H), 1.41 (brs, 1H), 1.30-1.36 (m, 2H). LCMS ($M+\text{H}$) = 155.18.

Intermediate 67



5 *1-Cyano-4-(hydroxymethyl)cyclohexylcarbamate.* Benzyl chloroformate (8.15 mL, 57.1 mmol) was added dropwise to a biphasic mixture of water (110 mL) and CH₂Cl₂ (55 mL) containing 1-amino-4-(hydroxymethyl)cyclohexanecarbonitrile (7g, 45.4 mmol) and Na₂CO₃, H₂O (5.63 g, 45.4 mmol) at 0 °C then stirred for 2h. The mixture was then allowed to warm to room temperature and stirred for 18 h. The
 10 mixture was diluted with CH₂Cl₂ (200 mL), washed with sat. NaHCO₃ (100 mL), water (100 mL) followed by brine (100 mL), then dried (Na₂SO₄), filtered and concentrated to give brown oil. Flash chromatography on silica gel (using 5% to 40% ethyl acetate/hexane gradient) afforded the title compound as a light yellow viscous oil (8 g, 61% yield). ¹H NMR (500 MHz, CDCl₃) δ: 7.33-7.36 (m, 5H), 5.13 (s, 2H), 5.08 (brs, 1H), 3.50 (t, 2H, J = 5.8 Hz), 2.52-2.56 (m, 2H), 1.88-1.91 (m, 2H), 1.61-1.63 (m, 1H), 1.36-1.48 (m, 4H). LCMS (M+H) = 289.19.


Intermediate 68



20 *Diethyl (2Z)-2-(((1-(((benzyloxy)carbonyl)amino)-4-(hydroxymethyl)cyclohexyl)(imino)methyl)amino)oxy)-2-butenedioate.* A mixture of 1-cyano-4-(hydroxymethyl)cyclohexylcarbamate (6.5 g, 22.54 mmol), 50% aq.
 25 hydroxylamine (1.66 mL, 27.05 mmol) in EtOH (150 mL) was heated at 70 °C for 18 h. The mixture was then cooled to room temperature and treated with diethyl

acetylenedicarboxylate (4.33 mL, 27.1 mmol). After 18 h at room temperature, the reaction mixture was concentrated and the resulting yellow residue was taken up in ethyl acetate (250 mL), washed with water (2 x 100 mL) followed by brine (100 mL), then dried (Na_2SO_4), filtered and concentrated to give yellow oil. Flash column chromatography on silica gel (using 5% to 50% ethyl acetate-hexane gradient) afforded the title compound as a mixture of isomers and as a yellow oil (8 g, 72% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.30-7.36 (m, 6H), 5.59 (s, 0.5 H), 5.75 (s, 0.5 H), 5.13 (s, 1H), 5.04 (s, 2H), 4.32-4.36 (m, 1H), 4.24-4.29 (m, 1H), 4.09-4.17 (m, 4H), 3.46-3.51 (m, 2H), 2.54-2.58 (m, 2H), 1.45-1.57 (m, 5H), 1.23-1.26 (m, 6H). LCMS ($\text{M}+\text{H}$) = 492.36.

Intermediate 69




15

Ethyl 2-(1-(benzyloxycarbonylamino)-4-(hydroxymethyl)cyclohexyl)-5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate. A solution of diethyl (2Z)-2-(((1-((benzyloxycarbonyl)amino)-4-(hydroxymethyl)cyclohexyl)(imino)methyl)amino)oxy)-2-butenedioate (7 g, 14.24 mmol) in xylenes (400 mL) was placed in a pre-heated oil bath and stirred for 16 h at 140 °C. The reaction mixture was then cooled, extracted with 0.2 M aq. NaOH (3 x 100 mL) and the combined aqueous layers washed with ethyl acetate (250 mL). The aqueous layer was acidified with conc. HCl and extracted with CH_2Cl_2 (3 x 100 mL). The CH_2Cl_2 extracts were dried (Na_2SO_4), filtered and concentrated to afford the title compound as a light yellow solid (3.1 g, 49% yield). ^1H NMR (500 MHz, DMSO-D6) δ : 12.00 (s, 1H), 10.77 (s, 1H), 7.24-7.36 (m, 6H), 5.45 (brs, 1H), 5.01 (s, 2H),

4.39-4.45 (m, 2H), 3.50 (d, 2H, , $J = 6.41$ Hz), 2.51-2.62 (m, 2H), 1.64-1.75 (m, 5H), 1.44-1.51 (m, 2H), 1.40-1.44 (m, 3H). LCMS ($M+H$) = 446.27.

Intermediate 70

5




Ethyl 2-(1-(benzyloxycarbonylamino)-4-

((methylsulfonyloxy)methyl)cyclohexyl)-5-(methylsulfonyloxy)-1,6-

10 *dihydropyrimidine-4-carboxylate.* To a stirred solution of ethyl 2-(1-(benzyloxycarbonylamino)-4-(hydroxymethyl)cyclohexyl)-5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate (1.85 g, 4.15 mmol) in THF (40 mL) cooled to 0 °C was added methanesulfonyl chloride (0.96 mmol, 12.45 mmol) and triethylamine (1.79 mL, 12.87 mmol). The mixture was allowed to warm to room temperature and stirred for 18 h. The resulting precipitate was filtered off and washed with THF. The filtrate and washings were concentrated to afford an amber oil. A solution of this oil in methanol (10 mL) was treated with K₂CO₃ (574 mg, 4.15 mmol) at room temperature for 3 h. The mixture was then concentrated and the residue was dissolved in ethyl acetate (200 mL), washed with 1N HCl (100 mL) followed by brine (100 mL), then dried (Na₂SO₄), filtered and concentrated to give 2.3 g of title compound as a yellow oil. LCMS ($M+H$) = 602.45.


Intermediate 64



- 5 *Ethyl 10-((benzyloxy)carbonyl)amino-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate.* A mixture of ethyl 2-(1-(benzyloxycarbonylamino)-4-((methylsulfonyloxy)methyl)cyclohexyl)-5-(methylsulfonyloxy)-1,6-dihydropyrimidine-4-carboxylate (2.25 g, 3.74 mmol) and cesium carbonate (1.22 g, 3.74 mmol) in DMF (40 mL) was heated at 60 °C for 18 h.
- 10 The reaction mixture was then cooled and concentrated *in vacuo*. The residue was taken up in ethyl acetate (200 mL) and extracted with 0.2 M aq. NaOH (3 x 50 mL). The combined aqueous phase was washed with ethyl acetate (250 mL). The aqueous layer was then acidified with conc. HCl and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ extracts were dried (Na₂SO₄), filtered and concentrated to afford the title compound as a brown solid (400 mg). ¹H NMR (500 MHz, CDCl₃) δ: 10.81 (s, 1H), 7.29-7.36 (m, 6H), 5.29 (s, 1H), 5.10 (s, 2H), 4.41 (q, 2H, J = 7.27 Hz), 4.13 (d, 2H, J = 3.66 Hz), 2.43-2.47 (m, 2H), 1.94-2.02 (m, 2H), 1.79-1.85 (m, 2H), 1.65-1.72 (m, 2H), 1.40 (t, 3H, J = 7.17). LCMS (M+H) = 428.26.
- 15 20 The reaction mixture was then cooled and concentrated *in vacuo*. The residue was taken up in ethyl acetate (200 mL) and extracted with 0.2 M aq. NaOH (3 x 50 mL). The combined aqueous phase was washed with ethyl acetate (250 mL). The aqueous layer was then acidified with conc. HCl and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ extracts were dried (Na₂SO₄), filtered and concentrated to afford the title compound as a brown solid (400 mg). ¹H NMR (500 MHz, CDCl₃) δ: 10.81 (s, 1H), 7.29-7.36 (m, 6H), 5.29 (s, 1H), 5.10 (s, 2H), 4.41 (q, 2H, J = 7.27 Hz), 4.13 (d, 2H, J = 3.66 Hz), 2.43-2.47 (m, 2H), 1.94-2.02 (m, 2H), 1.79-1.85 (m, 2H), 1.65-1.72 (m, 2H), 1.40 (t, 3H, J = 7.17). LCMS (M+H) = 428.26.

20

Intermediate 71




- 25 *Ethyl 10-amino-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate.* To a mixture of ethyl 10-((benzyloxy)carbonyl)amino-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-

ethanopyrimido[1,2-a]azepine-2-carboxylate (2 g, 4.68 mmol) in ethanol (75 mL) was added 1N HCl (5.15 mL, 5.15 mmol) followed by Pd/C (0.498 g, 0.468 mmol) and the mixture stirred under 1 atm of hydrogen for 18 h. The mixture was then filtered through a pad of CELITE® and the pad washed with dichloromethane. The 5 filtrate was then concentrated to afford the title compound (1.4 g, 91% yield) as an off-white solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 10.5 (1 H, brs), 4.3 (2 H, q, *J*=7.02 Hz), 4.0 (2 H, d, *J*=3.66 Hz), 2.4 (1 H, brs), 2.0 - 2.1 (4 H, m), 1.8 - 1.9 (2 H, m), 1.7 - 1.8 (2 H, m), 1.3 (3 H, t, *J*=7.02 Hz). LCMS (M+H) = 294.32.


10

Intermediate 72



15 *Ethyl 10-((dimethylamino)(oxo)acetyl)amino-3-hydroxy-4-oxo-4,6,7,8,9,10-*
hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate. To a stirred solution
of ethyl 10-amino-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-
ethanopyrimido[1,2-a]azepine-2-carboxylate (250 mg, 0.895 mmol) and 2-
20 (dimethylamino)-2-oxoacetic acid (157 mg, 1.343 mmol) in DMF (6 mL) was added
diisopropyl-ethylamine (0.498 mL, 2.86 mmol), 4-(dimethylamino)pyridine (DMAP)
25 (21.87 mg, 0.179 mmol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-
tetramethyluronium hexafluorophosphate (HATU) (511 mg, 1.343 mmol) and the
resulting mixture stirred at room temperature for 5 hr. Solvent was removed under
reduced pressure. The crude material was diluted with ethyl acetate (50 mL) and
washed with 1N HCl (20 mL), dried (Na₂SO₄) and concentrated to yield the title
compound.


Intermediate 73



5 *Ethyl 10-((benzyloxy)carbonyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-*
4,6,7,8,9,10-*hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate.* To a
solution of ethyl 10-((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-
hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (13.9 g, 32.5 mmol) in
pyridine (150 mL) was added benzoic anhydride (11.03 g, 48.8 mmol) and the
10 resulting mixture was stirred at room temp for 16 h. Solvent was removed under
reduced pressure and the residue was diluted with ethyl acetate (500 mL), washed
with 1N HCl (100 mL), sat. NaHCO₃ followed by brine, then dried (Na₂SO₄), filtered
and concentrated. The crude material was then triturated with ethyl acetate / hexane
and dried to afford the title compound (16 g, 30.1 mmol, 93 % yield) as a light
15 yellow solid. ¹H NMR (500 MHz, CDCl₃) δ: 8.16 (2 H, dd, *J*=8.28, 1.25 Hz), 7.59 -
7.64 (1 H, m), 7.48 (2 H, t, *J*=7.78 Hz), 7.26 - 7.38 (5 H, m), 5.27 (1 H, s), 5.08 (2 H,
s), 4.24 (2 H, q, *J*=7.03 Hz), 4.13 (2 H, d, *J*=3.76 Hz), 2.84 - 2.99 (2 H, m), 2.45 (1
H, brs), 1.85 - 2.05 (4 H, m), 1.67 - 1.77 (2 H, m), 1.11 (3 H, t, *J*=7.15 Hz). LCMS
(M+H) = 532.24.


20

Intermediate 74



5 *Ethyl 10-amino-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride.* To a mixture of ethyl 10-((benzyloxy)carbonyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (16 g, 30.1 mmol) in ethyl acetate (300 mL) and methanol (150 mL) was added 1N HCl (31.6 mL, 31.6 mmol) followed by Pd/C (3.20 g, 3.01 mmol) and the mixture was stirred under balloon hydrogen atmosphere for 2 h. The mixture was then filtered through a pad of CELITE® and the pad was thoroughly washed with ethyl acetate. The filtrate was concentrated under reduced pressure and dried under high vacuum overnight to afford the title compound (12.75 g, 98% yield), as an off-white solid. ¹H NMR (500 MHz, CDCl₃) δ: 8.11 - 8.16 (2 H, m), 7.62 (1 H, t, *J*=7.40 Hz), 7.48 (2 H, t, *J*=7.78 Hz), 4.19 (2 H, q, *J*=7.11 Hz), 4.13 (2 H, d, *J*=3.51 Hz), 2.71 - 2.83 (2 H, m), 2.53 (1 H, brs), 2.32 - 2.46 (2 H, m), 2.01 - 2.13 (2 H, m), 1.74 - 1.85 (2 H, m), 1.03 (3 H, t, *J*=7.03 Hz). LCMS (M+H) = 398.10.


Intermediate 75



5 *Ethyl 10-((dimethylamino)(oxo)acetyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate.* To a stirred solution of ethyl 10-amino-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride (2.4 g, 5.53 mmol) in DMF (50 mL) as added 2-(dimethylamino)-2-oxoacetic acid (1.295 g, 11.06 mmol), diisopropyl-ethylamine (5.80 mL, 33.2 mmol) and the resulting mixture was stirred at room temperature for 3 h. Water was then added and the mixture extracted with ethyl acetate (2 x 200 mL). The organic solution was washed with 1N HCl (50 mL), dried (Na_2SO_4), filtered and concentrated. The crude material was triturated with ethyl acetate / hexane to afford 15 the title compound (2.5 g, 5.04 mmol, 91% yield) as a light yellow solid. ^1H NMR (500 MHz, CDCl_3) δ : 9.33 (1 H, brs), 8.18 (2 H, d, $J=7.02$ Hz), 7.64 (1 H, t, $J=7.48$ Hz), 7.50 (2 H, t, $J=7.93$ Hz), 4.29 (2 H, q, $J=7.02$ Hz), 4.17 (2 H, d, $J=3.66$ Hz), 3.34 (3 H, s), 3.03 (3 H, s), 2.88 - 3.00 (2 H, m), 2.50 (1 H, brs), 1.93 - 2.10(4 H, m), 1.73 - 1.82 (2 H, m), 1.18 (3 H, t, $J=7.17$ Hz). LCMS ($\text{M}+\text{H}$) = 497.79.


20

Intermediate 76



5 *Ethyl 10-(benzyl(methyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-*
hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate. To a stirred solution
of ethyl 10-amino-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-
ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride (12.75 g, 29.4 mmol) in
1,2-dichloroethane (500 mL) was added triethylamine (4.10 mL, 29.4 mmol) and the
10 resulting mixture was stirred for 5 min. Benzaldehyde (5.94 mL, 58.8 mmol), AcOH
(1.682 mL, 29.4 mmol) and NaCNBH₃ (5.54 g, 88 mmol) were then added and the
mixture stirred at room temp for 6h. Formaldehyde (4.38 mL, 58.8 mmol) was then
added to the reaction and the mixture was stirred at room temperature for 16h. The
reaction was quenched with sat. NaHCO₃ and extracted with dichloromethane, dried
15 (Na₂SO₄), filtered and concentrated. The crude product was the recrystallized from
ethyl acetate to afford the title compound (11.6 g, 23.13 mmol, 79% yield) as an off-
white solid. ¹H NMR (500 MHz, CDCl₃) δ: 8.20 (2 H, d, *J*=7.32 Hz), 7.63 (1 H, t,
J=7.32 Hz), 7.50 (2 H, t, *J*=7.78 Hz), 7.39 (2 H, d, *J*=7.63 Hz), 7.31 (2 H, t, *J*=7.48
Hz), 7.20 - 7.23 (1 H, m), 4.28 - 4.34 (2 H, m), 4.09 (4 H, brs), 2.53 (3 H, s), 2.47 (1
20 H, brs), 2.32 - 2.42 (2 H, m), 2.02 - 2.10 (2 H, m), 1.89 - 1.99 (2 H, m), 1.66 - 1.74 (2
H, m), 1.22 (3 H, t, *J*=7.17 Hz). LCMS (M+H) = 502.37.


Intermediate 77



5 *Ethyl 10-(methylamino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-*
hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride. To a
mixture of ethyl 10-amino-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-
7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride (6 g, 11.96 mmol) in
ethyl acetate (100 mL) and ethanol (100 mL) was added 1N HCl (12.56 mL, 12.56
10 mmol) followed by Pd/C (1.273 g, 1.196 mmol) and the mixture stirred under 1 atm
of hydrogen for 2 h. The mixture was then filtered (filter paper) and the solid
thoroughly washed with ethyl acetate. The filtrate was concentrated under reduced
pressure and dried under high vacuum overnight to afford the title compound (4.8 g,
90% yield) as an off-white solid. ¹H NMR (500 MHz, CDCl₃) δ: 8.15 (2 H, d, *J*=7.28
15 Hz), 7.62 - 7.67 (1 H, m), 7.47 - 7.52 (2 H, m), 4.26 (2 H, q, *J*=7.03 Hz), 4.11 (2 H, d,
J=3.76 Hz), 2.76 (3 H, s), 2.56 (1 H, brs), 2.16 - 2.25 (2 H, m), 2.00 - 2.13 (4 H, m),
1.68 - 1.82 (3 H, m), 1.12 (3 H, t, *J*=7.03 Hz). LCMS (M+H)⁺ = 412.78.

Intermediate 78


20




Methyl 10-(((dimethylamino)(oxo)acetyl)(methyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate. To a solution of ethyl 10-(methylamino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride salt (4.7 g, 10.49 mmol) in dichloromethane (100 mL) at 0 °C was added diisopropyl-ethylamine (5.50 mL, 31.5 mmol) followed by methyl 2-chloro-2-oxoacetate (0.967 mL, 10.49 mmol) and the resulting mixture stirred at room temp for 2h. The reaction mixture was then washed with 1N HCl (50 mL) followed by brine (100 mL), then dried (Na_2SO_4), filtered, concentrated and dried under high vacuum for 2hr. The crude material was treated with 2M Me_2NH in methanol (52.5 mL, 105 mmol) and stirred at 50°C for 16h. The mixture was allowed to cool to room temperature and concentrated, diluted with ethyl acetate, washed with 0.2N NaOH (2 x 75 mL). The NaOH layer was acidified with conc. HCl and extracted with dichloromethane, then dried (Na_2SO_4), filtered and concentrated to provide the title compound (3.5 g, 85% yield). ^1H NMR (500 MHz, CDCl_3) δ : 10.09 (1 H, s), 3.93 (3 H, s), 3.16 (3 H, s), 3.11 (3 H, s), 3.02 (3 H, s), 2.52 (1 H, brs), 2.05 - 2.15 (2 H, m), 1.81 - 2.00 (2 H, m), 1.50 - 1.80 (6 H, m). LCMS ($\text{M}+\text{H}$) = 393.15.

Intermediate 79

20




*Ethyl 10-(((3*R*)-3-fluoro-1-pyrrolidinyl)(oxo)acetyl)(methyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate.* To a stirred solution of ethyl 10-(methylamino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride (120 mg, 0.268 mmol) in DMF (4 mL) was added (R)-2-(3-fluoropyrrolidin-1-yl)-2-oxoacetic acid (130 mg, 0.804 mmol), diisopropyl-ethylamine (0.281 mL, 1.607 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (306 mg, 0.804 mmol), 4-(dimethylamino)pyridine (DMAP) (9.82 mg, 0.080 mmol) and the resulting mixture stirred at room temperature for 16 h. The mixture was quenched with water and diluted with ethyl acetate, washed with 1N HCl, followed by sat.NaHCO₃, dried (Na₂SO₄) filtered and concentrated. The crude product was purified on Biotage flash chromatography system using 30-100% ethyl acetate/ hexane gradient to afford the title compound (80 mg, 0.144 mmol, 54% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ: 8.12 - 8.23 (2 H, m), 7.64 (1 H, t, *J*=7.48 Hz), 7.44 - 7.56 (2 H, m), 5.11 - 5.42 (1 H, m), 4.08 - 4.27 (2 H, m), 3.28 - 4.01 (9 H, m), 3.18 (3 H, d, *J*=3.05 Hz), 2.50 (1 H, brs), 1.72 - 2.37 (6 H, m), 0.99 - 1.11 (3 H, m). LCMS (M+H)⁺ = 555.76.



Ethyl 10-(benzyl(ethyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate. To a stirred solution of Ethyl 10-amino-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate hydrochloride (200 mg, 0.461 mmol) in 5 1,2-dichloroethane (10 mL) was added triethylamine (0.064 mL, 0.461 mmol) and the resulting mixture stirred for 5 min. Benzaldehyde (0.186 mL, 1.844 mmol), AcOH (0.053 mL, 0.922 mmol) and NaCNBH₃ (87 mg, 1.383 mmol) were then added and the mixture stirred for 6 h. Acetaldehyde (0.182 mL, 3.23 mmol) was then added to the reaction and the this was stirred at room temperature for 16 h. The 10 mixture was then quenched with sat. NaHCO₃ and extracted with dichloromethane, dried (Na₂SO₄), filtered and concentrated to afford title compound.


Intermediate 81

15



Ethyl 10-(ethylamino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate. To a mixture of ethyl 10-(benzyl(ethyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-20 7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (120 mg, 0.233 mmol) in methanol (5 mL) was added 1N HCl (0.256 mL, 0.256 mmol) followed by Pd/C (24.77 mg, 0.023 mmol) and the mixture stirred under 1atm hydrogen for 18 h. The mixture was filtered and the solids thoroughly washed with dichloromethane. The filtrate was concentrated under reduced pressure and dried under high vacuum overnight to 25 afford the title compound.


Intermediate 82



5 *Ethyl 8-methyl-1,4-dioxaspiro[4.5]decane-8-carboxylate.* Lithium diethylamide was prepared by adding (5 min) nBuLi in hexanes (94 mL, 150 mmol) to a solution of diisopropylamine (22.80 mL, 160 mmol) in THF (50 mL) at -30°C (dry ice/acetone) under N₂. After 20 min, ethyl 1,4-dioxaspiro[4.5]decane-8-carboxylate (21.43 g, 100 mmol) in THF (25 mL) was added over 5 min. After 20 10 min, iodomethane (28.4 g, 200 mmol) in THF (25 mL) was added over 5 min. The resulting solution was stirred at -30°C for 20 min and then allowed to slowly warm to room temperature overnight under N₂. The reaction was quenched with water (200 mL). The solution was concentrated to remove THF. The aqueous solution was extracted with CH₂Cl₂ (3 x 100 mL) and the organic layer dried (MgSO₄), filtered, 15 and concentrated to yield ethyl 8-methyl-1,4-dioxaspiro[4.5]decane-8-carboxylate (22 g, 96 mmol, 96 % yield) as a yellow oil. LCMS: m/z 229 (M+H). ¹H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 4.13 (2 H, q, *J*=7.22 Hz), 3.92 (4 H, s), 2.06 - 2.16 (2 H, m), 1.57 - 1.66 (4 H, m), 1.51 (2 H, dd, *J*=12.97, 4.12 Hz), 1.24 (3 H, t, *J*=7.17 Hz), 1.17 (3 H, s).

20

Intermediate 83




25 *(8-Methyl-1,4-dioxaspiro[4.5]decan-8-yl)methanol.* Yield: 100 % (clear oil). ¹H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 3.87 - 3.97 (4 H, m), 3.38 (2 H, s), 1.66 (1 H, d, *J*=3.66 Hz), 1.59 - 1.65 (3 H, m), 1.53 - 1.59 (2 H, m), 1.52 (1 H, d,

J=4.58 Hz), 1.45 (1 H, brs), 1.40 (2 H, t, *J*=5.80 Hz), 1.37 (1 H, s), 0.96 (3 H, s). LCMS: m/z 187 (M+H).

Intermediate 84

5



1-Amino-4-(hydroxymethyl)-4-methylcyclohexanecarbonitrile. Yield: 73.3 % (crude oil). LCMS: m/z 169.21 (M+H).

10


Intermediate 85



15 *Benzyl 1-cyano-4-(hydroxymethyl)-4-methylcyclohexylcarbamate.* Yield: 21 % (yellow oil; column-purified). LCMS: m/z 303.98 (M+H). HPLC: retention time = 1.90 min (area percent =:100%). • ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 7.26 - 7.59 (5 H, m) 5.13 (2 H, brs) 4.99 (1 H, d, *J*=3.05 Hz) 3.32 - 3.47 (2 H, m) 2.11 - 2.50 (2 H, m) 1.85 - 2.11 (1 H, m) 1.28 - 1.85 (8 H, m) 0.91 - 1.06 (3 H, m).


20

Intermediate 86




5 *Ethyl 2-(1-(benzyloxycarbonylamino)-4-(hydroxymethyl)cyclohexyl)-5-*
hydroxy-6-oxo-1,6-dihdropyrimidine-4-carboxylate. Yield: 42 % (crude glassy
solid). HPLC: retention time = 1.78 min (area percent = 100%). LCMS: m/z
460.21(M+H). ¹H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 0.97 (s, 3 H) 1.27 -
1.40 (m, 4 H) 1.42 (t, *J*=6.56 Hz, 3 H) 1.67 (d, *J*=9.16 Hz, 2 H) 1.97 - 2.03 (m, 2 H)
10 2.04 (s, 1 H) 2.06 (d, *J*=1.22 Hz, 1 H) 2.20 (br. s., 4 H) 3.35 (s, 1 H) 3.48 (s, 1 H)
4.41 - 4.47 (m, 2 H) 5.07 (br. s., 2 H) 7.33 (br. s., 6 H) 10.78 (br. s., 1 H).

Intermediate 87



15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxylic acid, 4,6,7,8,9,10-*
hexahydro-3-hydroxy-7-methyl-4-oxo-10-[(phenylmethoxy)carbonyl]amino]-, ethyl
ester. Yield: 18 % (crude oil), Used as is for next reaction. HPLC: retention time =
20 2.43 min (area percent =:15%). LCMS: m/z 441.94 (M+H).


Intermediate 88



5 *((1R,3S)-1,2,2-Trimethylcyclopentane-1,3-diyl)dimethanol.* A solution of (1R,3S)-1,2,2-trimethylcyclopentane-1,3-dicarboxylic acid (26.25 g, 131 mmol) in THF (400 mL) was cooled in an ice bath (10 °C) and to this was added 1M LiAlH₄/THF (328 mL, 328 mmol), dropwise over 1 h. The ice bath was removed and the white slurry was stirred for 5 h at room temperature and 17 h at reflux. After 10 cooling to room temperature, the mixture was diluted with Et₂O (500 mL), cooled in an ice bath and carefully quenched with water (12.2 mL), 15% NaOH (12.2 mL) and water (36.7 mL). After stirring for 30 min at room temperature, the white slurry was filtered through a pad of CELITE®, then dried (Na₂SO₄), filtered and concentrated to give ((1R,3S)-1,2,2-trimethylcyclopentane-1,3-diyl)dimethanol (21.665 g, 126 mmol, 15 96% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ: 3.73 (1 H, dd, *J* = 10.2, 5.3 Hz), 3.58 (1 H, d, *J* = 10.7 Hz), 3.51 (1 H, dd, *J* = 10.1, 8.5 Hz), 3.46 (1 H, d, *J* = 10.7 Hz), 2.04 - 2.12 (1 H, m), 1.89 - 1.99 (1 H, m), 1.55 - 1.64 (1 H, m), 1.31 - 1.41 (2 H, m), 1.22 (2 H, brs), 1.02 (3 H, s), 1.01 (3 H, s), 0.78 (3 H, s).

20

Intermediate 89




25 *((1R,3S)-3-((tert-Butyldiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentane-1,3-diyl)methanol.* To a stirred solution of ((1R,3S)-1,2,2-trimethylcyclopentane-1,3-diyl)dimethanol (18.19 g, 106 mmol) and imidazole (8.63 g, 127 mmol) in DMF (400 mL) at 0 °C (ice bath) was added tert-butylchlorodiphenylsilane (28.8 mL, 111 mmol) over 5 min. The reaction mixture

was stirred overnight (16 h) while slowly warming to room temperature. The reaction mixture was then diluted with Et₂O (1000 mL), washed with water (5 x 100 mL) followed by brine (100 mL), then dried (Na₂SO₄), filtered and concentrated to give a color less liquid. Flash column chromatography on silica gel column using 9:1 and 4:1 hexane/ethyl acetate as eluent provided ((1R,3S)-3-((tert-butylidiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentyl)methanol (40.5 g, 99 mmol, 93% yield) as a colorless viscous oil. ¹H NMR (500 MHz, CDCl₃) δ: 7.64 - 7.69 (4 H, m), 7.35 - 7.44 (6 H, m), 3.70 (1 H, dd, *J* = 10.1, 6.4 Hz), 3.54 (2 H, dt, *J* = 10.4, 6.7 Hz), 3.44 (1 H, dd, *J* = 10.7, 5.5 Hz), 2.09 - 2.18 (1 H, m), 1.82 - 1.92 (1 H, m), 1.49 - 1.58 (1 H, m), 1.24 - 1.36 (3 H, m), 1.04 (9 H, s), 0.98 (3 H, s), 0.97 (3 H, s), 0.75 (3 H, s). LCMS (M+H) calcd for C₂₆H₃₉O₂Si: 411.27; found: 411.31.

Intermediate 90

15




(1*R*,3*S*)-3-((tert-Butyldiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentanecarbaldehyde. A solution of 4-methylmorpholine (17.33 g, 148 mmol) in CH₂Cl₂ (400 mL) was dried over Mg₂SO₄ for 15 min and filtered into a round bottom flask containing ((1*R*,3*S*)-3-((tert-butylidiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentyl)methanol (40.5 g, 99 mmol). To the resulting clear solution was added 4A° powdered molecular sieves (20 g), tetrapropylammonium perruthenate (TPAP) (0.693 g, 1.972 mmol) and stirred for 4 h at room temperature. The reaction was concentrated to ~100 mL, diluted with hexane/Et₂O (9:1, 500 mL) and filtered through a plug of silica gel. The plug was washed with 500 mL of hexane/Et₂O (9:1). The filtrate was concentrated to afford (1*R*,3*S*)-3-((tert-butylidiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentanecarbaldehyde (32.61 g, 80 mmol, 81% yield) as a viscous pale yellow oil. ¹H NMR (500 MHz, CDCl₃) δ: 9.63 (1 H, s), 7.66 (4 H, dd, *J* = 7.9, 1.5 Hz), 7.35 - 7.45 (6 H, m), 3.66 - 3.71 (1 H, dd, *J* =

10.2, 6.9 Hz), 3.56 (1 H, dd, J = 10.4, 7.0 Hz), 2.29 (1 H, ddd, J = 13.7, 11.4, 6.0 Hz), 2.09 - 2.18 (1 H, m), 1.87 - 1.96 (1 H, m), 1.28 - 1.44 (2 H, m), 1.06 (3 H, s), 1.02 - 1.05 (12 H, s), 0.81 (3 H, s). LCMS (M+Na) calcd for $C_{26}H_{36}NaO_2Si$: 431.25; found: 431.52.

5

Intermediate 91




10 *Benzyl ((1R,3S)-3-((tert-butyldiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentyl)(cyano) ethylcarbamate.* To a stirred mixture of (1R,3S)-3-((tert-butyldiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentanecarbaldehyde (8.2 g, 20.07 mmol) and zinc(II) iodide (0.641 g, 2.007 mmol) was added trimethylsilyl cyanide (2.94 mL, 22.07 mmol) at room temperature. Note: The reaction was warm to touch within 5 min after the addition of TMSCN. After 15 min, 7N ammonia/methanol (40 mL, 1848 mmol) was added and stirring continued overnight (17 h) at room temperature. The reaction mixture was concentrated and the resulting residue taken up in toluene and concentrated to remove traces of methanol/NH₃. The resulting white slurry was dissolved in CH₂Cl₂ (50 mL) and 1N aq. Na₂CO₃ (25 mL), and treated with benzylchloroformate (3.15 mL, 22.07 mmol) at room temperature. After 3 h, the organic layer was separated and the aqueous layer extracted with CH₂Cl₂ (25 mL). The combined CH₂Cl₂ layers were dried (Na₂SO₄), filtered and concentrated to give a color less oil. Flash column chromatography on silica gel column using 10, 15 and 20% ethyl acetate/hexane gave benzyl ((1R,3S)-3-((tert-butyldiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentyl)(cyano) ethylcarbamate (9.4 g, 16.53 mmol, 82% yield) as a white foam. ¹H NMR (500 MHz, CDCl₃) δ: 7.61 - 7.67 (4 H, m), 7.31 - 7.46 (11 H, m), 5.09 - 5.19 (2 H, m), 4.97 (1 H, d, J = 10.1 Hz), 4.78 (1 H, d, J = 10.1 Hz), 3.62 - 3.69 (1 H, m), 3.52 (1 H, dd, J = 10.1, 7.0

Hz), 2.13 (1 H, tt, $J = 9.7, 6.7$ Hz), 1.83 - 1.93 (1 H, m), 1.65 - 1.74 (1 H, m), 1.58 - 1.64 (1 H, m), 1.28 - 1.38 (1 H, m), 1.14 (3 H, s), 1.01 - 1.05 (9 H, m), 0.95 (3 H, s), 0.81 (3 H, s). LCMS (M+Na) calcd for $C_{35}H_{44}NaN_2O_3Si$: 591.30 ; found: 591.63.

5


Intermediate 92



Benzyl cyano((1R,3S)-3-(hydroxymethyl)-1,2,2-


10 *trimethylcyclopentyl)methylcarbamate.* To a stirred colorless solution of benzyl ((1R,3S)-3-((tert-butyldiphenylsilyloxy)methyl)-1,2,2-trimethylcyclopentyl)(cyano)methylcarbamate (9.4 g, 16.53 mmol) in THF (50 mL) was added 1N tetrabutyl ammonium fluoride/THF (19.83 mL, 19.83 mmol) at room temperature. The resulting yellow mixture was stirred for 24 h and then concentrated. The residue was taken up in ethyl acetate (150 mL), washed with 1N aq. HCl (30 mL), water (50 mL) followed by brine (2 x 25 mL), then dried (Na_2SO_4), filtered and concentrated to give a colorless oil. Flash column chromatography on silica gel column using 3:7 and 3:2 ethyl acetate/hexane (product came out with 3:2 ethyl acetate/hexane) provided benzyl cyano((1R,3S)-3-(hydroxymethyl)-1,2,2-trimethylcyclopentyl)methylcarbamate (5.22 g, 15.80 mmol, 96% yield) as a white foam. 1H NMR (500 MHz, $CDCl_3$) δ : 7.31 - 7.41 (5 H, m), 5.09 - 5.20 (2 H, m), 5.01 (1 H, d, $J = 10.1$ Hz), 4.82 (1 H, d, $J = 10.1$ Hz), 3.66 - 3.75 (1 H, m), 3.48 - 3.56 (1 H, m), 2.06 - 2.15 (1 H, m), 1.93 - 2.04 (1 H, m), 1.72 - 1.81 (1 H, m), 1.64 - 1.71 (1 H, m), 1.37 - 1.48 (1 H, m), 1.18 - 1.22 (1 H, m), 1.17 (3 H, s), 1.00 (3 H, s), 0.84 (3 H, s). LCMS (M+H) calcd for $C_{19}H_{27}N_2O_3$: 331.20; found: 331.14.

Intermediate 93



5 *Diethyl 2-(1-amino-2-(benzyloxycarbonylamino)-2-((1R,3S)-3-(hydroxymethyl)-1,2,2-trimethylcyclopentyl)ethylideneaminoxy)but-2-enedioate.* A clear homogeneous solution benzyl cyano((1R,3S)-3-(hydroxymethyl)-1,2,2-trimethylcyclopentyl)methylcarbamate (5.22 g, 15.80 mmol) and 50% aq. hydroxylamine (1.484 mL, 24.22 mmol) in EtOH (50 mL) was stirred for 40 h at 50 °C. It was then concentrated and the resulting residue redissolved in EtOH (50 mL) and treated with diethyl acetylenedicarboxylate (3.03 mL, 18.96 mmol). After 4 h at 40 °C, the reaction mixture was concentrated and the resulting yellow residue purified by flash column chromatography on silica gel column using 20, 40, 60 and 75% ethyl acetate/hexane (product came out with 75% ethyl acetate/hexane) to provide diethyl 2-(1-amino-2-(benzyloxycarbonylamino)-2-((1R,3S)-3-(hydroxymethyl)-1,2,2-trimethylcyclopentyl)ethylideneaminoxy)but-2-enedioate (6.051 g, 11.34 mmol, 72% yield) as a white foam. ¹H NMR (500 MHz, CDCl₃) δ: 7.28 - 7.39 (5 H, m), 5.75 (0.5 H, s), 5.22 (0.5 H, brs), 5.06 - 5.16 (3 H, m), 4.91 (1 H, brs), 4.23 - 4.41 (3 H, m), 4.13 - 4.19 (2 H, m), 3.68 - 3.75 (1 H, m), 3.46 - 3.55 (1 H, m), 2.01 - 2.11 (1 H, m), 1.86 - 1.99 (1 H, m), 1.67 - 1.80 (1 H, m), 1.23 - 1.38 (7 H, m), 1.15 - 1.19 (1 H, m), 1.11 (2 H, s), 1.07 (1 H, s), 1.03 (2 H, s), 0.99 (1 H, s), 0.85 (3 H, s). LCMS (M+H) calcd for C₂₇H₄₀N₃O₈: 534.28; found: 534.26.

Intermediate 94




5 *Ethyl (7S,10R)-11-((benzyloxy)carbonyl)amino)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocine-2-carboxylate.* A solution of diethyl 2-(1-amino-2-(benzyloxycarbonylamino)-2-((1R,3S)-3-(hydroxymethyl)-1,2,2-trimethylcyclopentyl)ethylideneaminoxy)but-2-enedioate (6.05 g, 11.34 mmol) in xylene (250 mL) was heated at 125 - 130 °C for 8 h. The dark reaction mixture was concentrated and the resulting residue was used in the next step without purification. The above crude product was dissolved in DMF (40 mL) and stirred with K₂CO₃ (3 g, 21.71 mmol) and methanesulfonyl chloride (0.779 mL, 10 mmol) for 2 h at room temperature and 28 h at 90 °C. After cooling to room temperature it was diluted with ethyl acetate (200 mL), neutralized with 1N aq. HCl. The organic layer was separated and washed with water (2 x 50 mL) followed by brine (50 mL), then dried (Na₂SO₄), filtered and concentrated to give a brown paste. Purification by preparative HPLC on a C18 column using methanol/water (0.1% trifluoroacetic acid, gradient elution starting with 30% methanol/water) provided the title compound (1.4595 g, 3.11 mmol, 27% yield) as a brown solid.

10

15

20

Intermediate 95



25 *Bicycle[4.1.0]heptan-2-one.* The title compound was prepared by following the procedure described in O.O. Grygorenko et al., *Tetrahedron: Asymmetry*, **2006**,

17, 252. In a dry round bottom flask, 60% (in mineral oil) NaH (4.49 g, 112 mmol) was triturated with hexanes three times, and placed under high vacuum to further dry. To DMSO (100 mL) in an oven-dried three neck flask under nitrogen was added the concentrated NaH in small portions. The temperature was kept between 15-25°C.


5 Once gas evolution had subsided, trimethylsulfoxonium iodide (24.73 g, 112 mmol) was added via a powder addition funnel in small portions keeping the temperature between 15-25°C. The suspension was allowed to stir for 2 h during which time it became a milky solution. Cyclohex-2-enone (10 g, 104 mmol) in DMSO (25 mL) was then slowly added drop-wise. The solution became red and was stirred for 0.5 h

10 at room temperature and 2 h at 50°C. It was then poured into 100 mL of ice water. The water was extracted with Et₂O (4 x 100 mL). The ether extracts were combined, dried (MgSO₄) and concentrated under reduced pressure. The resulting red oil was purified by distillation. Fractions distilling ate 65 and 70°C were collected. Yield: 72% (liquid). HPLC: retention time = 1.05 min (area percent =:70%). ¹H NMR (500

15 MHz, CHLOROFORM-*d*) δ ppm 1.07 (ddd, *J*=9.84, 8.16, 5.19 Hz, 1 H) 1.19 (q, *J*=5.39 Hz, 1 H) 1.54 - 1.63 (m, 1 H) 1.63 - 1.77 (m, 3 H) 1.84 - 1.92 (m, 1 H) 1.92 - 1.99 (m, 1 H) 2.03 (ddd, *J*=18.31, 11.60, 6.71 Hz, 1 H) 2.27 (dt, *J*=18.31, 4.58 Hz, 1 H).

20

Intermediate 96




25 *3-(Chloromethyl)cyclohexanone*. The title compound was prepared following the procedure described in (N. Di Bello et al., *Synthesis*, **1978**, 227. To a solution of bicyclic[4.1.0]heptan-2-one (13.8 g, 125 mmol) in acetonitrile (300 mL) was added pyridine hydrochloride (43.3 g, 375 mmol) which had been triturated with Et₂O and dried under high vacuum. The solution was stirred under N₂ at 82°C, overnight. The reaction mixture was partially concentrated, poured into brine (200 mL) and extracted with Et₂O (4 x 100ML). The ether extracts were combined, washed with brine (100mL), dried and concentrated. The residual oil was purified on a Biotage flash

30

chromatography system (15% ethyl acetate in hexanes) to yield the title compound (13.4 g, 91 mmol, 73.1 % yield) as an oil. LCMS: 147.09 (M+H), 149.10 (M+3, 30% rel. intensity). ¹H NMR (500 MHz, CHLOROFORM-d) δ ppm 1.51 - 1.60 (m, 1 H) 1.62 - 1.74 (m, 1 H) 1.95 - 2.02 (m, 1 H) 2.06 - 2.13 (m, J=13.47, 9.80, 3.62, 3.62 Hz, 5 1 H) 2.13 - 2.20 (m, 1 H) 2.20 - 2.31 (m, 2 H) 2.35 - 2.43 (m, 1 H) 2.45 - 2.51 (m, 1 H) 3.45 - 3.55 (m, 2 H) . ¹³C NMR (126 MHz, CHLOROFORM-d) δ ppm 24.63, 28.75, 40.85, 41.15, 45.39, 210.25.

Intermediate 97

10




Benzyl 3-(chloromethyl)-1-cyanocyclohexylcarbamate. Yield: 46% (oil).

HPLC: retention time = 2.77 min. LCMS: m/z 307 (M+H).


15

Intermediate 98



20 *Ethyl 2-(1-(benzyloxycarbonylamino)-3-(chloromethyl)cyclohexyl)-5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate.* LCMS: m/z 464 (M+H) observed.


Intermediate 99



5 *7,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxylic acid, 6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-11-[(phenylmethoxy)carbonyl]amino-, ethyl ester.*
 LCMS: m/z 428 (M+H) observed.

Intermediate 100


10



15 *7,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxylic acid, 6,7,8,9,10,11-hexahydro-4-oxo-3-(phenylmethoxy)-11-[(phenylmethoxy)carbonyl]amino-, ethyl ester.* Yield: 1% HPLC: retention time = 3.03 min (area percent =: 75%). LCMS: m/z 518.16 (M+H). 1HNMR: In agreement with structure. Impurities present.

Intermediate 101


20



Ethyl 1-amino-5-hydroxy-6-oxo-3,7-diazatricyclo[7.3.1.0^{2,7}]trideca-2,4-diene-4-carboxylate. LCMS: m/z 294.46 (M+H).

Intermediate 102


5



Methyl 3,3-dimethoxycyclopentanecarboxylate. (Tetrahedron, (1977), 33, 1113) A solution of 3-oxocyclopentanecarboxylic acid (10.013 g, 78 mmol) and trimethyl orthoformate (60.5 mL, 547 mmol) in MeOH (120 mL) was treated with acetyl chloride (7.78 mL, 109 mmol) and stirred under a nitrogen atmosphere for 5 days. The reaction solution was poured into a stirred suspension of anhydrous sodium carbonate (32 g) in pentane (250 mL) and stirred for 3 hrs. The solids were removed by filtration (2x) and concentrated under reduced pressure to give the target compound (15.4 g, 82 mmol, 105 % yield) as a pale yellow oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 3.67 (3 H, s), 3.20 (3 H, s), 3.18 (3 H, s), 2.82 - 2.93 (1 H, m), 2.01 - 2.15 (2 H, m), 1.77 - 2.01 (4 H, m).

Intermediate 103

20




(3,3-Dimethoxycyclopentyl)methanol. A solution of methyl 3,3-dimethoxycyclopentanecarboxylate, methyl 3,3-dimethoxycyclopentanecarboxylate (14.68 g, 78 mmol), in THF (100 mL) was cooled (0°C ice bath) and treated with 1.0N lithium aluminum hydride in THF (74.1 mL, 74.1 mmol) by dropwise addition. Upon completion of the addition, the reaction was warmed to room temperature and

stirred for 6 hrs. The reaction was quenched by slow addition of Na₂SO₄·10H₂O (17g). The suspension was stirred for 1 hr, treated with anhyd. Na₂SO₄, and filtered. The filtrate was concentrated to afford the target compound (12.01 g, 75.0 mmol, 96 % yield) as a colourless oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 3.54 (2 H, s), 3.20 (5 H, s), 3.19 (3 H, s), 2.21 - 2.32 (1 H, m), 1.99 (1 H, dd, *J*=13.28, 9.00 Hz), 1.85 - 1.91 (1 H, m), 1.78 - 1.85 (2 H, m), 1.73 - 1.78 (1 H, m), 1.56 (1 H, dd, *J*=13.12, 7.32 Hz), 1.40 - 1.48 (1 H, m).


Intermediate 104

10




3-(Hydroxymethyl)cyclopentanone. A solution of (3,3-dimethoxycyclopentyl)methanol, (3,3-dimethoxycyclopentyl)methanol (12.01 g, 75.0 mmol), in Et₂O (80 mL), was treated with p-toluene sulfonic acid monohydrate (0.4 g, 2.103 mmol) and stirred for 2 hrs. The reaction was poured into water (50 mL) and the layers were separated. The aqueous layer was saturated with sodium chloride then extracted with CH₂Cl₂ (5 x 50 mL) and EtOAc (4 x 50 mL). The combined extracts were dried (MgSO₄), filtered and concentrated under reduced pressure to give the target compound (8.01 g, 70.2 mmol, 94 % yield), as a brownish-yellow oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.66 - 1.77 (m, 1 H) 2.01 (dd, *J*=18.01, 8.85 Hz, 1 H) 2.08 - 2.24 (m, 2 H) 2.27 - 2.40 (m, 2 H) 2.40 - 2.50 (m, 1 H) 3.63 - 3.68 (m, 2 H).

Intermediate 105



5 *7,10-Methanopyrimido[1,2-a]azepine-2-carboxylic acid, 4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[phenylmethoxy]carbonyl]amino]-, ethyl ester.* A solution of ethyl 2-(1-(benzyloxycarbonylamino)-3-((methylsulfonyloxy)methyl)cyclopentyl)-1-(methylsulfonyl)-5-(methylsulfonyloxy)-6-oxo-1,6-dihydropyrimidine-4-carboxylate (6.10 g, 9.16 mmol) in THF (50 mL) and
10 EtOH (50 mL) was treated with potassium carbonate (2.56 g, 18.52 mmol). The reaction was stirred for 20 hrs at room temp., then warmed (45°C oil bath) for 5 hrs. The solvent was removed by rotary evaporator. The residue was partitioned between ethyl acetate (100 mL) and water (100 mL), and after separation of the layers, the basic aqueous layer was further extracted with ethyl acetate (2 x 50 mL). The basic
15 aqueous layer was made acidic with 6.0 N HCl and extracted with ethyl acetate (2 x 75 mL). The combined organic extracts were dried ($MgSO_4$), filtered and concentrated under reduced pressure to give the target compound and unreacted bis-mesylate intermediate (1:1 mixture, 1.887 g) as an amber oil / glassy foam. LC/MS (M+H) = 414.24. This was used in the subsequent step without further purification
20 or characterization.


Intermediate 106



7,10-Methanopyrimido[1,2-a]azepine-2-carboxylic acid, 3-(benzoyloxy)-4,6,7,8,9,10-hexahydro-4-oxo-10-[[[(phenylmethoxy)carbonyl]amino]-, ethyl ester. A solution of 7,10-methanopyrimido[1,2-a]azepine-2-carboxylic acid, 4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[[(phenylmethoxy)carbonyl]amino]-, ethyl ester (1.887 g, 4.56 mmol) in pyridine (50 mL) was treated with benzoic anhydride (1.53 g, 6.76 mmol) and stirred at room temperature for 16 hrs. The solvent was removed under reduced pressure and the residue was diluted with ethyl acetate (75 mL), and washed with 1.0N HCl (75 mL), saturated NaHCO₃ solution (75 mL) followed by brine, then dried (MgSO₄) filtered and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography, affording a yellow oil after concentration. This was dried once from Et₂O, then the resulting solid was triturated with Et₂O. The target compound (0.466 g, 0.900 mmol, 19.7 % yield) was isolated as a white solid after filtration. ¹H NMR (500 MHz, CDCl₃) δ ppm 8.17 (2 H, d, *J*=7.02 Hz), 7.64 (1 H, t, *J*=7.48 Hz), 7.50 (2 H, t, *J*=7.78 Hz), 7.28 - 7.43 (5 H, m), 6.63 (1 H, br s), 5.12 (2 H, s), 4.26 (2 H, q, *J*=7.32 Hz), 3.84 - 4.01 (2 H, m), 2.93 - 3.05 (1 H, m), 2.76 - 2.85 (1 H, m), 2.54 - 2.68 (1 H, m), 2.23 - 2.37 (1 H, m), 1.98 - 2.10 (2 H, m), 1.65 - 1.77 (1 H, m), 1.11 (3 H, t, *J*=7.02 Hz). LC/MS (M+H)⁺ = 518.28. HPLC purity >98%.

20

Intermediate 107




Dimethyl 2-(3-oxocyclopentyl)malonate. Tetrahedron Letters 2005, 46, 6875-6878. To a solution of cyclopent-2-enone (99.82 g, 1216 mmol) and dimethyl malonate (560 mL, 4887 mmol) in dry toluene (1000 mL) was added 2,3,4,6,7,8-hexahydro-1H-pyrimido[1,2-a]pyrimidine (5.10 g, 36.6 mmol). The reaction was stirred at room temperature under nitrogen for 16 hrs, then concentrated under

reduced pressure to ~1/2 volume. The crude product was filtered through a short path of silica gel (3.5 x 4 cm), eluting with ethyl acetate. The filtrate was concentrated under vacuum pump rotary evaporation (70-75°C water bath) for 2 hrs, affording the target compound (247.7 g, 1098 mmol, 90 % yield). ¹H NMR (500 MHz, CDCl₃) δ ppm 3.76 (3 H, s), 3.74 (3 H, s), 3.37 (1 H, d, *J*=9.46 Hz), 2.80 - 2.91 (1 H, m), 2.49 (1 H, dd, *J*=18.31, 7.63 Hz), 2.29 - 2.37 (1 H, m), 2.15 - 2.28 (2 H, m), 2.00 (1 H, ddd, *J*=18.46, 10.99, 1.37 Hz), 1.58 - 1.71 (1 H, m).

Intermediate 108


10



Methyl 2-(3-oxocyclopentyl)acetate. A 1000 mL flask was charged with dimethyl 2-(3-oxocyclopentyl)malonate (329.1 g, 1536 mmol), and dodecanedioic acid (283 g, 1229 mmol). The mixture was heated (210°C, oil bath) under a nitrogen atmosphere for 20 hrs, then distilled (~1 torr, 180°C oil bath) to give the title compound (170.4 g, 1091 mmol, 71.0 % yield) as a pale yellow oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 3.68 (3 H, s), 2.55 - 2.66 (1 H, m), 2.40 - 2.51 (3 H, m), 2.27 - 2.35 (1 H, m), 2.13 - 2.26 (2 H, m), 1.87 (1 H, ddd, *J*=18.23, 10.30, 1.37 Hz), 1.53 - 1.63 (1 H, m). ¹³C NMR (126 MHz, CDCl₃) δ ppm 218.36, 172.58, 51.78, 44.70, 39.53, 38.40, 33.52, 29.31.

Intermediate 109

25




Methyl 2-(3,3-dimethoxycyclopentyl)acetate. Tetrahedron, 57 (2001), 5183.

A mixture of Montmorillonite K10 (188 g, 571 mmol) in trimethyl orthoformate (316 mL, 2856 mmol) was stirred for 2 hrs under nitrogen. To the mixture was slowly added methyl 2-(3-oxocyclopentyl)acetate (89.2 g, 571 mmol), as a suspension in pentane (500 mL). The mixture was stirred for 16 hrs, then filtered through a CELITE® pad rinsing with pentane. The filtrate was concentrated under reduced pressure to afford the title compound (104.9 g, 519 mmol, 91 % yield) as an amber oil. ¹H NMR (400 MHz, CDCl₃) δ ppm 3.67 (3 H, s) 3.20 (6 H, s) 2.35 - 2.46 (3 H, m) 2.11 (1 H, dd, *J*=13.18, 7.40 Hz) 1.86 - 1.96 (2 H, m) 1.74 - 1.83 (1 H, m) 1.44 (1 H, dd, *J*=13.05, 8.53 Hz) 1.34 (1 H, dd, *J*=12.55, 8.78 Hz). ¹³C NMR (101 MHz, CDCl₃) δ ppm 172.73, 110.91, 50.93, 48.80, 48.61, 40.27, 39.66, 33.51, 33.25, 29.37.

Intermediate 110


15



2-(3,3-Dimethoxycyclopentyl)ethanol. J. Org. Chem., 1986, 51 (21), 4000-4005. A 5000 mL large neck round bottom flask was charged with a solution of methyl 2-(3,3-dimethoxycyclopentyl)acetate (238 g, 1177 mmol) in THF (1100 mL).

20 To the solution was added LiBH₄, 2.0 M in THF (706 mL, 1412 mmol) over approximately 1 min, followed by dropwise addition of methanol (57.1 mL, 1412 mmol). Note: vigorous exotherm. The reaction was stirred for 16 hrs, monitoring by ¹H and ¹³C NMR. The reaction was quenched by the addition of sat'd ammonium chloride solution (600 mL), then diluted with Et₂O (~500 mL), and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (300 mL), and the combined organic extracts were concentrated. The resulting oil was diluted with CH₂Cl₂ (200 mL), dried (MgSO₄), filtered and concentrated to yield 2-(3,3-dimethoxycyclopentyl)ethanol (203.3 g, 1167 mmol, 99 % yield).


Intermediate 111



5 *3-(2-Hydroxyethyl)cyclopentanone.* A solution of 2-(3,3-dimethoxycyclopentyl)ethanol (203.3 g, 1167 mmol), in Et₂O (1000 mL) was treated with p-toluene sulfonic acid (6.66 g, 35.0 mmol) and stirred for 16 hrs. The solution was extracted with water (4 x 100 mL) and the combined extracts were concentrated under reduced pressure to give the title compound (142.33 g, 1110 mmol, 95 % yield), as a dark brown oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 3.64 - 3.76 (2 H, m), 2.41 (1 H, dd, *J*=17.85, 7.17 Hz), 2.24 - 2.35 (2 H, m), 2.13 - 2.23 (2 H, m), 1.82 (1 H, dd, *J*=18.01, 10.38 Hz), 1.64 - 1.77 (2 H, m), 1.49 - 1.60 (1 H, m). ¹³C NMR (126 MHz, CDCl₃) δ ppm 219.75, 61.30, 45.21, 38.54, 38.41, 34.01, 29.67.


15

Intermediate 112




20 *6-((tert-Butyldiphenylsilyloxy)methyl)dihydro-2H-pyran-3(4H)-one.* Prepared according to the general procedure described in Bioorganic Med. Chem. 2006, 14, 3953-3966.

Intermediate 113




5 *Cis*-3-Amino-6-((tert-butyldiphenylsilyloxy)methyl)tetrahydro-2*H*-pyran-3-carbonitrile and *trans*-3-amino-6-(*y*(tert-butyldiphenylsilyloxy)methyl)tetrahydro-2*H*-pyran-3-carbonitrile. To a stirred solution of NH₄OH (188 mL) and methanol (75mL) was added 6-((tert-butyldiphenylsilyloxy)methyl)dihydro-2*H*-pyran-3(4*H*)-one (35 g, 95 mmol), NH₄Cl (10.16 g, 190 mmol) followed by NaCN (9.31 g, 190 mmol). After 16 at room temperature, the reaction mixture was extracted with CH₂Cl₂ (250 mL x 2), washed with water (500 mL), then dried (Na₂SO₄), filtered and concentrated. The crude product was purified by silica gel column chromatography (10% to 50% ethyl acetate/ hexane) to afford a 1:1 mixture of C3-epimeric diastereomers, *cis* (14.9 g, 40% yield) and *trans* (15.1 g, 40% yield). *Trans*-isomer: ¹H NMR (500 MHz, CDCl₃) δ: 7.62 - 7.71 (4 H, m), 7.32 - 7.47 (6 H, m), 4.03 (1 H, dd, *J*=11.14, 2.90 Hz), 3.77 (1 H, dd, *J*=10.53, 5.34 Hz), 3.62 (1 H, dd, *J*=10.53, 5.65 Hz), 3.36 - 3.43 (1 H, m), 3.20 (1 H, d, *J*=11.29 Hz), 2.18 - 2.25 (1 H, m), 1.82 - 1.87 (1 H, m), 1.72 - 1.78 (1 H, m), 1.71 (2 H, s), 1.62-1.67 (1 H, m), 1.06 (9 H, s). LCMS (M+H) = 395.33. *Cis*-isomer: ¹H NMR (500 MHz, CDCl₃) δ: 7.61 - 7.69 (4 H, m), 7.33 - 7.48 (6 H, m), 3.86 (1 H, dd, *J*=11.44, 1.98 Hz), 3.72 (1 H, dd, *J*=10.68, 5.19 Hz), 3.68 (1 H, d, *J*=11.60 Hz), 3.61 (1 H, dd, *J*=10.68, 5.19 Hz), 3.44 - 3.51 (1 H, m), 2.05 - 2.08 (1 H, m), 1.73 (2 H, brs), 1.61 - 1.71 (3 H, m), 1.05 (9 H, s). LCMS (M+H) = 395.33.

Intermediate 114



5 *Benzyl cis-6-((tert-butyldiphenylsilyloxy)methyl)-3-cyanotetrahydro-2H-pyran-3-ylcarbamate.* Benzyl chloroformate (5.42 mL, 38 mmol) was added to a biphasic mixture of water (110 mL) and CH₂Cl₂ (55 mL) containing *cis*-3-amino-6-((tert-butyldiphenylsilyloxy)methyl)tetrahydro-2H-pyran-3-carbonitrile (15 g, 38.0 mmol) and Na₂CO₃•H₂O (4.71 g, 38.0 mmol) at 0°C and the mixture was stirred for
10 2 h. The mixture was then allowed to warm to room temperature and stirred for 18 h. The mixture was then diluted with CH₂Cl₂ (200 mL), washed with sat. NaHCO₃ (100 mL), water (100 mL) followed by brine (100 mL), then dried (Na₂SO₄), filtered and concentrated to give a brown oil. Flash column chromatography on silica gel, using a 5% to 40% ethyl acetate-hexane gradient, afforded the title compound as a light
15 yellow liquid (18.4 g, 92% yield). ¹H NMR (500 MHz, CDCl₃) δ: 7.60 - 7.72 (4 H, m), 7.30 - 7.48 (12 H, m), 5.14 (2 H, s), 4.93 (1 H, s), 3.77 (1 H, dd, *J*=10.53, 5.34 Hz), 3.62 (1 H, dd, *J*=10.53, 5.65 Hz), 3.37 - 3.46 (1 H, m), 3.21 (1 H, d, *J*=10.99 Hz), 2.45 (1 H, dd, *J*=12.51, 2.75 Hz), 1.79 - 1.92 (2 H, m), 1.63 - 1.74 (1 H, m), 1.05 (9 H, s). LCMS (M+H) = 527.1.


Intermediate 115



5 *Benzyl cis-3-cyano-6-(hydroxymethyl)tetrahydro-2H-pyran-3-ylcarbamate.*

To a solution of benzyl cis-6-((tert-butyldiphenylsilyloxy)methyl)-3-cyanotetrahydro-2H-pyran-3-ylcarbamate (11.6 g, 21.94 mmol) in THF (200 mL) at 0°C was added 1M tetrabutyl ammonium fluoride/THF (43.9 mL, 43.9 mmol) and the resulting mixture heated at 60°C for 2h. After cooling to room temperature, the solvent was removed under reduced pressure and the crude product was purified by Biotage flash chromatography using 30% to 100% ethyl acetate/hexane to afford the title compound (3.0 g, 47% yield) as a yellow oil. ¹H NMR (500 MHz, CDCl₃) δ: 7.32-7.38 (m, 6H), 5.13 (s, 1H), 4.99 (brs, 1H), 4.65-4.72 (m, 1H), 3.60-3.66 (m, 2H), 3.45-3.49 (m, 1H), 3.26 (d, 1H, J = 11.29 Hz), 2.44-2.47 (m, 1H), 1.83-1.91 (m, 1H), 1.69-1.77 (m, 2H). LCMS (M+H) = 291.4.

Intermediate 116




20

Ethyl 3-(cis-3-(benzyloxycarbonylamino)-6-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-5-(2-ethoxy-2-oxoethyl)-2,5-dihydro-1,2,4-oxadiazole-5-carboxylate: To a solution of benzyl cis-3-cyano-6-(hydroxymethyl)tetrahydro-2H-pyran-3-ylcarbamate (2.0 g, 6.89 mmol) in EtOH (60 mL) was added 50% hydroxylamine

(1.267 mL, 20.67 mmol) and the reaction mixture was stirred 80 °C for 3 h. The reaction mixture was then cooled and concentrated and then dissolved in EtOH (100 mL) and treated with diethyl acetylenedicarboxylate (1.654 mL, 10.34 mmol). After stirring for 18 h at room temperature, the reaction mixture was diluted with ethyl acetate (250 mL), washed with H₂O (2 x 100 mL) followed by brine (50 mL), then dried over Na₂SO₄, filtered and concentrated. The crude was then purified by column chromatography using a Biotage flash chromatography system (10% to 50% ethyl acetate) to give a yellow oil (2.7 g, 79% yield) which was ~2:1 mixture of two diastereomers. ¹H NMR (500 MHz, CDCl₃) δ: 7.31-7.37 (m, 6H), 6.04 (s, 0.5 H), 5.76 (s, 1H), 5.16 (s, 0.5 H), 5.07 (s, 2H), 4.80 (d, 1H, J = 10.98 Hz), 4.09-4.16 (m, 4H), 3.57-3.68 (m, 5H), 2.27-2.30 (m, 1H), 1.94-2.00 (m, 2H), 1.72-1.78 (m, 6H). LCMS (M+H) = 494.27.

Intermediate 117

15




Ethyl 2-(cis-3-(benzyloxycarbonylamino)-6-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate. A solution of 20 ethyl 3-(cis-3-(benzyloxycarbonylamino)-6-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-5-(2-ethoxy-2-oxoethyl)-2,5-dihydro-1,2,4-oxadiazole-5-carboxylate (2.5 g, 5.07 mmol) in xylene (100 mL) was placed in a pre-heated oil bath and stirred for 16 h at 140 °C. The reaction mixture was cooled and extracted with 0.2 M aq. NaOH (3 x 75 mL) and the combined aqueous phase was washed with ethyl acetate (100 mL). The aqueous layer was then acidified with conc. HCl and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ solution was dried (Na₂SO₄), filtered and concentrated to afford the title compound as a light yellow solid (1.6 g, 71% yield). ¹H NMR (500 MHz, CDCl₃) δ: 10.76 (s, 1H), 7.28-7.36 (m, 6H), 5.79 (s, 1H), 5.03 (s, 2H), 4.70 (d, 1H, J

= 11.9 Hz), 4.41 (q, 2H, J = 7.02 Hz), 3.85-3.92 (m, 1H), 3.72 (d, 1H, J = 9.46 Hz), 3.58-3.64 (m, 2H), 2.28-2.31 (m, 1H), 2.10-2.20 (m, 1H), 1.54-1.70 (m, 3H), 1.38 (t, 3H, J = 7.17 Hz). LCMS (M+H) = 448.28.

5


Intermediate 118

*Ethyl 2-(3-(benzyloxycarbonylamino)-6-*

10 *((methylsulfonyloxy)methyl)tetrahydro-2H-pyran-3-yl)-5-(methylsulfonyloxy)-6-oxo-1,6-dihydropyrimidine-4-carboxylate.* To a stirred solution of ethyl 2-(cis-3-(benzyloxycarbonylamino)-6-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxylate (1.7 g, 3.80 mmol) in THF (50 mL) at 0C was added methanesulfonyl chloride (0.623 mL, 8.05 mmol) and triethylamine (2.65 mL, 19.00 mmol) in an ice bath and the mixture stirred at room temperature for 18 h. Water (50 mL) was then added and the mixture extracted with ethyl acetate (2x100 mL), washed with 1N HCl (50 mL) followed by brine (100 mL), then dried (Na₂SO₄), filtered and concentrated to give 1.9 g (83% yield) of the title compound as a light yellow solid along with tris-mesylate as a minor product by LC-MS. LCMS 15 (M+H) = 604.25.

20


Intermediate 119



Ethyl 10-(((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepine-2-carboxylate. A mixture of ethyl 2-(3-(benzyloxycarbonylamino)-6-((methylsulfonyloxy)methyl)tetrahydro-2H-pyran-3-yl)-5-(methylsulfonyloxy)-6-oxo-1,6-dihydropyrimidine-4-carboxylate (360 mg, 0.596 mmol) and Cs₂CO₃ (194 mg, 0.596 mmol) in DMF (6 mL) was heated at 60 °C for 18 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The resulting residue was taken up in ethyl acetate (200 mL) and extracted with 0.2 M aq. NaOH (3 x 50 mL). The combined aqueous phase was washed with ethyl acetate (150 mL). The aqueous layer was then acidified with conc. HCl and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ extracts were dried (Na₂SO₄), filtered and concentrated to afford the title compound as a light yellow solid (140 mg, 55% yield). ¹H NMR (500 MHz, CDCl₃) δ: 10.62 (s, 1H), 7.31-7.36 (m, 5H), 6.76 (s, 1H), 5.10 (s, 2H), 4.60-4.64 (m, 1H), 4.41-4.48 (m, 1H), 4.40 (q, 2H, J = 7.02 Hz), 4.10-4.16 (mn, 2H), 2.91-2.95 (m, 1H), 2.35-2.40 (m, 1H), 1.82-1.97 (m, 1H), 1.58-1.64 (m, 1H), 1.37 (t, 3H, J = 7.02 Hz). LCMS (M+H) = 430.46.

Intermediate 120

20




4-(Benzylxymethyl)furan-2(3H)-one. To a stirred solution of 4-hydroxydihydrofuran-2(3H)-one (25 g, 245 mmol) and benzyl 2,2,2-trichloroacetimidate (68.6 mL, 367 mmol) in CH₂Cl₂ (500 mL) and cyclohexane (500 mL) at 0 °C was added, dropwise, trifluoromethanesulfonic acid (2.175 mL, 24.49 mmol) over 10 min. The reaction mixture was stirred for 2 h at 0 °C and 48 h at room temperature. After this, it was diluted with hexanes (0.75 L), filtered and the filtrate washed with sat. aq. NaHCO₃ (150 mL), dried (Na₂SO₄), filtered and concentrated. The resulting yellow oil was purified by flash column chromatography on silica gel column using 5-40% ethyl acetate/hexane (5% increment for every liter) to afford 4-(benzylxymethyl)furan-2(3H)-one (35.52 g, 185 mmol, 75% yield) as a

clear yellow oil. ^1H NMR (500 MHz, CDCl_3) δ : 7.34 - 7.39 (2 H, m), 7.29 - 7.34 (3 H, m), 4.53 (1 H, d, $J_{\text{AB}} = 11.9$ Hz), 4.52 (1 H, d, $J_{\text{AB}} = 11.90$ Hz), 4.34 - 4.41 (3 H, m), 2.64 - 2.68 (2 H, m).


5

Intermediate 121



4-(Benzylxy)tetrahydrofuran-2-yl acetate. To a stirred solution of 4-(benzylxy)dihydrofuran-2(3H)-one (3.15 g, 16.39 mmol) in CH_2Cl_2 (100 mL) was added, dropwise, 1M DIBAL-H/toluene (2.56 g, 18.03 mmol) over 10 min at -78 °C. After 2 h, pyridine (3.98 mL, 49.2 mmol), solid 4-(dimethylamino)pyridine (DMAP) (2.202 g, 18.03 mmol) and Ac_2O (6.19 mL, 65.6 mmol) were added and the solution was stirred for 6 h at -78 °C and slowly warmed to room temperature overnight (15 h). The resulting orange reaction mixture was stirred with sat. NH_4Cl (50 mL) for 30 min. The organic layer was separated and washed with 1N aq. NaHSO_4 (3 x 50 mL). The combined aq. layers were extracted with CH_2Cl_2 (2 x 50 mL) and the combined CH_2Cl_2 layers washed with 1N aq. NaHSO_4 (50 mL), sat. NaHCO_3 (50 mL) followed by brine (50 mL), then dried (Na_2SO_4), filtered and concentrated to afford a yellow oil. Flash column chromatography on a silica gel column using 15-30% ethyl acetate/hexane provided 4-(benzylxy)tetrahydrofuran-2-yl acetate (3.5769 g, 15.14 mmol, 92% yield) as a clear oil. ^1H NMR (500 MHz, CDCl_3) δ : 7.26 - 7.39 (5 H, m), 6.41 (0.3 H, dd, $J = 5.7, 2.0$ Hz), 6.24 - 6.30 (0.7 H, dd, $J = 4.9, 1.8$ Hz), 4.46 - 4.58 (2 H, m), 4.32 - 4.37 (0.3 H, m), 4.22 - 4.28 (0.7 H, m), 4.00 - 4.10 (2 H, m), 2.15 - 2.38 (2 H, m), 2.07 (3 H, s). LCMS (M-C₂H₃O) calcd for C₁₃H₁₆O₄: 177.09; found: 177.0.


Intermediate 122



5 *2-Allyl-4-(benzyloxy)tetrahydrofuran.* To a stirred solution of 4-(benzyloxy)tetrahydrofuran-2-yl acetate (3.57 g, 15.11 mmol) and allyltrimethylsilane (9.65 mL, 60.4 mmol) in CH₂Cl₂ (200 mL) at -78 °C was added a solution of tin(IV) bromide (7.95 g, 18.13 mmol) in CH₂Cl₂ (20 mL) over 15 min. After 1 h, the ice bath was removed and stirring continued at room temperature for 3
10 h. The CH₂Cl₂ was removed using a rotovap and the residue taken up in 1:1 Et₂O/hexane (200 mL), washed with sat. Na₂CO₃ (2 x 50 mL), water (2 x 50 mL) followed by brine (50 mL), then dried (Na₂SO₄), filtered and concentrated to give 2-allyl-4-(benzyloxy)tetrahydrofuran (3.15 g, 14.43 mmol, 95% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.27 - 7.39 (5 H, m), 5.76 - 5.90 (1 H, m), 5.03 - 5.15 (2 H, m), 4.49 (1.6 H, d, *J* = 2.3 Hz), 4.43 (0.4 H, d, *J* = 2.3 Hz), 4.17 - 4.23 (1 H, m), 4.01 (0.45 H, d, *J* = 1.8 Hz), 3.98 (0.55 H, d, *J* = 1.8 Hz), 3.85 - 3.94 (1 H, m), 3.73 (1 H, dd, *J* = 9.8, 5.3 Hz), 2.43 - 2.53 (1 H, m), 2.30 - 2.40 (1 H, m), 2.19 - 2.28 (1 H, m), 1.69 - 1.77 (1 H, m). LCMS (M+H) calcd for C₁₄H₁₉O₂: 219.14; found: 219.05.

20


Intermediate 123



25 *2-(4-(Benzoyloxy)tetrahydrofuran-2-yl)acetaldehyde.* To a stirred solution of 2-allyl-4-(benzyloxy)tetrahydrofuran (3.1 g, 14.20 mmol) in THF (100 mL) and water (100 mL) was added osmium tetroxide (2 mL, 0.327 mmol) at room temperature. After a few minutes, sodium periodate (7.59 g, 35.5 mmol) was added in small portions over 15 min. After 20 h, the reaction mixture was diluted with ether (200 mL), filtered and the filtrate washed with water (2 x 50 mL) followed by brine

(50 mL), then dried (Na_2SO_4), filtered and concentrated to give a gray oil. Flash column chromatography on a silica gel column using 15 - 30% ethyl acetate/hexane (5% increment) gave 2-(4-(benzyloxy)tetrahydrofuran-2-yl)acetaldehyde (2.4624 g, 11.18 mmol, 79% yield) as a colorless liquid consisting of a ~4:1 mixture of diastereomers. ^1H NMR (500 MHz, CDCl_3) δ : 9.81 (0.8 H, s), 9.77 (0.2 H, s), 7.23 - 7.37 (5 H, m), 4.48 (1.6 H, s), 4.41 (0.4 H, s), 4.29 - 4.31 (1 H, m), 4.23 - 4.20 (0.8 H, m), 4.12 - 4.09 (0.2 H, m), 4.04 - 4.06 (0.8 H, m), 3.90 - 3.94 (0.2 H, m), 3.73 (0.8 H, dd, J = 9.9, 4.7 Hz), 3.64 (0.2 H, dd, J = 9.9, 4.7 Hz), 2.82 - 2.94 (1 H, m), 2.60 - 2.73 (1 H, m), 2.25 - 2.39 (1 H, m), 1.77 (0.8 H, m), 1.59 - 1.69 (0.2 H, m). LCMS 10 (M+H) calcd for $\text{C}_{13}\text{H}_{17}\text{O}_3$: 221.12; found: 221.05.


Intermediate 124



15


2-(4-(Benzyl)tetrahydrofuran-2-yl)ethanol. To a stirred solution of 2-(4-(benzyloxy)tetrahydrofuran-2-yl)acetaldehyde (2.46 g, 11.17 mmol) in methanol (50 mL) was added NaBH_4 (0.423 g, 11.17 mmol), in small portions, at room temperature. After 2 h, the reaction mixture was concentrated and the residue was taken up in sat. Na_2CO_3 (25 mL) and extracted with CH_2Cl_2 (3 x 25 mL). The combined CH_2Cl_2 layers were dried (Na_2SO_4), filtered and concentrated to give 2-(4-(benzyloxy)tetrahydrofuran-2-yl)ethanol (2.3847 g, 10.73 mmol, 96% yield) as a colorless oil consisting of a ~4:1 mixture of diastereomers. ^1H NMR (500 MHz, CDCl_3) δ : 7.23 - 7.38 (5 H, m), 4.49 (1.6 H, s), 4.42 (0.4 H, d, J = 1.5 Hz), 4.17 - 4.29 (1 H, m), 3.99 - 4.11 (2 H, m), 3.74 - 3.82 (2 H, m), 3.70 (0.8 H, dd, J = 9.8, 4.9 Hz), 3.61 (0.2 H, dd, J = 9.9, 5.0 Hz), 2.52 (0.8 H, t, J = 5.6 Hz), 2.48 (0.2 H, t, J = 5.6 Hz), 2.30 (0.8 H, dt, J = 13.6, 7.2 Hz), 2.16 - 2.25 (0.2 H, m), 1.61 - 1.99 (3 H, m). LCMS (M+H) calcd for $\text{C}_{13}\text{H}_{19}\text{O}_3$: 223.13; found: 223.07.

Intermediate 125



5 *4-(Benzylxoy)-2-(2-chloroethyl)tetrahydrofuran.* To an ice-cold stirred solution of oxalyl chloride (1.005 mL, 11.48 mmol) in CH₂Cl₂ (30 mL) was added DMF (0.914 mL, 11.80 mmol), dropwise, over 10 min. The ice bath was removed and the white slurry stirred for 30 min at room temperature. The reaction was cooled to 0 °C in an ice bath and a solution of 2-(4-(benzyloxy)tetrahydrofuran-2-yl)ethanol (2.3847 g, 10.73 mmol) in CH₂Cl₂ (30 mL) was rapidly added via cannula. The resulting clear yellow solution was stirred for 2 h at room temperature and 3 h at reflux. After cooling to room temperature the mixture was concentrated and the resulting yellow residue taken up in Et₂O (100 mL), washed with water (3 x 50 mL) followed by brine (25 mL), then dried (MgSO₄), filtered and concentrated to give a colorless liquid. Flash column chromatography on a silica gel column using 9:1 hexane/ethyl acetate provided 4-(benzyloxy)-2-(2-chloroethyl)tetrahydrofuran (2.47 g, 10.26 mmol, 96% yield) as a colorless oil. ¹H NMR (500 MHz, CDCl₃) δ: 7.27 - 7.37 (5 H, m), 4.48 (1.6 H, s), 4.41 (0.4 H, s), 4.17 - 4.23 (1 H, m), 3.96 - 4.11 (2 H, m), 3.72 (0.8 H, dd, *J* = 10.1, 4.9 Hz), 3.59 - 3.70 (2.2 H, m), 2.30 (0.8 H, dt, *J* = 13.4, 7.0 Hz), 2.09 - 2.25 (1.2 H, m), 1.89 - 2.01 (1 H, m), 1.69 - 1.74 (0.8 H, m), 1.57 - 1.64 (0.2 H, m). LCMS (M+H) calcd for C₁₃H₁₈ClO₃: 241.1; found: 241.05.


Intermediate 126



Benzyl 5-(2-chloroethyl)-3-cyanotetrahydrofuran-3-ylcarbamate. A solution of 4-(benzyloxy)-2-(2-chloroethyl)tetrahydrofuran (2.47 g, 10.26 mmol) in ethyl acetate (50 mL) was deoxygenated for 5 min by bubbling N₂ through. 10% Pd/C


(0.145 g, 1.363 mmol) was added the flask evacuated then placed under 1 atm H₂ for 4 h. The reaction was filtered and concentrated to give a colorless oil (1.74 g) which was used in the next step without purification. A solution of 4-methylmorpholine N-oxide (1.803 g, 15.39 mmol) in CH₂Cl₂ (50 mL), intermediate 5-(2-chloroethyl)tetrahydrofuran-3-ol and 4Å powdered molecular sieves (4 g) was stirred for 30 min. To this mixture was added tetrapropylammonium perruthenate (0.036 g, 0.103 mmol) and stirring continued for 24 h at room temperature. The reaction was concentrated, then diluted with Et₂O (100 mL) and filtered through a plug of CELITE®. The filtrate was concentrated to afford 5-(2-chloroethyl)dihydrofuran-3(2H)-one as viscous colorless oil. To a stirred mixture of 5-(2-chloroethyl)dihydrofuran-3(2H)-one, aqueous ammonia (5 mL) and 1N HCl (10 mL) in methanol 20 mL was added NaCN (0.43 g, 8.77 mmol) at room temperature. The yellow solution was stirred for 42 h and concentrated. The solution was made basic with sat. Na₂CO₃ (20 mL) and extracted with CH₂Cl₂ (3 x 25 mL). To the combined CH₂Cl₂ extracts was added sat. Na₂CO₃ (10 mL), and the mixture cooled to 0 °C with an ice bath and treated with benzyl chloroformate (1.465 mL, 10.26 mmol). After 1 h, the ice bath was removed and the reaction was stirred at room temperature for 4 h. The organic layer was separated, dried (Na₂SO₄), filtered, concentrated and purified by flash column chromatography on silica gel column using 10, 20, 30 and 40% ethyl acetate/hexane to afford the title compound benzyl 5-(2-chloroethyl)-3-cyanotetrahydrofuran-3-ylcarbamate (0.863 g, 2.80 mmol, 27% yield), viscous pale yellow oil. ¹H NMR (500 MHz, CDCl₃) δ: 7.12 - 7.42 (5 H, m), 5.16 (2 H, s), 4.40 (0.2 H, d, *J*= 9.8 Hz), 4.25 - 4.33 (0.8 H, m), 4.10 - 4.19 (1 H, m), 3.95 (0.2 H, d, *J*= 9.8 Hz), 3.56 - 3.68 (1.8 H, m), 2.86 (1 H, dd, *J*= 13.4, 7.0 Hz), 2.53 - 2.62 (1 H, m), 2.27 (1 H, dd, *J*= 13.4, 8.9 Hz), 1.94 - 2.17 (2 H, m). LCMS (M+H) calcd for C₁₅H₁₈ClN₂O₃: 309.10; found: 309.10.

Intermediate 127



5 *Diethyl 2-(amino(3-(benzyloxycarbonylamino)-5-(2-chloroethyl)tetrahydrofuran-3-yl)methyleneaminoxy)but-2-enedioate.* A solution of benzyl 5-(2-chloroethyl)-3-cyanotetrahydrofuran-3-ylcarbamate (0.86 g, 2.79 mmol) and 50% aq. hydroxylamine (0.853 mL, 13.93 mmol) in THF (25 mL) was stirred at room temperature for 18 h. The mixture was concentrated and the residue was
10 redissolved in EtOH (20 mL) and treated with diethyl acetylenedicarboxylate (0.490 mL, 3.06 mmol) at room temperature. After 24 h, the reaction mixture was concentrated and the resulting pale yellow residue was taken up in Et₂O (75 mL), washed with 1% aq. NH₄OH (20 mL), water (20 mL) followed by brine (20 mL), then dried (Na₂SO₄), filtered, concentrated and purified by flash column
15 chromatography on silica gel column using 1:3, 2:3 and 1:1 ethyl acetate/hexane to afford diethyl 2-(amino(3-(benzyloxycarbonylamino)-5-(2-chloroethyl)tetrahydrofuran-3-yl)methyleneaminoxy)but-2-enedioate (1.1707 g, 2.287 mmol, 82% yield) as a pale yellow paste. ¹H NMR (500 MHz, CDCl₃) δ: 7.30 - 7.39 (5 H, m), 5.75 - 5.88 (2 H, m), 5.26 - 5.62 (2 H, m), 5.05 - 5.12 (2 H, m), 4.21 - 4.37 (3 H, m), 4.12 - 4.20 (3 H, m), 4.02 - 4.10 (1 H, m), 3.57 - 3.67 (2 H, m), 2.60 - 2.85 (1 H, m), 1.87 - 2.20 (3 H, m), 1.29 - 1.38 (3 H, m), 1.22 - 1.28 (3 H, m). LCMS (M+H) calcd for C₂₃H₃₁ClN₃O₈: 512.18; found: 512.25.

Intermediate 128




5 *Ethyl 11-((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-7,8,10,11-*
tetrahydro-4H,6H-8,11-methanopyrimido[2,1-d][1,5]oxazocine-2-carboxylate. A
 solution of diethyl 2-(amino(3-(benzyloxycarbonylamino)-5-(2-
 chloroethyl)tetrahydrofuran-3-yl)methyleneaminooxy)but-2-enedioate (1.15 g, 2.246
 mmol) in xylene (50 mL) was heated at 135 - 140 °C (oil bath temp.) for 4 h, then
 10 cooled, concentrated and the resulting residue dissolved in DMF (20 mL) and stirred
 with K₂CO₃ (0.621 g, 4.49 mmol) at 80-85 °C. After 9 h, the reaction was cooled,
 concentrated and the residue was acidified with 1M aq. HCl (20 mL) and extracted
 with CH₂Cl₂ (2 x 25 mL). The combined organic layers were washed with water (25
 mL) followed by brine (25 mL), then dried (Na₂SO₄), filtered and concentrated to
 15 give dark brown paste which was triturated with ethyl acetate to provide the a by-
 product, ethyl 3-hydroxy-4,6-dioxo-9,10-dihydro-4H,8H-10,12a-
 methanopyrimido[1',2':3,4]imidazo[5,1-c][1,4]oxazepine-2(12H)-carboxylate (0.1922
 g, 0.598 mmol, 27% yield) as a gray powder. The mother liquor was purified by
 prep-HPLC to afford the title compound (0.071 g, 0.165 mmol, 7% yield)as an off-
 20 white solid. Ethyl 3-hydroxy-4,6-dioxo-9,10-dihydro-4H,8H-10,12a-
 methanopyrimido[1',2':3,4]imidazo[5,1-c][1,4]oxazepine-2(12H)-carboxylate: ¹H
 NMR (500 MHz, CDCl₃) δ: 11.20 (1 H, brs), 4.85 (1 H, m), 4.49 (2 H, q, *J* = 7.3 Hz),
 4.28 (1 H, d, *J* = 8.9 Hz), 4.14 (1 H, dd, *J* = 13.4, 7.6 Hz), 4.08 (1 H, dd, *J* = 8.9, 1.0
 Hz), 3.42 (1 H, ddd, *J* = 13.4, 11.3, 6.1 Hz), 2.69 (1 H, ddd, *J* = 11.3, 6.3, 1.4 Hz),
 25 2.00 - 2.07 (1 H, m), 1.89 (1 H, dd, *J* = 11.3, 1.0 Hz), 1.71 (1 H, ddd, *J* = 13.2, 11.5,
 7.9 Hz), 1.44 (3 H, t, *J* = 7.3 Hz). LCMS (M+H) calcd for C₁₄H₁₆N₃O₆: 322.10;
 found: 322.08. Ethyl 11-((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-7,8,10,11-

tetrahydro-4H,6H-8,11-methanopyrimido[2,1-d][1,5]oxazocine-2-carboxylate. ¹H NMR (500 MHz, CDCl₃) δ: 10.48 (1 H, brs), 7.29 - 7.42 (5 H, m), 6.70 (1 H, brs), 5.45 (1 H, dd, *J* = 15.3, 3.4 Hz), 5.14 (2 H, d, *J_{AB}* = 12.4 Hz), 4.78 (1 H, t, *J* = 6.7 Hz), 4.57 (2 H, d, *J_{AB}* = 10.4 Hz), 4.37 (2 H, q, *J* = 7.2 Hz), 3.72 (1 H, t, *J* = 13.6 Hz), 5 3.02 (1 H, dd, *J* = 11.7, 8.4 Hz), 2.14 - 2.23 (1 H, m), 1.87 (1 H, d, *J* = 12.5 Hz), 1.58 - 1.67 (1 H, m), 1.35 (3 H, t, *J* = 7.2 Hz). LCMS (M+H) calcd for C₂₁H₂₄N₃O₇: 430.16; found: 430.12.

Intermediate 129


10



Ethyl 1,4-dioxaspiro[4.5]decane-8-carboxylate. This compound was prepared from ethyl cyclohexanone-4-carboxylate by following the procedure of 15 Pearson et al. (*J. Org. Chem.*, 1997, 62, 5284). A mixture of ethylene glycol (131 mL, 2350 mmol), ethyl 4-oxocyclohexanecarboxylate (100 g, 588 mmol), and p-toluene sulfonic acid monohydrate (2.012 g, 10.58 mmol) in benzene (125 mL) was stirred overnight at room temperature, under N₂. The solution was poured into Et₂O (1 L) and the mixture washed with water (3 x 300 mL), sat. NaHCO₃ (100 mL) followed by brine (100 mL). The organic layer was then dried (MgSO₄), filtered, and concentrated to yield the title compound (125.5 g, 586 mmol, 100 % yield) as a clear oil: LCMS m/z 215.13 (M+H).


Intermediate 130

25



1,4-Dioxaspiro[4.5]decan-8-ylmethanol. The procedure of Pearson et al., *J. Org. Chem.*, **1997**, *62*, 5284 was followed. To a stirred 1M solution of lithium aluminum hydride (148 mL, 148 mmol) cooled to 0 °C, in an ice-bath, was added a solution of ethyl 1,4-dioxaspiro[4.5]decane-8-carboxylate (31.8 g, 148 mmol) in THF (30 mL), dropwise, over a period of 50 min. The ice-bath was removed and the mixture stirred under N₂ overnight. It was then cooled and water (20 mL) slowly added under vigorous stirring [note: very vigorous reaction with heat and foam generation] followed by the addition of 15% NaOH (20 mL). The insoluble material was removed by filtration through CELITE®, washing with THF. The combined filtrate and washings were dried (Na₂SO₄), filtered and concentrated to provide the title compound, 1,4-dioxaspiro[4.5]decan-8-ylmethanol (26.3 g, 153 mmol, 103 % yield) as a clear oil: LCMS: m/z 173 (M+H) at 0.67 min; ¹H NMR (500 MHz, CDCl₃) δ ppm 1.11-1.25 (2H, m), 1.39-1.52 (3H, m), 1.70 (4H, d, J=9.5 Hz), 2.75 (1H, s), 3.39 (2H, d, J=6.4 Hz), 3.86-3.89 (4H, m); ¹³C NMR (126 MHz, CDCl₃) δ ppm 26.7, 34.2, 39.1, 64.3, 67.5, 109.2.

Intermediate 131




20

(1,4-Dioxaspiro[4.5]decan-8-ylmethoxy)(tert-butyl)dimethylsilane. To a solution of 1,4-dioxaspiro[4.5]decan-8-ylmethanol (8.60 g, 49.9 mmol) and imidazole (8.50 g, 125 mmol) in DMF (50 mL) was added tert-butyldimethylsilyl chloride (9.03 g, 59.9 mmol), and the mixture stirred at room temp. for 18 hours. The mixture was concentrated *in vacuo* and the residue redissolved in ethyl acetate (20 mL), washed with water followed by brine, then dried (Na₂SO₄), and concentrated *in vacuo* to provide the title compound (14.5 g, 50.6 mmol, 101 % yield) as a clear oil: LCMS: m/z 287 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 0.01 (6H, s), 0.86 (9H, s), 1.14-1.26 (2H, m), 1.45-1.54 (3H, m), 1.67-1.82 (4H, m),

3.40 (2H, d, $J=6.7$ Hz), 3.91 (4H, s); ^{13}C NMR (126 MHz, CDCl_3) δ ppm -5.25, 18.5, 26.1, 26.9, 34.3, 39.2, 64.3, 68.0, 109.3.

Intermediate 132


5



4-((tert-Butyldimethylsilyloxy)methyl)-1-(2-(trimethylsilyloxy)ethoxy)cyclohexanecarbonitrile. Reference: G. Olah et al., Synthesis, 498 (1983). Zinc iodide (0.034 g, 0.107 mmol) was added to a mixture of (1,4-dioxaspiro[4.5]decan-8-ylmethoxy)(tert-butyl)dimethylsilane (14.5 g, 50.6 mmol) and trimethylsilyl cyanide (6.79 mL, 50.6 mmol) at 10°C under an atmosphere of N_2 , and the mixture stirred at room temperature overnight. The reaction mixture was treated with additional aliquots of ZnI_2 and trimethylsilylcyanide and heated overnight until NMR (CDCl_3) and HPLC analysis indicated significant conversion to the title product. The mixture was then diluted with CH_2Cl_2 (25 mL) and washed with water followed by brine, then dried and concentrated to yield 15.7 g of a light red oil, which was purified by SiO_2 column chromatography (530 g, SiO_2 , 3-5-10-20-40% ethyl acetate/ CH_2Cl_2) to obtain the title compound (11.0 g, 28.5 mmol, 56 % yield) as an amber oil: SiO_2 -TLC: $R_f = 0.62$ (5% ethyl acetate/ CH_2Cl_2), molybdate stain; LCMS: m/z 386 (M+H, minor) and 359 (M-CN, major); ^1H NMR (500 MHz, CDCl_3) δ ppm 0.03, 0.03 (6H, 2s), 0.11, 0.11 (9H, 2s), 0.88 (9H, s), 1.21-1.32 (2H, m), 1.46-1.56 (3H, m), 1.87 (2H, dd, $J=14.2, 2.3$ Hz), 2.26 (2H, d, $J=11.9$ Hz), 3.43 (2H, d, $J=6.4$ Hz), 3.69 (2H, t, $J=4.9$ Hz), 3.75 (2H, t, $J=5.0$ Hz), ^{13}C NMR (126 MHz, CDCl_3) δ ppm -5.31, -0.23, 18.4, 26.0, 26.1, 35.1, 39.3, 61.9, 66.8, 67.5, 76.5, 119.8.


Reference: G. Olah et al., Synthesis, 498 (1983). Zinc iodide (0.034 g, 0.107 mmol) was added to a mixture of (1,4-dioxaspiro[4.5]decan-8-ylmethoxy)(tert-butyl)dimethylsilane (14.5 g, 50.6 mmol) and trimethylsilyl cyanide (6.79 mL, 50.6 mmol) at 10°C under an atmosphere of N_2 , and the mixture stirred at room temperature overnight. The reaction mixture was treated with additional aliquots of ZnI_2 and trimethylsilylcyanide and heated overnight until NMR (CDCl_3) and HPLC analysis indicated significant conversion to the title product. The mixture was then diluted with CH_2Cl_2 (25 mL) and washed with water followed by brine, then dried and concentrated to yield 15.7 g of a light red oil, which was purified by SiO_2 column chromatography (530 g, SiO_2 , 3-5-10-20-40% ethyl acetate/ CH_2Cl_2) to obtain the title compound (11.0 g, 28.5 mmol, 56 % yield) as an amber oil: SiO_2 -TLC: $R_f = 0.62$ (5% ethyl acetate/ CH_2Cl_2), molybdate stain; LCMS: m/z 386 (M+H, minor) and 359 (M-CN, major); ^1H NMR (500 MHz, CDCl_3) δ ppm 0.03, 0.03 (6H, 2s), 0.11, 0.11 (9H, 2s), 0.88 (9H, s), 1.21-1.32 (2H, m), 1.46-1.56 (3H, m), 1.87 (2H, dd, $J=14.2, 2.3$ Hz), 2.26 (2H, d, $J=11.9$ Hz), 3.43 (2H, d, $J=6.4$ Hz), 3.69 (2H, t, $J=4.9$ Hz), 3.75 (2H, t, $J=5.0$ Hz), ^{13}C NMR (126 MHz, CDCl_3) δ ppm -5.31, -0.23, 18.4, 26.0, 26.1, 35.1, 39.3, 61.9, 66.8, 67.5, 76.5, 119.8.

Intermediate 133



5 *4-((tert-Butyldimethylsilyloxy)methyl)-1-(2-hydroxyethoxy)cyclohexanecarbonitrile.* To a solution of 4-((tert-butyldimethylsilyloxy)methyl)-1-(2-(trimethylsilyloxy)ethoxy)cyclohexanecarbonitrile (9.70 g, 25.1 mmol) in methanol (25 mL) was added pyridine hydrochloride (2.91 g, 25.1 mmol) and the mixture stirred at room temperature for 30 min. The reaction was immediately quenched by careful addition of sat'd aqueous NaHCO₃ (30 mL), and extracted with ethyl acetate (25 mL x 2). The combined organic extracts were washed with brine, dried (Na₂SO₄) and concentrated to provide 5.82 g of a cloudy light brown oil, which was purified by column chromatography (SiO₂, 300 g, 20% ethyl acetate/CH₂Cl₂) to provide the title compound, 4-((tert-butyldimethylsilyloxy)methyl)-1-(2-hydroxyethoxy)cyclohexanecarbonitrile (3.45 g, 11.0 mmol, 44 % yield) as an amber oil: TLC: Rf 0.15 (5% ethyl acetate/CH₂Cl₂), 0.51 (20% ethyl acetate/CH₂Cl₂); molybdate stain; LC/MS: m/z 314 (M+H), 287 (M-CN), 331 (M+H₂O); ¹H NMR (500 MHz, CDCl₃) δ ppm 0.03 (6H, s), 0.88 (9H, s), 1.23 (1H, t, J=7.0 Hz), 1.25-1.33 (2H, m), 1.42-1.56 (3H, m), 1.88 (2H, d, J=11.9 Hz), 2.27 (2H, d, J=12.8 Hz), 3.43 (2H, d, J=6.4 Hz), 3.71-3.81 (4H, m); ¹³C NMR (126 MHz, CDCl₃) δ ppm -5.32, 18.4, 26.0, 26.1, 35.2, 39.3, 61.8, 66.6, 67.4, 76.6, 119.6.


Intermediate 134



5 *4-((tert-Butyldimethylsilyloxy)methyl)-N-hydroxy-1-(2-hydroxyethoxy)cyclohexane carboximidamide.* A mixture of 4-((tert-butylidimethylsilyloxy)methyl)-1-(2-hydroxyethoxy)cyclohexane carbonitrile (3.44 g, 10.97 mmol) and 50% aq. hydroxylamine (0.725 g, 10.97 mmol) in ethanol (17 mL) was heated at 80°C (oil bath temp) for 6 h. The mixture was concentrated to dryness
10 in *vacuo* and the residual solid triturated with ethanol to obtain the title compound, 4-((tert-butylidimethylsilyloxy)methyl)-N-hydroxy-1-(2-hydroxyethoxy)cyclohexane carboximidamide (3.30 g, 9.52 mmol, 87 % yield) as a tan crystalline powder: HPLC: retention time = 1.96 min (area percent = 91%); LCMS: m/z 347 (M+H), 369 (M+Na); ¹H NMR (500 MHz, CDCl₃) δ ppm 0.02 (6H, s), 0.87 (9H, s), 1.23 (1H, t, J=7.0 Hz), 1.25-1.34 (2H, m), 1.49-1.58 (2H, m), 1.57-1.63 (1H, m), 1.67-1.77 (2H, m), 2.05-2.16 (2H, m), 3.41-3.46 (4H, m), 3.69-3.72 (2H, m), 4.78 (2H, s), 7.95 (1H, brs); ¹³C NMR (126 MHz, CDCl₃) δ ppm -5.22, 18.5, 25.0, 26.1, 32.0, 38.6, 62.1, 64.1, 66.9, 77.9, 154.7.


20

Intermediate 135




Ethyl 5-(benzoyloxy)-2-(1-(2-(benzoylperoxy)ethoxy)-4-((tert-butyldimethylsilyloxy)methyl)cyclohexyl)-6-oxo-1,6-dihydropyrimidine-4-carboxylate. To a cold solution of 4-((tert-butyldimethylsilyloxy)methyl)-N-hydroxy-1-(2-hydroxyethoxy)cyclohexanecarboximidamide (3.80 g, 10.97 mmol) in ethanol (40 mL) was added diethyl but-2-ynedioate (2.05 g, 12.05 mmol) and the mixture stirred at room temp. under N₂ for 6 h and then kept in a freezer overnight. The mixture was concentrated to dryness *in vacuo* to provide diethyl 2-(4-((tert-butyldimethylsilyloxy)methyl)-1-(2-hydroxyethoxy)cyclohexanecarboxyimidamidoxy)fumarate (6.03 g, 11.67 mmol, 106 % yield) as a crude oil: HPLC: retention time = 3.81 min (area percent = 78%); LCMS: m/z 517 (M+H). A solution of this crude oil (6.00 g, 11.61 mmol) in xylene (220 mL) was heated at 150°C under N₂ for 4.5 h. The mixture was concentrated *in vacuo* to obtain 7.0 g of the crude pyrimidinone as a light brown oil. The crude oil (7.0 g) was dissolved in pyridine (50 mL) and added benzoic anhydride (5.52 g, 11 mmol) added after which the mixture stirred at room temp. overnight. Most of pyridine was removed *in vacuo*, and the residual oil dissolved in ethyl acetate (30 mL) and washed sequentially with 1N HCl, sat'd aq. NaHCO₃ followed by brine, then dried (Na₂SO₄) and concentrated to dryness to provide 8.4 g of a crude 1 oil, which was purified by column chromatography (SiO₂, 215 g, 5-10% ethyl acetate/CH₂Cl₂) to yield the title compound, ethyl 5-(benzoyloxy)-2-(1-(2-(benzoyloxy)ethoxy)-4-((tert-butyldimethylsilyloxy)methyl)cyclohexyl)-6-oxo-1,6-dihydropyrimidine-4-carboxylate (3.7 g, 5.45 mmol, 47 % yield) as an amber oil: HPLC: retention time = 4.48 min(area percent = 95%); TLC: RF 0.55 (10% ethyl acetate/CH₂Cl₂); LCMS: m/z 679 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 0.03 (6H, s), 0.87 (9H, s), 1.13 (3H, t, J=7.2 Hz), 1.50 (2H, m), 1.60-1.70 (1H, m), 1.71-1.84 (4H, m), 2.24 (2H, m), 3.49 (2H, d, J=7.0 Hz), 3.66 (2H, t, J=4.5 Hz), 4.26 (2H, q, J=7.3 Hz), 4.46 (2H, q, J=4.5 Hz), 7.41 (2H, t, J=7.8 Hz), 7.48 (2H, t, J=7.8 Hz), 7.50-7.55 (1H, m), 7.62 (1H, t, J=7.5 Hz), 8.03 (2H, d, J=7.3 Hz), 8.17 (2H, d, J=7.0 Hz), 11.20 (1H, br). ¹³C NMR (126 MHz, CDCl₃) δ ppm -5.22, 14.0, 18.4, 24.3, 26.1, 31.9, 37.5, 61.6, 62.2, 64.0, 66.4, 78.9, 128.4, 128.6, 128.7, 129.8, 129.9, 130.7, 133.2, 134.1, 138.4, 142.8, 158.8, 160.0, 162.9, 163.5, 166.7.

Intermediate 136



5 *Ethyl 5-(benzoyloxy)-2-(1-(2-(benzoylperoxy)ethoxy)-4-(hydroxymethyl)cyclohexyl)-6-oxo-1,6-dihydropyrimidine-4-carboxylate.* Journal of
Organic Chemistry 45:4797 (1980). To a solution of ethyl 5-(benzoyloxy)-2-(1-(2-(benzoylperoxy)ethoxy)-4-((tert-butyldimethylsilyloxy) methyl)cyclohexyl)-6-oxo-
1,6-dihydropyrimidine-4-carboxylate (4.28 g, 6.30 mmol) in EtOH (30 mL) was
10 added a dilute HCl solution in EtOH (5 mL; prepared from 6N-HCl 0.3 mL and 95%
EtOH 9.7 mL), and the mixture stirred at room temp for 2h. The mixture was
concentrated *in vacuo* and the residue taken up in ethyl acetate (20 mL), washed with
water, sat'd NaHCO₃ followed by brine, then dried (Na₂SO₄) and concentrated. The
residue was dissolved in CH₂Cl₂ and diluted with hexanes and concentrated to
15 dryness to obtain 3.26 g of a tan foam which was triturated with hexanes to provide
the title compound, ethyl 5-(benzoyloxy)-2-(1-(2-(benzoyloxy)ethoxy)-4-(hydroxymethyl)cyclohexyl)-6-oxo-1,6-dihydropyrimidine-4-carboxylate (3.21 g,
5.69 mmol, 90 % yield) as a tan colored foam: HPLC: retention time = 2.66 min (area
percent = 86%); TLC: RF 0.5 (50% ethyl acetate/CH₂Cl₂); LCMS: m/z 565 (M+H);
20 ¹H NMR (500 MHz, CDCl₃) δ ppm 1.12 (3H, t, J=7.2 Hz), 1.58-1.68 (2H, m), 1.69-
1.74 (1H, m), 1.74-1.86 (4H, m), 2.14-2.24 (2H, m), 3.58 (2H, d, J=6.4 Hz), 4.26
(2H, q, J=7.1 Hz), 4.48 (2H, t, J=5 Hz), 7.45 (2H, t, J=7.8 Hz), 7.50 (2H, t, J=7.6 Hz),
7.55 (1H, t, J=7.3 Hz), 7.64 (1H, t, J=7.5 Hz), 8.08 (2H, d, J=7.6 Hz), 8.18 (2H, d,
J=7.9 Hz), 10.29 (1H, br); ¹³C NMR (126 MHz, CDCl₃) δ ppm 14.0, 24.0, 31.7, 37.2,
25 61.7, 62.3, 64.0, 65.9, 78.6, 128.3, 128.7, 128.7, 129.6, 129.9, 130.7, 133.4, 134.2,
138.6, 142.6, 157.8, 159.9, 162.9, 163.5, 166.8.


Intermediate 137



5 *5-Benzoyloxy-1-(2-benzoyloxyethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2.7}]trideca-2,4-diene-4-carboxylic acid ethyl ester.* Under N₂, to a cold (ice-bath), stirred solution of ethyl 5-(benzoyloxy)-2-(1-(2-(benzoylperoxy)ethoxy)-4-(hydroxymethyl)cyclohexyl)-6-oxo-1,6-dihdropyrimidine-4-carboxylate (1.13 g, 2.001 mmol) in THF (10 mL) was added, 10 triethylamine (0.279 mL, 2.00 mmol), followed by methane sulfonylchloride (0.156 mL, 2.001 mmol), and the mixture stirred at room temperature overnight. Additional triethylamine (50 mg, 0.5 mmol) and methane sulfonylchloride (40 uL, 0.5 mmol) were added and the mixture stirred at room temperature 2 h. The mixture was diluted with ethyl acetate (15 mL) and washed with water followed by brine, then dried 15 (Na₂SO₄) and concentrated to provide ethyl 5-(benzoyloxy)-2-(1-(2-(benzoyloxy)ethoxy)-4-((methylsulfonyloxy) methyl)cyclohexyl)-6-oxo-1,6-dihdropyrimidine-4-carboxylate (1.27 g, 1.976 mmol, 99 %) as a tan foam: HPLC: retention time = 3.00 min (area percent = 71%); LCMS: m/z 643 (M+H). To a solution of this compound (1.27 g, 1.98 mmol) in DMF (10 mL) was added cesium 20 carbonate (0.644 g, 1.98 mmol) and the mixture stirred for 6 h at 60°C in an oil bath under N₂. The mixture was concentrated and the residue taken up in ethyl acetate (20 mL), washed with 1M-aq. K₂CO₃, water, followed by brine, then dried (Na₂SO₄), filtered and concentrated to provide 930 mg of a light brown crystalline powder. This was purified by silica gel column chromatography (60 g SiO₂, 5-10% ethyl 25 acetate /CH₂Cl₂) to provide the title compound (382 mg, 0.699 mmol, 35 % yield) as an off-white crystalline solid: HPLC: retention time = 3.24 min (area percent = 94%); TLC: RF 0.6 (10% ethyl acetate/CH₂Cl₂): LCMS: m/z 547 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 1.19 (3H, t, J=7.0 Hz), 1.66-1.79 (2H, m), 1.89-2.01 (2H, m), 2.07-2.16 (2H, m), 2.17-2.27 (2H, m), 2.46 (1H, brs), 4.00 (2H, t, J=5.2 Hz), 4.09

(2H, d, J=3.7 Hz), 4.27 (2H, q, J=7.0 Hz), 4.62 (2H, t, J=5.3 Hz), 7.43 (2H, t, J=7.8 Hz), 7.49 (2H, t, J=7.8 Hz), 7.55 (1H, t, J=7.5 Hz), 7.63 (1H, t, J=7.5 Hz), 8.06 (2H, dd, J=8.4, 1.1 Hz), 8.19 (2H, dd, J=8.4, 1.1 Hz); ^{13}C NMR (126 MHz, CDCl_3) δ ppm 14.0, 23.3, 27.7, 30.9, 54.2, 62.1, 62.2, 64.8, 78.0, 128.4, 128.6, 128.7, 129.8, 130.5, 5 130.7, 133.0, 134.0, 136.6, 140.7, 159.9, 161.2, 162.9, 163.8, 166.7.


Intermediate 138



10


5-Hydroxy-1-(2-hydroxyethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^2.7]trideca-2,4-diene-4-carboxylic acid ethyl ester. At room temperature, to a cold (ice-cooling) and stirred solution of 5-benzoyloxy-1-(2-benzoyloxy-ethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^2.7]trideca-2,4-diene-4-carboxylic acid ethyl ester (330 mg, 0.604 mmol) in THF (6 mL) and EtOH (3.00 mL) was added 21% sodium ethanolate in EtOH (0.676 mL, 1.81 mmol) diluted with EtOH (3 mL) and the mixture stirred under N_2 for 1 h.. The mixture was diluted with ethyl acetate (10 mL) and extracted with water (10 mL x 2). The aqueous extracts were acidified with 1N-HCl (3 mL) and extracted with ethyl acetate (10 mL x 5). The ethyl acetate extracts were washed 15 with brine, dried (Na_2SO_4) and concentrated to provide the title compound (130 mg, 0.38 mmol; 64% yield) as an oil: HPLC: retention time = 1.33 min (area percent = 95%); LCMS: m/z 339 (M+H); ^1H NMR (500 MHz, CDCl_3) δ ppm 1.43 (3H, t, J=7.2 Hz), 1.65 (2H, dt, J=12.3, 6.2 Hz), 1.85-1.97 (2H, m), 1.99-2.17 (4H, m), 2.46 (1H, brs), 3.60 (2H, t, J=4 Hz), 3.75 (2H, d, J=4.1 Hz), 4.06-4.15 (2H, m), 4.45 (2H, q, 20 J=7.0 Hz), 10.48 (1H, brs); ^{13}C NMR (126 MHz, CDCl_3) δ ppm 14.0, 23.2, 27.6, 30.5, 53.9, 62.0, 63.2, 65.3, 76.7, 124.1, 148.2, 154.3, 159.7, 169.1.

Intermediate 139



5 *5-Benzyl-1-(2-hydroxyethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2.7}]trideca-2,4-diene-4-carboxylic acid ethyl ester.* To a solution of 5-hydroxy-1-(2-hydroxyethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2.7}]trideca-2,4-diene-4-carboxylic acid ethyl ester (203 mg, 0.600 mmol) in DMF (1 mL) was added K₂CO₃ (249 mg, 1.80 mmol) and (bromomethyl)benzene (103 mg, 0.600 mmol) and the mixture stirred at room
10 temperature overnight under N₂. Additional (bromomethyl)benzene (206 mg, 1.2 mmol, 2 eq.) and K₂CO₃ (166 mg, 1.2 mmol, 2 eq.) were added and the mixture stirred overnight. The mixture was concentrated *in vacuo* to remove most of DMF, and the residue was diluted with ethyl acetate (15 mL), washed with water followed by brine, then dried (Na₂SO₄) and concentrated to provide an amber oil which was
15 purified by silica gel column chromatography (18gSiO₂, 45% ethyl acetate/CH₂Cl₂) to give the title compound (141 mg, 0.329 mmol, 55 % yield) as an amber oil; HPLC: retention time = 2.18 min (area percent = 92%); LCMS: m/z 429 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 1.28 (3H, t, J=7.2 Hz), 1.63 (2H, ddd, J=14.3, 7.5, 7.2 Hz), 1.83-1.94 (2H, m), 2.06 (4H, t, J=7.6 Hz), 2.42 (1H, brs), 3.59 (2H, t, J=4 Hz), 3.70 (2H, d, J=4 Hz), 4.05 (2H, d, J= 4.0 Hz), 4.31 (2H, q, J=7.2 Hz), 5.22 (2H, s), 7.27 (1H, t, J= 7.4 Hz), 7.32 (2H, t, J= 7.2 Hz), 7.46 (2H, d, J= 6.7 Hz); ¹³C NMR (126 MHz, CDCl₃) δ ppm 14.1, 23.2, 27.6, 30.5, 53.6, 62.1, 62.2, 65.3, 74.5, 76.7, 128.3, 128.5, 128.6, 136.8, 140.0, 142.4, 159.2, 161.3, 163.9.

Intermediate 140




5 *5-Benzyl-1-(2-methoxyethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2,7}]trideca-2,4-diene-4-carboxylic acid methyl ester.* To a solution of 5-benzyl-1-(2-hydroxyethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2,7}]trideca-2,4-diene-4-carboxylic acid ethyl ester (105 mg, 0.245 mmol) in DMF (1 mL) was added 60% NaH (15 mg, 0.37 mmol) and the mixture stirred at room temp. for 15 min.

10 under N₂. After H₂ gas evolution had ceased, MeI (0.031 mL, 0.490 mmol) was added and the mixture stirred at room temperature overnight. The mixture was concentrated and the residue was taken up in ethyl acetate (10 mL), washed with water followed by brine, then dried (Na₂SO₄) and concentrated to provide 85 mg of a crude oil which was purified by C-18 reverse phase column chromatography (YMC

15 AQ, 18-22-26-30-34% CH₃CN/H₂O with 01% trifluoroacetic acid) to provide 8 mg (0.019 mmol, 8 % yield) of the title compound as an amber film: HPLC: retention time = 2.30 min (area percent = 91%); LC/MS: m/z 429 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 1.66-1.70 (2H, m), 1.87-2.02 (2H, m), 2.12 (4H, t, J=7.5 Hz), 2.46 (1H, brs), 3.39 (3H, s), 3.62 (2H, t, J=4.7 Hz), 3.73 (2H, d, J=4.7 Hz), 3.84 (3H, s),

20 4.07 (2H, d, J=3.7 Hz), 5.26 (2H, s), 7.31 (1H, t, J=7.0 Hz), 7.36 (2H, t, J= 7.3 Hz), 7.48 (2H, d, J= 7.6 Hz).

Intermediate 141




25

5-Benzylxy-1-carboxymethoxy-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2,7}]trideca-2,4-diene-4-carboxylic acid ethyl ester. To a solution of 5-benzylxy-1-(2-methoxyethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2,7}]trideca-2,4-diene-4-carboxylic acid methyl ester (180 mg, 0.42 mmol) in DMF (3 mL) was added pyridinium dichromate (237 mg, 0.630 mmol) and the mixture vigorously stirred for 14 days at room temperature under N₂. The solvent was removed *in vacuo* and the residue was stirred in a mixture of ethyl acetate (10 mL) and water for 30 min. The insoluble materials were filtered off (CELITE®), and the organic filtrate was extracted with sat'd aq. NaHCO₃. The aqueous solution was acidified with 1N HCl, and extracted with ethyl acetate (10 mL). The ethyl acetate extract was washed with brine, dried (Na₂SO₄), and concentrated to provide the title compound (27 mg, 0.061 mmol, 15% yield) as a light yellow film. HPLC: retention time = 2.38 min (area percent = 80%); LCMS: m/z 443 (M+H); ¹H NMR (400 MHz, CDCl₃) δ ppm 1.34 (3H, t, J=7.0 Hz), 1.71 (2H, m), 1.97 (2H, m), 2.05-2.21 (4H, m), 2.51 (1H, brs), 4.05 (2H, s), 4.12 (2H, d, J= 2.7 Hz), 4.35 (2H, q, J=7.0 Hz), 5.28 (2H, s), 7.31-7.39 (3H, m), 7.48 (2H, d, J= 7.0 Hz); ¹³C NMR (101 MHz, CDCl₃) δ ppm 13.5, 22.6, 27.0, 30.1, 53.3, 62.1, 63.3, 74.2, 78.5, 128.0, 128.2, 128.7, 135.9, 139.4, 141.7, 158.9, 160.3, 162.5, 170.4.

Intermediate 142

20




5-Benzylxy-1-dimethylcarbamoylmethoxy-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2,7}]trideca-2,4-diene-4-carboxylic acid ethyl ester. To a solution of 5-benzylxy-1-carboxymethoxy-6-oxo-3,7-diaza-tricyclo[7.2.2.0^{2,7}]trideca-2,4-diene-4-carboxylic acid ethyl ester (25 mg, 0.057 mmol) in DMF (1 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (43 mg, 0.11 mmol) and 4-(dimethylamino)pyridine (DMAP) (14 mg, 0.11 mmol) and the mixture stirred at room temperature for 15 min under N₂. To this was added

a 2M THF solution of dimethylamine (0.057 mL, 0.11 mmol) and the mixture stirred at room temperature for 90 min. The mixture was concentrated and the residue taken up in ethyl acetate (10 mL), washed with 1N HCl followed by brine, then dried (Na_2SO_4) and concentrated to provide the title compound (36 mg, 0.057 mmol, 100 % yield) as a crude amber film: HPLC: retention time = 2.15 min; LCMS: m/z 470 ($\text{M}+\text{H}$); ^1H NMR (400 MHz, CDCl_3) δ ppm 1.23 (3H, t, $J=6.6$ Hz), 1.65-1.76 (2H, m), 1.88-2.01 (2H, m), 2.11-2.26 (4H, m), 2.46 (1H, brs), 2.90 (3H, s), 2.95 (3H, s), 4.10 (2H, t, $J=7.0$ Hz), 4.23-4.33 (4H, m), 5.24 (2H, s), 7.27-7.38 (3H, m), 7.46 (2H, d, $J=7.0$ Hz).

10

Intermediate 143



15 *2-Methylsulfanyl-1,6-dihydro-6-oxo-5-(phenylmethoxy)-4-pyrimidinecarboxylic acid, ethyl ester.* A solution of diethyl oxalate (21.06 g, 0.144 mol) and ethyl benzyloxyacetate (28.0 g, 0.144 mol) in dry THF (200 ml) was treated at 22 °C with sodium hydride (6.34 g of a 60 % dispersion in mineral oil, 0.158 mol). Ethanol (0.05 mL) was added and the resulting mixture was stirred at 22 °C for 18 hours. The THF was removed under reduced pressure and the residue was dissolved in a solution of sodium ethoxide (0.072 mol, prepared from 1.65 g of sodium) in ethanol (200 ml). Then powdered 2-methyl-2-thiopseudourea sulfate (20.1 g, 0.072 mol) was added and the resulting mixture was heated at 60 °C for 6 hours. Acetic acid (5 ml) was added and the ethanol was removed under reduced pressure. The residual paste was partitioned between water and dichloromethane and the aqueous phase was extracted twice with dichloromethane. The combined organic phases were washed with saturated sodium bicarbonate followed by brine, dried over anhydrous

20


25

magnesium sulfate, filtered, and concentrated under reduced pressure.

Chromatography on silica gel (elution with a gradient of ethyl acetate 0-20 % in toluene) followed by crystallization from ethyl acetate–hexane gave 8.34 g (18% yield) of the title compound; mp 109–110 °C. ^1H NMR 400 MHz (CDCl_3) δ : 1.35
 5 (3H, t, J = 7.1 Hz, CH_3), 2.62 (3H, s, SCH_3), 4.37 (2H, q, J = 7.1 Hz, CH_2), 5.28 (2H, s, OCH_2), 7.35 – 7.52 (5H, m, aromatics), 12.2 (1H, broad s, NH). Anal. Calcd for $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_4\text{S}$: C 56.23, H 5.03, N 8.74. Found: C 56.23, H 4.86, N 8.76.


Intermediate 144

10



Ethyl 6-(allyloxy)-5-(benzyloxy)-2-(methylthio)pyrimidine-4-carboxylate. To a solution of 2-methylsulfanyl-1,6-dihydro-6-oxo-5-(phenylmethoxy)-4-pyrimidinecarboxylic acid, ethyl ester (4.28 g, 13.36 mmol) in DMF (70 mL) was added Cs_2CO_3 (4.35 g, 13.36 mmol) followed by 3-bromoprop-1-ene (1.156 mL, 13.36 mmol). The resulting mixture was stirred at 80 °C for 1 h, cooled to room temperature and partitioned between ethyl acetate and hexane. The organic phase was washed with water followed by brine, dried (Na_2SO_4) and filtered.
 15 Concentration gave an oil that was purified by flash column chromatography (Biotage flash chromatography system, 0% - 40% ethyl acetate/hexane, 16 CV) to give the title compound as a yellow oil (1.8144 g, 38% yield). ^1H NMR (300 MHz, CDCl_3) δ : 7.41 - 7.37 (2H, m), 7.36 - 7.28 (3H, m), 6.10 - 5.97 (1H, m), 5.40 (2H, dq, J = 17.2, 1.5 Hz), 5.29 (2H, dq, J = 10.5, 1.2 Hz), 5.03 (2H, s), 4.93 (2H, dt, J = 5.6, 1.4 Hz), 4.32 (2H, q, J = 7.1 Hz), 4.32 (2H, q, J = 7.1 Hz), 2.50 (2H, q, J = 7.1 Hz), 1.28 (3H, t, J = 7.1 Hz). LCMS (M+H) calcd for $\text{C}_{18}\text{H}_{21}\text{N}_2\text{O}_4\text{S}$: 361.12; found: 361.21.


Intermediate 145



5 *Ethyl 6-(allyloxy)-5-(benzyloxy)-2-(methylsulfonyl)pyrimidine-4-carboxylate.*

To a solution of ethyl 6-(allyloxy)-5-(benzyloxy)-2-(methylthio)pyrimidine-4-carboxylate (1.813 g, 5.03 mmol) in CH₂Cl₂ (100 mL) was added 3-chloroperbenzoic acid (MCPBA) (2.59 g, 11.57 mmol). The resulting mixture was stirred at room temperature for 3 h. The mixture was washed with aqueous sodium bisulfite followed by saturated, aqueous sodium bicarbonate. The organic phase was dried (Na₂SO₄), filtered and concentrated affording the title compound as a colorless oil. (1.9127 g, 97% yield). ¹H NMR (300 MHz, CDCl₃) δ: 7.40 - 7.31 (5H, m), 6.13 - 6.00 (1H, m), 5.51 - 5.54 (1H, m), 5.39 - 5.34 (1H, m), 5.20 (2H, s), 5.08 - 5.05 (2H, m), 4.34 (2H, q, *J* = 7.2 Hz), 3.30 (3H, s), 1.30 (3H, t, *J* = 7.1 Hz). LCMS (M+H) calcd for C₁₈H₂₁N₂O₆S: 393.11; found: 393.19.


Intermediate 146



Ethyl 6-(allyloxy)-5-(benzyloxy)-2-(4-(hydroxymethyl)piperidin-1-yl)pyrimidine-4-carboxylate. A solution of ethyl 6-(allyloxy)-5-(benzyloxy)-2-(methylsulfonyl)pyrimidine-4-carboxylate (1.911 g, 4.87 mmol) in THF (40 mL) was stirred with molecular sieves for 30 min. To this mixture was added piperidin-4-ylmethanol (1.683 g, 14.61 mmol) and the resulting mixture was stirred overnight at 70 °C. The mixture was cooled to room temperature and diluted with ethyl acetate and washed with water followed by brine then dried (Na_2SO_4). Filtration followed by concentration gave a yellow oil that was purified by flash column chromatography (Biotage flash chromatography system; 0% - 100%, 10 CV) to afford the title compound as a yellow oil (1.2043 g, 58% yield). ^1H NMR (300 MHz, CDCl_3) δ : 7.42 - 7.39 (2H, m), 7.35 - 7.27 (3H, m), 6.10 - 5.97 (1H, m), 5.42 - 5.34 (1H, m), 5.28 - 5.22 (1H, m), 4.91 (2H, s), 4.87 - 4.84 (2H, m), 4.72 - 4.63 (2H, m), 4.30 (2H, q, J = 7.1 Hz), 3.50 (2H, t, J = 5.8 Hz), 2.80 (2H, td, J = 12.8, 1.8 Hz), 1.79 - 1.64 (3H, m), 1.28 (3H, t, J = 7.1 Hz), 1.23 - 1.10 (2H, m).

15

Intermediate 147




20 *Ethyl 5-(benzyloxy)-2-(4-(hydroxymethyl)piperidin-1-yl)-6-oxo-1,6-dihydropyrimidine-4-carboxylate.* A solution of ethyl 6-(allyloxy)-5-(benzyloxy)-2-(4-(hydroxymethyl)piperidin-1-yl)pyrimidine-4-carboxylate (1.206 g, 2.82 mmol) in n-propanol (10 mL) was sparged with N_2 . Rhodium(III) chloride (0.030 g, 0.141 mmol) was added and the resulting mixture was stirred overnight at 100 °C. The 25 mixture was cooled to room temperature and partitioned between water and CH_2Cl_2 .

The organic phase was dried (Na_2SO_4), filtered and concentrated to give a yellow solid that was triturated with Et_2O . The product was collected by filtration to give the title compound as a white solid (0.8252 g, 76% yield). ^1H NMR (300 MHz, CDCl_3) δ : 7.43 (2H, d, J = 7.3 Hz), 7.37 - 7.29 (3H, m), 5.05 (2H, s), 4.46 (2H, d, J = 13.1 Hz), 4.31 (2H, q, J = 7.1 Hz), 3.44 (2H, d, J = 5.8 Hz), 2.94 - 2.88 (2H, m), 1.74 (2H, d, J = 13.1 Hz), 1.73 - 1.66 (1H, m), 1.29 (3H, t, J = 7.2 Hz), 1.28 - 1.22 (2H, m). LCMS (M+H) calcd for $\text{C}_{20}\text{H}_{26}\text{N}_3\text{O}_5$: 388.18; found: 388.23.

Intermediate 148

10




Ethyl 5-(benzyloxy)-1-(methylsulfonyl)-2-(4-((methylsulfonyloxy)methyl)piperidin-1-yl)-6-oxo-1,6-dihdropyrimidine-4-carboxylate. To a solution of ethyl 5-(benzyloxy)-2-(4-(hydroxymethyl)piperidin-1-yl)-6-oxo-1,6-dihdropyrimidine-4-carboxylate (0.814 g, 2.1 mmol) in THF (20 mL) was added methanesulfonyl chloride (0.327 mL, 4.20 mmol) followed by Et_3N (0.732 mL, 5.25 mmol). The resulting mixture was stirred at room temperature for 2.5 h (white solids precipitated out over the reaction time). The mixture was diluted with ethyl acetate (50 mL) and washed with water followed by brine. The organic phase was dried (Na_2SO_4), filtered and concentrated to give the title compound as an amber oil (1.1320g, 99% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.44 (2H, d, J = 7.32 Hz), 7.39 - 7.32 (3H, m), 4.98 (2H, s), 4.67 - 4.64 (2H, m), 4.35 (2H, q, J = 7.1 Hz), 4.09 (2H, d, J = 6.4 Hz), 3.46 (3H, s), 3.02 (3H, s), 2.91 (2H, td, J = 12.9, 2.4 Hz), 2.08 -

2.00 (1H, m), 1.86 - 1.84 (2H, m), 1.35 - 1.28 (5H, m). LCMS (M+H) calcd for C₂₂H₃₀N₃O₉S₂: 544.14; found: 544.30.


Intermediate 149

5




Ethyl 3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylate. A mixture of ethyl 5-(benzyloxy)-1-(methylsulfonyl)-2-(4-((methylsulfonyloxy)methyl)piperidin-1-yl)-6-oxo-1,6-dihdropyrimidine-4-carboxylate and K₂CO₃ (0.051 g, 0.368 mmol) in ethanol (10 mL) was stirred at 60 °C for 4 h and cooled to room temperature. Water was added and the product was extracted with ethyl acetate. The organic phase was washed with brine then dried (Na₂SO₄), filtered and concentrated. The residue was purified by flash column chromatography (Biotage flash chromatography system; 0% - 10% methanol/CH₂Cl₂, 25 CV; 10% methanol/CH₂Cl₂, 5 CV) to give the title compound as a white glass (0.0248 g, 37% yield). ¹H NMR (500 MHz, CDCl₃) δ: 7.44 (2H, dd, J = 7.3 Hz), 7.30 - 7.22 (3H, m), 5.27 (2H, s), 4.22 (2H, q, J = 7.1 Hz), 4.01 - 3.94 (2H, m), 3.77 - 3.70 (2H, m), 3.67 (2H, brs), 3.02 (1H, brs), 2.29 - 2.24 (2H, m), 1.93 - 1.88 (2H, m), 1.22 (3H, t, J = 7.0 Hz). LCMS (M+H) calcd for C₂₀H₂₄N₃O₄: 370.17; found: 370.30.

Intermediate 150



5 *3-(Benzylxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylic acid.* A mixture of ethyl 3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylate (0.5491 g, 1.486 mmol) and LiOH.H₂O (0.071 g, 2.97 mmol) in ethanol (12 mL) and water (3.00 mL) was stirred at room temperature for 6 h. The mixture was acidified with concentrated HCl and the product was extracted with ethyl acetate. The organic phase was washed with brine, then dried (Na₂SO₄), filtered and concentrated to give the title compound as a white solid (0.4521 g, 89% yield). ¹H NMR (500 MHz, CD₃OD) δ: 7.53 - 7.49 (2H, m), 7.42 - 7.36 (3H, m), 5.27 (2H, d, *J* = 15.3 Hz), 4.11 - 4.00 (4H, m), 3.96 (2H, d, *J* = 14.0 Hz), 3.17 - 3.13 (1H, m), 2.45 - 2.39 (2H, m), 15 2.10 - 2.03 (2H, m). LCMS (M+H) calcd for C₁₈H₂₀N₃O₄: 342.14; found: 342.20.


Intermediate 151



3-(Benzylloxy)-N-(4-fluoro-2-(4,4,5-trimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)benzyl)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxamide. A mixture of 3-(benzylloxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylic acid (0.03 g, 5 0.088 mmol), 1-(4-fluoro-2-(4,4,5-trimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)phenyl)methanamine hydrochloride salt (0.043 g, 0.132 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.100 g, 0.264 mmol), 4-(dimethylamino)pyridine (DMAP) (1.074 mg, 8.79 µmol) and diisopropyl ethylamine (0.077 mL, 0.439 mmol) in DMF (2 mL) was stirred at 10 room temperature for 1 h. The mixture was diluted with ethyl acetate and washed with 0.5 N HCl, water followed by brine. The organic phase was dried (Na_2SO_4), filtered and concentrated to give the title compound as a crude yellow oil. LCMS (M+H) calcd for $\text{C}_{30}\text{H}_{36}\text{FN}_6\text{O}_5\text{S}$: 611.24; found: 611.33.


15

Intermediate 152



20 3-(Benzylloxy)-N-(4-fluorobenzyl)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxamide. Formed from 3-(benzylloxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylic acid and 4-fluorobenzylamine to give the title compound as a yellow residue LCMS (M+H) calcd for $\text{C}_{25}\text{H}_{26}\text{FN}_4\text{O}_3$: 449.19; found: 449.25.


Intermediate 153



5 *N*-Benzyl-3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxamide. Formed from 3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxylic acid and benzylamine to give the title compound as a yellow residue. LCMS (M+H) calcd for C₂₅H₂₇N₄O₃: 431.20; found: 431.30.


10

Intermediate 154




15 3-(Benzyl)-*N*-(4-fluoro-2-(3-(2-methoxyethyl)-2-oxo-1-imidazolidinyl)benzyl)-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxamide. Formed from 3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxylic acid and 1-(2-aminomethyl)-5-fluorophenyl-3-(2-methoxyethyl)imidazolidin-2-one to give the title compound as a yellow residue. LCMS (M+H) calcd for C₃₁H₃₆FN₆O₅: 591.27; found: 591.34.

Intermediate 155




5 *3-(Benzylxy)-N-(4-fluoro-2-(5-methyl-1H-1,2,4-triazol-1-yl)benzyl)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxamide.*
 Formed from 3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylic acid and (4-fluoro-2-(5-methyl-1H-1,2,4-triazol-1-yl)phenyl)methanamine to give the title compound as a white glass. LCMS (M+H)
 10 calcd for C₂₈H₂₉FN₇O₃: 530.23; found: 530.24.

Intermediate 156



15 *3-(Benzylxy)-N-(4-fluoro-2-(5-methyl-1H-1,2,3-triazol-1-yl)benzyl)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxamide.*
 Formed from 3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylic acid and (4-fluoro-2-(5-methyl-1H-1,2,3-triazol-1-yl)phenyl)methanamine to give the title compound as a yellow residue (5.5 mg, 15% yield). LCMS (M+H) calcd for C₂₈H₂₉FN₇O₃: 530.23; found: 530.17.

Intermediate 157




5 *3-(Benzylxy)-N-(4-fluoro-2-(3-methyl-2-oxo-1-imidazolidinyl)benzyl)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxamide.*
 Formed from 3-(benzyloxy)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxylic acid and 1-(2-(aminomethyl)-5-fluorophenyl)-3-methylimidazolidin-2-one to give the title compound as a yellow residue (4.7 mg,
 10 13% yield). LCMS (M+H) calcd for C₂₉H₃₂FN₆O₄: 547.24; found: 547.20.

Example 1 through can be synthesized according to Scheme III

Example 1


15



20 *7,11-Epoxy-4H-pyrimido[1,2-a]azocene-2-carboxamide, N-[(4-fluorophenyl)methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (7S,11R)-.* To ethyl (7S,11R)-3-hydroxy-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,11-epoxypyrimido[1,2-a]azocene-2-carboxylate (0.050mg, 0.18 mmol) in EtOH (10 mL) was added 4-fluorobenzylamine (0.081 g, 0.71 mmol). The reaction mixture was stirred at reflux for 4 hours in a sealed tube. The solvent was removed *in vacuo* and the product purified by preparative HPLC (C18 column) to afford 0.029g (45% yield)

of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 12.05 (1 H, s), 7.86 (1 H, m), 7.31 - 7.38 (2 H, m), 7.04 - 7.16 (2 H, m), 4.77 (1 H, d, $J=2.3$ Hz), 4.53-4.65 (3 H, m), 4.01 - 4.10 (2 H, m), 2.03 - 2.14 (2 H, m), 1.85 (1 H, dd, $J=13.6, 1.8$ Hz), 1.76 (2 H, d, $J=2.0$ Hz), 1.31 - 1.43 (1 H, m).). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 360. . HRMS (ESI $^+$) calculated for $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}_5$ [$\text{M}+\text{H}^+$] :360.1360; found : 360.1351.


Example 2



10


7,11-Epoxy-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[[4-fluoro-2-(1H-1,2,4-triazol-1-yl)phenyl]methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (7S,11R)-. To ethyl (7S,11R)-3-hydroxy-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,11-epoxypyrimido[1,2-a]azocine-2-carboxylate (0.050, 0.18 mmol) in EtOH (5 mL) was added (4-fluoro-2-(1H-1,2,4-triazol-1-yl)phenyl)methanamine hydrochloride (0.081 mg, 0.36 mmol) and diisopropylethylamine (0.063 mL, 0.36 mmol). The reaction mixture was heated in a microwave oven at 160 °C for 3h. LC-MS showed ~50% conversion to the target compound. The reaction mixture was stirred at 160 °C for an additional 3h. EtOH was removed *in vacuo* and the crude material was purified by preparative HPLC using a C18 column and trifluoroacetic acid in the mobile phase. Solvent from the purification was removed and CH_3CN (5 mL) and H_2O (1 mL) were added and the resulting solution frozen and lyophilized overnight (16h) to afford 21 mg (28% yield) of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 11.95 (1H, br s), 8.56 (1 H, s), 8.47 (1 H, t, $J=6.2$ Hz), 8.28 (1 H, s), 7.62 - 7.74 (1 H, m), 7.21 - 7.29 (1 H, m), 7.12 - 7.19 (1 H, m), 4.80 (1H, d, $J=2.3$ Hz), 4.47 - 4.56 (3 H, m), 3.98 - 4.10 (2 H, m), 2.06 - 2.18 (2 H, m), 1.86 (1 H, dd, $J=13.6, 1.8$ Hz), 1.75 (2 H, d, $J=1.5$ Hz), 1.29 - 1.41 (1 H, m).). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 427. HRMS (ESI $^+$) calculated for $\text{C}_{20}\text{H}_{19}\text{FN}_6\text{O}_4$ [$\text{M}+\text{H}^+$] :427.1530; found :427.1530.

Example 3




5 *7,11-Epoxy-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[{[4-fluoro-2-[(methylamino)carbonyl]phenyl}methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (7S,11R)-.* ^1H NMR (400 MHz, CDCl_3) δ ppm 8.55 (1 H, s), 7.38-7.45 (1 H, m),
7.24 - 7.31 (2 H, m), 4.72 (1 H, d, $J=2.3$ Hz), 4.47 - 4.54 (3 H, m), 3.83 – 3.98 (2 H, m),
2.78 (2 H, d, $J=4.6$ Hz), 1.89-1.99 (2 H, m), 1.73 (2 H,m), 1.49-1.54 (1 H, m),
10 1.34-1.17 (1 H, m). LCMS ($^+\text{ESI, M+H}^+$) m/z 417. HRMS (ESI $^+$) calculated for
 $\text{C}_{20}\text{H}_{21}\text{FN}_4\text{O}_5$ [M+H $^-$]: 417.1574; found: 417.1575.

Example 4



15 *7,11-Epoxy-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[{[4-fluoro-2-(5-methyl-1H-1,2,4-triazol-1-yl)phenyl}methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (7S,11R)-.* ^1H NMR (400 MHz, DMSO-D6) δ ppm 12.04 (1 H, s), 8.09 (1 H, s),
7.50 - 7.60 (2 H, m), 7.45 (1 H, td, $J=8.5, 2.4$ Hz), 4.70 (1 H, s), 4.47 (1 H, s), 4.11 -
4.22 (2 H, m), 3.81 - 3.92 (2 H, m), 2.33 (3 H, s), 1.85 - 1.96 (2 H, m), 1.68 - 1.80 (2 H, m), 1.54 (1 H, br d), 1.27 - 1.39 (1 H, m). LCMS ($^+\text{ESI, M+H}^+$) m/z 441.
HRMS (ESI $^+$) calculated for $\text{C}_{28}\text{H}_{27}\text{FN}_4\text{O}_6$ [M+H $^+$]: 441.1687; found: 441.1688.

Example 5




5 *7,11-Epoxy-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[{[4-fluoro-2-(2-oxo-3-oxazolidinyl)phenyl]methyl}-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (7S,11R)-.* ^1H NMR (400 MHz, CDCl_3) δ ppm 12.07 (1 H, s), 8.41 (1 H, s), 7.55 (1 H, dd, $J=8.5, 6.2$ Hz), 7.09 (1 H, td, $J=8.1, 2.4$ Hz), 7.01 (1 H, dd, $J=9.1, 2.5$ Hz), 4.80 (1 H, s), 4.56 - 4.63 (3 H, m), 4.01 - 4.11 (4 H, m), 2.01 - 2.13 (2 H, m), 1.90 (1 H, m), 1.59 - 1.75 (3 H, m), 1.25 - 1.37 (2 H, m). LCMS ($^+\text{ESI, M+H}^+$) m/z 445. HRMS (ESI^+) calculated for $\text{C}_{21}\text{H}_{21}\text{FN}_4\text{O}_6$ [M+H^+]: 445.1523; found: 445.1530.

Examples 6 and 7 can be synthesized according to Scheme IV

15

Example 6




20 *7,11-Methano-4H-pyrimido[1,2-a][1,5]diazocine-9(6H)-carboxylic acid, 2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,10,11-tetrahydro-3-hydroxy-4-oxo-, phenylmethyl ester.* To 9-benzyl 2-ethyl 3-hydroxy-4-oxo-7,8,10,11-tetrahydro-4H-

7,11-methanopyrimido[1,2-a][1,5]diazocene-2,9(6H)-dicarboxylate (0.035 g, 0.085 mmol) in ethanol (2.0 mL) was added 4-fluorobenzylamine (0.030 mL, 0.254 mmol) and triethylamine (0.035 mL, 0.254 mmol). The reaction mixture was stirred at reflux for 65 hours. HCl (1M, 20 mL) was added and the organic material extracted with ethyl acetate (3x 10 mL), and the combined organic fractions dried (MgSO_4) then concentrated *in vacuo*. The crude material was purified by preparative HPLC (C18 column) to afford 40 mg (76% yield) of the title compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 11.95 (1 H, br s), 7.83 (1 H, br s), 7.27-7.38 (5 H, m), 7.02-7.12 (4 H, m), 5.96 - 5.10 (2 H, m), 4.61 - 4.70 (1 H, m), 4.49 (2 H, d, $J=6.3$ Hz), 4.33 (1 H, s), 4.17 (1 H, d, $J=15.4$ Hz), 3.88 - 3.98 (1 H, m), 3.11 (2 H, s), 2.48 (1 H, s), 1.62 (3 H, br s). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 493. HRMS (ESI $^+$) calculated for $\text{C}_{26}\text{H}_{25}\text{FN}_4\text{O}_5$ [$\text{M}+\text{H}^-$]: 493.1887; found: 493.1870.

Example 7

15




7,11-Methano-4H-pyrimido[1,2-a][1,5]diazocene-2-carboxamide, N-[(4-fluorophenyl)methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-. To 7,11-methano-4H-pyrimido[1,2-a][1,5]diazocene-9(6H)-carboxylic acid, 2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,10,11-tetrahydro-3-hydroxy-4-oxo-, phenylmethyl ester (0.029 g, 0.059 mmol) dissolved in acetic acid (5 mL) was added Pd/C (10%, 0.025 g). The reaction mixture was shaken at 23 °C under 40 psi of hydrogen for 16 hours. Pd/C was removed by filtration on CELITE®. The acetic acid was removed *in vacuo* and remaining traces of acetic acid removed by azeotroping with toluene (2 x 25 mL). The crude material was purified by preparative HPLC (C18 column) to afford 0.0055 g (28% yield) of the title compound as a trifluoroacetic acid salt. ^1H NMR (400 MHz, MeOD) δ ppm 9.17 (1H,

br s), 7.40 (2 H, dd, $J=8.6, 5.6$ Hz), 7.06 (2 H, t, $J=8.8$ Hz), 4.53 (2 H, s), 4.33 - 4.44 (1 H, m), 4.05 (1 H, dd, $J=15.4, 6.3$ Hz), 3.54 - 3.66 (2 H, m), 3.41 - 3.52 (3 H, m), 2.81 (1 H, br s), 2.19 - 2.30 (2 H, m). LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z 359. HRMS (ESI $^+$) calculated for $\text{C}_{18}\text{H}_{19}\text{FN}_4\text{O}_3$ [$\text{M}+\text{H}^+$]: 359.1519 found: 359.1520.

5

Examples 8 and 9 can be synthesized according to Scheme V


Example 8



10

Carbamic acid, [2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-7,10-methano-4H-pyrimido[1,2-a]azocin-11-yl]methyl-phenylmethyl ester. To ethyl 3-(benzoyloxy)-11-((benzoyloxy)carbonyl)(methyl)amino)-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocene-2-carboxylate (0.026g, 0.048 mmol) in EtOH : DMF (3mL:1mL) was added p-fluorobenzylamine (0.018 g, 0.143 mmol). The reaction mixture was stirred at 85°C for 18 hours. HCl 1N (50 mL) was then added and the organic material extracted with ethyl acetate (3x 50 mL). The combined organic fractions were dried (MgSO_4) and concentrated *in vacuo*. The crude compound was purified by preparative HPLC (C18 column) to afford 0.012g (60% yield)of the target compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 7.50 - 7.65 (2 H, m), 7.16 - 7.37 (5 H,m), 6.98 - 7.09 (2 H, m), 5.40 - 5.00 (2 H, m), 4.31-4.62 (2 H, m), 2.92-2.98 (3 H, m), 2.62 - 2.69 (3 H, m), - 1.92 - 2.45 (2 H, m), 1.55 - 1.86 (3 H, m), 1.21-1.48 (3 H, m). LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z 521.

Example 9



5 *7,10-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[{(4-fluorophenyl)methyl}-6,7,8,9,10,11-hexahydro-3-hydroxy-11-(methylamino)-4-oxo-.*

To carbamic acid, [2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-7,10-methano-4H-pyrimido[1,2-a]azocin-11-yl]methyl-, phenylmethyl ester (0.068 g, 0.131 mmol) in ethyl acetate (2 mL) was added Pd/C (10%, 0.010g) and the reaction mixture was stirred at 23 °C for 3 hours under H₂.


10 The Pd/C was then removed by filtration and the solvent was evaporated *in vacuo* to afford, after crystallization in hot ethanol (2 mL), 0.004 g (8% yield) of the title compound. ¹H NMR (400 MHz, MeOD) δ ppm 7.36 (2 H, m), 7.05 (2 H, t, *J*=8.2 Hz), 4.51 - 4.62 (2 H, m), 2.25 - 2.93 (6 H, m), 1.60 - 2.06 (5 H, m), 1.10 - 1.45 (3 H, m). LCMS (ESI, M+H⁺) m/z 387.

15 LCMS (ESI, M+H⁺) m/z 387.

Examples 10 through 17 can be synthesized according to Scheme VIII

Example 10


20



Benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-7,8-dihydro-4H-6,9-ethanopyrido[1,2-a]pyrimidin-9(6H)-yl)carbamate. A mixture of 6,9-ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxylic acid, 6,7,8,9-tetrahydro-3-hydroxy-4-oxo-9-[(phenylmethoxy)carbonyl]amino-, ethyl ester (500 mg, 1.21 mmol) and 4-fluorobenzylamine (0.41 mL, 3.63 mmol) in EtOH (10 mL) was heated at 90 °C for 20 h. The reaction mixture was then cooled, concentrated and purified on a C-18 column using methanol/H₂O (containing 0.1% trifluoroacetic acid) as eluent to afford the title compound (295 mg, 45% yield) as a yellow solid. ¹H NMR (500 MHz, DMSO-D₆) δ: 10.25 (s, 1H), 7.48 (s, 1H), 7.34-7.40 (m, 7H), 7.15-7.21 (m, 3H), 5.08 (s, 2H), 4.51 (s, 2H), 3.85 (s, 1H), 2.84-2.91 (m, 2H), 1.92-2.00 (m, 2H), 1.65-1.73 (m, 2H), 1.41-1.49 (m, 2H). LCMS (M+H) = 493.24.


Example 11

15



6,9-Ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, 9-amino-N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-. To a mixture of benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-7,8-dihydro-4H-6,9-ethanopyrido[1,2-a]pyrimidin-9(6H)-yl)carbamate (135 mg, 0.27 mmol) in EtOH (5 mL) was added 10% Pd/C (50 mg) and the mixture was stirred under 1 atm of hydrogen for 18 h. The mixture was then filtered through a pad of CELITE® and the pad was washed with ethyl acetate. The filtrate was then concentrated to afford the title compound (80 mg, 82% yield) as a light yellow solid. ¹H NMR (500 MHz, CDCl₃) δ: 10.15 (s, 1H), 7.99 (brs, 1H), 7.32 (dd, 2H, J = 8.54, 5.19 Hz), 7.04 (t, 2H, J = 8.7 Hz), 5.33 (s, 1H), 4.58 (d, 2H, J = 6.41 Hz), 1.95-2.00 (m, 4H), 1.78-1.82 (m, 2H), 1.54-1.58 (m, 2H). LCMS (M+H) = 359.18.


Example 12



5 *6,9-Ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, 9-(acetylamino)-N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-*. To a stirred solution of 6,9-ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, 9-amino-N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo- (20 mg, 0.056 mmol) in CH₂Cl₂ (2 mL) was added triethylamine (0.1 mL) followed by acetyl chloride (0.1 mL). After 16 h at room temperature, the reaction mixture was concentrated to give a residue which was dissolved in methanol (2mL) and treated with 2M dimethylamine/methanol (0.5 mL). The resulting reaction mixture was stirred at 60 °C for 2 h, cooled and purified by preparative HPLC to afford the title compound as a white solid (10 mg, 45% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.30 (s, 1H), 7.61 (brs, 1H), 7.34 (dd, 2H, J = 8.7 and 5.34 Hz), 7.07 (t, 2H, J = 8.69 Hz), 6.79 (s, 1H), 5.34 (s, 1H), 4.62 (d, 2H, J = 6.1 Hz), 3.08-3.14 (m, 2H), 2.03 (s, 3H), 1.98-2.02 (m, 2H), 1.71-1.82 (m, 2H), 1.49-1.55 (m, 2H). LCMS (M+H) = 401.22. HPLC purity: > 95%.


20

Example 13



6,9-ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-9-[(methylsulfonyl)amino]-4-oxo-. White solid, ^1H NMR (500 MHz, CDCl_3) δ : 12.41 (s, 1H), 8.17 (s, 1H), 7.35 (dd, 2H, J = 8.7 and 5.34 Hz), 7.03 (t, 2H, J = 8.54 Hz), 5.78 (s, 1H), 5.34 (s, 1H), 5 4.61 (d, 2H, J = 6.41 Hz), 3.07 (s, 3H), 2.58-2.62 (m, 2H), 2.0-2.05 (m, 2H), 1.81-1.87 (m, 4H). LCMS ($M+\text{H}$) = 436.97. HPLC purity: > 95%.

Example 14




10

6,9-Ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, 9-[[[(dimethylamino)acetyl]amino]-N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-. Off-white solid (19% yield) ^1H NMR (500 MHz, CDCl_3) δ : 12.31 (s, 1H), 8.65 (s, 1H), 7.35 (dd, 2H, J = 8.55 and 5.49 Hz), 7.0 (, 2H, J = 8.69 Hz), 5.30 (s, 1H), 4.58 (d, 2H, J = 6.1 Hz), 3.80 (s, 1H), 2.93-3.10 (m, 2H), 2.89 (s, 6H), 1.96-2.01 (m, 2H), 1.80-1.86 (m, 2H), 1.52-1.57 (m, 2H). LCMS ($M+\text{H}$) = 444.29. HPLC purity: > 95%.

20


Example 15



6,9-Ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, 9-[[[3-(1,1-dimethylethyl)-1-methyl-1*H*-pyrazol-5-yl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-. Off-white solid (14% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.35 (s, 1H), 7.55 (s, 1H), 7.41 (s, 1H), 7.28-5 7.31 (m, 2H), 7.03 (t, 2H, $J = 7.93$ Hz), 6.23 (s, 1H), 5.38 (s, 1H), 4.61 (s, 2H), 4.09 (s, 3H), 3.11-3.16 (m, 2H), 2.04-2.09 (m, 2H), 1.84-1.90 (m, 2H), 1.64-1.69 (m, 2H), 1.22 (s, 9H). LCMS ($\text{M}+\text{H}$) = 523.42. HPLC purity: > 95%.

Example 16

10




Ethanediamide, N' -[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8-dihydro-3-hydroxy-4-oxo-6,9-ethano-4H-pyrido[1,2-a]pyrimidin-9(6*H*)-yl]-*N,N*-dimethyl-. To a stirred solution of *N,N*-dimethyloxamic acid (26 mg, 0.22 mmol) and *N*-methylmorpholine (0.05 mL) in CH_2Cl_2 (3 mL) was added isopropyl chloroformate (0.22 mL, 0.22 mmol, 1M in toluene). After 2 h at room temperature, 6,9-ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, 9-amino-*N*-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo- (20 mg, 0.056 mmol) in 15 CH_2Cl_2 (1 mL) was added in one portion and the mixture stirred at room temperature. After 16 h at room temperature, the reaction mixture was concentrated to give crude material which was dissolved in methanol (2mL) and treated with 2M dimethylamine/methanol (0.5 mL). The resulting reaction mixture stirred at 60 °C for 2 h, cooled and purified by preparative HPLC to afford the title compound as a white solid (9 mg, 35% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.31 (s, 1H), 8.73 (s, 1H), 8.02 (s, 1H), 7.36 (dd, 2H, $J = 8.39$ and 5.34 Hz), 7.05 (t, 2H, $J = 8.55$ Hz), 5.34 (s, 1H), 4.60 (d, 2H, $J = 6.1$ Hz), 3.38 (s, 3H), 3.01 (s, 3H), 2.98-3.04 (m, 2H), 2.01-20 2.01 (s, 6H).

2.06 (m, 2H), 1.82-1.87 (m, 2H), 1.59- 1.64 (m, 2H). LCMS (M+H) = 458.15.
HPLC purity: > 95%.

Example 17

5




6,9-Ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-9-[[5-methyl-1,3,4-oxadiazol-2-yl]carbonyl]amino]-4-oxo-. To a stirred solution of 6,9-ethano-4H-pyrido[1,2-a]pyrimidine-2-carboxamide, 9-amino-N-[(4-fluorophenyl)methyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo- (20 mg, 0.056 mmol) and triethylamine (0.1 mL) in CH₂Cl₂ (3 mL) was added a freshly prepared solution of 5-methyl-1,3,4-oxadiazole-2-carbonyl chloride (0.56 mmol) in CH₂Cl₂ (2 mL) at room temperature. After 16 h at room temperature, the reaction mixture was concentrated to give crude material which was dissolved in methanol (2mL) and treated with 2M dimethylamine/methanol (0.5 mL). The resulting reaction mixture was stirred at 60 °C for 2 h, then cooled and purified by preparative HPLC to afford the title compound as a white solid (6 mg, 23% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.34 (s, 1H), 8.77 (s, 1H), 7.89 (s, 1H), 7.41 (dd, 2H, J = 8.54 and 5.19 Hz), 7.11 (t, 2H, J = 8.7 Hz), 5.36 (s, 1H), 4.64 (d, 2H), 3.12-3.18 (m, 2H), 2.65 (s, 3H), 2.04-2.08 (m, 2H), 1.85-1.91 (m, 2H), 1.61-1.68 (m, 2H). LCMS (M+H) = 469.10. HPLC purity: > 95%.

Examples 18 through 56 can be synthesized according to Schemes IX through XI

Example 18


5



- Benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)carbamate.* A mixture of ethyl 10-(((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (5.00 g, 11.70 mmol), (4-fluorophenyl)methanamine (2.67 mL, 23.39 mmol) and Et₃N (3.26 mL, 23.39 mmol) in ethanol (78 mL) was stirred at 90 °C for 16 h, cooled and concentrated. The resulting pale yellow foam was dissolved in ethyl acetate and washed with 1N HCl. The organic phase was washed with brine, dried (Na₂SO₄), filtered and concentrated.
- 10 The residue was taken up in hot ethyl acetate and hexane was added until the solution became slightly cloudy. After cooling, the title compound was collected by filtration as a white solid (4.8203 g, 81 % yield). ¹H NMR (500 MHz, DMSO-D6) δ: 10.25 (s, 1H), 7.52 (s, 1H), 7.34-7.40 (m, 8H), 7.15-7.21 (m, 2H), 5.72 (s, 1H), 4.91 (s, 2H), 4.40 (d, 2H, J = 6.4 Hz), 4.15 (s, 2H), 2.48-2.52 (m, 2H), 2.02-2.08 (m, 2H), 1.94-20 1.98 (m, 2H), 1.63-1.70 (m, 2H). LCMS (M+H) = 507.42.

Example 19


25



10-Amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a mixture of benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)carbamate (300 mg, 0.59 mmol) in methanol (3 mL) was added 10% Pd/C (315 mg) and the mixture was stirred under balloon hydrogen atmosphere for 18 h. The mixture was then filtered through a pad of CELITE® and the pad washed with ethyl acetate. The filtrate was concentrated to afford the title compound (160 mg, 72% yield) as a light yellow solid. ¹H NMR (500 MHz, DMSO-d₆) δ: 12.22 (s, 1H), 9.50 (brs, 1H), 7.35-7.41 (m, 2H), 7.19-7.22 (m, 2H), 4.54 (d, 2H, J = 5.8 Hz), 4.01 (s, 2H), 2.42-2.48 (m, 3H), 2.08 (s, 2H), 1.98-2.03 (m, 2H), 1.84-1.90 (m, 2H), 1.70-1.78 (m, 2H). LCMS (M+H) = 373.28.

Example 20

15




10-(Acetylamino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a stirred solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (25 mg, 0.067 mmol) in CH₂Cl₂ (2 mL) was added triethylamine (0.05 mL) followed by acetyl chloride (0.05 mL). After 16 h at room temperature the reaction mixture was concentrated to give crude product which was dissolved in methanol (2mL) and treated with 2M dimethylamine/methanol (0.5 mL). The resulting reaction mixture stirred at 60 °C for 2 h, cooled and purified by preparative HPLC to afford the title compound as a white solid (3 mg, 11% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.00 (s, 1H), 7.32-7.35 (m, 3H), 7.04-7.09 (m, 2H), 6.69 (s, 1H), 4.57 (d, 2H, J = 5.8 Hz), 4.14 (d, 2H,

J = 3.66 Hz), 2.63-2.66 (m, 2H), 2.44-2.47 (m, 1H), 1.91-1.97 (m, 4H), 1.86 (s, 3H), 1.63-1.67 (m, 2H). LCMS (M+H) = 415.29. HPLC purity: > 95%.

Example 21


5



N-(4-Fluorobenzyl)-3-hydroxy-10-((methylsulfonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (11 mg, 36 % yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.99 (s, 1H), 8.85 (s, 1H), 7.34 (dd, 2H, J = 8.54 & 5.19 Hz), 6.98-7.04 (m, 2H), 4.84 (s, 1H), 4.53 (d, 2H, J = 6.41 Hz), 4.15 (d, 2H, J = 3.97 Hz), 3.04 (s, 3H), 2.51 (d, 1H, J = 3.05 Hz), 2.07-2.24 (m, 4H), 1.87-1.98 (m, 2H), 1.65-1.75 (m, 2H). LCMS (M+H) = 451.2. HPLC purity: > 95%.

15

Example 22




20 *N*-(4-Fluorobenzyl)-3-hydroxy-10-((4-morpholinylcarbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (10 mg, 31% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.00 (s, 1H), 8.70 (s, 1H), 7.37 (s, 1H), 7.31 (dd, 2H, J = 8.39 Hz & 5.34 Hz), 7.08 (t, 2H, J = 8.55 Hz),

4.58 (d, 2H, $J = 5.80$ Hz), 4.18 (d, 2H, $J = 3.97$ Hz), 3.46-3.48 (m, 4H), 3.11-3.16 (m, 4H), 2.46-2.57 (m, 3H), 1.89-2.05 (m, 4H), 1.62-1.71 (m, 2H). LCMS ($M+H$) = 486.23. HPLC purity: > 95%.

5


Example 23



N-(4-Fluorobenzyl)-3-hydroxy-10-(((3-(methylsulfonyl)-2-oxo-1-imidazolidinyl)carbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (9 mg, 24% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.30 (s, 1H), 10.08 (s, 1H), 9.00 (s, 1H), 7.38 (dd, 2H, $J = 8.39$ and 5.34 Hz), 7.00 (t, 2Hz, $J = 8.70$ Hz), 4.66 (d, 2H, $J = 6.71$ Hz), 4.18 (d, 2H, $J = 3.66$ Hz), 3.88-3.97 (m, 4H), 2.97-3.05 (m, 2H), 2.92 (s, 3H), 2.48 (brs, 1H), 1.97-2.06 (m, 2H), 1.70-1.78 (m, 4H). LCMS ($M+H$) = 563.19. HPLC purity: > 95%.

20

Example 24




*10-(((1,5-Dimethyl-1*H*-pyrazol-3-yl)carbonyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* White solid (12 mg, 36% yield). ^1H NMR (500 MHz, CDCl_3) δ :

11.89 (s, 1H), 8.37 (s, 1H), 7.97 (s, 1H), 7.20 (dd, 2H, $J = 8.39$ and 5.34 Hz), 7.00 (t, 5 2H, $J = 8.70$ Hz), 6.46 (s, 1H), 4.60 (d, 2H, $J = 6.71$ Hz), 4.19 (d, 2H, $J = 3.97$ Hz), 3.41 (s, 3H), 2.84-2.93 (m, 2H), 2.50 (brs, 1H), 2.19 (s, 3H), 1.97-2.09 (m, 4H), 1.68-1.78 (m, 2H). ($M+\text{H}$) = 495.2. HPLC purity: > 95%.

Example 25


10



*Methyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)carbamate.* White solid (4 mg, 14% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.99 (s, 1H), 7.59 (s, 1H), 7.29-7.35 (m, 2H), 7.02-7.08 (m, 2H), 5.57 (s, 1H), 4.56 (d, 2H, $J = 5.80$ Hz), 4.15 (d, 2H, $J = 3.66$ Hz), 3.39 (s, 3H), 2.49 (s, 1H), 2.03-2.10 (m, 4H), 1.88-1.96 (m, 2H), 1.64-1.72 (m, 2H). LCMS ($M+\text{H}$) = 451.2. HPLC purity: > 95%.

20


Example 26



N-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((2-pyridinylcarbonyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (6mg, 19% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.99 (s, 1H), 10.13(s, 1H), 8.08-8.17 (m, 2H), 7.79-7.84 (m, 1H), 7.76 (d, 1H, J = 4.58 Hz), 7.34 (dd, 2H, J = 8.39 & 5.34 Hz), 7.26-7.30 (m, 1H), 7.04 (t, 2H, J = 8.70 Hz), 4.67 (d, 2H, J = 6.10 Hz), 4.21 (d, 2H, J = 3.66 Hz), 3.04-3.14 (m, 2H), 2.48 (s, 1H), 2.02-2.14 (m, 2H), 1.89-1.98 (m, 2H), 1.72-1.82 (m, 2H). LCMS ($\text{M}+\text{H}$) = 478.24. HPLC purity: > 95%.

10


Example 27



10-((Dimethylcarbamoyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (7 mg, 24% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.89 (s, 1H), 7.29-7.35 (m, 3H), 7.08 (t, 2H, J = 8.55 Hz), 5.54 (s, 1H), 4.54 (d, 2H, J = 5.80 Hz), 4.16 (d, 2H, J = 3.66 Hz), 2.64 (s, 6H), 2.55-2.61 (m, 2H), 2.45 (s, 1H), 1.88-2.00 (m, 4H), 1.61-1.69 (m, 2H). LCMS ($\text{M}+\text{H}$) = 444.21. HPLC purity: > 95%.

20

Example 28




N-(4-Fluorobenzyl)-3-hydroxy-10-(((5-methyl-3-isoxazolyl)carbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.

White solid (12 mg, 37% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.01 (s, 1H), 9.05 (s, 1H), 7.97-8.05 (m, 1H), 7.40 (dd, 2H, $J = 8.55 \text{ & } 5.49 \text{ Hz}$), 7.03 (t, 2H, $J = 8.55 \text{ Hz}$), 6.38 (s, 1H), 4.60 (d, 2H, $J = 6.10 \text{ Hz}$), 4.17 (d, 2H, $J = 3.97 \text{ Hz}$), 2.91-3.01 (m, 2H), 2.48 (s, 1H), 2.46 (s, 3H), 2.00-2.11 (m, 2H), 1.87-1.96 (m, 2H), 1.71-1.79 (m, 2H). LCMS ($\text{M}+\text{H}$) = 482.24. HPLC purity: > 95%.

Example 29


10



10-((N,N-Dimethylglycyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (10 mg, 33% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.1 (1 H, brs), 9.0 (1 H, brs), 8.8 (1 H, s), 7.4 (2 H, dd, $J=8.55, 5.49 \text{ Hz}$), 7.0 (2 H, t, $J=8.55 \text{ Hz}$), 4.6 (2 H, d, $J=6.41 \text{ Hz}$), 4.1 (2 H, d, $J=3.66 \text{ Hz}$), 3.6 (2 H, s), 2.7 (6 H, s), 2.4 - 2.5 (3 H, m), 2.0 - 2.1 (2 H, m), 1.9 - 2.0 (2 H, m), 1.6 - 1.7 (2 H, m). LCMS ($\text{M}+\text{H}$) = 458.20. HPLC purity: > 95%.

20

Example 30




10-((2-(Dimethylamino)-2-oxoethyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.

Off-white solid (10 mg, 33% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.6 (1 H, brs), 8.8 (1 H, s), 7.4 (2 H, dd, $J=8.55, 5.19$ Hz), 7.0 - 7.0 (2 H, m), 4.5 (2 H, d, $J=6.41$ Hz), 4.1 (2 H, d, $J=3.66$ Hz), 4.0 (2 H, s), 3.0 (3 H, s), 2.8 (3 H, s), 2.5 - 2.6 (3 H, m), 2.2 - 2.2 (2 H, m), 2.0 - 2.1 (2 H, m), 1.7 - 1.8 (2 H, m). LCMS ($\text{M}+\text{H}$) = 458.17. HPLC purity: > 95%.

Example 31


10



N-(4-Fluorobenzyl)-3-hydroxy-10-((methoxyacetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (7 mg, 24% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.0 (1 H, s), 8.4 (1 H, s), 7.7 (1 H, brs), 7.3 (2 H, dd, $J=8.39, 5.34$ Hz), 7.1 (2 H, t, $J=8.55$ Hz), 4.6 (2 H, d, $J=6.10$ Hz), 4.2 (2 H, d, $J=3.66$ Hz), 3.8 (2 H, s), 3.1 (3 H, s), 2.9 - 3.0 (2 H, m), 2.5 (1 H, s), 2.0 - 2.1 (2 H, m), 1.8 - 1.9 (2 H, m), 1.6 - 1.7 (2 H, m). LCMS ($\text{M}+\text{H}$) = 445.4. HPLC purity: > 95%.

15
20


Example 32



N-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((tetrahydro-2H-pyran-4-ylcarbonyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (5 mg, 15% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.1 (1 H, brs), 7.3 - 7.4 (4 H, m), 7.1 (2 H, t, $J=8.55$ Hz), 4.6 (2 H, d, $J=6.10$ Hz), 5 4.2 (2 H, d, $J=3.66$ Hz), 3.8 - 3.9 (2 H, m), 3.3 (2 H, td, $J=11.44, 2.44$ Hz), 2.7 - 2.8 (4 H, m), 2.5 (1 H, brs), 2.2 - 2.3 (1 H, m, $J=11.37, 11.37, 4.12, 3.97$ Hz), 1.9 - 2.0 (2 H, m), 1.8 - 1.9 (2 H, m), 1.5 - 1.6 (4 H, m). LCMS ($\text{M}+\text{H}$) = 485.38. HPLC purity: > 95%.


10

Example 33




N-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((3-pyridinylcarbonyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (8 mg, 20% yield). ^1H NMR (500 MHz, CDCl_3) δ : 9.2 (1 H, s), 8.7 (1 H, d, $J=5.19$ Hz), 8.5 (1 H, d, $J=7.93$ Hz), 8.0 (1 H, s), 7.6 (1 H, dd, $J=7.93, 5.49$ Hz), 7.4 (1 H, brs), 7.1 (2 H, dd, $J=8.24, 5.49$ Hz), 6.9 (2 H, t, $J=8.55$ Hz), 4.5 (2 H, d, $J=5.80$ Hz), 4.2 (2 H, d, $J=3.66$ Hz), 2.7 - 2.8 (2 H, m), 2.5 (1 H, brs), 2.1 - 2.2 (2 H, m), 2.0 - 2.1 (2 H, m), 1.7 - 1.8 (2 H, m). LCMS ($\text{M}+\text{H}$) = 478.35. HPLC purity: > 95%.

Example 34




5 *N*-(4-*Fluorobenzyl*)-3-*hydroxy*-10-(((4-*methyl*-1,3-oxazol-5-*yl*)carbonyl)amino)-4-*oxo*-4,6,7,8,9,10-*hexahydro*-7,10-*ethanopyrimido*[1,2-*a*]azepine-2-carboxamide. Off-white solid (7 mg, 22% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.0 (1 H, brs), 8.2 (1 H, s), 7.5 (1 H, brs), 7.3 (2 H, dd, $J=8.39, 5.34$ Hz), 7.1 (2 H, t, $J=8.55$ Hz), 6.9 (1 H, s), 4.6 (2 H, d, $J=5.80$ Hz), 4.2 (2 H, d, $J=3.97$ Hz), 10 2.9 - 3.0 (2 H, m), 2.5 (1 H, brs), 2.4 (3 H, s), 2.0 - 2.1 (2 H, m), 1.9 - 2.0 (2 H, m), 1.7 - 1.8 (2 H, m). LCMS ($\text{M}+\text{H}$) = 482.33. HPLC purity: > 95%.

Example 35




15 *N*-(4-*Fluorobenzyl*)-3-*hydroxy*-4-*oxo*-10-((4-*pyridinylcarbonyl*)amino)-4,6,7,8,9,10-*hexahydro*-7,10-*ethanopyrimido*[1,2-*a*]azepine-2-carboxamide. Off-white solid (8 mg, 20% yield). ^1H NMR (500 MHz, CDCl_3) δ : 8.6 (2 H, d, $J=5.80$ Hz), 8.0 (1 H, s), 7.7 (2 H, d, $J=5.49$ Hz), 7.2 - 7.2 (3 H, m), 7.1 (2 H, t, $J=8.55$ Hz), 4.5 (2 H, d, $J=6.10$ Hz), 4.2 (2 H, d, $J=3.97$ Hz), 2.9 - 2.9 (2 H, m), 2.6 (1 H, brs), 2.0 - 2.1 (4 H, m), 1.7 - 1.8 (2 H, m). LCMS ($\text{M}+\text{H}$) = 478.35. HPLC purity: > 95%.

Example 36



5 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((phenylcarbonyl)amino)-
4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-
white solid (11 mg, 35% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.9 (1 H, brs), 7.7
(2 H, d, J =8.24 Hz), 7.5 (1 H, t, J =7.93 Hz), 7.4 (1 H, s), 7.3 - 7.3 (1 H, m), 7.2 - 7.3
(2 H, m), 7.1 (2 H, dd, J =8.55, 5.49 Hz), 7.0 (2 H, t, J =8.39 Hz), 4.4 (2 H, d, J =6.41
Hz), 4.2 (2 H, d, J =3.66 Hz), 2.8 - 2.9 (2 H, m), 2.5 (1 H, brs), 2.0 - 2.1 (4 H, m), 1.7
- 1.8 (2 H, m). LCMS ($\text{M}+\text{H}$) = 477.37. HPLC purity: > 95%.

Example 37



15


15 *N*-(4-Fluoro-2-(5-methyl-1*H*-1,2,4-triazol-1-yl)benzyl)-3-hydroxy-10-((5-methyl-3-isoxazolyl)carbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (18 mg, 29% yield).

20 ^1H NMR (500 MHz, CDCl_3) δ : 11.74 (1 H, brs), 8.94 (1 H, s), 8.00 (1 H, brs), 7.95 (1

H, s), 7.67 (1 H, dd, $J=8.24, 5.80$ Hz), 7.26 - 7.33 (1 H, m), 7.02 (1 H, dd, $J=8.09, 2.29$ Hz), 6.39 (1 H, s), 4.39 (2 H, d, $J=6.10$ Hz), 4.16 (2 H, d, $J=3.36$ Hz), 2.91 - 3.00 (2 H, m), 2.51 (1 H, brs), 2.45 (6 H, s), 2.00 - 2.09 (2 H, m), 1.89 - 1.98 (2 H, m), 1.69 - 1.79 (2 H, m). LCMS (M+H) = 563.72. HPLC purity: > 95%.


5

Example 38



10 *10-(Acetylamino)-N-(4-fluoro-2-(5-methyl-1H-1,2,4-triazol-1-yl)benzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* Off-white solid (6 mg, 11% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.86 (1 H, brs), 8.10 (1 H, brs), 8.08 (1 H, s), 7.71 (1 H, dd, $J=8.70, 6.26$ Hz), 7.57 (1 H, s), 7.28 - 7.35 (1 H, m), 7.00 - 7.08 (1 H, m), 4.30 (2 H, d, $J=6.10$ Hz), 4.15 (2 H, d, $J=3.66$ Hz), 2.83 - 2.92 (2 H, m), 2.52 (3 H, s), 1.99 (3 H, s), 1.93 - 2.03 (3 H, m), 1.78 - 1.88 (2 H, m), 1.57 - 1.70 (2 H, m). LCMS (M+H) = 496.66. HPLC purity: > 95%.


Example 39



5 *N*-(4-*Fluoro*-2-(5-methyl-1*H*-1,2,4-triazol-1-yl)benzyl)-3-hydroxy-10-((methylsulfonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide. Tan solid (12 mg, 21% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.70 (1 H, brs), 8.69 - 8.76 (1 H, m), 8.36 (1 H, s), 7.80 (1 H, dd, J =8.55, 5.80 Hz), 7.30 - 7.37 (1 H, m), 7.01 (1 H, dd, J =7.63, 2.14 Hz), 5.30 (1 H, brs), 4.32 (2 H, d, J =6.10 Hz), 4.13 (2 H, d, J =3.36 Hz), 3.03 (3 H, s), 2.54 (3 H, s), 2.49 (1 H, brs), 2.09 - 2.25 (4 H, m), 1.88 - 1.98 (2 H, m), 1.63 - 1.75 (2 H, m). LCMS (M+H)
= 532.63. HPLC purity: > 95%.

Example 40


15



20 10-((Dimethylsulfamoyl)amino)-*N*-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide. To a solution of 10-amino-*N*-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide (25 mg, 0.067 mmol) in CH_2Cl_2 (2 mL) was added Et_3N (0.047 mL, 0.336 mmol) followed by dimethylsulfamoyl

chloride (0.036 mL, 0.336 mmol) and the resulting mixture was stirred at room temperature overnight. The reaction mixture was concentrated to give crude product which was dissolved in methanol (2mL) and treated with sodium ethoxide (0.2 ml, 21 wt% in ethanol) and the heated at 70 °C overnight. The mixture was then cooled, 5 concentrated and purified by preparative HPLC to afford the title compound as a white solid (3.5 mg, 11% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.00 (s, 1H), 8.90 (s, 1H), 7.34 (dd, 2H, $J = 8.39 \text{ & } 5.34 \text{ Hz}$), 7.01 (t, 2H, $J = 8.55 \text{ Hz}$), 4.53 (d, 2H, $J = 6.41 \text{ Hz}$), 4.15 (d, 2H, $J = 3.97 \text{ Hz}$), 2.77 (s, 6H), 2.50 (s, 1H), 2.12 (t, 4H, $J = 7.63 \text{ Hz}$), 1.85-1.97 (m, 2H), 1.64-1.73 (m, 2H). LCMS ($\text{M}+\text{H}$) = 480.21. HPLC purity: 10 > 95%.


Example 41



15

10-((Cyclopropylsulfonyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid. (7.5 mg, 25% yield). ^1H NMR (500 MHz CDCl_3) δ : 11.9 (1 H, brs), 8.9 - 8.9 (1 H, m), 8.8 - 9.0 (1 H, m), 7.3 (2 H, dd, $J=8.55, 5.49 \text{ Hz}$), 7.0 (2 H, t, $J=8.70 \text{ Hz}$), 4.8 (1 H, brs), 4.5 (2 H, d, $J=6.41 \text{ Hz}$), 4.1 (2 H, d, $J=3.97 \text{ Hz}$), 2.5 - 2.5 (2 H, m), 2.2 (4 H, t, $J=7.78 \text{ Hz}$), 1.9 - 2.0 (2 H, m), 1.6 - 1.7 (2 H, m), 1.1 - 1.2 (2 H, m), 1.0 - 1.0 (2 H, m). LCMS ($\text{M}+\text{H}$) = 477.31. HPLC purity: > 95%.

Example 42




5 *10-((Cyclopropylsulfonyl)amino)-N-(4-fluoro-3-methylbenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.*

Off-white solid (15 mg, 25% yield). ^1H NMR (400 MHz, CDCl_3) δ : 11.94 (1 H, brs), 8.86 (1 H, brs), 7.20 (1 H, d, $J=7.28$ Hz), 7.12 - 7.18 (1 H, m), 4.83 (1 H, brs), 4.51 (2 H, d, $J=6.27$ Hz), 4.17 (2 H, d, $J=4.02$ Hz), 2.46 - 2.56 (2 H, m), 2.27 (3 H, d, $J=1.76$ Hz), 2.25 - 2.26 (1 H, m), 2.16 - 2.20 (3 H, m), 1.91 - 1.98 (2 H, m), 1.65 - 1.75 (2 H, m), 1.13 - 1.18 (2 H, m), 0.98 - 1.07 (2 H, m). LCMS ($\text{M}+\text{H}$) = 491.02. HPLC purity: > 95%.

Example 43

15




20 *N-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((2-oxo-1-pyrrolidinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* White crystalline solid (23 mg, 39% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.8 (1 H, s), 9.6 (1 H, brs), 8.5 (1 H, s), 7.4 (2 H, dd, $J=8.24, 5.49$ Hz), 7.0 (2 H, t, $J=8.70$ Hz), 4.6 (2 H, d, $J=6.41$ Hz), 4.1 (2 H, d, $J=3.66$ Hz), 3.9 (2 H, s), 3.5 (2 H, t, $J=7.17$ Hz), 2.9 - 3.1 (2 H, m), 2.4 (1 H, brs), 2.2 (2 H, t, $J=7.93$ Hz).

Hz), 2.0 - 2.1 (2 H, m), 1.9 - 2.0 (2 H, m), 1.6 - 1.7 (4 H, m). LCMS (M+H) = 498.24. HPLC purity: > 95%.

Example 44


5



N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylpropanediamide. White solid 10 (11 mg, 34% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.3 (1 H, s), 10.3 (1 H, s), 9.6 (1 H, brs), 7.3 - 7.4 (2 H, m), 7.0 - 7.0 (2 H, m), 4.6 (2 H, d, *J*=6.71 Hz), 4.2 (2 H, d, *J*=3.66 Hz), 3.3 (2 H, s), 3.1 - 3.1 (2 H, m), 3.1 (3 H, s), 2.8 (3 H, s), 2.5 (1 H, brs), 2.0 - 2.0 (2 H, m), 1.7 - 1.8 (4 H, m). LCMS (M+H) = 486.45. HPLC purity: > 95%.

15

Example 45




N-(4-Fluorobenzyl)-3-hydroxy-10-(((1-methyl-1*H*-pyrrol-2-yl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (19 mg, 31% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.91 (1 H, brs), 9.15 (1 H, s), 8.05 (1 H, brs), 7.91 (1 H, dd, *J*=4.39, 1.63 Hz), 7.31 - 7.37 (2 H, m), 6.93 - 7.00 (3 H, m), 6.19 (1 H, dd, *J*=4.39, 2.38 Hz), 4.60

(2 H, d, $J=6.27$ Hz), 4.16 (2 H, d, $J=3.76$ Hz), 3.77 (3 H, s), 2.79 - 2.90 (2 H, m), 2.44 - 2.53 (1 H, m), 1.86 - 2.07 (4 H, m), 1.68 - 1.79 (2 H, m). LCMS (M+H) = 508.10. HPLC purity: > 95%.

5


Example 46



- 10 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((5-oxopropyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (23 mg, 39% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.71 (1 H, brs), 8.50 (1 H, s), 7.63 (1 H, brs), 7.44 - 7.49 (2 H, m), 7.01 (2 H, t, $J=8.70$ Hz), 5.93 (1 H, s), 4.75 (1 H, dd, $J=14.19, 6.87$ Hz), 4.60 (1 H, dd, $J=14.19, 6.26$ Hz), 4.24 (1 H, dd), 4.04 - 4.13 (2 H, m), 2.95 - 3.09 (2 H, m), 2.53 - 2.64 (1 H, m), 2.47 (1 H, brs), 2.34 - 2.40 (2 H, m), 2.19 - 2.28 (1 H, m), 1.95 - 2.07 (2 H, m), 1.69 - 1.76 (2 H, m), 1.59 - 1.67 (2 H, m). LCMS (M+H) = 484.49. HPLC purity: > 95%.

20

Example 47




*10-(((7-Chloro-4-methoxy-1*H*-pyrrolo[2,3-*c*]pyridin-3-yl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* Off-white solid (23 mg, 29% yield).

5 ^1H NMR (500 MHz, CDCl_3) δ : 11.87 (1 H, brs), 9.24 - 9.34 (2 H, m), 8.97 (1 H, d, J =2.75 Hz), 8.09 (1 H, brs), 7.84 (1 H, s), 7.39 (2 H, dd, J =8.09, 5.65 Hz), 6.91 (2 H, t, J =8.39 Hz), 4.62 (2 H, d, J =6.10 Hz), 4.19 (2 H, d, J =3.36 Hz), 3.95 (3 H, s), 2.79 - 2.91 (2 H, m), 2.53 (1 H, brs), 1.95 - 2.10 (4 H, m), 1.73 - 1.80 (2 H, m). LCMS (M+H) = 609.60. HPLC purity: > 95%.


10

Example 48



*N-(4-Fluorobenzyl)-10-(((4-fluoro-7-(1*H*-1,2,4-triazol-1-yl)-1*H*-pyrrolo[2,3-*c*]pyridin-3-yl)(oxo)acetyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* Yellow solid (15 mg, 18% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.02 (1 H, brs), 11.16 (1 H, brs), 9.64 (1 H, s), 9.31 (1 H, s), 8.78 (1 H, s), 8.20 (1 H, s), 8.09 (1 H, brs), 7.94 (1 H, s), 7.44 - 7.53 (2 H, m), 6.95 - 7.02 (2 H, m), 4.66 (2 H, d, J =6.41 Hz), 4.19 (2 H, d, J =3.36 Hz), 2.89 - 3.00 (2 H, m), 2.54 (1 H, brs), 2.02 - 2.13 (2 H, m), 1.89 - 1.99 (2 H, m), 1.72 - 1.83 (2 H, m). LCMS (M+H) = 630.75. HPLC purity: > 95%.


Example 49



5 *N*-(4-Fluorobenzyl)-3-hydroxy-10-(((1-methyl-1*H*-imidazol-2-yl)carbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide. Off-white solid(10.5 mg, 18% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.95 (1 H, brs), 9.66 (1 H, s), 8.44 (1 H, t, $J=6.26$ Hz), 7.32 - 7.35 (2 H, m), Y7.00 - 7.03 (2 H, m), 6.87 (1 H, s), 6.56 (1 H, d, $J=1.22$ Hz), 4.61 (2 H, d, $J=6.41$ Hz), 4.18 (2 H, d, $J=3.97$ Hz), 3.99 (3 H, s), 2.89 - 2.97 (2 H, m), 2.50 (1 H, brs), 1.92 - 2.08 (4 H, m), 1.70 - 1.79 (2 H, m). ($\text{M}+\text{H}$) = 480.97. HPLC purity: > 95%.

Example 50

15




20 *N'*-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N*-methyl-*N*-(2,2,2-trifluoroethyl)ethanediamide. Crystalline white solid (22 mg, 33% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.02 (0.4 H, s), 11.98 (0.6 H, s), 8.95 (0.6 H, s), 8.55 (0.4 H, s), 8.27 (0.4 H, brs), 8.17 (0.6 H, brs), 7.33 - 7.40 (2 H, m), 6.99 - 7.06 (2 H, m), 4.67

(1.3 H, q, $J=8.85$ Hz), 4.58 (1.3 H, d, $J=6.41$ Hz), 4.54 (0.7 H, d, $J=6.41$ Hz), 4.16 (2 H, t, $J=3.51$ Hz), 3.89 - 3.95 (0.7 H, m), 3.44 (1.2 H, s), 3.02 (1.8 H, s), 2.59 - 2.70 (2 H, m), 2.50 (1 H, brs), 1.90 - 2.06 (4 H, m), 1.66 - 1.76 (2 H, m). LCMS (M+H) = 540.01. HPLC purity: > 95%.

5


Example 51



10 *N'*-(2-((4-Fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N*-methyl-*N*-(2,2,2-trifluoroethyl)ethanediamide. Crystalline off-white solid (26 mg, 40% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.06 (0.4 H, s), 12.02 (0.6 H, s), 8.97 (0.6 H, s), 8.51 (0.4 H, s), 8.26 (0.4 H, brs), 8.17 (0.6 H, m), 7.14 - 7.23 (2 H, m), 6.92 - 6.99 (1 H, m), 4.65 - 4.70 (1.3 H, m), 4.70 (1.3 H, s), 4.54 (1.3 H, d, $J=6.10$ Hz), 4.50 (0.7 H, d, $J=6.10$ Hz), 4.16 (2 H, t, $J=3.66$ Hz), 3.91 (0.7 H, q, $J=8.85$ Hz), 3.43 (1.2 H, s), 3.03 (1.8 H, s), 2.57 - 2.71 (2 H, m), 2.50 (1 H, brs), 2.26 (3 H, s), 1.91 - 2.07 (4 H, m), 1.67 - 1.76 (2 H, m). LCMS (M+H)+= 554.02. HPLC purity: > 95%.

15
20

Example 52




N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H-yl)-N-(2-fluoroethyl)-N-methylethanediamide.

A mixture of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (0.05 g, 0.122 mmol),

5 2-((2-fluoroethyl)(methyl)amino)-2-oxoacetic acid (0.036 g, 0.245 mmol), diisopropyl ethylamine (0.128 mL, 0.734 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.093 g, 0.245 mmol) and 4-(dimethylamino)pyridine (DMAP) (2.99 mg, 0.024 mmol) in DMF (1.2 mL) was stirred at room temperature for 16 h. The mixture was diluted with 1N HCl and extracted with ethyl acetate. Solids precipitated out of the biphasic mixture and were collected by filtration to give the title compound as white needles (0.0394 g, 60% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.92 - 12.05 (1 H, m), 8.16 - 8.68 (2 H, m), 7.28 - 7.42 (2 H, m), 6.91 - 7.10 (2 H, m), 4.37 - 4.75 (4 H, m), 4.16 (2 H, t, J = 4.58 Hz), 3.94 - 4.08 (1 H, m), 3.50 - 3.65 (1 H, m), 2.91 - 3.43 (3 H, m), 2.53 - 2.66 (2 H, m), 2.45 - 2.52 (1 H, m), 1.88 - 2.11 (4 H, m), 1.61 - 1.78 (2 H, m). LCMS (M+H) calcd for C₂₄H₂₈F₂N₅O₅: 504.20; found: 504.02.

Example 53

20




N-(4-Fluorobenzyl)-3-hydroxy-10-(((5-methyl-1,3,4-oxadiazol-2-yl)carbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a suspension of potassium 5-methyl-1,3,4-oxadiazole-2-carboxylate (112 mg, 0.671 mmol) in CH₂Cl₂ (2 mL, contains cat. DMF) was added oxalyl chloride (0.369 mL, 0.738 mmol) and the mixture was stirred at room temp for 2h.. Solvent was then removed under reduced pressure. The crude acid

chloride was then diluted with CH₂Cl₂ (1.000 mL) and was added to a pre-stirred solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (25 mg, 0.067 mmol) and triethylamine (0.103 mL, 0.738 mmol) in CH₂Cl₂ (2 mL) and the resulting solution 5 stirred at room temperature. After 16 h at room temperature, the reaction mixture was concentrated to give crude product which was dissolved in methanol (2mL) and treated with 2M dimethylamine/methanol (0.5 mL). The resulting reaction mixture stirred at 60 °C for 2 h, cooled and purified by preparative HPLC to afford the title compound as a white solid (8 mg, 25% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.09
10 (s, 1H), 8.08 (s, 1H), 7.47 (dd, 2H, J = 8.55 & 5.49 Hz), 7.00-7.06 (m, 2H), 4.63 (d, 2H, J = 6.41 Hz), 4.18 (d, 2H, J = 3.66 Hz), 2.94-3.03 (m, 2H), 2.64 (s, 3H), 2.48-
2.55 (m, 1H), 2.01-2.14 (m, 2H), 1.87-1.95 (m, 2H), 1.73-1.81 (m, 2H) . LCMS
(M+H) = 483.27. HPLC purity: > 95%.


15

Example 54




20 *N*-(4-Fluorobenzyl)-3-hydroxy-10-(((1-methyl-1*H*-1,2,4-triazol-3-yl)carbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (12 mg, 36% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.05 (1 H, brs), 9.74 (1 H, s), 8.28 (1 H, brs), 7.42 (2 H, dd, *J*=8.24, 5.19 Hz), 7.04 - 7.18 (2 H, m), 4.76 (2 H, d, *J*=5.80 Hz), 4.21 (2 H, d, *J*=3.66 Hz), 3.97 (3 H, s), 3.11 - 3.26 (2 H, m), 2.55 (1 H, brs), 2.04 - 2.18 (2 H, m), 1.81 - 1.96 (2 H, m),
25 1.72 - 1.87 (2 H, m). LCMS (M+H) = 482.1. HPLC purity: > 95%.

Example 55



5 *N*-(4-Fluorobenzyl)-3-hydroxy-10-((3-methyl-2-oxobutanoyl)amino)-4-oxo-
4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Thick
film (2.7 mg, 11% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.18 (1 H, brs), 9.24 (1 H,
s), 7.84 (1 H, brs), 7.43 - 7.56 (2 H, m), 7.06 - 7.14 (2 H, m), 4.64 (2 H, d, *J*=6.10
Hz), 4.27 (2 H, d, *J*=3.97 Hz), 3.64 (1 H, dt, *J*=13.96, 6.90 Hz), 2.88 - 2.98 (2 H, m),
10 2.54 (1 H, brs), 2.02 - 2.1 (2 H, m), 1.84-1.92 (2 H, m), 1.71 - 1.80 (2 H, m), 1.12 (6
H, d, *J*=7.02 Hz). LCMS (M+H) = 471.00. HPLC purity: > 95%.

Example 56



15 *N'*-(2-((2,5-Dibromo-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. A mixture of ethyl 10-(((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (0.08 g, 0.187 mmol), (2,5-dibromo-4-fluorophenyl)methanamine hydrochloride (0.080 g, 0.250 mmol) and Et₃N (0.139 mL, 1 mmol) in EtOH (3 mL) was heated at 90 °C for 22 h. Then, cooled, diluted with Ethyl acetate (50 mL),


washed with 1N aq. HCl (2 x 5 mL), water (10 mL) followed by brine (10 mL), dried (Na₂SO₄), filtered and concentrated to give light purple foam which was used in the next without purification. To CH₂Cl₂ (5 mL) solution of this crude material was added HBr/AcOH (1.5 mL, 45% w/v) and the mixture stirred at room temperature for 5 4 h. The reaction mixture was concentrated and resulting light brown solid was used in the next step without purification. A 10 mL flask was charged with this intermediate and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.190 g, 0.5 mmol). To this was added diisopropyl-ethylamine (0.3 mL, 1.718 mmol), N,N-dimethyloxamic acid (0.117 g, 1 mmol) and 10 DMF (3 mL). After 4 h at room temperature, 2M Me₂NH/methanol (2 mL) was added and this was stirred at room temperature overnight (15 h). The solution was diluted with Et₂O (50 mL), washed with 1N HCl (10 mL), water (2 x 5 mL) followed by brine (5 mL), then dried (Na₂SO₄), filtered, concentrated and purified by preparative-HPLC to afford the title compound (0.0136 g, 0.020 mmol, 11% yield) as 15 a white solid. ¹H NMR (500 MHz, CDCl₃) δ: 11.84 (1H, s), 8.69 (1H, t, J = 5.8 Hz), 7.98 (1H, s), 7.56 (1H, d, J = 7.3 Hz), 7.35 (1H, d, J = 7.9 Hz), 4.63 (2H, d, J = 6.1 Hz), 4.17 (2H, d, J = 3.7 Hz), 3.28 (3H, s), 2.89 (3H, s), 2.55-2.49 (3H, m), 2.16-2.10 (2H, m), 2.00-1.93 (2H, m), 1.76-1.69 (2H, m). LCMS (M+H) calcd for C₂₃H₂₅Br₂FN₅O₅: 628.02; found: 629.88.

20

Examples 57 through 105 can be synthesized according to Scheme XII

Example 57

25




N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. To a stirred

solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (50mg, 0.134 mmol) in CH₂Cl₂ (2 mL) at 0 °C was added Et₃N (0.094 mL, 0.671 mmol) followed by methyl 2-chloro-2-oxoacetate (0.1 mL, 0.671 mmol) and the resulting mixture was stirred at 5 room temperature. After 16 h, the reaction mixture was concentrated to give crude product which was dissolved in EtOH (2mL), treated with 2M dimethylamine/methanol (0.7 mL) and heated in a sealed tube at 85 °C for 2 h, cooled and purified by preparative HPLC to afford the title compound as a light yellow solid (9 mg, 13% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.00 (s, 1H), 8.60 (s, 1H), 8.17 (s, 1H), 7.36 (dd, 2H, J = 8.55 and 5.49 Hz), 7.00-7.04 (m, 2Hz), 4.54 (d, 2H, J = 6.1 Hz), 4.16 (d, 2H, J = 3.66 Hz), 3.28 (s, 3H), 2.91 (s, 3H), 2.50-2.56 (m, 3H), 2.06-2.10 (m, 2H), 1.92-1.99 (m, 2H), 1.70- 1.74 (m, 2H). LCMS (M+H) = 472.37. HPLC purity: > 95%.


15

Example 58




20 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(1-pyrrolidinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (14 mg, 32% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.00 (s, 1H), 9.25 (s, 1H), 8.22 (t, 1H, J = 6.10 Hz), 7.38 (dd, 2H, J = 8.55 & 6.19 Hz), 7.00-7.04 (m, 2H), 4.58 (d, 2H, J = 2.75 Hz), 4.16 (d, 2H, J = 3.66 Hz), 3.91 (t, 2H, J = 6.87 Hz), 3.43 (t, 2H, J = 7.02 Hz), 2.69-2.75 (m, 2H), 2.49 (brs, 1H), 1.91-2.03 (m, 6H), 1.82-25 1.88 (m, 2H), 1.67-1.76 (m, 2H), 1.67-1.76 (m, 2H). LCMS (M+H) = 498.31. HPLC purity: > 95%.

Example 59




5 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(1-piperidinyl)acetyl)amino)-
4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-
white solid (11 mg, 25% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.02 (s, 1H), 8.85
(s, 1H), 7.65 (s, 1H), 7.36 (dd, 2H, J = 8.24 & 5.49 Hz), 7.00 (t, 2H, J = 8.55 Hz),
4.53 (d, 2H, J = 6.41 Hz), 4.16 (d, 2H, J = 3.66 Hz), 3.69 (t, 2H, J = 7.12 Hz), 3.47 (t,
10 2H, J = 5.87 Hz), 2.39-2.53 (m, 3H), 2.10-2.20 (m, 2H), 1.89-1.98 (m, 2H), 1.53-1.74
(m, 8H). LCMS (M+H) = 512.32. HPLC purity: > 95%.

Example 60




15 *N*-(4-Fluorobenzyl)-3-hydroxy-10-((4-morpholinyl(oxo)acetyl)amino)-4-oxo-
4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-
white solid (8 mg, 24% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.01 (s, 1H), 8.48 (t,
20 1H, J = 5.80 Hz), 8.34 (s, 1H), 7.33-7.37 (m, 2H), 6.99-7.03 (m, 2H), 4.55 (d, 2H, J =
6.41 Hz), 4.17 (d, 2H, J = 3.97 Hz), 3.98-4.01 (m, 2H), 3.66-3.72 (m, 4H), 3.51-3.56
(m, 2H), 2.48-2.60 (m, 3H), 2.02-2.10 (m, 2H), 1.91-2.01 (m, 2H), 1.68-1.76 (m,
2H). LCMS (M+H) = 514.30. HPLC purity: > 95%.

Example 61




5 *N*-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N*'-methylethanediamide. Off-white solid (14 mg, 32% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.01 (s, 1H), 9.79 (s, 1H), 7.98 (s, 1H), 7.43 (dd, 2H, J = 8.39 & 5.34 Hz), 7.30-7.37 (m, 1H), 7.05 (t, 2H, J = 8.70 Hz), 4.61 (d, 2H, J = 6.41 Hz), 4.15 (d, 2H, J = 3.66 Hz), 2.88 (d, 3H, J = 5.19 Hz), 2.79-2.87 (m, 2H), 2.49 (s, 1H), 1.98-2.07 (m, 2H), 1.80-1.87 (m, 2H), 1.70-1.78 (m, 2H). LCMS ($\text{M}+\text{H}$) = 458.23. HPLC purity: > 95%.

Example 62



15 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide. Off-white solid (40 mg, 19% yield). ^1H NMR (500 MHz, MeOD) δ : 9.55 (1 H, brs), 8.93 (1 H, s), 7.48 (2 H, dd, J =8.24, 5.49 Hz), 7.06 (2 H, t, J =8.85 Hz), 4.65 (2 H, d, J =6.10 Hz), 4.23 (2 H, d, J =3.97 Hz), 3.83 - 3.94 (4 H, m), 3.21 - 3.34 (4 H, m), 2.55 (1 H, brs), 2.32 - 2.45 (2 H, m), 2.25 - 2.36 (2 H, m), 1.94 - 2.06 (2 H, m), 1.72 - 1.89 (2 H, m). LCMS ($\text{M}+\text{H}$) = 513.44. HPLC purity: > 95%.


Example 63



5 *N*-(4-Fluorobenzyl)-3-hydroxy-10-(((4-hydroxy-1-piperidinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Light brown solid (3 mg, 10% yield).
¹H NMR (500 MHz, CDCl₃) δ: 8.74 (1 H, brs), 7.92 (1 H, s), 7.34 - 7.43 (2 H, m), 7.01 - 7.09 (2 H, m), 4.54 (2 H, d, *J*=5.80 Hz), 4.13 - 4.24 (4 H, m), 4.01 - 4.08 (1 H, m), 3.85 - 3.94 (1 H, m), 3.62 - 3.69 (1 H, m), 3.33 - 3.38 (1 H, m), 2.45 - 2.53 (3 H, m), 1.92 - 2.06 (2 H, m), 1.84 - 1.92 (2 H, m), 1.73 - 1.86 (2 H, m). LCMS (M+H)⁺ = 528.40. HPLC purity: > 95%.

Example 64

15




15 10-(((2R,6S)-2,6-Dimethyl-4-morpholinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Thick film (2.2 mg, 3% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.25 (1 H, brs), 8.52 (1 H, brs), 8.48 (1 H, s), 7.46 (2 H, dd, *J*=8.55, 5.19 Hz), 7.12 (2 H, t, *J*=8.70 Hz), 4.78 (1 H, d, *J*=13.43 Hz), 4.63 (2 H, dd, *J*=5.95, 3.20 Hz), 4.28 (2 H, brs), 4.17 (1 H, s), 3.58 - 3.65 (1 H, m), 3.46 - 3.53 (1 H, m), 2.87 (1

H, dd, $J=13.43, 10.68$ Hz), 2.54 - 2.66 (2 H, m), 2.59 (1 H, brs), 2.43 (2 H, dd, $J=13.12, 10.68$ Hz), 2.08 - 2.19 (2 H, m), 1.89 - 2.05 (2 H, m), 1.72 - 1.88 (2 H, m), 1.27 (3 H, d, $J=6.10$ Hz), 1.22 (3 H, d, $J=6.41$ Hz). LCMS (M+H) = 542.44. HPLC purity: > 95%.

5


Example 65



10 *N*-(4-Fluorobenzyl)-3-hydroxy-10-((3-hydroxy-1-azetidinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (3.5 mg, 7% yield).
¹H NMR (500 MHz, CDCl₃) δ: 12.12 (1 H, brs), 9.64 (1 H, s), 8.14 (1 H, brs), 7.48 (2 H, dd, $J=8.55, 5.49$ Hz), 7.03 - 7.19 (2 H, m), 4.84 (1 H, ddd, $J=11.83, 6.49, 1.83$ Hz), 4.64 - 4.72 (1 H, m), 4.56 - 4.67 (2 H, m), 4.47 (1 H, ddd, $J=11.75, 4.12, 1.83$ Hz), 4.33 (1 H, ddd, $J=11.67, 6.79, 1.68$ Hz), 4.14 (2 H, d, $J=3.66$ Hz), 3.90 (1 H, ddd, $J=11.83, 4.20, 1.68$ Hz), 2.82 - 2.98 (2 H, m), 2.58 (1 H, brs), 2.03 - 2.18 (2 H, m), 1.84 - 1.98 (2 H, m), 1.64 - 1.84 (2 H, m). LCMS (M+H) = 500.39. HPLC purity: > 95%.

15
20


Example 66



N-(4-Fluorobenzyl)-3-hydroxy-10-(((3-methoxy-1-azetidinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (12 mg, 25% yield).
¹H NMR (500 MHz, CDCl₃) δ: 12.12 (1 H, brs), 9.64 (1 H, s), 8.12 (1 H, brs), 7.46 (2 H, dd, J=8.39, 5.34 Hz), 7.14 (2 H, t, J=8.70 Hz), 4.83 (1 H, dd, J=11.75, 6.26 Hz), 4.53 - 4.66 (2 H, m), 4.58 (1 H, dd, J=11.75, 2.29 Hz), 4.14 - 4.26 (2 H, m), 4.19 (2 H, d, J=3.66 Hz), 3.92 - 4.08 (1 H, m), 2.82 - 2.96 (2 H, m), 2.54 (1 H, brs), 1.98 - 2.17 (2 H, m), 1.84 - 1.98 (2 H, m), 1.76 - 1.88 (2 H, m). LCMS (M+H) = 514.23. HPLC purity: > 95%.


10

Example 67



15 *10-((1-Azetidinyl(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* Off-white solid (20 mg, 45% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.06 (1 H, brs), 9.67 (1 H, s), 8.12 (1 H, brs), 7.37 - 7.46 (2 H, m), 7.06 (2 H, t, J=8.55 Hz), 4.58 - 4.71 (4 H, m), 4.11 (2 H, d, J=3.66 Hz), 4.20 (2 H, t, J=7.78 Hz), 2.78 - 2.94 (2 H, m), 2.55 (1 H, brs), 2.33 - 2.47 (2 H, m), 1.89 - 2.02 (2 H, m), 1.76 - 1.94 (2 H, m), 1.71 - 1.88 (2 H, m). LCM S (M+H) = 484.17. HPLC purity: > 95%.


Example 68



5 *10-(((3S)-3-(Acetylamino)-1-pyrrolidinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* White solid (12 mg, 25% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.03 (1 H, brs), 9.46 (0.5 H, s), 9.35 (0.5 H, s), 8.13 (1 H, brs), 7.37 - 7.42 (2 H, m), 7.00 - 7.06 (2 H, m), 5.73 (1 H, brs), 4.45 - 4.64 (3 H, m), 4.00 - 4.24 (4 H, m), 3.85 (0.5 H, dd, *J*=13.43, 6.10 Hz), 3.71 (0.5 H, dd, *J*=13.43, 6.10 Hz), 3.49 - 3.58 (1 H, m), 2.70 - 2.81 (2 H, m), 2.49 (1 H, brs), 2.11 - 2.26 (2 H, m), 2.04 (1.6 H, s), 2.02 (1.4 H, s), 1.80 - 2.00 (4 H, m), 1.64 - 1.75 (2 H, m). LCMS (M+H)⁺ = 555.56. HPLC purity: > 95%.

15

Example 69




20 *N-(4-Fluorobenzyl)-3-hydroxy-10-(((2S)-2-(hydroxymethyl)-1-pyrrolidinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* Off-white solid (mixture of rotamers) (1.6 mg, 4% yield). ¹H NMR (500 MHz, CDCl₃) δ: 9.47 (0.5 H, s), 9.25

(0.5 H, s), 8.04 - 8.17 (1 H, m), 7.32 - 7.42 (2 H, m), 6.97 - 7.08 (2 H, m), 4.41 - 4.69 (3 H, m), 4.09 - 4.24 (4 H, m), 3.77 - 4.08 (3 H, m), 3.43 - 3.70 (2 H, m), 2.68 - 2.87 (2 H, m), 2.50 (1 H, brs), 1.81 - 2.11 (4 H, m), 1.67 - 1.78 (2 H, m). LCMS (M+H) = 528.49. HPLC purity: > 95%.

5

Example 70




10 *10-(((3,3-Difluoro-1-pyrrolidinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-*
hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-
carboxamide. White solid (11 mg, 24% yield). ^1H NMR (500 MHz, CDCl_3) δ :

11.99 (1 H, brs), 9.64 (0.6 H, s), 9.54 (0.4 H, s), 7.98 - 8.10 (1 H, m), 7.36 - 7.42 (2 H, m), 6.99 - 7.07 (2 H, m), 4.59 (2 H, t, $J=5.34$ Hz), 4.23 - 4.34 (2 H, m), 4.16 (2 H, d, $J=3.66$ Hz), 3.76 & 3.68 (2 H, t, $J=7.48$ Hz), 2.73 - 2.87 (2 H, m), 2.49 (1 H, brs), 2.27 - 2.46 (2 H, m), 1.96 - 2.05 (2 H, m), 1.82 - 1.91 (2 H, m), 1.65 - 1.77 (2 H, m).
15 LCMS (M+H) = 534.35. HPLC purity: > 95%.

Example 71


20



N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-methyl-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (17 mg, 30% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.03 (1 H, brs), 8.64 (1 H, s), 8.19 (1 H, t, J=5.95 Hz), 7.35 (2 H, dd, J=8.70, 5.34 Hz), 7.00 - 7.05 (2 H, m), 5.08 - 5.25 (1 H, m), 4.19 - 4.71 (4 H, m), 3.91 - 4.13 (1 H, m), 3.50 - 3.74 (3 H, m), 3.31 (1 H, brs), 2.88 - 3.01 (2 H, m), 2.85 (3 H, s), 2.55 - 2.69 (2 H, m), 2.51 (1 H, brs), 2.17 (1 H, brs), 1.66 - 2.05 (4 H, m). LCMS (M+H) = 527.53. HPLC purity: > 95%.


10

Example 72




N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-(2-hydroxyethyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (6 mg, 10% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.02 (1 H, brs), 8.67 (1 H, s), 8.17 (1 H, t, J=5.80 Hz), 7.33 - 7.37 (2 H, m), 7.01 - 7.05 (2 H, m), 4.52 - 4.62 (2 H, m), 4.11 - 4.24 (2 H, m), 4.00 - 4.06 (3 H, m), 3.15 - 3.20 (4 H, m), 2.56 - 2.66 (4 H, m), 2.47 - 2.54 (2 H, m), 1.92 - 2.04 (5 H, m), 1.68 - 1.77 (3 H, m). LCMS (M+H) = 557.55. HPLC purity: > 95%.

Example 73



5 *N*-(4-Fluorobenzyl)-3-hydroxy-10-(((4-(2-methoxyethyl)piperazinyl)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (14 mg, 23% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.01 (1 H, brs), 8.60 (1 H, s), 8.22 (1 H, brs), 7.32 - 7.38 (2 H, m), 7.02 (2 H, t, $J=8.24$ Hz), 4.49 - 4.66 (3 H, m), 4.07 - 4.30 (4 H, m), 3.73 - 3.79 (4 H, m), 3.36 (3 H, s), 3.22 - 3.27 (3 H, m), 2.48 - 2.65 (3 H, m), 1.92 - 2.03 (3 H, m), 1.67 - 1.78 (2 H, m). LCMS ($\text{M}+\text{H}$) = 571.58. HPLC purity: > 95%.

Example 74




15

15 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(4-(tetrahydro-2-furanylcarbonyl)piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (6.4 mg, 15% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.88 (1 H, brs), 8.23 - 8.52 (2 H, m), 7.35 (2 H, dd, $J=$

8.39, 5.34 Hz), 7.02 (2 H, t, J = 8.55 Hz), 4.42 - 4.66 (3 H, m), 4.05 - 4.24 (3 H, m), 3.61 - 4.00 (5 H, m), 3.33 - 3.60 (3 H, m), 2.74 - 3.32 (3 H, m), 2.42 - 2.65 (3 H, m), 2.41 - 2.65 (3 H, m), 1.80 - 2.18 (6 H, m), 1.62 - 1.80 (2 H, m). LCMS (M+H) calcd for C₃₀H₃₆FN₆O₇: 611.26; found: 611.73.

5


Example 75



10 *N*-(4-Fluorobenzyl)-3-hydroxy-10-(((4-(2-hydroxyethoxy)ethyl)-1-piperazinyl)(oxo)acetyl)amino-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (11.1 mg, 28% yield). LCMS (M+H) calcd for C₂₉H₃₈FN₆O₇: 601.27; found: 601.71.

15


Example 76



20 *N*-(4-Fluorobenzyl)-3-hydroxy-10-(((4-(methylsulfonyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (2.0 mg, 6% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.99 (1 H, brs), 8.45 (1 H, s), 8.37 (1 H, t, J = 5.65 Hz),

7.36 (2 H, dd, $J = 8.55, 5.19$ Hz), 6.94 - 7.09 (2 H, m), 4.56 (2 H, d, $J = 6.10$ Hz),
 4.16 (2 H, d, $J = 3.97$ Hz), 4.05 - 4.13 (2 H, m), 3.55 - 3.74 (2 H, m), 3.24 - 3.34 (2
 H, m), 3.12 - 3.24 (2 H, m), 2.79 (3 H, s), 2.54 - 2.65 (2 H, m), 2.51 (1 H, brs), 1.88 -
 2.21 (4 H, m), 1.67 - 1.78 (2 H, m). LCMS (M+H) calcd for $C_{26}H_{32}FN_6O_7S$: 591.20;
 5 found: 591.33.

Example 77




10

N-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(4-phenyl-1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (9.8 mg, 30% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 12.00 (1 H, s), 8.52 (1 H, t, $J = 5.65$ Hz), 8.36 (1 H, s), 7.38 (2 H, dd, $J = 8.55, 5.49$ Hz), 7.29 (2 H, t, $J = 7.93$ Hz), 7.02 (2 H, t, $J = 8.70$ Hz), 6.84 - 6.97 (3 H, m), 4.56 (2 H, d, $J = 6.10$ Hz), 4.17 (2 H, d, $J = 3.66$ Hz), 4.10 - 4.15 (2 H, m), 3.65 - 3.72 (2 H, m), 3.19 (4 H, dt, $J = 14.27, 5.07$ Hz), 2.58 (2 H, ddd, $J = 14.34, 9.31, 5.34$ Hz), 2.51 (1 H, brs), 2.02 - 2.13 (2 H, m), 1.87 - 2.02 (2 H, m), 1.66 - 1.77 (2 H, m). LCMS (M+H) calcd for $C_{31}H_{34}FN_6O_5$: 589.25; found: 589.77.

15
20


Example 78



5 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(4-(2-pyridinyl)-1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (8.1 mg, 18% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.93 (1 H, brs), 8.66 (1 H, s), 8.32 (1 H, t, J = 5.95 Hz), 8.11 - 8.26 (1 H, m), 7.86 - 8.08 (1 H, m), 7.36 (2 H, dd, J = 8.55, 5.49 Hz), 7.02 (2 H, t, J = 8.70 Hz),
10 6.89 - 6.99 (2 H, m), 4.57 (2 H, d, J = 6.10 Hz), 4.21 - 4.33 (2 H, m), 4.16 (2 H, d, J = 3.66 Hz), 3.58 - 3.95 (6 H, m), 2.58 - 2.78 (2 H, m), 2.51 (1 H, brs), 1.88 - 2.15 (4 H, m), 1.60 - 1.78 (2 H, m). LCMS (M+H) calcd for $\text{C}_{30}\text{H}_{33}\text{FN}_7\text{O}_5$: 590.25; found: 590.74.

15

Example 79




20

N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-(4-morpholinylcarbonyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Pale pink foam (7.3 mg, 21% yield).

1 ^1H NMR (500 MHz, CDCl_3) δ : 12.01 (1 H, brs), 8.45 (1 H, t, J = 5.34 Hz), 8.35 (1 H, s), 7.36 (2 H, dd, J = 8.39, 5.34 Hz), 7.02 (2 H, t, J = 8.55 Hz), 4.55 (2 H, d, J = 6.10 Hz), 4.16 (2 H, d, J = 3.66 Hz), 3.91 - 4.01 (2 H, m), 3.60 - 3.81 (4 H, m), 3.42 - 3.59 (2 H, m), 3.20 - 3.39 (8 H, m), 2.41 - 2.63 (3 H, m), 2.01 - 2.11 (2 H, m), 1.92 - 2.01 (2 H, m), 1.64 - 1.76 (2 H, m). LCMS (M+H) calcd for $\text{C}_{30}\text{H}_{37}\text{FN}_7\text{O}_7$: 626.27; found: 626.80.


10

Example 80




15 *10-(((4-(Dimethylcarbamoyl)-1-piperazinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* White solid (14.4 mg, 44% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.00 (1 H, s), 8.50 (1 H, t, J = 5.95 Hz), 8.30 (1 H, s), 7.36 (2 H, dd, J = 8.55, 5.19 Hz), 6.95 - 7.09 (2 H, m), 4.55 (2 H, d, J = 6.10 Hz), 4.16 (2 H, d, J = 3.97 Hz), 3.97 (2 H, dd, J = 5.95, 4.12 Hz), 3.47 - 3.58 (2 H, m), 3.25 (4 H, ddd, J = 18.77, 5.04, 4.88 Hz), 2.85 (6 H, s), 2.45 - 2.63 (3 H, m), 2.00 - 2.12 (2 H, m), 1.86 - 2.01 (2 H, m), 1.63 - 1.80 (2 H, m). LCMS (M+H) calcd for $\text{C}_{28}\text{H}_{35}\text{FN}_7\text{O}_6$: 584.26; found: 584.25.

Example 81



5 *N'*-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N*-*(2-hydroxyethyl)*-*N*-methylethanediamide. Thick film (2.5 mg, 6% yield). ^1H NMR (500 MHz, CDCl_3) δ : 13.00 (1 H, brs), 8.54 (1 H, s), 8.40 (1 H, brs), 7.32 - 7.38 (2 H, m), 6.99 - 7.05 (2 H, m), 4.53 - 4.60 (2 H, m), 4.17 (2 H, d, J =3.66 Hz), 3.74 - 3.85 (2 H, m), 3.35 (1.5 H, s), 2.94 (1.5 H, s), 2.54 - 2.66 (4 H, m), 2.50 (1 H, brs), 1.94 - 2.10 (4 H, m), 1.68 - 1.76 (2 H, m). LCMS ($\text{M}+\text{H}$) = 502.44. HPLC purity: > 95%.

Example 82




15

15 *N*-Cyclopropyl-*N'*-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N*-methylethanediamide. Off-white solid (1.3 mg, 3% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.15 (1 H, brs), 9.71 (1 H, brs), 7.67 (1 H, brs), 7.37 - 7.40 (2 H, m), 7.04 - 7.07 (2 H, m), 4.60 (2 H, d, J =6.41 Hz), 4.17 (2 H, d, J =3.66 Hz), 2.82 - 2.90 (2 H, m), 2.52 (1 H, brs), 1.99 - 2.09 (2 H, m), 1.82 - 1.89 (2 H, m), 1.73 - 1.80 (2 H, m). LCMS ($\text{M}+\text{H}$) = 498.43. HPLC purity: > 95%.

From this reaction Examples 83 and 84 were also isolated

Example 83


5



Ethyl ((2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)amino)(oxo)acetate. White solid (12 mg, 29% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.00 (1 H, brs), 9.50 (1 H, s), 7.70 (1 H, t, $J=5.34$ Hz), 7.36 - 7.41 (2 H, m), 7.02 - 7.08 (2 H, m), 4.60 (2 H, d, $J=6.10$ Hz), 4.27 (3 H, q, $J=7.02$ Hz), 4.17 (2 H, d, $J=3.66$ Hz), 2.87 - 2.96 (2 H, m), 2.50 (1 H, brs), 1.99 - 2.08 (2 H, m), 1.77 - 1.85 (2 H, m), 1.67 - 1.77 (2 H, m), 1.36 (3 H, t, $J=7.17$ Hz). LCMS ($\text{M}+\text{H}$) = 473.43. HPLC purity: > 95%.

15

Example 84




((2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)amino)(oxo)acetic acid. Off-white solid (2.6 mg, 7% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.05 (1 H, brs), 8.92 (1 H, brs), 7.33 - 7.38 (2 H, m), 6.98 - 7.05 (3 H, m), 4.53 (2 H, d, $J=6.41$ Hz), 4.17 (2 H, d, $J=3.97$ Hz), 2.98 - 3.03 (1 H, m), 2.88 (3 H, s), 2.51 (1 H, brs), 2.30 - 2.38 (2 H, m),

2.20 - 2.27 (2 H, m), 2.02 - 2.10 (1 H, m), 1.90 - 1.98 (2 H, m), 1.71 - 1.76 (4 H, m).
 LCMS (M+H) = 445.39. HPLC purity: > 95%.

Example 85


5



10 *10-(((4,4-Difluoro-1-piperidinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* Off-white solid (4 mg, 8). ^1H NMR (500 MHz, CDCl_3) δ : 12.01 (1 H, brs), 8.43 (1 H, brs), 8.33 (1 H, s), 7.36 (2 H, dd, $J=7.78, 5.65$ Hz), 7.02 (2 H, t, $J=8.55$ Hz), 4.55 (2 H, d, $J=6.10$ Hz), 4.16 (2 H, d, $J=3.66$ Hz), 4.03 (2 H, t, $J=5.49$ Hz), 3.64 (2 H, t, $J=5.80$ Hz), 2.47 - 2.61 (3 H, m), 2.01 - 2.11 (4 H, m), 1.91 - 2.01 (4 H, m), 1.68 - 1.76 (2 H, m). LCMS (M+H)= 548.49. HPLC purity: > 95%.

15

Example 86




20 *10-(((3R)-3-(Dcetylamino)-1-pyrrolidinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* Off-white solid (17 mg, 0.031 mmol, 28% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.03 (1 H, brs), 9.46 (0.4 H, s), 9.35 (0.6 H, s), 8.13 (1

H, brs), 7.37 - 7.42 (2 H, m), 7.00 - 7.06 (2 H, m), 5.73 (1 H, brs), 4.45 - 4.64 (3 H, m), 4.00 - 4.24 (4 H, m), 3.85 (0.6 H, dd, $J=13.43, 6.10$ Hz), 3.71 (0.4 H, dd, $J=13.43, 6.18$ Hz), 3.49 - 3.58 (1 H, m), 2.70 - 2.81 (2 H, m), 2.49 (1 H, brs), 2.11 - 2.26 (2 H, m), 2.04 & 2.02 (3 H, s), 1.80 - 2.00 (4 H, m), 1.64 - 1.75 (2 H, m).

5 LCMS (M+H) = 555.56. HPLC purity: > 95%.

Example 87




10

N-(4-Fluorobenzyl)-3-hydroxy-10-(((2*R*)-2-(hydroxymethyl)-1-pyrrolidinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide. Off-white solid (18 mg, 31% yield).
¹H NMR (500 MHz, CDCl₃) δ: 9.47 (0.5 H, s), 9.25 (0.5 H, s), 8.04 - 8.17 (1 H, m),
15 7.32 - 7.42 (2 H, m), 6.97 - 7.08 (2 H, m), 4.41 - 4.69 (3 H, m), 4.09 - 4.24 (4 H, m),
3.77 - 4.08 (3 H, m), 3.43 - 3.70 (2 H, m), 2.68 - 2.87 (2 H, m), 2.50 (1 H, brs), 1.81 -
2.11 (4 H, m), 1.67 - 1.78 (2 H, m). LCMS (M+H) = 528.49. HPLC purity: > 95%.

Example 88


20



*N-(4-Fluorobenzyl)-3-hydroxy-10-(((2*R*)-2-(methoxymethyl)-1-pyrrolidinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* Off-white solid (20 mg, 34% yield).
¹H NMR (500 MHz, CDCl₃) δ: 12.03 (1 H, brs), 9.35 (0.6 H, s), 8.93 (0.4 H, s), 8.23
5 - 8.37 (1 H, m), 7.34 - 7.42 (2 H, m), 7.02 (2 H, q, J=8.44 Hz), 4.91 - 4.98 (1.3 H, m),
4.07-4.26 (1.7 H, m), 4.58 (2 H, dd, J=10.07, 6.41 Hz), 3.97 (2 H, s), 3.89 - 3.94 (1
H, m), 3.35 - 3.53 (3 H, m), 3.32 (3 H, s), 2.62 - 2.80 (3 H, m), 2.49 (1 H, brs), 1.85 -
2.03 (5 H, m), 1.66 - 1.77 (2 H, m). LCMS (M+H) = 542.69. HPLC purity: > 95%.


10

Example 89



*N-(4-Fluorobenzyl)-3-hydroxy-10-(((2*S*)-2-(methoxymethyl)-1-pyrrolidinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* Off-white solid (21 mg, 36% yield).
¹H NMR (500 MHz, CDCl₃) δ: 12.03 (1 H, brs), 9.35 (0.6 H, s), 8.93 (0.4 H, s), 8.23
15 - 8.37 (1 H, m), 7.34 - 7.42 (2 H, m), 7.02 (2 H, q, J=8.44 Hz), 4.91 - 4.98 (1.3 H, m),
4.07-4.26 (1.7 H, m), 4.58 (2 H, dd, J=10.07, 6.41 Hz), 3.97 (2 H, s), 3.89 - 3.94 (1
H, m), 3.35 - 3.53 (3 H, m), 3.32 (3 H, s), 2.62 - 2.80 (3 H, m), 2.49 (1 H, brs), 1.85 -
2.03 (5 H, m), 1.66 - 1.77 (2 H, m). LCMS (M+H) = 542.69. HPLC purity: > 95%.


Example 90



5 *N*-(4-*Fluorobenzyl*)-3-*hydroxy*-10-(((3-*methoxy*-1-*pyrrolidinyl*)(*oxo*)*acetyl*)*amino*)-4-*oxo*-4,6,7,8,9,10-*hexahydro*-7,10-*ethanopyrimido*[1,2-*a*]*azepine*-2-*carboxamide*. Tan solid (12 mg, 21% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.01 (1 H, brs), 9.37 (1 H, s), 8.18 (1 H, brs), 7.35 - 7.42 (2 H, m), 6.98 - 7.06 (2 H, m), 4.52 - 4.64 (2 H, m), 4.09 - 4.25 (3 H, m), 3.83 - 10 4.04 (2 H, m), 3.40 - 3.70 (2 H, m), 3.34 (3 H, d, J=6.10 Hz), 2.66 - 2.82 (3 H, m), 2.49 (1 H, brs), 1.79 - 2.21 (5 H, m), 1.66 - 1.76 (2 H, m). LCMS (M+H)= 528.66. HPLC purity: > 95%.

Example 91

15




20 10-(((2*S*)-2-(*Dimethylcarbamoyl*)-1-*pyrrolidinyl*)(*oxo*)*acetyl*)*amino*)-*N*-(4-*fluorobenzyl*)-3-*hydroxy*-4-*oxo*-4,6,7,8,9,10-*hexahydro*-7,10-*ethanopyrimido*[1,2-*a*]*azepine*-2-*carboxamide*. Brown solid (14 mg, 23% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.24 (1 H, brs), 9.14 (0.4 H, s), 8.78 (0.6 H, s), 8.36 (0.4 H, br.s), 8.24 (0.6 H, brs), 7.30 - 7.41 (2 H, m), 6.96 - 7.04 (2 H, m), 4.62 - 4.70 (2 H, m), 4.45 - 4.53 (1 H, m), 4.05 - 4.28 (3 H, m), 3.97 (1 H, s), 3.67 - 3.75 (1 H, m), 3.48 - 3.58 (1

H, m), 3.11 (1.4 H, s), 3.06 (1.6 H, s), 2.93 (3 H, s), 2.53 - 2.70 (2 H, m), 2.47 (1 H, brs), 2.27 - 2.37 (1 H, m), 2.04 - 2.19 (1 H, m), 1.80 - 2.04 (5 H, m), 1.62 - 1.78 (2 H, m). LCMS (M+H) = 569.76. HPLC purity: > 95%.

5


Example 92



10 *Ethyl 1-(((2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)amino)(oxo)acetyl)-L-proline.* Off-white solid (mixture of isomers) (12 mg, 19% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.88 (1 H, brs), 9.49 (0.6 H, s), 9.29 (0.4 H, s), 8.06 - 8.21 (1 H, m), 7.34 - 7.44 (2 H, m), 6.98 - 7.08 (2 H, m), 5.15 (1 H, dd, *J*=8.85, 3.66 Hz) & 4.57 (1 H, d, *J*=5.80 Hz), 4.61 - 4.69 (1 H, m), 4.54 (1 H, d, *J*=5.49 Hz) & 4.39 - 4.50 (1 H, m), 4.12 - 4.22 (4 H, m), 3.65 - 3.74 (1 H, m), 3.46 - 3.57 (1 H, m), 2.64 - 2.79 (2 H, m), 2.26 - 2.36 (2 H, m), 2.17 (1 H, brs), 1.83 - 2.02 (6 H, m), 1.65 - 1.75 (2 H, m), 1.25 (3 H, t, *J*=7.17 Hz). LCMS (M+H) = 570.76. HPLC purity: > 95%.


20

Example 93



N-(2-(Dimethylamino)ethyl)-N'-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-methylethanediamide. Off-white solid (20 mg, 48% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.10 (1 H, brs), 9.14 (0.7 H, brs), 8.95 (0.3 H, s), 8.23 (0.3 H, brs), 8.07 (0.7 H, s), 7.37 (0.5 H, dd, $J=8.55, 5.49$ Hz), 7.29 (1.5 H, dd, $J=8.39, 5.34$ Hz), 6.98 - 7.05 (2 H, m), 4.56 (2 H, d, $J=6.41$ Hz), 4.17 (1.5 H, d, $J=3.66$ Hz), 4.15 (0.5 H, d, $J=3.97$ Hz), 4.04 - 4.09 (0.5 H, m), 3.55 - 3.62 (1.5 H, m), 3.33 - 3.38 (0.5 H, m), 3.22 - 3.28 (1.5 H, m), 3.11 (2.3 H, s), 2.96 (0.7 H, s), 2.76 (6 H, s), 2.46 - 2.64 (3 H, m), 2.09 - 2.18 (1.5 H, m), 1.91 - 2.01 (2.5 H, m), 1.65 - 1.77 (2 H, m). LCMS 10 (M+H) = 529.68. HPLC purity: > 95%.


Example 94



15

N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-methyl-N-(tetrahydro-2-furanylmethyl)ethanediamide. White solid (3.5 mg, 9% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.05 (1 H, brs), 8.55 - 8.69 (1 H, m), 8.25 (0.7 H, s), 8.02 (0.3 H, s), 7.31 - 7.39 (2 H, m), 6.96 - 7.05 (2 H, m), 4.55 (2 H, dd, $J=6.10, 2.75$ Hz), 4.01 - 4.29 (3 H, m), 3.70 - 3.89 (3 H, m), 3.33 (1.1 H, s), 2.98 (1.9 H, s), 2.42 - 2.60 (3 H, m), 1.83 - 2.27 (8 H, m), 1.62 - 1.78 (2 H, m), 1.41 - 1.56 (1 H, m). LCMS (M+H) = 542.68. HPLC purity: > 95%.


Example 95



5 *N-((1,5-Dimethyl-1H-pyrazol-4-yl)methyl)-N'-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-methylethanediamide.* Light pink solid (2.6 mg, 7% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.20 (1 H, brs), 8.55 (0.5 H, brs), 8.45 (1 H, brs), 8.22 (0.5 H, s), 7.62 (0.4 H, s), 7.53 (0.6 H, s), 7.32 - 7.39 (2 H, m), 6.95 - 7.02 (2 H, m), 4.76 (1.5 H, s), 4.57 (2.5 H, dd, J =10.22, 6.26 Hz), 4.18 (2 H, d, J =3.05 Hz), 3.88 (1.4 H, s), 3.86 (1.6 H, s), 3.19 (1.6 H, s), 2.79 (1.4 H, s), 2.47 - 2.61 (3 H, m), 2.28 (1.3 H, s), 2.23 (1.7 H, s), 2.03 - 2.12 (2 H, m), 1.91 - 2.01 (2 H, m), 1.64 - 1.77 (2 H, m). LCMS ($\text{M}+\text{H}$) = 566.63. HPLC purity: > 95%.

15

Example 96




20 *N-((1,3-Dimethyl-1H-pyrazol-4-yl)methyl)-N'-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-methylethanediamide.* Thick film (2.5

mg, 4% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.55 (1 H, brs), 8.49 - 8.56 (1 H, m), 8.28 (0.4 H, s), 8.26 (0.6 H, s), 7.48 (0.6 H, s), 7.32 - 7.39 (2 H, m), 7.19 (0.4 H, s), 6.92 - 7.02 (2 H, m), 4.62 (1 H, s), 4.52 - 4.58 (2 H, m), 4.27 (1 H, s), 4.17 (2 H, d, $J=3.66$ Hz), 3.90 (1.6 H, s), 3.87 (1.4 H, s), 3.20 (1.4 H, s), 2.82 (1.6 H, s), 2.49 - 5 2.59 (3 H, m), 2.30 (1.6 H, s), 2.28 (1.4 H, s), 2.04 - 2.15 (2 H, m), 1.92 - 2.02 (2 H, m), 1.67 - 1.77 (2 H, m). LCMS ($\text{M}+\text{H}$) = 566.65. HPLC purity: > 95%.


Example 97

10




N,N-Diethyl-N'-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)ethanediamide. To a stirred solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-15 7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (50 mg, 0.122 mmol) in DMF (3 mL) was added 2-(diethylamino)-2-oxoacetic acid (35.5 mg, 0.245 mmol), diisopropyl-ethylamine (0.128 mL, 0.734 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (93 mg, 0.245 mmol) and 4-(dimethylamino)pyridine (DMAP) (2.99 mg, 0.024 mmol) and the resulting mixture stirred at room temperature for 3h. After work-up, the crude was then purified by prep HPLC to afford the title compound as a white solid (30 mg, 49% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.0 (1 H, brs), 8.7 (1 H, brs), 8.2 (1 H, s), 7.4 (2 H, dd, $J=8.09, 5.65$ Hz), 7.0 (2 H, t, $J=8.55$ Hz), 4.6 (2 H, d, $J=6.41$ Hz), 4.2 (2 H, d, $J=3.66$ Hz), 3.6 (2 H, q, $J=6.82$ Hz), 3.3 (2 H, q, $J=7.02$ Hz), 2.5 - 2.6 (3 H, m), 2.0 - 2.1 (2 H, m), 1.9 - 2.0 (2 H, m), 1.7 - 1.7 (2 H, m), 1.2 (3 H, t, $J=7.02$ Hz), 1.1 (3 H, t, $J=7.02$ Hz). LCMS ($\text{M}+\text{H}$) = 500.45. HPLC purity: > 95%.

Example 98



5 *10-((3,3-Difluoro-1-azetidinyl)(oxo)acetyl)amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* White solid (25 mg, 39% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.0 (1 H, brs), 9.7 (1 H, s), 7.9 (1 H, brs), 7.4 (2 H, dd, $J=8.09, 5.65$ Hz), 7.0 (2 H, t, $J=8.55$ Hz), 4.9 (2 H, t, $J=11.75$ Hz), 4.6 (2 H, d, $J=6.10$ Hz), 4.4 (2 H, t, $J=11.60$ Hz), 4.2 (2 H, d, $J=3.66$ Hz), 2.8 - 2.9 (2 H, m), 2.5 (1 H, brs), 2.0 - 2.1 (2 H, m), 1.8 - 1.9 (2 H, m), 1.7 - 1.8 (2 H, m). LCMS ($\text{M}+\text{H}$) = 520.34. HPLC purity: > 95%.

Example 99




15 *N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-methyl-N-(2-methylpropyl)ethanediamide.* White solid (26 mg, 41% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.0 (1 H, brs), 8.7 (1 H, brs), 8.1 (0.5 H, s), 8.0 (0.5 H, s), 7.4 (2 H, dd, $J=7.78, 5.65$ Hz), 7.0 (2 H, t, $J=8.55$ Hz), 4.5 (2 H, t, $J=5.49$ Hz), 4.2 (2 H, brs), 3.5 (1 H, d, $J=7.63$ Hz), 3.2 (1.5 H, s), 2.9 (1.5 H, s), 3.1 (1 H, d, $J=7.63$ Hz), 2.4 - 2.6 (3

H, m), 2.1 - 2.1 (2 H, m), 1.9 - 2.0 (3 H, m), 1.7 - 1.8 (2 H, m), 0.9 (6 H, t, $J=6.56$ Hz). LCMS (M+H) = 514.48. HPLC purity: > 95%.

Example 100


5



N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-(2-methoxyethyl)-N-methylethanediamide. Off-white solid (20 mg, 32% yield). ^1H NMR (500 MHz, CDCl₃) δ: 11.95 (1 H, brs), 8.63 (1 H, brs), 8.16 (0.4 H, s), 8.17 (0.6 H, s), 7.34 – 7.48 (2 H, m), 7.04 (2 H, t, $J=8.55$ Hz), 4.63 (2 H, t, $J=5.95$ Hz), 4.26 (2 H, t, $J=3.51$ Hz), 3.85 (1 H, t, $J=5.04$ Hz), 3.67 (1 H, t, $J=5.19$ Hz), 3.51 – 3.59 (2 H, m), 3.30 (3 H, s), 3.35 (1.2H, s), 3.03 (1.8 H, s), 2.56 – 2.64 (3 H, m), 2.03 – 2.11 (2 H, m), 1.94 – 2.06 (2 H, m), 1.73 – 1.87 (2 H, m). LCMS (M+H) = 516.26. HPLC purity: > 95%.

Example 101

20




N-Ethyl-N'-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-methylethanediamide.

White solid (25 mg, 42% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.97 (1 H, brs), 8.57 & 8.62 (1 H, brs), 8.28 (0.4 H, s), 8.05 (0.6 H, s), 7.32 - 7.37 (2 H, m), 6.96 - 5 7.01 (2 H, m), 4.53 (2 H, d, $J=6.27$ Hz), 4.14 (2 H, d, $J=4.02$ Hz), 3.62 (1.4 H, q, $J=7.19$ Hz), 3.32 (1.6 H, q, $J=7.19$ Hz), 3.23 (1.4 H, s), 2.85 (1.6 H, s), 2.43 - 2.58 (3 H, m), 2.0 - 2.13 (2 H, m), 1.88 - 1.99 (2 H, m), 1.63 - 1.71 (2 H, m), 1.22 (1.3 H, t, $J=7.15$ Hz), 1.11 (1.7 H, t, $J=7.15$ Hz). LCMS ($\text{M}+\text{H}$) = 486.16. HPLC purity: > 95%.


10

Example 102



15 *N-(4-Fluorobenzyl)-10-(((4-fluoro-1-piperidinyl)(oxo)acetyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* White solid (22 mg, 34% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.98 (1 H, brs), 8.58 (1 H, brs), 8.06 (1 H, s), 7.32 - 7.36 (2 H, m), 6.96 - 7.01 (2 H, m), 4.79 - 4.96 (1 H, m), 4.52 (2 H, dd, $J=6.27, 3.01$ Hz), 4.17 - 4.25 (1 H, m), 4.04 - 20 4.15 (2 H, m), 3.68 - 3.85 (2 H, m), 3.31 - 3.39 (1 H, m), 2.43 - 2.57 (3 H, m), 2.11 - 2.21 (1 H, m), 1.83 - 2.02 (6 H, m), 1.66 - 1.81 (3 H, m). LCMS ($\text{M}+\text{H}$) = 530.15. HPLC purity: > 95%.


Example 103



5 *N*-(4-*Fluorobenzyl*)-10-(((3*S*)-3-*fluoro*-1-*pyrrolidinyl*)(*oxo*)*acetyl*)*amino*)-3-*hydroxy*-4-*oxo*-4,6,7,8,9,10-*hexahydro*-7,10-*ethanopyrimido*[1,2-*a*]*azepine*-2-*carboxamide*. White solid (15 mg, 24% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.93 (1 H, brs), 9.49 (1 H, br.s.), 8.11 (1 H, brs), 7.35 - 7.41 (2 H, m), 6.98 - 7.03 (2 H, m), 5.28 - 5.36 (1 H, m), 4.51 - 4.64 (2 H, m), 4.30 - 4.46 (1 H, m), 4.07 - 4.20 (2 H, m), 3.66 - 4.02 (2 H, m), 3.45 - 3.61 (1 H, m), 2.70 - 2.85 (2 H, m), 2.47 (1 H, brs), 2.19 - 2.41 (1 H, m), 1.80 - 2.04 (5 H, m), 1.67 - 1.76 (3 H, m). LCMS ($\text{M}+\text{H}$) = 516.10. HPLC purity: > 95%.

Example 104

15




20 *N*-(4-*Fluorobenzyl*)-10-(((3*R*)-3-*fluoro*-1-*pyrrolidinyl*)(*oxo*)*acetyl*)*amino*)-3-*hydroxy*-4-*oxo*-4,6,7,8,9,10-*hexahydro*-7,10-*ethanopyrimido*[1,2-*a*]*azepine*-2-*carboxamide*. White solid (16 mg, 25% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.93 (1 H, brs), 9.49 (1 H, br.s.), 8.11 (1 H, brs), 7.35 - 7.41 (2 H, m), 6.98 - 7.03 (2 H, m), 5.28 - 5.36 (1 H, m), 4.51 - 4.64 (2 H, m), 4.30 - 4.46 (1 H, m), 4.07 - 4.20 (2 H, m), 3.66 - 4.02 (2 H, m), 3.45 - 3.61 (1 H, m), 2.70 - 2.85 (2 H, m), 2.47 (1 H,

brs), 2.19 - 2.41 (1 H, m), 1.80 - 2.04 (5 H, m), 1.67 - 1.76 (3 H, m). LCMS (M+H) = 516.10. HPLC purity: > 95%.

Example 105

5




N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N-methyl-N-(1-methylethyl)ethanediamide.

- 10 White solid (25 mg, 39% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.97 (1 H, brs), 8.95 (0.6 H, brs), 8.76 (0.4 H, brs), 7.95 (0.6 H, brs), 7.42 (0.4 H, s), 7.32 - 7.39 (2 H, m), 6.96 - 7.03 (2 H, m), 4.56 - 4.6 (0.6 H, m), 4.36-4.42 (0.4 H, m), 4.54 (2 H, d, J =6.41 Hz), 4.16 (2 H, d, J =3.66 Hz), 3.05 (1.4 H, s), 2.71 (1.6 H, s), 2.45 - 2.53 (3 H, m), 2.07 - 2.24 (2 H, m), 1.90 - 1.99 (2 H, m), 1.65 - 1.75 (2 H, m), 1.21 (3 H, d, J =6.71 Hz), 1.12 (3 H, d, J =6.71 Hz). LCMS (M+H) = 500.24. HPLC purity: > 95%.
- 15

Examples 106 through 128 can be synthesized according to Scheme XIII

20


Example 106



N'-(2-((4-Fluoro-2-(5-methyl-1*H*-1,2,3-triazol-1-yl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. A mixture of ethyl 10-(((dimethylamino)(oxo)acetyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-5 ethanopyrimido[1,2-*a*]azepine-2-carboxylat (30 mg, 0.076 mmol), (4-fluoro-2-(5-methyl-1*H*-1,2,3-triazol-1-yl)phenyl)methanamine, bis-hydrochloride salt (42.7 mg, 0.153 mmol) and triethylamine (0.1 ml, 0.717 mmol) in ethanol (2 mL) was heated at 90 ° C for 18 h. The reaction mixture was then cooled, concentrated and purified on preparative reversed phase HPLC to afford the title compound (10 mg, 21% yield) as an off-white solid. ^1H NMR (500 MHz, CDCl_3) δ: 1.66 - 1.77 (m, 2 H) 1.90 - 2.01 (m, 2 H) 2.14 - 2.24 (m, 2 H) 2.27 (s, 3 H) 2.39 - 2.47 (m, 2 H) 2.50 (br. s., 1 H) 2.89 (s, 3 H) 3.19 (s, 3 H) 4.15 (d, J =3.97 Hz, 2 H) 4.23 (d, J =6.41 Hz, 2 H) 6.99 (dd, J =8.24, 2.44 Hz, 1 H) 7.26 - 7.33 (m, 2 H) 7.62 - 7.68 (m, 2 H) 8.86 (t, J =6.10 Hz, 1 H) 11.73 (br. s., 1 H). LCMS (M+H) = 553.22. HPLC purity: > 95%.


15

Example 107



20 *N'*-(2-((4-Fluoro-2-(1-methyl-1*H*-1,2,4-triazol-5-yl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. Off-white solid (10 mg, 12% yield). ^1H NMR (500 MHz, CDCl_3) δ: 11.90 (1 H, brs), 8.89 (1 H, brs), 8.05 (1 H, s), 8.03 (1 H, s), 7.62 (1 H, dd, J =8.55, 5.49 Hz), 7.19 - 7.24 (1 H, m), 7.02 - 7.08 (1 H, m), 4.45 (2 H, d, J =6.10 Hz), 4.15 (2 H, d, J =3.97 Hz), 3.85 (3 H, s), 3.24 (3 H, s), 2.88 (3 H, s), 2.46 - 2.61 (3 H, m), 2.03 - 2.13 (2 H, m), 1.91 - 2.01 (2 H, m), 1.65 - 1.75 (2 H, m). LCMS (M+H) = 553.25. HPLC purity: > 95%.

Example 108




5 *N'*-(2-((4-Fluoro-2-(5-methyl-1*H*-1,2,4-triazol-1-yl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. White solid (3mg, 6% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.77 (1 H, brs), 9.01 (1 H, s), 8.21 (1 H, s), 7.82 (1 H, dd, J =8.85, 5.80 Hz), 7.76 (1 H, s), 7.28 - 7.35 (1 H, m), 6.99 (1 H, dd, J =7.93, 2.75 Hz), 4.28 (2 H, d, J =5.80 Hz), 4.14 (2 H, d, J =3.97 Hz), 3.12 (3 H, s), 2.90 (3 H, s), 2.53 (3 H, s), 2.51 (1 H, brs), 2.30 - 2.39 (2 H, m), 2.19 - 2.29 (2 H, m), 1.90 - 2.01 (2 H, m), 1.65 - 1.74 (2 H, m). LCMS ($\text{M}+\text{H}$) = 553.24. HPLC purity: > 95%.

10

Example 109

15




20 *N'*-(2-((4-Fluoro-2-(4-methyl-4*H*-1,2,4-triazol-3-yl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. White solid (6 mg, 11% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.67 (1 H, brs), 9.15 (1 H, s), 8.33 (1 H, s), 7.82 (1 H, s), 7.61 (1 H, dd,

J=8.70, 5.34 Hz), 7.20 - 7.24 (1 H, m), 7.02 - 7.09 (1 H, m), 4.45 (2 H, d, *J*=6.41 Hz), 4.15 (2 H, d, *J*=3.66 Hz), 3.65 (3 H, s), 3.15 (3 H, s), 2.86 (3 H, s), 2.39 - 2.53 (3 H, m), 2.17 - 2.28 (2 H, m), 1.90 - 1.99 (2 H, m), 1.65 - 1.75 (2 H, m). LCMS (M+H)= 553.2. HPLC purity: > 95%.

5


Example 110



10 *N'*-(2-((4-Fluoro-2-(methylcarbamoyl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. Brown solid (10 mg, 26% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.89 (1 H, brs), 8.84 (1 H, s), 8.07 (1 H, s), 7.83 (1 H, brs), 7.44 (1 H, dd, *J*=8.70, 5.34 Hz), 7.28 (1 H, dd, *J*=8.70, 2.59 Hz), 7.12 (1 H, td, *J*=8.32, 2.59 Hz), 4.64 (2 H, d, *J*=5.80 Hz), 4.21 (2 H, d, *J*=3.66 Hz), 3.32 (3 H, s), 3.07 (3 H, d, *J*=4.58 Hz), 2.99 (3 H, s), 2.53 - 2.67 (2 H, m), 2.08 - 2.15 (3 H, m), 1.92 - 2.06 (2 H, m), 1.67 - 1.81 (2 H, m). LCMS (M+H)= 529.43. HPLC purity: > 95%.

20


Example 111



N'-(2-((3,4-Difluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. Thick film (2.5 mg, 6% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.93 (1 H, brs), 8.73 (1 H, brs), 8.12 (1 H, s), 7.27 - 7.38 (1 H, m), 7.06 - 7.18 (2 H, m), 4.55 (2 H, d, $J=6.10$ Hz), 4.22 (2 H, d, $J=3.66$ Hz), 3.31 (3 H, s), 2.92 (3 H, s), 2.51 - 2.63 (3 H, m), 2.06 - 2.16 (2 H, m), 1.92 - 2.06 (2 H, m), 1.72 - 1.88 (2 H, m). LCMS ($\text{M}+\text{H}$) = 490.37. HPLC purity: > 95%.


Example 112

10



N'-(2-((3-Chloro-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. Off-white solid (7 mg, 17% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.78 (1 H, brs), 8.73 (1 H, brs), 7.93 (1 H, s), 7.45 (1 H, dd, $J=7.02, 2.14$ Hz), 7.26 - 7.36 (1 H, m), 7.14 (1 H, t, $J=8.70$ Hz), 4.58 (2 H, d, $J=6.41$ Hz), 4.22 (2 H, d, $J=3.66$ Hz), 3.34 (3 H, s), 2.91 (3 H, s), 2.44 - 2.58 (3 H, m), 2.13 - 2.27 (2 H, m), 1.89 - 2.06 (2 H, m), 1.72 - 1.87 (2 H, m). LCMS ($\text{M}+\text{H}$) = 506.27. HPLC purity: > 95%.


Example 113



5 *N'*-(2-((4-Fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. White solid (13 mg, 33% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.03 (1 H, brs), 8.64 (1 H, brs), 8.18 (1 H, s), 7.11 - 7.29 (2 H, m), 6.93 (1 H, t, J =8.85 Hz), 4.57 (2 H, d, J =6.41 Hz), 4.25 (2 H, d, J =3.66 Hz), 3.33 (3 H, s), 10 2.92 - 3.00 (3 H, m), 2.85 (3 H, s), 2.45 - 2.57 (3 H, m), 2.34 (3 H, s), 2.07 - 2.19 (2 H, m), 1.88 - 2.11 (2 H, m), 1.74 - 1.88 (2 H, m). LCMS ($\text{M}+\text{H}$) = 486.43. HPLC purity: > 95%.

Example 114

15




20 *N'*-(2-((2,4-Difluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. Off-white solid (6 mg, 19% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.89 (1 H, brs), 8.56 (1 H, brs), 8.11 (1 H, s), 7.34 (1 H, td, J =8.47, 6.56 Hz), 6.78 - 6.89 (2 H, m), 4.56 (2 H, d, J =6.41 Hz), 4.22 (2 H, d, J =3.97 Hz), 3.35 (3 H, s), 2.89 (3 H, s), 2.45 - 2.62 (3

H, m), 2.11 - 2.28 (2 H, m), 1.89 - 2.04 (2 H, m), 1.72 - 1.84 (2 H, m). LCMS (M+H) = 490.1. HPLC purity: > 95%.

Example 115


5



N'-(2-((3-Bromo-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. Off-white solid (5.5 mg, 16% yield). ^1H NMR (500 MHz, CDCl₃) δ: 12.23 (1 H, brs), 8.67 (1 H, brs), 7.89 (1 H, s), 7.56 (1 H, dd, J=6.41, 2.14 Hz), 7.23 - 7.32 (1 H, m), 7.11 (1 H, t, J=8.39 Hz), 4.56 (2 H, d, J=6.10 Hz), 4.28 (2 H, d, J=3.66 Hz), 3.31 (3 H, s), 2.92 (3 H, s), 2.34 - 2.51 (3 H, m), 2.11 - 2.23 (2 H, m), 1.89 - 2.05 (2 H, m), 1.67 - 1.82 (2 H, m). LCMS (M+H) = 552.1. HPLC purity: > 95%.

Example 116


20



N'-(2-((4-Fluoro-2-methoxybenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. Yellow solid (2.7 mg, 11% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.22 (1 H, brs), 8.62 (1 H, s), 8.28 (1 H, brs), 7.23 - 7.35 (1 H, m), 6.56 - 6.68 (2 H, m), 4.62 (2 H, d, $J=6.10$ Hz), 4.28 (2 H, d, $J=3.66$ Hz), 3.78 (3 H, s), 3.33 (3 H, s), 2.94 (3 H, s), 2.56 - 2.72 (2 H, m), 2.54 (1 H, brs), 1.89 - 2.05 (4 H, m), 1.57 - 1.83 (2 H, m). LCMS ($\text{M}+\text{H}$) = 502.2. HPLC purity: > 95%.

Example 117


10



N'-(2-((4-Fluoro-2-(methylsulfonyl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. Off-white solid (9 mg, 24% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.89 (1 H, brs), 8.89 (1 H, t, $J=6.26$ Hz), 7.94 (1 H, s), 7.78 (1 H, dd, $J=8.24, 2.75$ Hz), 7.66 (1 H, dd, $J=8.70, 5.04$ Hz), 7.28 - 7.35 (1 H, m), 4.89 (2 H, d, $J=6.71$ Hz), 4.25 (2 H, d, $J=3.66$ Hz), 3.37 (3 H, s), 3.21 (3 H, s), 3.07 (3 H, s), 2.55 - 2.67 (3 H, m), 2.12 - 2.26 (2 H, m), 1.89 - 2.05 (2 H, m), 1.72 - 1.88 (2 H, m).
15 LCMS ($\text{M}+\text{H}$) = 550.15. HPLC purity: > 95%.

20 LCMS ($\text{M}+\text{H}$) = 550.15. HPLC purity: > 95%.


Example 118



5 *N'*-(2-((2-Chloro-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. Off-white solid (4 mg, 16% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.93 (1 H, brs), 8.54 (1 H, brs), 8.31 (1 H, s), 7.48 (1 H, dd, J =8.55, 6.10 Hz), 7.12 - 7.19 (1 H, m), 6.89 - 7.08 (1 H, m), 4.73 (2 H, d, J =6.10 Hz), 4.25 (2 H, d, J =3.36 Hz), 3.33 (3 H, s), 2.92 (3 H, s), 2.45 - 2.63 (2 H, m), 2.55 (1 H, brs), 2.05 - 2.15 (2 H, m), 1.93 - 2.08 (2 H, m), 1.74 - 1.85 (2 H, m). LCMS ($\text{M}+\text{H}$) = 506.38. HPLC purity: > 95%.

Example 119

15




20 *N'*-(2-((4-Fluoro-3-(trifluoromethyl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. Off-white solid (5.5 mg, 20% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.88 (1 H, brs), 8.89 (1 H, brs), 7.83 (1 H, s), 7.63 - 7.78 (1 H, m), 7.56 - 7.68 (1 H, m), 7.12 - 7.23 (1 H, m), 4.67 (2 H, d, J =6.71 Hz), 4.28 (2 H, d, J =3.66

Hz), 3.33 (3 H, s), 2.94 (3 H, s), 2.44 - 2.56 (3 H, m), 2.11 - 2.19 (2 H, m), 1.90 - 2.08 (2 H, m), 1.67 - 1.78 (2 H, m). LCMS (M+H) = 540.36. HPLC purity: > 95%.

Example 120


5



10 *N'*-(2-((4-Fluoro-2-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. White solid (7 mg, 23% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.21 (1 H, brs), 8.32 - 8.39 (2 H, m), 7.31 - 7.35 (1 H, m), 6.81 - 6.91 (2 H, m), 4.55 (2 H, d, J =6.10 Hz), 4.16 (2 H, d, J =3.97 Hz), 3.29 (3 H, s), 2.88 (3 H, s), 2.52 - 2.61 (2 H, m), 2.49 (1 H, brs), 2.39 (3 H, s), 2.01 - 2.09 (2 H, m), 1.92 - 2.00 (2 H, m), 1.64 - 1.75 (2 H, m). LCMS (M+H) = 486.11. HPLC purity: > 95%.

15


Example 121



20 *N'*-(2-((4-Fluoro-2-(trifluoromethyl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. White solid (9 mg, 26% yield). ^1H NMR (500 MHz,

CDCl_3 δ: 11.87 (1 H, brs), 8.61 (1 H, brs), 8.05 (1 H, s), 7.55 (1 H, dd, $J=8.55, 5.49$ Hz), 7.37 (1 H, dd, $J=8.85, 2.75$ Hz), 7.22 (1 H, td, $J=8.09, 2.75$ Hz), 4.75 (2 H, d, $J=6.10$ Hz), 4.17 (2 H, d, $J=3.97$ Hz), 3.26 (3 H, s), 2.86 (3 H, s), 2.47 - 2.57 (3 H, m), 2.06 - 2.15 (2 H, m), 1.91 - 2.01 (2 H, m), 1.68 - 1.76 (2 H, m). LCMS (M+H) = 540.06. HPLC purity: > 95%.

Example 122




10

N' -(2-((4-Fluoro-2-(hydroxymethyl)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. Light yellow solid (2 mg, 6% yield). ^1H NMR (500 MHz, CDCl_3) δ: ppm 12.36 (1 H, brs), 8.73 (1 H, brs), 8.70 (1 H, s), 7.43 (1 H, dd, $J=8.39, 5.65$ Hz), 7.03 - 7.06 (1 H, m), 6.99 (1 H, td, $J=8.39, 2.75$ Hz), 4.77 (2 H, s), 4.66 (2 H, d, $J=6.10$ Hz), 4.13 (2 H, d, $J=3.97$ Hz), 3.32 (3 H, s), 3.00 (3 H, s), 2.80 - 2.88 (2 H, m), 2.46 (1 H, brs), 1.94 - 2.02 (2 H, m), 1.77 - 1.85 (2 H, m), 1.64 - 1.71 (2 H, m). LCMS (M+H) = 502.11. HPLC purity: > 95%.

20


Example 123



N'-(2-((3,5-Difluoro-2-pyridinyl)methyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4H-*y*l)-*N,N*-dimethylethanediamide. Yellow film (1.2 mg, 4% yield). ^1H NMR (500 MHz, CDCl₃) δ: 12.14 (1 H, brs), 8.74 (1 H, s), 8.41 (1 H, d, *J*=2.44 Hz), 8.37 (1 H, s), 7.21 - 7.24 (1 H, m), 4.76 (2 H, d, *J*=5.49 Hz), 4.17 (2 H, d, *J*=3.66 Hz), 3.30 (3 H, s), 2.96 (3 H, s), 2.60 - 2.67 (2 H, m), 2.50 (1 H, brs), 2.02 - 2.08 (2 H, m), 1.93 - 2.00 (2 H, m), 1.66 - 1.76 (2 H, m). LCMS (M+H) = 491.39. HPLC purity: > 95%.


Example 124

10



N'-(2-((3-(Difluoromethyl)-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4H-*y*l)-*N,N*-dimethylethanediamide. Off-white solid (4 mg, 12% yield). ^1H NMR (500 MHz, CDCl₃) δ: 11.89 (1 H, brs), 8.83 (1 H, brs), 7.90 (1 H, s), 7.58 - 7.62 (1 H, m), 7.50 - 7.55 (1 H, m), 7.09 (1 H, t, *J*=9.31 Hz), 4.58 (2 H, d, *J*=6.41 Hz), 4.16 (2 H, d, *J*=3.97 Hz), 3.26 (3 H, s), 2.92 (3 H, s), 2.44 - 2.53 (3 H, m), 2.09 - 2.17 (2 H, m), 1.91 - 2.00 (2 H, m), 1.67 - 1.76 (2 H, m). LCMS (M+H) = 522.44. HPLC purity: > 95%.


Example 125



5 *N'*-(3-Hydroxy-4-oxo-2-((3-(1*H*-1,2,4-triazol-1-yl)benzyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. Thick film (1.5 mg, 5% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.81 (1 H, brs), 8.97 (1 H, s), 8.92 (1 H, brs), 8.19 (1 H, s), 7.81 (1 H, s), 7.77 (1 H, s), 7.67 (1 H, d, J =7.63 Hz), 7.45 - 7.54 (2 H, m), 4.68 (2 H, d, J =6.71 Hz), 4.17 (2 H, d, J =3.66 Hz), 3.21 (3 H, s), 2.85 (3 H, s), 2.52 (1 H, brs), 2.41 - 2.47 (2 H, m), 2.15 - 2.22 (2 H, m), 1.91 - 2.01 (2 H, m), 1.64 - 1.76 (2 H, m). LCMS ($\text{M}+\text{H}$) = 521.59. HPLC purity: > 95%.

Example 126


15



20 *N'*-(3-Hydroxy-4-oxo-2-((3-(1*H*-pyrrol-1-yl)benzyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N*-dimethylethanediamide. Light yellow solid (16 mg, 51% yield). ^1H NMR (500

MHz, CDCl₃) δ: 11.99 (1 H, brs), 8.77 (1 H, brs), 8.02 (1 H, s), 7.42 (1 H, s), 7.38 (1 H, t, *J*=7.63 Hz), 7.27 - 7.32 (2 H, m), 7.10 (2 H, s), 6.32 (2 H, brs), 4.63 (2 H, d, *J*=6.41 Hz), 4.16 (2 H, d, *J*=3.66 Hz), 3.23 (3 H, s), 2.81 (3 H, s), 2.45 - 2.54 (3 H, m), 2.08 - 2.16 (2 H, m), 1.91 - 2.00 (2 H, m), 1.67 - 1.76 (2 H, m). LCMS (M+H)= 5 519.61. HPLC purity: > 95%.

Example 127




10

15 *N'*-(3-Hydroxy-2-((3-(4-methyl-1-piperazinyl)benzyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. Light yellow solid (19 mg, 47% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.08 (1 H, brs), 8.87 (1 H, brs), 7.70 (1 H, s), 7.22 - 7.27 (1 H, m), 6.99 (1 H, d, *J*=7.32 Hz), 6.93 (1 H, s), 6.81 (1 H, d, *J*=7.93 Hz), 4.55 (2 H, d, *J*=6.41 Hz), 4.16 (2 H, d, *J*=3.36 Hz), 3.62 - 3.71 (4 H, m), 3.24 - 3.33 (2 H, m), 3.22 (3 H, s), 2.96 - 3.04 (2 H, m), 2.86 (6 H, s), 2.50 (1 H, brs), 2.39 - 2.47 (2 H, m), 2.13 - 2.20 (2 H, m), 1.92 - 1.99 (2 H, m), 1.67 - 1.76 (2 H, m). LCMS (M+H)= 552.80. HPLC purity: > 95%.

20

Example 128



5 *N'*-(2-((3-Cyclopropyl-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylmethanediamide. A suspension of ethyl 10-(((dimethylamino)(oxo)acetyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (0.025 g, 0.050 mmol)


10 and dimethylamine (0.050 mL, 0.101 mmol) (2 M in methanol) in ethanol (1 mL) was stirred at 50 °C for 1 h. Added to the mixture were (3-cyclopropyl-4-fluorophenyl)methanamine hydrochloride salt (0.025 g, 0.126 mmol) and triethylamine (0.070 mL, 0.504 mmol) and the mixture stirred at 90 °C for 5 h. The mixture was cooled to room temperature and concentrated. The brown residue was

15 purified by preparative HPLC: Solvent A = 10% ACN/90% H₂O/0.1% trifluoroacetic acid; Solvent B = 90% ACN/10% H₂O/0.1% trifluoroacetic acid; Start % B = 0; Final % B = 100; Gradient time = 20 min; Stop time = 23 min; Column = Sunfire 19 x 100 mm, C18, 5µm. The resulting purple solid was triturated with hot methanol to give the title compound as a white solid (6.0 mg, 21% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.03 (1 H, s), 8.56 (1 H, t, *J* = 5.49 Hz), 8.15 (1 H, s), 7.11 - 7.20 (1 H, m), 6.85 - 7.01 (2 H, m), 4.49 (2 H, d, *J* = 6.10 Hz), 4.16 (2 H, d, *J* = 3.97 Hz), 3.28 (3 H, s), 2.91 (3 H, s), 2.42 - 2.65 (3 H, m), 2.01 - 2.18 (3 H, m), 1.86 - 2.00 (2 H, m), 1.71 (2 H, ddd, *J* = 13.28, 6.56, 6.41 Hz), 0.90 - 1.02 (2 H, m), 0.67 - 0.80 (2 H, m).

20 LCMS (M+H) calcd for C₂₆H₃₁FN₅O₅: 512.23; found: 512.24.

Examples 129 through 135 can be synthesized according to Scheme XIV


Example 129



5

N'-(2-(Benzylcarbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. To a mixture of ethyl 10-(((dimethylamino)(oxo)acetyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (20 mg, 0.040 mmol) in ethanol (2 mL) was added 2M Me₂NH (0.040 mL, 0.081 mmol) and the mixture heated at 50 °C for 1 h (mixture became clear after 5-10 min at 50 °C and then precepitate formed). Phenylmethanamine (0.022 mL, 0.201 mmol) followed by triethylamine (0.028 mL, 0.201 mmol) were then added and the mixture heated at 90 °C for 18 h. After work-up the crude product was purified by preparative HPLC to afford the title compound (7 mg, 0.015 mmol, 38% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ: 12.04 (1 H, s), 8.53 (1 H, brs), 8.29 (1 H, s), 7.35 - 7.39 (2 H, m), 7.28 - 7.34 (2 H, m), 7.24 - 7.27 (1 H, m), 4.57 (2 H, d, J=6.27 Hz), 4.14 (2 H, d, J=4.02 Hz), 3.26 (3 H, s), 2.87 (3 H, s), 2.44 - 2.58 (3 H, m), 2.01 - 2.10 (2 H, m), 1.86 - 1.99 (2 H, m), 1.64 - 1.75 (2 H, m). LCMS (M+H)⁺ = 454.73. HPLC purity: > 95%.


Example 130



5 *N'*-(2-((3-(4-Fluorophenyl)propyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. White solid (19 mg, 47% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.15 (1 H, brs), 8.54 (1 H, s), 8.08 (1 H, brs), 7.17 (2 H, dd, J =8.24, 5.49 Hz), 6.94 (2 H, t, J =8.70 Hz), 4.17 (2 H, d, J =3.66 Hz), 3.45 (2 H, q, J =6.71 Hz), 10 3.34 (3 H, s), 2.95 (3 H, s), 2.69 (2 H, t, J =7.63 Hz), 2.57 - 2.65 (2 H, m), 2.50 (1 H, brs), 1.93 - 2.09 (6 H, m), 1.68 - 1.77 (2 H, m). LCMS ($\text{M}+\text{H}$) = 500.72. HPLC purity: > 95%.

Example 131

15




20 *N'*-(2-((3-(3,4-Dichlorophenyl)propyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. White solid (21 mg, 47% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.09 (1 H, s), 8.47 (1 H, s), 8.11 (1 H, brs), 7.29 - 7.33 (2 H, m), 7.06 (1 H, dd, J =8.09, 1.98 Hz), 4.16 (2 H, d, J =3.66 Hz), 3.43 - 3.49 (2 H, m), 3.34 (3 H, s),

2.97 (3 H, s), 2.66 - 2.70 (2 H, m), 2.57 - 2.65 (2 H, m), 2.50 (1 H, brs), 1.93 - 2.10 (6 H, m), 1.68 - 1.77 (2 H, m). LCMS (M+H) = 550.59. HPLC purity: > 95%.


Example 132

5



N'-(2-((3-Cyano-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. Crystalline white solid (16 mg, 32% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.74 (1 H, s), 9.77 (1 H, s), 8.71 (1 H, s), 7.86 (1 H, dd, $J=6.27, 2.26$ Hz), 7.70 - 7.76 (1 H, m), 7.49 (1 H, t, $J=9.03$ Hz), 4.51 (2 H, d, $J=6.27$ Hz), 4.02 (2 H, d, $J=3.76$ Hz), 2.94 (3 H, s), 2.85 (3 H, s), 2.40 (1 H, brs), 2.23 - 2.34 (2 H, m), 1.94 - 2.08 (2 H, m), 1.73 - 1.84 (2 H, m), 1.49 - 1.67 (2 H, m). LCMS (M+H) = 497.68. HPLC purity: > 95%.

Example 133




20

N'-(2-((4-Fluoro-3-methoxybenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N*-dimethylethanediamide. White solid (17 mg, 34% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.97 (1 H, brs), 8.60 (1 H, t, $J=5.80$ Hz), 8.05 (1 H, s), 6.97 - 7.04 (2 H, m), 6.88 - 6.93 (1 H, m), 4.54 (2 H, d, $J=6.41$ Hz), 4.16 (2 H, d, $J=3.66$ Hz), 3.88 (3 H, s), 3.26 (3 H, s), 2.87 (3 H, s), 2.45 - 2.55 (3 H, m), 2.07 - 2.15 (2 H, m), 1.91 - 2.00 (2 H, m), 1.70 - 1.75 (2 H, m). LCMS ($\text{M}+\text{H}$) = 502.60. HPLC purity: > 95%.


Example 134

10



N'-(2-((4-Fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N*-(2-methoxyethyl)-*N*-methylethanediamide. White solid (17 mg, 43% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.09 (1 H, brs), 8.59 (1 H, s), 8.20 & 8.16 (1 H, s), 7.13 - 7.22 (2 H, m), 6.92 - 6.97 (1 H, m), 4.51 (2 H, t, $J=6.26$ Hz), 4.16 (2 H, t, $J=3.97$ Hz), 3.84 (1 H, t, $J=5.19$ Hz), 3.56 (1 H, t, $J=5.19$ Hz), 3.46 - 3.51 (2 H, m), 3.33 (3 H, s), 3.32 & 2.96 (2 H, s), 2.45 - 2.58 (3 H, m), 2.25 (3 H, d, $J=1.53$ Hz), 2.05 - 2.13 (2 H, m), 1.90 - 2.00 (2 H, m), 1.67 - 1.76 (2 H, m). LCMS ($\text{M}+\text{H}$) = 530.56. HPLC purity: > 95%.


Example 135



5 *N'*-(2-((3-Chloro-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N*-(2-methoxyethyl)-*N*-methylethanediamide. White solid (13 mg, 32% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.89 (1 H, brs), 8.77 (1 H, brs), 7.98 & 7.94 (1 H, s), 7.41 - 7.48 (1 H, m), 7.26 - 7.30 (1 H, m), 7.09 (1 H, t, $J=8.70$ Hz), 4.52 (2 H, t, $J=6.87$ Hz), 4.16 (2 H, t, $J=3.81$ Hz), 3.82 (1 H, t, $J=5.19$ Hz), 3.57 (1 H, t, $J=5.19$ Hz), 3.47 - 3.54 (2 H, m), 3.33 (3 H, s), 3.31 & 2.98 (2 H, s), 2.39 - 2.55 (3 H, m), 2.07 - 2.17 (2 H, m), 1.91 - 2.01 (2 H, m), 1.65 - 1.75 (2 H, m). LCMS ($\text{M}+\text{H}$) = 550.13. HPLC purity: > 95%.


Examples 136 through 149 can be synthesized according to Scheme XVI
15

Example 136



*N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-((1-methyl-1*H*-imidazol-5-yl)methyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* To a solution of 7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide, 10-[[1,2-dioxo-2-(1-piperazinyl)ethyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo- (0.025 g, 0.040 mmol) in 1,2-dichloroethane (1 mL) and diisopropyl ethylamine (6.97 μ L, 0.040 mmol) cooled to 0 °C was added 1-methyl-1*H*-imidazole-5-carbaldehyde (4.39 mg, 0.040 mmol) and the mixture stirred at 0 °C for 15 min. Acetic acid (2.284 μ L, 0.040 mmol) and sodium triacetoxyborohydride (9.30 mg, 0.044 mmol) were added and the mixture stirred at room temperature for 4 h. The solvent was removed under a stream of air and the residue purified by preparative HPLC: Solvent A = 10% methanol/90% H₂O/0.1% trifluoroacetic acid; Solvent B = 90% methanol/10% H₂O/0.1% trifluoroacetic acid; Start % B = 0; Final % B = 100; Gradient time = 15 min; Stop time = 15 min; Column = Sunfire 19 x 100 mm, C18, 5 μ m to give the title compound as a trifluoroacetic acid salt as a purple solid (14.7 mg, 49% yield). ¹H NMR (500 MHz, MeOD) δ : 9.67 (1 H, brs), 8.94 (1 H, brs), 8.86 (1 H, brs), 7.60 (1 H, brs), 7.32 - 7.48 (2 H, m), 7.03 (2 H, t, *J* = 8.39 Hz), 4.56 (2 H, brs), 4.19 (2 H, brs), 3.77 - 4.08 (4 H, m), 3.50 - 3.77 (4 H, m), 2.69 (3 H, brs), 2.51 (1 H, brs), 2.39 (2 H, d, *J* = 5.80 Hz), 2.21 (2 H, brs), 1.96 (2 H, d, *J* = 4.88 Hz), 1.74 (2 H, brs). (M+H) calcd for C₃₀H₃₆FN₈O₅: 607.27; found: 607.44.

Example 137




*N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-((1-methyl-1*H*-imidazol-2-*y*l)methyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* Purple foam (16.0 mg, 56% yield).

5 ^1H NMR (500 MHz, MeOD) δ : 7.58 (1 H, s), 7.53 (1 H, brs), 7.41 (2 H, dd, $J=8.55$, 5.49 Hz), 7.03 (2 H, t, $J=8.70$ Hz), 4.56 (2 H, s), 4.19 (2 H, d, $J=3.66$ Hz), 3.85 - 4.03 (4 H, m), 3.66 (2 H, brs), 3.59 (2 H, brs), 2.54 - 2.69 (3 H, m), 2.51 (1 H, brs), 2.30 - 2.46 (2 H, m), 2.20 (2 H, ddd, $J=14.42$, 9.23, 5.65 Hz), 1.87 - 2.05 (2 H, m), 1.66 - 1.81 (2 H, m). LCMS (M+H) calcd for $\text{C}_{31}\text{H}_{33}\text{FN}_7\text{O}_6$: 607.27; found: 607.39.

10

Example 138



15 *10-(((4-Acetyl-1-piperazinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* To a solution of N-(4-fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide (25 mg, 0.049 mmol) in CH_2Cl_2 (2 mL) was added triethylamine (0.034 mL, 0.244 mmol), followed by acetyl chloride (0.017 mL, 0.244 mmol) and the resulting mixture stirred at room temperature. After 16 h, the reaction mixture was concentrated to give crude product which was dissolved in methanol (2 mL) and treated with 2M dimethylamine in methanol (0.5 mL). The resulting reaction mixture was stirred at 60 °C for 2 h then cooled and purified by preparative HPLC to afford the title compound as a brown solid (13 mg, 46% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.05 (1 H, brs), 8.53 (1 H, brs), 8.48 (1 H, $J=78$, 5.65 Hz), 7.07 (2 H, t, $J=8.24$ Hz), 4.66 (2 H, d, $J=5.80$ Hz), 4.24 (2 H, d, $J=3.05$ Hz), 4.18 (1 H, brs), 3.95 - 4.03 (1 H, m), 3.75 (2 H, brs), 3.46 - 3.63 (4 H, m), 2.64 - 2.73 (2 H, m), 2.51-


20 mmol) and the resulting mixture stirred at room temperature. After 16 h, the reaction mixture was concentrated to give crude product which was dissolved in methanol (2 mL) and treated with 2M dimethylamine in methanol (0.5 mL). The resulting reaction mixture was stirred at 60 °C for 2 h then cooled and purified by preparative HPLC to afford the title compound as a brown solid (13 mg, 46% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.05 (1 H, brs), 8.53 (1 H, brs), 8.48 (1 H, $J=78$, 5.65 Hz), 7.07 (2 H, t, $J=8.24$ Hz), 4.66 (2 H, d, $J=5.80$ Hz), 4.24 (2 H, d, $J=3.05$ Hz), 4.18 (1 H, brs), 3.95 - 4.03 (1 H, m), 3.75 (2 H, brs), 3.46 - 3.63 (4 H, m), 2.64 - 2.73 (2 H, m), 2.51-

25 mmol) and the resulting mixture stirred at room temperature. After 16 h, the reaction mixture was concentrated to give crude product which was dissolved in methanol (2 mL) and treated with 2M dimethylamine in methanol (0.5 mL). The resulting reaction mixture was stirred at 60 °C for 2 h then cooled and purified by preparative HPLC to afford the title compound as a brown solid (13 mg, 46% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.05 (1 H, brs), 8.53 (1 H, brs), 8.48 (1 H, $J=78$, 5.65 Hz), 7.07 (2 H, t, $J=8.24$ Hz), 4.66 (2 H, d, $J=5.80$ Hz), 4.24 (2 H, d, $J=3.05$ Hz), 4.18 (1 H, brs), 3.95 - 4.03 (1 H, m), 3.75 (2 H, brs), 3.46 - 3.63 (4 H, m), 2.64 - 2.73 (2 H, m), 2.51-

2.58 (1 H, m), 2.11 (3 H, d, $J=9.46$ Hz), 1.95 - 2.14 (4 H, m), 1.76 - 1.82 (2 H, m). LCMS ($M+H$) = 555.56. HPLC purity: > 95%.

Example 139

5




N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-((5-methyl-3-isoxazolyl)carbonyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a suspension N-(4-fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (0.024 g, 0.038 mmol) in dichloromethane (1 mL) was added diisopropyl ethylamine (0.027 mL, 0.153 mmol). After stirring for 5 min, 5-methylisoxazole-3-carbonyl chloride (0.011 g, 0.077 mmol) was added and the solution stirred at room temperature for 1 h. Dimethylamine (0.192 mL, 0.383 mmol) (2 M in methanol) was added and the solution stirred at room temperature for 1 h then concentrated. The residue was purified by preparative HPLC: Solvent A = 10% methanol/90% H₂O/0.1% trifluoroacetic acid; Solvent B = 90% methanol/10% H₂O/0.1% trifluoroacetic acid; Start % B = 10; Final % B = 100; Gradient time = 20 min; Stop time = 25 min; Column = XTERRA[®] 19 x 50 mm, C18, 5μm (product elutes at 12.6 min) to give the title compound as a pale lavender solid (12.1 mg, 49% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.00 (1 H, brs), 8.51 (1 H, d, *J* = 5.80 Hz), 8.40 (1 H, d, *J* = 5.19 Hz), 7.28 - 7.50 (2 H, m), 6.88 - 7.12 (2 H, m), 6.33 (1 H, d, *J* = 2.14 Hz), 4.55 (2 H, t, *J* = 6.26 Hz), 4.16 (2 H, brs), 4.02 - 4.14 (2 H, m), 3.89 - 4.00 (2 H, m), 3.80 (2 H, dt, *J* = 18.08, 5.00 Hz), 3.50 - 3.68 (2 H, m), 2.53 - 2.77 (2 H, m), 2.48 (4 H, d, *J* = 7.93 Hz),

1.87 - 2.15 (4 H, m), 1.72 (2 H, brs). LCMS (M+H) calcd for C₃₀H₃₃FN₇O₇: 622.24; found: 622.78.


Example 140

5



N-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(4-(2-pyridinylcarbonyl)-1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Purple foam (14.1 mg, 49% yield). ¹H NMR (500 MHz, CDCl₃) δ: 8.72 (1 H, d, *J* = 12.51 Hz), 8.44 (1 H, t, *J* = 5.80 Hz), 8.26 - 8.41 (1 H, m), 8.04 (1 H, d, *J* = 6.41 Hz), 7.73 (1 H, d, *J* = 7.93 Hz), 7.60 (1 H, brs), 7.34 (2 H, d, *J* = 18.62 Hz), 6.89 - 7.12 (2 H, m), 4.54 (2 H, d, *J* = 9.77 Hz), 3.96 - 4.23 (4 H, m), 3.75 - 3.94 (2 H, m), 3.39 - 3.74 (4 H, m), 2.37 - 2.71 (3 H, m), 1.87 - 2.16 (4 H, m), 1.72 (2 H, brs). LCMS (M+H) calcd for C₃₁H₃₃FN₇O₆: 618.24; found: 618.43.

Example 141




20

N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-(methoxyacetyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (2.7 mg, 11% yield).

LCMS (M+H) calcd for C₂₈H₃₄FN₆O₇: 585.24; found: 585.75. ¹H NMR (400 MHz, CDCl₃) δ: 11.97 (1 H, brs), 8.42 - 8.65 (1 H, m), 8.35 (1 H, brs), 7.34 (2 H, dd, *J* = 8.53, 5.27 Hz), 7.00 (2 H, t, *J* = 8.66 Hz), 4.54 (2 H, d, *J* = 6.27 Hz), 4.14 (2 H, d, *J* = 3.76 Hz), 4.10 (2 H, brs), 4.00 (2 H, brs), 3.45 - 3.72 (6 H, m), 3.40 (3 H, brs), 2.52 - 2.70 (2 H, m), 2.48 (1 H, brs), 1.86 - 2.10 (4 H, m), 1.62 - 1.80 (2 H, m).


10

Example 142



N-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(4-(2-thiophenylcarbonyl)-1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (13.1 mg, 51% yield). LCMS ($M+H$) calcd for $C_{30}H_{32}FN_6O_6S$: 623.20; found: 623.77. 1H NMR (400 MHz, $CDCl_3$) δ : 11.97 (1 H, s), 8.54 (1 H, s), 8.35 (1 H, t, J = 6.02 Hz), 7.48 (1 H, dd, J = 5.02, 1.25 Hz), 7.31 - 7.38 (2 H, m), 7.30 (1 H, dd, J = 3.64, 1.13 Hz), 7.06 (1 H, dd, J = 5.02, 3.76 Hz), 6.93 - 7.03 (2 H, m), 4.54 (2 H, d, J = 6.27 Hz), 4.14 (2 H, d, J = 3.76 Hz), 4.01 - 4.10 (2 H, m), 3.68 - 3.86 (4 H, m), 3.57 (2 H, dd, J = 6.27, 4.27 Hz), 2.52 - 2.68 (2 H, m), 2.48 (1 H, brs), 1.84 - 2.15 (4 H, m), 1.63 - 1.79 (2 H, m).


Example 143



5 *N*-(4-Fluorobenzyl)-10-(((4-(2-furanylcarbonyl)-1-piperazinyl)acetyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. White solid (13.9 mg, 52% yield). LCMS (M+H) calcd for C₃₀H₃₂FN₆O₇: 607.23; found: 607.76. ¹H NMR (400 MHz, CDCl₃) δ: 11.97 (1 H, s), 8.51 (1 H, s), 8.38 (1 H, t, *J* = 6.15 Hz), 7.48 (1 H, d, *J* = 0.75 Hz), 7.29 - 7.42 (2 H, m), 7.06 (1 H, dd, *J* = 3.51, 0.75 Hz), 6.95 - 7.04 (2 H, m), 6.49 (1 H, dd, *J* = 3.39, 1.88 Hz), 4.54 (2 H, d, *J* = 6.27 Hz), 4.14 (2 H, d, *J* = 3.76 Hz), 3.99 - 4.11 (2 H, m), 3.71 - 3.93 (4 H, m), 3.59 (2 H, dd, *J* = 6.27, 4.27 Hz), 2.53 - 2.66 (2 H, m), 2.48 (1 H, d, *J* = 2.26 Hz), 1.86 - 2.14 (4 H, m), 1.63 - 1.79 (2 H, m).

15

Example 144




N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-((3-(methylsulfonyl)-2-oxo-1-imidazolidinyl)carbonyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Pale purple solid (5.3 mg, 17% yield). LCMS (M+H) calcd for C₃₀H₃₆FN₈O₉S: 703.23; found: 703.88.

5 ¹H NMR (500 MHz, CDCl₃) δ: 12.00 (1 H, brs), 8.27 - 8.53 (2 H, m), 7.36 (2 H, dd, *J* = 8.55, 5.19 Hz), 6.95 - 7.09 (2 H, m), 4.55 (2 H, d, *J* = 6.10 Hz), 4.16 (2 H, d, *J* = 3.66 Hz), 4.01 - 4.12 (2 H, m), 3.81 - 3.99 (4 H, m), 3.59 - 3.69 (2 H, m), 3.43 - 3.58 (4 H, m), 3.32 (3 H, s), 2.57 (2 H, ddd, *J* = 14.19, 9.31, 5.49 Hz), 2.50 (1 H, d, *J* = 2.14 Hz), 1.89 - 2.15 (4 H, m), 1.69 (2 H, brs).

10

Example 145



15 *10-(((4-(Dimethylsulfamoyl)-1-piperazinyl)(oxo)acetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide.* To a suspension of N-(4-fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (0.025 g, 0.040 mmol) in dichloromethane (1 mL) was added diisopropyl ethylamine (0.028 mL, 0.160 mmol) and the solution was stirred at room temperature for 5 min. Added to this was dimethylsulfamoyl chloride (4.26 μL, 0.040 mmol) and the solution was stirred at room temperature for 16 h. The reaction solution was concentrated and purified by preparative HPLC: Solvent A = 10% methanol/90% H₂O/0.1% trifluoroacetic acid; Solvent B = 90% methanol/10% H₂O/0.1% trifluoroacetic acid; Start % B = 20; Final % B = 100; Gradient time = 20 min; Stop time = 30 min; Column = XTERRA[®]


20

25

19 x 50 mm, C18, 5 μ m to give the title compound as a white solid (12.2 mg, 45% yield). LCMS (M+H) calcd for C₂₇H₃₅FN₇O₇S: 620.23; found: 620.74. ¹H NMR (500 MHz, CDCl₃) δ : 12.00 (1 H, s), 8.45 (1 H, s), 8.39 (1 H, t, *J* = 5.95 Hz), 7.36 (2 H, dd, *J* = 8.55, 5.49 Hz), 7.01 (2 H, t, *J* = 8.55 Hz), 4.55 (2 H, d, *J* = 6.10 Hz), 4.16 (2 H, d, *J* = 3.66 Hz), 4.02 - 4.09 (2 H, m), 3.52 - 3.61 (2 H, m), 3.26 - 3.36 (2 H, m), 3.18 - 3.26 (2 H, m), 2.84 (6 H, s), 2.58 (2 H, ddd, *J* = 14.19, 9.16, 5.65 Hz), 2.50 (1 H, brs), 1.86 - 2.11 (4 H, m), 1.66 - 1.80 (2 H, m).


Example 146

10



10-(((4-((3,5-Dimethyl-4-isoxazolyl)sulfonyl)-1-piperazinyl)(oxoacetyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Purple solid (4.5 mg, 16% yield). LCMS (M+H) calcd for C₃₁H₃₃FN₇O₆: 672.22; found: 672.89. ¹H NMR (500 MHz, CDCl₃) δ : 11.99 (1 H, brs), 8.74 (1 H, brs), 8.22 (1 H, brs), 7.35 (2 H, dd, *J* = 8.09, 5.34 Hz), 7.00 (2 H, t, *J* = 8.55 Hz), 4.55 (2 H, d, *J* = 5.19 Hz), 4.03 - 4.31 (4 H, m), 3.63 (2 H, brs), 3.05 - 3.24 (4 H, m), 2.54 - 2.75 (5 H, m), 2.50 (1 H, brs), 2.40 (3 H, s), 1.97 (4 H, brs), 1.67 - 1.79 (2 H, m).


Example 147



5 *Methyl 3-((4-(((2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)amino)(oxo)acetyl)-1-piperazinyl)sulfonyl)-2-thiophenecarboxylate.* White solid (6.0 mg, 19% yield).
(M+H) calcd for C₃₁H₃₄FN₆O₉S₂: 717.18; found: 717.36. ¹H NMR (500 MHz, CDCl₃) δ: 11.98 (1 H, s), 8.28 - 8.53 (2 H, m), 7.40 - 7.60 (2 H, m), 7.34 (2 H, dd, *J*= 8.39, 5.34 Hz), 6.99 (2 H, t, *J*= 8.70 Hz), 4.53 (2 H, d, *J*= 6.41 Hz), 4.15 (2 H, d, *J*= 3.66 Hz), 3.99 - 4.10 (2 H, m), 3.92 (3 H, s), 3.51 - 3.74 (2 H, m), 3.38 (4 H, dt, *J*= 18.08, 5.00 Hz), 2.39 - 2.71 (3 H, m), 1.84 - 2.12 (4 H, m), 1.63 - 1.79 (2 H, m).

Example 148


15



*N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-((1-methyl-1*H*-imidazol-4-*y*l)sulfonyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* White solid (5.0 mg, 19% yield).
 (M+H) calcd for C₂₉H₃₄FN₈O₇S: 657.22; found: 657.27. ¹H NMR (400 MHz, CDCl₃) δ: 11.95 (1 H, brs), 8.03 - 8.67 (2 H, m), 7.44 (2 H, d, *J* = 17.32 Hz), 7.26 - 7.39 (2 H, m), 6.81 - 7.10 (2 H, m), 4.24 - 4.66 (2 H, m), 3.83 - 4.27 (5 H, m), 3.32 - 3.86 (6 H, m), 3.24 (2 H, brs), 2.27 - 2.75 (3 H, m), 1.81 - 2.16 (4 H, m), 1.69 (2 H, d, *J* = 4.52 Hz).

10

Example 149



N-(4-Fluorobenzyl)-3-hydroxy-10-(((4-((4-(4-

15 *morpholinylsulfonyl)phenyl)sulfonyl)-1-piperazinyl)(oxo)acetyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.* To a mixture N-(4-fluorobenzyl)-3-hydroxy-4-oxo-10-((oxo(1-piperazinyl)acetyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide (25 mg, 0.040 mmol) in dichloromethane (1 mL) was added diisopropyl ethylamine (0.028 mL, 0.160 mmol) followed by 4-(morpholinylsulfonyl)benzene-1-sulfonyl chloride (13.00 mg, 0.040 mmol) and the resulting solution stirred at room temperature for 24 h. The mixture was treated with sodium ethoxide (0.015 mL, 0.040 mmol) and stirred at room temperature for 1 h. The reaction was diluted with water and


dichloromethane. The aqueous phase was acidified with 1N HCl and the product extracted into the organic phase. The organic phase was concentrated and the residue was triturated with methanol. The white solids were collected by filtration to give the title compound (2.2 mg, 7% yield). (M+H) calcd for C₃₅H₄₁FN₇O₁₀S₂: 802.23; 5 found: 802.84. ¹H NMR (400 MHz, CDCl₃) δ: 11.94 (1 H, s), 8.70 (1 H, s), 8.17 (1 H, t, J = 6.02 Hz), 7.91 (4 H, s), 7.31 (2 H, dd, J = 8.41, 5.40 Hz), 6.96 (2 H, t, J = 8.66 Hz), 4.51 (2 H, d, J = 6.02 Hz), 4.04 - 4.22 (4 H, m), 3.67 - 3.80 (4 H, m), 3.44 - 3.65 (2 H, m), 2.88 - 3.20 (8 H, m), 2.50 - 2.67 (2 H, m), 2.46 (1 H, brs), 1.79 - 2.06 (4 H, m), 1.58 - 1.77 (2 H, m).

10

Examples 150 through 155 can be synthesized according to Scheme XVII

Example 150

15




10-(Dimethylamino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (50 mg, 0.134 mmol) in 1,2-dichloroethane (2 mL) was added acetic acid (0.038 mL, 0.671 mmol) and aq. formaldehyde (0.022 mL, 0.295 mmol) and the resulting mixture was stirred at room temperature for 3 h. Sodium triacetoxyborohydride (71.1 mg, 0.336 mmol) was then added and the resulting mixture stirred at room temperature overnight. Sat. NaHCO₃ was added and the mixture was extracted 3 times with chloroform, dried and filtered. The crude material was then purified by preparative HPLC to afford the title compound as a light yellow solid (7 mg, 13% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.17 (brs, 1H), 9.78 (brs, 1H), 7.36 (dd, 2H, J = 8.55 & 5.49 Hz), 6.96-7.00 (m, 2H),

4.52 (d, 2H, $J = 6.41$ Hz), 4.12 (d, 2H, $J = 3.36$ Hz), 2.83 (s, 6H), 2.59 (brs, 1H), 2.23-2.31 (m, 2H), 2.11-2.18 (m, 2H), 1.98-2.06 (m, 2H), 1.73-1.82 (m, 2H). LCMS ($M+H$) = 401.16. HPLC purity: > 95%.

5


Example 151



10-*(Benzyl(methyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide*. To a stirred solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (300 mg, 0.734 mmol) in methanol (7 mL), benzaldehyde (0.295 mL, 2.92 mmol), triethylamine (0.102 mL, 0.734 mmol), AcOH (0.420 mL, 7.34 mmol) and NaCNBH₃ (184 mg, 2.94 mmol) was added. The mixture was then stirred at room temperature for 72h. Aqueous formaldehyde (0.109 mL, 1.468 mmol) was then added to the reaction and the mixture stirred at room temperature for 4 h. The solvent was removed and the crude product purified by preparative HPLC to afford the title compound as the corresponding trifluoroacetic acid salt (280 mg, 65% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.6 (1 H, brs), 10.1 (1 H, brs), 7.4 - 7.5 (2 H, m), 7.3 (1 H, t, $J=7.63$ Hz), 7.3 (2 H, d, $J=7.63$ Hz), 7.0 - 7.1 (3 H, m), 4.8 (1 H, dd, $J=14.34, 6.71$ Hz), 4.4 - 4.5 (3 H, m), 3.8 (1 H, d, $J=15.56$ Hz), 3.7 (1 H, d, $J=12.82$ Hz), 2.8 (3 H, s), 2.6 (1 H, brs), 2.4 - 2.4 (2 H, m), 2.3 - 2.4 (1 H, m), 2.2 - 2.3 (1 H, m), 2.1 - 2.1 (1 H, m), 2.0 - 2.0 (2 H, m), 1.5 - 1.6 (1 H, m). LCMS ($M+H$) = 477.49.


25

Example 152



5 *N*-(4-*Fluorobenzyl*)-3-*hydroxy*-10-(*methylamino*)-4-*oxo*-4,6,7,8,9,10-
hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide. To a solution of 10-
(benzyl(methyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-
hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide (255 mg, 0.432 mmol)
in methanol (2 mL) was added Pd/C (32.2 mg, 0.030 mmol) and the mixture was
10 stirred under hydrogen atmosphere (balloon) for 1 h. The catalyst was then filtered
off, and the filtrate was evaporated under reduced pressure to afford the title
compound as a white solid (200 mg, 93% yield). ^1H NMR (300 MHz, DMSO-*d*₆) δ :
12.3 (1 H, s), 9.5 (1 H, s), 7.3 - 7.4 (2 H, m), 7.2 - 7.2 (2 H, m), 4.5 (2 H, d, *J*=6.22
Hz), 4.0 (2 H, d, *J*=3.29 Hz), 2.6 (3 H, brs), 2.0 - 2.1 (4 H, m), 1.7 - 1.9 (5 H, m).
15 LCMS (M+H) = 387.17.

Example 153




20 *N*-(4-*Fluorobenzyl*)-3-*hydroxy*-10-(*methyl(oxo(1-pyrrolidinyl)acetyl)amino*)-
4-*oxo*-4,6,7,8,9,10-*hexahydro*-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide.
To a stirred solution of N-(4-fluorobenzyl)-3-hydroxy-10-(methylamino)-4-oxo-

4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (50 mg, 0.100 mmol) in (DMF (2 mL) was added 2-oxo-2-(pyrrolidin-1-yl)acetic acid (57.2 mg, 0.400 mmol), diisopropyl-ethylamine (0.105 mL, 0.599 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (152 mg, 0.400 mmol) and 4-(dimethylamino)pyridine (DMAP) (1.221 mg, 9.99 µmol) and the resulting mixture was stirred at room temperature for 3h. The crude was then purified by preparative HPLC and recrystallized from methanol to afford the title compound as a glassy white crystals (20 mg, 40% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.1 (1 H, s), 9.6 (1 H, brs), 7.4 - 7.4 (2 H, m), 7.0 - 7.0 (2 H, m), 4.8 - 4.9 (1 H, m), 4.6 - 4.7 (1 H, m), 4.4 - 4.5 (1 H, m), 3.6 - 3.7 (2 H, m), 3.5 (2 H, t, J=6.71 Hz), 3.3 - 3.4 (2 H, m), 3.0 (3 H, s), 2.5 (1 H, brs), 1.9 - 2.1 (9 H, m), 1.7 - 1.8 (2 H, m). LCMS (M+H)= 512.24. HPLC purity: > 95%.

Example 154

15




N-(4-Fluorobenzyl)-3-hydroxy-10-(methyl((5-methyl-3-isoxazolyl)carbonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a solution of N-(4-fluorobenzyl)-3-hydroxy-10-(methylamino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (25 mg, 0.061 mmol) in CH₂Cl₂ (2 mL) was added triethylamine (0.051 mL, 0.365 mmol) followed by 5-methylisoxazole-3-carbonyl chloride (17.69 mg, 0.122 mmol) and the resulting mixture was stirred at room temperature. After 16 at room temperature the reaction mixture was concentrated to give crude product which was dissolved in methanol (2mL) and treated with 2M dimethylamine in methanol (0.5 mL). The resulting reaction mixture was stirred at 60 °C for 2 h, then cooled and

purified by preparative HPLC to afford the title compound (5 mg, 17% yield) as an off-white solid. ^1H NMR (500 MHz, CDCl_3) δ : 11.81 (1 H, brs), 8.70 (1 H, d, $J=5.19$ Hz), 7.31 (2 H, dd, $J=8.55, 5.49$ Hz), 6.92 - 7.01 (2 H, m), 6.09 (1 H, s), 4.75 - 4.86 (1 H, m), 4.58 - 4.67 (1 H, m), 4.38 - 4.47 (1 H, m), 3.67 (1 H, d, $J=15.56$ Hz), 3.35 - 5 3.46 (1 H, m), 3.13 (3 H, s), 2.50 (1 H, brs), 2.41 (3 H, s), 2.07 - 2.16 (3 H, m), 1.95 - 2.05 (1 H, m), 1.72 - 1.86 (2 H, m), 1.37 - 1.50 (1 H, m). LCMS ($\text{M}+\text{H}$) = 496.66. HPLC purity: > 95%.

Example 155

10




N-(4-Fluorobenzyl)-3-hydroxy-10-(methyl(methylsulfonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. Off-white solid (2.5 mg, 11% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.94 (1 H, brs), 9.03 (1 H, brs), 7.32 (2 H, dd, $J=8.39, 5.34$ Hz), 6.96 - 7.02 (2 H, m), 4.52 - 4.66 (2 H, m), 4.46 (1 H, d, $J=6.41$ Hz), 4.36 - 4.43 (1 H, m), 3.69 - 3.76 (1 H, m), 3.13 - 3.24 (1 H, m), 2.94 (3 H, s), 2.91 (3 H, s), 2.47 (1 H, brs), 2.04 - 2.20 (2 H, m), 1.93 - 15 2.01 (2 H, m), 1.52 - 1.63 (2 H, m). LCMS ($\text{M}+\text{H}$) = 465.60. HPLC purity: > 95%.

20

Examples 156 through 162 can be prepared according to Scheme XVIII


Example 156



5


N-(2-((4-Fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N',N'*-trimethylethanediamide. To a mixture of methyl 10-(((dimethylamino)(oxo)acetyl)(methyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (3 g, 7.65 mmol) in ethanol (80 mL) was added (4-fluoro-3-methylphenyl)methanamine (2.90 mL, 22.94 mmol) and the mixture was heated at 90 °C for 18 h. The reaction mixture was cooled, concentrated, diluted with ethyl acetate and washed twice with 0.2N NaOH (2 x 100 mL). The organic layer was discarded and the aqueous layer acidified with conc. HCl and extracted with dichloromethane (2 x 150mL), dried (Na_2SO_4), filtered and concentrated. The crude product was recrystallized from methanol to afford the title compound (2.5 g, 66% yield) as a crystalline white solid. ^1H NMR (500 MHz, CDCl_3) δ : 12.13 (1 H, s), 9.56 (1 H, brs), 7.20 - 7.23 (1 H, m), 7.15 - 7.20 (1 H, m), 6.88 - 6.93 (1 H, m), 4.83 (1 H, dd, J =15.56, 7.02 Hz), 4.55 - 4.62 (1 H, m), 4.39 - 4.45 (1 H, m), 3.60 (1 H, d, J =15.56 Hz), 3.33 - 3.47 (2 H, m), 3.01 (3 H, s), 2.99 (3 H, s), 2.97 (3 H, s), 2.49 (1 H, brs), 2.23 (3 H, d, J =1.83 Hz), 1.97 - 2.09 (4 H, m), 1.74 - 1.84 (2 H, m). LCMS (M+H) = 500.69. HPLC purity: > 95%.

Example 157



5 *N*-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N',N'*-trimethylethanediamide. White crystalline solid (27 mg, 40% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.12 (1 H, s), 9.58 (1 H, brs), 7.37 - 7.44 (2 H, m), 6.91 - 7.08 (2 H, m), 4.83 - 4.95 (1 H, m), 4.56 - 4.64 (1 H, m), 4.43 - 4.55 (1 H, m), 3.64 (1 H, d, $J=15.26$ Hz), 3.44 - 3.56 (2 H, m), 10 3.01 (3 H, s), 3.07 (3 H, s), 3.13 (3 H, s), 2.55 (1 H, brs), 2.01 - 2.18 (4 H, m), 1.72 - 1.83 (2 H, m). LCMS ($\text{M}+\text{H}$) = 486.16. HPLC purity: > 95%.

Example 158




15 *N*-(2-((3-Chloro-4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N',N'*-trimethylethanediamide. Crystalline white solid (25 mg, 49% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.00 (1 H, s), 9.63 (1 H, brs), 7.49 (1 H, dd, $J=7.02, 2.14$ Hz), 7.26 - 7.31 (1 H, m), 7.02 - 7.08 (1 H, m), 4.80 - 4.86 (1 H, m), 4.55 - 4.62 (1 H, m), 4.41 - 4.48 (1 H, m), 3.60 (1 H, d, $J=14.95$ Hz), 3.37 - 3.49 (2 H, m), 3.02 (3 H, s), 2.99 (3

H, s), 2.98 (3 H, s), 2.49 (1 H, brs), 1.93 - 2.11 (4 H, m), 1.73 - 1.85 (2 H, m). LCMS (M+H) = 520.64. HPLC purity: > 95%.

Example 159


5



N-(2-((3,4-Difluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N',N'-trimethylethanediamide.

- 10 Crystalline white solid (18 mg, 37% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.00 (1 H, s), 9.63 (1 H, brs), 7.23 - 7.28 (1 H, m), 7.02 - 7.14 (2 H, m), 4.83 (1 H, dd, J =15.26, 7.32 Hz), 4.55 - 4.63 (1 H, m), 4.41 - 4.48 (1 H, m), 3.60 (1 H, d, J =15.26 Hz), 3.36 - 3.47 (1 H, m), 3.02 (3 H, s), 2.99 (3 H, s), 2.98 (3 H, s), 2.49 (1 H, brs), 1.95 - 2.10 (4 H, m), 1.73 - 1.86 (2 H, m), 1.43 - 1.52 (1 H, m). LCMS (M+H) = 504.65. HPLC purity: > 95%.

Example 160




20

N-(2-((4-Fluoro-3-methoxybenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N'*-trimethylethanediamide. Crystalline white solid (18 mg, 36% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.10 (1 H, s), 9.55 (1 H, t, *J*=6.27 Hz), 7.04 (1 H, dd, *J*=8.16, 1.88 Hz), 6.93 - 6.99 (1 H, m), 6.86 - 6.92 (1 H, m), 4.81 (1 H, dd, *J*=15.31, 7.53 Hz), 4.53 - 4.61 (1 H, m), 4.40 - 4.48 (1 H, m), 3.83 (3 H, s), 3.57 (1 H, d, *J*=15.31 Hz), 3.31 - 3.45 (1 H, m), 2.99 (3 H, s), 2.96 (3 H, s), 2.94 (3 H, s), 2.47 (1 H, brs), 1.94 - 2.08 (4 H, m), 1.71 - 1.82 (2 H, m), 1.40 - 1.50 (1 H, m). LCMS (M+H) = 516.64. HPLC purity: > 95%.


10

Example 161



15 *N*-(2-((1,3-Benzodioxol-5-ylmethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N,N'*-trimethylethanediamide. White solid (32 mg, 40% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.14 (1 H, s), 9.58 (1 H, brs), 6.95 (1 H, d, *J*=1.53 Hz), 6.88 (1 H, dd, *J*=7.93, 1.53 Hz), 6.73 (1 H, d, *J*=7.93 Hz), 5.91 (2 H, s), 4.83 (1 H, dd, *J*=14.95, 7.32 Hz), 4.54 - 4.61 (1 H, m), 4.37 (1 H, dd, *J*=14.34, 5.80 Hz), 3.59 (1 H, d, *J*=15.26 Hz), 3.35 - 3.47 (1 H, m), 3.02 (3 H, s), 2.99 (6 H, s), 2.48 (1 H, brs), 1.95 - 2.09 (4 H, m), 1.72 - 1.85 (2 H, m), 1.43 - 1.52 (1 H, m). LCMS (M+H) = 512.12. HPLC purity: > 95%.


Example 162



5 *N*-(2-((2,6-Dibromo-4-fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-
6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N,N',N'*-
trimethylethanediamide. A solution of methyl 10-
(((dimethylamino)(oxo)acetyl)(methyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-
hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (0.10 g, 0.246 mmol),
10 (2,6-dibromo-4-fluoro-3-methylphenyl)methanamine (0.225 g, 0.758 mmol) and
Et₃N (0.279 mL, 2 mmol) in EtOH (5 mL) was heated at reflux for 48 h. Then,
cooled and purified by preparative HPLC to afford the title compound (0.0906 g,
0.138 mmol, 56% yield) as a white solid after crystallization from methanol/H₂O. ¹H
NMR (500 MHz, CDCl₃) δ: 12.18 (1 H, s), 9.08 (1 H, s), 7.32 (1 H, d, *J* = 8.5 Hz),
15 4.96 (2 H, d, *J* = 8.2 Hz), 4.78 - 4.90 (1 H, m), 3.57 - 3.72 (1 H, m), 3.36 - 3.49 (1 H,
m), 2.94 (3 H, s), 2.93 (3 H, s), 2.76 (3 H, s), 2.48 - 2.54 (1 H, m), 2.34 (3 H, d, *J*=
2.1 Hz), 2.00 - 2.10 (4 H, m), 1.71 - 1.87 (2 H, m), 1.45 - 1.53 (1 H, m). LCMS
(M+H) calcd for C₂₅H₂₉Br₂FN₅O₅: 656.05; found: 656.24.

Examples 163 through 172 can be prepared according to Scheme XIX


Example 163



5

N-(4-Fluorobenzyl)-10-(((3*R*)-3-fluoro-1-pyrrolidinyl)(oxo)acetyl)(methyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a mixture of ethyl 10-((3*R*)-3-fluoro-1-pyrrolidinyl)(oxo)acetyl)(methyl)amino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylat (35 mg, 0.063 mmol) in ethanol (2 mL) was added 2M dimethylamine in methanol (0.063 mL, 0.126 mmol) and the resulting mixture was heated at 50 °C for 1 h. (4-Fluorophenyl)methanamine (0.036 mL, 0.316 mmol) followed by triethylamine (0.044 mL, 0.316 mmol) were added and the mixture heated at 90 °C for 18 h. The reaction mixture was then cooled, concentrated and purified by preparative HPLC to afford the title compound (17 mg, 51% yield) as an off-white solid. ¹H NMR (500 MHz, CDCl₃) δ: 11.96 (1 H, brs), 9.45 (1 H, brs), 7.34 - 7.40 (2 H, m), 6.97 (2 H, t, J=8.70 Hz), 5.32 - 5.39 (0.5 H, m), 5.20 - 5.29 (0.5 H, m), 4.76 - 4.86 (1 H, m), 4.56 - 4.67 (1 H, m), 4.40 - 4.52 (1 H, m), 3.86 - 3.93 (0.5 H, m), 3.74 - 3.82 (0.5 H, m), 3.34 - 3.71 (6 H, m), 3.03 (1.2 H, s), 3.02 (1.8 H, s), 2.49 (1 H, brs), 2.29 - 2.41 (1 H, m), 1.99 - 2.09 (4 H, m), 1.76 - 1.83 (2 H, m), 1.44 - 1.51 (1 H, m). LCMS (M+H) = 530.61. HPLC purity: > 95%.


Example 164



5 *N*-(4-*Fluoro*-3-*methylbenzyl*)-10-(((3*R*)-3-*fluoro*-1-*pyrrolidinyl*)(*oxo*)*acetyl*)(*methyl*)*amino*)-3-*hydroxy*-4-*oxo*-4,6,7,8,9,10-*hexahydro*-7,10-*ethanopyrimido*[1,2-*a*]*azepine*-2-*carboxamide*. Off-white solid (17 mg, 50% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.17 (1 H, brs), 9.34 (1 H, brs), 7.14 - 7.23 (2 H, m), 6.91 (1 H, t), 5.32 - 5.39 (1 H, m), 5.20 - 5.29 (1 H, m), 4.75 - 4.86 (1 H, m), 4.54 - 4.63 (1 H, m), 4.37 - 4.47 (1 H, m), 3.87 - 3.93 (1 H, m), 3.75 - 3.82 (1 H, m), 3.34 - 3.71 (6 H, m), 3.03 (1 H, s), 3.02 (1 H, s), 2.49 (1 H, brs), 2.29 - 2.40 (1 H, m), 2.23 (3 H, d, J =1.83 Hz), 1.96 - 2.11 (4 H, m), 1.75 - 1.83 (2 H, m), 1.44 - 1.51 (1 H, m). LCMS ($\text{M}+\text{H}$) = 544.67. HPLC purity: > 95%.

15

Example 165




20 *N*-(2-((4-*Fluorobenzyl*)*carbamoyl*)-3-*hydroxy*-4-*oxo*-6,7,8,9-*tetrahydro*-7,10-*ethanopyrimido*[1,2-*a*]*azepin*-10(4*H*)-*yl*)-*N,N'*-*dimethyl*-*N'*-(*tetrahydro*-2-*furanyl**methyl*)*ethanediamide*. Off-white solid (7 mg, 18% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.18 (1 H, brs), 9.66 (0.4 H, brs), 9.58 (0.6 H, brs), 7.32 - 7.39 (2

H, m), 6.93 - 6.99 (2 H, m), 4.79 - 4.87 (1 H, m), 4.58 - 4.66 (1 H, m), 4.40 - 4.50 (1 H, m), 4.05 - 4.17 (1 H, m), 3.54 - 3.89 (5 H, m), 3.35 - 3.47 (2 H, m), 3.20 (0.5 H, d, *J*=2.75 Hz), 3.17 (0.5 H, d, *J*=2.75 Hz), 3.10 (1 H, s), 3.06 (1 H, s), 3.03 (2 H, s), 2.99 (1 H, s), 2.98 (1 H, s), 2.49 (1 H, brs), 1.84 - 2.08 (7 H, m), 1.42 - 1.58 (2 H, m).

5 LCMS (M+H) = 556.73. HPLC purity: > 95%.

Example 166




10

N-(2-((4-Fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N'-dimethyl-N'-(tetrahydro-2-furanyl methyl)ethanediamide. Off-white solid (12 mg, 31% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.29 (1 H, brs), 9.64 (0.4 H, brs), 9.56 (0.6 H, brs), 7.13 - 7.23 (2 H, m), 6.86 - 6.93 (1 H, m), 4.77 - 4.87 (1 H, m), 4.53 - 4.64 (1 H, m), 4.36 - 4.46 (1 H, m), 4.04 - 4.17 (1 H, m), 3.56 - 3.88 (5 H, m), 3.36 - 3.47 (2 H, m), 3.27 - 3.34 (0.5 H, m), 3.19 (0.5 H, dd, *J*=14.50, 2.59 Hz), 3.11 (1 H, s), 3.07 (1 H, s), 3.03 (2 H, s), 3.00 (1 H, s), 2.98 (1 H, s), 2.49 (1 H, brs), 2.23 (3 H, s), 1.86 - 2.09 (7 H, m), 1.43 - 1.59 (2 H, m). LCMS (M+H) = 570.73. HPLC purity: > 95%.

15 20


Example 167



5 *N*-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N'*-(2-methoxyethyl)-*N,N'*-dimethylethanediamide. Off-white solid (9 mg, 31% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.85 (1 H, brs), 9.53 - 9.66 (1 H, m), 7.30 - 7.37 (2 H, m), 6.91 - 6.98 (2 H, m), 4.75 - 4.86 (1 H, m), 4.55 - 4.64 (1 H, m), 4.38 - 4.48 (1 H, m), 3.70 - 3.81 (0.5 H, m), 3.49 - 3.66 (2.5 H, m), 3.34 - 3.45 (1 H, m), 3.33 (1.6 H, s), 3.30 (1.4 H, s), 3.04 (1.6 H, s), 2.97 (3 H, s), 2.94 (1.4 H, s), 2.47 (1 H, brs), 1.94 - 2.06 (4 H, m), 1.71 - 1.81 (4 H, m), 1.42-1.47 (1 H, m). LCMS (M+H) = 530.64. HPLC purity: > 95%.

15

Example 168




20 *N*-(2-((4-Fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-*N'*-(2-methoxyethyl)-*N,N'*-dimethylethanediamide. Off-white solid (14 mg, 48% yield). ¹H NMR (500 MHz, CDCl₃) δ: 11.81 (1 H, brs), 9.50 - 9.62 (1 H, m), 7.10 - 7.21 (2 H, m), 6.83 - 6.92 (1 H, m), 4.75 - 4.85 (1 H, m), 4.51 - 4.61 (1 H, m), 4.32 - 4.45 (1 H, m), 3.71 - 3.82 (1 H, m).

(0.5 H, m), 3.50 - 3.67 (2.5 H, m), 3.34 - 3.45 (1 H, m), 3.33 (1.2 H, s), 3.30 (1.8 H, s), 3.04 (1.6 H, s), 2.98 (3 H, s), 2.95 (1.4 H, s), 2.46 (1 H, brs), 2.21 (3 H, s), 1.94 - 2.07 (5 H, m), 1.69 - 1.80 (3 H, m), 1.37 - 1.51 (1 H, m). LCMS (M+H) = 544.67. HPLC purity: > 95%.


5

Example 169




10 *N-((1,3-Dimethyl-1H-pyrazol-4-yl)methyl)-N'-(2-((4-*
fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-
ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N'-dimethylethanediamide. Off-white
 solid (16 mg, 42% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.05 (1 H, brs), 9.49 -
 9.71 (1 H, m), 7.42 (0.5 H, s), 7.31 - 7.39 (2 H, m), 7.11 (0.5 H, s), 6.89 - 6.97 (2 H,
 m), 4.81 (1 H, dd, $J=15.43, 7.40$ Hz), 4.62 (1 H, dd, $J=14.68, 6.90$ Hz), 4.28 - 4.51
 (2.5 H, m), 4.04 (0.5 H, d, $J=15.31$ Hz), 3.77 (1.4 H, s), 3.76 (1.6 H, s), 3.57 (1 H, d,
 $J=14.56$ Hz), 3.31 - 3.47 (1 H, m), 3.02 (1.4 H, s), 2.94 (1.6 H, s), 2.86 (1.6 H, s),
 2.80 (1.4 H, s), 2.47 (1 H, brs), 2.22 (1.4 H, s), 2.19 (1.6 H, s), 1.93 - 2.08 (4 H, m),
 1.76 - 1.81 (2 H, m), 1.41 - 1.53 (1 H, m). LCMS (M+H) = 580.76. HPLC purity: >
 15 95%.
 20 95%.

Example 170



5 *N-((1,3-Dimethyl-1H-pyrazol-4-yl)methyl)-N'-(2-((4-fluoro-3-*
methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-
ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N'-dimethylethanediamide. Off-white
10 solid (17 mg, 43% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.96 (1 H, brs), 9.49 -
 9.60 (1 H, m), 7.47 (0.4 H, s), 7.13 - 7.21 (2 H, m), 7.11 (0.6 H, s), 6.83 - 6.91 (1 H,
 m), 4.81 (1 H, dd, $J=15.43, 7.65$ Hz), 4.54 - 4.62 (1 H, m), 4.26 - 4.47 (2.5 H, m),
 4.02 (0.5 H, d, $J=15.31$ Hz), 3.80 (1 H, s), 3.78 (2 H, s), 3.57 (1 H, d, $J=15.31$ Hz),
 3.31 - 3.44 (1 H, m), 3.02 (1 H, s), 2.94 (2 H, s), 2.86 (2 H, s), 2.81 (1 H, s), 2.47 (1
 H, brs), 2.24 (1 H, s), 2.21 (2 H, s), 2.19 (3 H, d, $J=1.76$ Hz), 1.95 - 2.07 (4 H, m),
 1.69 - 1.83 (2 H, m), 1.38 - 1.55 (1 H, m). LCMS ($\text{M}+\text{H}$) = 594.80. HPLC purity: >
15 95%.


Example 171



*N-((1,5-Dimethyl-1*H*-pyrazol-4-yl)methyl)-N'-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-N,N'-dimethylethanediamide.* Off-white solid (8 mg, 24% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.59 (1 H, brs), 9.56 (1 H, brs), 7.52 (0.3 H, s), 7.49 (0.7 H, s), 7.36 (2 H, dd, $J=8.39, 5.34$ Hz), 6.93 - 6.99 (2 H, m), 4.80 - 4.87 (1 H, m), 4.60 - 4.69 (1 H, m), 4.33 - 4.52 (3 H, m), 3.84 (1 H, s), 3.82 (2 H, s), 3.34 - 3.48 (2 H, m), 3.03 (1 H, s), 2.95 (2 H, s), 2.89 (2 H, s), 2.81 (1 H, s), 2.50 (1 H, brs), 2.28 (1 H, s), 2.17 (2 H, s), 1.96 - 2.10 (4 H, m), 1.71 - 1.84 (2 H, m), 1.44 - 1.55 (1 H, m). LCMS ($\text{M}+\text{H}$) = 580.17. HPLC purity: > 95%.

10

Example 172




15 *N-((1,5-Dimethyl-1*H*-pyrazol-4-yl)methyl)-N'-(2-((4-fluoro-3-methylbenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-N,N'-dimethylethanediamide.* Off-white solid (8 mg, 23% yield). ^1H NMR (500 MHz, CDCl_3) δ : 11.87 (1 H, brs), 9.50 - 9.57 (1 H, m), 7.51 (1 H, s), 7.48 (1 H, s), 7.13 - 7.22 (2 H, m), 6.90 (1 H, t, $J=8.85$ Hz), 4.79 - 4.87 (1 H, m), 4.56 - 4.65 (1 H, m), 4.32 - 4.48 (3 H, m), 3.83 (1 H, s), 3.81 (2 H, s), 3.34 - 3.47 (2 H, m), 3.03 (1 H, s), 2.96 (2 H, s), 2.89 (2 H, s), 2.82 (1 H, s), 2.49 (1 H, brs), 2.28 (1 H, s), 2.22 (3 H, d, $J=1.83$ Hz), 2.17 (2 H, s), 1.96 - 2.10 (4 H, m), 1.71 - 1.84 (2 H, m), 1.43 - 1.54 (1 H, m). LCMS ($\text{M}+\text{H}$) = 594.23. HPLC purity: > 95%.

20

25

Example 173 can be prepared according to Scheme XX.

Example 173




5

N-Ethyl-*N*-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4*H*)-yl)-*N*',*N*'-dimethylethanediamide. To a stirred precooled (0 °C) solution of 2-(dimethylamino)-10 oxoacetic acid (101 mg, 0.866 mmol) in CH₂Cl₂ (5 mL) was added oxalyl chloride (0.083 mL, 0.953 mmol) and 2 drops of DMF and the resulting solution stirred for 1 h at 0 °C, then 2 h at room temp. The mixture was then concentrated and dried under high vacuum to give crude acid chloride which was diluted with CH₂Cl₂ (3 mL) and added to a pre-stirred solution of ethyl 10-(ethylamino)-4-oxo-3-((phenylcarbonyl)oxy)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxylate (100 mg, 0.216 mmol) and diisopropyl-ethylamine (0.165 mL, 0.944 mmol) in CH₂Cl₂ (5 mL). The resulting solution was stirred at room temperature for 16 h. The reaction mixture was poured into sat. NaHCO₃ and extracted with dichloromethane (50 x 3), then dried (Na₂SO₄) and concentrated to give a yellow oil, which was diluted with ethanol (2 mL) and treated with 2M dimethylamine (0.095 mL, 0.191 mmol). The resulting mixture was heated at 50 °C for 1 h. (4-Fluorophenyl)methanamine (0.054 mL, 0.477 mmol) followed by triethylamine (0.066 mL, 0.477 mmol) was then added and the mixture heated at 90 °C for 18 h. The reaction mixture was cooled, concentrated and purified by preparative HPLC to afford the title compound (5 mg, 11% yield) as an off-white solid. ¹H NMR (500 MHz, CDCl₃) δ: 12.02 (1 H, brs), 9.33 (1 H, s), 7.35 - 7.38 (2 H, m), 6.95 - 6.99 (2 H, m), 4.81 (1 H, dd, J=15.26, 7.63 Hz), 4.66 (1 H, dd, J=14.34, 7.02 Hz), 4.42 (1 H, dd, J=14.65, 5.80 Hz), 3.53 - 3.62 (2 H, m), 3.39 - 3.47 (1 H, m), 3.26 - 3.36 (1 H, m),

2.98 (3 H, s), 2.97 (3 H, s), 2.49 (1 H, brs), 1.94 - 2.17 (4 H, m), 1.77 - 1.84 (2 H, m), 1.44 - 1.52 (1 H, m), 1.23 (3 H, t, $J=7.02$ Hz). LCMS (M+H) = 500.08. HPLC purity: > 95%.

5 Examples 174 through 176 can be prepared according to Scheme XXI


Example 174



10


N-(4-Fluorobenzyl)-3-hydroxy-10-((2-hydroxyethyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide. To a suspension of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (1.00 g, 2.446 mmol) 15 in methanol (40 mL) was added Et₃N (0.341 mL, 2.446 mmol) and the mixture was stirred at room temperature for 30 min. Added to this were glycolaldehyde dimer (0.323 g, 2.69 mmol), acetic acid (0.140 mL, 2.446 mmol) and sodium cyanoborohydride (0.307 g, 4.89 mmol). The suspension dissolved and the resulting solution was stirred at room temperature for 2.5 h then concentrated and purified by preparative HPLC: Solvent A = 10% methanol/90% H₂O/0.1% trifluoroacetic acid; 20 Solvent B = 90% methanol/10% H₂O/0.1% trifluoroacetic acid; Start % B = 10; Final % B = 100; Gradient time = 20 min; Stop time = 20min; Column = Sunfire 30 x 100 mm, C18, 5μm (product elutes at 12.5 min) to give the title compound as a trifluoroacetic acid salt as a white solid (1.0372 g, 80% yield). LCMS (M+H) calcd for C₂₁H₂₆FN₄O₄: 417.19; found: 417.19. ¹H NMR (500 MHz, MeOD) δ: 7.37 - 7.51 (2 H, m), 7.01 - 7.14 (2 H, m), 4.56 (2 H, s), 4.16 (2 H, d, $J=3.97$ Hz), 3.82 - 3.92 (2 H, m), 3.23 - 3.30 (2 H, m), 2.58 (1 H, brs), 2.27 - 2.38 (2 H, m), 2.16 - 2.26 (2 H, m), 1.95 - 2.11 (2 H, m), 1.85 (2 H, ddd, $J=13.96, 7.32, 7.10$ Hz).

Example 175



5 *N*-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-*a*]azepin-10(4*H*)-yl)-*N*-(2-hydroxyethyl)-*N'*,*N'*-dimethylethanediamide. A solution of *N*-(4-fluorobenzyl)-3-hydroxy-10-((2-hydroxyethyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-*a*]azepine-2-carboxamide (0.3358 g, 0.633 mmol) in CH₂Cl₂ (10 mL) with
10 diisopropyl ethylamine (0.553 mL, 3.17 mmol) was cooled to 0 °C. Methyl 2-chloro-2-oxoacetate (0.233 mL, 2.53 mmol) was added and the yellow solution stirred for 5 h, gradually warming to room temperature. The reaction was quenched with water and the organic phase was washed with 1N HCl, dried (Na₂SO₄), filtered and concentrated to give a yellow foam which was dissolved in hot ethanol (10 mL) and treated with 2M dimethylamine (1.583 mL, 3.17 mmol). The resulting yellow
15 solution was stirred in a sealed tube at 85 °C for 18h. The solution was concentrated and purified by preparative HPLC to give the title compound as a white solid (0.1692 g, 52% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.20 (1 H, brs), 9.71 (1 H, brs), 7.34 (2 H, dd, *J*=8.55, 5.49 Hz), 6.98 (2 H, t, *J*=8.70 Hz), 4.69 (1 H, dd, *J*=15.56, 7.63 Hz), 4.50 - 4.57 (1 H, m), 4.37 - 4.48 (2 H, m), 3.84 - 3.91 (1 H, m), 3.75 - 3.81 (1 H, m), 3.56 - 3.70 (3 H, m), 3.41 - 3.51 (1 H, m), 2.97 (3 H, s), 2.96 (3 H, s), 2.49 (1 H, brs), 1.98 - 2.14 (3 H, m), 1.71 - 1.91 (3 H, m), 1.38 - 1.47 (1 H, m). LCMS (M+H)
20 = 516.07. HPLC purity: > 95%.

Example 176



5 *2-((2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)amino)ethyl (dimethylamino)(oxo)acetate.*

To a stirred solution of N-(4-fluorobenzyl)-3-hydroxy-10-((2-hydroxyethyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (90 mg, 0.170 mmol) in DMF (3 mL) was added 2-(dimethylamino)-2-oxoacetic acid


10 (59.6 mg, 0.509 mmol), diisopropyl-ethylamine (0.178 mL, 1.018 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (194 mg, 0.509 mmol) and 4-(dimethylamino)pyridine (DMAP) (4.15 mg, 0.034 mmol) and the resulting mixture was stirred at room temperature for 16h, purified by preparative HPLC and crystallized from methanol/ water to afford the title

15 compound (37 mg, 42% yield) as a crystalline white solid. ¹H NMR (500 MHz, CDCl₃) δ: 12.50 (1 H, brs), 9.46 (1 H, brs), 7.33 - 7.38 (2 H, m), 6.97 - 7.01 (2 H, m), 4.55 (2 H, d, *J*=6.41 Hz), 4.43 - 4.48 (2 H, m), 4.10 (2 H, d, *J*=3.97 Hz), 3.28 (2 H, t, *J*=5.34 Hz), 3.06 (3 H, s), 2.96 (3 H, s), 2.56 (1 H, brs), 2.30 - 2.37 (2 H, m), 2.15 - 2.23 (2 H, m), 1.95 - 2.02 (2 H, m), 1.70 - 1.79 (2 H, m). LCMS (M+H) = 516.32.

20 HPLC purity: > 95%.

Example 177 can be synthesized according to Scheme XXII

Example 177




5

Ethyl N-(2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)glycinate. A mixture of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide (0.10 g, 0.245 mmol) and triethylamine (0.251 mL, 1.8 mmol) in 1,2-dichloroethane (10 mL) was stirred at room temperature for 15 min. Added to this was ethyl 2-oxoacetate (0.145 mL, 0.734 mmol) (50% in toluene) and the mixture stirred at 70 °C for 16 h. The mixture was cooled to room temperature. Acetic acid (0.2 mL, 3.49 mmol) followed by sodium cyanoborohydride (0.061 g, 0.978 mmol) were added and mixture stirred at room temperature for 6 h. The mixture was concentrated and the residue was purified by preparative HPLC: Solvent A = 10% methanol/90% H₂O/0.1% trifluoroacetic acid; Solvent B = 90% methanol/10% H₂O/0.1% trifluoroacetic acid; Start % B = 20; Final % B = 100; Gradient time = 20 min; Stop time = 20 min; Column = Waters-Atlantis 30 x 100 mm, C18, 5μm (product eluted at 9.0 min) to give the title compound as a trifluoroacetic acid salt as a white foam (0.0301 g, 22% yield). ¹H NMR (500 MHz, CDCl₃) δ: 12.23 (1 H, brs), 9.07 (1 H, brs), 8.69 (1 H, t, *J* = 6.26 Hz), 7.33 (2 H, dd, *J* = 8.24, 5.49 Hz), 6.98 (2 H, t, *J* = 8.55 Hz), 4.51 (2 H, d, *J* = 6.41 Hz), 3.98 - 4.07 (4 H, m), 3.91 (2 H, s), 2.54 (1 H, brs), 2.36 - 2.51 (2 H, m), 2.18 - 2.35 (2 H, m), 1.95 - 2.08 (2 H, m), 1.64 - 1.79 (2 H, m), 1.20 (3 H, t, *J* = 7.17 Hz) LCMS (M+H) calcd for C₂₃H₂₈FN₄O₅: 459.20; found: 459.04.

Examples 178 through 180 can be prepared according to Scheme XXIII

Example 178


5



Carbamic acid, [2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-7-methyl-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-phenylmethyl ester. Yield: 2% (white solid; C-18 prep-HPLC purified). HPLC:
10 retention time = 2.73 min. (area percent = 100%). LCMS: m/z 520.93 (M+H). ¹H NMR (500 MHz, CHLOROFORM-d) δ ppm 11.89 (1 H, brs) 7.60 (1 H, brs) 7.33 (3 H, brs) 7.18 (3 H, brs) 6.96 (3 H, brs) 4.91 (2 H, s) 4.41 (2 H, d, J=6.10 Hz) 3.92 (2 H, brs) 2.44 (2 H, brs) 2.23 (1 H, brs) 1.98 - 2.08 (2 H, m) 1.72 (2 H, brs) 1.58 (2 H, brs) 1.11 (3 H, s).


15

Example 179



20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-amino-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-7-methyl-4-oxo-.* Yield: 77 % (white solid). HPLC: retention time = 1.43 min (area percent = 100%). LCMS: m/z 386.90 (M+H).

Example 180




5 *Ethanediamide, N'-[2-[[[4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-7-methyl-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-*. Yield: 30 % (white crystalline solid; purified by preparative HPLC, C-18) HPLC: retention time = 8.32 min. LCMS: m/z 485.93 (M+H)¹H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 11.95 (1 H, s) 8.59 (1 H, t, *J*=5.80 Hz) 8.09 (1 H, s) 10 7.36 (2 H, dd, *J*=8.55, 5.49 Hz) 6.93 - 7.02 (2 H, m) 4.55 (2 H, d, *J*=6.41 Hz) 3.93 (2 H, s) 3.25 - 3.29 (3 H, m) 2.87 - 2.93 (3 H, m) 2.52 (2 H, ddd, *J*=14.50, 9.31, 5.49 Hz) 2.04 - 2.11 (2 H, m) 1.74 - 1.79 (2 H, m) 1.58 - 1.67 (2 H, m) 1.12 (3 H, s).

Examples 181 through 185 can be prepared according to Scheme XXIV

15

Example 181




20 *Benzyl ((7S,10R,11R)-2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocin-11-yl)carbamate*. A solution of ethyl (7S,10R)-11-(((benzyloxy)carbonyl)amino)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocine-2-carboxylate (1.4595 g, 3.11 mmol), (4-

fluorophenyl)methanamine (1.143 mL, 10.00 mmol) and Et₃N (1.394 mL, 10 mmol) in EtOH (25 mL) was heated at reflux for 14 h. the reaction mixture was then cooled, diluted with Et₂O (100 mL), washed with 1N aq. HCl (1 x 25 mL), water (2 x 25 mL) followed by brine (25 mL), then dried (Na₂SO₄), filtered and concentrated to give a
5 brown paste. The product was purified by preparative HPLC using a C18-column and 40-90% methanol/water (0.1 trifluoroacetic acid) to afford the title compound (0.367 g, 0.669 mmol, 22% yield) as a light brown solid. ¹H NMR (500 MHz, CDCl₃) δ: 12.21 (0.3 H, brs), 12.08 (0.7 H, s), 8.24 (0.3 H, brs), 7.71 (0.7 H, t, J = 6.1 Hz), 7.18 - 7.38 (8 H, m), 6.92 - 7.05 (2 H, m), 4.98 - 5.21 (4 H, m), 4.52 - 4.58 (1
10 H, m), 4.41 - 4.47 (1 H, m), 3.67 (0.7 H, d, J = 15.6 Hz), 3.56 (0.3 H, d, J = 15.6 Hz), 2.08 (0.7 H, t, J = 8.1 Hz), 2.01 - 2.06 (0.3 H, m), 1.73 - 1.87 (2 H, m), 1.40 - 1.49 (1 H, m), 1.37 (2 H, s), 1.28 - 1.35 (1 H, m), 1.25 (1 H, s), 1.00 (2 H, s), 0.99 (2 H, s), 0.95 (1 H, s), 0.92 (1 H, s). LCMS (M+H) calcd for C₃₀H₃₄FN₄O₅: 549.25; found:
549.28. Purity: >96%.

15

Example 182




20 (7S,10R,11R)-11-Amino-N-(4-fluorobenzyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocine-2-carboxamide. A stirred solution of benzyl ((7S,10R,11R)-2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocin-11-yl)carbamate (0.3566 g, 0.650 mmol) and 1N aq. HCl (0.650 mL, 0.650 mmol) in EtOH (25 mL) was sparged with N₂. Then, 10% Pd/C (0.1 g, 0.094 mmol) was added and the flask evacuated then replaced with a H₂ atmosphere three times. The reaction was allowed to stir for 3 h under 1 atm H₂. The reaction mixture was filtered, concentrated and purified by preparative HPLC on aC18 column using 20-50% methanol/water (trace HCl) to the
25

title compound as the corresponding hydrochloride salt (0.2277 g, 0.505 mmol, 78% yield) as a white solid. ^1H NMR (500 MHz, CDCl_3) δ : 12.43 (1 H, brs), 9.56 (1 H, brs), 8.51 (3 H, brs), 7.20 - 7.36 (3 H, m), 6.90 (2 H, brs), 4.90 - 5.01 (1 H, m), 4.65 - 4.76 (1 H, m), 4.35 - 4.53 (2 H, m), 3.53 - 3.64 (1 H, m), 2.04 - 2.12 (1 H, m), 1.76 - 5 1.91 (2 H, m), 1.36 - 1.46 (1 H, m), 1.20 (3 H, brs), 1.10 (3 H, brs), 0.94 (3 H, brs). LCMS ($\text{M}+\text{H}$) calcd for $\text{C}_{22}\text{H}_{28}\text{FN}_4\text{O}_3$: 415.21; found: 415.26. Purity: >97%.

Example 183

10




N'-($(7S,10R,11R)$ -2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocin-11-yl)-N,N-dimethylethanediamide. A solution of benzyl ($(7S,10R,11R)$ -2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4H-7,10-methanopyrimido[1,2-a]azocin-11-yl)carbamate hydrochloride salt (0.0367 g, 0.081 mmol), 2-(dimethylamino)-2-oxoacetic acid (9.53 mg, 0.081 mmol), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.031 g, 0.081 mmol) and diisopropyl-ethylamine (0.014 mL, 0.081 mmol) in DMF (3 mL) was stirred at room temperature for 24 h. After this the crude reaction mixture was purified by preparative-HPLC using methanol/water (0.1% trifluoroacetic acid) to afford the title compound (0.0301 g, 0.059 mmol, 72% yield) as a purple solid. ^1H NMR (500 MHz, CDCl_3) δ : 12.01 (1 H, s), 8.76 (1 H, t, J = 6.0 Hz), 8.21 (1 H, d, J = 9.8 Hz), 7.36 - 7.41 (2 H, m), 6.99 - 20 7.05 (2 H, m), 5.27 (1 H, d, J = 9.8 Hz), 5.18 (1 H, dd, J = 15.6, 8.9 Hz), 4.50 - 4.63 (2 H, m), 3.65 (1 H, d, J = 15.6 Hz), 3.34 (3 H, s), 2.95 (3 H, s), 2.14 (1 H, t, J = 8.2 Hz), 1.83 - 1.94 (1 H, m), 1.47 - 1.52 (1 H, m), 1.41 (1 H, dd, J = 9.5, 5.2 Hz), 1.36

(3 H, s), 1.06 (1 H, dd, 9.8, 4.6 Hz), 1.03 (3 H, s), 0.97 (3 H, s). LCMS (M+H) calcd for C₂₆H₃₃FN₅O₅: 514.25; found: 514.29.


Example 184

5



(7*S*,10*R*,11*R*)-11-(Acetylamino)-N-(4-fluorobenzyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4*H*-7,10-methanopyrimido[1,2-*a*]azocine-2-carboxamide. To a stirred solution of benzyl ((7*S*,10*R*,11*R*)-2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4*H*-7,10-methanopyrimido[1,2-*a*]azocin-11-yl)carbamate hydrochloride salt (0.025 g, 0.055 mmol) and diisopropyl-ethylamine (0.175 mL, 1 mmol) in CH₂Cl₂ (3 mL) was added acetyl chloride (0.036 mL, 0.5 mmol). After 4 h, 2M Me₂NH/methanol (1 mL) was added and stirring continued at room temperature. After 16 h, the reaction mixture was concentrated and purified by preparative-HPLC using methanol/water (0.1% trifluoroacetic acid) to afford the title compound (0.0188 g, 0.038 mmol, 69% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ: 11.99 (1 H, s), 7.56 (1 H, t, *J* = 5.8 Hz), 7.33 (2 H, dd, *J* = 8.5, 5.2 Hz), 7.04 - 7.09 (2 H, m), 6.42 (1 H, d, *J* = 9.2 Hz), 5.52 (1 H, d, *J* = 9.2 Hz), 5.17 (1 H, dd, *J* = 15.4, 8.7 Hz), 4.54 - 4.66 (2 H, m, *J* = 14.8, 14.6, 14.6, 6.0 Hz), 3.70 (1 H, d, *J* = 15.3 Hz), 3.48 (3 H, s, from CH₃OH), 2.09 (1 H, t, *J* = 8.1 Hz), 1.94 (3 H, s), 1.78 - 1.87 (1 H, m), 1.41 - 1.49 (1 H, m), 1.38 (3 H, s), 1.30 - 1.37 (1 H, m), 0.99 (3 H, s), 0.95 - 0.98 (1 H, m), 0.94 (3 H, s). LCMS (M+H) calcd for C₂₄H₃₀FN₄O₄: 457.23; found: 457.27. Purity: >98%.

Example 185




5 (7*S*,10*R*,11*R*)-*N*-(4-Fluorobenzyl)-3-hydroxy-10,12,12-trimethyl-11-((methylsulfonyl)amino)-4-oxo-6,7,8,9,10,11-hexahydro-4*H*-7,10-methanopyrimido[1,2-*a*]azocine-2-carboxamide. To a stirred solution of benzyl ((7*S*,10*R*,11*R*)-2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-10,12,12-trimethyl-4-oxo-6,7,8,9,10,11-hexahydro-4*H*-7,10-methanopyrimido[1,2-*a*]azocin-11-yl)carbamate
10 hydrochloride salt (0.025 g, 0.055 mmol) and diisopropyl-ethylamine (0.175 mL, 1 mmol) in CH₂Cl₂ (3 mL) was added methanesulfonyl chloride (0.039 mL, 0.5 mmol) at room temperature. After 4 h, 2M Me₂NH in methanol (1 mL) was added and stirring continued at room temperature. After 16 h, the reaction mixture was concentrated and purified by preparative-HPLC using methanol/water (0.1% trifluoroacetic acid) to afford the title compound (0.0186 g, 0.038 mmol, 68% yield) as a purple solid after crystallization from methanol/water. ¹H NMR (500 MHz, CDCl₃) δ: 12.39 (1 H, s), 8.60 (1 H, brs), 7.34 (2 H, dd, *J* = 8.5, 5.2 Hz), 7.01 (2 H, t, *J* = 8.7 Hz), 6.07 (1 H, d, *J* = 9.5 Hz), 5.19 (1 H, dd, *J* = 15.6, 8.5 Hz), 4.79 (1 H, d, *J* = 9.8 Hz), 4.52 - 4.65 (2 H, m), 3.63 (1 H, d, *J* = 15.6 Hz), 2.76 (3 H, s), 2.10 (1 H, t, *J* = 8.1 Hz), 1.80 - 1.89 (1 H, m), 1.43 - 1.51 (1 H, m), 1.30 - 1.35 (1 H, m), 1.29 (3 H, s), 1.04 (3 H, s), 1.01 (3 H, s), 0.96 - 1.01 (1 H, m). LCMS (M+H) calcd for C₂₃H₃₀FN₄O₅S: 493.19; found: 493.24. Purity: >98%.

Examples 186 and 187 can be synthesized according to Scheme XXV

Example 186


5



Ethanediamide, N'-[2-[[[4-fluorophenyl]methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-7,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-. LCMS: m/z 472.20 (M+H). ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 11.89 (1 H, brs) 8.16 (1 H, t, *J*=5.19 Hz) 7.65 (1 H, s) 7.32 (2 H, dd, *J*=8.55, 5.19 Hz) 7.02 (2 H, t, *J*=8.70 Hz) 4.58 (1 H, d, *J*=7.02 Hz) 4.50 (1 H, d, *J*=5.80 Hz) 4.08 - 4.14 (1 H, m) 3.97 (1 H, d, *J*=6.10 Hz) 3.21 (3 H, s) 2.92 (3 H, s) 2.71 - 2.81 (1 H, m) 2.61 (1 H, brs) 1.95 - 2.24 (2 H, m) 1.77 - 1.87 (3 H, m) 1.66 - 1.73 (2 H, m). ^{13}C NMR (126 MHz, CHLOROFORM-*d*) δ ppm 168.20, 163.35, 162.50, 161.86, 161.39, 158.31, 150.35, 146.51, 133.42, 129.77, 129.71, 125.38, 115.73, 115.57, 56.19, 50.52, 42.44, 38.22, 37.92, 36.71, 32.04, 31.33, 27.25, 20.04.

Example 187

20




7,11-Methano-4H-pyrimido[1,2-a]azocene-2-carboxamide, 11-(acetylamino)-N-[(4-fluorophenyl)methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-. glassy-white solid (8% yield). ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 10.70 (1 H, brs)

7.71 (1 H, brs) 7.29 (2 H, dd, $J=8.09, 5.34$ Hz) 7.04 (2 H, t, $J=8.55$ Hz) 6.07 (1 H, s)
 4.62 (1 H, dd, $J=14.80, 5.03$ Hz) 4.49 (1 H, dd, $J=14.80, 5.04$ Hz) 4.01 - 4.10 (1 H,
 m) 3.94 (1 H, dd, $J=15.26, 6.10$ Hz) 2.73 (1 H, d, $J=12.51$ Hz) 2.55 (1 H, brs) 2.08 (1
 H, d, $J=12.51$ Hz) 1.93 - 2.02 (1 H, m) 1.86 (3 H, s) 1.78 (2 H, brs) 1.69 (2 H, brs)
 5 1.67 (1 H, d, $J=4.58$ Hz) ^{13}C NMR (126 MHz, CHLOROFORM-d) δ ppm 169.37
 168.27, 163.47, 161.51, 158.36, 151.53, 146.33, 133.19, 133.17, 129.55, 129.48,
 125.28, 115.99, 115.83, 55.72, 50.75, 42.52, 38.13, 32.55, 31.20, 27.24, 23.96,
 20.15,. LCMS (M+H) = 415.20. HPLC purity: >98%.

10 Examples 188 through 196 can be synthesized according to Scheme XXVI


Example 188



15

Ethanediamide, N'-[2-[[[4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-. ^1H NMR (400 MHz, DMSO-D6) δ ppm 1.60 (m, 1 H), 2.00 - 2.12 (m, 4 H), 2.55 (d, $J=11.62$ Hz, 1 H), 2.75 (br s, 1 H), 2.86 (s, 3 H), 2.91 (s, 3 H), 3.79 (m, 2 H), 4.51 (m, 2 H), 7.15 - 7.21 (m, 2 H), 7.35 - 7.42 (m, 2 H), 9.09 (s, 1 H), 9.50 (t, $J=6.44$ Hz, 1 H), 11.84 (s, 1 H).


Example 189



5 *Carbamic acid, [2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-phenylmethyl ester.* LCMS retention time = 1.910 min. on a LCMS fitted with 4.6 x 30 mm, 5-micron Luna C18 column, mobile phase A = 0.5% trifluoroacetic acid in 10:90 acetonitrile/water; mobile phase B = 0.5% trifluoroacetic acid in 90:10 acetonitrile/water; gradient = 0 to 100 % mobile phase B over 2 min. at a flow rate of 10 mL/min.


Example 190

15



15 *7,10-Methanopyrimido[1,2-a]azepine-2-carboxamide, 10-amino-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-.* LCMS retention time = 1.18 min. on a LCMS fitted with 4.6 x 30 mm, 5-micron Luna C18 column, mobile phase A = 0.5% trifluoroacetic acid in 10:90 acetonitrile/water; mobile phase B = 0.5% trifluoroacetic acid in 90:10 acetonitrile/water; gradient = 0 to 100 % mobile phase B over 2 min. at a flow rate of 4 mL/min.


Example 191



5 *7,10-Methanopyrimido[1,2-a]azepine-2-carboxamide, 10-(acetylamino)-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ¹H NMR (500 MHz, CDCl₃) δ ppm 11.92 (1 H, s), 7.56 (1 H, t, *J*=5.19 Hz), 7.28 - 7.36 (2 H, m), 7.02 - 7.09 (2 H, m), 6.41 (1 H, br s), 4.49 - 4.65 (2 H, m), 3.93 - 3.99 (1 H, m), 3.83 - 3.90 (1 H, m), 2.75 - 2.83 (1 H, m), 2.56 (1 H, dd, *J*=11.90, 5.49 Hz), 2.40 (1 H, dt, *J*=12.74, 5.95 Hz), 2.18 - 2.30 (2 H, m), 1.91 - 1.99 (1 H, m), 1.90 (3 H, s), 1.60 - 1.66 (1 H, m). LCMS = 401.19 (M+H). HPLC: >97%.*


10 *15*

Example 192




15 *7,10-Methanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[(methylsulfonyl)amino]-4-oxo-. LCMS retention time = 1.60 min. on a LCMS fitted with 4.6 x 30 mm, 5-micron Luna C18 column, mobile phase A = 0.5% trifluoroacetic acid in 10:90 acetonitrile/water; mobile phase B = 0.5% trifluoroacetic acid in 90:10 acetonitrile/water; gradient = 0 to 100 % mobile phase B over 2 min. at a flow rate of 4 mL/min.*

Example 193



5 *Ethanediamide, N-[2-[[[4-fluorophenyl]methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N'-methyl-*. ^1H NMR (500 MHz, CDCl_3) δ ppm 12.09 (1 H, s), 8.86 (1 H, s), 7.87 (1 H, t, $J=5.65$ Hz), 7.32 - 7.39 (3 H, m), 7.01 - 7.09 (2 H, m), 4.54 - 4.64 (2 H, m), 3.97 - 4.03 (1 H, m), 3.83 - 3.91 (1 H, m), 2.88 (3 H, d, $J=5.19$ Hz), 2.80 - 2.86 (2 H, m),
10 2.53 - 2.62 (1 H, m), 2.24 - 2.35 (1 H, m), 2.03 (1 H, d, $J=10.99$ Hz), 1.93 - 2.01 (1 H, m), 1.65 - 1.72 (1 H, m). LCMS 444.17 ($\text{M}+\text{H}$). HPLC purity: >97%.

Example 194




15

15 *Carbamic acid, [2-[[[3-chloro-4-fluorophenyl]methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-, phenylmethyl ester.* HPLC: retention time = 2.90 min (100%) Start %B = 0%, Final %B = 100%. Gradient Time = 4 min, Total Time = 5 min. Flow Rate = 4 min, Wavelength = 254 nm. Solvent A = 10% CH₃CN - 90% H₂O - 0.1% trifluoroacetic acid. Solvent B = 90% CH₃CN - 10% H₂O - 0.1% trifluoroacetic acid. Column = YMC ODS-AQ 4.6 x 50 mm, 3 micron. LC/MS : 527.26 ($\text{M}+\text{H}$). Start %B = 0%, Final %B = 100%. Gradient Time = 4 min, Total Time = 5 min. Flow Rate = 4 min,

Wavelength = 220 nm. Solvent A = 5% CH₃CN - 95% H₂O - 10 mM Ammonium Acetate. Solvent B = 95% CH₃CN - 5% H₂O - 10 mM Ammonium Acetate. Column = XTERRA® 3.0 x 50 mm, 7μm. ¹H NMR (500 MHz, CDCl₃) δ ppm 11.94 (1 H, br s), 7.82 (1 H, br), 7.26 - 7.43 (5 H, m), 6.92 - 7.16 (3 H, m), 5.96 (1 H, br), 5.00 (2 H, br), 4.50 (1 H, dd, J=15.11, 6.26 Hz), 4.33 (1 H, m), 3.76 - 4.02 (2 H, m), 2.79 (1 H, br), 2.52 (1 H, br), 2.10 - 2.41 (2 H, m), 1.98 (1 H, m), 1.61 (1 H, m).

Example 195


10



7,10-Methanopyrimido[1,2-a]azepine-2-carboxamide, 10-(dimethylamino)-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. HPLC: retention time = 1.49 min (98.7%). Column = YMC ODS-AQ 4.6 x 50 mm, 3 micron. HPLC: retention time = 4.62 min (98.6%). Start %B = 0%, Final %B = 100%. Gradient Time = 23 min, Total Time = 25 min. Flow Rate = 2 min, Wavelength = 254 nm. Solvent A = 10% CH₃CN - 90% H₂O - 0.1% trifluoroacetic acid. Solvent B = 90% CH₃CN - 10% H₂O - 0.1% trifluoroacetic acid. Column = XTERRA® MS C-18, 4.6 x 50 mm, 5 micron. LC/MS: 387.29 (M+H). Column = XTERRA® 3.0 x 50 mm, 7μm. ¹H NMR (500 MHz, MeOD) δ ppm 7.42 - 7.49 (2 H, m), 7.06 - 7.15 (2 H, m), 4.65 - 4.71 (1 H, m), 4.52 - 4.60 (1 H, m), 4.00 - 4.06 (1 H, m), 3.84 - 3.91 (1 H, m), 3.06 (3 H, br), 2.99 (4 H, br), 2.50 - 2.59 (1 H, m), 2.42 - 2.50 (1 H, m), 2.31 - 2.43 (1 H, m), 2.25 (1 H, d, J=10.99 Hz), 2.08 - 2.17 (1 H, m), 1.73 - 1.86 (1 H, m).

25

Example 196




5 *Ethanediamide, N'-[2-[[[4-fluoro-3-methylphenyl]methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-*. ^1H NMR (400 MHz, DMSO-D6) δ ppm 9.03 (s, 1 H), 7.23 (d, 1 H), 7.15 - 7.20 (m, 1 H), 7.08 (t, 1 H), 4.35 - 4.52 (m, 2 H), 3.71 - 3.83 (m, 2 H), 2.91 (s, 3 H), 2.84 (s, 3 H), 2.71 - 2.78 (m, 1 H), 2.20 (s, 3 H), 2.06 (s, 4 H), 1.23 - 1.29 (m, 2 H). LCMS ($^+ \text{ESI}$, $\text{M}+\text{H}^+$) m/z 472.2. HPLC purity: 99%.

Examples 197 through 211 can be synthesized according to Scheme XXVIII


Example 197

15



20 *Carbamic acid, [2-[[[4-fluorophenyl]methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-phenylmethyl ester*. HPLC: retention time = 2.91 min (100%). Column = YMC ODS-AQ 4.6 x 50 mm, 3 micron, retention time = 11.56 min (97.8%). Column = XTERRA[®] MS C-18, 4.6 x 50 mm, 5 micron. LC/MS: 507.10 (M+H). Column = Phenomenex 4.6 x 30 mm, 10 μ .

Example 198




5 *8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 11-amino-N-[(4-fluorophenyl)methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-*. HPLC: retention time = 1.51 min (98.9%). Column = YMC ODS-AQ 4.6 x 50 mm, 3 micron. HPLC: retention time = 3.73 min (99.2%). Column = XTERRA[®] MS C-18, 4.6 x 50 mm, 5 micron. LC/MS: 373.40 (M+H). Column = Phenomenex 4.6 x 30 mm, 10 μ . ¹H NMR (500 MHz, MeOD) δ ppm 7.45 (2 H, br), 7.06 (2 H, br), 5.37 (1 H, br), 4.67 (1 H, br), 3.59 - 3.72 (1 H, m), 3.02 (1 H, br), 2.73 (2 H, br), 2.34 (2 H, br), 2.04 - 2.18 (1 H, m), 1.97 (2 H, br), 1.71 (1 H, br), 1.31 (1 H, br).

10

Example 199


15



20 *Ethanediame, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-*. HPLC: retention time = 2.12 min (100%). Column = YMC ODS-AQ 4.6 x 50 mm, 3 micron. HPLC: retention time = min (100%). Column = XTERRA[®] MS C-18, 4.6 x 50 mm, 5 micron. LC/MS: 742.41 (M+H). Column = Phenomenex 4.6 x 30 mm, 10 μ . ¹H NMR (500 MHz, CDCl₃) δ ppm 12.08 (1 H, s), 8.72 (1 H, t, J=5.80 Hz), 8.27 (1 H, s), 7.36 (2 H, dd, J=8.55, 5.49 Hz), 7.01 (2 H, t, J=8.70 Hz), 5.50 (1 H, dt, J=15.34, 3.78 Hz), 4.49 - 4.60 (2 H, m), 3.49 (1 H, t, J=14.04 Hz), 3.29

(3 H, s), 2.91 (3 H, s), 2.64 (1 H, q, $J=5.90$ Hz), 2.53 - 2.61 (1 H, m), 2.43 - 2.52 (2 H, m), 2.27 - 2.38 (1 H, m), 1.78 - 1.92 (3 H, m), 1.60 (1 H, t, $J=14.04$ Hz). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.22, 163.72, 163.29, 161.86, 161.33, 159.54, 152.06, 146.81, 133.74, 133.71, 129.95, 129.89, 123.78, 115.57, 115.40, 66.66, 44.27, 42.49, 5 40.02, 38.16, 37.19, 36.36, 35.21, 32., 30.24.


Example 200



10


8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 11-[[1,2-dioxo-2-(1-pyrrolidinyl)ethyl]amino]-N-[(4-fluorophenyl)methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo: Off-white crystalline solid (27% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 11.99 (1 H, s), 9.47 (1 H, s), 8.31 (1 H, t, $J=5.80$ Hz), 7.39 (2 H, dd, $J=8.39, 5.34$ Hz), 7.01 (2 H, t, $J=8.70$ Hz), 5.50 (1 H, ddd, $J=15.41, 3.51, 3.36$ Hz), 4.51 - 4.64 (2 H, m), 3.83 - 3.99 (2 H, m), 3.34 - 3.56 (3 H, m), 2.61 - 2.76 (3 H, m), 2.37 - 2.48 (1 H, m), 2.26 - 2.37 (1 H, m), 1.78 - 1.99 (5 H, m), 1.55 - 1.70 (3 H, m). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 167.97, 163.37, 161.42, 160.33, 159.82, 159.48, 152.62, 146.87, 133.38, 133.35, 130.05, 129.99, 123.34, 115.67, 115.50, 20 66.07, 48.55, 47.92, 43.08, 42.63, 40.24, 37.53, 34.18, 32.57, 30.31, 26.85, 23.50. LC/MS ($\text{M}+\text{H}$) = 498.46. HPLC purity >97%.

Example 201



5 *8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 11-[[1,2-dioxo-2-(1-pyrrolidinyl)ethyl]amino]-N-[(4-fluorophenyl)methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-*. White crystalline solid (31% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.08 (1 H, brs), 8.81 (1 H, t, $J=5.95$ Hz), 8.25 (1 H, s), 7.36 (2 H, dd, $J=8.55$, 5.49 Hz), 6.96 - 7.03 (2 H, m), 5.50 (1 H, dt, $J=15.56$, 3.66 Hz), 4.49 - 4.59 (2 H, m),
10 3.62 (2 H, q, $J=7.32$ Hz), 3.44 - 3.52 (1 H, m), 3.23 - 3.37 (2 H, m), 2.64 (1 H, q, $J=5.60$ Hz), 2.52 - 2.61 (1 H, m), 2.43 - 2.52 (2 H, m), 2.26 - 2.39 (1 H, m), 1.79 - 1.93 (3 H, m), 1.60 (1 H, t, $J=13.73$ Hz), 1.25 (3 H, t, $J=6.87$ Hz), 1.13 (2 H, t, $J=7.17$ Hz). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.25, 163.38, 163.27, 162.03, 161.33, 159.56, 152.08, 146.81, 133.76, 133.73, 129.91, 129.84, 123.82, 115.58,
15 115.41, 66.62, 44.27, 43.19, 42.50, 41.48, 40.01, 37.18, 35.22, 32.55, 30.25, 14.73, 12.48. LC/MS ($\text{M}+\text{H}$) = 500.15. HPLC purity >97%.

Example 202




20

8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 11-[[2-(3,3-difluoro-1-azetidinyl)-1,2-dioxoethyl]amino]-N-[{(4-fluorophenyl)methyl]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-. Off-white crystalline solid (21% yield).
¹H NMR (500 MHz, CDCl₃) δ ppm 12.04 (1 H, s), 9.97 (1 H, s), 7.91 (1 H, t, *J*=5.65 Hz), 7.39 (2 H, dd, *J*=8.55, 5.49 Hz), 7.00 - 7.06 (2 H, m), 5.52 (1 H, dt, *J*=15.34, 3.62 Hz), 4.92 (2 H, t, *J*=11.75 Hz), 4.60 (2 H, d, *J*=6.10 Hz), 4.37 (2 H, td, *J*=11.75, 3.66 Hz), 3.49 (1 H, t, *J*=13.89 Hz), 2.84 (1 H, td, *J*=6.49, 5.34 Hz), 2.64 - 2.73 (2 H, m), 2.42 - 2.53 (1 H, m), 2.22 (1 H, ddd, *J*=15.18, 9.00, 8.77 Hz), 1.79 - 1.94 (2 H, m), 1.62 (1 H, t, *J*=14.19 Hz), 1.57 (1 H, brs). ¹³C NMR (126 MHz, CDCl₃) δ ppm 167.72, 163.47, 161.52, 159.53, 159.32, 156.88, 152.48, 147.16, 133.01, 132.99, 130.14, 130.07, 122.89, 117.77, 115.83, 115.65, 115.60, 65.88, 65.64, 65.48, 65.41, 61.56, 61.32, 61.08, 42.88, 42.29, 40.42, 37.80, 33.58, 32.49, 30.32. LC/MS (M+H)⁺ = 520.2. HPLC purity >97%.

15

Example 203




20 8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-(4-fluoro-1-piperidinyl)-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-. Off-white crystalline solid (23% yield).
¹H NMR (500 MHz, CDCl₃) δ ppm 12.11 (1 H, brs), 8.64 - 8.84 (1 H, m), 8.33 (1 H, s), 7.35 (2 H, ddd, *J*=8.62, 5.42, 3.05 Hz), 6.97 - 7.04 (2 H, m), 5.50 (1 H, dt, *J*=15.56, 3.66 Hz), 5.29 (1 H, s), 4.82 - 4.98 (1 H, m), 4.48 - 4.60 (2 H, m), 4.06 - 4.18 (1 H, m), 3.77 - 3.91 (1 H, m), 3.65 - 3.75 (1 H, m), 3.49 (1 H, td, *J*=14.19, 6.41 Hz), 3.30 - 3.44 (1 H, m), 2.61 - 2.69 (1 H, m), 2.48 - 2.59 (2 H, m), 2.38 - 2.48 (1 H, m), 2.25 - 2.38 (1 H, m), 1.79 - 1.99 (6 H, m), 1.60 - 1.67 (1 H, m). ¹³C NMR (126

MHz, CDCl₃) δ ppm 168.26, 168.18, 163.28, 162.59, 162.33, 162.03, 161.43, 161.33, 159.53, 159.51, 152.01, 151.96, 146.90, 146.86, 133.77, 133.74, 133.62, 133.60, 130.00, 129.97, 129.93, 129.90, 123.82, 123.68, 115.62, 115.58, 115.45, 115.41, 87.71, 86.35, 66.74, 66.67, 53.53, 44.47, 44.05, 42.60, 42.53, 42.09, 40.13, 39.99, 5 39.03, 38.99, 38.84, 38.81, 37.38, 37.03, 35.27, 35.22, 32.55, 32.52, 32.02, 31.86, 31.83, 31.66, 30.92, 30.81, 30.75, 30.65, 30.33, 30.17. LC/MS (M+H) = 530.57. HPLC purity >97%.


Example 204

10



Ethanediamide, N-ethyl-N'-(2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N-methyl-. White crystalline solid (11% yield). ¹H NMR (500 MHz, CDCl₃) δ ppm 12.09 (1 H, s), 8.71 - 8.84 (1 H, m), 8.09 - 8.37 (1 H, m), 7.33 - 7.39 (2 H, m), 6.97 - 7.04 (2 H, m), 5.50 (1 H, dd, *J*=15.56, 3.36 Hz), 4.49 - 4.59 (2 H, m), 3.56 - 3.71 (1 H, m), 3.48 (1 H, t, *J*=14.34 Hz), 3.35 (1 H, q, *J*=7.32 Hz), 2.84 - 3.27 (3 H, m), 2.60 - 2.67 (1 H, m), 2.40 - 2.59 (3 H, m), 2.25 - 2.38 (1 H, m), 1.77 - 1.92 (3 H, m), 1.58 - 1.65 (1 H, m), 1.09 - 1.27 (3 H, m). ¹³C NMR (126 MHz, CDCl₃) δ ppm 168.25, 168.20, 163.82, 163.30, 163.27, 163.08, 162.04, 161.80, 161.35, 161.32, 159.55, 152.12, 152.00, 146.81, 133.81, 133.67, 130.00, 129.93, 129.87, 129.81, 123.84, 123.75, 115.57, 115.41, 66.72, 66.55, 45.59, 44.40, 44.12, 43.93, 42.57, 42.44, 40.06, 40.00, 37.24, 37.15, 35.59, 35.33, 35.07, 33.60, 32.55, 30.27, 30.24, 25 14.06, 11.74. LC/MS (M+H) = 486.25. HPLC purity >97%.


Example 205



5 *Ethanediamide, N'-(2-[[(4-fluorophenyl)methyl]amino]carbonyl)-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N-(2-methoxyethyl)-N-methyl-*. White crystalline solid (12% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.09 (1 H, brs), 8.71 - 8.77 (1 H, m), 8.23 - 8.33 (1 H, m), 7.32 - 7.40 (2 H, m), 6.97 - 7.04 (2 H, m), 5.45 - 5.54 (1 H, m), 4.50 - 4.59 (2 H, m), 3.90 (1 H, ddd, $J=14.65, 6.87, 4.43$ Hz), 3.73 - 3.79 (1 H, m), 3.55 - 3.62 (1 H, m), 3.43 - 3.55 (3 H, m), 3.33 (2 H, s), 3.32 - 3.33 (1 H, m), 2.95 (2 H, s), 2.59 - 2.67 (1 H, m), 2.52 - 2.59 (1 H, m), 2.42 - 2.51 (2 H, m), 2.26 - 2.38 (1 H, m), 1.78 - 1.92 (3 H, m), 1.58 - 1.65 (1 H, m). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.22, 168.20, 164.30, 163.72, 163.33, 163.30, 161.75, 161.65, 161.34, 159.55, 152.09, 152.02, 146.85, 146.81, 133.76, 133.73, 129.97, 129.94, 129.91, 129.87, 123.80, 123.74, 115.59, 115.42, 71.39, 70.03, 66.59, 66.57, 59.06, 59.01, 50.27, 48.92, 44.26, 44.14, 42.52, 42.50, 40.06, 40.02, 37.67, 37.24, 37.20, 35.42, 35.21, 35.11, 32.54, 30.28, 30.24. LC/MS ($\text{M}+\text{H}$) = 516.31. HPLC purity >97%.

20


Example 206



Ethanediamide, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N-methyl-N-(1-methylethyl)-. Off-white crystalline solid (9% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.12 (1 H, brs), 8.83 - 9.07 (1 H, m), 7.62 - 8.12 (1 H, m), 7.31 - 7.40 (2 H, m), 6.95 - 7.03 (2 H, m), 5.45 - 5.54 (1 H, m), 4.55 - 4.65 (1 H, m), 4.54 (2 H, d, $J=6.41$ Hz), 3.42 - 3.53 (2 H, m), 2.69 - 3.08 (3 H, m), 2.61 - 2.68 (1 H, m), 2.40 - 2.60 (3 H, m), 2.24 - 2.40 (1 H, m), 1.78 - 1.94 (2 H, m), 1.58 - 1.66 (1 H, m), 1.10 - 1.25 (6 H, m). LC/MS ($\text{M}+\text{H}$) = 500.5. HPLC purity >97%.


10

Example 207




8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3S)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-. Pale pink crystalline solid (18% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 11.97 - 12.04 (1 H, m), 9.55 - 9.90 (1 H, m), 8.11 - 8.23 (1 H, m), 7.37 - 7.43 (2 H, m), 7.03 (2 H, td, $J=8.70, 2.44$ Hz), 5.51 (1 H, dt, $J=15.56, 3.66$ Hz), 5.17 - 5.38 (1 H, m), 4.52 - 4.66 (2 H, m), 4.30 - 4.51 (1 H, m), 3.89 - 4.03 (1 H, m), 3.72 - 3.89 (1 H, m), 3.43 - 3.68 (2 H, m), 2.72 - 2.85 (1 H, m), 2.63 - 2.72 (2 H, m), 2.39 - 2.52 (1 H, m), 2.22 - 2.39 (2 H, m), 1.93 - 2.14 (1 H, m), 1.79 - 1.93 (2 H, m), 1.59 - 1.67 (2 H, m). LC/MS ($\text{M}+\text{H}$) = 516.19. HPLC purity >97%.

Example 208



5 *8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-*. Pale purple crystalline solid (13% yield).
¹H NMR (500 MHz, CDCl₃) δ ppm 12.01 (1 H, brs), 9.55 - 9.89 (1 H, m), 8.11 - 8.24 (1 H, m), 7.37 - 7.43 (2 H, m), 6.97 - 7.06 (2 H, m), 5.52 (1 H, ddd, *J*=15.41, 3.51, 3.36 Hz), 5.17 - 5.38 (1 H, m), 4.52 - 4.66 (2 H, m), 4.31 - 4.51 (1 H, m), 3.89 - 4.04 (1 H, m), 3.73 - 3.89 (1 H, m), 3.41 - 3.69 (2 H, m), 2.62 - 2.87 (3 H, m), 2.39 - 2.52 (1 H, m), 2.22 - 2.40 (2 H, m), 1.93 - 2.14 (1 H, m), 1.78 - 1.93 (2 H, m), 1.58 - 1.67 (2 H, m). LC/MS (M+H) = 516.53. HPLC purity >97%.

15 Example 209




20 *Ethanediamide, N-(2-fluoroethyl)-N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N-methyl*. Pale purple crystalline solid (31% yield). ¹H NMR (500 MHz, CDCl₃) δ ppm 12.06 (1 H, brs), 8.41 - 8.75 (2 H, m), 7.29 - 7.43 (2 H, m), 6.95 - 7.07 (2 H, m), 5.49 (1 H, brs), 4.43 - 4.74 (3 H, m), 3.88 - 4.16 (1 H, m), 3.87 - 4.16 (1 H, m), 3.42 - 3.77 (1 H, m), 3.35 - 3.41 (1 H,

m), 2.96 - 3.04 (2 H, m), 2.27 - 2.71 (4 H, m), 1.71 - 1.96 (2 H, m), 1.58 (4 H, brs). LC/MS (M+H) = 504.18. HPLC purity >97%.

Example 210


5



Acetic acid, [[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]amino]oxo-, methyl ester. Pale purple solid (58% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.07 (1 H, s), 9.66 (1 H, s), 7.69 (1 H, t, $J=5.80$ Hz), 7.37 - 7.41 (2 H, m), 7.02 - 7.08 (2 H, m), 5.53 (1 H, dt, $J=15.49, 3.85$ Hz), 4.61 (2 H, d, $J=6.10$ Hz), 3.83 (3 H, s), 3.47 - 3.55 (1 H, m), 2.90 (1 H, ddd, $J=12.89, 6.79, 1.68$ Hz), 2.66 - 2.75 (2 H, m), 2.48 - 2.59 (1 H, m), 2.22 (1 H, ddd, $J=15.11, 9.00, 8.85$ Hz), 1.86 - 1.95 (1 H, m), 1.77 - 1.85 (1 H, m), 1.49 - 1.56 (2 H, m). LC/MS (M+H) = 458.88. HPLC purity >87%.

Example 211

20



Ethanediamide, N-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N',N'-trimethyl-. White crystalline solid (4% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.35 - 12.54 (1 H, m), 9.72 - 9.91 (1 H, m), 7.33 - 7.41 (2 H, m), 6.91 - 7.00


(2 H, m), 5.46 (1 H, dt, $J=15.03, 3.78$ Hz), 4.44 - 4.64 (2 H, m), 3.46 (1 H, t, $J=13.58$ Hz), 3.01 - 3.06 (3 H, m), 2.98 - 3.01 (3 H, m), 2.94 - 2.98 (3 H, m), 2.59 - 2.67 (1 H, m), 2.42 - 2.53 (1 H, m), 2.29 (1 H, dd, $J=11.60, 6.71$ Hz), 2.08 - 2.20 (1 H, m), 1.78 - 1.90 (3 H, m), 1.57 - 1.63 (2 H, m). LC/MS (M+H) = 486.48. HPLC purity >97%.

5

Examples 212 through 216 can be synthesized according to Scheme XXIX


Example 212

10




Ethanediamide, N'-[2-[[[4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-. White crystalline solid (46% yield). ^1H NMR (500 MHz, CDCl₃) δ ppm 12.11 (1 H, s), 8.71 (1 H, t, $J=5.95$ Hz), 8.26 (1 H, s), 7.13 - 7.21 (2 H, m), 6.94 (1 H, t, $J=8.85$ Hz), 5.50 (1 H, dt, $J=15.26, 3.66$ Hz), 4.50 (2 H, d, $J=6.10$ Hz), 3.48 (1 H, t, $J=13.89$ Hz), 3.28 (3 H, s), 2.91 (3 H, s), 2.64 (1 H, q, $J=5.29$ Hz), 2.50 - 2.60 (1 H, m), 2.42 - 2.50 (2 H, m), 2.27 - 2.39 (1 H, m), 2.25 (3 H, s), 1.78 - 1.93 (3 H, m), 1.57 (1 H, brs). ^{13}C NMR (126 MHz, CDCl₃) δ ppm 168.16, 163.66, 161.81, 161.78, 159.86, 159.57, 152.03, 146.81, 133.39, 133.36, 131.34, 131.30, 127.11, 127.05, 125.07, 124.94, 123.82, 115.18, 115.00, 66.65, 44.23, 42.50, 40.04, 38.19, 37.20, 36.40, 35.17, 32.53, 30.26, 14.66, 14.64. LC/MS (M+H) = 486.28. HPLC purity >97%.

Example 213



5 *Ethanediamide, N'-[2-[[[3-chloro-4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-*. Off-white crystalline solid (62% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 11.97 (1 H, s), 8.86 (1 H, t, $J=5.95$ Hz), 8.09 (1 H, s), 7.44 (1 H, dd, $J=7.02$, 2.14 Hz), 7.26 - 7.29 (1 H, m), 7.09 (1 H, t, $J=8.70$ Hz), 5.50 (1 H, dt, $J=15.56$, 3.81 Hz), 4.45 - 4.56 (2 H, m), 3.44 - 3.54 (1 H, m), 3.28 (3 H, s), 2.94 (3 H, s), 2.64 (1 H, q, $J=5.59$ Hz), 2.48 - 2.60 (2 H, m), 2.40 - 2.47 (1 H, m), 2.26 - 2.37 (1 H, m), 1.81 - 1.94 (3 H, m), 1.59 - 1.66 (1 H, m). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.28, 163.88, 161.88, 159.46, 158.62, 156.64, 152.13, 146.85, 135.14, 135.11, 130.52, 128.15, 128.09, 123.68, 121.13, 120.99, 116.77, 116.60, 66.71, 44.32, 42.09, 15.03, 38.14, 37.19, 36.34, 35.23, 32.55, 30.23. LC/MS ($\text{M}+\text{H}$) = 506.19. HPLC purity >97%.

Example 214




20 *Ethanediamide, N'-[2-[[[3,4-difluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl*. Pale yellow crystalline solid (61% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 11.98 (1 H, s), 8.82 (1 H, t, $J=5.34$ Hz), 8.15 (1 H, s), 7.21 -

7.29 (1 H, m), 7.05 - 7.16 (2 H, m), 5.45 - 5.55 (1 H, m), 4.46 - 4.57 (2 H, m), 3.49 (1 H, t, $J=14.34$ Hz), 3.29 (3 H, s), 2.95 (3 H, s), 2.61 - 2.69 (1 H, m), 2.42 - 2.61 (3 H, m), 2.26 - 2.39 (1 H, m), 1.79 - 1.94 (3 H, m), 1.60 - 1.67 (1 H, m). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.28, 163.88, 161.92, 159.50, 152.11, 151.44, 151.34, 150.93, 5 150.83, 149.46, 149.36, 148.96, 148.86, 146.84, 135.02, 134.98, 134.95, 124.33, 124.30, 124.28, 124.25, 123.69, 117.43, 117.40, 117.29, 117.26, 66.69, 44.32, 42.23, 40.04, 38.18, 37.17, 36.31, 35.24, 32.52, 30.24. LC/MS ($\text{M}+\text{H}$) = 490.26. HPLC purity >95%.


10

Example 215



15 *8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(3-chloro-4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-*. Off-white crystalline solid (54% yield).
 ^1H NMR (500 MHz, CDCl_3) δ ppm 11.84 - 11.90 (1 H, m), 9.49 - 9.90 (1 H, m), 8.17 - 8.31 (1 H, m), 7.46 - 7.54 (1 H, m), 7.27 - 7.33 (1 H, m), 7.08 - 7.14 (1 H, m), 5.51 (1 H, dt, $J=15.49, 3.70$ Hz), 5.18 - 5.39 (1 H, m), 4.48 - 4.65 (2 H, m), 4.30 - 4.48 (1 H, m), 3.56 - 4.05 (3 H, m), 3.46 - 3.55 (1 H, m), 2.64 - 2.86 (3 H, m), 2.39 - 2.52 (1 H, m), 2.23 - 2.40 (2 H, m), 1.79 - 1.94 (3 H, m), 1.60 - 1.67 (2 H, m). LC/MS ($\text{M}+\text{H}$) = 550.23. HPLC purity >95%.


Example 216



5 *8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-*. White crystalline solid (44% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.02 - 12.07 (1 H, m), 9.54 - 9.89 (1 H, m), 8.09 - 8.22 (1 H, m), 7.18 - 7.24 (2 H, m), 6.91 - 6.99 (1 H, m), 5.51 (1 H, dt, $J=15.49, 3.70$ Hz), 5.17 - 5.38 (1 H, m), 4.48 - 4.62 (2 H, m), 4.31 - 4.48 (1 H, m), 3.54 - 4.03 (3 H, m), 3.46 - 3.54 (1 H, m), 2.63 - 2.85 (3 H, m), 2.39 - 2.50 (1 H, m), 2.27 - 2.36 (1 H, m), 2.26 (3 H, s), 1.79 - 1.92 (3 H, m), 1.60 - 1.66 (2 H, m). LC/MS ($\text{M}+\text{H}$) = 486.15. HPLC purity >97%.

15 Examples 217 and 218 can be synthesized according to Scheme XXX

Example 217




20

Benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-7,8-dihydro-4H,6H-8,11-methanopyrimido[2,1-d][1,5]oxazocin-11(10H)-yl)carbamate. A mixture of

ethyl 11-(((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-7,8,10,11-tetrahydro-4H,6H-8,11-methanopyrimido[2,1-d][1,5]oxazocine-2-carboxylate (0.070 g, 0.163 mmol), (4-fluorophenyl)methanamine (0.125 g, 1 mmol) and Et₃N (0.418 mL, 3 mmol) in EtOH (5 mL) was heated at reflux for 7 h. The reaction was cooled, 5 concentrated and the residue purified by preparative-HPLC to afford title compound (0.0405 g, 0.080 mmol, 49%) as an off-white solid. ¹H NMR (500 MHz, CDCl₃) δ: 11.99 (0.6 H, brs), 11.92 (0.4 H, brs), 7.82 (0.4 H, t, J = 5.8 Hz), 7.58 (0.6 H, t, J = 5.8 Hz), 7.32 – 7.39 (3 H, m), 7.17 – 7.28 (2 H, m), 7.09 (1 H, dd, J = 8.1, 5.6 Hz), 10 7.01 – 7.06 (1 H, m), 6.91 – 6.99 (2 H, m), 5.60 (0.6 H, s), 5.41 – 5.51 (1 H, m), 5.10 5.02 (0.6 H, d, J = 12.5 Hz), 4.89 (1 H, d, J_{AB} = 12.4 Hz.), 4.79 (0.6 H, d, J = 10.7 Hz), 4.69 – 4.75 (2 H, m), 4.60 (0.4 H, d, J = 10.4 Hz), 4.37 – 4.49 (2.8 H, m), 4.12 (0.6 H, dd, J = 18.8, 6.0 Hz), 3.70 (0.6 H, t, J = 13.7 Hz), 3.63 (0.4 H, t, J = 13.7 Hz), 2.47 – 2.60 (1 H, m), 2.13 – 2.21 (1 H, m.), 1.97 – 2.05 (1 H, m). LCMS (M+H) calcd for C₂₆H₂₆FN₄O₆: 509.18; found: 509.14.

15

Example 218



20 *N'*-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-7,8-dihydro-4H,6H-8,11-methanopyrimido[2,1-d][1,5]oxazocin-11(10H)-yl)-N,N-dimethylethanediamide. A solution of benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-7,8-dihydro-4H,6H-8,11-methanopyrimido[2,1-d][1,5]oxazocin-11(10H)-yl)carbamate (0.039 g, 0.077 mmol) and 1N aq. HCl (0.5 mL, 0.500 mmol) 25 in THF (10 mL) was sparged for 5 min by with N₂. The flask was evacuated and replaced with 1 atm H₂ after which the reaction was stirred for 5 h. The mixture was filtered and concentrated to give amine hydrochloride salt (33 mg) as a white solid which was used in the next step without purification. To a stirred mixture of this


compound in DMF (3 mL) was added, 2-(dimethylamino)-2-oxoacetic acid (0.027 g, 0.230 mmol), diisopropyl-ethylamine (0.134 mL, 0.767 mmol) and 4-(dimethylamino)pyridine (DMAP) (3 mg, 0.025 mmol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.087 g, 0.230 mmol) at room temperature. After 18 h, the reaction mixture was purified by preparative-HPLC to afford the title compound (0.0214 g, 0.045 mmol, 59% yield) as a crystalline off-white solid after crystallization from methanol/H₂O. ¹H NMR (500 MHz, CDCl₃) δ: 12.23 (1 H, s), 9.06 (1 H, t, *J* = 6.0 Hz), 7.43 (1 H, s), 7.33 (2 H, dd, *J* = 8.4, 5.3 Hz), 6.97 - 7.03 (2 H, t, *J* = 8.7 Hz), 5.48 (1 H, td, *J* = 15.6, 3.7 Hz), 4.88 (1 H, d, *J* = 10.7 Hz), 4.72 (1 H, t, *J* = 6.7 Hz), 4.51 (2 H, qd, *J* = 14.7, 6.4 Hz), 4.42 (1 H, d, *J* = 10.7 Hz), 3.69 (1 H, td, *J* = 13.3, 1.2 Hz), 3.17 (3 H, s), 2.89 (3 H, s), 2.59 (1 H, dd, *J* = 11.9, 7.6 Hz), 2.15 - 2.22 (1 H, m), 2.09 (1 H, d, *J* = 12.2 Hz), 1.60 (1 H, td, *J* = 13.9, 3.4 Hz). LCMS (M+H) calcd for C₂₂H₂₅FN₅O₆: 474.18; found: 474.10.

15

Examples 219 through 222 can be synthesized according to Scheme XXXI

Example 219

20




Benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepin-10(4H)-yl)carbamate. A mixture of ethyl 10-(((benzyloxy)carbonyl)amino)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepine-2-carboxylate (150 mg, 0.349 mmol) and 4-fluorobenzylamine (0.200 mL, 1.75 mmol) in EtOH (3 mL) was heated at 90°C for 20 h. The reaction mixture was cooled, concentrated and purified by preparative reversed phase HPLC using a C-18 column (10% to 70% methanol/water) to afford the title compound (110 mg, 62% yield) as a yellow solid. ¹H NMR (500 MHz,

CDCl_3 δ : 7.29-7.38 (m, 4H), 7.11-7.18 (m, 2H), 6.96-7.04 (m, 4H), 5.48 (s, 1H), 4.93-4.98 (m, 1H), 4.81 (d, 1H, J = 11.6 Hz), 4.55-4.65 (m, 2H), 4.45-4.52 (m, 2H), 4.27-4.32 (m, 1H), 3.86 (s, 2H), 2.44-2.48 (m, 1H), 2.28-2.33 (m, 1H), 1.77-1.82 (m, 1H), 1.46-1.52 (m, 1H). LCMS (M+H) = 509.32.

5


Example 220



10 *10-Amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepine-2-carboxamide.* To a mixture of benzyl (2-((4-fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepin-10(4H)-yl)carbamate (110 mg, 0.216 mmol) in methanol (3 mL) was added 10% Pd/C (115 mg, 0.108 mmol) and the mixture
15 stirred under 1 atm H₂ for 18 h. The mixture was then filtered through a pad of CELITE® and the pad washed with ethyl acetate. The filtrate was concentrated to afford the title compound (70 mg, 86% yield) as a light yellow solid. ¹H NMR (500 MHz, CDCl_3) δ : 7.84 (brs, 1H), 7.29 (dd, 2H, J = 8.09 & 5.34 Hz), 7.02 (t, 2H, J = 8.55 Hz), 4.56 (d, 2H, J = 6.1 Hz), 4.43-4.50 (m, 2H), 4.04-4.11 (m, 2H), 3.87 (d, 1H, J = 10.99 Hz), 2.28-2.32 (m, 2H), 1.98-2.09 (m, 3H), 1.55-1.61 (m, 1H). LCMS (M+H) = 375.2.

20

Example 221




5 *10-(Acetylamino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-*
hexahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepine-2-carboxamide. To a
 solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-
 7,10-(epoxymethano)pyrimido[1,2-a]azepine-2-carboxamide (30 mg, 0.080 mmol) in
 CH₂Cl₂ (2 mL) was added Et₃N (0.056 mL, 0.401 mmol) followed by acetyl chloride
10 (0.05 mL, 0.401 mmol) and the resulting mixture stirred at room temperature for 16
 h. The reaction mixture was concentrated and the resulting residue dissolved in
 methanol (2mL) and treated with 2M dimethylamine in methanol (0.5 mL). The
 resulting reaction mixture was stirred at 60 °C for 2 h, cooled and purified by
 preparative HPLC to afford the title compound as a white solid (9 mg, 27 % yield).

15 ¹H NMR (500 MHz, CDCl₃) δ: 11.92 (s, 1H), 7.39 (brs, 1H), 7.29-7.33 (m, 2H),
 7.05-7.09 (m, 2H), 6.13 (s, 1H), 4.59-4.65 (m, 2H), 4.48-4.55 (m, 3H), 4.30 (d, 1H, J
 = 11.9 Hz), 4.10 (d, 1H, J = 16.17 Hz), 2.52-2.58 (m, 1H), 2.30-2.36 (m, 1H), 1.82 (s,
 3H), 1.75-1.82 (m, 1H), 1.50-1.55 (m, 1H). LCMS (M+H)⁺ = 417.25. HPLC purity
 >95%.

20

Example 222




N'-(2-((4-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide. To a stirred solution of 10-amino-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-(epoxymethano)pyrimido[1,2-a]azepine-2-carboxamide (70 mg, 0.187 mmol) and 2-(dimethylamino)-2-oxoacetic acid (32.8 mg, 0.280 mmol) in DMF (3 mL) was added diisopropyl-ethylamine (0.175 mL, 1.002 mmol), 4-(dimethylamino)pyridine (DMAP) (4.57 mg, 0.037 mmol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (107 mg, 0.280 mmol) and the resulting mixture stirred at room temperature for 3 h. After work-up the crude was purified by preparative HPLC to afford the title compound as an off-white solid (35 mg, 40% yield). ^1H NMR (500 MHz, CDCl_3) δ : 12.11 (s, 1H), 8.73 (brs, 1H), 7.47 (s, 1H), 7.34 (dd, 2H, J = 8.39 & 5.34 Hz), 6.98-7.03 (m, 2H), 4.62-4.70 (m, 2H), 4.53 (d, 2H, J = 6.71 Hz), 4.49-4.52 (m, 1H), 4.29 (d, 1H, J = 11.6 Hz), 4.08 (d, 1H, J = 16.79 Hz), 3.21 (s, 3H), 2.89 (s, 3H), 2.46-2.53 (m, 1H), 2.33-2.38 (m, 1H), 1.86-1.92 (m, 1H), 1.53-1.58 (m, 1H). LCMS ($M+\text{H}^+$) = 474.24. HPLC purity >95%.

Examples 223 and 224 can be synthesized according to Scheme XXXII

20

Example 223




25 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[2-(benzoyloxy)ethoxy]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. A mixture of 5-benzoyloxy-1-(2-benzoyloxy-ethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0_{2,7}]trideca-2,4-diene-4-carboxylic acid ethyl ester (33 mg, 0.060 mmol) and (4-fluorophenyl)methanamine (60.4 mg, 0.483 mmol) in DMF (1 mL) was stirred at 100 °C for 2 h under N₂. The mixture was concentrated *in vacuo* and

the residue was dissolved in CH₂Cl₂ (10 mL), washed with 1N-HCl followed by brine, then dried (Na₂SO₄) and concentrated to provide 54 mg of an amber oil which was purified by C-18 reverse phase silica gel, eluting with 34-38-42-50% CH₃CN/H₂O (0.1% trifluoroacetic acid) to obtain the title compound (16 mg, 0.031 mmol, 51 % yield) after crystallization from 95% EtOH as a light-pink crystalline powder: HPLC: retention time = 2.97 min (area percent = 100%); LCMS: m/z 522 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 1.60-1.72 (2H, m), 1.84-1.96 (2H, m), 2.05 (4H, t, J=7.5 Hz), 2.45 (1H, brs), 3.81 (2H, t, J=5.8 Hz), 4.08 (2H, d, J=3.7 Hz), 4.38 (2H, t, J=5.8 Hz), 4.49 (2H, d, J=6.4 Hz), 6.97 (2H, t, J=8.7 Hz), 7.26 (2H, dd, J=8.6, 5.5 Hz), 7.43 (1H, t, J=7.8 Hz), 7.57 (1H, t, J=7.5 Hz), 7.98 (2H, d, J=7.3 Hz), 8.24 (1H, t, J=5.8 Hz), 11.99 (1H, s); ¹³C NMR (126 MHz, CDCl₃) δ ppm 23.3, 27.7, 30.9, 42.5, 53.8, 60.7, 64.0, 77.3, 115.7, 124.9, 128.6, 129.6, 129.7, 130.1, 133.3, 133.5, 146.6, 153.0, 160.0, 162.4, 166.7, 168.6. Anal. calcd for C₂₈N₂₈FN₃O₆: C64.48, H5.41, N8.05; found: C64.45, H5.44, N7.92.

15

Example 224



20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-(2-hydroxyethoxy)-4-oxo-*. To a solution of 7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[2-(benzoyloxy)ethoxy]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo- (8 mg, 0.015 mmol) in a mixture of methanol (0.5 mL) and THF (0.500 mL)


25 was added 0.5 M sodium methoxide (0.092 mL, 0.046 mmol) in methanol at room temp. and the mixture left under N₂ for 15 min. The mixture diluted with ethyl acetate (10 mL) and washed with 1N HCl (2 x 5 mL) followed by brine, then dried (Na₂SO₄), and concentrated to provide a light-pink solid which was triturated with 95% EtOH to yield the title compound (4 mg, 0.01 mmol, Y 63% yield) as a light-

pink crystalline powder: HPLC: retention time = 2.01 min (area percent = 100%); LCMS: m/z 418 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 1.66-1.71 (2H, m), 1.89-1.96 (2H, m), 2.04 (4H, t, J=7.8 Hz), 2.45 (1H, br.s), 2.83 (1H, br), 3.69 (4H, s), 4.09 (2H, d, J=4.0 Hz), 4.50 (2H, d, J=6.1 Hz), 7.00 (2H, t, J=8.7 Hz), 7.33 (2H, dd, J=8.4, 5.3 Hz), 9.35 (H, t, J=5.2 Hz), 12.28 (1H, s); ¹³C NMR (126 MHz, CDCl₃) δ ppm 23.3, 27.7, 30.9, 42.4, 53.8, 61.8, 62.9, 76.5, 115.5, 125.2, 129.9, 134.0, 146.8, 152.1, 160.1, 162.3, 168.8.

Examples 225 through 228 can be synthesized according to Scheme XXXIII

10

Example 225




15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-(2-methoxyethoxy)-4-oxo-*. A solution of 5-benzyloxy-1-(2-methoxy-ethoxy)-6-oxo-3,7-diaza-tricyclo[7.2.2.0_{2,7}]trideca-2,4-diene-4-carboxylic acid methyl ester (8 mg, 0.019 mmol) in trifluoroacetic acid (1 mL, 13 mmol) was kept at room temperature for 3 h.

20 The trifluoroacetic acid was removed *in vacuo* to obtain a light-pink film: HPLC: retention time = 1.35 min (area percent = 100%); LCMS: m/z 339 (M+H). ¹H NMR (500 MHz, CDCl₃) δ ppm 1.74 (2H, br.s), 1.97 (2H, br.s), 2.15 (4H, br.s), 2.50 (1H, br.s), 3.46 (3H, s), 3.69 (2H, br.s), 3.73 (2H, br.s), 4.01 (3H, s), 4.17 (2H, br.s). A mixture of this compound and (4-fluorophenyl)methanamine (11 mg, 0.089 mmol) in

25 DMF (1 mL) was heated in an oil bath at 100 °C under N₂ for 5 h. The mixture was concentrated and the residue purified by C-18 reverse phase column (30% CH₃CN/H₂O (0.1% trifluoroacetic acid)) to obtain the title compound (2.8 mg, 6.4 μmol, 36% yield) as a film: HPLC: retention time = 2.41 min (area percent = 100%); LCMS: m/z 432 (M+H); ¹H NMR (500 MHz, CDCl₃) δ ppm 1.62-1.76 (2H, m),

1.86-1.97 (2H, m), 1.98-2.15 (4H, m), 2.45 (1H, d, J=1.2 Hz), 3.25 (3H, s), 3.52 (2H, t, J=3.5 Hz), 3.75 (2H, d, J=3.5 Hz), 4.10 (2H, d, J=3.4 Hz), 4.57 (2H, d, J=6.4 Hz), 7.01 (2H, t, J=8.4 Hz), 7.28 (2H, dd, J= 7.6, 5.5 Hz), 9.00 (1H, t, J= 5.2 Hz); ^{13}C NMR (126 MHz, CDCl_3) δ ppm 23.3, 27.7, 30.8, 42.4, 53.9, 59.9, 62.6, 72.9, 76.4, 5 115.6, 125.2, 129.0, 133.8, 146.5, 152.3, 160.3, 162.2, 169.1.

Example 226




*7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[2-(dimethylamino)-2-oxoethoxy]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-A solution of 5-benzyloxy-1-dimethylcarbamoylmethoxy-6-oxo-3,7-diazatricyclo[7.2.2.02,7]trideca-2,4-diene-4-carboxylic acid ethyl ester (36 mg, 0.057 mmol) in trifluoroacetic acid (1 mL, 12.98 mmol) was left at room temperature overnight. The mixture was concentrated *in vacuo* to provide an amber film: HPLC: retention time = 1.39 min; LCMS: m/z 380 (M+H); ^1H NMR (400 MHz, CDCl_3) δ ppm 1.44 (3H, t, J=7.2 Hz), 1.68-1.80 (2H, m), 1.99 (2H, m), 2.15-2.33 (4H, m), 2.53 (1H, br.s), 3.00 (3H, s), 3.01 (3H, s), 4.23 (2H, d, J=3.1 Hz), 4.48-4.56 (4H, m). A solution of this film (40 mg, 0.057 mmol) and (4-fluorophenyl) methanamine (93 mg, 0.740 mmol) in DMF (1 mL) was heated at 100 °C (oil bath temp.) for 2 h under N_2 . The mixture was concentrated to remove DMF and the residue purified by C-18 reverse phase column (22% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (0.1% trifluoroacetic acid)) to yield the title compound (11 mg, 0.024 mmol, 42 % yield) after crystallization from 95% EtOH as an off-white crystal: HPLC: retention time = 2.05 min (area percent = 100%); LCMS: m/z 459 (M+H); ^1H NMR (500 MHz, CDCl_3) δ ppm 1.60-1.78 (2H, m), 1.87-1.98 (2H, m), 1.98-2.09 (2H, m), 2.09-2.19 (2H, m), 2.43 (1H, br.s), 2.87 (3H, s), 3.04 (3H, s), 4.09 (2H, d, J=3.7 Hz), 4.28 (2H, s), 4.56 (2H, d, J=6.4 Hz), 6.98 (2H, t, J=6.6 Hz), 7.41 (2H, dd, J=8.7, 5.3 Hz), 9.55 (1H, t, J=5.8 Hz), 12.33 (1H, br);*

¹³C NMR (126 MHz, CDCl₃) δ ppm 23.3, 27.6, 30.7, 35.4, 37.1, 42.2, 53.8, 63.8, 77.5, 115.2, 125.5, 130.1, 134.4, 146.8, 151.7, 160.1, 162.2, 168.8, 169.5. Anal. Calcd for C₂₃H₂₇FN₄O₅: C 60.25, H 5.93, N 12.22; found C 60.08, H 5.80, N 12.13.

5 Examples 227 through 233 can be synthesized according to Scheme XXXIV


Example 227



10

N-(4-Fluoro-2-(4,4,5-trimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)benzyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxamide. A solution of 3-(benzyloxy)-N-(4-fluoro-2-(4,4,5-trimethyl-1,1-dioxido-1,2,5-thiadiazolidin-2-yl)benzyl)-4-oxo-6,7,8,9-tetrahydro-4H-7,10-ethanopyrimido[1,2-a][1,3]diazepine-2-carboxamide (0.054 g, 0.088 mmol) in CH₂Cl₂ (5 mL) and trifluoroacetic acid (0.7 mL, 9.09 mmol) was stirred at room temperature for 18 h. The solution was concentrated to give a yellow oil that was purified by preparative HPLC (Solvent A = 10% methanol/90% H₂O/0.1% trifluoroacetic acid; Solvent B = 90% methanol/10% H₂O/0.1% trifluoroacetic acid; 15 Start % B = 20; Final % B = 100; Gradient time = 20min; Stop time = 25 min; Column = Sunfire 19 x 100 mm, C18, 5μm) to give the title compound (as a trifluoroacetic acid salt) as a pale yellow solid (0.0075 g, 13% yield). LCMS (M+H) calcd for C₂₃H₃₀FN₆O₅S: 521.19; found: 521.25. ¹H NMR (500 MHz, CDCl₃) δ: 8.64 (1H, t, J = 5.6 Hz), 7.60 (1H, dd, J = 8.2, 6.4 Hz), 7.22 (1H, dd, J = 9.2, 2.4 Hz), 7.11 (1H, td, J = 8.1, 2.2 Hz), 4.64 (2H, d, J = 5.8 Hz), 4.00 - 3.94 (2H, m), 3.79 - 3.76 (2H, m), 3.72 (2H, s), 3.58 (2H, s), 3.03 (1H, brs), 2.73 (3H, s), 2.32 - 2.28 (2H, m), 1.91 - 1.87 (2H, m), 1.46 (6H, s).


Example 238



5 *N*-Benzyl-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxamide. Pale yellow solid (3.3 mg, 14% yield). LCMS (M+H) calcd for C₁₈H₂₁N₄O₃: 341.16; found: 341.20. ¹H NMR (500 MHz, CD₃OD) δ: 7.39 - 7.33 (5H, m), 7.29 - 7.27 (1H, m), 4.62 (2H, s), 4.00 - 3.94 (4H, m), 3.88 (2H, s), 3.10 - 3.08 (1H, m), 2.38 - 2.34 (2H, m), 2.02 - 1.97 (2H, m).

10


Example 229



15 *N*-(4-Fluorobenzyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxamide. Pale yellow solid (2.1 mg, 9% yield). LCMS (M+H) calcd for C₁₈H₂₀FN₄O₃: 359.15; found: 359.20. ¹H NMR (500 MHz, CD₃OD) δ: 7.42 - 7.39 (2H, m), 7.09 - 7.04 (3H, m), 4.59 (2H, s), 3.99 - 3.91 (4H, m), 3.86 (2H, s), 3.09 - 3.07 (1H, m), 2.38 - 2.33 (2H, m), 2.02 - 1.97 (2H, m).


20

Example 230



5 *N*-(4-*Fluoro*-2-(3-(2-methoxyethyl)-2-oxo-1-imidazolidinyl)benzyl)-3-
hydroxy-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-
carboxamide. Amber solid (2.1 mg, 6% yield). LCMS (M+H) calcd for
C₂₄H₃₀FN₆O₅: 501.22; found: 501.33. ¹H NMR (500 MHz, CD₃OD) δ: 7.54 - 7.51
(1H, m), 7.39 (1H, s), 4.56 (2H, s), 3.98 - 3.84 (8H, m), 3.69 (2H, t, *J* = 7.8 Hz), 3.58
10 (2H, t, *J* = 5.3 Hz), 3.47 - 3.44 (2H, m), 3.38 (3H, s), 3.06 - 3.04 (1H, m), 2.36 - 2.29
(2H, m), 2.00 - 1.94 (2H, m).


Example 231



15 *N*-(4-*Fluoro*-2-(5-methyl-1*H*-1,2,3-triazol-1-yl)benzyl)-3-hydroxy-4-oxo-
6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxamide.
White solid (1.4 mg, 23% yield). LCMS (M+H) calcd for C₂₁H₂₃FN₇O₃: 440.18;
20 found: 440.23. ¹H NMR (500 MHz, CD₃OD) δ: 9.02 (1H, brs), 7.75 (1H, s), 7.74
(1H, t, *J* = 7.2 Hz), 7.43 (1H, td, *J* = 8.4, 2.6 Hz), 7.35 (1H, dd, *J* = 8.5, 2.9 Hz, 4.33
(2H, s), 4.02 (4H, t, *J* = 7.9 Hz), 3.14 (1H, t, *J* = 4.3 Hz), 2.44 - 2.37 (2H, m), 2.33

(3H, s), 2.07 - 2.02 (2H, m).

Example 232



5


N-(4-Fluoro-2-(3-methyl-2-oxo-1-imidazolidinyl)benzyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxamide.

Pale orange solid (2.0 mg, 40% yield). LCMS (M+H) calcd for C₂₂H₂₆FN₆O₄:

10 457.20; found: 457.24. ¹H NMR (500 MHz, CD₃OD) δ: 9.30 (1H, brs), 7.52 (1H, dd, *J* = 8.5, 6.4 Hz), 7.17 (1H, dd, *J* = 9.8, 2.7 Hz), 7.09 (1H, td, *J* = 8.3, 2.5 Hz), 4.58 (2H, s), 4.02 - 3.98 (4H, m), 3.92 (2H, s), 3.87 (2H, dd, *J* = 8.8, 7.0 Hz), 3.61 (2H, t, *J* = 7.9 Hz), 3.10 (1H, t, *J* = 4.0 Hz), 2.87 (3H, s), 2.38 - 2.33 (2H, m), 2.04 - 1.98 (2H, m).

15

Example 233




20 *N*-(4-Fluoro-2-(5-methyl-1*H*-1,2,4-triazol-1-yl)benzyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-4*H*-7,10-ethanopyrimido[1,2-*a*][1,3]diazepine-2-carboxamide.

Pale orange solid (5.2 mg, 13% yield) LCMS (M+H) calcd for C₂₁H₂₃FN₇O₃: 440.18; found: 440.10. ¹H NMR (500 MHz, CD₃OD) δ: 8.14 (1H, s), 7.70 (1H, dd, *J*

= 8.5, 5.8 Hz), 7.42 - 7.36 (2H, m), 4.41 (2H, s), 4.02 (4H, t, J = 7.8 Hz), 3.93 (2H, m), 3.15 (1H, t, J = 4.3 Hz), 2.44 (3H, s), 2.42 - 2.37 (2H, m), 2.08 - 2.03 (2H, m).

Example 234


5



Ethanediamide, N,N-dimethyl-N'-(6,7,8,9-tetrahydro-3-hydroxy-2-[(1H-indol-3-ylmethyl)amino]carbonyl]-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl). ^1H NMR (500 MHz, DMSO- d_6) δ ppm 11.97 (1 H, s), 9.58 (1 H, t, J =6.10 Hz), 8.75 (1 H, s), 6.18 (1 H, d, J =3.05 Hz), 5.99 (1 H, d, J =2.14 Hz), 4.43 (2 H, d, J =6.10 Hz), 4.04 (2 H, d, J =3.66 Hz), 3.26 - 3.47 (3 H, m), 2.94 (3 H, s), 2.88 (3 H, s), 2.42 (1 H, brs), 2.25 - 2.34 (2 H, m), 2.04 (2 H, ddd, J =14.34, 9.16, 5.80 Hz), 1.73 - 1.86 (2 H, m), 1.55 - 1.66 (2 H, m). LCMS (M+H) $+=$ 458.2.

15

Example 235




Ethanediamide, N,N-dimethyl-N'-(6,7,8,9-tetrahydro-3-hydroxy-2-[(1H-indol-3-ylmethyl)amino]carbonyl]-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl). ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.32 (1 H, s), 10.96 (1 H, brs), 9.61 (1 H, t, J =6.26 Hz), 8.70 (1 H, s), 7.70 (1 H, d, J =7.93 Hz), 7.30 - 7.39 (2 H, m), 7.07 (1

H, t, $J=7.17$ Hz), 6.98 (1 H, t, $J=7.17$ Hz), 4.65 (2 H, d, $J=6.10$ Hz), 3.99 - 4.07 (2 H, m), 2.88 - 2.94 (6 H, m), 2.39 (1 H, brs), 2.20 - 2.30 (2 H, m), 2.00 (2 H, ddd, $J=14.27, 9.08, 5.65$ Hz), 1.71 - 1.81 (2 H, m), 1.55 - 1.65 (2 H, m). LCMS (M+H) $+=$ 493.3.

5


Example 236



10 *Ethanediamide, N'-(2-[[[1-(1,1-dimethylethyl)-5-methyl-1H-pyrazol-3-yl]methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-*. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.15 (1 H, brs), 9.51 (1 H, t, $J=6.41$ Hz), 8.74 (1 H, s), 5.98 (1 H, s), 4.36 (2 H, d, $J=6.41$ Hz), 4.04 (2 H, d, $J=3.97$ Hz), 2.94 (3 H, s), 2.87 (3 H, s), 15 2.42 (1 H, brs), 2.34 - 2.39 (3 H, m), 2.23 - 2.31 (2 H, m), 2.07 (2 H, ddd, $J=14.19, 9.00, 5.49$ Hz), 1.75 - 1.84 (2 H, m), 1.59 - 1.65 (2 H, m), 1.51 - 1.55 (9 H, m). LCMS (M+H) $+=$ 514.3.

Example 237


20



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[5-methyl-2-furanyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.84 (1 H, brs), 8.86 (1 H, t, J=6.26 Hz), 8.37 (1 H, s), 7.24 - 7.30 (2 H, m), 7.11 - 7.19 (2 H, m), 6.93 (1 H, d, J=3.36 Hz), 6.20 (1 H, d, J=2.44 Hz), 4.44 (2 H, d, J=6.41 Hz), 4.04 (2 H, d, J=3.66 Hz), 2.52 - 2.60 (2 H, m), 2.43 (1 H, brs), 2.17 (3 H, s), 2.03 - 2.11 (2 H, m), 1.79 - 1.89 (2 H, m), 1.61 - 1.71 (2 H, m). LCMS (M+H)⁺= 481.36.


Example 238

10



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[tetrahydro-3-furanyl]carbonyl]amino]-. . ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.98 (1 H, brs), 8.92 (1 H, t, J=6.41 Hz), 8.04 (1 H, s), 7.33 - 7.43 (2 H, m), 7.15 - 7.21 (2 H, m), 4.44 - 4.60 (2 H, m), 4.01 (2 H, d, J=3.66 Hz), 3.68 - 3.75 (1 H, m), 3.60 - 3.67 (2 H, m), 3.52 - 3.59 (1 H, m), 2.98 - 3.10 (1 H, m), 2.39 (1 H, brs), 2.30 - 2.38 (2 H, m), 1.94 - 2.06 (2 H, m), 1.83 - 1.92 (2 H, m), 1.72 - 1.79 (2 H, m), 1.54 - 1.65 (2 H, m). LCMS (M+H)⁺= 471.2.


Example 239



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[imidazo[1,2-a]pyridin-6-ylcarbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.82 (1 H, brs), 9.41 (1 H, s), 8.79 (1 H, s), 8.66 (1 H, t, J=6.26 Hz), 8.26 (1 H, d, J=1.83 Hz), 8.20 (1 H, s), 8.09 (1 H, d, J=9.46 Hz), 7.84 (1 H, d, J=9.46 Hz), 7.05 (2 H, dd, J=8.70, 5.65 Hz), 6.86 - 6.91 (2 H, m), 4.36 (2 H, d, J=6.41 Hz), 4.08 (2 H, d, J=3.66 Hz), 2.45 - 2.48 (1 H, m), 2.32 - 2.39 (2 H, m), 2.24 - 2.31 (2 H, m), 1.83 - 1.91 (2 H, m), 1.65 - 1.73 (2 H, m). LCMS (M+H)+= 517.37.


10

Example 240




7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[1-(1,1-dimethylethyl)-5-methyl-1H-pyrazol-3-yl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.82 (1 H, brs), 8.78 (1 H, t, J=6.41 Hz), 8.56 (1 H, s), 7.27 (2 H, dd, J=8.70, 5.65 Hz), 7.10 - 7.17 (2 H, m), 6.41 (1 H, s), 4.44 (2 H, d, J=6.41 Hz), 4.04 (2 H, d, J=3.66 Hz), 2.75 (2 H, ddd, J=14.04, 9.00, 5.65 Hz), 2.42 - 2.45 (1 H, m), 2.41 (3 H, s), 1.92 - 2.02 (2 H, m), 1.81 - 1.91 (2 H, m), 1.62 - 20 1.71 (2 H, m), 1.48 - 1.53 (9 H, m). LCMS (M+H)+= 537.3.

Example 241




5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1,3-dimethyl-1H-pyrazol-5-yl)carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 8.64 (1 H, t, *J*=6.41 Hz), 8.32 (1 H, s), 7.16 - 7.21 (2 H, m), 7.09 - 7.15 (2 H, m), 6.57 (1 H, s), 4.40 (2 H, d, *J*=6.41 Hz), 4.05 (2 H, d, *J*=3.66 Hz), 3.83 (3 H, s), 2.44 (1 H, brs), 2.36 (2 H, ddd, *J*=14.19, 10.00, 5.80 Hz), 2.15 - 2.23 (2 H, m), 2.07 (3 H, s), 1.78 - 1.88 (2 H, m), 1.60 - 1.69 (2 H, m). LCMS (M+H)⁺= 495.3.

Example 242




15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1,5-dimethyl-1H-imidazol-4-yl)carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.75 (1 H, brs), 9.23 (1 H, brs), 8.80 (1 H, t, *J*=6.10 Hz), 7.79 (1 H, brs), 7.37 (2 H, dd, *J*=8.70, 5.65 Hz), 7.19 (2 H, t, *J*=8.85 Hz), 4.55 (2 H, d, *J*=6.41 Hz), 4.04 (2 H, d, *J*=3.66 Hz), 3.55 (3 H, s), 2.69 - 2.80 (2 H, m), 2.43 (1 H, brs), 2.38 (3 H, s), 1.81 - 1.96 (4 H, m), 1.64 - 1.74 (2 H, m). LCMS (M+H)⁺= 495.3.

Example 243



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1,4-dimethyl-1H-imidazol-5-yl)carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-.* ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 8.64 (1 H, brs), 8.31 (1 H, brs), 7.99 (1 H, brs), 7.23 - 7.31 (2 H, m), 6.98 (2 H, t, *J*=8.24 Hz), 4.50 (2 H, d, *J*=6.10 Hz), 4.18 (2 H, d, *J*=3.36 Hz), 3.75 (3 H, s), 2.77 - 2.88 (2 H, m), 2.53
10 (1 H, brs), 2.41 (3 H, s), 1.97 - 2.09 (4 H, m), 1.69 - 1.79 (2 H, m). LCMS (M+H) $+=$ 495.3.


Example 244



15 *Ethanediamide, N,N-dimethyl-N'-[6,7,8,9-tetrahydro-3-hydroxy-2-[[imidazo[1,2-a]pyridin-6-ylmethyl]amino]carbonyl]-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl].* ^1H NMR (500 MHz, DMSO-*d*₆) δ ppm 11.70 (1 H, brs), 9.85 (1 H, t, *J*=6.26 Hz), 8.89 (1 H, s), 8.75 (1 H, s), 8.37 (1 H, d, *J*=1.53 Hz), 8.18 (1 H, d, *J*=1.83 Hz), 7.96 - 8.00 (1 H, m), 7.91 - 7.94 (1 H, m), 4.65 (2 H, d, *J*=6.41 Hz), 4.05 (3 H, d, *J*=3.97 Hz), 2.95 (3 H, s), 2.82 (3 H, s), 2.43 (1 H, brs), 2.25 - 2.34 (2 H, m), 2.00 - 2.09 (2 H, m), 1.75 - 1.85 (2 H, m), 1.57 - 1.67 (2 H, m). LCMS (M+H) $+=$ 494.3.

25


Example 245



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[1-methyl-1H-pyrazol-3-yl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.77 (1 H, brs), 8.69 (1 H, s), 8.47 (1 H, t, *J*=6.56 Hz), 7.73 (1 H, d, *J*=2.14 Hz), 7.28 (2 H, dd, *J*=8.55, 5.49 Hz), 7.10 - 7.21 (2 H, m), 6.56 (1 H, d, *J*=2.14 Hz), 4.53 (2 H, d, *J*=6.71 Hz), 4.06 (2 H, d, *J*=3.66 Hz), 3.67 (3 H, s), 2.66 (2 H, ddd, *J*=14.19, 9.16, 5.65 Hz), 2.41 - 2.47 (1 H, m), 1.98 - 2.08 (2 H, m), 1.79 - 1.92 (2 H, m), 1.62 - 1.74 (2 H, m). LCMS (M+H) $^{+}$ = 481.28.


Example 246

15



15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[1-methyl-1H-pyrazol-4-yl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.62 (1 H, brs), 8.56 (1 H, t, *J*=6.26 Hz), 8.08 (1 H, s), 7.96 (1 H, s), 7.80 (1 H, s), 7.09 - 7.19 (4 H, m), 4.36 (2 H, d, *J*=6.41 Hz), 4.04 (2 H, d, *J*=3.66 Hz), 3.79 (3 H, s), 2.43 (1 H, brs), 2.33 (2 H, ddd, *J*=14.34, 9.00, 5.95 Hz), 2.20 (2 H, ddd, *J*=13.96, 7.32, 7.10 Hz), 1.77 - 1.86 (2 H, m), 1.59 - 1.68 (2 H, m). LCMS (M+H) $^{+}$ = 481.28.


Example 247



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[1-methyl-1H-pyrazol-5-yl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.72 (1 H, brs), 8.66 (1 H, t, *J*=6.26 Hz), 8.41 (1 H, s), 7.39 - 7.43 (1 H, m), 7.17 - 7.23 (2 H, m), 7.10 - 7.17 (2 H, m), 6.76 (1 H, d, *J*=2.14 Hz), 4.38 (2 H, d, *J*=6.41 Hz), 4.05 (2 H, d, *J*=3.66 Hz), 3.90 (3 H, s), 2.36 - 2.48 (3 H, m), 2.14 - 2.22 (2 H, m), 1.84 (2 H, td, *J*=14.19, 5.80 Hz), 1.59 - 1.71 (2 H, m). LCMS (M+H) $^{+}$ = 481.28.

10


Example 248



15


15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[1-methyl-1H-1,2,3-triazol-4-yl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.65 (1 H, brs), 9.20 (1 H, s), 8.71 (1 H, t, *J*=6.56 Hz), 8.51 (1 H, s), 7.34 - 7.42 (2 H, m), 7.12 - 7.19 (2 H, m), 4.48 (2 H, d, *J*=6.41 Hz), 4.10 (3 H, s), 4.05 (2 H, d, *J*=3.66 Hz), 2.60 - 2.68 (2 H, m), 2.42 - 2.47 (1 H, m), 2.03 - 2.12 (2 H, m), 1.82 - 1.93 (2 H, m), 1.69 (2 H, dd, *J*=7.32, 6.10 Hz). LCMS (M+H) $^{+}$ = 482.27.

Example 249




5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[1-methyl-1H-tetrazol-5-yl)carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 12.12 (1 H, brs), 10.12 (1 H, s), 8.26 - 8.33 (1 H, m), 7.50 - 7.55 (2 H, m), 7.00 - 7.06 (2 H, m), 4.68 (2 H, d, *J*=6.41 Hz), 4.38 (3 H, s), 4.18 (2 H, d, *J*=3.97 Hz), 2.95 - 3.04 (2 H, m), 2.50 - 2.56 (1 H, m), 2.03 - 2.12 (2 H, m), 1.87 - 1.97 (2 H, m), 1.74 - 1.84 (2 H, m). LCMS (M+H) $^{+}$ = 483.3.

Example 250




15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[2-methyl-2H-tetrazol-5-yl)carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, DMSO-*d*₆) δ ppm 11.83 (1 H, s), 9.26 (1 H, s), 8.87 (1 H, t, *J*=6.26 Hz), 7.32 - 7.38 (2 H, m), 7.13 - 7.20 (2 H, m), 4.47 (2 H, d, *J*=6.41 Hz), 4.38 (3 H, s), 4.06 (2 H, d, *J*=3.66 Hz), 2.59 (2 H, td, *J*=9.38, 4.73 Hz), 2.54 - 2.55 (1 H, m), 2.08 - 2.17 (2 H, m), 1.83 - 1.92 (2 H, m), 1.65 - 1.74 (2 H, m). LCMS (M+H) $^{+}$ = 483.3.

Example 251




5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-10-[(3-isoxazolylcarbonyl)amino]-3-methoxy-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.75 (1 H, s), 9.08 (1 H, d, *J*=1.53 Hz), 8.96 (1 H, s), 8.80 (1 H, t, *J*=6.41 Hz), 7.29 - 7.34 (2 H, m), 7.13 - 7.19 (2 H, m), 6.83 (1 H, d, *J*=1.83 Hz), 4.45 (2 H, d, *J*=6.71 Hz), 4.05 (2 H, d, *J*=3.66 Hz), 2.40 - 2.48 (3 H, m), 2.14 - 2.22 (2 H, m), 1.84 (2 H, td, *J*=14.11, 5.34 Hz), 1.63 - 1.71 (2 H, m). LCMS (M+H)⁺= 486.29.

Example 252



15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(2R)-2-pyrrolidinylcarbonyl]amino-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.95 (1 H, s), 9.16 (1 H, d, *J*=5.49 Hz), 9.02 (1 H, t, *J*=6.26 Hz), 8.53 (1 H, s), 7.36 - 7.43 (2 H, m), 7.15 - 7.22 (2 H, m), 4.63 (1 H, dd, *J*=14.80, 6.87 Hz), 4.43 (1 H, dd, *J*=14.95, 5.49 Hz), 4.26 - 4.34 (1 H, m), 3.98 - 4.10 (2 H, m), 3.14 - 3.23 (2 H, m), 2.41 - 2.45 (1 H, m), 2.29 - 2.40 (2 H, m), 2.07 - 2.16 (2 H, m), 1.99 - 2.07 (1 H, m), 1.72 - 1.86 (5 H, m), 1.58 - 1.69 (2 H, m). LCMS (M+H)⁺= 470.3.


Example 253



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2R)-1-acetyl-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo.* ¹H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 10.10 (1 H, brs), 9.24 (1 H, s), 7.33 - 7.39 (2 H, m), 6.97 - 7.04 (2 H, m), 4.65 - 4.72 (2 H, m), 4.55 (1 H, dd, *J*=14.65, 6.10 Hz), 4.23 (1 H, dd, *J*=15.26, 3.05 Hz), 4.08 (1 H, dd, *J*=15.26, 1.83 Hz), 3.48 - 3.54 (1 H, m), 3.39 (1 H, td, *J*=10.07, 7.02 Hz), 2.92 - 3.01 (2 H, m), 2.53 (1 H, dd, *J*=12.51, 6.41 Hz), 2.44 (1 H, brs), 2.07 - 2.19 (1 H, m), 1.91 - 2.06 (3 H, m), 1.88 (3 H, s), 1.66 - 1.80 (3 H, m), 1.56 - 1.66 (2 H, m). LCMS (M+H)⁺= 512.55.

15


Example 254



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2R)-1-(methylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-4-oxo-. ^1H NMR (500 MHz, CHLOROFORM-d) δ ppm 8.65 (1 H, s), 8.37 (1 H, t, $J=6.41$ Hz), 7.37 (2 H, dd, $J=8.55$, 5.19 Hz), 6.98 - 7.03 (2 H, m), 4.61 (2 H, d, $J=6.41$ Hz), 4.25 (1 H, dd, $J=15.26$, 3.05 Hz), 4.13 (1 H, dd, $J=8.70$, 3.20 Hz), 4.08 (1 H, dd, $J=15.41$, 1.98 Hz), 3.38 (1 H, ddd, $J=10.61$, 6.94, 4.12 Hz), 3.22 (1 H, td, $J=9.46$, 7.02 Hz), 2.96 (1 H, ddd, $J=14.04$, 9.61, 6.87 Hz), 2.80 - 2.87 (1 H, m), 2.79 (3 H, s), 2.47 (1 H, brs), 2.35 - 2.41 (1 H, m), 1.75 - 2.10 (7 H, m), 1.68 - 1.75 (1 H, m), 1.58 - 1.66 (1 H, m).

10 LCMS (M+H) $+=$ 548.60.

Example 255




15

7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-1-acetyl-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ^1H NMR (500 MHz, CHLOROFORM-d) δ ppm 10.10 (1 H, brs), 9.24 (1 H, s), 7.33 - 7.39 (2 H, m), 6.97 - 7.04 (2 H, m), 4.65 - 4.72 (2 H, m), 4.55 (1 H, dd, $J=14.65$, 6.10 Hz), 4.23 (1 H, dd, $J=15.26$, 3.05 Hz), 4.08 (1 H, dd, $J=15.26$, 1.83 Hz), 3.48 - 3.54 (1 H, m), 3.39 (1 H, td, $J=10.07$, 7.02 Hz), 2.92 - 3.01 (2 H, m), 2.53 (1 H, dd, $J=12.51$, 6.41 Hz), 2.44 (1 H, brs), 2.07 - 2.19 (1 H, m), 1.91 - 2.06 (3 H, m), 1.88 (3 H, s), 1.66 - 1.80 (3 H, m), 1.56 - 1.66 (2 H, m). LCMS (M+H) $+=$ 512.55.

25


Example 256



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S)-1-(methylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 8.65 (1 H, s), 8.37 (1 H, t, *J*=6.41 Hz), 7.37 (2 H, dd, *J*=8.55, 5.19 Hz), 6.98 - 7.03 (2 H, m), 4.61 (2 H, d, *J*=6.41 Hz), 4.25 (1 H, dd, *J*=15.26, 3.05 Hz), 4.13 (1 H, dd, *J*=8.70, 3.20 Hz), 4.08 (1 H, dd, *J*=15.41, 1.98 Hz), 3.38 (1 H, ddd, *J*=10.61, 6.94, 4.12 Hz), 3.22 (1 H, td, *J*=9.46, 7.02 Hz), 2.96 (1 H, ddd, *J*=14.04, 9.61, 6.87 Hz), 2.80 - 2.87 (1 H, m), 2.79 (3 H, s), 2.47 (1 H, brs), 2.35 - 2.41 (1 H, m), 1.75 - 2.10 (7 H, m), 1.68 - 1.75 (1 H, m), 1.58 - 1.66 (1 H, m). LCMS (M+H) $^+$ = 548.60.

15

Example 257




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2R)-1-methyl-2-pyrrolidinyl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, DMSO-*d*₆) δ ppm 11.86 (1 H, brs), 9.49 (1 H, brs), 8.92 (1 H, t, *J*=6.26 Hz), 8.76 (1 H, s), 7.38 (2 H, dd, *J*=8.55, 5.49 Hz), 7.19 (2 H, t, *J*=8.85 Hz), 4.53 - 4.60 (1 H, m), 4.45 (1 H, dd, *J*=14.80, 5.95 Hz), 4.13 (1 H, q, *J*=8.14 Hz), 3.98 - 4.09 (2 H, m), 3.50 (1 H, d,

J=3.36 Hz), 3.05 - 3.13 (1 H, m), 2.67 (3 H, s), 2.43 (1 H, brs), 2.28 - 2.38 (1 H, m), 2.10 - 2.26 (4 H, m), 1.89 - 1.99 (1 H, m), 1.70 - 1.85 (4 H, m), 1.58 - 1.67 (2 H, m). LCMS (M+H)⁺ = 484.32.

5


Example 258



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S)-1-methyl-2-pyrrolidinyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.86 (1 H, brs), 9.49 (1 H, brs), 8.92 (1 H, t, *J*=6.26 Hz), 8.76 (1 H, s), 7.38 (2 H, dd, *J*=8.55, 5.49 Hz), 7.19 (2 H, t, *J*=8.85 Hz), 4.53 - 4.60 (1 H, m), 4.45 (1 H, dd, *J*=14.80, 5.95 Hz), 4.13 (1 H, q, *J*=8.14 Hz), 3.98 - 4.09 (2 H, m), 3.50 (1 H, d, *J*=3.36 Hz), 3.05 - 3.13 (1 H, m), 2.67 (3 H, s), 2.43 (1 H, brs), 2.28 - 2.38 (1 H, m), 2.10 - 2.26 (4 H, m), 1.89 - 1.99 (1 H, m), 1.70 - 1.85 (4 H, m), 1.58 - 1.67 (2 H, m). LCMS (M+H)⁺ = 484.32.

20

Example 259




Carbamic acid, [2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-, 2-methoxyethyl ester. ¹H NMR (500 MHz, CHLOROFORM-d) δ ppm 7.76 (1 H, brs),

7.31 (2 H, d, $J=17.70$ Hz), 7.04 (2 H, t, $J=8.55$ Hz), 4.58 (2 H, d, $J=5.80$ Hz), 4.15 (2 H, d, $J=3.66$ Hz), 3.98 - 4.02 (2 H, m), 3.45 (2 H, brs), 3.34 (3 H, brs), 2.48 (1 H, brs), 2.01 - 2.09 (2 H, m), 1.86 - 1.97 (4 H, m), 1.64 - 1.72 (2 H, m). LCMS (M+H) $= 475.33$.


5

Example 260



10 *Carbamic acid, [2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-, 2-hydroxyethyl ester.* ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 7.72 (1 H, brs), 7.29 - 7.35 (2 H, m), 7.05 (2 H, t, $J=8.55$ Hz), 4.58 (2 H, d, $J=5.19$ Hz), 4.16 (2 H, d, $J=3.66$ Hz), 3.98 - 4.02 (2 H, m), 3.67 (2 H, brs), 2.49 (1 H, brs), 2.03 - 2.09 (4 H, m), 1.88 - 1.97 (2 H, m), 1.64 - 1.73 (2 H, m). LCMS (M+H) $= 461.29$.

Example 261




20

20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[(2-hydroxy-2-methyl-1-oxopropyl)amino]-4-oxo-*. ^1H NMR (500 MHz, DMSO-*d*₆) δ ppm 11.80 (1 H, brs), 8.73 (1 H, s), 8.61 (1 H, t, $J=6.41$ Hz), 7.39 (2 H, dd, $J=8.55, 5.49$ Hz), 7.14 - 7.21 (2 H, m), 5.70 (1 H, brs), 4.51 (2 H, d, $J=6.41$ Hz), 4.02 (2 H, d, $J=3.66$ Hz), 2.56 - 2.65

(2 H, m), 2.41 (1 H, brs), 1.77 - 1.91 (4 H, m), 1.61 - 1.71 (2 H, m), 1.21 - 1.26 (6 H, m). LCMS (M+H)⁺ = 459.5.

Example 262


5



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-1-acetyl-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.50 (1 H, brs), 9.89 (1 H, t, J=6.41 Hz), 8.68 (1 H, s), 7.25 (1 H, d, J=7.32 Hz), 7.17 - 7.22 (1 H, m), 7.07 - 7.12 (1 H, m), 4.49 - 4.55 (3 H, m), 3.96 - 4.07 (2 H, m), 3.48 - 3.54 (1 H, m), 3.41 - 3.47 (1 H, m), 2.40 - 2.48 (2 H, m), 2.38 (1 H, brs), 2.21 (3 H, s), 2.07 - 2.13 (1 H, m), 1.92 (3 H, s), 1.85 - 1.90 (3 H, m), 1.73 - 1.82 (3 H, m), 1.55 - 1.64 (2 H, m). LCMS (M+H)⁺ = 526.2.

Example 263

20




7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S)-1-(methylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.34 (1 H, brs), 8.62 (1 H, t, J=6.41 Hz), 8.43 (1 H, s), 7.25 (1 H, d, J=7.63 Hz), 7.16 - 7.21 (1 H, m), 7.09 (1 H, t, J=9.00 Hz), 4.49 - 4.55 (1 H, m),

4.43 - 4.48 (1 H, m), 4.29 (1 H, t, $J=5.95$ Hz), 4.09 (1 H, dd, $J=15.26, 3.66$ Hz), 3.95 (1 H, dd, $J=15.11, 2.59$ Hz), 3.32 - 3.42 (1 H, m), 3.26 - 3.32 (1 H, m), 3.02 (3 H, s), 2.57 - 2.65 (1 H, m), 2.36 - 2.42 (1 H, m), 2.21 (3 H, s), 2.00 - 2.06 (2 H, m), 1.88 - 1.97 (2 H, m), 1.70 - 1.86 (5 H, m), 1.56 - 1.68 (2 H, m). LCMS (M+H) $+=$ 562.2.

5


Example 264



10 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[[(2S)-4-oxo-2-azetidinyl]carbonyl]amino]-.* ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.04 (1 H, brs), 8.96 - 9.05 (1 H, m), 8.28 (1 H, s), 8.19 (1 H, s), 7.37 - 7.43 (2 H, m), 7.14 - 7.21 (2 H, m), 4.51 - 4.55 (2 H, m), 4.10 (1 H, dd, $J=5.49, 2.44$ Hz), 4.00 - 4.04 (2 H, m), 3.02 - 3.08 (1 H, m), 2.83 (1 H, dd, $J=17.55, 6.26$ Hz), 2.63 - 2.72 (1 H, m), 2.35 - 2.43 (2 H, m), 1.97 (2 H, dd, $J=13.58, 5.34$ Hz), 1.75 - 1.84 (2 H, m), 1.58 - 1.68 (2 H, m). LCMS (M+H) $+=$ 470.1.

20

Example 265




25 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-1-acetyl-2-piperidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-.* ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.44 (1 H, brs), 9.47 (1 H, t,

J=6.41 Hz), 8.27 (1 H, s), 7.37 - 7.43 (2 H, m), 7.14 - 7.20 (2 H, m), 5.09 (1 H, d, *J*=4.58 Hz), 4.61 - 4.67 (1 H, m), 4.52 - 4.59 (1 H, m), 4.05 - 4.11 (1 H, m), 3.94 - 3.99 (1 H, m), 3.75-3.72 (1 h, m), 3.05 (1 H, td, *J*=12.97, 2.75 Hz), 2.57 - 2.65 (1 H, m), 2.45 - 2.48 (1 H, m), 2.36 - 2.41 (1 H, m), 1.97 - 2.02 (1 H, m), 1.96 (3 H, s),
5 1.88 - 1.94 (1 H, m), 1.73 - 1.86 (3 H, m), 1.58 - 1.66 (3 H, m), 1.53 - 1.58 (2 H, m), 1.38 - 1.47 (2 H, m). LCMS (M+H)⁺= 526.4.

Example 266


10



*7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S)-1-(methylsulfonyl)-2-piperidinyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.24 (1 H, brs), 8.79 (1 H, t, *J*=6.41 Hz), 8.30 (1 H, s), 7.36 - 7.41 (2 H, m), 7.15 - 7.20 (2 H, m), 4.68 (1 H, dd, *J*=14.80, 7.48 Hz), 4.62 (1 H, d, *J*=4.58 Hz), 4.40 (1 H, dd, *J*=14.80, 5.34 Hz), 4.13 (1 H, dd, *J*=15.11, 3.81 Hz), 3.93 (1 H, dd, *J*=15.11, 2.59 Hz), 3.54 - 3.59 (1 H, m), 3.22 (1 H, td, *J*=12.82, 2.75 Hz), 2.95 (3 H, s), 2.54 - 2.57 (1 H, m), 2.38 - 2.48 (2 H, m), 1.99 - 2.14 (2 H, m), 1.74 - 1.83 (3 H, m), 1.53 - 1.69 (5 H, m), 1.37 - 1.47 (1 H, m), 1.23 - 1.33 (1 H, m). LCMS (M+H)⁺= 562.5.*

Example 267


25



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-1-acetyl-2-azetidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 12.66 (1 H, brs), 10.04 (1 H, t, *J*=6.26 Hz), 8.99 (1 H, s), 7.38 (2 H, dd, *J*=8.70, 5.65 Hz), 7.14 - 7.21 (2 H, m), 5 4.80 (1 H, dd, *J*=9.31, 6.26 Hz), 4.56 - 4.63 (1 H, m), 4.50 - 4.56 (1 H, m), 4.03 - 4.10 (2 H, m), 3.96 - 4.02 (2 H, m), 2.72 - 2.80 (1 H, m), 2.66 - 2.72 (1 H, m), 2.36 - 2.43 (2 H, m), 2.24 - 2.33 (1 H, m), 1.75 - 1.85 (4 H, m), 1.74 (3 H, s), 1.58 - 1.67 (2 H, m). LCMS (M+H)⁺= 498.4.


10

Example 268



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S)-1-(methylsulfonyl)-2-azetidinyl]carbonyl]amino]-4-oxo-. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 12.35 (1 H, brs), 8.78 (1 H, t, *J*=6.41 Hz), 8.51 (1 H, s), 7.38 (2 H, dd, *J*=8.55, 5.80 Hz), 7.16 (2 H, t, *J*=8.85 Hz), 4.81 (1 H, dd, *J*=9.77, 7.63 Hz), 4.57 (1 H, dd, *J*=14.65, 7.02 Hz), 4.44 (1 H, dd, *J*=14.80, 5.95 Hz), 4.12 (1 H, dd, *J*=14.95, 20 3.66 Hz), 3.97 - 4.02 (1 H, m), 3.91 - 3.97 (1 H, m), 3.67 (1 H, td, *J*=8.62, 4.73 Hz), 3.21 (3 H, s), 2.75 (1 H, ddd, *J*=13.20, 9.23, 6.87 Hz), 2.63 (1 H, ddd, *J*=13.73, 9.31, 4.12 Hz), 2.38 - 2.48 (2 H, m), 2.19 - 2.28 (1 H, m), 1.93 - 2.02 (1 H, m), 1.76 - 1.85 (2 H, m), 1.65 - 1.75 (2 H, m), 1.57 - 1.64 (1 H, m). LCMS (M+H)⁺= 534.5.


Example 269



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-1-[2-(dimethylamino)-1,2-dioxoethyl]-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 12.29 (1 H, brs), 9.31 (1 H, t, *J*=6.41 Hz), 8.44 (1 H, s), 7.33 (2 H, dd, *J*=8.55, 5.49 Hz), 7.15 (2 H, t, *J*=8.85 Hz), 4.54 - 4.63 (2 H, m), 4.37 (1 H, dd, *J*=14.65, 6.10 Hz), 4.17 (1 H, dd, *J*=15.11, 4.12 Hz), 3.86 - 3.92 (1 H, m), 3.33 - 3.42 (2 H, m), 2.84 (3 H, s), 2.79 - 2.82 (3 H, m), 2.36 - 2.41 (1 H, m), 2.13 - 2.33 (3 H, m), 1.99 - 2.07 (1 H, m), 1.92 - 1.99 (1 H, m), 1.82 - 1.93 (3 H, m), 1.70 - 1.82 (2 H, m), 1.53 - 1.66 (2 H, m). LCMS (M+H) $^{+}$ = 569.2.

10

15 Example 270




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1R,3S,5R)-2-acetyl-2-azabicyclo[3.1.0]hex-3-yl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 9.86 (1 H, brs), 8.60 (1 H, brs), 7.38 (2 H, dd, *J*=8.55, 5.49 Hz), 7.17 (2 H, t, *J*=8.85 Hz), 4.56 (2 H, d, *J*=6.41 Hz), 4.46 - 4.49 (1 H, m), 4.02 (2 H, t, *J*=3.51 Hz), 3.37 -

3.41 (1 H, m), 3.17 (1 H, d, $J=5.19$ Hz), 2.47 (2 H, brs), 2.37 - 2.40 (1 H, m), 2.05 (3 H, s), 1.83 - 1.93 (3 H, m), 1.71 - 1.83 (3 H, m), 1.56 - 1.65 (2 H, m), 0.92 - 0.98 (1 H, m), 0.54 (1 H, td, $J=4.81$, 2.29 Hz). LCMS (M+H) $^+=$ 524.2.

5


Example 271



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S,4S)-1-acetyl-4-methoxy-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. 1 H NMR (500 MHz, DMSO- d_6) δ ppm 12.40 (1 H, brs), 9.71 (1 H, t, $J=6.41$ Hz), 8.26 (1 H, s), 7.39 (2 H, dd, $J=8.55$, 5.49 Hz), 7.14 - 7.21 (2 H, m), 4.63 (1 H, dd, $J=14.80$, 6.87 Hz), 4.52 (1 H, dd, $J=14.95$, 6.10 Hz), 4.37 (1 H, dd, $J=9.46$, 2.14 Hz), 4.02 - 4.05 (1 H, m), 3.98 - 4.00 (1 H, m), 3.94 - 3.97 (1 H, m), 3.67 (1 H, dd, $J=11.14$, 4.73 Hz), 3.42 (1 H, d, $J=10.99$ Hz), 3.17 (3 H, s), 2.35 - 2.42 (3 H, m), 1.96 - 2.03 (1 H, m), 1.71 - 1.90 (5 H, m), 1.55 - 1.62 (2 H, m). LCMS (M+H) $^+=$ 542.1.


Example 272

20



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S,4S)-4-methoxy-1-(methylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-4-oxo-. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.34 (1 H, s), 8.63 (1 H, t, $J=6.56$ Hz), 8.48 (1 H, s), 7.39 (2 H, dd, $J=8.39, 5.65$ Hz), 7.16 (2 H, t, $J=8.85$ Hz), 4.49 - 4.61 (2 H, m), 4.26 (1 H, d, $J=9.16$ Hz), 4.02 - 4.08 (1 H, m), 3.97 - 4.02 (1 H, m), 3.95 (1 H, brs), 3.47 (1 H, dd, $J=10.99, 3.97$ Hz), 3.24 (1 H, d, $J=10.68$ Hz), 3.08 (3 H, s), 3.06 (3 H, s), 2.74 - 2.83 (1 H, m), 2.60 - 2.67 (1 H, m), 2.43 - 2.48 (1 H, m), 2.37 (1 H, brs), 2.09 (1 H, ddd, $J=13.81, 9.99, 3.97$ Hz), 1.70 - 1.83 (4 H, m), 1.57 - 1.68 (2 H, m). LCMS (M+H) $=$ 10 578.1.


Example 273



15

7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S,4R)-1-acetyl-4-methoxy-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.43 (1 H, brs), 9.77 (1 H, t, $J=6.26$ Hz), 8.61 (1 H, s), 7.35 - 7.41 (2 H, m), 7.16 (2 H, t, $J=8.70$ Hz), 4.50 - 4.64 (3 H, m), 4.05 - 4.11 (1 H, m), 3.96 - 4.02 (2 H, m), 3.56 - 3.61 (1 H, m), 3.50 - 3.54 (1 H, m), 3.25 (3 H, s), 2.29 - 2.41 (3 H, m), 2.17 - 2.25 (1 H, m), 1.92 - 2.04 (2 H, m), 1.71 - 1.81 (3 H, m), 1.55 - 1.64 (2 H, m). LCMS (M+H) $=$ 20 542.1.


Example 274



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S,4R)-4-methoxy-1-(methylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.34 (1 H, s), 8.63 (1 H, t, $J=6.56$ Hz), 8.48 (1 H, s), 7.39 (2 H, dd, $J=8.39, 5.65$ Hz), 7.16 (2 H, t, $J=8.85$ Hz), 4.49 - 4.61 (2 H, m), 4.26 (1 H, d, $J=9.16$ Hz), 4.02 - 4.08 (1 H, m), 3.97 - 4.02 (1 H, m), 3.95 (1 H, brs), 3.47 (1 H, dd, $J=10.99, 3.97$ Hz), 3.24 (1 H, d, $J=10.68$ Hz), 3.08 (3 H, s), 3.06 (3 H, s), 2.74 - 2.83 (1 H, m), 2.60 - 2.67 (1 H, m), 2.43 - 2.48 (1 H, m), 2.37 (1 H, brs), 2.09 (1 H, ddd, $J=13.81, 9.99, 3.97$ Hz), 1.70 - 1.83 (4 H, m), 1.57 - 1.68 (2 H, m). LCMS (M+H) $+=$ 578.1.

15


Example 275



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-1-acetyl-2-azetidinyl]carbonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.68 (1 H, brs), 10.01 (1 H, t, J=6.26 Hz), 8.98 (1 H, s), 7.24 (1 H, d, J=7.32 Hz), 7.18 (1 H, d, J=5.19 Hz), 7.09 (1 H, t, J=9.16 Hz), 4.80 (1 H, dd, J=8.85, 6.71 Hz), 4.46 - 4.57 (2 H, m), 4.03 - 4.10 (2 H, m), 3.95 - 4.02 (2 H, m), 2.71 - 2.80 (1 H, m), 2.62 - 2.71 (1 H, m), 2.52 - 2.57 (1 H, m), 2.35 - 2.42 (2 H, m), 2.24 - 2.33 (1 H, m), 2.22 (3 H, s), 1.75 - 1.84 (3 H, m), 1.74 (3 H, s), 1.58 - 1.66 (2 H, m). LCMS (M+H)⁺= 512.1.


10

Example 276



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S)-1-(methylsulfonyl)-2-azetidinyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.37 (1 H, brs), 8.76 (1 H, t, J=6.41 Hz), 8.50 (1 H, s), 7.24 (1 H, d, J=7.93 Hz), 7.16 - 7.20 (1 H, m), 7.08 (1 H, t, J=9.00 Hz), 4.78 - 4.83 (1 H, m), 4.53 (1 H, dd, J=14.65, 7.02 Hz), 4.40 (1 H, dd, J=14.50, 5.95 Hz), 4.12 (1 H, dd, J=15.11, 4.12 Hz), 3.90 - 4.02 (2 H, m), 3.64 - 3.70 (1 H, m), 3.21 (3 H, s), 2.70 - 2.78 (1 H, m), 2.58 - 2.66 (1 H, m), 2.53 - 2.58 (1 H, m), 2.38 - 2.48 (2 H, m), 2.23 - 2.28 (1 H, m), 2.21 (3 H, s), 1.93 - 2.02 (1 H, m), 1.76 - 1.86 (2 H, m), 1.65 - 1.76 (1 H, m), 1.57 - 1.64 (1 H, m). LCMS (M+H)⁺= 548.1.


Example 277



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1R,2S)-1-(acetylamino)-2-ethenylcyclopropyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 12.60 (1 H, brs), 9.98 (1 H, t, *J*=6.10 Hz), 8.95 (1 H, s), 8.67 (1 H, s), 7.41 (2 H, dd, *J*=8.24, 5.80 Hz), 7.13 - 7.21 (2 H, m), 5.39 - 5.50 (1 H, m), 5.13 (1 H, d, *J*=17.09 Hz), 4.98 (1 H, d, *J*=10.68 Hz), 4.55 - 4.59 (2 H, m), 4.17 (1 H, dd, *J*=15.11, 4.73 Hz), 3.85 (1 H, d, *J*=14.65 Hz), 2.61 - 2.71 (1 H, m), 2.47 (1 H, m), 2.37 (1 H, brs), 1.89 - 2.04 (2 H, m), 1.71 - 1.82 (4 H, m), 1.60 - 1.69 (2 H, m), 1.51 - 1.60 (2 H, m), 1.48 (1 H, dd, *J*=7.02, 5.49 Hz), 1.09 (1 H, dd, *J*=9.46, 5.19 Hz). LCMS (M+H)⁺= 524.1.

15


Example 278



*7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1*R*,2*S*)-2-ethenyl-1-[(methylsulfonyl)amino]cyclopropyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.45 (1 H, brs), 8.78 - 8.83 (1 H, m), 8.65 (1 H, s), 8.51 (1 H, s), 7.41 (2 H, dd, J=8.09, 5.95 Hz), 7.16 (2 H, t, J=8.70 Hz), 5.58 - 5.68 (1 H, m), 5.24 (1 H, d, J=17.70 Hz), 5.03 (1 H, d, J=10.68 Hz), 4.48 - 4.51 (2 H, m), 4.08 - 4.14 (1 H, m), 3.95 (1 H, brs), 3.00 (3 H, s), 2.78 (1 H, brs), 2.57 - 2.65 (1 H, m), 2.35 - 2.42 (1 H, m), 2.14 - 2.23 (1 H, m), 1.75 - 1.85 (3 H, m), 1.65 - 1.75 (3 H, m), 1.61 (2 H, dd, J=9.61, 5.04 Hz). LCMS (M+H)⁺= 524.1.*


10

Example 279



*15 7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2*S*,4*S*)-4-methoxy-1-[(1-methyl-1*H*-pyrrol-2-yl)carbonyl]-2-pyrrolidinyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.20 (1 H, brs), 9.29 (1 H, brs), 7.87 (1 H, s), 7.09 - 7.20 (2 H, m), 7.02 - 7.09 (1 H, m), 6.87 (1 H, brs), 6.56 (1 H, brs), 6.04 (1 H, brs), 4.86 (1 H, brs), 4.56 (1 H, brs), 4.27 (1 H, d, J=6.41 Hz), 4.04 - 4.10 (2 H, m), 3.95 (1 H, brs), 3.76 (1 H, brs), 3.64 (1 H, brs), 3.60 (3 H, s), 3.27 (3 H, brs), 2.39 (1 H, brs), 2.21 - 2.29 (2 H, m), 2.19 (3 H, brs), 2.13 (1 H, brs), 1.99 (2 H, brs), 1.87 - 1.94 (1 H, m), 1.76 (2 H, brs), 1.59 (2 H, brs). LCMS (M+H)⁺= 621.1.*


Example 280



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S,4S)-1-(5-isoxazolylcarbonyl)-4-methoxy-2-pyrrolidinyl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, DMSO-*d*₆) δ ppm 12.28 (1 H, brs), 9.21 (1 H, t, *J*=6.26 Hz), 8.77 (1 H, d, *J*=1.53 Hz), 8.26 (1 H, s), 7.12 - 7.22 (2 H, m), 7.03 - 7.09 (2 H, m), 4.94 (1 H, d, *J*=7.02 Hz), 4.66 (1 H, dd, *J*=14.50, 7.17 Hz), 4.40 (1 H, dd, *J*=14.50, 5.34 Hz), 4.24 (1 H, dd, *J*=15.11, 4.43 Hz), 4.06 - 4.13 (1 H, m), 3.84 (2 H, d, *J*=11.60 Hz), 3.72 - 3.80 (1 H, m), 3.33 (3 H, s), 2.30 - 2.43 (2 H, m), 2.15 - 2.26 (5 H, m), 2.06 (2 H, q, *J*=6.61 Hz), 1.87 - 1.95 (1 H, m), 1.70 - 1.82 (2 H, m), 1.60 - 1.68 (1 H, m), 1.55 (1 H, brs). LCMS (M+H)⁺= 609.2.

15

Example 281




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[2-methyl-1-oxo-2-(1H-pyrazol-1-yl)propyl]amino]-4-oxo-*. ^1H NMR (500 MHz, DMSO-*d*₆) δ ppm 12.08 (1 H, brs), 9.49 (1 H, brs), 7.97 (1 H, s), 7.48 (1 H, s), 7.22 (1 H, d, *J*=7.32 Hz), 7.16 (2

H, brs), 7.08 (1 H, t, $J=9.00$ Hz), 6.34 (1 H, s), 4.60 (2 H, d, $J=6.10$ Hz), 3.99 (2 H, d, $J=3.36$ Hz), 3.31 (3 H, s), 2.37 (1 H, brs), 2.20 - 2.29 (2 H, m), 2.18 (3 H, s), 1.68 - 1.82 (4 H, m), 1.64 (6 H, s), 1.53 - 1.61 (2 H, m). LCMS (M+H) $+=$ 523.2.

5


Example 282



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(3S)-1-acetyl-3-pyrrolidinyl]carbonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. 1 H NMR (500 MHz, DMSO- d_6) δ ppm 11.93 (1 H, brs), 8.82 - 8.93 (1 H, m), 8.03 (1 H, s), 7.26 (1 H, d, $J=5.80$ Hz), 7.16 - 7.23 (1 H, m), 7.07 - 7.15 (1 H, m), 4.50 - 4.60 (1 H, m), 4.41 - 4.49 (1 H, m), 3.99 - 4.04 (2 H, m), 3.41 - 3.54 (2 H, m), 3.25 - 3.39 (2 H, m), 3.20 (1 H, dd, $J=11.29$, 7.63 Hz), 3.08 - 3.15 (1 H, m), 2.98 - 3.05 (1 H, m), 2.38 - 2.43 (1 H, m), 2.24 - 2.33 (2 H, m), 2.22 (3 H, brs), 1.97 - 2.11 (2 H, m), 1.90 - 1.93 (3 H, m), 1.73 - 1.80 (2 H, m), 1.55 - 1.64 (2 H, m). LCMS (M+H) $+=$ 526.1.

20


Example 283



*7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[{(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(3S)-1-(methylsulfonyl)-3-pyrrolidinyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.93 (1 H, brs), 8.90 (1 H, t, *J*=6.41 Hz), 8.04 (1 H, s), 7.24 (1 H, d, *J*=7.32 Hz), 7.16 - 7.21 (1 H, m), 7.09 - 7.14 (1 H, m), 4.56 (1 H, dd, *J*=14.95, 6.71 Hz), 4.41 (1 H, dd, *J*=14.65, 5.80 Hz), 3.95 - 4.08 (2 H, m), 3.38 - 3.45 (1 H, m), 3.10 - 3.27 (4 H, m), 2.86 (3 H, s), 2.36 - 2.43 (1 H, m), 2.25 - 2.33 (2 H, m), 2.23 (3 H, s), 1.97 - 2.10 (2 H, m), 1.84 - 1.95 (2 H, m), 1.72 - 1.82 (2 H, m), 1.55 - 1.65 (2 H, m). LCMS (M+H)⁺ = 562.2.*


10

Example 284



15 *1H-1,2,3-Triazole-4-carboxylic acid, 1-[2-[[2-[[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]amino]-1,1-dimethyl-2-oxoethyl]-, methyl ester. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.93 - 12.06 (1 H, m), 9.03 (1 H, t, *J*=6.56 Hz), 8.93 (1 H, s), 7.91 (1 H, s), 7.29 (1 H, d, *J*=7.63 Hz), 7.21 (1 H, brs), 7.07 (1 H, t, *J*=9.16 Hz), 4.61 (2 H, d, *J*=6.41 Hz), 4.00 (2 H, d, *J*=3.05 Hz), 3.87 (3 H, s), 2.39 (1 H, brs), 2.19 (3 H, s), 2.08 - 2.16 (2 H, m), 2.04 (2 H, ddd, *J*=13.96, 7.32, 7.10 Hz), 1.68 - 1.81 (8 H, m), 1.53 - 1.62 (2 H, m). LCMS (M+H)⁺ = 582.2.*


Example 285



5 *1H-1,2,3-Triazole-4-carboxylic acid, 1-[2-[2-[[[4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]amino]-1,1-dimethyl-2-oxoethyl]-.* ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 12.04 (1 H, brs), 9.07 (1 H, t, *J*=6.41 Hz), 8.82 (1 H, s), 7.94 (1 H, s), 7.31 (1 H, d, *J*=7.32 Hz), 7.22 (1 H, t, *J*=5.65 Hz), 7.06 (1 H, t, *J*=9.16 Hz), 4.62 (2 H, d, *J*=6.41 Hz), 4.00 (2 H, d, *J*=3.66 Hz), 2.38 (1 H, brs), 2.19 (3 H, s), 2.10 - 2.17 (2 H, m), 1.98 - 2.08 (2 H, m), 1.70 - 1.81 (8 H, m), 1.53 - 1.62 (2 H, m). LCMS (M+H)⁺= 568.0.

Example 286

15




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[2-methyl-2-[4-[(methylamino)carbonyl]-1H-1,2,3-triazol-1-yl]-1-oxopropyl]amino]-4-oxo-.* ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.99 (1 H, brs), 9.03 (1 H, t, *J*=6.41 Hz), 8.93 (1 H, s), 7.91 (1 H, s), 7.29 (1 H, d, *J*=7.32 Hz), 7.20 (1 H, d, *J*=5.80 Hz), 7.07 (1 H, t,

J=9.16 Hz), 4.61 (2 H, d, *J*=6.41 Hz), 4.00 (2 H, d, *J*=3.36 Hz), 3.87 (3 H, s), 2.38 (1 H, brs), 2.19 (3 H, s), 2.09 - 2.16 (2 H, m), 2.00 - 2.09 (2 H, m), 1.70 - 1.81 (8 H, m), 1.54 - 1.62 (2 H, m). LCMS (M+H)⁺= 581.1.

5


Example 287



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[2-[4-[(dimethylamino)carbonyl]-1H-1,2,3-triazol-1-yl]-2-methyl-1-oxopropyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 12.03 (1 H, brs), 9.13 (1 H, t, *J*=6.41 Hz), 8.65 (1 H, s), 7.96 (1 H, s), 7.24 (1 H, d, *J*=7.63 Hz), 7.15 - 7.21 (1 H, m), 7.06 (1 H, t, *J*=9.00 Hz), 4.61 (2 H, d, *J*=6.41 Hz), 4.00 (2 H, d, *J*=3.66 Hz), 3.30 (3 H, s), 3.00 (3 H, s), 2.39 (1 H, brs), 2.13 - 2.21 (5 H, m), 1.98 - 2.06 (2 H, m), 1.71 - 1.80 (8 H, m), 1.55 - 1.63 (2 H, m). LCMS (M+H)⁺= 595.2.

Example 288

20




7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[2-methyl-2-[4-[(4-methyl-1-piperazinyl)carbonyl]-1H-1,2,3-triazol-1-yl]-1-oxopropyl]amino]-4-oxo-

1 ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.92 (1 H, brs), 9.82 (1 H, brs), 9.00 (1 H, t, J=6.41 Hz), 8.76 (1 H, s), 8.01 (1 H, s), 7.23 (1 H, d, J=7.63 Hz), 7.15 - 7.21 (1 H, m), 7.09 (1 H, t, J=9.00 Hz), 4.59 (2 H, d, J=6.41 Hz), 4.01 (2 H, d, J=3.66 Hz), 3.52 (3 H, brs), 3.11 (3 H, brs), 2.85 (3 H, s), 2.38 - 2.42 (1 H, m), 2.14 - 2.23 (5 H, m), 1.99 - 2.06 (2 H, m), 1.72 - 1.81 (8 H, m), 1.55 - 1.64 (2 H, m). LCMS (M+H)⁺= 650.2.


10

Example 289



15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-1-acetyl-4,4-difluoro-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.40 (1 H, brs), 9.59 (1 H, t, J=6.41 Hz), 8.54 (1 H, s), 7.24 (1 H, d, J=7.32 Hz), 7.17 - 7.21 (1 H, m), 7.06 - 7.13 (1 H, m), 4.76 (1 H, dd, J=9.16, 4.27 Hz), 4.47 - 20 4.60 (2 H, m), 4.04 - 4.16 (2 H, m), 3.89 - 4.02 (2 H, m), 2.57 - 2.67 (2 H, m), 2.30 - 2.43 (3 H, m), 2.21 (3 H, s), 1.89 - 2.01 (5 H, m), 1.72 - 1.83 (2 H, m), 1.56 - 1.64 (2 H, m). LCMS (M+H)⁺= 562.0.


Example 290



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S)-4,4-difluoro-1-(methylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-N-[4-fluoro-3-methylphenyl]methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.29 (1 H, s), 8.64 (1 H, t, J =6.71 Hz), 8.48 (1 H, s), 7.24 (1 H, d, J =7.63 Hz), 7.17 - 7.21 (1 H, m), 7.06 - 7.11 (1 H, m), 4.67 (1 H, dd, J =9.31, 4.73 Hz), 4.51 - 4.57 (1 H, m), 4.40 - 4.46 (1 H, m), 4.06 - 4.12 (1 H, m), 3.90 - 3.99 (2 H, m), 3.65 - 3.75 (1 H, m), 3.14 (3 H, s), 2.56 - 2.65 (2 H, m), 2.45 - 2.48 (2 H, m), 2.38 - 2.42 (1 H, m), 2.21 (3 H, s), 1.90 - 1.99 (1 H, m), 1.75 - 1.90 (3 H, m), 1.57 - 1.68 (2 H, m). LCMS (M+H) $+=$ 598.1.

15


Example 291



20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[1-oxo-2-(1H-1,2,4-triazol-1-yl)propyl]amino]*. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 12.08 (1 H, brs), 9.03 (1 H, t, J =6.41 Hz), 8.61 (1 H, s), 8.35 (1 H, s), 7.94 (1 H, s), 7.26 (1 H, d,

J=7.63 Hz), 7.20 (1 H, td, *J*=5.42, 2.59 Hz), 7.08 - 7.14 (1 H, m), 5.33 (1 H, q, *J*=7.12 Hz), 4.62 (1 H, dd, *J*=14.95, 7.02 Hz), 4.50 (1 H, dd, *J*=14.95, 6.10 Hz), 4.10 (1 H, dd, *J*=15.26, 3.97 Hz), 3.93 (1 H, dd, *J*=14.95, 2.75 Hz), 2.39 (1 H, brs), 2.23 - 2.32 (2 H, m), 2.21 (3 H, s), 2.07 - 2.16 (1 H, m), 1.87 - 1.96 (1 H, m), 1.71 - 1.82 (2 H, m), 1.54 - 1.66 (5 H, m). LCMS (M+H)⁺ = 510.1.


Example 292



10

7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[5-(1-methyl-1H-pyrazol-4-yl)-3-isoxazolyl]carbonyl]amino]-4-oxo. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.77 (1 H, brs), 8.92 (1 H, s), 8.74 (1 H, t, *J*=6.41 Hz), 8.33 (1 H, s), 7.96 (1 H, s), 7.14 (1 H, d, *J*=7.32 Hz), 7.06 - 7.11 (1 H, m), 6.99 (1 H, t, *J*=9.00 Hz), 6.89 (1 H, s), 4.42 (2 H, d, *J*=6.41 Hz), 4.05 (2 H, d, *J*=3.66 Hz), 3.93 (3 H, s), 2.39 - 2.48 (3 H, m), 2.16 - 2.24 (2 H, m), 2.14 (3 H, s), 1.80 - 1.89 (2 H, m), 1.62 - 1.72 (2 H, m). LCMS (M+H)⁺ = 562.1.

Example 293




5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[1,2,3,4-tetrahydro-1,3-dimethyl-2,4-dioxo-5-pyrimidinyl]sulfonyl]amino]*. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 11.85 (1 H, brs), 9.25 (1 H, t, $J=6.56$ Hz), 8.40 (1 H, s), 8.02 (1 H, s), 7.28 (1 H, d, $J=7.32$ Hz), 7.17 - 7.22 (1 H, m), 7.07 - 7.15 (1 H, m), 4.50 (2 H, d, $J=6.41$ Hz), 3.96 (2 H, d, $J=3.66$ Hz), 3.41 (3 H, s), 2.96 (3 H, s), 2.39 (1 H, brs), 2.31 (2 H, ddd, $J=14.11, 8.93, 5.65$ Hz), 2.22 (3 H, s), 2.07 (2 H, ddd, $J=13.66, 7.17, 6.94$ Hz), 1.69 - 1.80 (2 H, m), 1.52 - 1.62 (2 H, m). LCMS ($\text{M}+\text{H})+= 589.0$.

10

Example 294


15



20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(1R,3S,5R)-2-(methylsulfonyl)-2-azabicyclo[3.1.0]hex-3-yl]carbonyl]amino]-4-oxo*. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 12.23 (1 H, brs), 8.53 (1 H, t, $J=6.56$ Hz), 8.33 (1 H, s), 7.39 (2 H, dd, $J=8.55, 5.49$ Hz), 7.14 - 7.21 (2 H, m), 4.56 - 4.65 (1 H, m), 4.46 - 4.56 (1

H, m), 4.15 (1 H, dd, $J=15.26, 4.27$ Hz), 3.89 - 3.95 (2 H, m), 3.07 (3 H, s), 2.99 (1 H, td, $J=5.95, 2.44$ Hz), 2.57 - 2.66 (1 H, m), 2.43 - 2.48 (1 H, m), 2.38 - 2.42 (1 H, m), 2.31 - 2.37 (1 H, m), 2.06 (1 H, dt, $J=13.20, 6.68$ Hz), 1.92 - 2.01 (1 H, m), 1.72 - 1.83 (3 H, m), 1.63 - 1.70 (1 H, m), 1.54 - 1.63 (2 H, m), 0.79 (1 H, ddd, $J=7.71, 2.98, 2.75$ Hz), 0.63 (1 H, dt, $J=8.55, 5.80$ Hz). LCMS (M+H) $= 560.2$.


Example 295



10

*7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(2S,4S)-1-[(1,5-dimethyl-1H-pyrazol-3-yl)carbonyl]-4-methoxy-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 12.30 (1 H, brs), 9.29 (1 H, t, $J=6.26$ Hz), 8.08 - 8.17 (1 H, m), 7.04 - 7.19 (3 H, m), 6.24 (1 H, s), 4.83 - 4.94 (1 H, m), 4.64 (1 H, dd, $J=14.65, 7.32$ Hz), 4.48 (1 H, dd, $J=14.65, 5.49$ Hz), 4.26 (1 H, dd, $J=14.95, 4.58$ Hz), 4.01 - 4.11 (2 H, m), 3.94 (2 H, dd, $J=7.48, 4.12$ Hz), 3.75 (3 H, s), 3.31 (3 H, s), 2.34 - 2.41 (2 H, m), 2.16 - 2.24 (6 H, m), 2.09 - 2.16 (2 H, m), 1.99 - 2.06 (1 H, m), 1.90 - 1.97 (1 H, m), 1.68 - 1.86 (3 H, m), 1.48 - 1.67 (2 H, m). LCMS (M+H) $= 636.1$.*


Example 296



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S,4S)-4-methoxy-1-[(5-methyl-3-isoxazolyl)carbonyl]-2-pyrrolidinyl]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 12.28 (1 H, brs), 9.20 (1 H, t, $J=6.26$ Hz), 8.24 (1 H, s), 7.03 - 7.18 (3 H, m), 6.33 (1 H, s), 4.93 (1 H, d, $J=7.02$ Hz), 4.61 (1 H, dd, $J=14.65$, 7.02 Hz), 4.37 (1 H, dd, $J=14.95$, 5.80 Hz), 4.26 (1 H, dd, $J=15.26$, 5.19 Hz), 4.04 - 4.11 (1 H, m), 3.99 - 4.04 (1 H, m), 3.74 - 3.81 (1 H, m), 3.65 - 3.73 (1 H, m), 3.31 (3 H, s), 2.31 - 2.44 (5 H, m), 2.15 - 2.23 (4 H, m), 2.02 (2 H, q, $J=8.04$ Hz), 1.91 (1 H, brs), 1.70 - 1.84 (3 H, m), 1.61 - 1.69 (1 H, m), 1.49 - 1.60 (1 H, m). LCMS (M+H) $+=$ 623.2.

15

Example 297




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S,4S)-4-methoxy-1-[(5-methyl-1,3,4-oxadiazol-2-yl)carbonyl]-2-pyrrolidinyl]carbonyl]amino]-4-oxo-*.
HPLC(A): Solvent(A): 0.1%trifluoroacetic acid/95%water/5%Acetonitirle:

Solvent(B): 0.1%trifluoroacetic acid/95%Acetonitrile/5%water: Column: Sunfire C18 3.5 μ m, 4.5 \times 150mm: Gradient time: 15min: Stop time: 18min: Start B%=10% Final B%=100%: Flow rate: 2ml/min: UV Det: 220/254nm.. HPLC(B): Solvent(A): 10 mM amm. Bicarb (pH=9.5) 95%water/5%Methanol: Solvent(B): 10 mM amm. Bicarb (pH=9.5) 5%water/95%Methanol: Column: Xbridge Phenyl 3.5 μ m, 4.5 \times 150mm: Gradient time: 15min: Stop time: 18min: Start B%=10% Final B%=100%: Flow rate: 2ml/min: UV Det: 220/254nm. LCMS (M+H) $+=$ 624.0. HPLC(A): Rt=6.22min, HPLC(B): Rt=6.99.


10

Example 298



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S,4S)-4-methoxy-1-[(1-methyl-1H-imidazol-2-yl)carbonyl]-2-pyrrolidinyl]carbonyl]amino]-4-oxo-.¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 12.29 (1 H, brs), 9.24 (1 H, t, *J*=6.10 Hz), 8.14 (1 H, s), 7.46 (1 H, s), 7.24 (1 H, brs), 7.01 - 7.18 (3 H, m), 4.93 (1 H, d, *J*=6.71 Hz), 4.57 (1 H, dd, *J*=14.65, 6.71 Hz), 4.34 (1 H, dd, *J*=15.26, 5.80 Hz), 4.13 - 4.20 (1 H, m), 4.07 (1 H, q, *J*=6.21 Hz), 4.01 (1 H, d, *J*=11.29 Hz), 3.91 - 3.98 (1 H, m), 3.79 - 3.87 (1 H, m), 3.71 - 3.76 (3 H, m), 3.27 (3 H, s), 2.37 - 2.42 (1 H, m), 2.27 - 2.36 (2 H, m), 2.18 (3 H, s), 1.90 - 2.00 (2 H, m), 1.74 - 1.89 (2 H, m), 1.51 - 1.71 (4 H, m). LCMS (M+H) $+=$ 622.0.


Example 299



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[{(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(2S,4S)-4-methoxy-1-[(1-methyl-1H-1,2,4-triazol-3-yl)carbonyl]-2-pyrrolidinyl]carbonyl]amino]-4-oxo-}.*
¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.42 (1 H, brs), 9.22 (1 H, t, J=6.41 Hz), 8.61 (1 H, s), 8.31 (1 H, s), 7.17 - 7.22 (2 H, m), 7.01 - 7.08 (1 H, m), 4.91 (1 H, d, J=7.32 Hz), 4.67 - 4.71 (2 H, m), 4.34 (1 H, dd, J=15.11, 5.65 Hz), 4.08 (1 H, q, J=7.63 Hz), 3.93 - 4.00 (1 H, m), 3.89 - 3.93 (3 H, m), 3.79 - 3.87 (2 H, m), 3.35 (3 H, s), 2.39 (1 H, brs), 2.16 - 2.24 (4 H, m), 2.03 - 2.11 (4 H, m), 1.62 - 1.92 (4 H, m), 1.44 - 1.54 (1 H, m). LCMS (M+H)⁺= 623.0.

15

Example 300




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1S,2R)-1-(acetylamino)-2-ethenylcyclopropyl]carbonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-}.*¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.62 (1 H, brs), 9.96 (1 H, t, J=6.10 Hz), 8.95 (1 H, s), 8.68 (1 H, brs), 7.28 (1 H, d, J=7.63 Hz), 7.21 (1 H, t, J=5.80 Hz), 7.08 (1 H, t, J=9.16 Hz), 5.40 - 5.50 (1 H, m), 5.13 (1 H, d, J=17.40 Hz), 4.98 (1 H, d, J=11.29 Hz), 4.47

- 4.59 (2 H, m), 4.17 (1 H, dd, $J=15.11, 4.73$ Hz), 3.85 (1 H, d, $J=14.95$ Hz), 2.62 - 2.71 (1 H, m), 2.37 (1 H, brs), 2.21 (3 H, s), 1.89 - 2.01 (2 H, m), 1.72 - 1.80 (4 H, m), 1.46 - 1.69 (4 H, m), 1.16 - 1.20 (1 H, m), 1.06 - 1.14 (2 H, m). LCMS (M+H) $=$ 538.2.

5


Example 301



10 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(3S)-1-[(1,5-dimethyl-1H-pyrazol-3-yl)carbonyl]-3-pyrrolidinyl]carbonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 11.95 (1 H, brs), 8.83 - 8.96 (1 H, m), 8.06 (1 H, d, $J=9.77$ Hz), 7.01 - 7.29 (3 H, m), 6.40 (1 H, d, $J=2.44$ Hz), 4.48 - 4.64 (1 H, m), 4.30 - 4.48 (1 H, m), 3.98 - 4.04 (2 H, m), 3.90 (1 H, ddd, $J=11.37, 7.71, 4.12$ Hz), 3.73 - 3.79 (3 H, m), 3.60 - 3.68 (1 H, m), 3.52 (1 H, ddd, $J=11.83, 8.01, 3.97$ Hz), 3.28 - 3.45 (1 H, m), 3.01 - 3.18 (1 H, m), 2.40 (1 H, brs), 2.24 - 2.35 (5 H, m), 2.20 (3 H, d, $J=5.19$ Hz), 1.82 - 2.14 (4 H, m), 1.73 - 1.82 (2 H, m), 1.54 - 1.65 (2 H, m). LCMS (M+H) $=$ 606.2.

15
20


Example 302



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[1-(methoxymethyl)cyclopropyl]carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.13 (1 H, s), 8.92 (1 H, t, J=6.26 Hz), 8.69 (1 H, s), 7.23 (1 H, d, J=7.02 Hz), 7.14 - 7.19 (1 H, m), 7.11 (1 H, t, J=9.00 Hz), 4.48 (2 H, d, J=6.10 Hz), 4.02 (2 H, d, J=3.66 Hz), 3.53 (2 H, s), 3.18 (3 H, s), 2.63 - 2.73 (2 H, m), 2.39 (1 H, brs), 2.22 (3 H, s), 1.74 - 1.85 (4 H, m), 1.58 - 1.68 (2 H, m), 0.94 - 0.99 (2 H, m), 0.59 - 0.64 (2 H, m). LCMS (M+H)⁺= 499.2.


10

Example 303



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[[1-(trifluoromethyl)cyclopropyl]carbonyl]amino]-. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 9.26 (1 H, t, J=6.41 Hz), 7.22 (1 H, d, J=7.32 Hz), 7.14 - 7.18 (1 H, m), 7.10 (1 H, t, J=9.16 Hz), 4.43 (2 H, d, J=6.10 Hz), 4.05 (2 H, d, J=3.05 Hz), 2.48 (1 H, brs), 2.16 - 2.26 (5 H, m), 2.02 - 2.11 (2 H, m), 1.82 - 1.92 (2 H, m), 1.73 - 1.82 (4 H, m), 20 1.63 - 1.68 (2 H, m). LCMS (M+H)⁺= 523.1.


Example 304



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[1-(2-pyridinyl)cyclopropyl]carbonyl]amino]*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.85 (1 H, brs), 9.35 (1 H, t, *J*=5.80 Hz), 8.45 (1 H, d, *J*=4.58 Hz), 7.89 (1 H, t, *J*=7.63 Hz), 7.48 (1 H, d, *J*=7.93 Hz), 7.34 - 7.43 (2 H, m), 7.12 (1 H, d, *J*=7.32 Hz),
10 7.01 - 7.09 (2 H, m), 4.47 (2 H, d, *J*=6.41 Hz), 3.99 - 4.04 (2 H, m), 2.39 (1 H, brs), 2.17 (3 H, s), 2.08 - 2.14 (2 H, m), 2.01 (2 H, ddd, *J*=14.50, 7.17, 7.02 Hz), 1.67 - 1.78 (2 H, m), 1.52 - 1.61 (2 H, m), 1.28 - 1.33 (2 H, m), 1.01 - 1.07 (2 H, m). LCMS (M+H)⁺ = 532.16.

15

Example 305




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(1,4,5,6-tetrahydro-6-oxo-3-pyridazinyl)carbonyl]amino]*. ^1H NMR (500 MHz, *DMSO-d*₆) δ ppm 11.79 (1 H, brs), 11.04 (1 H, s), 8.47 (1 H, s), 8.43 (1 H, t, *J*=6.26 Hz), 7.19 (1 H, d, *J*=7.02 Hz), 7.04 - 7.15 (2 H, m), 4.53 (2 H, d, *J*=6.10 Hz), 4.03 (2 H, d, *J*=3.05

Hz), 2.57 (2 H, t, $J=8.24$ Hz), 2.40 - 2.47 (3 H, m), 2.27 (2 H, t, $J=8.39$ Hz), 2.20 (3 H, s), 2.00 - 2.10 (2 H, m), 1.77 - 1.87 (2 H, m), 1.61 - 1.70 (2 H, m). LCMS (M+H) $+=$ 511.18.


5

Example 306



- 7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-
 10 [(cyclopropylmethyl)sulfonyl]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-
 4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ^1H NMR (500 MHz, DMF) δ ppm 9.56 (1
 H, t, $J=6.26$ Hz), 8.98 (2 H, brs), 7.68 (1 H, d, $J=7.63$ Hz), 7.58 - 7.63 (1 H, m), 7.53
 (1 H, t, $J=9.16$ Hz), 4.87 (2 H, d, $J=6.10$ Hz), 4.48 (2 H, d, $J=3.66$ Hz), 4.08 (2 H, d,
 $J=7.32$ Hz), 2.89 (1 H, brs), 2.60 - 2.68 (5 H, m), 2.43 - 2.51 (2 H, m), 2.26 - 2.35 (2
 H, m), 2.19 - 2.26 (2 H, m), 1.62 - 1.71 (1 H, m), 1.05 - 1.11 (2 H, m), 0.90 (2 H, q,
 $J=4.88$ Hz). LCMS (M+H) $+=$ 505.1.

Example 307




20

Ethanediamide, N'-[2-[[[5-fluoro[1,1'-biphenyl]-2-yl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-. ^1H NMR (400 MHz, CDCl_3) δ ppm 12.01 (s, 1 H), 8.30 (s, 2 H), 7.31 - 7.46 (m, 6 H), 7.03 (td, $J=8.47$, 5 2.64 Hz, 1 H), 6.97 (dd, $J=9.54$, 2.76 Hz, 1 H), 4.50 (d, $J=6.02$ Hz, 2 H), 4.15 (d, $J=3.76$ Hz, 2 H), 3.27 (s, 3 H), 2.79 (s, 3 H), 2.46 - 2.62 (m, 3 H), 1.90 - 2.11 (m, 4 H), 1.64 - 1.76 (m, 2 H). LCMS ($^+$ ESI, $\text{M}+\text{H}^+$) m/z 548.2. HPLC purity: 98%.

Example 308


10



Butanoic acid, 4-[[2-[[[4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-methoxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]amino]-4-oxo-. ^1H NMR (500 MHz, CDCl_3) δ ppm 8.37 (t, 1 H), 8.29 (s, 1 H), 7.30 (dd, $J=8.70$, 5.34 Hz, 2 H), 7.01 (t, 2 H), 4.58 (d, $J=5.80$ Hz, 2 H), 4.06 (d, $J=3.66$ Hz, 2 H), 3.96 (s, 3 H), 2.86 - 2.97 (m, 2 H), 2.58 - 2.64 (m, 2 H), 2.51 - 2.58 (m, 2 H), 2.42 (s, 1 H), 1.91 - 2.00 (m, 2 H), 1.76 - 1.86 (m, 2 H), 1.59 - 1.68 (m, 2 H). LCMS ($^+$ ESI, $\text{M}+\text{H}^+$) m/z 487.3. HPLC purity: 96%.

15


Example 309



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(4-pyridinylcarbonyl)amino]-. ^1H NMR (500 MHz, DMSO-D6) δ ppm 8.83 (s, 1 H), 8.71 (d, $J=5.49$ Hz, 2 H), 8.66 (t, $J=6.26$ Hz, 1 H), 7.85 (d, $J=4.88$ Hz, 2 H), 7.02 (dd, $J=18.92, 9.16$ Hz, 2 H), 6.93 - 6.98 (m, 1 H), 4.33 (d, $J=6.10$ Hz, 2 H), 4.02 - 4.07 (m, 2 H), 2.42 - 2.46 (m, 1 H), 2.33 - 2.42 (m, 2 H), 2.18 - 2.25 (m, 2 H), 2.16 (s, 3 H), 1.79 - 1.89 (m, 2 H), 1.66 (d, $J=5.49$ Hz, 2 H). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 492.1. HPLC purity: 91%.

10


Example 310



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[methyl(4-pyridinylcarbonyl)amino]-4-oxo-. ^1H NMR (500 MHz, DMSO-d6) δ ppm (major rotamer) 8.49 (d, $J=5.19$ Hz, 2 H), 7.36 (dd, $J=8.24, 5.80$ Hz, 2 H), 7.30 (br. s., 2 H), 7.16 (t, $J=8.70$ Hz, 2 H), 4.01 - 4.09 (m, 4 H), 2.96 - 3.01 (m, 3 H), 2.39 - 2.46 (m, 1 H), 1.23 - 2.19 (m, 8 H). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 492.2. HPLC purity: 95%.

20


Example 311



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(2-pyrimidinylcarbonyl)amino]-. ^1H NMR (500 MHz, DMSO-d6) δ ppm 11.88 (br. s., 1 H), 9.62 (br. s., 1 H), 8.87 (br. s., 1 H), 8.74 (d, $J=2.75$ Hz, 2 H), 7.61 (br. s., 1 H), 5 7.35 (br. s., 2 H), 7.12 - 7.23 (m, 2 H), 4.58 (d, $J=4.58$ Hz, 2 H), 4.07 (br. s., 2 H), 2.64 (br. s., 2 H), 2.42 - 2.47 (m, 1 H), 2.07 (br. s., 2 H), 1.88 (br. s., 2 H), 1.70 (br. s., 2 H). LCMS ($^+\text{ESI, M+H}^+$) m/z 479.1. HPLC purity: 96%.

Example 312


10



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(3-pyridinylcarbonyl)amino]-. ^1H NMR (500 MHz, CHLOROFORM-d) δ ppm 9.00 (d, $J=1.83$ Hz, 1 H), 8.70 (dd, $J=4.88, 1.53$ Hz, 1 H), 8.10 (d, $J=8.24$ Hz, 1 H), 7.75 (s, 1 H), 7.32 (dd, $J=8.09, 5.04$ Hz, 1 H), 7.02 - 7.07 (m, 1 H), 6.96 - 7.01 (m, 1 H), 6.89 (t, $J=8.85$ Hz, 1 H), 4.46 (d, $J=6.10$ Hz, 2 H), 4.21 (d, $J=3.66$ Hz, 2 H), 3.49 (s, 1 H), 2.80 - 2.90 (m, 2 H), 2.53 (br. s., 1 H), 2.22 (d, $J=1.83$ Hz, 3 H), 1.99 - 2.12 (m, 4 H), 20 1.71 - 1.87 (m, 2 H). LCMS ($^+\text{ESI, M+H}^+$) m/z 492.1. HPLC purity: 97%.

Example 313


25



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(2-pyridinylcarbonyl)amino]-. ^1H NMR (500 MHz, CHLOROFORM-d) δ ppm 12.03 (br. s., 1 H), 10.14 (s, 1 H), 8.12 (d, $J=7.63$ Hz, 1 H), 8.08 (t, $J=5.95$ Hz, 1 H), 7.80 (5 td, $J=7.71$, 1.68 Hz, 1 H), 7.74 (d, $J=4.88$ Hz, 1 H), 7.23 - 7.29 (m, 1 H), 7.17 (d, $J=7.32$ Hz, 1 H), 7.10 - 7.15 (m, 1 H), 6.97 (t, $J=8.85$ Hz, 1 H), 4.62 (d, $J=6.10$ Hz, 2 H), 4.21 (d, $J=3.97$ Hz, 2 H), 3.05 - 3.13 (m, 2 H), 2.47 - 2.58 (m, 1 H), 2.23 (d, $J=1.83$ Hz, 3 H), 2.03 - 2.12 (m, 2 H), 1.88 - 1.97 (m, 2 H), 1.70 - 1.81 (m, 2 H). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 492.1. HPLC purity: 92%.


10

Example 314



15 7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(3-pyridazinylcarbonyl)amino]-. ^1H NMR (500 MHz, CHLOROFORM-d) δ ppm 12.03 (br. s., 1 H), 10.84 (s, 1 H), 9.30 (dd, $J=5.04$, 1.68 Hz, 1 H), 8.49 (t, $J=6.56$ Hz, 1 H), 8.30 (dd, $J=8.55$, 1.83 Hz, 1 H), 7.69 (dd, $J=8.55$, 4.88 Hz, 1 H), 7.33 (dd, $J=7.17$, 1.68 Hz, 1 H), 7.26 - 7.30 (m, 1 H), 6.85 - 6.94 (m, 1 H), 4.67 (d, $J=6.71$ Hz, 2 H), 4.20 (d, $J=3.66$ Hz, 2 H), 3.08 (ddd, $J=14.42$, 9.54, 5.34 Hz, 2 H), 2.53 (br. s., 1 H), 2.22 (d, $J=1.83$ Hz, 3 H), 2.05 - 2.16 (m, 2 H), 1.91 - 2.00 (m, 2 H), 1.73 - 1.84 (m, 2 H). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 493.0. HPLC purity: 95%.

Example 315




5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(3-quinolinylcarbonyl)amino]-.* ^1H NMR (500 MHz, CHLOROFORM-d) δ ppm 9.62 (br. s., 1 H), 9.11 (s, 1 H), 8.39 (d, $J=8.85$ Hz, 1 H), 8.09 - 8.16 (m, 2 H), 8.04 (d, $J=7.93$ Hz, 1 H), 7.91 (t, $J=7.63$ Hz, 1 H), 7.50 (br. s., 1 H), 6.95 (dd, $J=8.24, 5.49$ Hz, 2 H), 6.58 (t, $J=8.55$ Hz, 2 H), 4.41 (d, $J=6.10$ Hz, 2 H), 4.23 (d, $J=3.66$ Hz, 2 H), 2.62 - 2.72 (m, 2 H), 2.56 (br. s., 1 H), 2.28 (br. s., 2 H), 1.98 - 2.08 (m, 2 H), 1.75 (br. s., 2 H). LCMS ($^+\text{ESI}, \text{M}+\text{H}^+$) m/z 528.0. HPLC purity: 91%.

10

Example 316

15




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[2-(2,2,2-trifluoroethoxy)-4-pyridinyl]carbonyl]amino]-.* ^1H NMR (500 MHz, DMSO-d6) δ ppm 11.78 (br. s., 1 H), 8.74 (t, $J=6.26$ Hz, 1 H), 8.58 (d, $J=2.14$ Hz, 1 H), 8.55 (s, 1 H), 8.10 (dd, $J=8.55, 2.44$ Hz, 1 H), 7.14 - 7.20 (m, 2 H), 7.04 - 7.11 (m, 2 H), 6.94

(d, J=8.55 Hz, 1 H), 5.05 (q, J=9.05 Hz, 2 H), 4.38 (d, J=6.10 Hz, 2 H), 4.05 (d, J=3.66 Hz, 2 H), 2.45 (br. s., 1 H), 2.36 - 2.44 (m, 2 H), 2.15 - 2.24 (m, 2 H), 1.84 (td, J=13.89, 5.49 Hz, 2 H), 1.61 - 1.69 (m, 2 H). LCMS (⁺ESI, M+H⁺) m/z 575.9. HPLC purity: 91%.

5


Example 317



10 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[2-(4-morpholinyl)-4-pyridinyl]carbonyl]amino]-4-oxo-*. ¹H NMR (500 MHz, MeOD) δ ppm 7.94 (br. s., 1 H), 7.48 (br. s., 1 H), 7.20 (br. s., 2 H), 7.10 (br. s., 1 H), 6.96 - 7.04 (m, 2 H), 4.49 (br. s., 2 H), 4.17 - 4.23 (m, 2 H), 3.85 (br. s., 4 H), 3.65 (br. s., 4 H), 2.55 (br. s., 1 H), 2.36 - 2.47 (m, 4 H), 2.02 (d, J=5.49 Hz, 2 H), 1.78 (d, J=5.49 Hz, 2 H). LCMS (⁺ESI, M+H⁺) m/z 563.0. HPLC purity: 93%.

15

Example 318




20

7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[2-(acetylamino)-4-pyridinyl]carbonyl]amino]-N-[{(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ^1H NMR (500 MHz, MeOD) δ ppm 8.45 (br. s., 1 H), 8.38 (br. s., 1 H), 7.48 (br. s., 1 H), 7.24 (br. s., 2 H), 6.98 (br. s., 2 H), 4.53 (br. s., 2 H), 4.16 (br. s., 2 H), 2.74 (br. s., 2 H), 2.49 (br. s., 1 H), 2.20 (s, 3 H), 2.08 - 2.16 (m, 2 H), 2.03 (br. s., 2 H), 1.70 - 1.80 (m, 2 H). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 535.1. HPLC purity: 97%.


Example 319

10



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[methyl(3-pyridinylcarbonyl)amino]-4-oxo-. ^1H NMR (400 MHz, DMSO-d6) (spectrum complicated by rotamers) δ ppm 11.91 (s, 1 H), 8.58 - 8.64 (m, 2 H), 8.49 - 8.57 (m, 1 H), 7.72 (d, $J=7.28$ Hz, 1 H), 7.31 (dd, $J=7.53, 5.02$ Hz, 1 H), 7.10 - 7.18 (m, 2 H), 7.04 (t, $J=8.91$ Hz, 1 H), 4.28 - 4.67 (m, 3 H), 3.76 (br. s., 1 H), 3.09 (s, 3 H), 2.44 - 2.49 (m, 1 H), 2.16 (s, 3 H), 2.10 - 2.24 (m, 2 H), 1.93 (br. s., 2 H), 1.76 (br. s., 2 H), 1.35 - 1.53 (m, 2 H). LCMS (^+ESI , $\text{M}+\text{H}^+$) m/z 506.1. HPLC purity: 91%.


Example 320



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[6-(acetylamino)-3-pyridinyl]carbonyl]amino]-N-[{(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 11.19 (br. s., 1 H), 8.51 (d, *J*=2.14 Hz, 1 H), 8.46 (d, *J*=8.85 Hz, 1 H), 8.14 (dd, *J*=9.00, 2.29 Hz, 1 H), 5 7.53 (s, 1 H), 7.15 - 7.21 (m, 3 H), 6.94 - 7.02 (m, 2 H), 4.52 (d, *J*=5.80 Hz, 2 H), 4.21 (d, *J*=3.66 Hz, 2 H), 2.71 - 2.81 (m, 2 H), 2.54 (br. s., 1 H), 2.36 (s, 3 H), 1.95 - 2.13 (m, 4 H), 1.74 (br. s., 2 H). LCMS (^+ESI , M+H $^+$) m/z 535.0. HPLC purity: 81%.


10

Example 321




7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[5-(acetylamino)-3-pyridinyl]carbonyl]amino]-N-[{(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-. ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 9.93 (s, 1 H), 9.44 (d, *J*=1.53 Hz, 1 H), 9.22 (s, 1 H), 8.80 (d, *J*=1.22 Hz, 1 H), 8.52 (s, 1 H), 7.89 (t, *J*=6.10 Hz, 1 H), 7.12 (dd, *J*=8.70, 5.34 Hz, 2 H), 6.85 - 6.92 (m, 2 H), 4.51 (d, *J*=6.10 Hz, 2 H), 4.22 (d, *J*=3.66 Hz, 2 H), 2.70 (ddd, *J*=14.19, 9.00, 5.80 Hz, 2 H), 20 2.56 (br. s., 1 H), 2.26 (s, 3 H), 2.19 - 2.26 (m, 2 H), 1.99 - 2.09 (m, 2 H), 1.69 - 1.80 (m, 2 H). LCMS (^+ESI , M+H $^+$) m/z 535.0. HPLC purity: 98%.

Example 322




5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[methyl(3-pyridinylcarbonyl)amino]-4-oxo.* LCMS (⁺ESI, M+H⁺) m/z 492.0. HPLC purity: 97%.

10 Example 323



15 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[6-(trifluoromethyl)-3-pyridinyl]carbonyl]amino].* ¹H NMR (500 MHz, DMSO-d6) δ ppm 11.53 - 11.83 (m, 1 H), 9.11 (br. s., 2 H), 8.36 (br. s., 1 H), 7.89 (d, *J*=7.02 Hz, 1 H), 7.19 (br. s., 2 H), 7.06 (br. s., 2 H), 4.41 (d, *J*=3.05 Hz, 2 H), 4.04 (br. s., 2 H), 2.44 (br. s., 1 H), 2.12 (br. s., 2 H), 1.80 - 1.94 (m, 4 H), 1.67 (br. s., 2 H). ¹⁹F NMR (471 MHz, DMSO-d6) δ ppm -67.12 (br. s., 3 F), -116.29 (br. s., 1 F). LCMS (⁺ESI, M+H⁺) m/z 545.9. HPLC purity: 93%.


Example 324



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(2-pyridinylamino)carbonyl]amino]-.* ^1H NMR (400 MHz, DMF) δ ppm 12.32 (br. s., 1 H), 9.27 (br. s., 1 H), 9.04 (br. s., 1 H), 8.52 (br. s., 1 H), 8.33 (d, $J=4.27$ Hz, 1 H), 7.76 - 7.90 (m, 2 H), 7.40 - 7.50 (m, 2 H), 7.24 (t, $J=8.66$ Hz, 2 H), 7.08 (t, $J=5.90$ Hz, 1 H), 4.60 (d, $J=6.27$ Hz, 2 H), 4.29 (d, $J=3.26$ Hz, 2 H), 2.76 - 2.87 (m, 2 H), 2.63 (br. s., 1 H), 2.19 - 2.33 (m, 2 H), 2.00 - 2.14 (m, 2 H), 1.90 (br. s., 2 H). LCMS ($^{\pm}$ ESI, $\text{M}+\text{H}^+$) m/z 493.2. HPLC purity: 98%.

Example 325

15




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(2-pyridinylamino)carbonyl]amino]-.* ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 11.92 (br. s., 1 H), 9.39 (s, 1 H), 8.92 (br. s., 1 H), 8.14 (d, $J=4.27$ Hz, 1 H), 7.63 - 7.68 (m, 1 H), 7.52 - 7.59 (m, 1 H), 7.09 (d, $J=7.02$ Hz, 1 H), 6.96 - 7.06 (m, 2 H), 6.86 - 6.94

(m, 1 H), 4.38 (d, $J=6.41$ Hz, 2 H), 4.05 (d, $J=3.36$ Hz, 2 H), 2.55 - 2.65 (m, 2 H), 2.42 (br. s., 1 H), 2.17 (s, 3 H), 1.94 - 2.03 (m, 2 H), 1.82 (td, $J=13.89$, 5.49 Hz, 2 H), 1.65 (d, $J=5.80$ Hz, 2 H). LCMS ($^+ \text{ESI}$, $\text{M}+\text{H}^+$) m/z 507.1. HPLC purity: 99%.

5


Example 326



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[(3-pyridinylamino)carbonyl]amino]. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 12.16 (br. s., 1 H), 10.15 (br. s., 1 H), 9.17 - 9.24 (m, 1 H), 9.02 (br. s., 1 H), 8.43 (d, $J=4.88$ Hz, 1 H), 8.18 (d, $J=8.24$ Hz, 1 H), 7.82 (dd, $J=8.24$, 5.49 Hz, 1 H), 7.35 (s, 1 H), 7.15 (d, $J=7.32$ Hz, 1 H), 7.03 - 7.11 (m, 1 H), 6.92 (t, $J=9.00$ Hz, 1 H), 4.45 (d, $J=5.80$ Hz, 2 H), 4.05 (br. s., 2 H), 2.46 - 2.53 (m, 2 H), 2.43 (br. s., 1 H), 2.14 (s, 3 H), 1.97 - 2.07 (m, 2 H), 1.82 (d, $J=5.19$ Hz, 2 H), 1.65 (br. s., 2 H). LCMS ($^+ \text{ESI}$, $\text{M}+\text{H}^+$) m/z 507.0. HPLC purity: 94%.

20


Example 327



*7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[(methylamino)carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, CHLOROFORM-d) δ ppm 11.97 (br. s., 1 H), 8.03 - 8.12 (m, 1 H), 7.10 (d, *J*=7.32 Hz, 1 H), 7.05 (td, *J*=5.26, 2.29 Hz, 1 H), 6.91 (t, *J*=8.85 Hz, 1 H), 4.46 (d, *J*=6.10 Hz, 2 H), 4.07 (d, *J*=3.36 Hz, 2 H), 2.46 (s, 3 H), 2.44 (br. s., 1 H), 2.34 (ddd, *J*=14.11, 8.93, 5.65 Hz, 2 H), 2.22 (d, *J*=1.53 Hz, 3 H), 2.05 - 2.15 (m, 2 H), 1.82 - 1.94 (m, 2 H), 1.54 - 1.67 (m, 2 H). LCMS (ESI, M+H⁺) m/z 444.1. HPLC purity: 98%.*


10

Example 328



*7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[methyl[(methyl-2-pyridinylamino)carbonyl]amino]-4-oxo-. ¹H NMR (500 MHz, CHLOROFORM-d) δ ppm 8.31 (d, *J*=4.27 Hz, 1 H), 7.98 (t, *J*=5.95 Hz, 1 H), 7.73 (t, *J*=7.93 Hz, 1 H), 7.19 (d, *J*=8.55 Hz, 1 H), 7.15 (dd, *J*=8.39, 5.34 Hz, 2 H), 7.09 (t, *J*=6.10 Hz, 1 H), 6.94 (t, *J*=8.70 Hz, 2 H), 4.46 (br. s., 2 H), 3.49 (s, 3 H), 3.24 (s, 2 H), 2.91 (s, 3 H), 2.50 (br. s., 1 H), 1.43 - 2.04 (m, 8 H). LCMS (ESI, M+H⁺) m/z 521.2. HPLC purity: 93%.*


Example 329



5 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[(methyl-2-pyridinylamino)carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 11.71 (s, 1 H), 9.80 (s, 1 H), 8.09 - 8.17 (m, 2 H), 7.66 - 7.72 (m, 1 H), 6.94 (dd, *J*=7.32, 4.88 Hz, 1 H), 6.84 (d, *J*=8.55 Hz, 1 H), 6.81 (d, *J*=6.41 Hz, 1 H), 6.70 - 6.78 (m, 2 H), 4.24 (d, *J*=6.41 Hz, 2 H), 4.18 (d, *J*=3.97 Hz, 2 H), 3.07 (s, 3 H), 2.48 (d, *J*=2.44 Hz, 1 H), 2.21 - 2.36 (m, 4 H), 2.16 (d, *J*=1.83 Hz, 3 H), 1.84 - 1.94 (m, 2 H), 1.61 - 1.72 (m, 2 H). LCMS († ESI, M+H $^+$) m/z 521.0. HPLC purity: 96%.

Example 330

15




20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[[(phenylsulfonyl)amino]carbonyl]amino]-4-oxo-*. ^1H NMR (500 MHz, CHLOROFORM-*d*) δ ppm 12.56 (br. s., 1 H), 9.06 (s, 1 H), 8.76 (t, *J*=6.10 Hz, 1 H), 7.67 - 7.70 (m, 2 H), 7.63 (t, *J*=7.48 Hz, 1 H), 7.45 (t, *J*=7.93 Hz, 2 H), 7.18 - 7.25 (m, 3 H), 6.75 - 6.82 (m, 2 H), 4.53 (d, *J*=6.41 Hz, 2 H), 4.16 (d, *J*=3.97 Hz, 2 H), 2.80 - 2.90 (m, 2

H), 2.45 (d, $J=1.83$ Hz, 1 H), 1.89 - 1.99 (m, 2 H), 1.59 - 1.72 (m, 4 H). LCMS ($^+$ ESI, M+H $^+$) m/z 556.2. HPLC purity: 90%.

Example 331


5



7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[
10 (methylamino)thioxomethyl]amino]-4-oxo-. 1 H NMR (500 MHz, DMSO-d6) δ ppm 11.98 (br. s., 1 H), 8.97 (br. s., 1 H), 7.82 (br. s., 1 H), 7.39 (dd, $J=8.24, 5.80$ Hz, 3 H), 7.15 - 7.20 (m, 2 H), 4.51 (d, $J=6.41$ Hz, 2 H), 4.00 (d, $J=3.66$ Hz, 2 H), 2.72 - 2.88 (m, 5 H), 2.41 (br. s., 1 H), 2.05 (d, $J=4.88$ Hz, 2 H), 1.79 (td, $J=14.04, 5.80$ Hz, 2 H), 1.57 - 1.65 (m, 2 H). LCMS ($^+$ ESI, M+H $^+$) m/z 446.1. HPLC purity: 92%.

15

Example 332




20 7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[
21 (benzoylamino)thioxomethyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo- 1 H NMR (500 MHz, CHLOROFORM-d) δ ppm 12.51 (s, 1 H), 12.48 (s, 1 H), 9.45 (t, $J=6.26$ Hz, 1 H), 8.86 (s, 1 H), 7.63 (t, $J=7.48$ Hz, 1 H), 7.57 (d, $J=7.32$ Hz, 2 H), 7.46 (t, $J=7.93$ Hz, 2 H), 7.33 (dd, $J=8.55, 5.19$ Hz, 2

H), 6.90 - 6.98 (m, 2 H), 4.70 (d, $J=6.41$ Hz, 2 H), 4.22 (d, $J=3.66$ Hz, 2 H), 3.81 - 3.92 (m, 2 H), 2.53 (br. s., 1 H), 1.99 - 2.12 (m, 2 H), 1.69 - 1.83 (m, 4 H). LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z 536.2. HPLC purity: 90%.

5


Example 333



- 10 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[(aminothioxomethyl)amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-*. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 9.10 (br. s., 1 H), 7.98 (br. s., 1 H), 7.41 (dd, $J=8.39, 5.65$ Hz, 2 H), 7.15 - 7.23 (m, 2 H), 4.52 (d, $J=6.41$ Hz, 2 H), 3.99 (br. s., 2 H), 3.37 - 3.55 (m, 4 H), 2.38 - 2.43 (m, 1 H), 1.71 - 1.84 (m, 2 H), 1.60 (d, $J=1.53$ Hz, 2 H). LCMS ($^+ \text{ESI}, \text{M}+\text{H}^+$) m/z 432.1. HPLC purity: 94%.

15

Example 334




- 20 *7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-(1H-imidazol-2-ylamino)-4-oxo-*. ^1H NMR (400 MHz, DMF) δ ppm 11.40 - 11.91 (m, 2 H), 9.81 (br. s., 1 H), 7.55 (t, $J=6.27$ Hz, 2 H), 7.32 (t, $J=8.41$ Hz, 2 H), 6.71 (s, 2 H), 6.45 (br. s., 1 H), 4.69 (d, $J=6.02$ Hz, 2 H), 4.24 (d, $J=3.51$ Hz, 2 H), 2.56 - 2.75 (m, 3 H), 2.20 -

2.34 (m, 2 H), 2.04 (dd, $J=13.80, 8.78$ Hz, 2 H), 1.85 (d, $J=6.78$ Hz, 2 H). LCMS ($^+$ ESI, $M+H^+$) m/z 439.1. HPLC purity: 98%.

Example 335

5





7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[(1Z)-(cyanoamino)(methylamino)methylene]amino]-N-[(4-fluoro-3-methylphenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-.



10 1 H NMR (500 MHz, DMSO- d_6) δ ppm 11.95 (br. s., 1 H), 8.95 (t, $J=6.26$ Hz, 1 H), 7.34 (br. s., 1 H), 7.21 - 7.25 (m, 1 H), 7.14 - 7.20 (m, 1 H), 7.08 - 7.14 (m, 1 H), 6.86 (br. s., 1 H), 4.44 (d, $J=6.41$ Hz, 2 H), 4.00 (d, $J=3.66$ Hz, 2 H), 2.68 (d, $J=4.58$ Hz, 3 H), 2.49 - 2.53 (m, 2 H), 2.42 (br. s., 1 H), 2.22 (d, $J=1.53$ Hz, 3 H), 1.94 - 2.03 (m, 2 H), 1.84 (d, $J=5.19$ Hz, 2 H), 1.62 (d, $J=6.71$ Hz, 2 H). LCMS ($^+$ ESI, $M+H^+$) m/z 468.0. HPLC purity: 85%.



The compounds in Table 5 were synthesized using methods similar to those described previously. After synthesis the compounds were analyzed reverse phase LCMS on an HPLC system coupled to Waters LCT mass spectrometer. LC
20 Conditions; 50 x 4.6 mm, 5 micron Supelco Ascentis C18 column; mobile phase A = 10 mM NH₄OAc in 5:95 acetonitril/water mobile phase B = 10 mM NH₄OAc in 95:5 acetonitril/water; 2 mL/min flow rate; gradient = 0 to 100% mobile phase B over 8 min.



25

Table 5.

Example	Structure	Observed MS	HPLC retention time
336	 <p><i>N'</i>-(3-Hydroxy-2-((2-(1-methyl-2-pyrrolidinyl)ethyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	475.2	1.255
337	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((3-(2-oxo-1-pyrrolidinyl)propyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	489.2	1.686



Example	Structure	Observed MS	HPLC retention time
338	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((tetrahydro-2-furanylmethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	448.2	1.914
339	 <p><i>N'</i>-(3-Hydroxy-2-((1-methyl-3-phenylpropyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	496.3	3.786



Example	Structure	Observed MS	HPLC retention time
340	 <p><i>N'</i>-(2-((2,2-Diphenylethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	544.3	3.914
341	 <p><i>N'</i>-(3-Hydroxy-2-((2-methylpropyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	420.3	2.592



Example	Structure	Observed MS	HPLC retention time
342	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-phenylamino)ethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	483.2	1.947
343	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-phenylethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	468.2	3.032



Example	Structure	Observed MS	HPLC retention time
344	<p><i>N'</i>-(2-((2-(2-chlorophenyl)ethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	502.2	3.391
345	<p><i>N'</i>-(2-((2-(4-chlorophenyl)ethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	502.2	3.464



Example	Structure	Observed MS	HPLC retention time
346	<p><i>N'</i>-(3-Hydroxy-2-((2-(4-methoxyphenyl)ethyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	498.3	2.983
347	<p><i>N'</i>-(3-Hydroxy-2-((2-(4-hydroxyphenyl)ethyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	484.2	2.195



Example	Structure	Observed MS	HPLC retention time
348	 <p><i>N'</i>-(2-((3,3-Dimethylbutyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-<i>N,N</i>-dimethylethanediamide</p>	448.3	3.434
349	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((3-phenylpropyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-<i>N,N</i>-dimethylethanediamide</p>	482.2	3.387

Example	Structure	Observed MS	HPLC retention time
350	 <p><i>N'</i>-(3-Hydroxy-2-((3-(1<i>H</i>-imidazol-1-yl)propyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	472.2	1.219
351	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-(4-sulfamoylphenyl)ethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	547.3	1.95

Example	Structure	Observed MS	HPLC retention time
352	 <p><i>N'</i>-(2-((Cyclopropylmethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	418.2	2.356
353	 <p><i>N'</i>-(2-((2-(3-Chlorophenyl)ethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	502.2	3.433

Example	Structure	Observed MS	HPLC retention time
354	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-(2-thiophenyl)ethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	474.2	2.859
355	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-phenoxyethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	484.2	2.98



Example	Structure	Observed MS	HPLC retention time
356	 <p><i>N'</i>-(2-((1-Benzyl-3-pyrrolidinyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	523.3	1.868
357	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-(((1<i>S</i>,2<i>R</i>)-2-phenylcyclopropyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	480.2	3.366



Example	Structure	Observed MS	HPLC retention time
358	 <p>Ethanediamide, <i>N'</i>-[2-[[2-(4-fluorophenyl)ethyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl]-<i>N,N</i>-dimethyl-</p>	486.2	3.108
359	 <p>Ethanediamide, <i>N,N</i>-dimethyl-<i>N'</i>-[6,7,8,9-tetrahydro-3-hydroxy-4-oxo-2-[[[(2-phenyl-4-thiazolyl)methyl]amino]carbonyl]-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl]-</p>	537.2	3.263


Example	Structure	Observed MS	HPLC retention time
360	<p>Ethanediamide, <i>N'</i>-[2-[(2-benzothiazolylmethyl)amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl]-<i>N,N</i>-dimethyl-</p>	511.2	2.715



The compounds in Table 2 were synthesized using methods similar to those described previously. After synthesis the compounds were analyzed reverse phase LCMS on a Waters ZQ HPLC system coupled to a ESI mass spectrometer. LC
5 Conditions; 50 x 4.66 mm, 2.7 micron Supelco Ascentis C18 columns; mobile phase A = 0.1% trifluoroacetic acid in water, mobile phase B = acetonitrile; 2 mL/min flow rate; gradient = 0 to 95% mobile phase B over 7 min.



Table 2

Example	Structure	Observed MS	HPLC retention time
361	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-thiophenylmethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-<i>N,N</i>-dimethylethanediamide</p>	458.5	3.21
362	 <p><i>N'</i>-(2-((1,3-Benzodioxol-5-ylmethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-<i>N,N</i>-dimethylethanediamide</p>	496.6	3.51



Example	Structure	Observed MS	HPLC retention time
363	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-pyridinylmethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	453.6	2.42
364	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((3-pyridinylmethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-N,N-dimethylethanediamide</p>	453.5	2.26



Example	Structure	Observed MS	HPLC retention time
365	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((4-pyridinylmethyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl)-<i>N,N</i>-dimethylethanediamide</p>	453.6	2.21
366	<p>Ethanediamide, <i>N,N</i>-dimethyl-<i>N'</i>-[6,7,8,9-tetrahydro-3-hydroxy-4-oxo-2-[(1-phenylethyl)amino]carbonyl]-7,10-ethanopyrimido[1,2-<i>a</i>]azepin-10(4<i>H</i>)-yl]-</p>	466.5	3.79



Example	Structure	Observed MS	HPLC retention time
367	 <p><i>Ethanediamide, N'-[2-[[[2-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-</i></p>	470.7	3.46
368	 <p><i>N'-(2-((2,4-Dichlorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethyl-ethanediamide</i></p>	520.5	4.42



Example	Structure	Observed MS	HPLC retention time
369	 <p><i>N'</i>-(3-Hydroxy-2-((2-methoxybenzyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	482.6	3.48
370	 <p><i>N'</i>-(3-Hydroxy-2-((2-methylbenzyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	466.7	3.74



Example	Structure	Observed MS	HPLC retention time
371	<p data-bbox="477 720 949 934"><i>N'</i>-(2-((3-Fluorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	470.6	3.54
372	<p data-bbox="477 1260 949 1473"><i>N'</i>-(2-((3,4-Dichlorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	522.3	4.46



Example	Structure	Observed MS	HPLC retention time
373	 <p><i>N'</i>-(3-Hydroxy-2-((3-methoxybenzyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	482.6	3.42
374	 <p><i>N'</i>-(2-((4-Chlorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	486.6	3.88



Example	Structure	Observed MS	HPLC retention time
375	 <p><i>N'</i>-(3-Hydroxy-2-((4-methoxybenzyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	482.7	3.4
376	 <p><i>N'</i>-(3-Hydroxy-2-((4-methylbenzyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	466.6	3.77



Example	Structure	Observed MS	HPLC retention time
377	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-(trifluoromethyl)benzyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	520.6	4.01
378	 <p><i>N'</i>-(2-((4-(dimethylamino)benzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	495.7	3.58



Example	Structure	Observed MS	HPLC retention time
379	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((4-sulfamoylbenzyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	531.6	2.4
380	 <p><i>N'</i>-(2-((3-Chlorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	486.6	3.97

Example	Structure	Observed MS	HPLC retention time
381	 <p><i>N'</i>-(2-((2,3-dimethoxybenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	512.7	3.3
382	 <p><i>N'</i>-(2-((2,4-dimethoxybenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	512.7	3.61

Example	Structure	Observed MS	HPLC retention time
383	 <p><i>N'</i>-(2-((3,5-Dichlorobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	520.6	4.43
384	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((3-(trifluoromethoxybenzyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	536.6	4.28

Example	Structure	Observed MS	HPLC retention time
385	 <p><i>N'</i>-(2-((4-Aminobenzyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	467.6	2.49
386	 <p><i>N'</i>-(3-Hydroxy-4-oxo-2-((2-trifluoromethoxy)benzyl)carbamoyl)-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	536.7	4.17

Example	Structure	Observed MS	HPLC retention time
387	 <p><i>N'</i>-(2-((2-Biphenylylmethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	528.6	4.55
388	 <p><i>N'</i>-(2-((3-Biphenylylmethyl)carbamoyl)-3-hydroxy-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	530.5	4.59



Example	Structure	Observed MS	HPLC retention time
389	 <p><i>N'</i>-(3-Hydroxy-2-(((5-methyl-2-pyrazinyl)methyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	468.7	2.15
390	 <p><i>N'</i>-(3-Hydroxy-2-((4-hydroxybenzyl)carbamoyl)-4-oxo-6,7,8,9-tetrahydro-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl)-N,N-dimethylethanediamide</p>	468.6	2.59



The compounds in Table 3 were synthesized using methods similar to those described previously. After synthesis the compounds were analyzed reverse phase LCMS on a Shimadzu HPLC system coupled to a mass spectrometer. LC Conditions;

50 x 4.6 mm, 5 micron SunfireC18 column; mobile phase A 10 mM NH₄OAc in 5:95 acetonitrile/water, mobile phase B = 10 mM NH₄OAc in 95:5 acetonitrile/water; 4 mL/min flow rate; gradient = 0 to 100% mobile phase B over 4 min.

5 Table 3

Example	Structure	Observed MS ion	HPLC retention time
391	<p><i>N</i>-(4-Fluorobenzyl)-3-hydroxy-10-((3-methylphenyl)sulfonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide</p>	525.7	4.96
392	<p>10-(((3,5-Dimethyl-4-isoxazolyl)sulfonyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide</p>	530.5	3.98



Example	Structure	Observed MS ion	HPLC retention time
393	 <p><i>N</i>-(4-Fluorobenzyl)-3-hydroxy-10-(((5-methyl-4-isoxazolyl)sulfonyl)amino)-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide</p>	518.3	3.22
394	 <p>10-(((2-(Acetylamino)-4-methyl-1,3-thiazol-5-yl)sulfonyl)amino)-N-(4-fluorobenzyl)-3-hydroxy-4-oxo-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-a]azepine-2-carboxamide</p>	591.7	3.12



Example	Structure	Observed MS ion	HPLC retention time
395	 <p><i>N</i>-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((phenylsulfonyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepine-2-carboxamide</p>	513.6	4.61
396	 <p><i>N</i>-(4-Fluorobenzyl)-3-hydroxy-4-oxo-10-((2-(trifluoromethoxy)phenyl)sulfonyl)amino)-4,6,7,8,9,10-hexahydro-7,10-ethanopyrimido[1,2-<i>a</i>]azepine-2-carboxamide</p>	597.4	4.54



The compounds in Table 4 were synthesized using methods similar to those described previously. After synthesis the compounds were analyzed reverse phase LCMS on a Shimadzu HPLC system coupled to a mass spectrometer. LC Conditions;



- 5 50 x 4.6 mm, 5 micron Phenomenex Luna C18 column; mobile phase A 10 mM NH₄OAc in 5:95 acetonitrile/water, mobile phase B = 10 mM NH₄OAc in 95:5 acetonitrile/water; 4 mL/min flow rate; gradient = 0 to 100% mobile phase B over 4 min.



Table 4



Example	Structure	Observed MS	HPLC retention time
397	 <p>1-Pyrrolidinecarboxylic acid, 2-[{2-[{[(4-fluorophenyl)methyl]amino}carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl}amino}carbonyl]-, ethyl ester</p>	542.26	6.62
398	 <p>1-Pyrrolidinecarboxylic acid, 2-[{2-[{[(4-fluorophenyl)methyl]amino}carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl}amino}carbonyl]-, propyl ester</p>	556.29	6.95



Example	Structure	Observed MS	HPLC retention time
399	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, <i>N</i>-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[[1-(2-methyl-1-oxopropyl)-2-pyrrolidinyl]carbonyl]amino]-4-oxo-</p>	540.3	6.69
400	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[1-(cyclobutylcarbonyl)-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-</p>	552.3	6.83

Example	Structure	Observed MS	HPLC retention time
401	 <p>1,2-Pyrrolidinedicarboxamide, <i>N</i>~2~-{2-[{[(4-fluorophenyl)methyl]amino}carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4<i>H</i>)-yl}-<i>N</i>~1~-methyl-</p>	527.25	5.97
402	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, <i>N</i>-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-10-[[1-(1-methylethyl)sulfonyl]-2-pyrrolidinyl]carbonyl]amino]-4-oxo-</p>	576.25	6.42

Example	Structure	Observed MS	HPLC retention time
403	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[1-(ethylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-N-[{(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-</p>	562.27	6.14
404	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[{(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[[1-(propylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-</p>	576.28	6.49


Example	Structure	Observed MS	HPLC retention time
405	 <p><i>1-Pyrrolidinecarboxylic acid, 2-[{2-[{[(4-fluorophenyl)methyl]amino}carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl}amino}carbonyl]-, 1-methylethyl ester</i></p>	556.3	6.92
406	 <p><i>1-Pyrrolidinecarboxylic acid, 2-[{2-[{[(4-fluorophenyl)methyl]amino}carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl}amino}carbonyl]-, methyl ester</i></p>	528.27	6.21

Example	Structure	Observed MS	HPLC retention time
407	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[1-(cyclopropylcarbonyl)-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-</p>	538.26	6.5
408	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, 10-[[[1-(cyclopentylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-</p>	602.31	6.82

Example	Structure	Observed MS	HPLC retention time
409	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-10-[[[1-(1H-1,2,4-triazol-3-ylsulfonyl)-2-pyrrolidinyl]carbonyl]amino]-</p>	601.23	5.82
410	 <p>7,10-Ethanopyrimido[1,2-a]azepine-2-carboxamide, N-[(4-fluorophenyl)methyl]-10-[[[1-(2-furanyl sulfonyl)-2-pyrrolidinyl]carbonyl]amino]-4,6,7,8,9,10-hexahydro-3-hydroxy-4-oxo-</p>	600.24	6.54

ISOLATION OF SINGLE DIASTEREOMERS

Scheme XXXV




5

The isolation of the single diastereomers of certain compounds of this invention is illustrated in Scheme XXXV.

METHOD A

10

Separation N^t-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl into its individual enantiomers,




15

Ethanediamide, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-4-oxo-3-(phenylmethoxy)-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-. A solution of ethanediamide, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl (0.160 g, 0.350 mmol) in dry DMF (5 mL) was treated with benzyl bromide (0.050 mL, 0.420 mmol) and potassium carbonate (0.276 g, 2 mmol) and stirred with heating (60°C oil bath) overnight. The solvent was removed under reduced pressure and the residue was dissolved in EtOAc (20 mL) and washed with 1.0 N HCl (2 x 20 mL). The organic layer was dried (MgSO_4), filtered, and concentrated under reduced pressure to give the crude product as a light brown oil, which partially solidified upon standing. The solid was triturated with Et_2O several times then loaded onto a Biotage 25+M samplet and purified with Biotage flash chromatography system, eluting with a solvent gradient of 40% to 100% hexanes in EtOAc. The individual diastereomers of this compound were separated by chiral SFC chromatography (5 micron 30 x 250 mm Chiralcel OD-H column eluting with 70% CO_2 30% MeOH and 0.1% diethylamine at 35 °C and 150 Bar). Two separately eluting compounds were isolated, a faster eluting compound (isomer A) and a slower eluting compound (isomer B).

20

Example 411




25 *Ethanediamide, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-.* Removal of the benzyl group (H_2 , Pd/C) from the faster eluting isomer (isomer A) provided the title compound white crystalline solid (54% yield). ^1H NMR (500 MHz, MeOD) δ ppm 1.65 - 1.74 (m, 1 H) 1.90 - 1.99 (m, 1 H) 2.02 - 2.12 (m, 1

H) 2.21 - 2.32 (m, 1 H) 2.62 (td, $J=12.51, 5.49$ Hz, 1 H) 2.75 - 2.82 (m, 1 H) 2.91 - 2.98 (m, 1 H) 2.99 - 3.03 (m, 3 H) 3.11 (s, 3 H) 3.78 - 3.95 (m, 2 H) 4.59 (s, 1 H) 4.89 (br. s., 4 H) 7.04 (t, $J=8.70$ Hz, 2 H) 7.39 (dd, $J=8.55, 5.49$ Hz, 2 H). LC/MS ($M+H$) = 458.16. HPLC purity >97%.

5

Example 412




10 *Ethanediamide, N'-[2-[[[4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-methanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-*. Removal of the benzyl group (H_2 , Pd/C) from the slower eluting isomer (isomer B) provided the title compound as an off-white crystalline solid (52% yield). LC/MS ($M+H$) = 458.08. HPLC purity >97%.

15

METHOD B

Separation of Ethanediamide, N'-[2-[[[4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-20 11(6H)-yl]-N,N-dimethyl. into its individual enantiomers




Ethanediamide, N'-[3-[(4-chlorophenyl)methoxy]-2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-*. A solution of ethanediamide, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl (0.367 g, 0.779 mmol) in dry DMF (10 mL) was treated with cesium carbonate (0.512 g, 1.571 mmol) and 4-chlorobenzyl bromide (0.218 g, 1.040 mmol), and the reaction stirred with heating (60°C oil bath) for 1 hr. Additional 4-chlorobenzyl bromide (0.219 g, 1.040 mmol) and cesium carbonate (0.501 g, 1.55 mmol) were added and the reaction was stirred for and 45 min, then cooled and concentrated under reduced pressure.*

The crude residue, was dissolved in CH₂Cl₂, and washed with 1.0N HCl, then dried (MgSO₄), filtered, and concentrated under reduced pressure to give a yellow oil. The oil crystallized upon standing for two days at room temperature. Trituration with ether (2 times), followed by crystallization from 95% EtOH-H₂O yielded the title compound (0.242 g, 0.406 mmol, 52.1 % yield), as a white crystalline solid. ¹H NMR (500 MHz, CDCl₃) δ ppm 8.87 (1 H, s), 8.22 (1 H, t, *J*=5.95 Hz), 7.50 (2 H, d, *J*=8.55 Hz), 7.27 - 7.35 (4 H, m), 6.98 (2 H, t, *J*=8.70 Hz), 5.44 (1 H, dt, *J*=15.49, 3.70 Hz), 5.21 (2 H, s), 4.48 - 4.59 (2 H, m), 3.48 - 3.56 (1 H, m), 3.31 (3 H, s), 2.89 (3 H, s), 2.57 - 2.70 (3 H, m), 2.36 - 2.48 (2 H, m), 1.80 - 1.93 (2 H, m), 1.76 (1 H, m), 1.61 (1 H, m). The individual diastereomers of this compound were separated on by chiral SFC chromatography (5 micron 30 x 250 mm Chiraldak AS-H column eluting with 85% CO₂ 15% MeOH and 0.1% diethylamine at 35 °C and 150 Bar). Two separately eluting compounds were isolated., the faster eluting compound (isomer A, Intermediate 159) and a slower eluting compound (isomer B, Intermediate 160).

25


Intermediate 159



Isomer A. Earlier eluting peak. 0.08 g (34% yield) was isolated as a glassy solid.

Intermediate 159


5



Isomer B. The later eluting peak. 0.08 g (34% yield) was isolated as a crystalline solid from 95% EtOH. This material was analyzed by chiral SFC chromatography, retention time = 12.97 min; Chiraldak AS-H column, 4.6 x 250mm, 5 microns solvent: 85% CO₂, 15% MeOH with 0.1%diethylamine, Temp: 35 °C, Pressure: 150bar, flow rate at 2.0 ml/min, UV at 210 nm.

Example 413


15



Ethanediame, N'-[2-[[[4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-. Hydrogenolysis (H₂, Pd/C) of Intermediate 158 provided the title compound as a single diastereomer. Off-white crystalline solid (86% yield). ¹H NMR (500 MHz, CDCl₃) δ ppm 12.08 (1 H, s), 8.71 (1 H, t, *J*=6.10 Hz), 8.27 (1 H, s), 7.36 (2 H, dd, *J*=8.55, 5.49 Hz), 6.98 - 7.04 (2 H, m), 5.50 (1 H, dt, *J*=15.56, 3.81 Hz), 4.49 - 4.59 (2 H, m), 3.49 (1 H, t, *J*=13.58 Hz), 3.29 (3 H, s), 2.91 (3 H, s), 2.64

(1 H, q, $J=5.70$ Hz), 2.52 - 2.61 (1 H, m), 2.43 - 2.52 (2 H, m), 2.27 - 2.38 (1 H, m), 1.78 - 1.93 (3 H, m), 1.57 - 1.65 (1 H, m). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.20, 163.58, 163.33, 161.55, 161.38, 159.49, 152.11, 146.86, 133.72, 133.69, 129.96, 129.89, 123.72, 115.57, 115.40, 66.61, 44.13, 42.52, 40.05, 38.17, 37.26, 5 36.51, 35.09, 32.57, 30.25. LC/MS ($\text{M}+\text{H}$) = 472.12. HPLC purity >97%.

Example 414



10


Ethanediamide, N'-(2-[[[4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-l. Hydrogenolysis (H_2 , Pd/C) of Intermediate 159 provided the title compound as a single diastereomer White crystalline solid (96% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.08 (1 H, s), 8.70 (1 H, t, $J=5.95$ Hz), 8.29 (1 H, s), 7.33 - 7.39 (2 H, m), 6.97 - 7.04 (2 H, m), 5.50 (1 H, dt, $J=15.56, 3.81$ Hz), 4.49 - 4.60 (2 H, m), 3.44 - 3.53 (1 H, m), 3.29 (3 H, s), 2.91 (3 H, s), 2.64 (1 H, q, $J=5.49$ Hz), 2.53 - 2.61 (1 H, m), 2.42 - 2.52 (2 H, m), 2.33 (1 H, td, $J=12.97, 7.02$ Hz), 1.77 - 1.93 (3 H, m), 1.57 - 1.66 (1 H, m). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.20, 163.58, 163.33, 161.55, 161.38, 159.49, 152.11, 146.86, 133.72, 133.69, 129.96, 129.89, 123.72, 115.56, 115.40, 66.61, 44.13, 42.52, 40.05, 38.16, 37.26, 36.51, 20 35.09, 32.57, 30.25. LCMS ($\text{M}+\text{H}$) = 472.35.

METHOD C

25

Separation of Ethanediamide, N'-($8\text{S},11\text{R}$)-2-[[[4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-into its individual enantiomers


Intermediate 160



5 Isomer A was isolated as the faster eluting compound from a racemic mixture
of ethanediamide, N'-[2-[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-
7,8,9,10-tetrahydro-4-oxo-3-(phenylmethoxy)-8,11-methano-4H-pyrimido[1,2-
a]azocin-11(6H)-yl]-N,N-dimethyl- by chiral SCF chromatography. LCMS (M+H) =
576.42 min. α_D = +16.60 (5.02 mg/mL in CHCl₃). Chiral SCG analysis; Chiralcel
10 OD-H column, 4.6 x 250mm, 5microns; solvent: 75% CO₂, 25% MeOH with 0.1%
diethylamine; Temp: 35 °C, Pressure: 150 bar ; Flow rate at 1.0 ml/min, UV at 300
nm.

Intermediate 170

15




Isomer B was isolated from a as the slower moving compound from a racemic
mixture of ethanediamide, N'-[2-[[(4-fluoro-3-
20 methylphenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-4-oxo-3-
(phenylmethoxy)-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-

dimethyl- by chiral SCF chromatography. LCMS (M+H) = 576.21 min. $\alpha_D = -16.60$ (5.02 mg/mL in CHCl₃). Chiral SCG analysis; Chiralcel OD-H column, 4.6 x 250mm, 5microns; solvent: 75% CO₂, 25% MeOH with 0.1% diethylamine; Temp: 35 °C, Pressure: 150 bar; Flow rate at 1.0 ml/min, UV at 300 nm.


5

Example 415



10 *Ethanediamide, N'-(8S,11R)-2-[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-.*: A solution of Intermediate 160 (35 mg, 0.061 mmol) in ethyl acetate (2 mL) and methanol (2 mL) was treated with 10% palladium on carbon (65 mg) and stirred under hydrogen for 16 hrs. The reaction was filtered and the filtrate concentrated. The residue was dissolved in 95% EtOH (2mL) and the solution was allowed to stand for 16 hrs. Solids were collected by vacuum filtration to afford the title compound as an off-white crystalline solid (24 mg, 0.049 mmol, 81 % yield). ¹H NMR (500 MHz, CDCl₃) δ ppm 12.14 (1 H, s), 8.72 (1 H, t, *J*=6.10 Hz), 8.32 (1 H, s), 7.17 - 7.25 (2 H, m), 6.92 - 7.01 (1 H, m), 5.54 (1 H, ddd, *J*=15.41, 3.81, 3.66 Hz), 4.54 (2 H, d, *J*=6.41 Hz), 3.52 (1 H, t, *J*=13.73 Hz), 3.32 (3 H, s), 2.94 (3 H, s), 2.67 (1 H, q, *J*=5.49 Hz), 2.56 - 2.64 (1 H, m), 2.46 - 2.55 (2 H, m), 2.30 - 2.42 (1 H, m), 2.28 (3 H, d, *J*=1.83 Hz), 1.81 - 1.96 (3 H, m), 1.60 - 1.68 (1 H, m). ¹³C NMR (126 MHz, CDCl₃) δ ppm 168.15, 163.62, 161.82, 161.70, 159.88, 159.56, 152.04, 146.83, 133.38, 133.35, 131.35, 131.31, 127.13, 127.06, 125.07, 124.94, 123.79, 115.19, 115.00, 109.69, 66.63, 44.20, 42.51, 40.04, 38.19, 37.22, 36.44, 35.14, 32.55, 30.27, 14.66, 14.63. LCMS (M+H) = 486.15. HPLC purity: >97%.

Example 416



5 *Ethanediamide, N'-(8R,11S)-2-[[[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-*. White crystalline solid (71% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 12.11 (1 H, brs), 8.70 (1 H, t, $J=6.10$ Hz), 8.27 (1 H, s), 7.12 - 7.21 (2 H, m), 6.90 - 6.98 (1 H, m), 5.50 (1 H, dt, $J=15.49, 3.70$ Hz), 4.50 (2 H, d, $J=6.41$ Hz), 3.48 (1 H, t, $J=13.58$ Hz), 3.29 (3 H, s), 2.91 (3 H, s), 2.64 (1 H, q, $J=5.49$ Hz), 2.52 - 2.61 (1 H, m), 2.43 - 2.53 (2 H, m), 2.27 - 2.39 (1 H, m), 2.25 (3 H, d, $J=1.83$ Hz), 1.78 - 1.93 (3 H, m), 1.57 - 1.65 (1 H, m). ^{13}C NMR (126 MHz, CDCl_3) δ ppm 168.15, 163.65, 161.81, 161.75, 159.87, 159.56, 152.03, 146.82, 133.39, 133.36, 131.35, 131.31, 127.12, 127.06, 125.07, 124.94, 123.81, 115.19, 115.00, 66.64, 44.22, 42.50, 40.03, 38.19, 37.21, 36.42, 35.16, 32.54, 30.26, 14.67, 14.64. LC/MS ($\text{M}+\text{H}$) = 486.12. HPLC purity >97%.

METHOD D


20 Separation of 8,11-methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo- into its individual diastereomers.

A solution of 8,11-methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo- (0.196 g, 0.38 mmol) in dry DMF (5 mL) was treated with 4-chlorobenzyl bromide (0.094 g, 0.46 mmol) and potassium carbonate (0.079 g, 0.57 mmol), and the reaction was heated (60°C) for 90 min. Solvent was removed under reduced pressure, and the residue, dissolved

in CH_2Cl_2 , was washed with 1.0 N HCl. The organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure to give an oily solid which was triturated with ether (2x). The residue was subjected to chiral SCF chromatography (Chiralcel OD-H column, 4.6 x 250mm, 5 microns, solvents: 75% CO_2 , 25% methanol with 0.1%diethylamine, Temp: 35 °C, Pressure: 150 bar, Flow rate at 2.0 ml/min UV at 300nm) for separation to the component diastereomers.


Intermediate 171

10



8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 3-[(4-chlorophenyl)methoxy]-N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-4-oxo-, (8S,11R)- First eluting compound isolated from the chiral SCF chromatographic purification, was recrystallized from EtOH and water, resulting in the title compound (0.047 g, 0.073 mmol, 38 % yield) as a crystalline white solid. ^1H NMR (500 MHz, CDCl_3) δ ppm 9.71 - 9.90 (1 H, m), 7.48 (2 H, dd, J =8.24, 2.75 Hz), 7.32 - 7.39 (2 H, m), 7.29 (2 H, d, J =8.55 Hz), 6.96 - 7.02 (2 H, m), 5.46 (1 H, dt, J =15.26, 3.66 Hz), 5.16 - 5.37 (3 H, m), 4.38 - 4.63 (3 H, m), 3.74 - 4.02 (2 H, m), 3.42 - 3.63 (2 H, m), 2.78 - 2.87 (1 H, m), 2.65 - 2.74 (2 H, m), 2.49 (1 H, brs), 2.21 - 2.39 (2 H, m), 1.80 - 2.01 (3 H, m), 1.60 - 1.68 (2 H, m). LC/MS ($\text{M}+\text{H}$) = 640.09. HPLC purity >98%.

Intermediate 172



5 *8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 3-[(4-chlorophenyl)methoxy]-N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-4-oxo-, (8R,11S)-*


Second eluting compound isolated from the chiral SCF chromatographic purification.

White crystalline solid (37% yield). ^1H NMR (500 MHz, CDCl_3) δ ppm 1.57 - 1.61 (m, 3 H), 1.62 - 1.70 (m, 2 H), 1.78 - 2.12 (m, 3 H), 2.20 - 2.41 (m, 2 H), 2.41 - 2.55 (m, 1 H), 2.64 - 2.78 (m, 2 H), 2.81 (dd, $J=13.12, 6.71$ Hz, 1 H), 3.46 - 3.69 (m, 2 H), 3.78 - 4.02 (m, 1 H), 4.29 - 4.42 (m, 1 H), 4.47 - 4.66 (m, 2 H), 5.17 - 5.36 (m, 3 H), 5.42 - 5.51 (m, 1 H), 6.99 (t, $J=8.70$ Hz, 2 H), 7.29 (d, $J=8.24$ Hz, 2 H), 7.32 - 7.39 (m, 2 H), 7.49 (dd, $J=7.17, 5.04$ Hz, 2 H), 7.92 - 8.05 (m, 1 H), 9.55 - 9.82 (m, 1 H).

10 LC/MS ($\text{M}+\text{H}$) = 640.11. HPLC purity >97%.

15


Example 417



8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (8R,11S)-. The 4-chloro-benzyl group of 8,11-methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 3-[(4-chlorophenyl)methoxy]-N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-4-oxo-, (8S,11R)- was removed under hydrogenolytic conditions (H_2 , Pd/C) to yield the title compound as a white crystalline solid. 1H NMR (500 MHz, $CDCl_3$) δ ppm 11.98 - 12.03 (1 H, m), 9.54 - 9.71 (1 H, m), 8.14 - 8.23 (1 H, m), 7.39 (2 H, dd, $J=8.24, 5.49$ Hz), 7.02 (2 H, t, $J=8.70$ Hz), 5.51 (1 H, ddd, $J=15.41, 3.66, 3.51$ Hz), 5.17 - 5.38 (1 H, m), 4.52 - 4.66 (2 H, m), 4.30 - 4.41 (1 H, m), 3.89 - 4.03 (1 H, m), 3.45 - 3.87 (3 H, m), 2.63 - 2.78 (3 H, m), 1.79 - 2.50 (6 H, m), 1.60 - 1.66 (2 H, m). ^{13}C NMR (126 MHz, $CDCl_3$) δ ppm 167.91, 167.87, 160.43, 160.40, 159.45, 159.12, 158.75, 152.59, 152.57, 146.96, 146.94, 133.30, 130.11, 130.06, 130.00, 123.26, 123.20, 115.76, 115.71, 115.58, 115.54, 94.04, 92.63, 90.93, 89.53, 66.00, 65.91, 55.20, 55.02, 54.55, 54.35, 46.23, 45.62, 43.07, 42.91, 42.75, 42.67, 40.33, 40.27, 37.61, 37.47, 33.92, 33.88, 33.61, 33.45, 32.55, 30.37, 30.35, 30.23, 30.05. LC/MS (M+H) = 515.91. HPLC purity >97%.

20

Example 418



25 8,11-Methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (8S,11R)-. A solution of 8,11-methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, 3-[(4-chlorophenyl)methoxy]-N-[(4-

fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-4-oxo-, (8R,11S)- (0.045 g, 0.070 mmol) in methanol (5 mL) was treated with palladium on carbon (10 mg, 9.40 μ mol), and hydrogen, with stirring for 16 hrs. The reaction was filtered, then concentrated under reduced pressure. The film residue recrystallized from EtOH and water over several days. The crystals were isolated and vacuum dried, resulting in the title compound (10.1 mg, 0.0196 mmol, 28% yield) as a purple crystalline solid (28% yield). The crystal structure of 8,11-methano-4H-pyrimido[1,2-a]azocine-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (8S,11R)- was determined by crystallography to elucidate the absolute configuration of the two chiral centers at the bicyclic ring. Crystal system: monoclinic; space group: C2; a = 18.1035(3) Å (α = 90), b = 27.1111(4) Å (β = 98.947(1) $^\circ$), c = 10.0563(5) Å (γ = 90 $^\circ$); V = 4875.6(3) Å³; no. of molecules/cell: Z = 8, calculated crystal density: d_x = 1.405 g cm⁻³; absorption coefficient: μ = 0.110 mm⁻¹; θ range for lattice parameters ($^\circ$): 2.53 to 27.30. The prismatic crystal from which the X-ray data was collected is chiral, and there are two molecules per asymmetric unit. Based on the known **R** configuration of C(1), the absolute configuration at C(7) and C(11) are established to be **R** and **S**, respectively. Intramolecular H-bonding was observed between the hydroxyl O(4) and the amide oxygen O(5) [O4···O5 = 2.590/2.570 Å, H···O5 = 1.889/1.862 Å, O4-H···O5 = 143/144 $^\circ$]. ¹H NMR (500 MHz, CDCl₃) δ ppm 11.98 - 12.05 (1 H, m), 9.66 - 9.90 (1 H, m), 8.10 - 8.23 (1 H, m), 7.36 - 7.44 (2 H, m), 6.99 - 7.06 (2 H, m), 5.47 - 5.56 (1 H, m), 5.16 - 5.40 (1 H, m), 4.52 - 4.65 (2 H, m), 4.39 - 4.51 (1 H, m), 3.71 - 4.04 (2 H, m), 3.39 - 3.61 (2 H, m), 2.80 (1 H, td, J =12.89, 6.56 Hz), 2.62 - 2.73 (2 H, m), 2.40 - 2.51 (1 H, m), 2.21 - 2.39 (2 H, m), 1.95 - 2.14 (1 H, m), 1.79 - 1.93 (3 H, m), 1.63 (1 H, brs). ¹³C NMR (126 MHz, CDCl₃) δ ppm 167.87, 167.83, 163.43, 163.40, 161.27, 160.45, 160.17, 159.46, 158.73, 158.43, 152.61, 146.98, 146.95, 133.26, 133.24, 133.18, 133.15, 130.16, 130.12, 130.10, 130.06, 123.20, 123.14, 115.76, 115.73, 115.58, 115.56, 94.11, 92.71, 90.95, 89.55, 65.91, 65.82, 55.22, 55.03, 54.53, 54.34, 46.08, 45.78, 42.79, 42.74, 42.66, 42.48, 40.38, 40.32, 37.80, 37.64, 34.05, 34.00, 33.58, 33.41, 32.55, 30.33, 30.29, 30.23, 30.05. HPLC purity >97%.


It will be evident to one skilled in the art that the present disclosure is not limited to the foregoing illustrative examples, and that it can be embodied in other specific forms without departing from the essential attributes thereof. It is therefore desired that the examples be considered in all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing examples, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

5

CLAIMS

We claim:

5 1. A compound of Formula I



I

10 wherein:

R¹ is (Ar¹)alkyl, alkyl, (cycloalkyl)alkyl, diphenylalkyl, phenoxyalkyl, (PhNH)alkyl, (methylpyrrolidinyl)alkyl, (imidazolyl)alkyl (valerolactamyl)alkyl, (tetrahydrofuranyl)alkyl, ((fluoro)(methyl)pyridinyl)methyl, phenylcyclopropyl, or 15 benzylpyrrolidinyl;

R² is hydrogen, alkyl, hydroxy or alkoxy;

R³ is hydrogen, halo, hydroxy, alkoxy, cyano, alkyl, cycloalkyl, haloalkyl, 20 hydroxyalkyl, alkoxyalkyl, alkoxy, haloalkoxy, N(R⁸)(R⁹), N(R⁶)COR⁷, N(R⁶)SO₂R⁷, N(R⁶)CO₂R⁷, N(R⁶)SO₂N(R⁸)(R⁹), CO₂R⁶, CON(R⁸)(R⁹), SOR⁷, SO₂R⁷, SO₂N(R⁸)(R⁹), PO(OR⁶)₂, R¹², or Ar²;

R⁴ is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or 25 N(R⁶)(R⁶);

R⁵ is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or N(R⁶)(R⁶);

30 R⁶ is hydrogen, alkyl, haloalkyl, or cycloalkyl;

R⁷ is alkyl, haloalkyl, or cycloalkyl;

R⁸ is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl;

5 R⁹ is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl; or

N(R⁸)(R⁹) taken together is azetidinyl, pyrrolidinyl, (R¹⁰)-piperidinyl,


N-(R¹¹)-piperazinyl, morpholinyl, thiomorpholinyl, or dioxothiazinyl;

10 R¹⁰ is hydrogen, alkyl, hydroxy, or hydroxyalkyl;

R¹¹ is hydrogen, alkyl, cyclolkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, phenyl, pyridinyl, (methylimidazolyl)methyl, COR⁶, CO₂R⁶, (hydroxyalkyl)CO, (alkoxyalkyl)CO, (tetrahydrofuranyl)CO, (methylisoxazolyl)CO, (thienyl)CO,

15 (furanyl)CO, (pyridinyl)CO, CON(R⁸)(R⁹), SO₂R⁷, SO₂N(R⁸)(R⁹),

(dimethylisoxazolyl)SO₂, ((carboethoxy)thienyl)SO₂, (methylimidazolyl)SO₂,



R¹² is azetidinonyl, pyrrolidinonyl, valerolactamyl, caprolactamyl, maleimido,

20 oxazolidinonyl, imidazolidinonyl, triazolonyl, N-(R⁶)-dioxothiazolidinyl or dioxothiazinyl, and is substituted with 0-2 substituents selected from the group consisting of alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, and aminoalkyl;

R¹³ is hydrogen, alkyl, alkylcarbonyl, alkoxycarbonyl, benzyloxycarbonyl, or

25 alkylsulfonyl;

R¹⁴ is N(R¹⁵)(R¹⁵), N(R¹⁵)(benzyloxycarbonyl), N(R¹⁵)(alkyloxycarbonyl),

N(R¹⁵)((hydroxyalkyl)oxycarbonyl), N(R¹⁵)((alkyloxy)alkyloxycarbonyl),


N(R¹⁵)CO(R¹⁶), N(R¹⁵)((CO(N(R¹⁵)₂)alkyl), N(R¹⁵)((CO₂CO(N(R¹⁵)₂)alkyl),


30 N(R¹⁵)CO((N(R¹⁵)₂)alkyl), N(R¹⁵)CO((CON(R¹⁵)(R¹⁵)alkyl), N(R¹⁵)COCO₂(R¹⁵),

$\text{N}(\text{R}^{15})\text{CO}((\text{CO}_2(\text{R}^{15})\text{alkyl}), \text{N}(\text{R}^{15})\text{COCON}(\text{R}^{15})(\text{R}^{15}),$
 $\text{N}(\text{R}^{15})\text{COCO}((\text{N}(\text{R}^{15})(\text{R}^{15}))\text{alkyl}), \text{N}(\text{R}^{15})\text{CO}(\text{N}(\text{R}^{15})_2), \text{N}(\text{R}^{15})\text{SO}_2\text{R}^{16},$
 $\text{N}(\text{R}^{15})\text{SO}_2(\text{N}(\text{R}^{15})_2), \text{or } \text{N}(\text{R}^{15})\text{COAr}^3;$

5 or R^{14} is hydroxy, alkoxy, (hydroxy)alkoxy, (alkoxy)alkoxy, $(\text{R}^{16}\text{CO}_2)\text{alkoxy}$, or
 $(\text{PhCO}_2)\text{alkoxy};$

or R^{14} is $\text{N}(\text{R}^{15})\text{COCON}(\text{R}^{15})(\text{alkoxyalkyl}), \text{N}(\text{R}^{15})\text{COCON}(\text{R}^{15})(\text{hydroxyalkyl}),$
 $\text{N}(\text{R}^{15})\text{COCON}(\text{R}^{15})((\text{tetrahydrofuran})\text{methyl}), \text{N}(\text{R}^{15})\text{COCON}(\text{R}^{15})((\text{N,N-})$
10 $\text{dimethylpyrazolyl})\text{methyl}), \text{or } \text{N}(\text{R}^{15})\text{CO}((\text{Ar}^4)\text{alkyl});$





R^{15} is hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, haloalkyl, or cycloalkyl;


5

or $N(R^{15})(R^{15})$ taken together is azetidinyl, pyrrolidinyl, (R^{10})-piperidinyl, $N-(R^{11})$ -piperazinyl, morpholinyl, or thiomorpholinyl, and is substituted with 0-2 substituents selected from halo, hydroxy, alkoxy, alkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkyloxyalkyl, acetamido, CO_2R^6 , and $CON(R^8)(R^9)$;

10

R^{16} is alkyl, haloalkyl, cycloalkyl, (cycloalkyl)alkyl, hydroxyalkyl, alkoxyalkyl, tetrahydropyranyl, or Ar^3 ;

R^{17} is hydrogen, alkyl, alkylCO, cycloalkylCO, alkyloxyCO, $CON(R^{15})(R^{15})$,
15 $COCON(R^{15})(R^{15})$, $COAr_3$, alkylSO₂, cycloalkylSO₂, furanylSO₂, triazolylSO₂, or N -methylpyrrolylSO₂;




Ar² is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyrimidinyl, pyrazinyl, pyridinyl, or hydroxypyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, cyano, benzyl, alkyl, alkoxy, N(R⁸)(R⁹), CO₂R⁶, and CON(R⁸)(R⁹);

Ar³ is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, imidazopyridinyl, quinolinyl, isoquinolinyl, or phenyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, alkoxy, haloalkoxy, cyano, benzyl, alkyl, haloalkyl, N(R⁸)(R⁹), and N(R¹⁵)CO(R¹⁶);

Ar⁴ is pyrazolyl or triazolyl, and is substituted with 0-1 substituents selected from CO₂R⁶ and CON(R⁸)(R⁹); and

15

X-Y-Z is



20

or a pharmaceutically acceptable salt thereof.

2. A compound of claim 1 where

25

R¹ is (Ar¹)alkyl;

R² is hydrogen, alkyl, hydroxy or alkoxy;

5 R³ is hydrogen, halo, hydroxy, cyano, alkyl, haloalkyl, alkoxy, haloalkoxy, N(R⁸)(R⁹), N(R⁶)COR⁷, N(R⁶)SO₂R⁷, N(R⁶)CO₂R⁷, N(R⁶)SO₂N(R⁸)(R⁹), CO₂R⁶, CON(R⁸)(R⁹), SOR⁷, SO₂R⁷, SO₂N(R⁸)(R⁹), PO(OR⁶)₂, R¹², or Ar²;

10 R⁴ is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or N(R⁶)(R⁶);

R⁵ is hydrogen, halo, hydroxy, cyano, alkyl, alkoxy, haloalkyl, haloalkoxy, or N(R⁶)(R⁶);

15 R⁶ is hydrogen, alkyl, haloalkyl, or cycloalkyl;

R⁷ is alkyl, haloalkyl, or cycloalkyl;

R⁸ is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl;

20 R⁹ is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or dialkylaminoalkyl; or N(R⁸)(R⁹) taken together is azetidinyl, pyrrolidinyl, (R¹⁰)-piperidinyl, N-(R¹¹)-piperazinyl, morpholinyl, thiomorpholinyl, or dioxothiazinyl;

25 R¹⁰ is hydrogen, alkyl, hydroxy, or hydroxyalkyl;


R¹¹ is hydrogen, alkyl, cyclolkyl, COR⁶, or CO₂R⁶;

30 R¹² is azetidinonyl, pyrrolidinonyl, valerolactamyl, caprolactamyl, maleimido, oxazolidinonyl, imidazolidinonyl, triazolonyl, N-(R⁶)-dioxothiazolidinyl or dioxothiazinyl, and is substituted with 0-2 substituents selected from the group consisting of alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, and aminoalkyl;

R^{13} is hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, benzyloxycarbonyl, or alkylsulfonyl;

5 R^{14} is $N(R^{15})(R^{15})$, $N(R^{15})(\text{benzyloxycarbonyl})$, $N(R^{15})(\text{alkyloxycarbonyl})$,
 $N(R^{15})CO(R^{16})$, $N(R^{15})CO(\text{alkyl}(N(R^{15})_2))$, $N(R^{15})COCO_2(R^{15})$,
 $N(R^{15})COCON(R^{15})(R^{15})$, $N(R^{15})CO(N(R^{15})_2)$, $N(R^{15})SO_2R^{16}$, $N(R^{15})SO_2(N(R^{15})_2)$,
or $N(R^{15})COAr^3$;

10 or R^{14} is hydroxy, alkoxy, (hydroxy)alkoxy, (alkoxy)alkoxy, $(R^{16}CO_2)\text{alkoxy}$, or
 $(PhCO_2)\text{alkoxy}$;




15 R^{15} is hydrogen, alkyl, haloalkyl, or cycloalkyl;

or $N(R^{15})(R^{15})$ taken together is azetidinyl, pyrrolidinyl, $(R^{10})\text{-piperidinyl}$,
 $N-(R^{11})\text{-piperazinyl}$, morpholinyl, or thiomorpholinyl;

20 R^{16} is alkyl, haloalkyl, or cycloalkyl;


Ar¹ is




Ar² is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyrimidinyl, pyrazinyl, pyridinyl, or hydroxypyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, cyano, benzyl, alkyl, alkoxy, N(R⁸)(R⁹), CO₂R⁶, and CON(R⁸)(R⁹);

Ar³ is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, or pyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, alkoxy, cyano, benzyl, and alkyl; and

X-Y-Z is




or a pharmaceutically acceptable salt thereof.



3. A compound of claim 1 where R¹ is .

20



4. A compound of claim 1 where R¹ is and R³ is other than hydrogen or halo.

5. A compound of claim 1 where R³ is CON(R⁸)(R⁹), SO₂N(R⁸)(R⁹), azetidinonyl, pyrrolidinonyl, valerolactamyl, caprolactamyl, oxazolidinonyl, imidazolidinonyl, dioxothiazinyl, N-(R⁶)-dioxothiazolidinyl, or triazolyl where triazolyl is substituted with 0-2 alkyl or oxo substituents.

5


6. A compound of claim 1 where R³ is halo, CON(R⁸)(R⁹), oxazolidinonyl, or triazolyl where triazolyl is substituted with 0-2 alkyl or oxo substituents.

8. A compound of claim 1 where $R^{14}N(R^{15})(R^{15})$, $N(R^{15})(benzyloxycarbonyl)$,
 $N(R^{15})(alkyloxycarbonyl)$, $N(R^{15})(hydroxylalkyloxycarbonyl)$,
 $N(R^{15})(alkyloxyalkyloxycarbonyl)$, $N(R^{15})CO(R^{16})$, $N(R^{15})(CO(N(R^{15})_2)alkyl)$,
 $N(R^{15})(CO_2CO(N(R^{15})_2)alkyl)$, $N(R^{15})CO((N(R^{15})_2)alkyl)$,
15 $N(R^{15})CO((CON(R^{15})(R^{15})alkyl)$, $N(R^{15})COCO_2(R^{15})$, $N(R^{15})CO((CO_2(R^{15})alkyl)$,
 $N(R^{15})COCON(R^{15})(R^{15})$, $N(R^{15})COCO((N(R^{15})(R^{15}))alkyl)$, $N(R^{15})CO(N(R^{15})_2)$,
 $N(R^{15})SO_2R^{16}$, $N(R^{15})SO_2(N(R^{15})_2)$, or $N(R^{15})COAr^3$;

or R¹⁴ is hydroxy, alkoxy, (hydroxy)alkoxy, (alkoxy)alkoxy, (R¹⁶CO₂)alkoxy, or
20 (PhCO₂)alkoxy;

or R¹⁴ is N(R¹⁵)COCON(R¹⁵)(alkoxyalkyl), N(R¹⁵)COCON(R¹⁵)(hydroxyalkyl), N(R¹⁵)COCON(R¹⁵)(tetrahydrofuryl)methyl), N(R¹⁵)COCON(R¹⁵)(N,N-dimethylpyrazolyl)methyl), or N(R¹⁵)CO((Ar⁴)alkyl);

25



9. A compound of claim 1 where R¹ is (Ar¹)alkyl, alkyl, (cycloalkyl)alkyl, diphenylalkyl, phenoxyalkyl, (PhNH)alkyl, (methylpyrrolidinyl)alkyl, (imidazolyl)alkyl (valerolactamyl)alkyl, (tetrahydrofuranyl)alkyl, ((fluoro)(methyl)pyridinyl)methyl, phenylcyclopropyl, or benzylpyrrolidinyl;

5

R² is hydrogen;

R³ is hydrogen, halo, hydroxy, alkoxy, cyano, alkyl, cycloalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, alkoxy, haloalkoxy, N(R⁸)(R⁹), N(R⁶)COR⁷, N(R⁶)SO₂R⁷,

10 N(R⁶)CO₂R⁷, N(R⁶)SO₂N(R⁸)(R⁹), CO₂R⁶, CON(R⁸)(R⁹), SO₂R⁷, SO₂N(R⁸)(R⁹), or Ar²;

R⁴ is hydrogen, halo, or alkyl;

15 R⁵ is hydrogen, halo, or alkyl;

R⁶ is hydrogen, alkyl, haloalkyl, or cycloalkyl;

R⁷ is alkyl, haloalkyl, or cycloalkyl;

20

R⁸ is hydrogen or alkyl;


R⁹ is hydrogen or alkyl; or

25 N(R⁸)(R⁹) taken together is azetidinyl, pyrrolidinyl, (R¹⁰)-piperidinyl, N-(R¹¹)-piperazinyl, morpholinyl, thiomorpholinyl, or dioxothiazinyl;

R¹⁰ is hydrogen, alkyl, hydroxy, or hydroxyalkyl;

30 R¹¹ is hydrogen, alkyl, cycloalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyalkoxyalkyl, phenyl, pyridinyl, (methylimidazolyl)methyl, COR⁶, CO₂R⁶, (hydroxyalkyl)CO, (alkoxyalkyl)CO, (tetrahydrofuranyl)CO, (methylisoxazolyl)CO, (thienyl)CO, (furanyl)CO, (pyridinyl)CO, CON(R⁸)(R⁹), SO₂R⁷, SO₂N(R⁸)(R⁹),

(dimethylisoxazolyl)SO₂, ((carboethoxy)thienyl)SO₂, (methylimidazolyl)SO₂,



R¹³ is hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, benzyloxycarbonyl, or

5 alkylsulfonyl;

R¹⁴ is N(R¹⁵)(R¹⁵), N(R¹⁵)(benzyloxycarbonyl), N(R¹⁵)(alkyloxycarbonyl),

N(R¹⁵)((hydroxyalkyl)oxycarbonyl), N(R¹⁵)((alkyloxy)alkyloxycarbonyl),

N(R¹⁵)CO(R¹⁶), N(R¹⁵)((CO(N(R¹⁵)₂)alkyl), N(R¹⁵)((CO₂CO(N(R¹⁵)₂)alkyl),

10 N(R¹⁵)CO((N(R¹⁵)₂)alkyl), N(R¹⁵)CO((CON(R¹⁵)(R¹⁵)alkyl), N(R¹⁵)COCO₂(R¹⁵),

N(R¹⁵)CO((CO₂(R¹⁵)alkyl), N(R¹⁵)COCON(R¹⁵)(R¹⁵),

N(R¹⁵)COCO((N(R¹⁵)(R¹⁵)alkyl), N(R¹⁵)CO(N(R¹⁵)₂), N(R¹⁵)SO₂R¹⁶,


N(R¹⁵)SO₂(N(R¹⁵)₂), or N(R¹⁵)COAr³;


15 or R¹⁴ is hydroxy, alkoxy, (hydroxy)alkoxy, (alkoxy)alkoxy, (R¹⁶CO₂)alkoxy, or
(PhCO₂)alkoxy;

or R¹⁴ is N(R¹⁵)COCON(R¹⁵)(alkoxyalkyl), N(R¹⁵)COCON(R¹⁵)(hydroxyalkyl),

N(R¹⁵)COCON(R¹⁵)((tetrahydrofuran-2-yl)methyl), N(R¹⁵)COCON(R¹⁵)((N,N-

20 dimethylpyrazolyl)methyl), or N(R¹⁵)CO((Ar⁴)alkyl);






R¹⁵ is hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, haloalkyl, or cycloalkyl;

10 or N(R¹⁵)(R¹⁵) taken together is azetidinyl, pyrrolidinyl, (R¹⁰)-piperidinyl, N-(R¹¹)-piperazinyl, morpholinyl, or thiomorpholinyl, and is substituted with 0-2 substituents selected from halo, hydroxy, alkoxy, alkyl, hydroxyalkyl, alkoxyalkyl, hydroxalkyloxyalkyl, acetamido, CO₂R⁶, and CON(R⁸)(R⁹);

15 R¹⁶ is alkyl, haloalkyl, cycloalkyl, (cycloalkyl)alkyl, hydroxyalkyl, alkoxyalkyl, tetrahydropyranyl, or Ar³;

R^{17} is hydrogen, alkyl, alkylCO, cycloalkylCO, alkyloxyCO, CON(R^{15})(R^{15}), COCON(R^{15})(R^{15}), COAr₃, alkylSO₂, cycloalkylSO₂, furanylSO₂, triazolylSO₂, or N-methylpyrrolylSO₂;




10 Ar² is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyrimidinyl, pyrazinyl, pyridinyl, or hydroxypyridinyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, cyano, benzyl, alkyl, alkoxy, N(R^8)(R^9), CO₂R⁶, and CON(R^8)(R^9);

15 Ar³ is tetrazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furanyl, thienyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, imidazopyridinyl, quinolinyl, isoquinolinyl, or phenyl, and is substituted with 0-3 substituents selected from the group consisting of oxo, halo, alkoxy, haloalkoxy, cyano, benzyl, alkyl, haloalkyl, N(R^8)(R^9), and N(R^{15})CO(R^{16});

20 Ar⁴ is pyrazolyl or triazolyl, and is substituted with 0-1 substituents selected from CO₂R⁶ and CON(R^8)(R^9); and

X-Y-Z is

25



5 or a pharmaceutically acceptable salt thereof.

10. A compound of claim 9 where R¹ is (Ar¹)alkyl and R² is hydrogen.


11. A compound of claim 10 where

10

R¹⁴ is N(R¹⁵)(R¹⁵), N(R¹⁵)(benzyloxycarbonyl), N(R¹⁵)(alkyloxycarbonyl), N(R¹⁵)((hydroxyalkyl)oxycarbonyl), N(R¹⁵)((alkyloxy)alkyloxycarbonyl), N(R¹⁵)CO(R¹⁶), N(R¹⁵)((CO(N(R¹⁵)₂)alkyl), N(R¹⁵)((CO₂CO(N(R¹⁵)₂)alkyl), N(R¹⁵)CO((N(R¹⁵)₂)alkyl), N(R¹⁵)CO((CON(R¹⁵)(R¹⁵)alkyl), N(R¹⁵)COCO₂(R¹⁵),
15 N(R¹⁵)CO((CO₂(R¹⁵)alkyl), N(R¹⁵)COCON(R¹⁵)(R¹⁵), N(R¹⁵)COCO((N(R¹⁵)(R¹⁵))alkyl), N(R¹⁵)CO(N(R¹⁵)₂), N(R¹⁵)SO₂R¹⁶, N(R¹⁵)SO₂(N(R¹⁵)₂), or N(R¹⁵)COAr³.

12. A compound of claim 11 where X-Y-Z is

20



or a pharmaceutically acceptable salt thereof.

13. A compound of claim 1 selected from the group consisting of
ethanediamide, N-[2-[[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N',N'-trimethyl-;
5 ethanediamide, N-[2-[[[(3-chloro-4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N',N'-trimethyl-;
10 ethanediamide, N-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N',N'-trimethyl-;
15 ethanediamide, N'-[2-[[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-;
20 ethanediamide, N'-[2-[[[(3-cyclopropyl-4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-;
ethanediamide, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-6,7,8,9-tetrahydro-3-hydroxy-4-oxo-7,10-ethanopyrimido[1,2-a]azepin-10(4H)-yl]-N,N-dimethyl-;
25 8,11-methano-4H-pyrimido[1,2-a]azocene-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-, (8R,11S)-;
ethanediamide, N-(2-fluoroethyl)-N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-30 7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N-methyl-;

- ethanediamide, N'-(8R,11S)-2-[[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-;
- 5 ethanediamide, N'-[2-[[[(4-fluorophenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-;
- 10 ethanediamide, N'-[2-[[[(4-fluoro-3-methylphenyl)methyl]amino]carbonyl]-7,8,9,10-tetrahydro-3-hydroxy-4-oxo-8,11-methano-4H-pyrimido[1,2-a]azocin-11(6H)-yl]-N,N-dimethyl-;
- 15 8,11-methano-4H-pyrimido[1,2-a]azocene-2-carboxamide, N-[(4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-;
- 20 8,11-methano-4H-pyrimido[1,2-a]azocene-2-carboxamide, N-[(3-chloro-4-fluorophenyl)methyl]-11-[[2-[(3R)-3-fluoro-1-pyrrolidinyl]-1,2-dioxoethyl]amino]-6,7,8,9,10,11-hexahydro-3-hydroxy-4-oxo-; and
- 25 or a pharmaceutically acceptable salt thereof.

14. A composition useful for treating HIV infection comprising a therapeutic amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

30 15. The composition of claim 14 further comprising a therapeutically effective amount at least one other agent used for treatment of AIDS or HIV infection selected from the group consisting of nucleoside HIV reverse transcriptase inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, HIV protease inhibitors, HIV fusion

inhibitors, HIV attachment inhibitors, CCR5 inhibitors, CXCR4 inhibitors, HIV budding or maturation inhibitors, and HIV integrase inhibitors, and a pharmaceutically acceptable carrier.

5 16. A method for treating HIV infection comprising administering a therapeutically effective amount of a compound of claim 1, or a pharmaceutically acceptable salt thereof, to a patient in need thereof.

10 17. The method of claim 16 further comprising administering a therapeutically effective amount of at least one other agent used for treatment of AIDS or HIV infection selected from the group consisting of nucleoside HIV reverse transcriptase inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, HIV protease inhibitors, HIV fusion inhibitors, HIV attachment inhibitors, CCR5 inhibitors, CXCR4 inhibitors, HIV budding or maturation inhibitors, and HIV integrase
15 inhibitors.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/037596

A. CLASSIFICATION OF SUBJECT MATTER			
INV.	C07D498/18	A61K31/5386	C07D471/18
	A61K31/55	C07D471/08	A61P31/18
			A61K31/529 C07D487/08
			C07D493/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2005/118593 A (SQUIBB BRISTOL MYERS CO [US]; NAIDU B NARASIMHULU [US]; BANVILLE JACQU) 15 December 2005 (2005-12-15) cited in the application the whole document	1-17
Y	WO 2004/058756 A (ANGELETTI P IST RICERCHE BIO [IT]; CRESCENZI BENEDETTA [IT]; KINZEL O) 15 July 2004 (2004-07-15) cited in the application the whole document	1-17
Y	WO 2007/027754 A (SQUIBB BRISTOL MYERS CO [US]; BANVILLE JACQUES [CA]; REMILLARD ROGER []) 8 March 2007 (2007-03-08) claims 1,12-14	1-17

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
22 July 2009	29/07/2009
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Seymour, Liza

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2009/037596

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2007/061714 A (SQUIBB BRISTOL MYERS CO [US]; NAIDU B NARASIMHULU [US]; UEDA YASUTSUGU) 31 May 2007 (2007-05-31) the whole document -----	1-17
Y	WO 2006/103399 A (ISTITUTO DE RICERCHE DI BIOLOG [IT]; CECERE GIUSEPPE [IT]; FERREIRA MA) 5 October 2006 (2006-10-05) the whole document in particular compounds 158 and 173 -----	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

 International application No
 PCT/US2009/037596

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2005118593	A	15-12-2005	AR 049124 A1 AT 412656 T AU 2005250356 A1 BR PI0511623 A CA 2568356 A1 DK 1749011 T3 EP 1749011 A1 ES 2315875 T3 HR 20090005 T3 JP 2008501018 T KR 20070039502 A NZ 551282 A PT 1749011 E US 2005267105 A1 US 2006199956 A1 US 2006276466 A1	28-06-2006 15-11-2008 15-12-2005 11-03-2008 15-12-2005 16-02-2009 07-02-2007 01-04-2009 28-02-2009 17-01-2008 12-04-2007 31-05-2009 27-11-2008 01-12-2005 07-09-2006 07-12-2006
WO 2004058756	A	15-07-2004	AU 2003292436 A1 AU 2003292437 A1 BR 0317749 A CA 2509554 A1 CN 1753892 A EC SP055890 A EP 1578748 A1 WO 2004058757 A1 HR 20050593 A2 IS 7892 A JP 2006513200 T KR 20050087865 A MA 27583 A1 MX PA05007010 A NZ 540729 A RU 2329265 C2 US 2008176869 A1 US 2006046985 A1	22-07-2004 22-07-2004 22-11-2005 15-07-2004 29-03-2006 20-09-2005 28-09-2005 15-07-2004 31-03-2006 13-06-2005 20-04-2006 31-08-2005 03-10-2005 18-08-2005 28-03-2008 20-07-2008 24-07-2008 02-03-2006
WO 2007027754	A	08-03-2007	CN 101253177 A EP 1919921 A1 JP 2009508817 T	27-08-2008 14-05-2008 05-03-2009
WO 2007061714	A	31-05-2007	AR 057896 A1 AU 2006316582 A1 EP 1948666 A1 JP 2009515981 T KR 20080069254 A US 2007111984 A1	26-12-2007 31-05-2007 30-07-2008 16-04-2009 25-07-2008 17-05-2007
WO 2006103399	A	05-10-2006	AU 2006228278 A1 CA 2600832 A1 CN 101146811 A EP 1866313 A1 JP 2008534561 T	05-10-2006 05-10-2006 19-03-2008 19-12-2007 28-08-2008