



US 20130143856A1

(19) **United States**(12) **Patent Application Publication****Kim et al.**(10) **Pub. No.: US 2013/0143856 A1**(43) **Pub. Date: Jun. 6, 2013**(54) **NOVEL ANTIBIOTIC COMPOSITION
COMPRISING FLUFENAMIC ACID AS AN
ACTIVE INGREDIENT**(75) Inventors: **Kyeong Kyu Kim**, Seoul (KR); **Sung
Wook Kang**, Suwon-si (KR)(73) Assignee: **SUNGKYUNKWAN UNIVERSITY
FOUNDATION FOR CORPORATE
COLLABORATION**, Suwon-si (KR)(21) Appl. No.: **13/522,764**(22) PCT Filed: **Jan. 21, 2011**(86) PCT No.: **PCT/KR2011/000444**

§ 371 (c)(1),


(2), (4) Date: **Oct. 9, 2012**(30) **Foreign Application Priority Data**

Jan. 22, 2010 (KR) 10-2010-0006217

Publication Classification(51) **Int. Cl.****A61K 31/196** (2006.01)**C12N 9/04** (2006.01)**A61K 31/496** (2006.01)**A61K 31/431** (2006.01)**A61K 31/546** (2006.01)**A61Q 19/00** (2006.01)**A61K 8/44** (2006.01)**A61K 8/49** (2006.01)**A61Q 5/02** (2006.01)**A61Q 5/06** (2006.01)**A61Q 5/10** (2006.01)**A61Q 19/10** (2006.01)**C07C 229/58** (2006.01)**A61P 31/04** (2006.01)(52) **U.S. Cl.**USPC **514/196**; 562/456; 435/190; 514/567;
514/253.08; 514/202(57) **ABSTRACT**


The present invention relates to a novel antibiotic, and more specifically relates to an antibiotic composition comprising flufenamic acid as an active ingredient, which can reduce drug toxicity and side effects and the problem of antibiotic resistance caused by excessive use of antibiotics since the composition exhibits a good therapeutic effect, even in a small dose, when it is administered either alone or together with another antibiotic of the prior art.

[Fig. 1]




Flufenamic acid


[Fig. 2]



[Fig. 3]



[Fig. 4]



[Fig. 5]

```
AKR1C3      MDSKQQCVKLNDCGFMPVLGFTYAPPEVPRSKALEVTKLAIEAGFRHIDSABLYNNEEQ
SA0658      --MLNEIQILNNGYFMPVGLCVYKIS---DEDMTKVVNAIDAGYRAFDYFYDN S
SA1606      ----MEVKTFYNGNTMPQIGLGTFRVEN--DENCMESVKYAIQQGYRSIDTAKVYGNNEEQ
SA2001      ----MNHTEISKDVKIPVLGFGVFIPIQ---EQTAEA VKAIAKAGYRHIDTAQSYLNETE
              :  :  . .  :* :*:*. :  . .  :  :  **  :* :*:  * ** .

AKR1C3      VGLAIRSKIADGSGVKREDIFYTSKLWSTFHRPELVREPALENSLXKAQLDYVDLYLIHSPM
SA0658      LGR---ALKDNGVDREDLFITTKLWNDYQGYEKTFEYFNKSIENLQTDYLDLFLIHWPC
SA1606      VGAGIRAGLESTGIAREDLFITSKLYFEDFGRENVAAAYEASLSRLGLKYLDLYLVHWPG
SA2001      VGQG----IEASGIORSELFITTKVWIENVNYEDTIKSIERSLQRLNLDYLDLVLIHQPY
              :*      :  :  :*:** :*: :  * .  :  :*. . . :*:** :*:


AKR1C3      SLKPGEELSPTDENGKVIDIVDLCTTWEAMEKCKDAGLAKSIGVSNFNRRLQEMILNKP
SA0658      EADG-----LPLETYKAMEELYEQGKVKAGVCNFNHLEKLMQAS
SA1606      TNEA-----VMVDTWKGMEDLYKNNKAKNIGVSNFEPEHLEALLAQV
SA2001      NDVYG-----SWRALEELKENGKIKAGVSNFGVDRIVDLGIHN
              : : : :*. . . * ***,** : : : :

AKR1C3      GLKYKPCVNQVECHPYFNRSKLLDFCKSKDIVLVAYSALGSQRDKRWVDPNSFVLLEDPV
SA0658      SIK--P VNQIEVHPYFNQQLQEFCDRHDIKVTAWMPLMR-----NRGLLDNPV
SA1606      SIK--PVINQVEYHPYLTQHKLKLYLAAQHIVMESWSPL -----NAQILNDET
SA2001      QIQ--PQVNQIEINPFHQEEQVAALQQENVVVEAWAPFAE-----GKNQLFQNL
              : :  *  **:* :*: :  :  . .  :  :  :  :  :  :  :  :

AKR1C3      LCATAKKKXRTPALIALRYQLQRCVVVIAKSYNEQRIRQNVQVFEFQLTAEOMKAIDGLD
SA0658      IVKIAEKYBKTPAQVVLRWBLAHNRILIPKSQTPKRIQENIDILDFNLELTEVAEIDALD
SA1606      IKDIAQELGKSPAQVVLRWNVQHGCVIIPKSVTPNRISENFQIFDFELSDEQMTRIDGLN
SA2001      LQAIADKYNKSIQVILRWLVERDIVVLAKSVNPERMAQNLDIFDFELTEEDKQOIATLE
              :  :*: :  :  * : ** :  :  :  : ** . :*: :*: :*: :* :  *  * :


AKR1C3      RNLHYFNSDSFASHPNYPYSDEYLEHHHHHH
SA0658      R-----NARQGKNPDDVKIGDLK-----
SA1606      Q-----DKRIGPDPKTFEG-----
SA2001      ES-----NSQFFSHADPEMIKALTSRELDV-
              . . . . .
```

[Fig. 6]




[Fig. 7]


A M 1 2 3 4 5





B M 1 2 3 4 5 6 7



[Fig. 8]



[Fig. 9]



NOVEL ANTIBIOTIC COMPOSITION COMPRISING FLUFENAMIC ACID AS AN ACTIVE INGREDIENT

TECHNICAL FIELD

[0001] The present invention is to provide a novel antibiotic material effective for killing or inhibiting the growth or metabolism of Gram-positive bacteria.

BACKGROUND ART

[0002] Gram staining is a method of differentiating bacterial species into two large groups which are the Gram-positive and Gram-negative bacteria, which was devised by Danish physician H. C. J. Gram (1853-1938) in 1884 not for the purpose of distinguishing one type of bacterium from another but to enable bacteria to be more visible in stained sections. These bacterial groups are generally different in various aspects including sensitivity to chemical agents, nutrients necessary for their growth, response to physical and chemical stimuli, toxins produced thereby, and lesions caused thereby. Representative among Gram-positive bacteria, characterized by a purple color stain, are *Staphylococcus* spp., *Pneumococcus* spp., *Mycobacterium leprae*, *Corynebacterium diphtheria*, *Clostridium tetani*, *Bacillus anthracis*, and *Actinomyces*. Most of these bacteria are leading pathogens responsible for serious disease such as diphtheria, tuberculosis, pneumonia, etc.

[0003] For example, *Staphylococcus aureus*, a Gram-positive bacterium, is a pathogen which causes various diseases including food poisoning, impetigo, cellulitis, scalded skin syndrome, mastitis, bacteremia, sepsis, *staphylococcal* pneumonia, endocarditis, heart failure, osteomyelitis, *Staphylococci sepsis*, circulatory collapse, and toxic shock syndrome. In the United States, *staphylococcal* infection is diagnosed in more than two million people every year and accounts for the death of more than ninety thousand people.

[0004] Most bacterial infections are readily cured by antibiotics such as penicillin, etc., but some have recently been reported to be difficult to treat. Bacteria become resistant to antibiotic drugs through a spontaneous DNA mutation or the uptake of foreign DNA so that they are able to survive exposure to an antibiotic. This situation is called "antibiotic resistance" and the bacteria with antibiotic resistance are called "drug resistant bacteria." In addition, drug-resistant bacteria show multidrug resistance in which, for example, only 10% and 50% of them can be regulated with penicillin and methicillin, respectively.

[0005] Antibiotic resistance has been found over a broad spectrum of Gram-positive pathogenic bacteria including *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacillus* and *Listeria*. Of them, MRSA (Methicillin-Resistant *Staphylococcus Aureus*), VRSA (Vancomycin-Resistant *Staphylococcus Aureus*), MRE (multiple-drug resistant enterococci) and VRE (Vancomycin-Resistant enterococci) attract social interest because they are more dangerous.

[0006] For example, MRSA is treated with vancomycin, more potent than methicillin, but some show resistance to the drug, as well. Because such resistant pathogens are poor in susceptibility to drugs, it is often necessary to apply a large dose of antibiotics which is, however, noxious to the patient. In recent years, the increasing spread of drug resistance by *S. aureus* has been recognized as serious threat to the health of mankind. Therefore, there is a pressing need for a novel

antibiotic that works through a new mechanism that confers a potent therapeutic effect even at a small dose thereof on an infection of the drug-resistant *S. aureus*.

[0007] Classically, new antibiotics have been developed by chemically modifying bioactive compounds produced by microorganisms. With the great progress of genomic and drug discovery techniques such as high-throughput screening, many new drug targets are secured from which various drug candidates have been separated. The time and cost which it takes for such new drug candidates to undergo clinical trials and to acquire the approval of the FDA is currently acting as the main hindrance to the development of novel drugs. Hence, if found, new uses of commercially available drugs which have passed clinical trials and have had their safety verified in vivo will provide the advantage of significantly reducing the time and cost required to bring novel drugs to market.

[0008] Given this background, the present inventors have searched for finding a drug that works in a manner different from already known antibiotic mechanisms in the art in order to overcome the problem with multidrug resistance and, in addition, for finding a compound that exhibits antibiotic activity from among drugs currently used in clinics in order to make it easier to develop a novel antibiotic agent.

DISCLOSURE

Technical Problem

[0009] The present invention is intended to provide a novel antibiotic destructive of or inhibitory of the growth or metabolism of Gram-positive bacteria which does not confer drug resistance on the bacteria but is effective on drug resistant bacteria, and to reduce the time and cost taken for the development of a novel antibiotic by taking advantage of a preexisting drug currently used in clinics. More particularly, the present invention provides an antibiotic composition with potent inhibitory activity against Gram-positive bacteria, comprising flufenamic acid.

Technical Solution

[0010] The present invention newly discovers the use of flufenamic acid, currently used for anti-inflammation, as a novel antibiotic in the treatment of bacterial infections. More particularly, the present invention provides an antibiotic composition comprising flufenamic acid, a derivative thereof, an isomer thereof, or a salt thereof as an active ingredient, and various uses thereof, that is, a pharmaceutical or cosmetic composition with antibiotic activity.

[0011] In addition, the present invention is directed to an aldo-keto reductase, present in Gram-positive bacteria, which can be utilized as a target protein of an antibiotic for destroying or inhibiting the growth or metabolism of Gram-positive bacteria.

[0012] Leading to the present invention, intensive and thorough research was done regarding an antibiotic which has a working mechanism that is different from classical antibiotics. This research resulted in the finding that flufenamic acid, currently used as an analgesic, can inhibit the growth and metabolism of Gram-positive bacteria through a novel antibiotic mechanism.

[0013] Below, a detailed description will be given of the present invention.

[0014] The present invention addresses an antibiotic composition comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt as an active ingredient.

[0015] In the present invention, flufenamic acid is used to regulate the growth or metabolism of Gram-positive bacteria, that is, it has potent inhibitory activity against Gram-positive bacteria.

[0016] No special limitations are imparted to species of the Gram-positive bacteria against which the antibiotic composition of the present invention exhibits inhibitory activity. Preferred examples include *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes*, and *Enterococcus faecium*. More preferred is methicillin-resistant *Staphylococcus aureus* (MRSA).

[0017] Particularly, the present invention relates to an antibiotic composition with potent destructive and inhibitory activity against drug-resistant bacteria, comprising flufenamic acid as an active ingredient. Included among the drug-resistant bacteria are MRSA (Methicillin-Resistant *Staphylococcus aureus*), VRSA (Vancomycin-Resistant *Staphylococcus aureus*), MRE (multiple-drug resistant enterococci) and VRE (Vancomycin-Resistant enterococci).

[0018] Any flufenamic acid, whether naturally occurring or artificially synthesized, may be used in the present invention. If artificially synthesized, flufenamic acid may be prepared from any material using any method. For example, flufenamic acid may be synthesized through substitution and fraction (Herbert O. House: Modern Synthetic Reactions, 2nd ED., The Benjamin/Cummings Publishing Co., 1972).

[0019] Instead of flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt may be used as an active ingredient of the composition according to the present invention.

[0020] It is naturally understood that all stereoisomers, diastereomers, enantiomers, and racemates of flufenamic acid fall within the scope of the present invention.

[0021] The salt of flufenamic acid useful in the present invention may be an acid salt or a base salt. By way of example, pharmaceutically acceptable salts of flufenamic acid include sodium, calcium and potassium salts for the hydroxy group and hydrobromide, sulfate, hydrogen sulfate, phosphate, hydrogen phosphate, dihydrogen phosphate, acetate, succinate, citrate, tartrate, lactate, mandelate, methanesulfonate (mesylate) and p-toluenesulfonate (tosylate) for the amino group, but are not limited thereto. They may be prepared using methods or techniques well known in the art.

[0022] flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt act as an active ingredient in the antibiotic composition provided by the present invention, targeting the aldo-keto reductase of Gram-positive bacteria.

[0023] As stated above, flufenamic acid inhibits the activity of aldo-keto reductase in Gram-positive bacteria. For example, the aldo-keto reductase which flufenamic acid target may be the enzymes present in *Staphylococcus aureus*, and may preferably be SA0658, SA1606 or SA2001.

[0024] In one embodiment of the present invention, the antibiotic composition may further comprise a conventional antibiotic against Gram-positive bacteria in addition to flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt thereof. So long as it is approved for use in the treatment of the infection of Gram-positive

bacteria, any conventional antibiotic may be employed in the composition of the present invention. Preferred is norfloxacin, oxacillin or cephalothin.

[0025] Also, the present invention addresses a Gram-positive bacterial aldo-keto reductase as a target protein of an antibiotic that is destructive or inhibitory of the growth or metabolism of Gram-positive bacteria.

[0026] The Gram-positive bacteria affected by the present invention are not limited specifically to species so long as they contain aldo-keto reductase therein. Preferred examples include *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes* and *Enterococcus faecium*. More preferred is methicillin-resistant *S. aureus* (MRSA).

[0027] In the present invention, the aldo-keto reductase functions as a target protein on which an inhibitory agent acting against the growth or metabolism of Gram-positive bacteria directly acts.

[0028] No particular limitations are imposed on the kinds of the inhibitory agent, whether natural or synthetic. The aldo-keto reductase which is regulated by the inhibitory agent may be an enzyme which is found in Gram-positive bacteria.

[0029] For example, the aldo-keto reductase may be found in *Staphylococcus aureus* and preferably SA0658 with the amino acid sequence of SEQ ID NO: 1, SA1606 with the amino acid sequence of SEQ ID NO: 2 or SA2001 with the amino acid sequence of SEQ ID NO: 3.

[0030] Further, the present invention addressed an antibiologically active, pharmaceutical composition comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt as an active ingredient.

[0031] Showing inhibitory activity against Gram-positive bacteria without side effects and toxicity in the body, the composition is useful for preventing and treating diseases caused by Gram-positive bacteria. No particular limitations are imparted on the type of inhibition exhibited against Gram-positive bacteria. Preferably, it destroys or inhibits the growth or metabolism of Gram-positive bacteria.

[0032] The pharmaceutical composition of the present invention may be administered orally or parenterally to mammals such as rats, dogs, domestic animals, and humans. For example, it may be administered orally or injected intrarectally, intramuscularly or subcutaneously, or via the route of the dura mater within the uterine or via a cerebrovascular route. Preferable are parenteral routes, with greater preference for transdermal administration. Thus, the composition may be applied topically.

[0033] For practical use in the destruction or inhibition of growth or metabolism of Gram-positive bacteria, the composition in accordance with the present invention may be formulated into typical forms, for example, oral dosage forms such as tablets, capsules, troches, liquids, suspensions, etc., injection forms such as injectable solutions or suspensions, powder that can be reconstituted with distilled water for injection, and ready-to-use solutions, topical forms such as ointments, creams, lotions, liquids, sprays, etc.

[0034] So long as it is pharmaceutically accepted, any vehicle may be used for formulating the composition of the present invention. By way of example, the composition may be formulated with a binder, a suspension agent, a disintegrant, an excipient, a solubilizer, a dispersant, a stabilizer, a colorant, and/or a flavorant for oral dosage forms, or with a preservative, an appeaser, a solubilizer and/or a stabilizer for injection forms, or with a base, an excipient, a lubricant, a

preservative and/or a propellant for topical forms. The pharmaceutical formulations thus prepared may be administered orally or applied topically.

[0035] The effective dosage of the pharmaceutical composition in accordance with the present invention depends on various factors, including dosage form, the patient's age, weight, gender, state of health, diet, the time of administration, the route of administration, excretion rate, sensitivity, etc. In general, it may be administered in a single dose or in multiple doses per day at a daily dose ranging from 0.01 to 200 mg/day and preferably at a daily dose ranging from 5 to 50 mg/kg.

[0036] The antibiotically active, pharmaceutical composition comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt in accordance to the present invention may further comprise an antibiotic inhibitory of Gram-positive bacteria. Any antibiotic, if approved for use in inhibiting Gram-positive bacteria, may be used in the composition of the present invention. Preferred examples of the antibiotic include norfloxacin, oxacillin and cephalothin, but are not limited thereto.

[0037] In addition, the present invention addresses an antibiotically active, cosmetic composition comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt.

[0038] The cosmetic composition may further comprise an additive in addition to flufenamic acid as an active ingredient. The additive, if conventionally used, may be contained in the cosmetic composition. For example, a supplement and an excipient such as an antioxidant, a stabilizer, a solubilizer, vitamin, a colorant, a flavorant, etc. may be used. In the cosmetic composition, a transdermal delivery enhancer may be employed.

[0039] The composition of the present invention may contain flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt in an amount of from 0.0001 to 99 wt %, and preferably in an amount of from 0.001 to 90 wt %.

[0040] The cosmetic composition of the present invention may be prepared into any conventional formulation. For example, the cosmetic composition of the present invention may be formulated with a vehicle into cream, lotion, tonic, spray, aerosol, oil, solution, suspension, gel, ointment, emulsion, paste, or hair tonic. The pharmaceutical or cosmetic formulations may be prepared using any apparatus or method well known in the art (refer to Remington's Pharmaceutical Science, latest version).

[0041] In the composition of the present invention, a base may be properly combined with, for example, a cosmetically acceptable additive. So long as it is accepted in the art, any base may be used in the composition. Exemplary among them are distilled water, mineral water, ethanol, glycerin, squalene, 1,3-propyleneglycol, 1,3-butyleneglycol, castor oil, tsubaki oil, and liquid petrolatum. Examples of the additive include a surfactant, an emulsifier, a thickener, a preservative, an antioxidant, and a flavorant, but are not limited thereto. The surfactant may be cationic, anionic or non-ionic. Examples of the cationic surfactant include alkaline salts of higher fatty acids such as lauric acid, myristic acid, palmitic acid, stearic acid and oleic acid (e.g., sodium, potassium, ammonium, triethanolamine salt); esters of alkyl sulfonic acid such as sodium laurylsulfate, and triethanol amine laurylsulfonate; and alkyl ether sulfonate esters such as sodium polyoxyethylene laurylether sulfonate and triethanolamine polyoxyethylene laurylether sulfonate. Included among the non-ionic surfactants are polyoxyethylene alkyl ethers such as polyoxyethylene laurylether and polyoxyethylene oleylether; and alkanolamine such as coconut fatty acid diethanolamide and diethanolamide laurate. Stearyl trimethyl ammonium chloride, cetyl trimethyl ammonium chloride and stearyl bis(diethylalcohol)hydroxy ethyl ammonium chloride are examples of the anionic surfactants. Examples of the emulsifier include cetanol, stearyl alcohol and behenyl alcohol. Examples of the thickener include sodium alginate, methyl cellulose, carboxymethyl cellulose, polyvinyl alcohol and polyvinyl pyrrolidone.

[0042] Examples of the preservative include ethyl parahydroxybenzoate, butyl parahydroxybenzoate and benzalkonium chloride. Examples of the antioxidant include butyl hydroxytoluene, propyl galate and butyl hydroxyanisole. The flavorant may be perfume for general use and may be exemplified by citrus, lavender and floral.

[0043] Various factors including the patient's sex, age and health state, the severity of disease, etc., must be considered in determining the quantity of a dose and the number of doses of the composition according to the present invention.

[0044] The antibiotically active, cosmetic composition comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt in accordance to the present invention may further comprise an antibiotic inhibitory of Gram-positive bacteria. Any antibiotic, if approved for use in inhibiting Gram-positive bacteria, may be used in the composition of the present invention. Preferred examples of the antibiotic include norfloxacin, oxacillin and cephalothin, but are not limited thereto.

Advantageous Effects

[0045] Because the inhibition mechanism used is different from those of conventional antibiotics, flufenamic acid can exert inhibitory activity effectively on bacteria resistant to conventional antibiotics. This novel antibiotic may be used alone or in combination with a conventional antibiotic. In the latter case, high therapeutic effects can be obtained at a low dose, with the concomitant reduction of side effects and toxicity. Because its intrinsic medical efficacy is newly discovered, the antibiotic of the present invention, which has been used as an anti-inflammatory drug, does not need to undergo separate clinical trials for the FDA's approval, thus remarkably reducing the time and cost of drug development and having a great advantage in market entry.

DESCRIPTION OF DRAWINGS

[0046] FIG. 1 shows a chemical structure of flufenamic acid.

[0047] FIG. 2 shows inhibitory concentrations of flufenamic acid against *Staphylococcus aureus*.

[0048] FIG. 3 shows inhibitory concentrations of flufenamic acid against *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes* and *Enterococcus faecium*.

[0049] FIG. 4 shows inhibitory concentration of flufenamic acid used in combination with Norfloxacin, oxacillin, and cephalothin against *Staphylococcus aureus*.

[0050] FIG. 5 shows comparison of sequence alignments of AKR1C3, SA0658, SA1606, SA2001, and 2,5-diketo-D-glucanate reductase A.

[0051] FIG. 6 shows three-dimensional structures and essential residues of AKR1C3, SA0658, SA1606, and SA2001.

[0052] FIG. 7 shows purification of recombinant proteins (SA1606 and SA2001) for examining the function of SaAKRs.

[0053] FIG. 8 shows binding affinity of FLF for SA1606 and SA2001.

[0054] FIG. 9 shows enzymatic activities of AKR on methylglyoxal in the presence and absence of FLF, confirming the inhibitory activity of FLF against SaAKRs.

MODE FOR INVENTION

[0055] The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

EXAMPLES

Example 1

Bacterial Strain

[0056] CCARM3080, a clinically identified species of methicillin-resistant *S. aureus* (MRSA), was purchased from the Culture Collection of antibiotic Resistant Microbes (www.ccar.m.or.kr) and used for screening compounds. *Listeria monocytogenes* (KCTC3586), *Staphylococcus epidermidis* (KCTC1917) and *Bacillus subtilis* (KCTC2217) were obtained from the Korean Collection for Type Cultures (http://kctc.kribb.re.kr). *Enterococcus faecium* was granted from Dr. Ko, K. S. Sungkyunkwan University.

Example 2

Identification of Novel Compounds

[0057] To screen compounds having inhibitory activity against bacteria, MRSA was 200-fold diluted in 0.2 mL of LB in 96-well microplates and cultured overnight at 37° C. with agitation at 180 RPM. Prestwick chemical library was added at a final concentration of 100 µg/mL per well. The microplates were incubated at 37° C. for 24 hours with agitation at 180 RPM and the optical density at 600 nm was measured.

[0058] Of the candidate compounds, flufenamic acid (FLF), shown in FIG. 1, was identified to be the most inhibitory of the bacteria.

Example 3

Assay for Inhibition against Methicillin-Resistant *Staphylococcus aureus*

[0059] FLF was assayed for inhibitory activity against Methicillin-resistant *staphylococcus aureus* (MRSA). In this regard, Methicillin-resistant *Staphylococcus aureus* (MRSA)

was cultured overnight and 200-fold diluted in 2 mL of LB containing various concentrations of FLF. After additional incubation at 37° C. for 8 hours, cell densities were measured by recoding OD at 600 nm.

[0060] As can be seen in FIG. 2 and Table 1, the growth of MRSA was 90% inhibited after incubation with 40 µg/mL FLF for 8 hours (FIG. 2) and the MIC of FLF against MRSA was found to be 64 µg/mL when measured after 24 hours of incubation (Table 1).

Example 4

Assay for Inhibition against *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes*, and *Enterococcus faecium*

[0061] FLF was examined for inhibitory activity against various bacteria. *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes* and *Enterococcus faecium* were cultured overnight and 200-fold diluted in 2 mL of LB containing various concentrations of FLF. Cell densities were determined by measuring the OD at 600 nm after incubation at 37° C. for 8 hours. The MICs of FLF were determined using a broth dilution method. In brief, bacteria were inoculated into LB containing serially diluted concentrations of FLF in 96-well microplates. After 24 hours of incubation, bacterial growth was examined. Antibacterial activity of FLF was tested over various Gram-positive bacteria including *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes*, and *Enterococcus faecium*.

TABLE 1

MICs of Flufenamic Acid against Gram-Positive Bacteria	
Bacteria	MICs (µg/ml)
<i>S. aureus</i> MSSA	64
<i>S. aureus</i> MRSA	64
<i>S. epidermidis</i>	64
<i>B. subtilis</i>	32
<i>L. monocytogenes</i>	32
<i>E. faecium</i>	32

[0062] As shown in FIG. 3 and Table 1, *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes*, and *Enterococcus faecium* were susceptible to FLF. Their growth was 90% inhibited after incubation with 20–40 µg/mL FLF for 8 hours (FIG. 3). The MICs of FLF against these bacteria were found to range from 32 to 64 µg/mL when measured after 24 hours of incubation (Table 1).

Example 5

Synergistic Effect of FLF with Other Antibiotics

[0063] To examine the synergistic effect of FLF with other antibiotics, Methicillin-resistant *Staphylococcus aureus* (MRSA) was cultured overnight and 200-fold diluted in 2 mL of LB containing 20 µg/ml FLF together with 0, 10, 20, 30, or 40 µg/ml Norfloxacin, Oxacillin or Cephalocin. Cell densities were determined by measuring OD at 600 nm after incubation at 37° C. for 8 hours.

TABLE 2

MICs of Antibiotics against MRSA in the Presence of Flufenamic Acid					
Antibiotics (mg/ml)	MICs (μ g/ml)				
	No FLF	10 μ g/ml FLF	20 μ g/ml FLF	30 μ g/ml FLF	40 μ g/ml FLF
Norfloxacin	128	64	32	16	8
Oxacillin	>256	256	128	32	8
Cephalothin	128	128	64	16	4

[0064] As can be seen in Table 2 and FIG. 4, the MIC of Norfloxacin, a representative quinolone antibiotic targeting DNA gyrase, was reduced from 128 to 32 μ g/ml in the presence of 20 μ g/ml FLF (FIG. 4A). Similarly, FLF allows MRSA to be far more susceptible to oxacillin (FIG. 4B) and cephalothin (FIG. 4C), both β -lactam antibiotics targeting the biosynthesis of the cell wall. When used in combination with classical antibiotics, FLF synergistically inhibited MRSA infection and thus showed effective therapeutic effects (Table 2).

Example 6

Analysis of Bacterial Target Protein of Flufenamic Acid Using Bioinformation

[0065] In order to identify the bacterial protein target of FLF, human AKR1C3, known as a protein to be associated with FLF, was compared with a library or database of sequences using the BLAST bioinformatics program.

[0066] As shown in FIG. 5, a Blast search identified ORFs—SA0658, SA1606, and SA2001 as being comparable with the sequence of human AKR1C3. The AKR analogs were found to share sequence homology of 34, 36 and 31%, respectively, with AKR1C3. In addition, these three proteins showed sequence homology of 41~45% with 2,5-diketo-D-gluconate reductase A (DkgA), one member of the AKR family. To examine whether FLF is associated with the ORFs (SA0658, SA1606, and SA2001) identified in *S. aureus* MRSA, DkgA of *E. coli* the three-dimensional structure of which had been established was used as a template for remodeling the three-dimensional structures of the proteins of interest. Primary modeling was performed with DS MODELLER, followed by optimization using the protocol of Discovery Studio (Accelrys, Inc). The models thus built were qualitatively checked using PROCHECK. Sequence alignments among AKR1C3, SA0658, SA1606, SA2001, and DkgA were established using the CLUSTALW program with a default parameter.

[0067] As can be seen in FIG. 6, analysis of the crystal structure of human AKR1C3 complexed with FLF exhibited that FLF binds to the active site of AKR1C3. The inhibitory activity of FLF is due to the occupation of the oxyanion hole of AKR1C3 having a carboxylic group by FLF through hydrogen bonds with His117 and Tyr55 (FIG. 6). Each of the three AKR analogs SA0658, SA1606 and SA2001, identified in *S. aureus*, has four essential residues Asp50, Tyr55, Lys84, and His117, which are highly conserved in the AKR family and play an important role in enzymatic activity.

[0068] Analysis results showed that SaAKRs were significantly similar to human AKR1C3 in entire structure (FIG. 6A) as well as active site structure (FIG. 6B) and that the docking of FLF into the active site was successfully achieved (FIG. 6C). In this docking model, the oxygen atoms of the

carboxylic acid group form hydrogen bonds with Tyr49 and His111, allowing FLF to occupy the oxyanion hole of the active site. Consequently, the three AKR analogs SA0658, SA1606 and SA2001, identified in *S. aureus*, were observed to have the same structure and working mechanism as those of ARK.

Example 7

Identification of the Binding of FLF to Target Proteins SA1606 and SA2001 by Gene Cloning and Protein Purification

[0069] To examine the practical binding affinity of FLF for *S. aureus* protein targets (SA0658, SA1606, SA2001) and inhibitory activity against *S. aureus*, the targets were cloned. In the case of SA0658, although it was cloned and expressed, experiments could not be completed because the expressed protein precipitated.

[0070] Genes of SA1606 (GenBank number: BAB42874) and SA2001 (GenBank number: BAB43292) were amplified using PCR with the genomic DNA of *S. aureus* N315 serving as a template. PCR primers were 5'-GCGCGCGCCATG-GAGGTT AAAACATTTTAT-3' and 5'-CGGCGCCTC-GAGTCCTTCAAAAGTTTTTGG-3' for SA1606, and 5'-GCGCGCGCCATGGATCATATTGAAATAAGT-3' and 5'-CGGCGCCTCGAGTTCTCTGCTTGTAAGTGC-3' for SA2001. NcoI and XhoI restriction sites were incorporated into the forward primer sequences and the reverse primer sequences, respectively. The PCR products were digested with NcoI and XhoI and inserted into NcoI/XhoI-digested pET28a vectors (Novagen). The DNAs thus ligated were transformed into *E. coli* BL21 DE3 (Invitrogen). Coding sequences of the cloned genes SA1606 and SA2001 were confirmed by nucleotide sequencing. Recombinant SA1606 and SA2001 with histidine tag at the C-terminus were expressed in *E. coli* and then loaded into Hi-Trap chelating columns (GE Healthcare) equilibrated with buffer A (20 mM HEPES, pH 7.5, 50 mM imidazole and 500 mM NaCl). The bound proteins were eluted with a linear gradient of 0.01-0.5 M imidazole in buffer A. Active fractions were concentrated and gel filtrated using a Superdex 200 column (GE Healthcare) in a buffer containing 20 mM HEPES (pH 7.5) and 2 mM DTT. The purified proteins are as shown in FIG. 7.

Example 8

Evaluation of Binding Affinity of FLF for Target Proteins by Isothermal Titration calorimetry

[0071] Binding affinity for SA1606 and SA2001 of FLF was measured using VP-ITC Micro calorimeter (MicroCal Inc). All of the proteins and FLF were prepared in 10 mM HEPES, pH 7.0 and 3% DMSO, followed by removing gas from the samples. In each titration experiment, 8 μ l of FLF was injected into 1.5 mL of sample cells containing the proteins 20~30 times at regular intervals of 4 min. Thereafter, the affinity of FLF for the proteins were calculated on the basis of the heat generated upon the binding of FLF to the proteins. The data was analyzed with ORIGIN software (MicroCal Inc).

[0072] As is apparent from the data of FIG. 8, dissociation constants (Kd) between FLF and SA1606 and between FLF and SA2001 were 14.5 μ M and 38.1 μ M, respectively, indi-

cating that FLF can practically bind to SA1606 and SA2001 and that the protein targets of FLF in *S. aureus* MRSA are SA1606 and SA2001.

Example 9

Enzymatic Assay for Inhibitory Activity of FLF Against Protein Targets

[0073] Aldo-keto reductase (AKR) activity was measured to examine whether FLF inhibits the enzymatic activities of SA1606 and SA2001 on the basis of the data of Example 8 showing the binding of FLF to SA1606 and SA2001. On the whole, bacterial AKR catalyzes the reduction of various chemicals with NADPH. Hence, the enzymatic activity of AKR can be determined by monitoring the absorbance of NADPH at 340 nm. As is evidenced in Example 6, SA1606 and SA2001 have structural and functional similarity with the human ald-keto reductase AKR1C3 and the bacterial ald-keto reductase DkgA. Thus, the enzymatic activity of DkgA was measured to analyze the inhibitory activity of FLF. The reaction for evaluating enzymatic activity was comprised of 0.5 mM NADPH, 5 mM methylglyoxal and 10 μ M ald-keto reductase in 50 mM sodium phosphate buffer, pH 7.0. Absorbance at 340 nm of the reaction was measured for 10 min to calculate the consumption of NADPH.

[0074] As can be seen in FIG. 9, SA1606 and SA2001 reduced methylglyoxal with NADPH (yellow lines), but the enzymatic activity was inhibited in the presence of FLF (brown lines), demonstrating that FLF is bound directly to the active sites of the enzymes SA1606 and SA2001, thus inhibiting the enzymatic activity of the *S. aureus* ald-keto reductase.

FORMULATION EXAMPLES

[0075] A better understanding of the present invention may be obtained through the following formulation examples of the antibiotically active, pharmaceutical or cosmetic composition comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt which are set forth to illustrate, but are not to be construed as the limit of the present invention.

EXAMPLES OF COSMETIC FORMULATION

Formulation Example 1

Skin Lotion (Content: wt %)

[0076] The composition: 0.001
[0077] Glycerin: 3.0
[0078] Butylene glycol: 2.0
[0079] Propylene glycol: 2.0
[0080] Carboxyvinyl polymer: 0.1
[0081] Ethanol: 10.0
[0082] Triethanolamine: 0.1
[0083] Preservative, pigment, flavorant, distilled water: q.s.
[0084] Total: 100.0

Formulation Example 2

Nutrient Cream (Content: wt %)

[0085] The composition: 0.001
[0086] Beeswax: 10.0
[0087] Polysorbate 60: 1.5

[0088] Sorbitan sesquiolate: 0.5
[0089] Liquid paraffin: 10.0
[0090] Squalane: 5.0
[0091] Caprylic/capric triglyceride: 5.0
[0092] Triethanolamine: 0.2
[0093] Preservative, pigment, flavorant, distilled water: q.s.
[0094] Total: 100.0

Formulation Example 3

Hair Soap (Content: wt %)

[0095] The composition: 0.001
[0096] Titanium dioxide: 0.2
[0097] Polyethylene glycole: 0.8
[0098] Glycerin: 0.5
[0099] Ethylene diaminetetraacetic acid: 0.05
[0100] Sodium: 1.0
[0101] Pigment: q.s.
[0102] Soap flavor: q.s.
[0103] Cosmetic soap base (water content 13, wt parts): q.s.
[0104] Total: 100.0

EXAMPLES OF PHARMACEUTICAL FORMULATION

Formulation Example 1

Ointment

[0105] The composition: 3 g
[0106] Dexpanthenol: 1.5 g
[0107] Stearic acid: 1.0 g
[0108] Liquid paraffin: 5.0 g
[0109] Spermaceti: 4.0 g
[0110] Cetanol: 3.0 g
[0111] Propylene glycol: 13.0 g
[0112] Triethanolamine: 1.5 g
[0113] Dibutylhydroxytoluene: 0.025 g
[0114] Ethyl benzoate: 0.0225 g
[0115] Propyl benzoate: 0.015 g
[0116] Polysorbate: 0.1 g
[0117] Purified water: 65.0 g
[0118] An ointment was prepared from these ingredients according to a conventional method.
[0119] The ointment was applied at a dose of 0.5~1 g twice a day constantly for 3~6 months to the lesion.

Formulation Example 2

Lotion

[0120] The composition: 1.0 g
[0121] Dexpanthenol: 1.5 g
[0122] Glycerin: 0.6 g
[0123] Hydroxypropylcellulose: 0.085 g
[0124] Beegum: 0.0255 g
[0125] Purified water: q.s.
[0126] A lotion was prepared from these ingredient according to a conventional method.
[0127] The lotion was applied at a dose of 0.1~1 mL twice a day constantly for 3~6 months to the lesion.

Formulation Example 3

Capsule

- [0128] The composition: 10 g
 [0129] Crystalline cellulose: 3 g
 [0130] Lactose: 14.8 g
 [0131] Magnesium stearate: 0.2 g
 [0132] These ingredients were mixed and loaded into gelatin capsules according to a conventional method to form capsules.

Formulation Example 4

Injection

- [0133] The composition: 10 g
 [0134] Mannitol: 180 g

- [0135] Sterile water for injection: 2974 g
 [0136] $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$: 26 g
 [0137] These ingredients were dissolved to form a total volume of 2 liters according to a conventional method.

Formulation Example 5

Liquid

- [0138] The composition: 20 mg
 [0139] Isomerase: 10 g
 [0140] Mannitol: 5 g
 [0141] Purified water: q.s.
 [0142] Using a conventional method, the ingredients were added to distilled water to which lemon flavor was then added in a suitable amount. After the ingredients were mixed, distilled water was added to bring the total volume to 100 mL, which was then loaded into a brown bottle and sterilized.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 8

<210> SEQ ID NO 1

<211> LENGTH: 279

<212> TYPE: PRT

<213> ORGANISM: *Staphylococcus aureus*

<400> SEQUENCE: 1

Met Leu Asn Glu Ile Gln Ile Leu Asn Asn Gly Tyr Pro Met Pro Ser
 1 5 10 15

Val Gly Leu Gly Val Tyr Lys Ile Ser Asp Glu Asp Met Thr Lys Val
 20 25 30

Val Asn Ala Ala Ile Asp Ala Gly Tyr Arg Ala Phe Asp Thr Ala Tyr
 35 40 45

Phe Tyr Asp Asn Glu Ala Ser Leu Gly Arg Ala Leu Lys Asp Asn Gly
 50 55 60

Val Asp Arg Glu Asp Leu Phe Ile Thr Thr Lys Leu Trp Asn Asp Tyr
 65 70 75 80

Gln Gly Tyr Glu Lys Thr Phe Glu Tyr Phe Asn Lys Ser Ile Glu Asn
 85 90 95

Leu Gln Thr Asp Tyr Leu Asp Leu Phe Leu Ile His Trp Pro Cys Glu
 100 105 110

Ala Asp Gly Leu Phe Leu Glu Thr Tyr Lys Ala Met Glu Glu Leu Tyr
 115 120 125

Glu Gln Gly Lys Val Lys Ala Ile Gly Val Cys Asn Phe Asn Val His
 130 135 140

His Leu Glu Lys Leu Met Ala Gln Ser Ser Ile Lys Pro Met Val Asn
 145 150 155 160

Gln Ile Glu Val His Pro Tyr Phe Asn Gln Gln Glu Leu Gln Glu Phe
 165 170 175

Cys Asp Arg His Asp Ile Lys Val Thr Ala Trp Met Pro Leu Met Arg
 180 185 190

Asn Arg Gly Leu Leu Asp Asn Pro Val Ile Val Lys Ile Ala Glu Lys
 195 200 205

Tyr His Lys Thr Pro Ala Gln Val Val Leu Arg Trp His Leu Ala His
 210 215 220

Asn Arg Ile Ile Ile Pro Lys Ser Gln Thr Pro Lys Arg Ile Gln Glu

-continued

```

225             230             235             240
Asn Ile Asp Ile Leu Asp Phe Asn Leu Glu Leu Thr Glu Val Ala Glu
             245             250             255
Ile Asp Ala Leu Asp Arg Asn Ala Arg Gln Gly Lys Asn Pro Asp Asp
             260             265             270
Val Lys Ile Gly Asp Leu Lys
             275

```

```

<210> SEQ ID NO 2
<211> LENGTH: 277
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus

```

```

<400> SEQUENCE: 2

```

```

Met Glu Val Lys Thr Phe Tyr Asn Gly Asn Thr Met Pro Gln Ile Gly
1             5             10             15
Leu Gly Thr Phe Arg Val Glu Asn Asp Glu Asn Cys Met Glu Ser Val
             20             25             30
Lys Tyr Ala Ile Glu Gln Gly Tyr Arg Ser Ile Asp Thr Ala Lys Val
             35             40             45
Tyr Gly Asn Glu Glu Gln Val Gly Ala Gly Ile Arg Ala Gly Leu Glu
             50             55             60
Ser Thr Gly Ile Ala Arg Glu Asp Leu Phe Ile Thr Ser Lys Leu Tyr
65             70             75             80
Phe Glu Asp Phe Gly Arg Glu Asn Val Ala Ala Ala Tyr Glu Ala Ser
             85             90             95
Leu Ser Arg Leu Gly Leu Lys Tyr Leu Asp Leu Tyr Leu Val His Trp
             100            105            110
Pro Gly Thr Asn Glu Ala Val Met Val Asp Thr Trp Lys Gly Met Glu
             115            120            125
Asp Leu Tyr Lys Asn Asn Lys Ala Lys Asn Ile Gly Val Ser Asn Phe
             130            135            140
Glu Pro Glu His Leu Glu Ala Leu Leu Ala Gln Val Ser Ile Lys Pro
145            150            155            160
Val Ile Asn Gln Val Glu Tyr His Pro Tyr Leu Thr Gln His Lys Leu
             165            170            175
Lys Leu Tyr Leu Ala Ala Gln His Ile Val Met Glu Ser Trp Ser Pro
             180            185            190
Leu Met Asn Ala Gln Ile Leu Asn Asp Glu Thr Ile Lys Asp Ile Ala
             195            200            205
Gln Glu Leu Gly Lys Ser Pro Ala Gln Val Val Leu Arg Trp Asn Val
             210            215            220
Gln His Gly Val Val Ile Ile Pro Lys Ser Val Thr Pro Asn Arg Ile
225            230            235            240
Ser Glu Asn Phe Gln Ile Phe Asp Phe Glu Leu Ser Asp Glu Gln Met
             245             250             255
Thr Arg Ile Asp Gly Leu Asn Gln Asp Lys Arg Ile Gly Pro Asp Pro
             260            265            270
Lys Thr Phe Glu Gly
             275

```

```

<210> SEQ ID NO 3
<211> LENGTH: 282
<212> TYPE: PRT

```

-continued

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 3

Met Asn His Ile Glu Ile Ser Lys Asp Val Lys Ile Pro Val Leu Gly
 1 5 10 15

Phe Gly Val Phe Gln Ile Pro Gln Glu Gln Thr Ala Glu Ala Val Lys
 20 25 30

Glu Ala Ile Lys Ala Gly Tyr Arg His Ile Asp Thr Ala Gln Ser Tyr
 35 40 45

Leu Asn Glu Thr Glu Val Gly Gln Gly Ile Glu Ala Ser Gly Ile Asp
 50 55 60

Arg Ser Glu Leu Phe Ile Thr Thr Lys Val Trp Ile Glu Asn Val Asn
 65 70 75 80

Tyr Glu Asp Thr Ile Lys Ser Ile Glu Arg Ser Leu Gln Arg Leu Asn
 85 90 95

Leu Asp Tyr Leu Asp Leu Val Leu Ile His Gln Pro Tyr Asn Asp Val
 100 105 110

Tyr Gly Ser Trp Arg Ala Leu Glu Glu Leu Lys Glu Asn Gly Lys Ile
 115 120 125

Lys Ala Ile Gly Val Ser Asn Phe Gly Val Asp Arg Ile Val Asp Leu
 130 135 140

Gly Ile His Asn Gln Ile Gln Pro Gln Val Asn Gln Ile Glu Ile Asn
 145 150 155 160

Pro Phe His Gln Gln Glu Glu Gln Val Ala Ala Leu Gln Gln Glu Asn
 165 170 175

Val Val Val Glu Ala Trp Ala Pro Phe Ala Glu Gly Lys Asn Gln Leu
 180 185 190

Phe Gln Asn Gln Leu Leu Gln Ala Ile Ala Asp Lys Tyr Asn Lys Ser
 195 200 205

Ile Ala Gln Val Ile Leu Arg Trp Leu Val Glu Arg Asp Ile Val Val
 210 215 220

Leu Ala Lys Ser Val Asn Pro Glu Arg Met Ala Gln Asn Leu Asp Ile
 225 230 235 240

Phe Asp Phe Glu Leu Thr Glu Glu Asp Lys Gln Gln Ile Ala Thr Leu
 245 250 255

Glu Glu Ser Asn Ser Gln Phe Phe Ser His Ala Asp Pro Glu Met Ile
 260 265 270

Lys Ala Leu Thr Ser Arg Glu Leu Asp Val
 275 280

<210> SEQ ID NO 4

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 4

gcgcgcgccca tggagggttaa aacattttat

30

<210> SEQ ID NO 5

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 5

cggcgcctcg agtccttcaa aagtttttgg 30

<210> SEQ ID NO 6

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 6

gcgcgcgccca tggatcatat tgaaataagt 30

<210> SEQ ID NO 7

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 7

cggcgcctcg agttctctgc ttgtaagtgc 30

<210> SEQ ID NO 8

<211> LENGTH: 331

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Met Asp Ser Lys Gln Gln Cys Val Lys Leu Asn Asp Gly His Phe Met
1 5 10 15

Pro Val Leu Gly Phe Gly Thr Tyr Ala Pro Pro Glu Val Pro Arg Ser
20 25 30

Lys Ala Leu Glu Val Thr Lys Leu Ala Ile Glu Ala Gly Phe Arg His
35 40 45

Ile Asp Ser Ala His Leu Tyr Asn Asn Glu Glu Gln Val Gly Leu Ala
50 55 60

Ile Arg Ser Lys Ile Ala Asp Gly Ser Val Lys Arg Glu Asp Ile Phe
65 70 75 80

Tyr Thr Ser Lys Leu Trp Ser Thr Phe His Arg Pro Glu Leu Val Arg
85 90 95

Pro Ala Leu Glu Asn Ser Leu Lys Lys Ala Gln Leu Asp Tyr Val Asp
100 105 110

Leu Tyr Leu Ile His Ser Pro Met Ser Leu Lys Pro Gly Glu Glu Leu
115 120 125

Ser Pro Thr Asp Glu Asn Gly Lys Val Ile Phe Asp Ile Val Asp Leu
130 135 140

Cys Thr Thr Trp Glu Ala Met Glu Lys Cys Lys Asp Ala Gly Leu Ala
145 150 155 160

Lys Ser Ile Gly Val Ser Asn Phe Asn Arg Arg Gln Leu Glu Met Ile
165 170 175

Leu Asn Lys Pro Gly Leu Lys Tyr Lys Pro Val Cys Asn Gln Val Glu
180 185 190

-continued

Cys	His	Pro	Tyr	Phe	Asn	Arg	Ser	Lys	Leu	Leu	Asp	Phe	Cys	Lys	Ser
	195						200					205			
Lys	Asp	Ile	Val	Leu	Val	Ala	Tyr	Ser	Ala	Leu	Gly	Ser	Gln	Arg	Asp
	210					215					220				
Lys	Arg	Trp	Val	Asp	Pro	Asn	Ser	Pro	Val	Leu	Leu	Glu	Asp	Pro	Val
	225				230					235				240	
Leu	Cys	Ala	Leu	Ala	Lys	Lys	His	Lys	Arg	Thr	Pro	Ala	Leu	Ile	Ala
			245						250					255	
Leu	Arg	Tyr	Gln	Leu	Gln	Arg	Gly	Val	Val	Val	Leu	Ala	Lys	Ser	Tyr
			260					265						270	
Asn	Glu	Gln	Arg	Ile	Arg	Gln	Asn	Val	Gln	Val	Phe	Glu	Phe	Gln	Leu
		275					280						285		
Thr	Ala	Glu	Asp	Met	Lys	Ala	Ile	Asp	Gly	Leu	Asp	Arg	Asn	Leu	His
	290					295					300				
Tyr	Phe	Asn	Ser	Asp	Ser	Phe	Ala	Ser	His	Pro	Asn	Tyr	Pro	Tyr	Ser
	305					310				315					320
Asp	Glu	Tyr	Leu	Glu	His	His	His	His	His	His	His				
			325							330					

1. An antibiotically active composition, comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt as an active ingredient.

2. The antibiotically active composition of claim 1, functioning to destroy or inhibit growth or metabolism of Gram-positive bacteria.

3. The antibiotically active composition of claim 2, wherein the Gram-positive bacteria is selected from the group consisting of *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Bacillus subtilis*, *Listeria monocytogenes*, and *Enterococcus faecium*.

4. The antibiotically active composition of claim 2, wherein the active ingredient targets Gram-positive bacterial aldo-keto reductase.

5. The antibiotically active composition of claim 4, wherein the reductase is selected from the group consisting of SA0658, SA1606, and SA2001, which are *Staphylococcal* aldo-keto reductases.

6. The antibiotically active composition of claim 1, functioning to destroy or inhibit growth or metabolism of antibiotic-resistant bacteria.

7. The antibiotically active composition of claim 6, wherein the antibiotic-resistant bacteria is selected from the group consisting of MRSA (Methicillin-Resistant *Staphylococcus aureus*), VRSA (Vancomycin-Resistant *Staphylococcus aureus*), MRE (multiple-drug resistant enterococci), and VRE (Vancomycin-Resistant enterococci).

8. An aldo-keto reductase, present in Gram-positive bacteria, as a target protein of an antibiotic functioning to destroy or inhibit growth or metabolism of Gram-positive bacteria.

9. The aldo-keto reductase of claim 8, wherein the Gram-positive bacteria is *Staphylococcus aureus*.

10. The aldo-keto reductase of claim 9, having an amino acid sequence of SEQ ID NO: 1 (SA0658), serving as a target

of an antibiotic destructive or inhibitory of growth or metabolism of *Staphylococcus aureus*.

11. The aldo-keto reductase of claim 9, having an amino acid sequence of SEQ ID NO: 2 (SA1606), serving as a target of an antibiotic destructive or inhibitory of growth or metabolism of *Staphylococcus aureus*.

12. The aldo-keto reductase of claim 9, having an amino acid sequence of SEQ ID NO: 3 (SA2001), serving as a target of an antibiotic destructive or inhibitory of growth or metabolism of *Staphylococcus aureus*.

13. An antibiotically active, pharmaceutical composition, comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt as an active ingredient.

14. The antibiotically active, pharmaceutical composition of claim 13, further comprising an antibiotic.

15. The antibiotically active, pharmaceutical composition of claim 14, wherein the antibiotic is selected from the group consisting of norfloxacin, oxacillin, and cephalothin.

16. An antibiotically active, cosmetic composition, comprising flufenamic acid, flufenamic acid derivative, flufenamic acid isomer, or flufenamic acid salt as an active ingredient.

17. The antibiotically active, cosmetic composition of claim 16, being in a form selected from the group consisting of hand wash, face wash, cleansing agent, shampoo, rinse, soap, lotion, essence, cream, wax, nutrient pack, a dyeing agent, a waving agent, and spray.

18. The antibiotically active, cosmetic composition of claim 16, further comprising an antibiotic.

19. The antibiotically active, cosmetic composition of claim 18, wherein the antibiotic is selected from the group consisting of norfloxacin, oxacillin, and cephalothin.

* * * * *