

(19)



(11)

EP 1 991 526 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

12.02.2014 Bulletin 2014/07

(51) Int Cl.:

C07D 209/80 <small>(2006.01)</small>	C07D 495/04 <small>(2006.01)</small>
A61K 31/404 <small>(2006.01)</small>	A61K 31/407 <small>(2006.01)</small>
A61P 25/00 <small>(2006.01)</small>	A61P 35/00 <small>(2006.01)</small>
A61P 11/00 <small>(2006.01)</small>	A61K 51/04 <small>(2006.01)</small>

(21) Application number: **06808621.4**

(22) Date of filing: **20.11.2006**

(86) International application number:

PCT/GB2006/004342

(87) International publication number:

WO 2007/057705 (24.05.2007 Gazette 2007/21)

(54) **Tetracyclic indole derivatives as in vivo imaging agents and having peripheralbenzodiazepine receptor affinity (PBR)**

Tetracyclische Indolderivate als Mittel zur In-Vivo-Bilddarstellung und mit Affinität zum Peripheralbenzodiazepinrezeptor (PBR)

Dérivés d'indole tetracyclique en tant qu' agents de formation d' images in vivo et ayant une affinité pour les récepteurs de peripheralbenzodiazepine

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**

• **ROBINS, Edward, George**

London W12 0NN (GB)

• **SHAN, Bo**

London W12 0NN (GB)

(30) Priority: **18.11.2005 GB 0523506**

(43) Date of publication of application:

19.11.2008 Bulletin 2008/47

(73) Proprietor: **Hammersmith Imanet, Ltd**

London W12 0NN (GB)

(74) Representative: **Canning, Lewis R. et al**

GE Healthcare Limited

Amersham Place

Little Chalfont

Buckinghamshire HP7 9NA (GB)

(72) Inventors:

- **ARSTAD, Erik**
London W12 0NN (GB)
- **WILSON, Ian**
Buckinghamshire HP7 9LL (GB)
- **LUTHRA, Sajinder, Kaur**
London W12 0NN (GB)
- **BRADY, Frank**
London SW8 2AB (GB)
- **LANGSTRÖM, Bengt**
S-751 85 Uppsala (SE)
- **KARIMI, Farhad**
S-751 85 Uppsala (SE)

(56) References cited:

FR-A1- 2 788 776 JP-A- 7 165 721

- **T-OKUBO ET AL: "Design, synthesis, and structure-activity relationships of novel tetracyclic compounds as peripheral benzodiazepine receptor ligands" BIOORGANIC & MEDICINAL CHEMISTRY, vol. 12, 2004, pages 3569-3580, XP002416574 cited in the application**

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 991 526 B1

Description**Technical Field of the Invention**

5 [0001] The present invention relates to compounds having a high affinity for peripheral benzodiazepine receptors (PBR). Novel compounds are provided which find use in both diagnosis and therapy. In particular, the compounds of the invention are useful for *in vivo* imaging and for treatment of disease states in which expression of PBR is upregulated.

Description of Related Art

10

[0002] PBR are known to be mainly localised in peripheral tissues and glial cells but their physiological function remains to be clearly elucidated. Subcellularly, PBR are known to localise on the outer mitochondrial membrane. Their presence on the outer membrane of mitochondria indicates a potential role in the modulation of mitochondrial function and in the immune system. It has furthermore been postulated that PBR are involved in cell proliferation, steroidogenesis, calcium

15 flow and cellular respiration. Altered expression of PBR has been observed in a variety of conditions including acute and chronic stress, anxiety, depression, Parkinson's disease, Alzheimer's disease, brain damage, cancer [Gavish et al 1999 Pharm. Rev. 51 p629], Huntington's disease [Neurosci. Lett. 1998 24(1) pp53-6], asthma [Gen. Pharmacol. 1997 28(4) pp495-8], rheumatoid arthritis [Eur. J. Pharmacol. 2002 452(1) pp111-22], atherosclerosis [J. Nucl. Med. 2004 45 pp1898-1907] and multiple sclerosis [Banati et al 2000 Brain 123 p2321]. PBR may also be associated with neuropathic pain, Tsuda et al having observed activated microglia in subjects with neuropathic pain [2005 TINS 28(2) pp101-7].

20

[0003] Ligands having affinity for PBR are known in the art. A class of indole compounds having affinity for PBR [IC_{50} values for most active compounds of between 0.2nM and 5.0nM] is disclosed in US 6451795. The patent states that the compounds are useful for the prevention or treatment of peripheral neuropathies and for the treatment of central neurodegenerative diseases. Okubu et al [Bioorganic & Medicinal Chemistry 2004 12 3569-80] describe the design, synthesis and structure of a group of tetracyclic indole compounds having affinity for PBR [IC_{50} values as low as about 0.4nM]. No particular application of the compounds is discussed. Campiani et al [2002 J. Med. Chem. 45 4276-81] disclose a class of pyrrolobenzoxazepine derivatives that bind to PBR with high affinity [K_i against PK11195 as low as about 0.1nM].

25

[0004] Positron emission tomography [PET] imaging using the PBR selective ligand, (R)-[^{11}C]PK11195 provides a generic indicator of central nervous system (CNS) inflammation. Despite the successful use of (R)-[^{11}C]PK11195, it has its limitations. It is known to have high protein binding, and low specific to non-specific binding. The role of its radiolabelled metabolites is not known and quantification of binding requires complex modelling.

30

[0005] Radioiodinated and radiobrominated isoquinoline carboxamide derivatives having high affinity for PBR are disclosed in JP 07165721 as novel *in vivo* imaging agents or radiotherapeutic agents for PBR.

[0006] There is therefore scope for alternative *in vivo* imaging agents and therapeutic agents that target PBR.

35

Summary of the Invention

[0007] The present invention provides novel tetracyclic indole compounds suitable for use either as *in vivo* imaging agents or as therapeutic agents. A method for the preparation of the *in vivo* imaging agent compound is also provided by the invention, as well as a precursor for use in said method. Pharmaceutical compositions comprising the compounds of the invention are additionally provided. Where the pharmaceutical composition comprises a compound suitable for *in vivo* imaging, a kit is provided for the preparation of the pharmaceutical composition. In a further aspect, use of the compound for *in vivo* imaging or treatment of conditions associated with PBR is provided.

40


45

Detailed Description of the Invention

[0008] In one aspect, the present invention provides a compound of Formula I:

50

55



or a salt or solvate thereof, wherein said compound is labelled with an imaging moiety, and wherein:

X¹ and X² are independently selected from hydrogen, halogen, C₁₋₆ alkoxy, C₁₋₆ alkyl;

R¹ and R² are independently selected from C₁₋₆ alkyl, C₁₋₆ aminoalkyl, C₁₋₆ alkoxyalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, a polyethylene glycol (PEG) group, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloethers, and C₃₋₁₀ cycloamines;

R³ is the group -A-R⁵ wherein:

A is the optional group -(CH₂)_z-R⁶- wherein z=0-6, R⁶ is a 5- or 6-membered heterocycle having 1-3 heteroatoms selected from N, S and O, and wherein R⁵ is a substituent selected from hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ alkanoyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkenyl, C₁₋₆ haloalkynyl, C₁₋₆ haloalkoxy, C₁₋₆ haloalkylthiol, C₁₋₆ haloalkylsulfinyl, C₁₋₆ haloalkylsulfonyl, C₁₋₆ haloalkyl ketone, C₁₋₆ haloalkyl sulfinyl, C₁₋₆ haloalkyl sulfonyl, a polyethylene glycol (PEG) group, C₁₋₆ hydroxyalkyl, a nitrogen-containing C₂₋₁₀ alkyl and hydroxy;

R⁴ is hydrogen, C₁₋₆ alkyl, C₁₋₆ alkanoyl, C₁₋₆ cycloalkyl, C₁₋₆ fluoroalkyl, hydroxy, or halogen;

Y¹ is S, SO, SO₂, or CH₂; and,

n is 0 to 10

[0009] "Alkyl", alone or in combination, means a straight-chain or branched-chain alkyl radical. Suitable alkyls include, but are not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl.

[0010] As used herein, the term "halogen" or "halo" includes iodine, fluorine, which are preferred, and chlorine and bromine.

[0011] "Heterocycle" means an aliphatic or aromatic saturated or partially unsaturated monocycle. Suitable heterocycles include, but are not limited to, furan, pyrrole, imidazole, triazole, thiophene, pyridine, piperidine, pyran, and piperazine.

[0012] Where a substituent is a PEG group, the group suitably comprises between 5 and 20 ethylene glycol units and preferably between 5 and 10 ethylene glycol units.

[0013] For the compounds of Formula I defined above, there are potentially a number of chiral centres. The present invention therefore encompasses racemic and optically pure forms of any compounds of the invention with one chiral centre, as well as racemic, diastereomer and optically pure forms of any compounds of the invention with two chiral centres. Optically pure forms of these compounds are preferred.

[0014] Suitable salts according to the invention, include physiologically acceptable acid addition salts such as those derived from mineral acids, for example hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric and sulphuric acids, and those derived from organic acids, for example tartaric, trifluoroacetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic, methanesulphonic, and paratoluenesulphonic acids.

[0015] Suitable solvents according to the invention include ethanol, water, saline, physiological buffer and glycol.

[0016] The term "labelled with an imaging moiety" means either (i) that one of the atoms of the compound of Formula I itself is an imaging moiety, or (ii) that a group comprising an imaging moiety is conjugated to the compound of Formula I. Where one of the atoms of the compound of Formula I itself is an imaging moiety, for example any of the carbons could be ¹¹C, or any F could be ¹⁸F. Where a group comprising an imaging moiety is conjugated to the compound of Formula I, it may be incorporated directly via any suitable atom present in the compound. Examples of groups comprising an imaging moiety include metal complexes which comprise a metallic imaging moiety, or radioiodophenyl. Alternatively,

where a group comprising an imaging moiety is conjugated to the compound of Formula I, it may be incorporated indirectly via a linker group of Formula $-(L^1)_m-$ wherein:

5 each L is independently $-\text{CO}-$, $-\text{CR}_2-$, $-\text{CR}=\text{CR}-$, $-\text{C}\equiv\text{C}-$, $-\text{CR}_2\text{CO}_2-$, $-\text{CO}_2\text{CR}_2-$, $-\text{NR}-$, $-\text{NRCO}-$, $-\text{CONR}-$, $\text{NR}(\text{C}=\text{O})\text{NR}-$, $-\text{NR}(\text{C}=\text{S})\text{NR}-$, $-\text{SO}_2\text{NR}-$, $-\text{NRSO}_2-$, $-\text{CR}_2\text{OCR}_2-$, $-\text{CR}_2\text{SCR}_2-$, $-\text{CR}_2\text{NRCR}_2-$, a C_{4-8} cycloheteroalkylene group, a C_{4-8} cycloalkylene group, a C_{5-12} arylene group, a C_{3-12} heteroarylene group, an amino acid residue, a polyalkyleneglycol, polylactic acid or polyglycolic acid moiety;

10 m is an integer of value 1 to 5;

each R group is independently H or C_{1-6} alkyl, C_{3-10} alkylaryl, C_{2-6} alkoxyalkyl, C_{1-6} hydroxyalkyl, C_{1-6} fluoroalkyl, or 2 or more R groups, together with the atoms to which they are attached form a carbocyclic, heterocyclic, saturated or unsaturated ring.

15 **[0017]** More detail relating to imaging moieties and groups comprising imaging moieties is provided below.

[0018] The following is a recitation of the preferred substituents for Formula I:

X^1 and X^2 are preferably both hydrogen.

20 R^1 and R^2 are preferably independently selected from C_{1-6} alkyl, C_{1-6} methoxyalkyl or C_{1-6} alkoxy, are most preferably both C_{1-6} alkyl, and are most especially preferably both ethyl.

25 R^3 is preferably hydrogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkenyl, C_{1-6} alkynyl, C_{1-6} alkanoyl, C_{1-6} fluoroalkyl, C_{1-6} fluoroalkoxy, or is A- C_{1-6} alkyl or A- C_{1-6} fluoroalkyl wherein A is as defined previously, most preferably hydrogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkenyl, C_{1-6} alkynyl, acetyl, C_{1-6} fluoroalkyl, or is A- C_{1-6} alkyl or A- C_{1-6} fluoroalkyl wherein for A, R^6 is a 5- or 6-membered N-containing heterocycle, and most especially preferably C_{1-3} alkyl, C_{1-6} alkoxy, C_{1-3} alkenyl, C_{1-3} alkynyl, acetyl or C_{1-6} fluoroalkyl, or is A- C_{1-6} fluoroalkyl wherein for A, R^6 is a 5-membered N-containing heterocycle.

30 R^4 is preferably hydrogen, C_{1-6} alkyl, C_{1-6} alkanoyl, C_{1-6} cycloalkyl or C_{1-6} fluoroalkyl, most preferably hydrogen, C_{1-6} alkyl or C_{1-6} alkanoyl and most especially preferably hydrogen or C_{1-3} alkyl.

Y^1 is preferably S, SO_2 or CH_2 , most preferably S or SO_2 , and most especially preferably S.

35 n is preferably 0 to 5 and most preferably 0.

[0019] For preferred compounds of Formula I of the invention:

40 X^1 and X^2 are both hydrogen;

R^1 and R^2 are independently selected from C_{1-6} alkyl, C_{1-6} methoxyalkyl or C_{1-6} alkoxy;

45 R^3 is hydrogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkenyl, C_{1-6} alkynyl, or C_{1-6} fluoroalkyl, or is A- C_{1-6} alkyl, or A- C_{1-6} fluoroalkyl wherein A is as defined previously;

R^4 is hydrogen, C_{1-6} alkyl, C_{1-6} acetyl, C_{1-6} cycloalkyl or C_{1-6} fluoroalkyl;

Y^1 is S, SO_2 , or CH_2 ; and,

50 n is 0

[0020] For most preferred compounds of Formula I of the invention:

55 X^1 and X^2 are both independently hydrogen;

R^1 and R^2 are both C_{1-6} alkyl;

R^3 is hydrogen, C_{1-3} alkyl, C_{1-3} alkoxy, C_{1-3} alkanoyl, C_{1-3} alkenyl, C_{1-3} alkynyl, or C_{1-6} fluoroalkyl, or is A- C_{1-6} alkyl

or A-C₁₋₆ fluoroalkyl wherein for A, R⁶ is a 5- or 6-membered N-containing heterocycle;

R⁴ is hydrogen, C₁₋₆ alkyl or C₁₋₆ acetyl;

5 Y¹ is S or SO₂; and,

n is 0

[0021] For most especially preferred compounds of Formula I of the invention:

10

X¹ and X² are both hydrogen;

R¹ and R² are both ethyl;

15

R³ is hydrogen, methyl, ethyl, 2-methoxy-ethyl, prop-2-ynyl, isopropyl, isobutyl, 2-methyl-allyl, acetyl, or 4-fluorobutyl, or is or is A-C₁₋₆ fluoroalkyl wherein for A, R⁶ is a 5-membered N-containing heterocycle;

R⁴ is hydrogen;

20

Y¹ is S; and,

n is 0

[0022] An alternative embodiment of this aspect of the invention is a compound of Formula I or a salt or solvate thereof, wherein said compound is labelled with an imaging moiety, and wherein:

25

X¹ and are independently selected from hydrogen, halogen, C₁₋₆ alkoxy, C₁₋₆ alkyl;

30

R¹ and R² are independently selected from C₁₋₆ alkyl, C₁₋₆ aminoalkyl, C₁₋₆ methoxyalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl alcohols, a polyethylene glycol (PEG) group, C₃₋₁₀ cycloalkanes, C₃₋₁₀ cycloethers, and C₃₋₁₀ cycloamines;

R³ is an optional substituent selected from hydrogen, C₁₋₆ alkyl, C₁₋₆ acetyl, C₁₋₆ fluoroalkyl, C₁₋₆ fluoroalkoxy, C₁₋₆ fluoroalkylsulfinyl, C₁₋₆ fluoroalkylsulfonyl, trifluoromethyl ketone, trifluoromethyl sulfinyl, trifluoromethyl sulfonyl, a polyethylene glycol (PEG) group, C₁₋₆ alkyl alcohols and hydroxy;

35

R⁴ is hydrogen, C₁₋₆ alkyl, C₁₋₆ acetyl, C₁₋₆ cycloalkyl, C₁₋₆ fluoroalkyl, hydroxy, or halogen;

Y¹ is S, SO, SO₂, or CH₂; and,

40

n is 0 to 10


[0023] The synthesis of compounds of Formula I that are not labelled with an imaging moiety may be carried out by methods described by Okubo et al [Bioorganic and Medicinal Chemistry 2004 12 3569-80].


[0024] Preferred sites for incorporation of an imaging moiety into a compound of Formula I and in the more specific aspects of the invention are any of R¹-R⁴ and either of X¹ or X², as in compounds of Formulas li-lvi:

45

50

55






wherein IM is an imaging moiety or a group comprising an imaging moiety and R¹-R⁴, X¹, X² and Y¹ are as defined above for Formula I.


[0025] Most preferred sites for the incorporation of an imaging moiety into a compound of Formula I are R¹⁻³, and especially preferably R³.

[0026] For ¹¹C, a further preferred site of incorporation is at the carbonyl group bound to NR¹R², as in compounds of Formula Ivii:



wherein R¹-R⁴, X¹, X² and Y¹ are as defined above for Formula I.

[0027] For radioiodine, a further preferred site of incorporation is at phenyl ring A' of Formula Iviii:



[0028] Methods for incorporation of the preferred imaging moieties of the invention into a compound of Formula I are discussed below in relation to a further aspect of the invention.

[0029] The "imaging moiety" allows the compound of the invention to be detected using a suitable imaging modality following its administration to a mammalian body *in vivo* and are chosen from:

- (i) a radioactive metal ion;
- (ii) a paramagnetic metal ion;
- (iii) a gamma-emitting radioactive halogen;

(iv) a positron-emitting radioactive non-metal;

(v) a hyperpolarised NMR-active nucleus;

5 (vi) a reporter suitable for *in vivo* optical imaging;

(vii) a β -emitter suitable for intravascular detection.

10 **[0030]** When the imaging moiety is a radioactive metal ion, i.e. a radiometal, suitable radiometals can be either positron emitters such as ^{64}Cu , ^{48}V , ^{52}Fe , ^{55}Co , $^{94\text{m}}\text{Tc}$ or ^{68}Ga ; γ -emitters such as $^{99\text{m}}\text{Tc}$, ^{111}In , $^{113\text{m}}\text{In}$, or ^{67}Ga . Preferred radiometals are $^{99\text{m}}\text{Tc}$, ^{64}Cu , ^{68}Ga and ^{111}In . Most preferred radiometals are γ -emitters, especially $^{99\text{m}}\text{Tc}$.

[0031] When the imaging moiety is a paramagnetic metal ion, suitable such metal ions include: Gd(III) , Mn(II) , Cu(II) , Cr(III) , Fe(III) , Co(II) , Er(II) , Ni(II) , Eu(III) or Dy(III) . Preferred paramagnetic metal ions are Gd(III) , Mn(II) and Fe(III) , with Gd(III) being especially preferred.

15 **[0032]** When the imaging moiety is a gamma-emitting radioactive halogen, the radiohalogen is suitably chosen from ^{123}I , ^{131}I or ^{77}Br . A preferred gamma-emitting radioactive halogen is ^{123}I .

[0033] When the imaging moiety is a positron-emitting radioactive non-metal, suitable such positron emitters include: ^{11}C , ^{13}N , ^{15}O , ^{17}F , ^{18}F , ^{75}Br , ^{76}Br or ^{124}I . Preferred positron-emitting radioactive non-metals are ^{11}C , ^{13}N , ^{18}F and ^{124}I , especially ^{11}C and ^{18}F , most especially ^{18}F .

20 **[0034]** When the imaging moiety is a hyperpolarised NMR-active nucleus, such NMR-active nuclei have a non-zero nuclear spin, and include ^{13}C , ^{15}N , ^{19}F , ^{29}Si and ^{31}P . Of these, ^{13}C is preferred. By the term "hyperpolarised" is meant enhancement of the degree of polarisation of the NMR-active nucleus over its' equilibrium polarisation. The natural abundance of ^{13}C (relative to ^{12}C) is about 1%, and suitable ^{13}C -labelled compounds are suitably enriched to an abundance of at least 5%, preferably at least 50%, most preferably at least 90% before being hyperpolarised.

25 **[0035]** When the imaging moiety is a reporter suitable for *in vivo* optical imaging, the reporter is any moiety capable of detection either directly or indirectly in an optical imaging procedure. The reporter might be a light scatterer (e.g. a coloured or uncoloured particle), a light absorber or a light emitter. More preferably the reporter is a dye such as a chromophore or a fluorescent compound. The dye can be any dye that interacts with light in the electromagnetic spectrum with wavelengths from the ultraviolet light to the near infrared. Most preferably the reporter has fluorescent properties. Preferred organic chromophoric and fluorophoric reporters include groups having an extensive delocalized electron system, e.g. cyanines, merocyanines, indocyanines, phthalocyanines, naphthalocyanines, triphenylmethines, porphyrins, pyrilium dyes, thiapyrilium dyes, squarylium dyes, croconium dyes, azulenium dyes, indoanilines, benzophenoxazinium dyes, benzothiaphenothiazinium dyes, anthraquinones, naphthoquinones, indathrenes, phthaloylacridones, tri-sphenoquinones, azo dyes, intramolecular and intermolecular charge-transfer dyes and dye complexes, tropones, tetrazines, bis(dithiolene) complexes, bis(benzene-dithiolate) complexes, iodoaniline dyes, bis(S,O-dithiolene) complexes. Fluorescent proteins, such as green fluorescent protein (GFP) and modifications of GFP that have different absorption/emission properties are also useful. Complexes of certain rare earth metals (e.g., europium, samarium, terbium or dysprosium) are used in certain contexts, as are fluorescent nanocrystals (quantum dots).

30 **[0036]** Particular examples of chromophores which may be used include: fluorescein, sulforhodamine 101 (Texas Red), rhodamine B, rhodamine 6G, rhodamine 19, indocyanine green, Cy2, Cy3, Cy3.5, Cy5, Cy5.5, Cy7, Marina Blue, Pacific Blue, Oregon Green 88, Oregon Green 514, tetramethylrhodamine, and Alexa Fluor 350, Alexa Fluor 430, Alexa Fluor 532, Alexa Fluor 546, Alexa Fluor 555, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 633, Alexa Fluor 647, Alexa Fluor 660, Alexa Fluor 680, Alexa Fluor 700, and Alexa Fluor 750.

45 **[0037]** When the imaging moiety is a β -emitter suitable for intravascular detection, suitable such β -emitters include the radiometals ^{67}Cu , ^{89}Sr , ^{90}Y , ^{153}Sm , ^{186}Re , ^{188}Re or ^{192}Ir , and the non-metals ^{32}P , ^{33}P , ^{38}S , ^{38}Cl , ^{39}Cl , ^{82}Br and ^{83}Br .


[0038] The most preferred imaging moieties of the invention are radioactive, especially radioactive metal ions, gamma-emitting radioactive halogens and positron-emitting radioactive non-metals, particularly those suitable for imaging using SPECT or PET. Most especially preferred imaging moieties of the invention are suitable for imaging using PET, i.e. ^{11}C and ^{18}F .

50 **[0039]** Preferably, compounds of the invention have K_i values for binding to PBR (determined by method of Le Fur et al 1983 Life Sci. USA 33: 449-57) of between 0.01nM and 10nM, most preferably between 0.1nM and 5.0nM and most especially preferably between 0.1nM and 1.0nM.

[0040] Examples of most preferred compounds of Formula I labeled with an imaging moiety, are as follows:


55

5




Compound 1

10




Compound 2




Compound 3

15




Compound 4

20




Compound 5

25




Compound 6

30




Compound 7

35



Compound 8

40




Compound 9

45

50

55




[0041] K_i values for the best compounds, tested using the method of Example 18 below, were found to be between 1.0nM and 0.1nM.

[0042] Preferably, compounds of the invention do not undergo facile metabolism *in vivo*, and hence most preferably exhibit a half-life *in vivo* of 60 to 240 minutes in humans. The compound is preferably excreted *via* the kidney (i.e. exhibits urinary excretion). The compound preferably exhibits a signal-to-background ratio at diseased foci of at least 1.5, most preferably at least 5, with at least 10 being especially preferred. Where the compound comprises a radioisotope, clearance of one half of the peak level of compound which is either non-specifically bound or free *in vivo*, preferably occurs over a time period less than or equal to the radioactive decay half-life of the radioisotope of the imaging moiety.

[0043] The molecular weight of the compound is preferably up to 5000 Daltons. Most preferably, the molecular weight is in the range 150 to 3000 Daltons, most especially preferably 200 to 1500 Daltons, with 300 to 800 Daltons being ideal.

[0044] In a further aspect, the present invention provides a compound of Formula II:



or a salt or solvate thereof wherein:

R^7 is as defined previously for R^3 , with the proviso that R^7 is not hydrogen, C_{1-5} alkyl or a nitrogen-containing C_{2-10} alkyl group; and,

Y^2 is as defined for Y^1 above.

[0045] The synthesis of compounds of Formula II may be carried out by adapting the methods described by Okubo et al [Bioorganic and Medicinal Chemistry 2004 12 3569-80].

[0046] Examples of preferred compounds of Formula II are as follows:

5

10

15

20

25

30


35

40

45


50

55



[0047] K_i values for the best compounds, tested using the method of Example 18 below, were found to be between 1.0nM and 0.1nM.

[0048] In another further aspect, the present invention provides a method for the preparation of a compound of Formula I labelled with an imaging moiety comprising reaction of a convenient chemical form of an imaging moiety with a precursor of Formula Ia:



wherein:

X^{11} , X^{12} , R^{13} , R^{14} and Y^{11} are as defined for X^1 , X^2 , R^3 , R^4 and Y^{11} , respectively, of Formula I above, or independently

comprise a suitable protecting group;

R¹⁵ is the group -(CH₂)_o-C(=O)-NR¹¹R¹² wherein o, R¹¹ and R¹² are as defined for n, R¹ and R² respectively for Formula I, or comprises a suitable protecting group; and,

R¹⁶ is hydrogen,

and provided that at least one of X¹¹, X¹² and R¹¹-R¹⁶ comprises a chemical group capable of reacting with a suitable source of said imaging moiety wherein:

(i) when said imaging moiety is a metal ion, said chemical group is a chelating agent where 2-6 donor atoms selected from amine, thiol, amide, oxime and phosphine are arranged such that 5- or 6-membered chelate rings result, or is a monodentate ligand comprising donor atoms selected from isonitrile, phosphine and diazenide;

(ii) when said imaging moiety is radioiodine, said chemical group is a trialkylstannane, a trialkylsilane, an organoboron compound, alkyl bromide, alkyl tosylate, alkyl mesylate, alkyl triflate, phenol, aryl iodonium salt, aryl diazonium, aryl trialkylammonium or nitroaryl;

(iii) when said imaging moiety is a radioactive isotope of fluorine, said chemical group is alkyl bromide, alkyl tosylate, alkyl mesylate,

(iv) hydroxyl, N-haloacetyl, aryl diazonium salt, aryl nitro, aryl quaternary ammonium salt, chloro, P(O)Ph₃ or an activated ester;

wherein said imaging moiety is ¹¹C, said chemical group is hydrogen, halide, boronic acid, triflate or organotin.

[0049] A "precursor" comprises a derivative of the compound of Formula I, designed so that chemical reaction with a convenient chemical form of the imaging moiety occurs site-specifically; can be conducted in the minimum number of steps (ideally a single step); and without the need for significant purification (ideally no further purification), to give the desired imaging agent. Such precursors are synthetic and can conveniently be obtained in good chemical purity. The "precursor" may optionally comprise a protecting group for certain functional groups of the compound of Formula Ia.

[0050] By the term "protecting group" is meant a group which inhibits or suppresses undesirable chemical reactions, but which is designed to be sufficiently reactive that it may be cleaved from the functional group in question under mild enough conditions that do not modify the rest of the molecule. After deprotection the desired product is obtained. Protecting groups are well known to those skilled in the art and are suitably chosen from, for amine groups: Boc (where Boc is *tert*-butoxycarbonyl), Fmoc (where Fmoc is fluorenylmethoxycarbonyl), trifluoroacetyl, allyloxycarbonyl, Dde [i.e. 1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl] or Npys (i.e. 3-nitro-2-pyridine sulfonyl); and for carboxyl groups: methyl ester, *tert*-butyl ester or benzyl ester. For hydroxyl groups, suitable protecting groups are: methyl, ethyl or *tert*-butyl; alkoxymethyl or alkoxyethyl; benzyl; acetyl; benzoyl; trityl (Trt) or trialkylsilyl such as tetrabutyltrimethylsilyl. For thiol groups, suitable protecting groups are: trityl and 4-methoxybenzyl. The use of further protecting groups are described in 'Protective Groups in Organic Synthesis', Theodor W. Greene and Peter G. M. Wuts, (Third Edition, John Wiley & Sons, 1999).


[0051] When the imaging moiety comprises a metal ion, the precursor is derivatised to include a chemical group capable of complexing the metal ion to form a metal complex. By the term "metal complex" is meant a coordination complex of the metal ion with one or more ligands. It is strongly preferred that the metal complex is "resistant to transchelation", i.e. does not readily undergo ligand exchange with other potentially competing ligands for the metal coordination sites. Potentially competing ligands include the compound of Formula I itself plus other excipients in the preparation *in vitro* (e.g. radioprotectants or antimicrobial preservatives used in the preparation), or endogenous compounds *in vivo* (e.g. glutathione, transferrin or plasma proteins).

[0052] Suitable ligands for use in the present invention which form metal complexes resistant to transchelation include: chelating agents, where 2-6, preferably 2-4, metal donor atoms are arranged such that 5- or 6-membered chelate rings result (by having a non-coordinating backbone of either carbon atoms or non-coordinating heteroatoms linking the metal donor atoms); or monodentate ligands which comprise donor atoms which bind strongly to the metal ion, such as isonitriles, phosphines or diazenides. Examples of donor atom types which bind well to metals as part of chelating agents are: amines, thiols, amides, oximes and phosphines. Phosphines form such strong metal complexes that even monodentate or bidentate phosphines form suitable metal complexes. The linear geometry of isonitriles and diazenides is such that they do not lend themselves readily to incorporation into chelating agents, and are hence typically used as monodentate ligands. Examples of suitable isonitriles include simple alkyl isonitriles such as *tert*-butylisonitrile, and ether-substituted isonitriles such as mibi (i.e. 1-isocyano-2-methoxy-2-methylpropane). Examples of suitable phosphines include Tetrofosmin, and monodentate phosphines such as *tris*(3-methoxypropyl)phosphine. Examples of suitable dia-

zenides include the HYNIC series of ligands i.e. hydrazine-substituted pyridines or nicotinamides.

[0053] Examples of suitable chelating agents for technetium which form metal complexes resistant to transchelation include, but are not limited to:

(i) diaminedioximes of Formula III:



where E¹-E⁶ are each independently an R* group;

each R* is H or C₁₋₁₀ alkyl, C₃₋₁₀ alkylaryl, C₂₋₁₀ alkoxyalkyl, C₁₋₁₀ hydroxyalkyl, C₁₋₁₀ fluoroalkyl, C₂₋₁₀ carboxyalkyl or C₁₋₁₀ aminoalkyl, or two or more R* groups together with the atoms to which they are attached form a carbocyclic, heterocyclic, saturated or unsaturated ring, and wherein one or more of the R* groups is conjugated to the vector;

and Q is a bridging group of formula -(J)_f;

where f is 3, 4 or 5 and each J is independently -O-, -NR*- or -C(R*)₂-, wherein R* is as previously defined, provided that -(J)_f contains a maximum of one J group which is -O- or -NR*-.

[0054] Preferred Q groups are as follows:


Q = -(CH₂)(CHR*)(CH₂)- i.e. propyleneamine oxime or PnAO derivatives;

Q = -(CH₂)₂(CHR*)(CH₂)₂- i.e. pentyleneamine oxime or PentAO derivatives;

Q = -(CH₂)₂NR*(CH₂)₂-.

[0055] E¹ to E⁶ are preferably chosen from: C₁₋₃ alkyl, alkylaryl alkoxyalkyl, hydroxyalkyl, fluoroalkyl, carboxyalkyl or aminoalkyl. Most preferably, each E¹ to E⁶ group is CH₃.

[0056] The compound of the invention is preferably conjugated at either the E¹ or E⁶ R* group, or an R* group of the Q moiety. Most preferably, it is conjugated to an R* group of the Q moiety. When it is conjugated to an R* group of the Q moiety, the R* group is preferably at the bridgehead position. In that case, Q is preferably -(CH₂)(CHR*)(CH₂)-, -(CH₂)₂(CHR*)(CH₂)₂- or -(CH₂)₂NR*(CH₂)₂-, most preferably -(CH₂)₂(CHR*)(CH₂)₂-. An especially preferred bifunctional diaminedioxime chelator has the Formula IV:



where:

E⁷-E²⁰ are each independently an R* group as previously defined;


G is N or CE²¹ wherein E²¹ is an R* group as previously defined;

5 Z is the site of linkage to a compound of Formula I and may comprise a linker group -(L²)_r-wherein each L² is independently -O-, -NR*, -C(R*)₂-, or a C₅₋₁₂ arylene group wherein R* is as previously defined, and r is an integer between 1 and 5.

10 [0057] A preferred chelator of Formula IV is of Formula IVa:

15

20



where G is as defined above;

25

such that the compound of Formula I is conjugated via the bridgehead -CH₂CH₂NH₂ group.

[0058] Further suitable chelators of the invention include:

30

(ii) N₃S ligands having a thioltriamide donor set such as MAG₃ (mercaptoacetyltriglycine) and related ligands; or having a diamidepyridinethiol donor set such as Pica;

(iii) N₂S₂ ligands having a diaminedithiol donor set such as BAT or ECD (i.e. ethylcysteinate dimer), or an amide-aminedithiol donor set such as MAMA;

35

(iv) N₄ ligands which are open chain or macrocyclic ligands having a tetramine, amidetriamine or diamidediamine donor set, such as cyclam, monoxocyclam or dioxocyclam.

(v) N₂O₂ ligands having a diaminediphenol donor set.

40

[0059] The above described ligands are particularly suitable for complexing technetium e.g. ^{94m}Tc or ^{99m}Tc, and are described more fully by Jurisson et al [Chem.Rev., 99, 2205-2218 (1999)]. The ligands are also useful for other metals, such as copper (⁶⁴Cu or ⁶⁷Cu), vanadium (e.g. ⁴⁸V), iron (e.g. ⁵²Fe), or cobalt (e.g. ⁵⁵Co). Other suitable ligands are described in Sandoz WO 91/01144, which includes ligands which are particularly suitable for indium, yttrium and gadolinium, especially macrocyclic aminocarboxylate and aminophosphonic acid ligands. Ligands which form nonionic (i.e. neutral) metal complexes of gadolinium are known and are described in US 4885363. When the radiometal ion is technetium, the ligand is preferably a chelating agent which is tetradentate. Preferred chelating agents for technetium are the diaminedioximes, or those having an N₂S₂ or N₃S donor set as described above.

45

[0060] It is envisaged that the role of the linker group -(L₂)_r- is to distance the relatively bulky technetium complex, which results upon metal coordination, from the active site of the compound of Formula I so that e.g. receptor binding is not impaired.


50

[0061] Preferred linker groups -(L₂)_r- in the context of these chelators have a backbone chain (i.e. the linked atoms which make up the -(L₂)_r- moiety) which contains 2 to 5 atoms, with 2 or 3 atoms being most preferred. A minimum linker group backbone chain of 2 atoms confers the advantage that the aza-diaminedioxime chelator is well-separated from the biological targeting moiety so that any interaction is minimised. Furthermore, the vector is unlikely to compete effectively with the coordination of the chelator to the metal ion. In this way, both the biological targeting characteristics of the vector, and the metal complexing capability of the chelator is maintained. It is strongly preferred that the compound of Formula I is bound to the chelator in such a way that the linkage does not undergo facile metabolism in blood. That is because such metabolism would result in the imaging metal complex being cleaved off before the labelled compound

55

reaches the desired *in vivo* target site. The compound of Formula I is therefore preferably covalently bound to the metal complexes of the present invention *via* $-(L_2)_r$ - linker groups which are not readily metabolised.

[0062] Non-peptide linker groups such as alkylene groups or arylene groups have the advantage that there are no significant hydrogen bonding interactions with the conjugated compound of Formula I so that the linker does not wrap round onto the compound. Preferred alkylene spacer groups are $-(CH_2)_q$ - where q is an integer of value 2 to 5. Preferably q is 2 or 3. Preferred arylene spacers are of formula:



where: a and b are each independently 0, 1 or 2.

[0063] When the imaging metal is technetium, the usual technetium starting material is pertechnetate, i.e. TcO_4^- which is technetium in the Tc(VII) oxidation state. Pertechnetate itself does not readily form metal complexes, hence the preparation of technetium complexes usually requires the addition of a suitable reducing agent such as stannous ion to facilitate complexation by reducing the oxidation state of the technetium to the lower oxidation states, usually Tc(I) to Tc(V). The solvent may be organic or aqueous, or mixtures thereof. When the solvent comprises an organic solvent, the organic solvent is preferably a biocompatible solvent, such as ethanol or DMSO. Preferably the solvent is aqueous, and is most preferably isotonic saline.

[0064] Where the imaging moiety is radioiodine, preferred precursors are those which comprise a derivative which either undergoes electrophilic or nucleophilic iodination or undergoes condensation with a labelled aldehyde or ketone. Examples of the first category are:


(a) organometallic derivatives such as a trialkylstannane (e.g. trimethylstannyl or tributylstannyl, or a trialkylsilane (e.g. trimethylsilyl) or an organoboron compound (e.g. boronate esters or organotrifluoroborates);

(b) a non-radioactive alkyl bromide for halogen exchange or alkyl tosylate, mesylate or triflate for nucleophilic iodination;


(c) aromatic rings activated towards electrophilic iodination (e.g. phenols) and aromatic rings activated towards nucleophilic iodination (e.g. aryl iodonium salt aryl diazonium, aryl trialkylammonium salts or nitroaryl derivatives).

[0065] For radioiodination, the precursor preferably comprises: an aryl iodide or bromide (to permit radioiodine exchange); an activated precursor aryl ring (e.g. a phenol group); an organometallic precursor compound (e.g. trialkyltin, trialkylsilyl or organoboron compound); or an organic precursor such as triazenes or a good leaving group for nucleophilic substitution such as an iodonium salt. Precursors and methods of introducing radioiodine into organic molecules are described by Bolton [J.Lab.Comp.Radiopharm., 45, 485-528 (2002)]. Precursors and methods of introducing radioiodine into proteins are described by Wilbur [Bioconj.Chem., 3(6), 433-470 (1992)]. Suitable boronate ester organoboron compounds and their preparation are described by Kabalaka et al [Nucl.Med.Biol., 29, 841-843 (2002) and 30, 369-373 (2003)]. Suitable organotrifluoroborates and their preparation are described by Kabalaka et al [Nucl.Med.Biol., 31, 935-938 (2004)]. Preferred precursors for radioiodination comprise an organometallic precursor compound, most preferably a trialkyltin.


[0066] Examples of aryl groups to which radioactive iodine can be attached are given below:



[0067] Both contain substituents which permit facile radioiodine substitution onto the aromatic ring. Alternative substituents containing radioactive iodine can be synthesised by direct iodination via radiohalogen exchange, e.g.




[0068] Examples of precursor compounds of Formula I derivatised to include the above aryl groups are as follows:



[0069] The radioiodine atom is preferably attached via a direct covalent bond to an aromatic ring such as a benzene ring, or a vinyl group since it is known that iodine atoms bound to saturated aliphatic systems are prone to *in vivo* metabolism and hence loss of the radioiodine.

[0070] When the imaging moiety is a radioactive isotope of fluorine the radiofluorine atom may form part of a fluoroalkyl or fluoroalkoxy group, since alkyl fluorides are resistant to *in vivo* metabolism. Alternatively, the radiofluorine atom may attach via a direct covalent bond to an aromatic ring such as a benzene ring. Radiofluorination may be carried out via direct labelling using the reaction of ^{18}F -fluoride with a suitable chemical group in the precursor having a good leaving group, such as an alkyl bromide, alkyl mesylate or alkyl tosylate. ^{18}F can also be introduced by O-alkylation of hydroxyl groups with $^{18}\text{F}(\text{CH}_2)_3\text{OMs}$ or $^{18}\text{F}(\text{CH}_2)_3\text{Br}$. ^{18}F can also be introduced by alkylation of N-haloacetyl groups with a $^{28}\text{F}(\text{CH}_2)_3\text{OH}$ reactant, to give $-\text{NH}(\text{CO})\text{CH}_2\text{O}(\text{CH}_2)_3^{18}\text{F}$ derivatives. For aryl systems, ^{18}F -fluoride nucleophilic displacement from an aryl diazonium salt, aryl nitro compound or an aryl quaternary ammonium salt are suitable routes to aryl- ^{18}F derivatives.

[0071] A ^{18}F -labelled compound of the invention of Formula Ib may be obtained by formation of ^{18}F fluorodialkylamines and subsequent amide formation as shown in the following reaction scheme:



wherein X^{21} , X^{22} , Y^{21} and s are the same as X^1 , X^2 , Y^1 and n , respectively, described previously for Formula I; and

LG is a suitable leaving group, e.g. Cl, Br, I, OTs, OMs, or OTf;

Z* is e.g. Cl, P(O)Ph₃, or an activated ester; and,


x = 0-6.

[0072] Alternatively, labeling with ¹⁸F can be achieved by nucleophilic displacement of a leaving group (LG as defined above) from a derivative of Formula I. Such derivatives are precursors for the preparation of *in vivo* imaging compounds of the invention.

[0073] Another strategy would be to have a leaving group (LG, as defined above) in place on an alkylamide group present on the precursor. In this way, the precursor compound may be labeled in one step by reaction with a suitable source of [¹⁸F]-fluoride ion (¹⁸F⁻), which is normally obtained as an aqueous solution from the nuclear reaction ¹⁸O(p, n)¹⁸F and is made reactive by the addition of a cationic counterion and the subsequent removal of water. For this method, the precursor compounds are normally selectively chemically protected so that radiofluorination takes place at a particular site on the compound. Suitable protecting groups are those already mentioned previously.

[0074] ¹¹C-labelled PET tracer compounds may be synthesised by reacting a precursor with ¹¹C methyl iodide. As the half-life of ¹¹C is only 20.4 minutes, it is important that the intermediate ¹¹C methyl iodide has high specific activity and, consequently, that it is produced using a reaction process which is as rapid as possible. A thorough review of such ¹¹C-labelling techniques may be found in Antoni et al "Aspects on the Synthesis of ¹¹C-Labelled Compounds" in Handbook of Radiopharmaceuticals, Ed. M.J. Welch and C.S. Redvanly (2003, John Wiley and Sons).

[0075] A ¹¹C-labelled compound of Formula Ic may be obtained by employing the following reaction scheme:



wherein X³¹, X³², Y³¹ and t are as described previously for X¹, X², Y¹ and n, respectively of Formula I; and

Z** is a substrate suitable for transition metal catalysts, e.g. hydrogen, halide, boronic acid, OTf, organotin; and,

y = 0-6.

[0076] The precursor is ideally provided in sterile, apyrogenic form. The precursor can accordingly be used for the preparation of a pharmaceutical composition and is also suitable for inclusion as a component in a kit for the preparation of a pharmaceutical composition. These aspects are discussed in more detail below in relation to additional aspects of the invention.

[0077] In a further preferred embodiment of the method of the invention, the precursor is bound to a solid phase. The precursor is preferably supplied covalently attached to a solid support matrix. In this way, the desired product forms in solution, whereas starting materials and impurities remain bound to the solid phase. As an example of such a system, precursors for solid phase electrophilic fluorination with ¹⁸F-fluoride are described in WO 03/002489, and precursors for solid phase nucleophilic fluorination with ¹⁸F-fluoride are described in WO 03/002157.


[0078] The Examples section below describes some methods for preparation of compounds of the invention where certain precursors are used:

[0079] Example 5 uses 6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid propylamide as a precursor in the synthesis of Compound 12.

[0080] Example 14 uses 11-(2-Tosyloxy-ethyl)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide as a precursor in the synthesis of Compound 10.

[0081] Example 17 uses 11-(2-hydroxy-ethyl)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide as a precursor in the synthesis of Compound 11.

[0082] In a further aspect, the invention provides a precursor for the preparation of the *in vivo* imaging compound of the invention wherein said precursor is selected from :



where Compound 8P is a precursor for the preparation of Compound 8, Compounds 9P¹ and 9P² (also non-radioactive Compound 6) are precursors for the preparation of Compound 9, and Compound 10P is a precursor for the preparation of Compound 10.

[0083] In another further aspect, the present invention provides a pharmaceutical composition which comprises a compound of the invention together with a biocompatible carrier in a form suitable for mammalian administration.

[0084] Where the pharmaceutical composition comprises a compound of Formula I labelled with an imaging moiety, the "biocompatible carrier" is a fluid, especially a liquid, in which the compound is suspended or dissolved, such that the composition is physiologically tolerable, i.e. can be administered to the mammalian body without toxicity or undue discomfort. The biocompatible carrier medium is suitably an injectable carrier liquid such as sterile, pyrogen-free water for injection; an aqueous solution such as saline (which may advantageously be balanced so that the final product for injection is either isotonic or not hypotonic); an aqueous solution of one or more tonicity-adjusting substances (e.g. salts of plasma cations with biocompatible counterions), sugars (e.g. glucose or sucrose), sugar alcohols (e.g. sorbitol or mannitol), glycols (e.g. glycerol), or other non-ionic polyol materials (e.g. polyethyleneglycols, propylene glycols and the like). The biocompatible carrier medium may also comprise biocompatible organic solvents such as ethanol. Such organic solvents are useful to solubilise more lipophilic compounds or formulations. Preferably the biocompatible carrier medium is pyrogen-free water for injection, isotonic saline or an aqueous ethanol solution. The pH of the biocompatible carrier medium for intravenous injection is suitably in the range 4.0 to 10.5.

[0085] Such pharmaceutical compositions comprising a compound of Formula I labelled with an imaging moiety are suitably supplied in either a container which is provided with a seal which is suitable for single or multiple puncturing with a hypodermic needle (e.g. a crimped-on septum seal closure) whilst maintaining sterile integrity. Such containers may contain single or multiple patient doses. Preferred multiple dose containers comprise a single bulk vial (e.g. of 10 to 30 cm³ volume) which contains multiple patient doses, whereby single patient doses can thus be withdrawn into clinical grade syringes at various time intervals during the viable lifetime of the preparation to suit the clinical situation. Pre-filled syringes are designed to contain a single human dose, or "unit dose" and are therefore preferably a disposable or other syringe suitable for clinical use.

[0086] Preferably, where the compound is a compound of Formula I labelled with an imaging moiety, the pharmaceutical composition is a radiopharmaceutical composition. For radiopharmaceutical compositions, the pre-filled syringe may optionally be provided with a syringe shield to protect the operator from radioactive dose. Suitable such radiopharmaceutical syringe shields are known in the art and preferably comprise either lead or tungsten.

[0087] The radiopharmaceuticals may be administered to patients for SPECT or PET imaging in amounts sufficient to yield the desired signal, typical radionuclide dosages of 0.01 to 100 mCi, preferably 0.1 to 50 mCi will normally be sufficient per 70kg bodyweight.

[0088] The pharmaceutical composition comprising a compound of Formula I labelled with an imaging moiety may be prepared from kits, as is described below. Alternatively, such a pharmaceutical composition may be prepared under aseptic manufacture conditions to give the desired sterile product. This pharmaceutical composition may also be prepared under non-sterile conditions, followed by terminal sterilisation using e.g. gamma-irradiation, autoclaving, dry heat or chemical treatment (e.g. with ethylene oxide). Preferably, the pharmaceutical composition comprising the compound of

Formula I labelled with an imaging moiety is prepared from a kit.

[0089] Where the pharmaceutical composition comprises the compound of Formula II the biocompatible carrier may be a solid or liquid pharmaceutically acceptable nontoxic carrier. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerols solutions are also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatine, malt, rice, flour, chalk, silica gel, magnesium carbonate, magnesium stearate, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol and the like. These compositions can take the form of solutions, suspensions, tablets, pills, capsules, powders, sustained release formulations and the like. Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E. W. Martin. Such compositions will contain an effective therapeutic amount of the compound together with a suitable amount of carrier so as to provide the form for proper administration to the host. While intravenous injection is a very effective form of administration, other modes can be employed, e.g. oral administration.

[0090] The posology depends on the effects required and the method of administration used. For example, by the oral route, it may be between 20 and 100 mg of active substance per day, with unit doses of from 5 to 200 mg.

[0091] In a yet further aspect, the present invention provides kits for the preparation of the pharmaceutical compositions wherein the compound is a compound of Formula I labelled with an imaging moiety. Such kits comprise a precursor selected from compounds 8P, 9P¹, 9P² and 10P (defined above), preferably in sterile non-pyrogenic form, so that reaction with a sterile source of an imaging moiety gives the desired pharmaceutical with the minimum number of manipulations. Such considerations are particularly important for radiopharmaceuticals, in particular where the radioisotope has a relatively short half-life, and for ease of handling and hence reduced radiation dose for the radiopharmacist. Hence, the reaction medium for reconstitution of such kits is preferably a "biocompatible carrier" as defined above, and is most preferably aqueous.

[0092] Suitable kit containers comprise a sealed container which permits maintenance of sterile integrity and/or radioactive safety, plus optionally an inert headspace gas (e.g. nitrogen or argon), whilst permitting addition and withdrawal of solutions by syringe. A preferred such container is a septum-sealed vial, wherein the gas-tight closure is crimped on with an overseal (typically of aluminium). Such containers have the additional advantage that the closure can withstand vacuum if desired e.g. to change the headspace gas or degas solutions.

[0093] In the case of precursors bound to a solid phase, the sealed container may be a cartridge provided as part of the kit, which can be plugged into a suitably adapted automated synthesizer. The cartridge may contain, apart from the solid support-bound precursor, a column to remove unwanted fluoride ion, and an appropriate vessel connected so as to allow the reaction mixture to be evaporated and allow the product to be formulated as required. These cartridges are especially useful for the preparation of compounds of the invention labeled with short-lived radioisotopes such as ¹¹C or ¹⁸F.

[0094] Preferred embodiments of the precursor when employed in the kit are as described above. The precursors for use in the kit may be employed under aseptic manufacture conditions to give the desired sterile, non-pyrogenic material. The precursors may also be employed under non-sterile conditions, followed by terminal sterilisation using e.g. gamma-irradiation, autoclaving, dry heat or chemical treatment (e.g. with ethylene oxide). Preferably, the precursors are employed in sterile, non-pyrogenic form. Most preferably the sterile, non-pyrogenic precursors are employed in the sealed container as described above.

[0095] The kits may optionally further comprise additional components such as a radioprotectant, antimicrobial preservative, pH-adjusting agent or filler.

[0096] By the term "radioprotectant" is meant a compound which inhibits degradation reactions, such as redox processes, by trapping highly-reactive free radicals, such as oxygen-containing free radicals arising from the radiolysis of water. The radioprotectants of the present invention are suitably chosen from: ascorbic acid, para-aminobenzoic acid (i.e. 4-aminobenzoic acid), gentisic acid (i.e. 2,5-dihydroxybenzoic acid) and salts thereof with a biocompatible cation. The "biocompatible cation" and preferred embodiments thereof are as described above. By the term "antimicrobial preservative" is meant an agent which inhibits the growth of potentially harmful micro-organisms such as bacteria, yeasts or moulds. The antimicrobial preservative may also exhibit some bactericidal properties, depending on the dose. The main role of the antimicrobial preservative(s) of the present invention is to inhibit the growth of any such micro-organism in the pharmaceutical composition post-reconstitution, i.e. in the radioactive imaging product itself. The antimicrobial preservative may, however, also optionally be used to inhibit the growth of potentially harmful micro-organisms in one or more components of the non-radioactive kit of the present invention prior to reconstitution. Suitable antimicrobial preservative(s) include: the parabens, i.e. methyl, ethyl, propyl or butyl paraben or mixtures thereof; benzyl alcohol; phenol; cresol; cetrimide and thiomersal. Preferred antimicrobial preservative(s) are the parabens.

[0097] The term "pH-adjusting agent" means a compound or mixture of compounds useful to ensure that the pH of the reconstituted kit is within acceptable limits (approximately pH 4.0 to 10.5) for human or mammalian administration.

Suitable such pH-adjusting agents include pharmaceutically acceptable buffers, such as tricine, phosphate or TRIS [i.e. *tris*(hydroxymethyl)aminomethane], and pharmaceutically acceptable bases such as sodium carbonate, sodium bicarbonate or mixtures thereof. When the conjugate is employed in acid salt form, the pH adjusting agent may optionally be provided in a separate vial or container, so that the user of the kit can adjust the pH as part of a multi-step procedure.

[0098] By the term "filler" is meant a pharmaceutically acceptable bulking agent which may facilitate material handling during production and lyophilisation. Suitable fillers include inorganic salts such as sodium chloride, and water soluble sugars or sugar alcohols such as sucrose, maltose, mannitol or trehalose.

[0099] The compounds of the invention are useful for *in vivo* imaging. Accordingly, in an even further aspect, the present invention provides a compound of the invention for use in an *in vivo* imaging method, such as SPECT or PET, which are preferred, and magnetic resonance imaging (MRI) or optical imaging. The imaging method may be used to study PBR in healthy subjects, or in subjects known or suspected to have a pathological condition associated with abnormal expression of PBR (a "PBR condition"). Preferably, said method relates to the *in vivo* imaging of a subject suspected to have a PBR condition, and therefore has utility in the diagnosis of said condition. Examples of such PBR conditions where *in vivo* imaging would be of use include neuropathologies such as Parkinson's disease, multiple sclerosis, Alzheimer's disease and Huntington's disease where neuroinflammation is present. Other PBR conditions that may be usefully imaged with the compounds of the invention include neuropathic pain, arthritis, asthma, atherosclerosis and cancer.

[0100] This aspect of the invention also provides a method for the *in vivo* diagnosis or imaging of a PBR condition in a subject, comprising administration of a pharmaceutical composition comprising a compound of the invention. Said subject is preferably a mammal and most preferably a human. In an alternative embodiment, this aspect of the invention furthermore provides for the use of the compound of the invention for imaging *in vivo* in a subject of a PBR condition wherein said subject is previously administered with the pharmaceutical composition comprising a compound of Formula I labelled with an imaging moiety.

[0101] By "previously administered" is meant that the step involving the clinician, wherein the imaging agent is given to the patient e.g., intravenous injection, has already been carried out. This aspect of the invention includes the use of a compound of Formula I labelled with an imaging moiety for the manufacture of diagnostic agent for the diagnostic imaging *in vivo* of a PBR condition.

[0102] Furthermore, this aspect of the invention provides for use of the compound of the invention in the manufacture of a pharmaceutical for the *in vivo* diagnosis or imaging of a PBR condition.

[0103] The compounds of Formula I may also be used for *in vivo* imaging of PBR in human and animal subjects in the context of their use as research tools. For example, for the performance of competition studies which allow the interaction of a drug with PBR to be studied. Such studies include dose-occupancy studies, determination of optimal therapeutic dose, drug candidate selection studies, and determination of PBR distribution in the tissue of interest.

[0104] In another aspect the invention provides a compound of Formula I as defined herein for use in a method of monitoring the effect of treatment of a human or animal body with a drug to combat a PBR condition, said method comprising administering to said body a compound of Formula I labelled with an imaging moiety and detecting the uptake of said compound, said administration and detection optionally but preferably being effected repeatedly, e.g. before, during and after treatment with said drug.

[0105] In a further aspect, the present invention provides a method for the treatment of a PBR condition in a mammal, preferably a human, by administration of a pharmaceutical composition comprising the compound of Formula II. Details of the nature of such a pharmaceutical composition, its administration and dosage are described above.

Brief Description of the Examples

[0106] Example 1 describes an alternative synthetic route to Okubo *et al* for obtaining 6,11-Dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (this is a non-radioactive version of Compound 1 of the invention). The alternative synthesis could be easily adapted to obtain other compounds of Formula I. An increased yield of product (54%) was obtained using this method compared with the method of Okubo *et al*.

[0107] Example 2 describes the synthesis of a chelate of Formula IIIa wherein G is C. This chelate is suitable for forming a complex with ^{99m}Tc .

[0108] Example 3 describes how to conjugate the chelate of Example 1 to a compound of Formula I to obtain a precursor compound of the invention.

[0109] Example 4 describes how the precursor compounds obtained by the method of Example 2 can be labelled with ^{99m}Tc in order to obtain compounds of the invention.

[0110] Example 5 describes the synthesis of a precursor compound suitable for reaction with ^{18}F and how to label the precursor compound with ^{18}F to form a Compound 12.

[0111] Example 6 describes the synthesis of a precursor compound suitable for reaction with ^{11}C and how to label the precursor compound with ^{11}C to form a compound of the invention.

[0112] Examples 7-13 describe how to obtain non-radioactive versions of Compounds 3-9 of the invention.

[0113] Example 14 describes how to obtain a non-radioactive version of Compound 10 *via* a route which may by analogy be used to obtain the radioactive version.

[0114] Example 15 describes the synthetic route used to obtain Compound 10.

5 [0115] Example 16 describes how to obtain a non-radioactive version of Compound 11 *via* a route which by analogy may be used to obtain the radioactive version.


[0116] Example 17 is a prophetic example describing a method suitable for the preparation of Compound 11.

[0117] Example 18 describes the method used to screen the compounds of the invention for their affinity to PBR.

10 Examples

Example 1: Synthesis of 6,11-Dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (non-radioactive Compound 1)

15 [0118]



30 a) 3-Phenylsulfanyl-dihydro-furan-2,5-dione:

[0119] Triethylamine (0.8ml) was added dropwise to a solution of benzenethiol (9.3ml, 91mmol) and maleic anhydride (8.9g, 91mmol) in toluene (125ml). After stirring at room temperature for 12hr the solvent was evaporated to leave 20g of crude 3-phenylsulfanyl-dihydro-furan-2,5-dione as a brown oil. ¹H NMR (CDCl₃) δ 7.20-7.70(5H, m), 4.20(1H, dd), 3.40(1H, dd), 2.90(1H, dd).

b) 4-Oxo-thiochroman-2-carboxylic acid:

[0120] Crude 3-phenylsulfanyl-dihydro-furan-2,5-dione (20g, 91mmol) was dissolved in CH₂Cl₂ (30ml) and cooled to 0°C. Aluminium chloride (18.16g, 136mmol) was added and the mixture was stirred at room temperature for 1.5hr. The reaction mixture was diluted with CH₂Cl₂ (1000ml) and poured into ice-cooled conc HCl (1000ml). The phases were separated and the aqueous phase was extracted with CH₂Cl₂(x3). The combined organic phases were washed with water, dried (MgSO₄) and evaporated to a brown solid. The solid was triturated with Et₂O to give 8.98g of 4-oxo-thiochroman-2-carboxylic acid as a pale brown solid. ¹H NMR (DMSO) δ 7.96(1H, dd), 7.20-7.60(3H, m), 4.40(1H, dd), 3.20-3.33(2H, m).

c) 6,11-Dihydro-5-thia-11-azo-benzo[α]fluorene-6-carboxylic acid ethyl ester:

[0121] To a solution of 4-oxo-thiochroman-2-carboxylic acid (3g, 14mmol) and phenyl hydrazine (1.4ml, 14mmol) in EtOH (14ml), H₂SO₄ (1.8ml) was added, and the mixture was heated at reflux for 5hr. The reaction was cooled to room temperature and the solid, which formed overnight, was filtered, washed with cold EtOH and cold Et₂O to give 2.26g (51%) of 6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid ethyl ester as a crème solid. ¹H NMR (CDCl₃) δ 8.47(1H, br s), 7.53-7.58(1H, m), 7.10-7.40(7H, m), 5.00(1H, s), 4.09(2H, q), 1.15(3H, t).

55 d) 6,11-Dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid:

[0122] A solution of KOH (1.64g, 29mmol) in water (6ml) was added to 6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid ethyl ester (2.26g, 7mmol) in EtOH (16ml) and the mixture heated to reflux for 2hr. The reaction mixture

was acidified with 2N HCl and extracted with EtOAc. The organic phase was washed with brine, dried (MgSO₄) and evaporated to give 1.65g (80%) of 6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid as a yellow foam. ¹H NMR (DMSO) δ 12.55(1H, br s), 11.70(1H, s), 7.74-7.82(1H, m), 7.00-7.57(7H, m), 5.17(1H, s).

5 e) 6,11-Dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide:

[0123] To 6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid (1.65g, 6mmol) in CH₂Cl₂ (15ml) were added diethylamine (0.7ml, 7mmol), bromotripyrrolidinophosphonium hexafluorophosphate (2.75g, 6mmol) and diisopropylethylamine (3.1ml, 18mmol). The reaction was stirred at room temperature for 3 days. The reaction mixture was diluted with CH₂Cl₂ (100ml), washed with 1N HCl, saturated NaHCO₃ soln, brine, dried (MgSO₄) and evaporated. The residue was purified by silica gel chromatography eluting with 50% EtOAc/petroleum ether to give 1.07g (54%) of 6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide as a yellow solid. ¹H NMR (CDCl₃) δ 9.23(1H, s), 7.33(1H, dd), 7.20(1H, dd), 7.69-7.08(5H, m), 6.66-6.72(1H, m), 5.33(1H, s), 3.34-3.76(4H, m), 1.39(3H, t), 1.37(3H, t).

15 **Example 2: Synthesis of chelate of Formula IIIa (where G = C)**

(Step a): Preparation of tris(methyloxycarbonylmethyl)methane

[0124] 3-(methoxycarbonylmethylene)glutaric acid dimethylester (89g, 267mmol) in methanol (200ml) was shaken with (10% palladium on charcoal: 50% water) (9 g) under an atmosphere of hydrogen gas (3.5 bar) for (30h). The solution was filtered through kieselguhr and concentrated *in vacuo* to give 3-(methoxycarbonylmethyl)glutaric acid dimethylester as an oil, yield (84.9g, 94 %).

NMR ¹H(CDCl₃), δ 2.48 (6H, d, J=8Hz, 3xCH₂), 2.78 (1H, hextet, J=8Hz CH), 3.7(9H, s, 3xCH₃).

NMR ¹³C(CDCl₃), δ 28.6, CH; 37.50, 3xCH₃; 51.6, 3xCH₂, 172.28, 3xCOO.

(Step b): Amidation of trimethylester with p-methoxy-benzylamine

[0125] Tris(methyloxycarbonylmethyl)methane [2 g, 8.4 mmol] was dissolved in p-methoxybenzylamine (25 g, 178.6 mmol). The apparatus was set up for distillation and heated to 120 °C for 24 hrs under nitrogen flow. The progress of the reaction was monitored by the amount of methanol collected. The reaction mixture was cooled to ambient temperature and 30 ml of ethyl acetate was added, then the precipitated triamide product stirred for 30 min. The triamide was isolated by filtration and the filter cake washed several times with sufficient amounts of ethyl acetate to remove excess p-methoxybenzylamine. After drying 4.6 g, 100 %, of a white powder was obtained. The highly insoluble product was used directly in the next step without further purification or characterisation.

(Step c): Preparation of 1,1,1-tris[2-(p-methoxybenzylamino)ethyl]methane.

[0126] To a 1000 ml 3-necked round bottomed flask cooled in a ice-water bath the triamide from step 2(a) (10 g, 17.89 mmol) was carefully added to 250 ml of 1M borane solution (3.5 g, 244.3 mmol) borane. After complete addition the ice-water bath was removed and the reaction mixture slowly heated to 60 °C. The reaction mixture was stirred at 60 °C for 20 hrs. A sample of the reaction mixture (1 ml) was withdrawn, and mixed with 0.5 ml 5N HCl and left standing for 30 min. To the sample 0.5 ml of 50 NaOH was added, followed by 2 ml of water and the solution was stirred until all of the white precipitate dissolved. The solution was extracted with ether (5 ml) and evaporated. The residue was dissolved in acetonitrile at a concentration of 1 mg/ml and analysed by MS. If mono- and diamide (M+H/z = 520 and 534) are seen in the MS spectrum, the reaction was not complete. To complete the reaction, a further 100 ml of 1M borane THF solution was added and the reaction mixture stirred for 6 more hrs at 60 °C and a new sample withdrawn following the previous sampling procedure. Further addition of the 1M borane in THF solution was continued as necessary until there was complete conversion to the triamine.

[0127] The reaction mixture was cooled to ambient temperature and 5N HCl is slowly added, [CARE: vigorous foam formation occurs!]. HCl was added until no more gas evolution was observed. The mixture was stirred for 30 min and then evaporated. The cake was suspended in aqueous NaOH solution (20-40 %; 1:2 w/v) and stirred for 30 minutes. The mixture was then diluted with water (3 volumes). The mixture was then extracted with diethylether (2 x 150 ml) [CARE: do not use halogenated solvents]. The combined organic phases were then washed with water (1x 200 ml), brine (150 ml) and dried over magnesium sulphate. Yield after evaporation: 7.6 g, 84 % as oil.

NMR ¹H(CDCl₃), δ : 1.45, (6H, m, 3xCH₂); 1.54, (1H, septet, CH); 2.60 (6H, t, 3xCH₂N); 3.68 (6H, s, ArCH₂); 3.78 (9H, s, 3xCH₃O); 6.94(6H, d, 6xAr). 7.20(6H, d, 6xAr).

NMR ¹³C(CDCl₃), δ : 32.17, CH; 34.44, CH₂; 47.00, CH₂; 53.56, ArCH₂; 55.25, CH₃O; 113.78, Ar; 129.29, Ar; 132.61; Ar; 158.60, Ar.

(Step d): Preparation of 1,1,1-*tris*(2-aminoethyl)methane.

[0128] 1,1,1-*tris*[2-(*p*-methoxybenzylamino)ethyl]methane (20.0 gram, 0.036 mol) was dissolved in methanol (100 ml) and Pd(OH)₂ (5.0 gram) was added. The mixture was hydrogenated (3 bar, 100 °C, in an autoclave) and stirred for 5 hours. Pd(OH)₂ was added in two more portions (2 x 5gram) after 10 and 15 hours respectively.

[0129] The reaction mixture was filtered and the filtrate was washed with methanol. The combined organic phase was evaporated and the residue was distilled under vacuum

(1 x 10⁻², 110 °C) to give 2.60 gram (50 %) of 1,1,1-*tris*(2-aminoethyl)methane.

NMR ¹H(CDCl₃), δ 2.72 (6H, t, 3xCH₂N), 1.41 (H, septet, CH), 1.39 (6H, q, 3xCH₂).

NMR ¹³C(CDCl₃), δ 39.8 (CH₂NH₂), 38.2(CH₂), 31.0(CH).

(Step e): Preparation of Formula IIIa (where G = C)

[0130] To a solution of *tris*(2-aminoethyl)methane (4.047g, 27.9mmol) in dry ethanol (30ml) was added potassium carbonate anhydrous (7.7g, 55.8mmol, 2eq) at room temperature with vigorous stirring under a nitrogen atmosphere. A solution of 3-chloro-3-methyl-2-nitrosobutane (7.56g, 55.8mol, 2eq) was dissolved in dry ethanol (100ml) and 75ml of this solution was dripped slowly into the reaction mixture. The reaction was followed by TLC on silica [plates run in dichloromethane, methanol, concentrated (0.88sg) ammonia; 100/30/5 and the TLC plate developed by spraying with ninhydrin and heating]. The mono-, di- and tri-alkylated products were seen with RF's increasing in that order. Analytical HPLC was run using RPR reverse phase column in a gradient of 7.5-75% acetonitrile in 3% aqueous ammonia. The reaction was concentrated *in vacuo* to remove the ethanol and resuspended in water (110ml). The aqueous slurry was extracted with ether (100ml) to remove some of the trialkylated compound and lipophilic impurities leaving the mono and desired dialkylated product in the water layer. The aqueous solution was buffered with ammonium acetate (2eq, 4.3g, 55.8mmol) to ensure good chromatography. The aqueous solution was stored at 4°C overnight before purifying by automated preparative HPLC.

Yield (2.2g, 6.4mmol, 23%).

Mass spec; Positive ion 10 V cone voltage. Found: 344; calculated M+H= 344.

NMR ¹H(CDCl₃), δ 1.24(6H, s, 2xCH₃), 1.3(6H, s, 2xCH₃), 1.25-1.75(7H, m, 3xCH₂,CH), (3H, s, 2xCH₂), 2.58 (4H, m, CH₂N), 2.88(2H, t CH₂N₂), 5.0 (6H, s, NH₂, 2xNH, 2xOH).

NMR ¹H ((CD₃)₂SO) δ 1.1 4xCH; 1.29, 3xCH₂; 2.1 (4H, t, 2xCH₂);

NMR ¹³C((CD₃)₂SO), δ 9.0 (4xCH₃), 25.8 (2xCH₃), 31.0 2xCH₂, 34.6 CH₂, 56.8 2xCH₂N; 160.3, C=N.

HPLC conditions: flow rate 8ml/min using a 25mm PRP column

A=3% ammonia solution (sp.gr = 0.88) /water; B = Acetonitrile

	Time	%B
	0	7.5
	15	75.0
	20	75.0
	22	7.5
	30	7.5


Load 3ml of aqueous solution per run, and collect in a time window of 12.5-13.5 min.

Example 3: Synthesis of precursors of the invention for ^{99m}Tc labelling


[0131] The chelate produced in Example 2 may be conjugated at R¹ or R² of a compound of Formula I via the bridgehead -CH₂CH₂NH₂ group of the chelate in order to form a precursor compound.

Example 4: ^{99m}Tc labelling of the precursors of Example 3

[0132] For ^{99m}Tc labelling, 50□g of a precursor compound will be added to a nitrogen filled vial and dissolved in 50□L water, 150□L of sodium gluconate solution (25mg in 6mL H₂O), 100□L ammonium acetate (pH 4.0, 50 mM), 1 mL TcO₄ soln (500 MBq) and 50□L SnCl₂ soln (20 mg in 100 mL H₂O). The mixture will be heated at 75 °C for 20 minutes before analysis by ITLC and HPLC.


Example 5: Preparation of (\pm)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid N-1-[^{18}F]-propyl-N-propylamide [Compound 12]*(i) Preparation of (\pm)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid propylamide***[0133]**

[0134] This compound can be made using the synthetic route described for the dipropylamide version (compound 12f of Okubo et al Bioorg. Med. Chem. 2004 12 3569-3580) replacing dipropylamine with a N-propyl-N-propyltosylate amine.

*(ii) Preparation of (\pm)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid N-1-[^{18}F]-propyl-N-propylamide***[0135]**

[0136] To 5.1.4. (\pm)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid propylamide in a suitable solvent (acetonitrile, DMSO, DMF, THF, Dioxane) is added a base (LDA, NaH, or similar). To the mixture is added ^{18}F fluoropropyl bromide (or other leaving group like tosylate etc) and the mixture is heated to 50-100°C for 5-30 min, followed by H PLC purification. Protection of the indole amine is required during radiofluorination.


Example 6: Preparation of (\pm)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-[^{11}C]carboxylic acid dipropylamide**[0137]**



[0138] 6,11-dihydro-5-thia-11-aza-benzo[α]fluorene is deprotonated using a strong base, such as potassium carbonate, and the resultant deprotonated intermediate is reacted with 1mM [¹¹C]CO₂ to produce 5.1.4. (±)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-[¹¹C]-carboxylic acid. Reaction of this reagent with a coupling reagent [such as described by Christensen Molecules 2001; 6; pp47-51] to produce an activated ester or mixed anhydride followed by addition of dipropylamine to provide the title compound.

Example 7: Preparation of 11-methyl-6,11-dihydro-5-thia-11-azabenz[α]fluorene-6-carboxylic acid diethylamide (non-radioactive Compound 3)

[0139]



[0140] This method is an adaption of "Method D" described by Okubo *et al, supra*.


[0141] 100mg (0.3mmol) of 6,11-Dihydro-5-thia-11-azo-benzo[α]fluorene-6-carboxylic acid diethylamide (synthesis described in Example 1) was dissolved in 3ml of DMSO. 84mg (1.5mmol) potassium hydroxide was added and the reaction mixture stirred at room temperature for 30 minutes. 0.04ml (0.6mmol) methyl iodide was added dropwise at room temperature and the reaction mixture was stirred for 1 hour. The reaction was quenched with 20ml water, and extracted with 2 x 20ml ether. The organic was dried (MgSO₄) and removed *in vacuo*. The residue was purified by column chromatography over silica using a 3:7 mixture of ethyl acetate and hexane as eluant. Appropriate fractions were combined and the solvent removed *in vacuo* to give the title compound as a yellow solid.

HPLC: 94.3%

500 MHz ¹H-NMR (CDCl₃) δ (ppm): 1.10 (3H, m), 1.34 (3H, m), 3.39-3.70 (4H, m), 4.01 (3H, s), 5.14 (1H, s), 7.12-7.7.41 (7H, m), 7.72 (1H, d).

Example 8: Preparation of 11-isobutyl-6,11-dihydro-5-thia-11-azabenz[α]fluorene-6-carboxylic acid diethylamide (non-radioactive Compound 4)

[0142]




[0143] The method of Example 7 was used but replacing methyl iodide with 1-bromo-2-methylpropane in order to obtain the isobutyl derivative. The product was obtained as a white solid.

HPLC 91.2%

500 MHz $^1\text{H-NMR}$ (CDCl_3) δ (ppm): 0.76 (6H, dd), 1.10 (3H, m), 1.34 (3H, m), 2.16 (1H, m), 3.39-3.70 (4H, m), 4.24-4.38 (2H, m), 7.12-7.741 (7H, m), 7.72 (1H, d).

Example 9: Preparation of 11-isopropyl-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (non-radioactive Compound 5)

[0144]




[0145] The method of Example 7 was used but replacing methyl iodide with 1-bromopropane in order to obtain the isopropyl derivative. The product was obtained as a white solid.

HPLC 90% Yield 44%

500 MHz $^1\text{H-NMR}$ (CDCl_3) δ (ppm): 1.10 (3H, m), 1.34 (3H, m), 1.60 (3H, m), 1.88 (3H, m), 3.39-3.70 (4H, m), 5.07 (1H, s), 5.11 (1H, m), 7.10-7.60 (8H, m).

Example 10: Preparation of 11-prop-2-ynyl-6,11-dihydro-5-thia-11-azobenzofluorene-6-carboxylic acid diethylamide (non-radioactive Compound 6/Compound 9P²)

[0146]




[0147] The method of Example 7 was used but replacing methyl iodide with propargyl bromide 80% in toluene in order to obtain the prop2ynyl derivative. The product was obtained as a yellow solid.

HPLC 99.3%

500 MHz ^1H -NMR (CDCl_3) δ (ppm): 1.10 (3H, m), 1.34 (3H, m), 2.50 (1H, s), 3.39-3.70 (4H, m), 4.91-5.06 (2H, m), 5.20 (1H, s), 7.11-7.50 (7H, m), 7.89 (1H, d).

Example 11: Preparation of 11-(2-methyl-allyl)-6,11-dihydro-5-thia-11-azabenzofluorene-6-carboxylic acid diethylamide (non-radioactive Compound 7)

[0148]




[0149] The method of Example 7 was used but replacing methyl iodide with 3-bromo-2-methylpropene in order to obtain the 2-methyl-allyl derivative. The product was obtained as a yellow solid.

HPLC 99.2% Yield 41%

500 MHz ^1H -NMR (CDCl_3) δ (ppm): 1.10 (3H, m), 1.34 (3H, m), 1.91 (3H, s), 3.31-3.66 (4H, m), 4.61-4.82 (3H, m), 5.08 (1H, s), 5.21 (1H, s), 7.12-7.53 (8H, m).

Example 12: Preparation of 11-(4-fluoro-butyl)-6,11-dihydro-5-thia-11-azabenzofluorene-6-carboxylic acid diethylamide (non-radioactive Compound 8)

[0150]




[0151] The method of Example 7 was used but replacing methyl iodide with bromofluorobutane in order to obtain the 4-fluoro-butyl derivative. The product was obtained as a yellow solid.

HPLC 96% Yield 33%

500 MHz $^1\text{H-NMR}$ (CDCl_3) δ (ppm): 1.10 (3H, m), 1.34 (3H, m), 1.78 (2H, m), 2.08 (1H, m), 3.31-3.66 (4H, m), 4.40-4.50 (4H, m), 5.10 (1H, s), 7.11-7.50 (7H, m), 7.89 (1H, d).

Example 13: Preparation of 11-[1-(2-fluoro-ethyl)-1H-[1,2,3]triazol-4-ylmethyl]-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (nonradioactive Compound 9)

[0152]




[0153] 1-Azido-2-fluoro ethane was synthesized by adding to a solution of fluoroethyltosylate (100mg, 0.46mmol) in anhydrous dimethylformamide 2ml sodium azide (90mg, 1.4mmol) at room temperature under nitrogen. The mixture was stirred for 24 hours. The product was prepared as a solution in dimethylformamide after filtering off the solid. The solution was used without further purification to the next step.

[0154] To a solution of copper (II) sulfate pentahydrate (1.6mg, 0.007mmol) and L-ascorbic acid (2.5mg, 0.014mmol) in water 0.1ml was added a solution of non-radioactive Compound 9P² as prepared in Example 10 (50mg, 0.134mmol) in anhydrous dimethylformamide 1ml and a solution of 1-Azido-2-fluoro ethane in dimethylformamide (0.16mmol) at room temperature under nitrogen. The reaction mixture was stirred at 90 °C for 10 hours. After cooling, the reaction was then quenched with 20ml water, extracted with dichloromethane 2 x 20ml. The organic was dried (MgSO_4) and removed *in vacuo*. The residue was purified by column chromatography over silica using a 3:7 mixture of ethyl acetate and hexane as eluant. Appropriate fractions were combined and the solvent was removed *in vacuo* to give the product as a yellow solid. 500 MHz $^1\text{H-NMR}$ (CDCl_3) δ (ppm): 1.12 (3H, m), 1.40 (3H, m), 2.19 (2H, s), 3.38-3.60 (4H, m), 5.12 (1H, s), 5.45 (2H,

m), 5.56 (2H, m), 7.12-7.28 (5H, m), 7.44 (3H, m).
 MS: m/z = 462 (M^+) found (calculated mass 462)

Example 14: Preparation of 11-(2-fluoro-ethyl)-6, 11-dihydro-5-thia-11-azabenz[α]fluorene-6-carboxylic acid diethylamide [non-radioactive Compound 10]

[0155]



Synthesis of acetyl-2-(2-hydroxyethyl)-2-phenylhydrazine (i) :

[0156] To a solution of 1-acetyl-2-phenylhydrazine (2g, 13.3mmol) in anhydrous toluene 20ml was added bromoethanol (1.04ml, 14.6mmol), diisopropylethylamine (2.54ml, 14.6ml) at room temperature. The reaction mixture was refluxed under nitrogen for 40 hours. After cooling, the reaction was then quenched with 100ml water, extracted with dichloromethane 2 x 100ml. The organic was dried ($MgSO_4$) and removed *in vacuo*. The residue was purified by flash column chromatography over silica using a gradient 50% mixture of ethyl acetate and hexane to 100% ethyl acetate as eluant. Appropriate fractions were combined and the solvent was removed *in vacuo* to give the product as a yellow oil 0.9g (Yield 35%).
 GC/MS: m/z = 194 (M^+).

Synthesis of 2-(N-phenyl-hydrazino)-ethanol (ii) :

[0157] A solution of acetyl-2-12-hydroxyethyl-2-phenylhydrazine (i) (0.5g, 2.57mmol) in 6N hydrogen chloride 5ml was refluxed for 2 hours. The reaction was cooled to room temperature and basified with 6N sodium hydroxide to PH 8. The solution was extracted with DCM 2 x 50ml, washed with brine 2 x 30ml. The organic was dried ($MgSO_4$) and removed *in vacuo*. The residue was used to next step without further purification.
 GC/MS: m/z = 152 (M^+).

Synthesis of 11-(2-Hydroxy-ethyl)-6, 11-dihydro-5-thia-11-azo-benzo[α]fluorene-6-carboxylic acid diethylamide (iv):


[0158] To a solution of 2-(N-phenyl-hydrazino)-ethanol (ii) (0.59g, 1.9mmol) and 4-oxo-thiochroman-2-carboxylic acid diethylamide (iii) (0.5g, 1.9mmol) in ethanol 10ml was added conc. sulfuric acid 0.5ml. The reaction mixture was stirred and refluxed for 20 hours. It was then cooled to room temperature and poured onto water 50ml. The resulting precipitate was collected by filtration and washed with water (3 x 30ml). It was then dried in desiccators under vacuum with P₂O₅ as the drying agent to give the product 11-(2-Hydroxy-ethyl)-6,11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (iv) as a yellow solid 0.62g, which was used to next step without further purification. Yield: 86%. 500 MHz ¹H-NMR (CDCl₃) δ (ppm): 1.12 (3H, m), 1.40 (3H, m), 3.38-3.70 (4H, m), 4.05-4.15 (2H, m), 4.40 (1H, m), 4.60 (1H, m), 5.31 (1H, s), 7.12-7.47 (7H, m), 7.90 (1H, d).

Synthesis of 11-(2-Tosyloxy-ethyl)-6,11-dihydro-5-thio-11-azo-benzo[α]fluorene-6-carboxylic acid diethylamide (Compound 10P):

[0159] To a solution of 11-(2-Hydroxy-ethyl)-6, 11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (iv) (100mg, 0.26mmol) in anhydrous dichloromethane 3ml was added toluenesulfonyl chloride (100mg, 0.52mmol) and pyridine (0.2ml, 2.6mmol) at 0 °C under nitrogen. The reaction mixture was then stirred at room temperature for 48 hours. After quenched with 20ml water, the mixture was extracted with dichloromethane 2 x 20ml. The organic was dried (MgSO₄) and removed *in vacuo*. The residue was purified by flash column chromatography over silica using a gradient 20% mixture of ethyl acetate and hexane to 40% ethyl acetate as eluant. Appropriate fractions were combined and the solvent was removed *in vacuo* to give the product 11-(2-Tosyloxy-ethyl)-6, 11-dihydro-5-thia-11-azo-benzo[α]fluorene-6-carboxylic acid diethylamide (Compound 10P) as a white solid 98mg. Yield 48%. 500 MHz ¹H-NMR (CDCl₃) δ (ppm): 1.12 (3H, m), 1.40 (3H, m), 2.12 (3H, s), 3.38-3.60 (4H, m), 4.40-4.71 (4H, m), 5.02 (1H, s), 7.12-7.27 (7H, m), 7.38 (1H, d), 7.42 (1H, d), 7.53 (1H, d), 7.7 5 (2 H, d).

Synthesis of 11-(2-fluoroethyl)-6, 11-dihydro-5-thio-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (non-radioactive Compound 10):


[0160] To a solution of 11-(2-Tosyloxy-ethyl)-6, 11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (Compound 10P) (100mg, 0.19mmol) in anhydrous acetonitrile 5ml was added tetrabutylammonium fluoride 1.0 M in tetrahydrofuran (0.4ml) at room temperature under nitrogen. The reaction was stirred for 3 hours. After quenched with 20ml water, the mixture was extracted with dichloromethane 2 x 20ml. The organic was dried (MgSO₄) and removed *in vacuo*. The residue was purified by flash column chromatography over silica using a gradient 20% mixture of ethyl acetate and hexane to 40% ethyl acetate as eluant. Appropriate fractions were combined and the solvent was removed *in vacuo* to give the product 11-(2-fluoroethyl)-6, 11-dihydro-5-thia-11-aza-benzo[α]fluorene-6-carboxylic acid diethylamide (non-radioactive Compound 10) as a yellow solid 41mg. Yield 55%. 500 MHz ¹H-NMR (CDCl₃) δ (ppm): 1.12 (3H, m), 1.40 (3H, m), 3.38-3.60 (4H, m), 4.56-4.78 (2H, m), 4.85-4.98 (2H, m), 5.12 (1H, s), 7.12-7.48 (7H, m), 7.64 (1H, d). MS: m/z 383 (M⁺) found (calculated mass 383)

Example 15: Preparation of 11-(2-[¹⁸F]Fluoroethyl)-6,11-dihydro-5-thia-11-azabenzofluorene-6-carboxylic acid diethylamide [Compound 10]**[0161]**

[0162] Anhydrous potassium [^{18}F]fluoride was prepared by azeotropic drying of aqueous [^{18}F]fluoride in the presence of potassium carbonate (0.3mg) and Kryptofix® 2.2.2. (1.8mg) and anhydrous acetonitrile (3 x 0.5ml). A solution of 11-(2-Tosyloxyethyl)-6,11-dihydro-5-thia-11-aza-benzo[a]fluorene-6-carboxylic acid diethylamide (1mg; Compound 10P) in anhydrous DMSO (0.4 mL) was added to the vial containing anhydrous Potassium-Kryptofix® 2.2.2. [^{18}F]fluoride complex and the reaction vial sealed and heated to 130°C with rapid stirring for 10 min. The resulting reaction mixture was cooled to room temperature, then diluted with 0.5 mL of HPLC mobile phase (acetonitrile:water, 60:40). An aliquot of the resulting solution was injected onto a analytical HPLC column (Phenomenex Luna 5u, C18(2) 100A 150mm x 4.6mm) and eluted with a flow rate of 1mL/min. The title compound was eluted from the column with a retention time of 8.5 minutes with a labelling yield of 50% by HPLC integration. The identity of the product was confirmed by co-elution with an authentic, unlabelled sample of 11-(2-Fluoroethyl)-6,11-dihydro-5-thia-11-azabenz[a]fluorene-6-carboxylic acid diethylamide.

Example 16: Preparation of 11-(2-methoxy-ethyl)-6,11-dihydro-5-thia-11-azabenz[a]fluorene-6-carboxylic acid diethylamide (non-radioactive Compound 11)


[0163]



[0164] To a stirred solution of intermediate iv (from the synthesis described in Example 15) (60mg, 0.158mmol) in anhydrous DMSO 3ml was added potassium hydroxide (84mg, 1.58mmol) and iodomethane (1.58mmol, 0.12ml) at room temperature. The mixture was then stirred under nitrogen for 30 minutes. The reaction was poured onto water 50ml, extracted with dichloromethane 2 x 20ml, washed with brine 2 x 20ml. The organic was dried (MgSO_4) and removed *in vacuo*. The residue was recrystallised from ethylacetate/hexane. The product was filtered and dried under vacuum to give a yellow solid 34mg. Yield: 55%.

Example 17: Synthesis of 11-(2-[^{11}C]methoxy-ethyl)-6,11-dihydro-5-thia-11-azabenz[a]fluorene-6-carboxylic acid diethylamide [Compound 11]

[0165]



[0166] [^{11}C]Methyl iodide is trapped in a stirred solution of 11-(2-hydroxy-ethyl)-6,11-dihydro-5-thia-11-aza-benzo[a]fluorene-6-carboxylic acid diethylamide (intermediate iv of Example 14) and a suitable base (e.g. potassium hydroxide, 10 equivalents) in anhydrous DMSO (0.4 mL). The reaction mixture is stirred rapidly at room temperature or elevated temperature to effect the [^{11}C]methylation reaction. After consumption of the [^{11}C]methyl iodide the reaction mixture is cooled to room temperature and purified by HPLC chromatography.

Example 18: Screening method for Compounds of the Invention


[0167] The compounds were screened for their affinity for PBR using a method adapted from Le Fur et al 1983 [Life Sci. USA 33 pp449-57].

[0168] The compounds to be tested [dissolved in 50mM Tris-HCl, pH 7.4, 10mM MgCl₂ containing 1%DMSO] competed for binding to Wistar rat heart PBR to the receptor against 0.3 nM [³H] PK-11195. The reaction was carried out in 50mM Tris-HCl, pH 7.4 10mM MgCl₂ for 15 minutes at 25°C.

[0169] K_i values for the best compounds tested were found to be between 1.0nM and 0.1nM.

Claims

1. A compound of Formula I:



or a salt or solvate thereof, wherein said compound is labelled with an imaging moiety, and wherein:

X¹ and X² are independently selected from hydrogen, halogen, C₁₋₆ alkoxy, C₁₋₆ alkyl;

R¹ and R² are independently selected from C₁₋₆ alkyl, C₁₋₆ aminoalkyl, C₁₋₆ alkoxyalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, a polyethylene glycol (PEG) group, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloethers, and C₃₋₁₀ cycloamines; R³ is the group -A-R⁵ wherein:

A is the optional group -(CH₂)_z-R⁶- wherein z=0-6, R⁶ is a 5- or 6-membered heterocycle having 1-3 heteroatoms selected from N, S and O, and wherein R⁵ is a substituent selected from hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkenyl, C₁₆ alkynyl, C₁₋₆ alkanoyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkenyl, C₁₋₆ haloalkynyl, C₁₋₆ haloalkoxy, C₁₋₆ haloalkylthiol, C₁₋₆ haloalkylsulfinyl, C₁₋₆ haloalkylsulfonyl, C₁₋₆ haloalkyl ketone, C₁₋₆ haloalkyl sulfinyl, C₁₋₆ haloalkyl sulfonyl, a polyethylene glycol (PEG) group, C₁₋₆ hydroxyalkyl, a nitrogen-containing C₂₋₁₀ alkyl and hydroxy;

R⁴ is hydrogen, C₁₋₆ alkyl, C₁₋₆ alkanoyl, C₁₋₆ cycloalkyl, C₁₋₆ fluoroalkyl, hydroxy, or halogen;

Y¹ is S, SO, SO₂, or CH₂;

wherein said imaging moiety is selected from:

- (i) a radioactive metal ion;
- (ii) a paramagnetic metal ion;
- (iii) a gamma-emitting radioactive halogen;
- (iv) a positron-emitting radioactive non-metal;
- (v) a hyperpolarised NMR-active nucleus;
- (vi) a reporter suitable for *in vivo* optical imaging; and,
- (vii) a β-emitter suitable for intravascular detection.

2. The compound of claim 1 wherein:

X¹ and X² are both hydrogen;

R¹ and R² are independently selected from C₁₋₆ alkyl, C₁₋₆ methoxyalkyl or C₁₋₆ alkoxy;

R³ is hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkanoyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, or C₁₆ fluoroalkyl, or is A-C₁₋₆ alkyl, or A-C₁₋₆ fluoroalkyl wherein A is as defined previously;

R⁴ is hydrogen, C₁₋₆ alkyl, C₁₋₆ acetyl, C₁₋₆ cycloalkyl or C₁₋₆ fluoroalkyl;
Y¹ is S, SO₂, or CH₂; and,
n is 0.


3. The compound of claim 2 wherein:

R¹ and R² are both C₁₋₆ alkyl;
R³ is hydrogen, C₁₋₃ alkyl, C₁₋₃ alkoxy, C₁₋₃ alkanoyl, C₁₋₃ alkenyl, C₁₋₃ alkynyl, or C₁₋₆ fluoroalkyl, or is A-C₁₋₆ alkyl or A-C₁₋₆ fluoroalkyl wherein for A, R⁶ is a 5- or 6-membered N-containing heterocycle;
R⁴ is hydrogen, C₁₋₆ alkyl or C₁₋₆ acetyl; and,
Y¹ is S or SO₂.

4. The compound of claim 3 wherein:

R¹ and R² are both ethyl;
R³ is hydrogen, methyl, ethyl, 2-methoxy-ethyl, prop-2-ynyl, isopropyl, isobutyl, 2-methyl-allyl, acetyl, or 4-fluorobutyl, or is or is A-C₁₋₆ fluoroalkyl wherein for A, R⁶ is a 5-membered N-containing heterocycle;
R⁴ is hydrogen; and,
Y¹ is S.


5. A compound of Formula II:



or a salt or solvate thereof wherein:

R⁷ is as defined in claims 1-4 for R³, with the proviso that R⁷ is not hydrogen, C₁₋₅ alkyl or a nitrogen-containing C₂₋₁₀ alkyl group; and,
Y² is as defined for Y¹ in claims 1-4.

6. A method for the preparation of the compound of any of claims 1-4 comprising reaction of a convenient chemical form of an imaging moiety with a precursor of Formula Ia:



wherein:

X¹¹, X¹², R¹³, R¹⁴ and Y¹¹ are as defined for X¹, X², R³, R⁴ and Y¹¹, respectively, of Formula I of claims 1-4, or independently comprise a suitable protecting group;

R^{15} is the group $-(CH_2)_o-C(=O)-NR^{11}R^{12}$ wherein o , R^{11} and R^{12} are as defined for n , R^1 and R^2 for Formula I of claims 1-4, or comprises a suitable protecting group; and,

R^{16} is hydrogen,

and provided that at least one of X^{11} , X^{12} and $R^{11}-R^{16}$ comprises a chemical group capable of reacting with a suitable source of said imaging moiety wherein:


(i) when said imaging moiety is a metal ion, said chemical group is a chelating agent where 2-6 donor atoms selected from amine, thiol, amide, oxime and phosphine are arranged such that 5- or 6-membered chelate rings result, or is a monodentate ligand comprising donor atoms selected from isonitrile, phosphine and diazenide;

(ii) when said imaging moiety is radioiodine, said chemical group is a trialkylstannane, a trialkylsilane, an organoboron compound, alkyl bromide, alkyl tosylate, alkyl mesylate, alkyl triflate, phenol, aryl iodonium salt, aryl diazonium, aryl trialkylammonium or nitroaryl;

(iii) when said imaging moiety is a radioactive isotope of fluorine, said chemical group is alkyl bromide, alkyl tosylate, alkyl mesylate, hydroxyl, N-haloacetyl, aryl diazonium salt, aryl nitro, aryl quaternary ammonium salt, chloro, $P(O)Ph_3$ or an activated ester;

(iv) when said imaging moiety is ^{11}C , said chemical group is hydrogen, halide, boronic acid, triflate or organotin.

7. A precursor for the preparation of the compound of any of claims 1-4 wherein said precursor is selected from:



8. A pharmaceutical composition which comprises the compound of any of claims 1-5, together with a biocompatible carrier in a form suitable for mammalian administration.

9. The pharmaceutical composition of claim 8 which is an *in vivo* imaging agent comprising the compound of any of claims 1-4.

10. The pharmaceutical composition of claim 8 which is a therapeutic composition comprising the compound of claim 5.

11. A kit comprising the precursor of claim 7 wherein said kit is suitable for the preparation of the pharmaceutical composition of claim 8.

12. A compound of any of claims 1-4 for use in an *in vivo* diagnostic or imaging method.

13. Use of the compound of any of claims 1-4 in the manufacture of a pharmaceutical for the *in vivo* diagnosis or imaging of a PBR condition.

14. A compound of any of claims 1-4 for use in a method of monitoring the effect of treatment of a human or animal body with a drug to combat a PBR condition, said method comprising administering to said body the pharmaceutical composition of claim 9, and detecting the uptake of said pharmaceutical composition.

5


Patentansprüche

1. Verbindung der Formel I:

10

15

20



(I)

25

oder ein Salz oder Solvat derselben, wobei die Verbindung mit einer bildgebenden Einheit markiert ist und wobei

X¹ und X² unabhängig voneinander aus Wasserstoff, Halogen, C₁₋₆-Alkoxy, C₁₋₆-Alkyl ausgewählt sind;
 R¹ und R² unabhängig voneinander aus C₁₋₆-Alkyl, C₁₋₆-Aminoalkyl, C₁₋₆-Alkoxyalkyl, C₁₋₆-Alkoxy, C₁₋₆-Hydroxyalkyl, einer Polyethylenglykol(PEG)-Gruppe, C₃₋₁₀-Cycloalkyl, C₃₋₁₀-Cycloethern und C₃₋₁₀-Cycloaminen ausgewählt sind;
 R³ für die Gruppe -A-R⁵ steht, wobei:

30

35

A für die optionale Gruppe -(CH₂)_z-R⁶- steht, wobei z = 0 bis 6, R⁶ ein 5- oder 6-gliedriger Heterozyklus mit 1 bis 3 Heteroatomen ist, die aus N, S und O ausgewählt sind, und wobei R⁵ ein Substituent ist, der aus Wasserstoff, C₁₋₆-Alkyl, C₁₋₆-Alkoxy, C₁₋₆-Alkenyl, C₁₋₆-Alkinyl, C₁₋₆-Alkanoyl, C₁₋₆-Halogenalkyl, C₁₋₆-Halogenalkenyl, C₁₋₆-Halogenalkinyl, C₁₋₆-Halogenalkoxy, C₁₋₆-Halogenalkylthiol, C₁₋₆-Halogenalkylsulfanyl, C₁₋₆-Halogenalkylsulfonyl, C₁₋₆-Halogenalkylketon, C₁₋₆-Halogenalkylsulfinyl, C₁₋₆-Halogenalkylsulfonyl, einer Polyethylenglykol (PEG) -Gruppe, C₁₋₆-Hydroxyalkyl, einem stickstoffhaltigen C₂₋₁₀-Alkyl und Hydroxy ausgewählt ist;

40

R⁴ für Wasserstoff, C₁₋₆-Alkyl, C₁₋₆-Alkanoyl, C₁₋₆-Cycloalkyl, C₁₋₆-Fluoralkyl, Hydroxy oder Halogen steht;
 Y¹ für S, SO, SO₂ oder CH₂ steht;
 n 0 bis 10 ist und

45

wobei die bildgebende Einheit aus dem Folgenden ausgewählt ist:

- (i) einem radioaktiven Metallion;
- (ii) einem paramagnetischen Metallion;
- (iii) einem gamma-emittierenden radioaktiven Halogen;
- (iv) einem positron-emittierenden radioaktiven Nichtmetall;
- (v) einem hyperpolarisierten NMR-aktiven Atomkern;
- (vi) einem für die optische *In-vivo*-Bildgebung geeigneten Reporter und
- (vii) einem β-Emitter, der für die intravaskuläre Erfassung geeignet ist.

50

55

2. Verbindung nach Anspruch 1, wobei:

X¹ und X² beide für Wasserstoff stehen;
 R¹ und R² unabhängig voneinander aus C₁₋₆-Alkyl, C₁₋₆-Methoxyalkyl und C₁₋₆-Alkoxy ausgewählt sind;

R³ für Wasserstoff, C₁₋₆-Alkyl, C₁₋₆-Alkoxy, C₁₋₆-Alkanoyl, C₁₋₆-Alkenyl, C₁₋₆-Alkynyl oder C₁₋₆-Fluoralkyl oder für A-C₁₋₆-Alkyl oder A-C₁₋₆-Fluoralkyl steht, wobei A wie vorstehend definiert ist;
 R⁴ für Wasserstoff, C₁₋₆-Alkyl, C₁₋₆-Acetyl, C₁₋₆-Cycloalkyl oder C₁₋₆-Fluoralkyl steht;
 Y¹ für S, SO₂ oder CH₂ steht und
 n 0 ist.


3. Verbindung nach Anspruch 2, wobei:

R¹ und R² beide für C₁₋₆-Alkyl stehen;
 R³ für Wasserstoff, C₁₋₃-Alkyl, C₁₋₃-Alkoxy, C₁₋₃-Alkanoyl, C₁₋₃-Alkenyl, C₁₋₃-Alkynyl oder C₁₋₆-Fluoralkyl oder für A-C₁₋₆-Alkyl oder A-C₁₋₆-Fluoralkyl steht, wobei für A R⁶ für einen 5- oder 6-gliedrigen N-haltigen Heterozyklus steht;
 R⁴ für Wasserstoff, C₁₋₆-Alkyl oder C₁₋₆-Acetyl steht und
 Y¹ für S oder SO₂ steht.

4. Verbindung nach Anspruch 3, wobei


R¹ und R² beide für Ethyl stehen;
 R³ für Wasserstoff, Methyl, Ethyl, 2-Methoxyethyl, Prop-2-ynyl, Isopropyl, Isobutyl, 2-Methylallyl, Acetyl oder 4-Fluorbutyl oder A-C₁₋₆-Fluoralkyl steht, wobei für A R⁶ für einen 5-gliedrigen N-haltigen Heterozyklus steht;
 R⁴ für Wasserstoff steht und
 Y¹ für S steht.

5. Verbindung der Formel II:



oder ein Salz oder Solvat derselben, wobei
 R⁷ wie in Ansprüchen 1 bis 4 für R³ definiert ist, mit der Vorgabe, dass R⁷ nicht für Wasserstoff, C₁₋₅-Alkyl oder eine stickstoffhaltige C₂₋₁₀-Alkylgruppe steht; und
 Y² wie in Ansprüchen 1 bis 4 für Y¹ definiert ist.

6. Verfahren zur Herstellung der Verbindung nach einem der Ansprüche 1 bis 4, umfassend die Reaktion einer geeigneten chemischen Form einer bildgebenden Einheit mit einer Vorstufe der Formel Ia:



wobei:

X^{11} , X^{12} , R^{13} , R^{14} und Y^{11} wie für X^1 , X^2 , R^3 , R^4 bzw. Y^{11} der Formel I in Ansprüche 1 bis 4 definiert sind oder unabhängig eine geeignete Schutzgruppe umfassen;

R^{15} für die Gruppe $-(CH_2)_o-C(=O)-NR^{11}R^{12}$ steht, wobei o , R^{11} und R^{12} wie für n , R^1 und R^2 der Formel I in Ansprüche 1 bis 4 definiert sind oder eine geeignete Schutzgruppe umfassen; und

R^{16} für Wasserstoff steht,

und unter der Voraussetzung, dass mindestens eine Gruppe aus X^{11} , X^{12} und R^{11} bis R^{16} eine chemische Gruppe umfasst, die in der Lage ist, mit einer geeigneten Quelle der bildgebenden Einheit zu reagieren, wobei:


(i) wenn die bildgebende Einheit ein Metallion ist, die chemische Gruppe ein Chelatbildner ist, wobei 2 bis 6 Donoratome, ausgewählt aus Amin, Thiol, Amid, Oxim und Phosphan, so angeordnet sind, dass 5- oder 6-gliedrige Chelatringe entstehen, oder ein Monodentatligand ist, welcher Donoratome umfasst, die aus Isonitril, Phosphan und Diazenid ausgewählt sind;

(ii) wenn die bildgebende Einheit radioaktives Iod ist, die chemische Gruppe ein Trialkylstannan, ein Trialkylsilan, eine Organoboranverbindung, Alkylbromid, Alkyltosylat, Alkylmesylat, Alkyltriflat, Phenol, ein Aryliodoniumsalz, Aryldiazonium, Aryltrialkylammonium oder Nitroaryl ist;

(iii) wenn die bildgebende Einheit ein radioaktives Fluorisotop ist, die chemische Gruppe Alkylbromid, Alkyltosylat, Alkylmesylat, Hydroxyl, N-Halogenacetyl, ein Aryldiazoniumsalz, Arylnitro, ein quartäres Arylammoniumsalz, Chlor, $P(O)Ph_3$ oder ein aktivierter Ester ist;

(iv) wenn die bildgebende Einheit ^{11}C ist, die chemische Gruppe Wasserstoff, Halogenid, Boronsäure, Triflat oder eine zinnhaltige organische Gruppe ist.


7. Vorstufe für die Herstellung der Verbindung nach einem der Ansprüche 1 bis 4, wobei die Vorstufe aus den folgenden ausgewählt ist:



8. Pharmazeutische Zusammensetzung, welche die Verbindung nach einem der Ansprüche 1 bis 5 zusammen mit einem biokompatiblen Träger in einer Form umfasst, die für die Verabreichung an ein Säugetier geeignet ist.
9. Pharmazeutische Zusammensetzung nach Anspruch 8, welche ein *in-vivo*-bildgebendes Mittel ist, das die Verbindung nach einem der Ansprüche 1 bis 4 umfasst.
10. Pharmazeutische Zusammensetzung nach Anspruch 8, welche eine therapeutische Zusammensetzung ist, die die Verbindung nach Anspruch 5 umfasst.
11. Kit, welches die Vorstufe nach Anspruch 7 umfasst, wobei das Kit für die Herstellung der pharmazeutischen Zusammensetzung nach Anspruch 8 geeignet ist.
12. Verbindung nach einem der Ansprüche 1 bis 4 zur Verwendung in einem diagnostischen oder bildgebenden *In-vivo*-Verfahren.
13. Verwendung der Verbindung nach einem der Ansprüche 1 bis 4 bei der Herstellung eines Pharmazeutikums für die *In-vivo*-Diagnose oder Bildgebung eines PBR-Zustands.
14. Verbindung nach einem der Ansprüche 1 bis 4 zur Verwendung in einem Verfahren zur Überwachung der Wirkung einer Behandlung eines menschlichen Körpers oder eines Tierkörpers mit einem Medikament zur Bekämpfung eines PBR-Zustands, wobei das Verfahren das Verabreichen der pharmazeutischen Zusammensetzung nach Anspruch 9 an den Körper und das Erfassen der Aufnahme der pharmazeutischen Zusammensetzung umfasst.

Revendications

1. Composé de Formule I :



ou sel ou solvate de celui-ci, dans lequel ledit composé est marqué avec un fragment pour imagerie, et dans lequel :

X¹ et X² sont indépendamment sélectionnés parmi l'hydrogène, un halogène, un groupe alcoxy en C₁₋₆, alkyle en C₁₋₆ ;

R¹ et R² sont indépendamment sélectionnés parmi un groupe alkyle en C₁₋₆, aminoalkyle en C₁₋₆, alcoxyalkyle en C₁₋₆, alcoxy en C₁₋₆, hydroxyalkyle en C₁₋₆, un groupe polyéthylène glycol (PEG), cycloalkyle en C₃₋₁₀, des cycloéthers en C₃₋₁₀ et des cycloamines en C₃₋₁₀ ;

R³ est le groupe -A-R⁵ dans lequel :

A est le groupe facultatif - (CH₂)_z-R⁶- dans lequel z = 0 à 6, R⁶ est un composé hétérocyclique à 5 ou à 6 éléments ayant 1 à 3 hétéroatomes sélectionnés parmi N, S et O, et dans lequel R⁵ est un substituant sélectionné parmi l'hydrogène, un groupe alkyle en C₁₋₆, alcoxy en C₁₋₆, alcényle en C₁₋₆, alcynyle en C₁₋₆, alcanoyl en C₁₋₆, halogénoalkyle en C₁₋₆, halogénoalcényle en C₁₋₆, halogénoalcynyle en C₁₋₆, halogénoalcoxy en C₁₋₆, halogénoalkylthiol en C₁₋₆, halogénoalkylsulfinyle en C₁₋₆, halogénoalkylsulfonyl en C₁₋₆, halogénoalkyle cétone en C₁₋₆, halogénoalkyle sulfinyle en C₁₋₆, halogénoalkyle sulfonyl en C₁₋₆, un groupe polyéthylène glycol (PEG), hydroxyalkyle en C₁₋₆, un groupe alkyle en C₂₋₁₀ contenant de l'azote et hydroxy ;

R⁴ est l'hydrogène, un groupe alkyle en C₁₋₆, alcanoyl en C₁₋₆, cycloalkyle en C₁₋₆, fluoroalkyle en C₁₋₆, hydroxy, ou un halogène ;

Y¹ est S, SO, SO₂, ou CH₂ ;

n est 0 à 10 ; et

dans lequel ledit fragment pour imagerie est sélectionné parmi :

- (i) un ion métallique radioactif ;
- (ii) un ion métallique paramagnétique ;
- (iii) un halogène radioactif émettant des gammas ;
- (iv) un métalloïde radioactif émettant des positrons ;
- (v) un noyau hyperpolarisé sensible à la RMN ;
- (vi) un rapporteur approprié pour l'imagerie optique *in vivo* ; et
- (vii) un β -émetteur approprié pour la détection intravasculaire.

2. Composé selon la revendication 1, dans lequel :

X¹ et X² sont tous les deux l'hydrogène ;

R¹ et R² sont indépendamment sélectionnés parmi un groupe alkyle en C₁₋₆, méthoxyalkyle en C₁₋₆ ou alcoxy en C₁₋₆ ;

R³ est l'hydrogène, un groupe alkyle en C₁₋₆, alcoxy en C₁₋₆, alcanoyl en C₁₋₆, alcényle en C₁₋₆, alcynyle en C₁₋₆, ou fluoroalkyle en C₁₋₆, ou est alkyle en A-C₁₋₆, ou fluoroalkyle en A-C₁₋₆ où A est tel que précédemment défini ;

R⁴ est l'hydrogène, un groupe alkyle en C₁₋₆, acétyl en C₁₋₆, cycloalkyle en C₁₋₆ ou fluoroalkyle en C₁₋₆ ;

Y¹ est S, SO₂, ou CH₂ ; et

n est 0.

3. Composé selon la revendication 2, dans lequel :

R¹ et R² sont tous les deux un groupe alkyle en C₁₋₆ ;

R³ est l'hydrogène, un groupe alkyle en C₁₋₃, alcoxy en C₁₋₃, alcanoyle en C₁₋₃, alcényle en C₁₋₃, alcynyle en C₁₋₃, ou fluoroalkyle en C₁₋₆, ou est un groupe alkyle en A-C₁₋₆ ou fluoroalkyle en A-C₁₋₆ où pour A, R⁶ est un composé hétérocyclique à 5 ou à 6 éléments contenant N ;

R⁴ est l'hydrogène, un groupe alkyle en C₁₋₆ ou acétyle en C₁₋₆ ; et

Y¹ est S ou SO₂.

4. Composé selon la revendication 3, dans lequel :


R¹ et R² sont tous les deux un groupe éthyle ;

R³ est l'hydrogène, un groupe méthyle, éthyle, 2-méthoxy-éthyle, prop-2-ynyle, isopropyle, isobutyle, 2-méthylallyle, acétyle, ou 4-fluorobutyle, ou est un groupe fluoroalkyle en A-C₁₋₆ où pour A, R⁶ est un composé hétérocyclique à 5 éléments contenant N ;

R⁴ est l'hydrogène ; et

Y¹ est S.

5. Composé de Formule II :




(II)

ou sel ou solvate de celui-ci dans lequel :

R⁷ est tel que défini dans les revendications 1 à 4 pour R³, avec la clause restrictive selon laquelle R⁷ n'est pas l'hydrogène, un groupe alkyle en C₁₋₅ ou un groupe alkyle en C₂₋₁₀ contenant de l'azote ; et,

Y² est tel que défini pour Y¹ dans les revendications 1 à 4.

6. Procédé pour la préparation du composé selon l'une quelconque des revendications 1 à 4 comprenant la réaction d'une forme chimique commode d'un fragment pour imagerie avec un précurseur de Formule Ia :



(Ia)

dans laquelle :

X¹¹, X¹², R¹³, R¹⁴ et Y¹¹ sont tels que respectivement définis pour X¹, X², R³, R⁴ et Y¹¹, de Formule I des revendications 1 à 4, ou comprennent indépendamment un groupe de protection approprié ;


R¹⁵ est le groupe - (CH₂)_o-C(=O)-NR¹¹R¹² dans lequel o, R¹¹ et R¹² sont tels que définis pour n, R¹ et R² pour la Formule I des revendications 1 à 4, ou comprend un groupe de protection approprié ; et,

R¹⁶ est l'hydrogène,

et à condition qu'au moins un de X^{11} , X^{12} et R^{11} à R^{16} comprenne un groupe chimique susceptible de réagir avec une source appropriée dudit fragment pour imagerie dans lequel :

- (i) lorsque ledit fragment pour imagerie est un ion métallique, ledit groupe chimique est un agent chélatant où 2 à 6 atomes donneurs sélectionnés parmi une amine, un thiol, un amide, une oxime et une phosphine sont agencés de sorte qu'il en résulte des anneaux chélate à 5 ou 6 éléments, ou est un ligand monodentate comprenant des atomes donneurs sélectionnés parmi un isonitrile, une phosphine et un diazénide ;
- (ii) lorsque ledit fragment pour imagerie est de l'iode radioactif, ledit groupe chimique est un trialkylstannane, un trialkylsilane, un composé organoboré, du bromure d'alkyle, du tosylate d'alkyle, du mésylate d'alkyle, du triflate d'alkyle, un phénol, un sel d'aryle iodonium, de l'aryle diazonium, de l'aryle trialkylammonium ou du nitroaryle ;
- (iii) lorsque ledit fragment pour imagerie est un isotope radioactif du fluor, ledit groupe chimique est du bromure d'alkyle, du tosylate d'alkyle, du mésylate d'alkyle, de l'hydroxyle, du N-haloacétyle, un sel d'aryle diazonium, du nitroaryle, un sel d'aryle ammonium quaternaire, du chloro, $P(O)Ph_3$ ou un ester activé ;
- (iv) lorsque ledit fragment pour imagerie est ^{11}C , ledit groupe chimique est l'hydrogène, un halogénure, l'acide borique, un triflate ou un organoétain.

7. Précurseur pour la préparation du composé selon l'une quelconque des revendications 1 à 4, dans lequel ledit précurseur est sélectionné à partir de :



Composé 8P Composé 9P1 Composé 9P2 Composé 10P

8. Composition pharmaceutique qui comprend le composé selon l'une quelconque des revendications 1 à 5, ainsi qu'un vecteur compatible d'un point de vue biologique sous une forme appropriée pour l'administration à un mammifère.
9. Composition pharmaceutique selon la revendication 8, qui est un agent d'imagerie *in vivo* comprenant le composé selon l'une quelconque des revendications 1 à 4.
10. Composition pharmaceutique selon la revendication 8 qui est une composition thérapeutique comprenant le composé selon la revendication 5.
11. Kit comprenant le précurseur selon la revendication 7, dans lequel ledit kit est approprié pour la préparation de la composition pharmaceutique selon la revendication 8.
12. Composé selon l'une quelconque des revendications 1 à 4 pour utilisation dans un procédé d'imagerie ou de diagnostic *in vivo*.

13. Utilisation du composé selon l'une quelconque des revendications 1 à 4 dans la fabrication d'une substance pharmaceutique pour le diagnostic ou l'imagerie *in vivo* d'une affection liée au PBR.

5 14. Composé selon l'une quelconque des revendications 1 à 4, pour une utilisation dans un procédé de contrôle de l'effet d'un traitement d'un corps humain ou animal avec un médicament pour combattre une affection liée au PBR, ledit procédé comprenant l'administration audit corps de la composition pharmaceutique selon la revendication 9, et la détection de l'assimilation de ladite composition pharmaceutique.

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6451795 B [0003]
- JP 07165721 B [0005]
- WO 9101144 A [0059]
- US 4885363 A [0059]
- WO 03002489 A [0077]
- WO 03002157 A [0077]

Non-patent literature cited in the description

- **GAVISH et al.** *Pharm. Rev.*, 1999, vol. 51, 629 [0002]
- *Neurosci. Lett.*, 1998, vol. 24 (1), 53-6 [0002]
- *Gen. Pharmacol.*, 1997, vol. 28 (4), 495-8 [0002]
- *Eur. J. Pharmacol.*, 2002, vol. 452 (1), 111-22 [0002]
- *J. Nucl. Med.*, 2004, vol. 45, 1898-1907 [0002]
- **BANATI et al.** *Brain*, 2000, vol. 123, 2321 [0002]
- *TINS*, 2005, vol. 28 (2), 101-7 [0002]
- **OKUBU et al.** *Bioorganic & Medicinal Chemistry*, 2004, vol. 12, 3569-80 [0003]
- **CAMPIANI et al.** *J. Med. Chem.*, 2002, vol. 45, 4276-81 [0003]
- **OKUBO et al.** *Bioorganic and Medicinal Chemistry*, 2002, vol. 12, 3569-80 [0023]
- **LE FUR et al.** *Life Sci. USA*, 1983, vol. 33, 449-57 [0039] [0167]
- **OKUBO et al.** *Bioorganic and Medicinal Chemistry*, 2004, vol. 12, 3569-80 [0045]
- **THEORODORA W. GREENE ; PETER G. M. WUTS.** *Protective Groups in Organic Synthesis*. John Wiley & Sons, 1999 [0050]
- **JURISSON et al.** *Chem. Rev.*, 1999, vol. 99, 2205-2218 [0059]
- **BOLTON.** *J. Lab. Comp. Radiopharm.*, 2002, vol. 45, 485-528 [0065]
- **WILBUR.** *Bioconj. Chem.*, 1992, vol. 3 (6), 433-470 [0065]
- **KABALAKA et al.** *Nucl. Med. Biol.*, 2002, vol. 29, 841-843 [0065]
- *NUCL. MED. BIOL.*, 2003, vol. 30, 369-373 [0065]
- **KABALAKA et al.** *Nucl. Med. Biol.*, 2004, vol. 31, 935-938 [0065]
- Aspects on the Synthesis of C-Labelled Compounds. **ANTONI et al.** *Handbook of Radiopharmaceuticals*. John Wiley and Sons, 2003 [0074]
- **OKUBO et al.** *Bioorg. Med. Chem.*, 2004, vol. 12, 3569-3580 [0134]
- **CHRISTENSEN.** *Molecules*, 2001, vol. 6, 47-51 [0138]