



(11) **EP 2 777 701 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**17.09.2014 Bulletin 2014/38**

(21) Application number: **14001638.7**

(22) Date of filing: **31.05.2007**

(51) Int Cl.:  
**A61K 31/22** <sup>(2006.01)</sup> **A61K 31/232** <sup>(2006.01)</sup>  
**A61K 31/351** <sup>(2006.01)</sup> **A61K 45/06** <sup>(2006.01)</sup>  
**A61P 3/10** <sup>(2006.01)</sup> **A61P 9/00** <sup>(2006.01)</sup>  
**A61P 9/10** <sup>(2006.01)</sup>

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE  
SI SK TR**

(30) Priority: **31.05.2006 JP 2006152740**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**07744487.5 / 2 022 495**

(71) Applicant: **MOCHIDA PHARMACEUTICAL CO.,  
LTD.  
Shinjuku-ku  
Tokyo 160-8515 (JP)**

(72) Inventors:  
• **Yokoyama, Mitsuhiro  
Kobe-shi Hyogo, 651-1112 (JP)**

- **Origasa, Hideki  
Toyama-shi Toyama, 930-0044 (JP)**
- **Matsuzaki, Masunori  
Ube-shi Yamaguchi, 755-0152 (JP)**
- **Matsuzawa, Yuji  
Takarazuka-shi Hyogo, 665-0804 (JP)**
- **Saito, Yasushi  
Chiba-shi Chiba, 260-0853 (JP)**

(74) Representative: **Grünecker, Kinkeldey,  
Stockmair & Schwanhäusser  
Anwaltssozietät  
Leopoldstrasse 4  
80802 München (DE)**

Remarks:

This application was filed on 08-05-2014 as a  
divisional application to the application mentioned  
under INID code 62.

(54) **Composition for preventing the occurrence of a cardiovascular event in multiple risk patient  
comprising the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid**

(57) Disclosed is a composition comprising all-cis-  
5,8,11,14,17-icosapentaenoic acid for use in the preven-  
tion of a cardiovascular event in a hypercholesterolemia

patient undergoing treatment with HMG-CoA reductase  
inhibitor and suffering from additional risk factors.

**EP 2 777 701 A1**

**Description**

## TECHNICAL FIELD

- 5   **[0001]** This invention relates to a composition for preventing occurrence of cardiovascular events (primary prevention) in multiple risk patients, the composition containing at least ethyl icosapentate (hereinafter abbreviated as EPA-E).

## BACKGROUND ART

- 10   **[0002]** Westernization of diet has resulted in the increase of patients suffering from lifestyle-related diseases such as diabetes, hyperlipidemia, and hypertension. Some of these diseases finally lead to arteriosclerotic diseases such as myocardial infarction, angina pectoris, and cerebral infarction. Treatment of the lifestyle-related diseases is based on the improvement of lifestyle, and more specifically, on the alimentary therapy and kinesitherapy. However, such improvement of the dietary life or the lack of exercise is often difficult in the patients suffering from the "lifestyle-related diseases", and they usually transfer to pharmacotherapy in order to prevent poor prognosis, for example, onset of myocardial infarction or cerebral infarction.

- 15   **[0003]** An exemplary compound having the action of improving such lifestyle-related diseases is polyunsaturated fatty acid. The polyunsaturated fatty acid is defined as a fatty acid including two or more carbon-carbon double bonds in one molecule, and the polyunsaturated fatty acids are categorized by the position of the double bond into  $\omega$ 3 fatty acid,  $\omega$ 6 fatty acid, and the like. The  $\omega$ 3 polyunsaturated fatty acids include  $\alpha$ -linolenic acid, icosapentaenoic acid (EPA), and docosahexaenoic acid (DHA), and the  $\omega$ 6 polyunsaturated fatty acids include linoleic acid,  $\gamma$ -linolenic acid, and arachidonic acid. Polyunsaturated fatty acids are derived from natural products, and exhibit various actions including antiarteriosclerotic action, platelet aggregation inhibitory action, hypolipidemic action, antiinflammatory action, antitumor action, and central action, and due to the high safety, polyunsaturated fatty acids are incorporated in various kinds of food, or sold as a health food or drug.

- 20   **[0004]** Decrease in the death rate in the patients who have history of suffering from myocardial infarction has been reported for the administration of a mixture of ethyl ester of an  $\omega$ -3 polyunsaturated fatty acid EPA (EPA-E) and ethyl ester of an  $\omega$ -3 polyunsaturated fatty acid DHA (DHA-E) for 3.5 years (see Patent Document 1). However, the results disclosed in Patent Document 1 relates to the secondary prevention, that is, prevention of recurrence, and the drug which is effective in the secondary prevention is not always effective in the primary prevention.

- 30   **[0005]** Based on the results of animal experiments and small scale clinical observations, many large scale clinical trials have been recently planned and conducted for the purpose of confirming whether various drugs which are effective in improving the lifestyle-related diseases can also prevent arteriosclerotic diseases in human. The results, however, have not necessarily been as intended, and the situation is still severe for the prevention of the occurrence of cardiovascular events in the case of patients suffering from a plurality of risk factors.

- 35   **[0006]** High purity EPA-E is commercially available in the trade names of Epadel<sup>TM</sup> and Epadel S<sup>TM</sup> (manufactured by Mochida Pharmaceutical Co., Ltd.) as therapeutic drugs for hyperlipidemia. There has been reported that when such high purity EPA-E is orally administered at 600 mg per administration and 3 times a day immediately after meal (when TG is abnormal, the dose is increased to the level of 900 mg per administration and 3 times a day), serum T-Cho concentration can be reduced by 3 to 6%, and serum TG can be reduced by 14 to 20% (see Non-Patent Document 1). There has also been reported in The Heart Failure Society of America 2005 Annual Meeting that, based on such action, such high purity EPA-E was expected to have the effects of improving cardiovascular events in hyperlipidemia patients, and combined use with HMG-CoA RI was effective in inhibiting cardiac events in a large scale clinical trial. In this large scale clinical trial (JELIS, Japan EPA Lipid Intervention Study), statistically significant suppression of the cardiac events by the EPA-E was confirmed for the total of the primary prevention patients and secondary prevention patients, and for the secondary prevention patients. On the other hand, in the analysis limited to the primary prevention patients, the incidence of the events was lower in the EPA-E group (the group administered with EPA-E in combination with HMG-CoA RI) compared to the control group (the group administered with solely with HMG-CoA RI), while this difference was not statistically significant. This trial also revealed that after 5 years from the start of the trial, the LDL-cholesterol value reduced by 26% in both of the EPA-E group and control group, that no significant difference was found between these groups, and that change of the HDL-cholesterol value was slight in both groups (see Non-Patent Document 2). This trial also revealed that the total cholesterol and the LDL-cholesterol decreased by 19% and 25%, respectively, in both of the EPA-E group and the control group, and that triglyceride decreased by 9% (significant) and 4% in the EPA-E group and the control group, respectively, while little change in HDL-C was noted in both of the EPA-E group and the control group (see Non-Patent Document 3). There is so far no report that has analyzed prevention of the occurrence of the cardiovascular events in the case of patients having two or more risk factors.

Non-Patent Document 1: Drug Interview Form "EPA preparation, Epadel capsule 300", revised in July, 2002 and February, 2004, version 21 issued in December, 2004; pp. 21-22.

Non-Patent Document 2: Medical Tribune, issue of November 17, 2005, Feature article 3, pp. 75-76.

Non-Patent Document 3: Lancet, vol. 369, pages 1090 to 1098 (2007).

5

## DISCLOSURE OF THE INVENTION

### PROBLEMS TO BE SOLVED BY THE INVENTION

10 **[0007]** In view of the situation that there is a serious problem that death from the cardiovascular disease is still a major cause of the death, and many cases of cardiovascular events are still impossible to prevent by the HMG-CoA RI therapy, an object of the present invention is to provide a composition for preventing onset of the cardiovascular events.

### MEANS TO SOLVE THE PROBLEMS

15

**[0008]** In order to solve the problems as described above, the inventors of the present invention made an extensive study on a therapy of hypercholesterolemia patients and found that EPA-E has the effect of preventing occurrence of the cardiovascular events in patients suffering from multiple risk factors, and in particular, the effect of preventing occurrence of the cardiovascular events in male patients suffering from multiple risk factors. The present invention has been completed on the bases of such finding. Accordingly, the present invention is directed to the following:

20

(1) A composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

25

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

30

(2) A composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the hypercholesterolemia patient is a patient also suffering from two or more of the risk factors.

35

(3) A composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one of risk factors as defined by a body mass index (BMI) of at least 25 for the obesity; by a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg for the hypertension or the prehypertension; by a fasting blood glucose (FBS) of at least 126 mg/dL or a hemoglobin A1c (HbA1c) of at least 6.5% for the diabetes, the prediabetes, or the abnormal glucose tolerance; and by triglyceride (TG) of at least 150 mg/dL and/or a HDL-C of less than 40 mg/dL for the hypertriglyceridemia and/or the low HDL cholesterolemia.

40

(4) The composition according to any one of (1) to (3) above, wherein the content of the EPA-E is at least 96.5% by weight in relation to the total content of fatty acid and derivatives thereof.

(5) The composition according to any one of (1) to (4) above, wherein the EPA-E is orally administered at a dose of 1.8 g/day to 2.7 g/day.

45

(6) The composition according to any one of (1) to (5) above, wherein the composition is used in combination with HMG-CoA RI.

(7) The composition according to any one of (1) to (6) above, wherein the hypercholesterolemia patient is a male patient.

50

(8) The composition according to any one of (1) to (7) above, wherein the hypercholesterolemia patient is a patient also suffering from hypertriglyceridemia and low HDL cholesterolemia.

(9) A method for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient by administering the patient with the composition according to any one of (1) to (8) above.

(10) Use of the composition according to any one of (1) to (8) above for the manufacture of an agent for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient.

55

### MERITS OF THE INVENTION

**[0009]** The above-mentioned composition of the present invention containing at least EPA-E as its effective component

is effective in preventing occurrence of cardiovascular events in hypercholesterolemia patients, and in particular, in preventing occurrence of cardiovascular events in hypercholesterolemia patients who have been treated with HMG-CoA RI but still suffer from the risk of the cardiovascular events, or more particularly, in preventing occurrence of cardiovascular events in hypercholesterolemia patients also suffering from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

**[0010]** The effect of the composition of the present invention will be synergistically improved by combined use with the HMG-CoA RI, and such use of the composition of the present invention with the HMG-CoA RI has clinical utility since the effect of preventing the cardiovascular event occurrence is expected to be improved.

#### BRIEF DESCRIPTION OF THE DRAWINGS

##### **[0011]**

[FIG. 1] FIG. 1 is a graph prepared by plotting the incidence of the cardiovascular events in Y-axis and the time after the start of the trial in X-axis for male patients having at least 2 risk factors.

[FIG. 2] FIG. 2 is a graph prepared by plotting the incidence of the cardiovascular event in Y-axis and the time after the start of the trial in X-axis for patients having the risk factors of a triglyceride (TG) of at least 150 mg/dL and a HDL-C of less than 40 mg/dL.

#### BEST MODE FOR CARRYING OUT THE INVENTION

**[0012]** Next, the present invention is described in detail. A first aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia. Alternatively, the first aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of
- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

**[0013]** The prevention of the occurrence of the cardiovascular events include all cases of primary prevention, and exemplary cases include prevention of cardiovascular death, fatal myocardial infarction, sudden cardiac death, nonfatal myocardial infarction, cardiovascular angioplasty, new occurrence of rest angina and exercise-induced angina, and destabilization of the angina. The composition of the present invention may be administered to any person who needs prevention of the occurrence of the cardiovascular events, and typical such patients are hypercholesterolemia patients.

**[0014]** A second aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient undergoing a HMG-CoA RI therapy, the composition containing at least EPA-E, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia. Alternatively, the second aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient undergoing a HMG-CoA RI therapy, the composition containing at least EPA-E and/or DHA-E, wherein the patient

also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

**[0015]** While HMG-CoA RI includes all those having inhibitory action for 3-hydroxy-3-methylglutaryl coenzyme A reductase, the one used in the present invention is preferably a pharmaceutically administrable inhibitor which is preferably at least one member selected from the group consisting of pravastatin, simvastatin, lovastatin, fluvastatin, cerivastatin, atorvastatin, pitavastatin, rosuvastatin, and salts and derivatives thereof, and more preferably, pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, pitavastatin, or rosuvastatin, and most preferably, pravastatin or simvastatin. All salts are included as long as they are pharmaceutically administrable, and preferred are sodium and calcium salts such as pravastatin sodium, fluvastatin sodium, cerivastatin sodium, atorvastatin calcium, pitavastatin calcium, and rosuvastatin calcium. In the present invention, "pravastatin", for example, also includes the pravastatin in the form of a salt unless otherwise noted.

**[0016]** A third aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least two risk factors selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia; namely; obesity, and hypertension or prehypertension; obesity, and diabetes, prediabetes, or abnormal glucose tolerance; obesity, and hypertriglyceridemia and/or low HDL cholesterolemia; hypertension or prehypertension, and diabetes, prediabetes or abnormal glucose tolerance; diabetes, prediabetes, or abnormal glucose tolerance, and hypertriglyceridemia and/or low HDL cholesterolemia; obesity, and hypertension or prehypertension, and diabetes, prediabetes, or abnormal glucose tolerance; obesity, and hypertension or prehypertension, and hypertriglyceridemia and/or low HDL cholesterolemia; obesity, and diabetes, prediabetes, or abnormal glucose tolerance, and hypertriglyceridemia and/or low HDL cholesterolemia; hypertension or prehypertension, and diabetes, prediabetes, or abnormal glucose tolerance, and hypertriglyceridemia and/or low HDL cholesterolemia. Alternatively, the third aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least two risk factors selected from the group consisting of
- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

**[0017]** A fourth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one, and more preferably, at least two risk factors selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia. In this case, the hypercholesterolemia patient is preferably a male patient. Alternatively, the fourth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one, and more preferably, at least two risk factors selected from the group consisting of
- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia. In this case, the hypercholesterolemia patient is preferably a male patient.

**[0018]** A fifth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from risk factors of hypertriglyceridemia and low HDL cholesterolemia, and more specifically, hypertriglyceridemia and low HDL cholesterolemia with a serum triglyceride (TG) concentration of at least 150 mg/dl and a serum HDL-C concentration of less than 40 mg/dl, or serum TG/HDL-C ratio of at least 3.75. In this case, the hypercholesterolemia patient is preferably a male patient. Alternatively, the fifth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from risk factors of hypertriglyceridemia and low HDL cholesterolemia, and more specifically, hypertriglyceridemia and low HDL cholesterolemia with a serum triglyceride (TG) concentration of at least 150 mg/dl and a serum HDL-C concentration of less than 40 mg/dl, or a serum TG/HDL-C ratio of at least 3.75. In this case, the hypercholesterolemia patient is preferably a male patient.

**[0019]** A sixth aspect of the present invention provides a composition containing at least EPA-E as its effective component, the composition exhibiting an excellent effect of preventing occurrence of a cardiovascular event in a patient suffering from multiple risk factors who has been administered with this composition for at least 2 years since the start of the administration. Alternatively, the sixth aspect of the present invention provides a composition containing at least EPA-E and/or DHA-E as its effective component, the composition exhibiting an excellent effect of preventing recurrence of a cardiovascular event in a patient suffering from multiple risk factors who has been administered with this composition for at least 2 years since the start of the administration. The hypercholesterolemia patient is preferably a male patient.

**[0020]** A seventh aspect of the present invention provides a method for preventing occurrence of a cardiovascular event in a patient suffering from multiple risk factors by continuously administering the patient with a composition containing at least EPA-E as its effective component for at least 2 years. Alternatively, the seventh aspect of the present invention provides a method for preventing occurrence of a cardiovascular event in a patient suffering from multiple risk factors by continuously administering the patient with a composition containing at least EPA-E and/or DHA-E as its effective component for at least 2 years. The hypercholesterolemia patient is preferably a male patient.

**[0021]** An eighth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a dyslipidemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia. Alternatively, the eighth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a dyslipidemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of
- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

**[0022]** A ninth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient to be able to administered with HMG-CoA RI, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia. Alternatively, the ninth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient to be able to administered with HMG-CoA RI, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of
- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

**[0023]** While the EPA-E content in the total fatty acid and dosage are not particularly limited as long as intended effects of the present invention are attained, the EPA-E used is preferably the one having a high purity, for example, the one having the proportion of the EPA-E in the total fatty acid and derivatives thereof of preferably 40% by weight or higher, more preferably 90% by weight or higher, and still more preferably 96.5% by weight or higher. The daily dose in terms of EPA-E is typically 0.3 to 6 g/day, preferably 0.9 to 3.6 g/day, and still more preferably 1.8 to 2.7 g/day. Another preferable daily dose is 0.3 to 2.7 g/day, and 0.3 to 1.8 g/day. Another preferable fatty acid included is DHA-E. While the compositional ratio of EPA-E/DHA-E, content of EPA-E and DHA-E (hereinafter referred to as (EPA-E + DHA-E)) in the total fatty acid, and dosage of (EPA-E + DHA-E) are not particularly limited as long as intended effects of the present invention are attained, the composition is preferably the one having a high purity of EPA-E and DHA-E, for example, the one having a proportion of the (EPA-E + DHA-E) in the total fatty acid and derivatives thereof of preferably 40% by weight or higher, more preferably 80% by weight or higher, and still more preferably 90% by weight or higher. The daily dose in terms of EPA-E + DHA-E is typically 0.3 to 10 g/day, preferably 0.5 to 6 g/day, and still more preferably 1 to 4 g/day. Another preferable daily dose is 0.3 to 6 g/day, 0.3 to 4 g/day, and 0.3 to 1 g/day. The content of other long chain saturated fatty acids is preferably low, and among the long chain unsaturated fatty acids, the content of  $\omega$ 6 fatty acids, and in particular, the content of arachidonic acid is preferably as low as less than 2% by weight, and more preferably less than 1% by weight.

**[0024]** The composition of the present invention contains EPA-E and/or DHA-E, and has the effect of preventing occurrence of cardiovascular events in healthy people or those suffering from the risk factors of hyperlipidemia, diabetes, and hypertension when the composition is orally administered, and in particular, of preventing occurrence of cardiovascular events in hypercholesterolemia patients who have been treated with HMG-CoA RI but still suffering from the risk of the cardiovascular events. The composition of the present invention may also be used in combination with the HMG-CoA RI, and such combination may further prevent the occurrence of the cardiovascular events.

**[0025]** The composition of the present invention may be used with other drugs, for example, antiplatelet drugs such as aspirin, ticlopidine, clopidogrel, prasugrel, and cilostazol; anticoagulants such as warfarin, heparin, and ximelagatran; antihypertensive drugs such as angiotensin II receptor antagonists (candesartan, losartan, valsartan, etc.), angiotensin converting enzyme inhibitors, calcium channel antagonists (amlodipine, cilnidipine, etc.), and  $\alpha$ 1 blockers; diabetes drugs or abnormal glucose tolerance stimulants such as  $\alpha$ -glucosidase inhibitors (voglibose, acarbose, etc.), biguanide drugs, thiazolidinedione drugs (pioglitazone, rosiglitazone, rivoglitazone, etc.), and prompt insulin release promoters (mitiglinide, nateglinide, etc.); antilipotropic drugs and antiarteriosclerotic drugs such as HMG-CoA RI as described above, fibrate drugs, squalene synthetase inhibitors (TAK-475, etc.), and cholesterol absorption inhibitors (ezetimibe, etc.), probucol, anion exchange resin, nicotinic acid drugs, phytosterol, elastase, dextran sulfate sodium sulfur, pantothenic acid, and polyenephosphatidylcholine.

**[0026]** The composition of the present invention contains smaller amounts of impurities such as saturated fatty acids and arachidonic acid which are unfavorable for cardiovascular events compared to fish oil or fish oil concentrate, and accordingly, the intended effects can be attained without causing problems like overnutrition or excessive intake of vitamin A. In addition, since the effective component of the present composition is in the form of an ester, the effective component is more stable to oxidation compared to the case of fish oil in which the effective component is in the form of a triglyceride, and a sufficiently stable composition can be produced by adding a conventional antioxidant. In other words, it is the use of the EPA-E that has for the first time enabled production of a composition for preventing onset of cardiovascular events which can be used in clinical practice.

**[0027]** In the present invention, the term "icosapentaenoic acid" designates all-cis-5,8,11,14,17-icosapentaenoic acid.

**[0028]** In the present invention, the term "hypercholesterolemia patient" means the patient with increased serum T-Cho concentration or serum LDL-Cho concentration. In a narrower sense, this term means the patient suffering from hypercholesterolemia (serum T-Cho concentration of at least about 220 mg/dl, and in more strict sense, at least 250 mg/dl) or high LDL cholesterol (serum LDL-Cho concentration of at least 140 mg/dL).

**[0029]** In the present invention, the term "dyslipidemia" is the condition which satisfies at least one of high LDL cholesterol (i.e. fasting serum LDL cholesterol value of at least 140 mg/dL), low HDL cholesterol (i.e. fasting serum HDL cholesterol value of less than 40 mg/dL), and hypertriglyceridemia (i.e. fasting serum triglyceride value of at least 150 mg/dL) according to the diagnostic criteria described in "Guideline for Preventing Arteriosclerotic diseases, 2007" (edited and published by Japan Atherosclerosis Society).

**[0030]** Of the risk factors treated in the present invention, "obesity" is the state with excessive accumulation of fats in the body. For example, non-limiting examples of the obesity include a body mass index (BMI) of at least 25, a waist measurement of at least 85 cm in male and at least 90 cm in female. "Hypertension" is the state with an abnormal increase in resting arterial blood pressure of the greater circulatory system. For example, in the criteria proposed by Japanese Society of Hypertension at the time of the filing of this application, hypertension is defined as a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg. "Prehypertension" is the condition with the blood pressure between the normal blood pressure (or optimal blood pressure) and the blood pressure in the hypertension, and this condition is also referred to as "mild elevated blood pressure" or "borderline hypertension".

Non-limiting exemplary criteria for such condition include a systolic blood pressure (SBP) of 120 to 139 mmHg or a diastolic blood pressure (DBP) of 80 to 89 mmHg. In the present invention, "hypertension or prehypertension" means a condition with a systolic blood pressure (SBP) of at least 120 mmHg or a diastolic blood pressure (DBP) of at least 80 mmHg, more strictly, a systolic blood pressure (SBP) of at least 135 mmHg or a diastolic blood pressure (DBP) of at least 85 mmHg, even more strictly a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg. "Diabetes" is the glucose metabolism disorder caused by hyposecretion of insulin from the insulin-producing cell ( $\beta$  cell) in the pancreas or insufficient action of the insulin in the target cell. Exemplary non-limiting criteria proposed by Japan Diabetes Society at the time of the filing of this application is one of 1) fasting blood glucose of at least 126 mg/dL, 2) 75 g glucose tolerance test at 2 hours of at least 200 mg/dL, and 3) casual blood glucose level of at least 200 mg/dL; or a hemoglobin A1c (HbA1c) of at least 6.5%. The criteria, however, are not limited to these. "Prediabetes" is the condition in which the blood glucose level is between the normal value and the value in the diabetes. "Abnormal glucose tolerance" is the condition in which the blood glucose level in the glucose tolerance test is between the normal value and the value in the diabetes. These conditions are also referred to as the borderline diabetes, prediabetic state, and the diabetic high-risk group. For these conditions, exemplary non-limiting criteria include a fasting blood glucose of 110 to 125 mg/dL, a 75 g glucose tolerance test at 2 hours of 140 to 199 mg/dL, and a hemoglobin A1c (HbA1c) of 5.6 to 6.4%. In the present invention, "diabetes, prediabetes, or abnormal glucose tolerance" means a condition with a fasting blood glucose (FBS) of at least 110 mg/dL or a hemoglobin A1c (HbA1c) of at least 5.6%, more strictly, a fasting blood glucose (FBS) of at least 110 mg/dL or a hemoglobin A1c (HbA1c) of at least 5.9%, and even more strictly with a fasting blood glucose (FBS) of at least 126 mg/dL or a hemoglobin A1c (HbA1c) of at least 6.5%. "Hypertriglyceridemia" is the condition with an increased serum triglyceride (TG) concentration, and strictly, with the serum TG concentration of at least 150 mg/dL. "Low HDL cholesterolemia" is the condition with a reduced serum HDL-C concentration, and strictly, with the serum HDL-C concentration of less than 40 mg/dL. In the present invention, "hypertriglyceridemia and/or low HDL cholesterolemia" means the state with a serum TG concentration of at least 150 mg/dL and/or a serum HDL-C concentration of less than 40 mg/dL. The hypertriglyceridemia and the low HDL cholesterolemia are both diseases included in the category of dyslipidemia, and they are mutually independent risk factors. Combination of these risk factors, however, is known to result in an increased risk of the occurrence of an arteriosclerotic disease. In the present invention, "the hypertriglyceridemia and/or the low HDL cholesterolemia" is treated as a single risk factor.

**[0031]** In the present invention, the term "combined use of EPA-E with HMG-CoA RI" include both the embodiment in which the EPA-E and the HMG-CoA RI are simultaneously administered and the embodiment in which both agents are separately administered. When these agents are simultaneously administered, they may be formulated either as a single combined drug or separate drugs. When these agents are separately administered, EPA-E may be administered either before or after the HMG-CoA RI. The doses and ratio of the EPA-E and the HMG-CoA RI may be adequately selected.

**[0032]** In the present invention, the term "combined use of EPA-E and/or DHA-E with HMG-CoA RI" include both the embodiment in which the EPA-E and/or DHA-E and the HMG-CoA RI are simultaneously administered and the embodiment in which these agents are separately administered. When these agents are simultaneously administered, they may be formulated either as a single combined drug or separate drugs. When these agents are separately administered, EPA-E and/or DHA-E may be administered either before or after the HMG-CoA RI. The doses and ratio of the EPA-E and/or DHA-E and the HMG-CoA RI may be adequately selected.

**[0033]** The composition of the present invention has the action of preventing onset of the cardiovascular events by the sole administration of the composition, and in particular, the present composition is expected to have the effect of preventing onset of the cardiovascular events which could not be prevented by the sole administration of the HMG-CoA RI. In addition, EPA-E has not only the action of reducing the serum T-Chol concentration and the serum TG, but also the action of suppressing platelet aggregation based on inhibition of arachidonic acid cascade, which is a pharmacological action different from the HMG-CoA RI. Therefore, the action as described above can also be exerted by combined administration with the HMG-CoA RI.

**[0034]** Since EPA-E and DHA-E are highly unsaturated, inclusion of an effective amount of an antioxidant such as butylated hydroxytoluene, butylated hydroxyanisole, propyl gallate, gallic acid, and pharmaceutically acceptable quinone, or  $\alpha$ -tocopherol is preferable.

**[0035]** The preparation may be orally administered to the patient in the dosage form of tablet, capsule, microcapsule, granules, fine granules, powder, oral liquid preparation, syrup, or jelly. Preferably, the preparation is orally administered by filling in a capsule such as soft capsule or microcapsule.

**[0036]** The soft capsules containing high purity EPA-E (Epadel™ and Epadel S™) are commercially available in Japan as safe therapeutic agents for arteriosclerosis obliterans and hyperlipidemia with reduced side effects, and in such products, proportion of EPA-E in total fatty acid is at least 96.5% by weight. The soft capsule (Omacor™, Ross products, Reliant, and Pronova) containing about 46% by weight of EPA-E and about 38% by weight of DHA-E is commercially available in the U.S., Europe, and other countries as a drug applied for hypertriglyceridemia. These drugs may be purchased for use in the present invention.



**[0037]** The dose and administration period of the composition of the present invention for preventing the onset of the cardiovascular events is the dose and period sufficient for the expression of the intended action, and the dose and administration period may be adequately adjusted depending on the dosage form, administration route, daily frequency, severity of the symptoms, body weight, age, and the like. When orally administered, the composition may be administered at a dose in terms of EPA-E of 0.3 to 6 g/day, preferably 0.9 to 3.6 g/day, and more preferably 1.8 to 2.7 g/day, and while such dose is typically administered in 3 divided doses, if desired, such dose may be administered in a single dose or in several divided doses. The composition is preferably administered during or after the meal, and more preferably, immediately (within 30 minutes) after the meal. When such dose is orally administered, the administration period is typically at least 1 year, preferably at least 2 years, more preferably at least 3 years and still more preferably at least 5 years. The administration, however, is preferably continued as long as there is a considerable risk of onset of the cardiovascular events. If necessary, drug holidays of about 1 day to 3 months, and preferably about 1 week to 1 month may be given.

**[0038]** The HMG-CoA RI is preferably used according to the dosage regimen recommended for the particular drug used, and the dose may be adequately adjusted depending on the type, dosage form, administration route, daily frequency, severity of the symptoms, body weight, gender, age, and the like. When orally administered, the HMG-CoA RI is typically administered at a dose of 0.05 to 200 mg/day, and preferably 0.1 to 100 mg/day in a single dose or in two divided doses. If necessary, the total dose may be administered in several divided doses. The dose of the HMG-CoA RI may be reduced depending on the dose of the EPA-E.

**[0039]** It is to be noted that pravastatin sodium (Mevalotin™ tablets and fine granules, Daiichi Sankyo Co., Ltd.), simvastatin (Lipovas™ tablets, Banyu Pharmaceutical Co., Ltd.), fluvastatin sodium (Lochol™ Tablets, Novartis Pharma K.K. and Tanabe Seiyaku Co., Ltd.), atorvastatin calcium hydrate (Lipitor™ tablets, Astellas Pharma Inc. and Pfizer Inc.), pitavastatin calcium (Livalo™, Kowa Company, Ltd. and Daiichi Sankyo Co., Ltd.), and rosuvastatin calcium (Crestor™ tablets, AstraZeneca and Shionogi & Co., Ltd.) are commercially available in Japan as drugs for treating hyperlipidemia, and lovastatin (Mevacor™ tablets, Merck) is commercially available in the U.S. as a drug for treating hyperlipidemia. These drugs may be purchased and used according to the prescribed dosing schedules.

**[0040]** In the case of pravastatin sodium, the preferable daily dose is 5 to 60 mg, and more preferably 10 to 20 mg, and in the case of simvastatin, the preferable daily dose is 2.5 to 60 mg, and more preferably 5 to 20 mg. In the case of fluvastatin sodium, the preferable daily dose is 10 to 180 mg, and more preferably 20 to 60 mg, and in the case of atorvastatin calcium hydrate, the preferable daily dose is 5 to 120 mg, and more preferably 10 to 40 mg. In the case of pitavastatin calcium, the preferable daily dose is 0.5 to 12 mg, and more preferably 1 to 4 mg, and in the case of rosuvastatin calcium, the preferable daily dose is 1.25 to 60 mg, and more preferably 2.5 to 20 mg. In the case of lovastatin, the preferable daily dose is 5 to 160 mg, and more preferably 10 to 80 mg, and in the case of cerivastatin sodium, the preferable daily dose is 0.075 to 0.9 mg, and more preferably 0.15 to 0.3 mg. The dose, however, is not limited to those as described above.

## EXAMPLES

**[0041]** Next, the effects of the composition of the present invention are demonstrated by referring to Examples, which by no means limit the scope of the present invention.

Example 1: Effect of the EPA-E in preventing occurrence of cardiovascular events in patients having multiple risk factors

### Trial procedure

**[0042]** This trial corresponds to a partial analysis of the results obtained in JELIS (Japan EPA Lipid Intervention Study) which is a large scale clinical trial of high purity EPA preparation which was presented in The Heart Failure Society of America 2005 Annual Meeting (for general information on JELIS, see Medical Tribune, issue of November 17, 2005, Feature article 3, pp. 75-76).

**[0043]** More specifically, for the EPA-E group (7503 cases) and the control group (7478 cases) evaluated for the primary prevention effect in the 18,645 subject patients of the JELIS trial (EPA-E group (9,326 cases) and control group (9,319 cases)), occurrence of the cardiovascular events was observed and analysed for 5 years from the start of the administration in relation to the number of risk factors at the registration as defined by the following (1) to (4):

- (1) obesity: body mass index (BMI) of at least 25;
- (2) hypertension or prehypertension: systolic blood pressure (SBP) of at least 140 mmHg or diastolic blood pressure (DBP) of at least 90 mmHg;
- (3) diabetes, prediabetes, or abnormal glucose tolerance: fasting blood glucose (FBS) of at least 126 mg/dL or hemoglobin A1c (HbA1c) of at least 6.5%;

(4) hypertriglyceridemia or low HDL cholesterolemia: a triglyceride (TG) of at least 150 mg/dL or a HDL-C of less than 40 mg/dL. The EPA-E group was orally administered with Epadel (Mochida Pharmaceutical Co., Ltd.) typically at an adult dose of 600 mg per administration and 3 times a day after the meal. However, in the case of abnormal serum TG, the dose could be increased to 900 mg per administration and 3 times a day. In both groups, pravastatin sodium (Mevalotin™ tablets and fine granules, Daiichi Sankyo Co., Ltd.), simvastatin (Lipovas™ tablets, Banyu Pharmaceutical Co., Ltd.), or atorvastatin calcium hydrate (Lipitor™ tablets, Astellas Pharma Inc. and Pfizer Inc.) was used for the base drug, and these drugs were orally administered according to the prescribed dosage regimen. More specifically, pravastatin sodium was orally administered at a daily dose of 10 to 20 mg in a single dose or two divided doses; simvastatin was orally administered at a daily dose of 5 to 20 mg in a single dose; atorvastatin calcium hydrate was orally administered at a daily dose of 10 to 40 mg in a single dose.

## Results

**[0044]** The number of occurrence of cardiovascular events in the observation period of 5 years, incidence (%), and rate of suppression of the incidence of the cardiovascular events in the EPA-E group with respect to the control group are shown in Table 1 for each number of risk factors. The rate of suppression of the incidence of the cardiovascular events was calculated by the formula:  $\{[(\text{incidence in the control group}) - (\text{incidence in the EPA-E group})] / \text{incidence in the control group}\} \times 100$ .

**[0045]** [Table 1]

Table 1

Number of risk factors	Incidence in the control group (cases of occurrence/ all cases, %)	Incidence in the EPA-E group (cases of occurrence/ all cases, %)	Rate of suppression (%)
0	14/1309 (1.1)	11/1326 (0.8)	22
1	29/2424 (1.2)	25/2468 (1.0)	15
2	46/2324 (2.0)	34/2238 (1.5)	23
3	29/1205 (2.4)	28/1229 (2.3)	5
4	9/216 (4.2)	6/242 (2.5)	40
1-2	75/4748 (1.6)	59/4706 (1.3)	21
1-3	104/5953 (1.7)	87/5935 (1.5)	18
1-4	113/6169 (1.8)	93/6177 (1.5)	18
2-3	75/3529 (2.1)	62/3467 (1.8)	16
2-4	84/3745 (2.2)	68/3709 (1.8)	18
3-4	38/1421 (2.7)	34/1471 (2.3)	14

**[0046]** The incidence (%) of cardiovascular events was found to increase with the increase in the number of risk factors. While the incidence was 1.1% for the risk factor number of 0 and 4.2% for the risk factor number 4 in the control group, the incidence was 0.8% in the risk factor number 0 and 2.5% for the case of risk factor number 4 in the group administered with the EPA-E. As evident from Table 1, for all cases of both groups with 1 to 4 risk factors, the cardiovascular event incidence was lower in the group administered with the EPA-E compared to the control group, and the cardiovascular events were suppressed by 5 to 40%. The effect of preventing occurrence of the cardiovascular events by the administration of the EPA-E was thereby confirmed for the hypercholesterolemia patients having the risk factors.

**[0047]** From the results of the trial as described above, the number of occurrence of cardiovascular events in the observation period of 5 years, incidence (%), and rate of suppression of the incidence of the cardiovascular events in the EPA-E group with respect to the control group were calculated for the male patients having at least two risk factors. The results are shown in Table 2. (The calculation was conducted by the same procedure as described above.)

**[0048]** [Table 2]

Table 2

Number of risk factors	Incidence in the control group (cases of occurrence/ all cases, %)	Incidence in the EPA-E group (cases of occurrence/ all cases, %)	Rate of suppression (%)
2-4	43/1053 (4.1)	19/1065 (1.8)	56

**[0049]** FIG. 1 is a graph prepared by plotting the incidence of the cardiovascular events in Y-axis and time after the start of the trial in X-axis.

**[0050]** As evident from Table 2 and FIG. 1, in the case of male patients having two or more risk factors, EPA-E significantly suppressed the occurrence of cardiovascular events. It was also confirmed that decrease in the incidence of the cardiovascular events was significant after 2 years or more from the start of the administration. At the end of the trial, the rate of suppression of the cardiovascular event occurrence was 56% compared to the control group (the value after correcting the dispersion between groups was 55%; see FIG. 1).

**[0051]** From the results of the trial as described above, the number of occurrence of cardiovascular events in the observation period of 5 years, incidence (%), and rate of suppression of the incidence of the cardiovascular events in the EPA-E group with respect to the control group were calculated for the patients exhibiting a triglyceride (TG) of at least 150 mg/dL and a HDL-C of less than 40 mg/dL as the risk factors. The results are shown in Table 3. (The calculation was conducted by the same procedure as described above.)

**[0052]** [Table 3]

Table 3

Risk factor	Incidence in the control group (cases of occurrence/ all cases, %)	Incidence in the EPA-E group (cases of occurrence/ all cases, %)	Rate of suppression (%)
TG of at least 150 mg/dL and HDL-C of less than 40 mg/dL	21/475 (4.4)	11/482 (2.3)	48

**[0053]** FIG. 2 is a graph prepared by plotting the incidence of the cardiovascular events in Y-axis and time after the start of the trial in X-axis.

**[0054]** As evident from Table 3 and FIG. 2, EPA-E significantly suppressed occurrence of cardiovascular events in the patients having the risk factors of the triglyceride (TG) of at least 150 mg/dL and the HDL-C of less than 40 mg/dL. It was also confirmed that decrease in the incidence of the cardiovascular events was significant after 2 years or more from the start of the administration. At the end of the trial, the rate of suppression of the cardiovascular event occurrence was 48% compared to the control group (the value after correcting the dispersion between groups was 53%; see FIG. 2). This suggests that the composition containing EPA-E as its effective component effectively prevents the occurrence of the cardiovascular event in the patient having the serum TG/HDL-C ratio of at least 3.75. It is also to be noted that, while the events that occurred in the control group were fatal myocardial infarction, nonfatal myocardial infarction, new occurrence of angina and cardiovascular angioplasty, the events that occurred in the EPA-E group were either nonfatal myocardial infarction or new occurrence of angina, and occurrence of fatal events was not found in the EPA-E group.

**[0055]** In addition, in the group of patients having the risk factor of the triglyceride of at least 150 mg/dL, the occurrence of the cardiovascular events was suppressed by 15% in the EPA-E group compared to the control group; and in the group of patients having the risk factor of HDL-C of less than 40 mg/dL, the occurrence of the cardiovascular events was suppressed by 35% compared to the control group (both values are uncorrected values).

**[0056]** As described above, a significant effect of the EPA-E administration was confirmed for the prevention of the occurrence of the cardiovascular events in the hypercholesterolemia patients having the risk factors.

**[0057]** The present invention is also directed to the following embodiments.

1. A composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition comprising ethyl icosapentate as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and

(4) hypertriglyceridemia and/or low HDL cholesterolemia.

2. The composition according to item 1, wherein the hypercholesterolemia patient is a patient also suffering from two or more of the risk factors.

3. The composition according to item 1 or 2, wherein the obesity is defined by a body mass index (BMI) of at least 25; the hypertension or the prehypertension is defined by a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg; the diabetes, the prediabetes, or the abnormal glucose tolerance is defined by a fasting blood glucose (FBS) of at least 126 mg/dL or a hemoglobin A1c (HbA1c) of at least 6.5%; and the hypertriglyceridemia and/or the low HDL cholesterolemia is defined by triglyceride (TG) of at least 150 mg/dL and/or a HDL-C of less than 40 mg/dL.

4. The composition according to any one of items 1 to 3, wherein the content of the ethyl icosapentate is at least 96.5% by weight in relation to the total content of fatty acid and derivatives thereof.

5. The composition according to any one of items 1 to 4, wherein the ethyl icosapentate is orally administered at a dose of 1.8 g/day to 2.7 g/day.

6. The composition according to any one of items 1 to 5, wherein the composition is used in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor.

7. The composition according to any one of items 1 to 6, wherein the hypercholesterolemia patient is a male patient.

8. The composition according to any one of items 1 to 7, wherein the hypercholesterolemia patient is a patient also suffering from hypertriglyceridemia and low HDL cholesterolemia.

## Claims

1. A composition for use in the primary prevention of a cardiovascular event in a hypercholesterolemia patient, wherein the composition comprises the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid as its effective component and is used in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor, wherein the cardiovascular event is cardiovascular death, fatal myocardial infarction, sudden cardiac death, nonfatal myocardial infarction, cardiovascular angioplasty, or new occurrence of rest angina and exercise-induced angina, and wherein said patient is a male patient and suffers from two or more of the risk factors selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

2. The composition for use according to claim 1, wherein the obesity is defined by a body mass index (BMI) of at least 25; the hypertension or the prehypertension is defined by a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg; the diabetes, the prediabetes, or the abnormal glucose tolerance is defined by a fasting blood glucose (FBS) of at least 126 mg/dL or a hemoglobin A1c (HbA1c) of at least 6.5%; and the hypertriglyceridemia and/or the low HDL cholesterolemia is defined by triglyceride (TG) of at least 150 mg/dL and/or a HDL-C of less than 40 mg/dL.

3. The composition for use according to claim 1 or 2, wherein the content of the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid is at least 96.5% by weight in relation to the total content of fatty acid and derivatives thereof.

4. The composition for use according to any one of claims 1 to 3, wherein the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid is orally administered at a dose of 0.3 g/day to 6 g/day.

5. The composition for use according to any one of claims 1 to 4, wherein the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid is orally administered at a dose of 1.8 g/day to 2.7 g/day.

6. A composition for use in the primary prevention of a cardiovascular event in a hypercholesterolemia patient, wherein the composition comprises the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid as its effective component and is used in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor, wherein the cardiovascular event is cardiovascular death, fatal myocardial infarction, sudden cardiac death, nonfatal myocardial infarction, cardiovascular angioplasty, or new occurrence of rest angina and exercise-induced angina, and wherein said patient also suffers from the following risk factors:
  - (1) obesity,
  - (2) hypertension or prehypertension,
  - (3) diabetes, prediabetes, or abnormal glucose tolerance, and
  - (4) hypertriglyceridemia and/or low HDL cholesterol.
7. The composition for use according to claim 6, wherein the obesity is defined by a body mass index (BMI) of at least 25; the hypertension or the prehypertension is defined by a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg; the diabetes, the prediabetes, or the abnormal glucose tolerance is defined by a fasting blood glucose (FBS) of at least 126 mg/dL or a hemoglobin A1c (HbA1c) of at least 6.5%; and the hypertriglyceridemia and/or the low HDL cholesterol is defined by triglyceride (TG) of at least 150 mg/dL and/or a HDL-C of less than 40 mg/dL.
8. The composition for use according to claim 6 or 7, wherein the content of the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid is at least 96.5% by weight in relation to the total content of fatty acid and derivatives thereof.
9. The composition for use according to any one of claims 6 to 8, wherein the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid is orally administered at a dose of 0.3 g/day to 6 g/day.
10. The composition for use according to any one of claims 6 to 9, wherein the ethyl ester of all-cis-5,8,11,14,17-icosapentaenoic acid is orally administered at a dose of 1.8 g/day to 2.7 g/day.
11. The composition for use according to any one of claims 6 to 10, wherein the hypercholesterolemia patient is a male patient.

FIG. 1





FIG. 2





## EUROPEAN SEARCH REPORT

 Application Number  
EP 14 00 1638

5

10

15

20

25

30

35

40

45

50

55

## DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	"AHA Hatsu Sokuho JELIS Kekka Happyo", INTERNET CITATION, 17 November 2005 (2005-11-17), XP003020147, Retrieved from the Internet: URL:http://www.medical-tribune.jp/congress/ /jelis/jelis.html [retrieved on 2008-01-01] * the whole document *	1-11	INV. A61K31/22 A61K31/232 A61K31/351 A61K45/06 A61P3/10 A61P9/00 A61P9/10
X	YOKOYAMA M ET AL: "Effects of eicosapentaenoic acid on cardiovascular events in Japanese patients with hypercholesterolemia: rationale, design, and baseline characteristics of the Japan EPA Lipid Intervention Study (JELIS)", AMERICAN HEART JOURNAL, MOSBY- YEAR BOOK INC, US, vol. 146, no. 4, 1 January 2003 (2003-01-01), pages 613-620, XP003020148, ISSN: 0002-8703, DOI: 10.1016/S0002-8703(03)00367-3 * the whole document *	1-11	TECHNICAL FIELDS SEARCHED (IPC) A61K
Y,D	WO 00/48592 A1 (PHARMACIA & UPJOHN SPA [IT]; PAMPARANA FRANCO [IT]) 24 August 2000 (2000-08-24) * abstract * * page 1, line 18 - line 21 * * claim 1 *	1-11	
Y	EP 1 157 692 A1 (QUATEX NV [AN] PRO APARTS INVESTIMENTOS E CON [PT]) 28 November 2001 (2001-11-28) * abstract * * claims 1-5,20-27 *	1-11	
The present search report has been drawn up for all claims			

Place of search

Date of completion of the search

Examiner

The Hague

8 August 2014

Langer, Oliver

## CATEGORY OF CITED DOCUMENTS

X : particularly relevant if taken alone  
 Y : particularly relevant if combined with another  
 document of the same category  
 A : technological background  
 O : non-written disclosure  
 P : intermediate document

T : theory or principle underlying the invention  
 E : earlier patent document, but published on, or  
 after the filing date  
 D : document cited in the application  
 L : document cited for other reasons

& : member of the same patent family, corresponding  
 document

EPO FORM 1503 03.82 (P04C01)



## EUROPEAN SEARCH REPORT

Application Number  
EP 14 00 1638

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	ADAN Y ET AL: "Effects of docosahexaenoic and eicosapentaenoic acid on lipid metabolism, eicosanoid production, platelet aggregation and atherosclerosis in hypercholesterolemic rats", BIOSCIENCE BIOTECHNOLOGY BIOCHEMISTRY, JAPAN SOCIETY FOR BIOSCIENCE, BIOTECHNOLOGY, AND AGROCHEMISTRY, TOKYO, JAPAN, vol. 63, no. 1, 1 January 1999 (1999-01-01), pages 111-119, XP002613171, ISSN: 0916-8451, DOI: 10.1271/BBB.63.111 * abstract * * page 111 *	1-11	
X,P	EP 1 790 339 A1 (MOCHIDA PHARM CO LTD [JP]) 30 May 2007 (2007-05-30) * abstract * * claims 1,7-11 *	1-11	
			TECHNICAL FIELDS SEARCHED (IPC)
The present search report has been drawn up for all claims			
Place of search <b>The Hague</b>		Date of completion of the search <b>8 August 2014</b>	Examiner <b>Langer, Oliver</b>
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 (P04C01)



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 14 00 1638

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-08-2014

10

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0048592 A1	24-08-2000	AT	216230 T	15-05-2002
		AU	775078 B2	15-07-2004
		BR	0008153 A	06-11-2001
		CA	2362271 A1	24-08-2000
		CN	1343120 A	03-04-2002
		CZ	20012947 A3	17-04-2002
		DE	60000133 D1	23-05-2002
		DE	60000133 T2	21-11-2002
		DK	1152755 T3	22-07-2002
		EP	1152755 A1	14-11-2001
		EP	1247523 A1	09-10-2002
		ES	2174814 T3	16-11-2002
		HK	1044720 A1	18-08-2006
		HU	0200111 A2	29-05-2002
		ID	30205 A	15-11-2001
		IL	144377 A	10-12-2006
		IT	MI990313 A1	17-08-2000
		JP	5441287 B2	12-03-2014
		JP	2002537252 A	05-11-2002
		JP	2010116414 A	27-05-2010
		KR	20070098954 A	05-10-2007
		NO	20013938 A	14-08-2001
		NZ	513093 A	28-02-2003
		PL	350851 A1	10-02-2003
		PT	1152755 E	30-09-2002
		SK	10802001 A3	03-12-2001
		US	7462643 B1	09-12-2008
		US	2009018193 A1	15-01-2009
		WO	0048592 A1	24-08-2000
		ZA	200106029 A	25-09-2002
-----				
EP 1157692 A1	28-11-2001	AT	305810 T	15-10-2005
		DE	60022987 D1	10-11-2005
		DE	60022987 T2	19-10-2006
		DK	1157692 T3	06-02-2006
		EP	1157692 A1	28-11-2001
		ES	2246769 T3	01-03-2006
		PL	347655 A1	03-12-2001
-----				
EP 1790339 A1	30-05-2007	BR	PI0613704 A2	01-02-2011
		CA	2570763 A1	08-01-2007
		CN	101217952 A	09-07-2008
		CN	102526733 A	04-07-2012
		EP	1790339 A1	30-05-2007
		JP	2012193211 A	11-10-2012
		KR	20080024511 A	18-03-2008

55

EPO FORM P0458

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 14 00 1638

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-08-2014

10

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		KR 20130137055 A	13-12-2013
		TW 1412361 B	21-10-2013
		US 2007021504 A1	25-01-2007
		US 2013065956 A1	14-03-2013
		WO 2007007686 A1	18-01-2007
-----			

15

20

25

30

35

40

45

50

55

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Patent documents cited in the description**

- WO 0048592 A [0006]
- JP 2002537252 A [0006]

**Non-patent literature cited in the description**

- EPA preparation, Epadel capsule 300. *Drug Interview Form*, July 2002, 21-22 [0006]
- *Medical Tribune*, 17 November 2005, vol. 3, 75-76 [0006] [0042]
- *Lancet*, 2007, vol. 369, 1090-1098 [0006]
- Guideline for Preventing Arteriosclerotic diseases. Japan Atherosclerosis Society, 2007 [0029]