



US 20110160132A1

(19) **United States**(12) **Patent Application Publication**  
**Morishita et al.**(10) **Pub. No.: US 2011/0160132 A1**(43) **Pub. Date: Jun. 30, 2011**(54) **LYMPHANGIOGENESIS-PROMOTING  
AGENTS****Publication Classification**(75) Inventors: **Ryuichi Morishita**, Osaka (JP);  
**Yasufumi Kaneda**, Osaka (JP);  
**Hironori Nakagami**, Osaka (JP);  
**Yukihiro Saito**, Hokkaido (JP)(51) **Int. Cl.**  
**A61K 38/18** (2006.01)  
**A61K 31/7052** (2006.01)  
**G01N 33/53** (2006.01)  
**C12Q 1/02** (2006.01)  
**A61P 9/00** (2006.01)  
**A61P 43/00** (2006.01)(73) Assignees: **AnGes MG, Inc.**, Ibaraki-shi (JP);  
**Osaka University**, Suita-shi (JP)(52) **U.S. Cl. .... 514/9.5; 514/44 R; 436/501; 435/29**(21) Appl. No.: **11/996,898**(57) **ABSTRACT**(22) PCT Filed: **Jul. 28, 2006**(86) PCT No.: **PCT/JP2006/315010**§ 371 (c)(1),  
(2), (4) Date: **Feb. 3, 2009**(30) **Foreign Application Priority Data**Jul. 28, 2005 (JP) ..... 2005-219410  
May 29, 2006 (JP) ..... 2006-148970

Described herein is the discovery that HGFs activate the growth and migration of lymphatic endothelial cells and thereby promote lymphangiogenesis. The present invention is based on this finding, and provides lymphangiogenesis-promoting agents comprising as active ingredients HGFs, or proteins or compounds functionally equivalent thereto. Based on the finding described above, the present invention also provides methods for promoting lymphangiogenesis which comprise the step of locally administering HGFs or proteins functionally equivalent thereto to affected areas in patients with lymphedema.




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7






FIG. 8




FIG. 9




FIG. 10





FIG. 11



## LYMPHANGIOGENESIS-PROMOTING AGENTS

### TECHNICAL FIELD

**[0001]** The present invention relates to novel lymphangiogenesis-promoting agents comprising, as active ingredients, HGFs or nucleic acids encoding HGFs, and/or methods for promoting lymphangiogenesis. The present invention also relates to agents or methods for preventing or treating lymphedema. The present invention further relates to methods of screening for compounds having lymphangiogenesis-promoting activity or compounds having preventive or therapeutic effects on lymphedema.

### BACKGROUND ART

**[0002]** Lymphedema refers to a condition characterized by the occlusion of lymphatic vessels which, in turn, causes abnormal congestion of tissue fluid which results in swelling, chronic inflammation, and/or fibrosis. There are primary and secondary lymphedemas. Known instances of primary lymphedema include Milroy's disease, Meige's disease, distal hypoplasia, proximal obstructive lymphadenopathy, and lymphangiectasia. Secondary lymphedema arises from another disease; for example, it often appears as an aftereffect of surgical treatment of cancers. In particular, secondary lymphedema often occurs after surgery or radiation therapy for breast cancer, uterine cancer, prostatic cancer, Kaposi's sarcoma and such. Particularly, after breast cancer surgery, lymphedema often appears in the upper limbs. 80% or more cases of lymphedema of the upper limbs arise after breast cancer surgery. Furthermore, lymphedema of the upper limbs is presumed to develop at a frequency of several percent or more after breast cancer surgery. In contrast, lymphedema of the lower limbs is often observed subsequent to uterine cancer surgery. In addition to the above, infections, injuries such as burn injury, inflammations and the like can also result in secondary lymphedema.

**[0003]** Lymphedema significantly impairs motor functions and increases the risk of infection in the affected areas, both of which result in reduction of patients' QOL. However, common therapies for lymphedema, such as massage, exercise therapy, and wearing a supporter, merely treat symptoms; neither radical treatment methods nor therapeutic agents are available at present and almost no therapeutic agent is known. While a compound called "guaifenesin" has been reported to be effective in treating lymphedema, its therapeutic effect still remains unclear (Patent Document 1).

**[0004]** VEGF-C, a member of the VEGF family, has been reported to be a peptidic factor that enhances lymphangiogenesis (Patent Documents 2 and 3; Non-Patent Documents 1 and 2). However, the other VEGF family members have no lymphangiogenesis-promoting activity. VEGF-C is the only lymphangiogenesis-promoting factor that is currently known.

**[0005]** Meanwhile, VEGF<sub>165</sub> (also referred to as "VEGF-A" or more simply "VEGF"; hereinafter also referred to as "VEGF"), is a member of the VEGF family that, like VEGF-C, is also well known as an angiogenic factor, the factor has been also known as a factor that enhances vascular permeability (for example, see Senger, D. et al., *Science* 219, 983-6 (1983)). In fact, VEGF has been reported to induce edema by its vascular permeability-enhancing activity when overexpressed in tissues (for example, see Isner, J M., et al., *Circ.*

*Res.* 89: 389-400 (2001)). In fact, edema has been reported to be observed at a frequency of 30% or more in human patients with ischemic diseases of lower limbs when the VEGF gene is intraarterially or intramuscularly introduced into the ischemic areas in lower limbs for the therapeutic purpose (Baumgartner, I. et al., *Ann. Intern. Med.* 132: 880-884 (2000)). In addition, there are many reports of the increased risk of edema accompanying VEGF gene therapy, including a report showing that edema of the lower limbs was observed in six of nine patients in which a plasmid encoding naked VEGF was administered to ischemic areas of the lower limbs by intramuscular injection (Baumgartner, I. et al., *Circulation* 97: 1114-1123 (1998)) and another report showing that edema appeared in three limbs when a plasmid encoding naked VEGF was intramuscularly administered to ischemic areas in seven limbs of six patients with lower limb ischemia due to thromboangiitis obliterans (TAO) (Isner, J M., et al., *J. Vasc. Surg.* 28: 964-975 (1998)).

**[0006]** In addition, VEGF-C is a ligand for VEGF receptor 3 (VEGFR-3), and the signal from VEGFR-3 alone is known to be sufficient for lymphangiogenesis. VEGF-C also binds to VEGFR-2. Since VEGFR-2-deficient mice die earlier than VEGF-A-deficient mice, VEGF-C and other VEGF members are presumed to be able to complement the VEGF activity (Scavelli, C. et al., *J. Anat.* 433-449 (2004)). VEGF exerts its function through binding to VEGFR-1 and VEGFR-2. Thus, the activation of VEGFR-2 is thought to induce the enhancement of vascular permeability (for example, see Issbrucker, K. et al., *FASEB J. express article* 10.1096/fj.02-0329fje. Published online Dec. 18, 2002).

**[0007]** HGF is hepatocyte growth factor, and has not only the activity of promoting the growth of hepatocytes but also other various physiological activities, including angiogenic activity (for example, see Non-Patent Document 3). HGFs are presently applied to the treatment of ischemic disease based on their angiogenic activity (Patent Documents 4 and 5; Non-Patent Document 4).

**[0008]** [Patent Document 1] WO03/000242

**[0009]** [Patent Document 2] U.S. Pat. No. 6,818,220

**[0010]** [Patent Document 3] U.S. Pat. No. 6,689,352

**[0011]** [Patent Document 4] WO97/07824

**[0012]** [Patent Document 5] WO01/32220

**[0013]** [Non-patent Document 1] Szuba, A. et al., *FASEB J. express article* 10.1096/fj.02-0401fje. Published online Oct. 18, 2002

**[0014]** [Non-patent Document 2] Young-sup, Yoo et al., *J. Clin. Invest.* 111:717-725 (2003)

**[0015]** [Non-patent Document 3] Nakamura Y. et al., *J. Hypertens.* September; 14 (9):1067-72 (1996)

**[0016]** [Non-patent Document 4] Taniyama Y., et al., *Gene Therapy* 8, 181-189 (2001)

### DISCLOSURE OF THE INVENTION

#### Problems to be Solved by the Invention

**[0017]** The present invention was achieved in view of the circumstances described above. An objective of the present invention is to provide novel uses of HGFs and genes encoding the same. More specifically, an objective of the present invention is to provide novel therapeutic agents and treatment

methods for lymphedema, a condition for which no effective therapeutic agent or treatment method is presently available.

#### Means for Solving the Problems

**[0018]** The present inventors conducted dedicated studies to achieve the objectives described above. The present inventors searched for factors that could reduce or eliminate lymphedema, and found that HGFs significantly reduced lymphedema. Then, the present inventors discovered that HGFs had lymphangiogenic activity to induce the growth of lymphatic endothelial cells.

**[0019]** As described above, there are many reports associating VEGF gene therapy with an increased risk of inducing edema. Thus, the present inventors considered that VEGF to be unsuitable for lymphedema treatment. Furthermore, since it is suggested that VEGF-C may enhance cell permeability by binding to VEGF receptors other than VEGF-3, the present inventors considered that the application of VEGF-C to lymphedema treatment also constituted a high risk.

**[0020]** HGF has been known as an angiogenic factor. VEGF is also known as an angiogenic factor, but is presumed to have neither lymphangiogenic activity (see, for example, Dev. Biol. 1997, 188: 96-109) nor the effect of reducing lymphedema, as described above. In view of the above, the novel findings by the present inventors that HGFs have lymphangiogenic activity and the effect of reducing lymphedema is a remarkable discovery.

**[0021]** To confirm the lymphangiogenic activity of HGFs based on its action mechanism, the present inventors demonstrated that the HGF receptor c-met was expressed in lymphatic endothelial cells and that phosphorylation of MAPK and Akt, which are intracellular signaling proteins whose phosphorylation is known to be induced by HGFs, was also induced in lymphatic endothelial cells.

**[0022]** These findings demonstrate that HGFs and their genes are effective as therapeutic or preventive agents for lymphedema, and as lymphangiogenesis-promoting agents. Specifically, the present invention provides the following [1] to [44]:

**[0023]** [1] a lymphangiogenesis-promoting agent comprising as an active ingredient an HGF or a protein or compound functionally equivalent to an HGF;

**[0024]** [2] the agent of [1], wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):

**[0025]** (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

**[0026]** (b) a protein encoded by a nucleic acid comprising the coding sequence in the nucleotide sequence of SEQ ID NO: 1,

**[0027]** (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

**[0028]** (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

**[0029]** [3] a lymphangiogenesis-promoting agent, which comprises as an active ingredient a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF;

**[0030]** [4] the agent of [3], wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

**[0031]** (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

**[0032]** (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

**[0033]** (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

**[0034]** (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

**[0035]** [5] the agent of [3] or [4], wherein the nucleic acid has been inserted into a mammalian expression vector;

**[0036]** [6] the agent of any one of [3] to [5], wherein the nucleic acid is a naked nucleic acid;

**[0037]** [7] the agent of any one of [1] to [6], which is used as a pharmaceutical agent for preventing or treating lymphedema;

**[0038]** [8] a method for promoting lymphangiogenesis, which comprises the step of administering to a subject an HGF or a protein or compound functionally equivalent to an HGF;

**[0039]** [9] the method of [8], wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):

**[0040]** (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

**[0041]** (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

**[0042]** (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

**[0043]** (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

**[0044]** [10] a method for promoting lymphangiogenesis, which comprises the step of administering to a subject a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF;

**[0045]** [11] the method of [10], wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

**[0046]** (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

**[0047]** (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

**[0048]** (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID

NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

[0049] (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

[0050] [12] the method of [10] or [11], wherein the nucleic acid has been inserted into a mammalian expression vector,

[0051] [13] the method of any one of [10] to [12], wherein the nucleic acid is a naked nucleic acid;

[0052] [14] a method for inducing activation of an HGF receptor and promoting lymphangiogenesis through the activation;

[0053] [15] a method for preventing or treating lymphedema, which comprises the step of administering to a subject an HGF or a protein or compound functionally equivalent to an HGF;

[0054] [16] the method of [15], wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):

[0055] (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

[0056] (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

[0057] (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

[0058] (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

[0059] [17] a method for preventing or treating lymphedema, which comprises the step of administering to a subject a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF;

[0060] [18] the method of [17], wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

[0061] (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

[0062] (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

[0063] (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

[0064] (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

[0065] [19] the method of [17] or [18], wherein the nucleic acid has been inserted into a mammalian expression vector;

[0066] [20] the method of any one of [17] to [19], wherein the nucleic acid is a naked nucleic acid;

[0067] [21] use of an HGF or a protein or compound functionally equivalent to an HGF for producing a lymphangiogenesis-promoting agent or a pharmaceutical agent to be used to prevent or treat lymphedema;

[0068] [22] the use of [21], wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):

[0069] (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

[0070] (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

[0071] (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

[0072] (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

[0073] [23] use of a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF for producing a lymphangiogenesis-promoting agent or a pharmaceutical agent to be used to prevent or treat lymphedema;

[0074] [24] the use of [23], wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

[0075] (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

[0076] (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

[0077] (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

[0078] (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

[0079] [25] the use of [23] or [24], wherein the nucleic acid has been inserted into a mammalian expression vector;

[0080] [26] the use of any one of [23] to [25], wherein the nucleic acid is a naked nucleic acid;

[0081] [27] a method of screening for a compound having lymphangiogenesis-promoting activity or a compound having an effect of preventing or treating lymphedema, wherein the method comprises the following steps:

[0082] (a) contacting a test compound with an HGF receptor or a protein functionally equivalent to an HGF receptor,

[0083] (b) detecting the binding between the protein and test compound, and

[0084] (c) selecting a test compound that binds to the protein,

[0085] [28] the method of [27], wherein the HGF receptor or protein functionally equivalent to an HGF receptor is selected from the following (i) to (iv):

[0086] (i) a protein comprising the amino acid sequence of SEQ ID NO: 4,

**[0087]** (ii) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 3,

**[0088]** (iii) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 4, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4, and

**[0089]** (iv) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 3, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4;

**[0090]** [29] a method of screening for a compound having lymphangiogenesis-promoting activity or a compound having an effect of preventing or treating lymphedema, wherein the method comprises the following steps:

**[0091]** (a) contacting a test compound with a cell expressing an HGF receptor or a protein functionally equivalent to an HGF receptor,

**[0092]** (b) measuring the growth capacity or migratory activity of the cell, or phosphorylation of a signaling molecule, and

**[0093]** (c) selecting a test compound that increases the growth capacity or migratory activity of the cell, or causes phosphorylation of the signaling molecule, as compared to when the test compound is not contacted;

**[0094]** [30] the method of [29], wherein the HGF receptor or protein functionally equivalent to an HGF receptor is selected from the following (i) to (iv):

**[0095]** (i) a protein comprising the amino acid sequence of SEQ ID NO: 4,

**[0096]** (ii) a protein encoded by a nucleic acid comprising the coding region of the nucleotide of SEQ ID NO: 3,

**[0097]** (iii) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 4, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4, and

**[0098]** (iv) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 3, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4;

**[0099]** [31] a vector for preventing or treating lymphedema, which is a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted;

**[0100]** [32] the vector of [31], wherein the nucleic acid encoding an HGF or the protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

**[0101]** (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

**[0102]** (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

**[0103]** (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

**[0104]** (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising

the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

**[0105]** [33] the vector of [31] or [32], which comprises the nucleotide sequence of SEQ ID NO: 5 or 6;

**[0106]** [34] the vector of any one of [31] to [33], which is administered in a naked state;

**[0107]** [35] the vector of any one of [31] to [34], which is administered by intramuscular injection to or around an affected area in a subject;

**[0108]** [36] a pharmaceutical agent for preventing or treating lymphedema, which comprises as an active ingredient a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted, wherein the vector is administered in a naked state by intramuscular injection to or around an affected area in a subject;

**[0109]** [37] the pharmaceutical agent of [36], wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

**[0110]** (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

**[0111]** (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

**[0112]** (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

**[0113]** (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;

**[0114]** [38] the pharmaceutical agent of [36] or [37], wherein the vector comprises the nucleotide sequence of SEQ ID NO: 5 or 6;

**[0115]** [39] a method for preventing or treating lymphedema, which comprises the step of administering, in a naked state, a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted, to or around an affected area in a subject by intramuscular injection;

**[0116]** [40] the method of [39], wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

**[0117]** (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

**[0118]** (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

**[0119]** (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

**[0120]** (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2;



[0121] [41] the method of [40], wherein the vector comprises the nucleotide sequence of SEQ ID NO: 5 or 6;

[0122] [42] use of a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted, for producing a pharmaceutical agent for preventing or treating lymphedema, wherein the vector is administered in a naked state to or around an affected area in a subject by intramuscular injection;

[0123] [43] the use of [42], wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

[0124] (a) a protein comprising the amino acid sequence of SEQ ID NO: 2,

[0125] (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1,

[0126] (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2, and

[0127] (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and

[0128] [44] the use of [42] or [43], wherein the vector comprises the nucleotide sequence of SEQ ID NO: 5 or 6.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0129] FIG. 1 depicts a series of photographs showing the results of immunofluorescence staining of lymphatic endothelial cells. The top panels correspond to staining with anti-c-met antibody; the middle panels correspond to nuclear staining with DAPI; and the bottom panels constitute merged images of the two. The results confirm that all cells expressed c-met.

[0130] FIG. 2 is a graph depicting the results on an MTS assay examining the effect of a recombinant human HGF on growth capacity. The vertical axis indicates the measured values. The results demonstrate that the HGF promotes the growth of lymphatic endothelial cells in a concentration-dependent manner.

[0131] FIG. 3 is a graph depicting the results of a migration assay examining the effect of a recombinant human HGF on migratory capacity. The vertical axis indicates the number of cells. The results demonstrate that the HGF promotes the migration of lymphatic endothelial cells.

[0132] FIG. 4 is a graph depicting the results of an MTS assay examining the effect of introducing a naked cDNA plasmid on growth capacity. The vertical axis indicates the measured values. The results demonstrate that the introduction of the HGF gene is effective to promote the growth of lymphatic endothelial cells.

[0133] FIG. 5 is a graph depicting the results of a c-fos promoter assay examining the effect of introducing a naked cDNA plasmid on growth capacity. The vertical axis indicates the measured values for luciferase activity. The results demonstrate that the introduction of the HGF gene is effective to promote the growth of lymphatic endothelial cells.

[0134] FIG. 6 is a graph depicting the results of a c-fos promoter assay examining the effect of introducing a naked cDNA plasmid on growth capacity. The vertical axis indicates

the measured values for luciferase activity. These results demonstrate that the expression vector for HGF cDNA significantly enhances the c-fos promoter activity while the expression vector for VEGF cDNA does not enhance the c-fos promoter activity.

[0135] FIG. 7 is a series of graphs demonstrating the effects of HGF on the growth and migratory activity of human lymphatic endothelial cells. FIG. 7A depicts the results of an MTS assay examining cell growth; and FIG. 7B depicts the results of a migration assay examining migratory activity. Each assay was carried out after the recombinant human HGF was added at a concentration of 0, 2, 10, or 50 ng/ml to the culture media of human lymphatic endothelial cells. In FIG. 7A, the vertical axis indicates the absorbance at 490 nm; and in FIG. 7B the vertical axis indicates the number of cells. These results demonstrate that the addition of HGF promotes the growth and migratory activity. \* $p < 0.001$  (relative to 0 ng/ml of the recombinant human HGF); † $p < 0.001$  (relative to 2 ng/ml of the recombinant human HGF).

[0136] FIG. 8 is a graph depicting the lymphedema-improving effect of the expression plasmid for HGF cDNA in rat models of lymphedema which have lymphedema at the base of their tail. The horizontal axis indicates time elapsed after surgery (days), and the vertical axis indicates the thickness of the base of tail (mm). The Venus group as a control, HGF group, and VEGF group were injected with 0.1 ml of 200  $\mu$ g expression plasmids for GFP, HGF, and VEGF, respectively. The physiological saline group was injected with 0.1 ml of physiological saline. The results demonstrate that the reduction of tail thickness is promoted by introducing the naked HGF cDNA and thus lymphedema is improved.

[0137] FIG. 9 is a graph obtained by calculating the area under the curve of the graph shown in FIG. 8. The results demonstrate that the tail thickness was significantly reduced only in the group of rats into which the HGF gene was introduced. The vertical axis indicates the area ratio (%) relative to the control. \*\* $p < 0.0001$  (relative to each of the group introduced with the VEGF gene, the Venus group, the physiological saline group, and the group treated with surgery alone).

[0138] FIG. 10 is a series of photographs and graphs depicting the results of examination on the lymphangiogenesis at the surgical sites after the introduction of the HGF or VEGF gene in rat models of lymphedema. FIG. 10A is composed of micrographs of immunostained sections of the tissue near the gene injection site in each rat. Immunostaining was carried out using antibodies against PECAM-1, LYVE-1, Prox1, and c-Met. The top panels are from rats into which the HGF gene was introduced; the middle panels are from rats into which the VEGF gene was introduced; and the bottom panels are from control rats injected with physiological saline. FIG. 10B is composed of graphs depicting the number of vessels that were positive for immunostaining with each antibody. HGF: rats into which the HGF gene was introduced; VEGF: rats into which the VEGF gene was introduced; control: rats injected with physiological saline. The vertical axis indicates the number of positive vessels. These results demonstrate that the number of vessels positive for the lymphatic endothelial cell markers (LYVE-1 and Prox1) is significantly increased only in the rats introduced with the HGF gene; whereas the number of vessels positive for the endothelial cell marker PECAM-1 is increased in both the rats introduced with the HGF gene and the rats introduced with the VEGF gene as compared to the control, but there is no significant difference. Furthermore, it is shown that the number of vessels positive for c-Met in the

rats introduced with the HGF gene is significantly increased as compared with that in the rats introduced with the VEGF gene and tends to be increased as compared to the control.

**[0139]** FIG. 11 is a series of photographs depicting phosphorylation of MAPK and Akt after HGF stimulation of canine thoracic duct-derived lymphatic endothelial cells. FIGS. 11A and 11B depict results of Western blotting using antibodies specific to phosphorylated MAPK and phosphorylated Akt, respectively (each upper panel). The lower panels show results of Western blotting using antibodies specific to MAPK and Akt regardless of their phosphorylation state. These antibodies were used as internal controls to demonstrate that samples contain almost equal amounts of MAPK or Akt.

#### BEST MODE FOR CARRYING OUT THE INVENTION

**[0140]** The present invention relates to lymphangiogenesis-promoting agents comprising, as active ingredients HGFs, or proteins or compounds (hereinafter sometimes referred to as “HGF”) functionally equivalent thereto. The present invention is based on the above-mentioned finding of the present inventors that HGFs activate the growth and migration of lymphatic endothelial cells and thereby promote lymphangiogenesis.

**[0141]** The present invention is further based on the finding that HGFs activate the growth and migration of lymphatic endothelial cells isolated from neither fetal nor neonatal but adult animals and thereby promote lymphangiogenesis. Since lymphatic endothelial cells are differentiated from fetal veins at an early stage of embryogenesis, lymphatic endothelial cell marker proteins are also expressed in venous endothelial cells at the early fetal stage. Lymphatic endothelial cells are also known to further differentiate into venous endothelial cells (Wigle J T and Oliver G Cell, 98: 769-778 (1999)). Therefore, fetal or neonatal cells are highly likely to be at a stage before terminal differentiation even when expressing lymphatic endothelial cell markers. Thus, fetal or neonatal lymphatic endothelial cells are unlikely to accurately reflect lymphatic endothelial cells of adult animals. The present invention demonstrated the effect of HGFs on lymphatic endothelial cells of adult animals. Thus, HGFs and their genes were found to be effective as therapeutic agents for patients (more particularly adults) suffering from lymphedema after the surgical removal of cancer tissues and/or lymph nodes for cancer treatment and for other lymphedema patients.

**[0142]** In one embodiment of the present invention, HGFs include proteins comprising the amino acid sequence of SEQ ID NO: 2. In another embodiment of the present invention, HGFs include proteins encoded by the “HGF gene”. Such a gene includes, for example, nucleic acids comprising the coding region of the nucleotide sequence of SEQ ID NO: 1.

**[0143]** The term “protein functionally equivalent to an HGF” in the context of the present invention encompasses proteins isolated from humans or nonhuman animals which have a biological activity equivalent to that of a protein comprising the amino acid sequence of SEQ ID NO: 2. For example, in a more specific embodiment, the proteins functionally equivalent to HGFs in the present invention include proteins comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2; and proteins encoded by nucleic acids that hybridize under stringent conditions to nucleic acids comprising the nucleotide

sequence of SEQ ID NO: 1, which are functionally equivalent to proteins comprising the amino acid sequence of SEQ ID NO: 2.

**[0144]** HGFs in the present invention may be human or nonhuman HGFs; however, human HGFs are most preferred. Those skilled in the art can obtain the amino acid sequences of nonhuman HGFs and the nucleotide sequences encoding them from various databases.

**[0145]** Furthermore, “proteins and compounds functionally equivalent to HGFs” can also include proteins and compounds that induce intracellular signaling by binding to c-met, an HGF receptor, and then activating intramolecular kinases of c-met. Thus, such proteins and compounds also include HGF agonists, for example, antibodies agonistically acting on c-met, which are disclosed in Prat, M., et al. (J. Cell Sci. 111 237-247 (1998)), and HGF mutants that bind to HGF receptors, which are disclosed in WO 98/51798.

**[0146]** Biological activity equivalent to that of a protein comprising the amino acid sequence of SEQ ID NO: 2 includes phosphorylation of tyrosines (Y<sup>1349</sup> and Y<sup>1356</sup>) in the cytoplasmic domain of c-met and/or phosphorylation of signaling molecules downstream thereof (see, for example, Graziani, A. et al., J. Biol. Chem. 266, 22087-22090 (1991); Graziani, A. et al., J. Biol. Chem. 268, 9165-9168 (1993); Nakagami, H. et al., Hypertension, 37 [part2]: 581-586 (2001)), and promotion of growth and migration of vascular endothelial cells and/or lymphatic endothelial cells; but is not limited thereto. The activity includes all biological activities of HGFs on cells.

**[0147]** The “proteins and compounds functionally equivalent to HGFs” in the present invention include all proteins and compounds having such an activity. Whether such compounds have a biological activity equivalent to that of HGFs can be determined by methods known to those skilled in the art. See, for example, Graziani, A. et al., J. Biol. Chem. 266, 22087-22090 (1991); Graziani, A. et al., J. Biol. Chem. 268, 9165-9168 (1993); and Nakagami, H. et al., Hypertension 37 [part2]: 581-586 (2001). Such determination as described above can be achieved by methods for assaying the growth capacity, migratory activity, and c-fos promoter activity of lymphatic endothelial cells herein described in the Examples; however, the methods are not limited thereto. Such methods also include methods in which cells expressing c-met are contacted with an HGF and then the phosphorylation of c-met or other downstream molecules in the HGF-c-met signaling pathway is detected in lysates of the cells. These methods are known to those skilled in the art. Furthermore, those skilled in the art can readily make appropriate modifications and improvements to known methods.

**[0148]** For example, HGF production in cells is promoted through the increase in the intracellular cyclic AMP concentration, and as a result the biological activity of HGFs is enhanced (Morishita R., et al., Diabetologia 40 (9):1053-61 (1997)). Accordingly, compounds and proteins such as cilostazol, a type 3 phosphodiesterase inhibitor, that increase the intracellular cyclic AMP concentration are also encompassed in the “proteins and compounds functionally equivalent to HGFs” of the present invention. In addition, antagonists to angiotensin II (Nakano N., et al., Hypertension 32 (3): 444-51 (1998)) that suppress HGF production (angiotensin II receptor antagonists) are also included.

**[0149]** Stringent hybridization conditions necessary to isolate the above-described nucleic acids that hybridize under stringent conditions to a nucleic acid comprising the nucle-

otide sequence of SEQ ID NO: 1 include the conditions of 6 M urea, 0.4% SDS, 0.5×SSC, and 37° C., and hybridization conditions of stringency equivalent thereto. Nucleic acids with higher homology can be expected to be isolated through use of more stringent conditions, for example, the conditions of 6 M urea, 0.4% SDS, 0.1×SSC, and 42° C. The sequences of isolated nucleic acids can be determined by the known methods described below. Homology of isolated nucleic acids over the entirety of their nucleotide sequences is at least 50% or higher, preferably 70% or higher, more preferably 90% or higher (for example, 95%, 96%, 97%, 98%, or 99% or higher) sequence identity.

**[0150]** As an alternative to the above-described methods using hybridization techniques, nucleic acids which hybridize under stringent conditions to nucleic acids comprising the nucleotide sequence of SEQ ID NO: 1, and which encode proteins functionally equivalent to proteins comprising the amino acid sequence of SEQ ID NO: 2, can be isolated by gene amplification method, for example, polymerase chain reaction (PCR) method, using primers synthesized based on the sequence information of a nucleic acid encoding an HGF (SEQ ID NO: 1).

**[0151]** Methods well known to those skilled in the art for preparing proteins functionally equivalent to a certain protein, include methods for introducing mutations into proteins. For example, those skilled in the art can prepare mutants functionally equivalent to HGFs by introducing appropriate mutations into the amino acid sequence of a human HGF using site-directed mutagenesis (Hashimoto-Gotoh, T, Mizuno, T, Ogasahara, Y, and Nakagawa, M. (1995) An oligodeoxyribonucleotide-directed dual amber method for site-directed mutagenesis. *Gene* 152, 271-275; Zoller, M J, and Smith, M. (1983) Oligonucleotide-directed mutagenesis of DNA fragments cloned into M13 vectors. *Methods Enzymol.* 100, 468-500; Kramer, W, Drutsa, V, Jansen, H W, Kramer, B, Pflugfelder, M, and Fritz, H J (1984) The gapped duplex DNA approach to oligonucleotide-directed mutation construction. *Nucleic Acids Res.* 12, 9441-9456; Kramer W, and Fritz H J (1987) Oligonucleotide-directed construction of mutations via gapped duplex DNA *Methods. Enzymol.* 154, 350-367; Kunkel, T A (1985) Rapid and efficient site-specific mutagenesis without phenotypic selection. *Proc Natl Acad Sci USA.* 82, 488-492) or the like. Amino acid mutations in proteins may also occur naturally. Thus, the proteins of the present invention also include proteins comprising an amino acid sequence with one or more amino acid mutations in the amino acid sequence of an HGF (SEQ ID NO: 2), which are functionally equivalent to HGFs.

**[0152]** When an amino acid residue is altered, the amino acid is preferably mutated to a different amino acid(s) that conserves the properties of the amino acid side-chain. Examples of amino acid side chain properties are the following: hydrophobic amino acids (A, I, L, M, F, P, W, Y, and V), hydrophilic amino acids (R, D, N, C, E, Q, G, H, K, S, and T), amino acids having aliphatic side chains (G, A, V, L, I, and P), amino acids having hydroxyl group-containing side chains (S, T, and Y), amino acids having sulfur-containing side chains (C and M), amino acids having carboxylic acid- and amide-containing side chains (D, N, E, and Q), amino acids having basic side chains (R, K, and H), and amino acids having aromatic side chains (H, F, Y, and W) (amino acids are represented by one-letter codes in parentheses). Amino acid substitutions within each group are called conservative substitutions. It is already known that a polypeptide comprising a

modified amino acid sequence in which one or more amino acid residues in a given amino acid sequence are deleted, added, and/or substituted with other amino acids can retain the original biological activity (Mark, D. F. et al., *Proc. Natl. Acad. Sci. USA* (1984) 81: 5662-6; Zoller, M. J. and Smith, M., *Nucleic Acids Res.* (1982) 10: 6487-500; Wang, A. et al., *Science* (1984) 224: 1431-3; Dalbadie-McFarland, G et al., *Proc. Natl. Acad. Sci. USA* (1982) 79: 6409-13). Such mutants have at least 70% amino acid sequence identity to the amino acid sequence of an HGF of the present invention, more preferably at least 75%, even more preferably at least 80%, still more preferably at least 85%, yet more preferably at least 90%, and most preferably at least 95% amino acid sequence identity thereto. Herein, sequence identity is defined as the percentage of residues identical to those in the original amino acid sequence of HGF, determined after the sequences are aligned as needed and gaps are appropriately introduced to maximize the sequence identity. The identity of amino acid sequences can be determined by the method described above.

**[0153]** Nucleotide sequence identity and amino acid sequence identity can be determined using the algorithm BLAST, by Karlin and Altschul (*Proc. Natl. Acad. Sci. USA* (1993) 90, 5873-7). Programs such as BLASTN and BLASTX were developed based on this algorithm (Altschul et al., *J. Mol. Biol.* (1990) 215, 403-10). To analyze nucleotide sequences according to BLASTN based on BLAST, the parameters are set, for example, as score=100 and wordlength=12. On the other hand, parameters used for the analysis of amino acid sequences by BLASTX based on BLAST include, for example, score=50 and wordlength=3. Default parameters for each program are used when using the BLAST and Gapped BLAST programs. Specific techniques for such analyses are known (see the website of the National Center for Biotechnology Information (NCBI), Basic Local Alignment Search Tool (BLAST); <http://www.ncbi.nlm.nih.gov>).

**[0154]** HGFs of the present invention can be isolated from natural sources including various biological samples, for example, cells or tissues expressing HGFs, based on their physical properties and the like. Alternatively, HGFs may be chemically synthesized based on known sequence information. In addition, HGFs can be obtained by using genetic recombination techniques to transform host cells with vectors carrying the genes encoding HGFs, then culturing the resulting transformed cells that produce the recombinant HGFs, and collecting the HGFs from the cells or culture supernatant.

**[0155]** Vectors suitable for producing HGFs using genetic engineering methods include various vectors using viruses, cosmid, plasmids, bacteriophages, and the like (*Molecular Cloning 2<sup>nd</sup> ed.*, Cold Spring Harbor Press (1989); *Current Protocols in Molecular Biology*, John Wiley & Sons (1987)). Such vectors include appropriate regulatory sequences, and an HGF-encoding nucleic acid is inserted so as to maintain the correct reading frame relative to the regulatory sequence, so that the HGF is expressed when the vectors are introduced into desired host cells. Any nucleic acids encoding HGF can be used, so long as they can be expressed by the selected vector and host. Such nucleic acids preferably include cDNAs; however, RNAs or the like may be used in some cases. When the host cell is a prokaryotic cell, the "regulatory sequence" includes a promoter, a ribosome-binding site, and a terminator. Alternatively, when the host is a eukaryotic cell, the regulatory sequence includes a promoter and terminator,

and as necessary, an enhancer, splicing signal, transcription factor, transactivator, poly A signal and/or polyadenylation signal, and so on. Such expression vectors for HGFs may include selection markers for easily selecting transformed host cells, as necessary. Furthermore, signal peptides may be inserted into vectors to be attached to HGFs so that intracellularly expressed HGFs are translocated into the lumen of the endoplasmic reticulum or translocated extracellularly, or alternatively translocated into the periplasm when the host cells are Gram-negative bacteria. Such signal peptides may be inherent in HGFs or may be derived from different proteins, as long as they are properly recognized in selected host cells. Furthermore, linkers, start codons, stop codons and such may be added, if required.

**[0156]** Genes can be inserted into vectors by ligase reactions using restriction enzyme sites (Molecular Cloning 2<sup>nd</sup> ed., Cold Spring Harbor Press (1989) Section 5.61-5.63; Current Protocols in Molecular Biology, John Wiley & Sons (1987) 11.4-11.11). Such vectors may be designed by considering codon usage in the host cells to be used, and selecting nucleotide sequences that allow high efficiency expression (Grantham et al., Nucleic Acids Res. (1981) 9, r43-74).

**[0157]** When vectors are introduced into adequate hosts to produce HGFs, the above expression vectors and appropriate hosts can be used in combination. Animal cells, plant cells, and fungal cells may be used as eukaryotic host cells.

**[0158]** Host cells can be transformed using methods suitable for selected hosts and vectors. For example, when prokaryotic cells are used as the hosts, known methods include calcium treatment and electroporation. Examples also include the *Agrobacterium* method for plant cells, and the calcium phosphate precipitation method for mammalian cells. The present invention is not particularly limited to these methods. Rather, the present invention can use various known methods, including nuclear microinjection, cell fusion, electroporation, protoplast fusion, lipofectamine methods (GIBCO BRL), DEAE-dextran methods, and methods using FuGENE6 reagent (Boehringer-Mannheim).

**[0159]** Host cells can be cultured by known methods suitable for selected cells. For example, when animal cells are used as the host, the cells may be cultured using a medium such as DMEM, MEM, RPMI-1640, 199, or IMDM, if required, supplemented with fetal calf serum (FCS) and such, at pH of about 6 to 8 at 30° C. to 40° C. for about 15 to 200 hours.

**[0160]** HGFs are preferably used after purification by known methods. HGFs can be purified to homogeneity by conventional protein purification methods. HGFs can be separated and purified, for example, by appropriately selecting and combining chromatographic columns, filters, ultrafiltration, salting out, dialysis, preparative polyacrylamide gel electrophoresis, isoelectric focusing and such (Strategies for Protein Purification and Characterization, A Laboratory Course Manual, Daniel R. Marshak et al., eds., Cold Spring Harbor Laboratory Press (1996)), but the present invention is not limited thereto.

**[0161]** Furthermore, as described above, HGFs of the present invention also include proteins and polypeptides in which amino acid residues are added to HGFs. Examples of proteins and polypeptides in which amino acid residues are added to HGFs include fusion proteins. To prepare a polynucleotide encoding such a fusion protein, for example, a DNA encoding an HGF may be linked in frame with a nucleic acid encoding another protein or polypeptide. The protein or

polypeptide to be fused with an HGF is not particularly limited. Any nucleic acid encoding an appropriate protein or polypeptide may be linked depending on the purposes.

**[0162]** The present invention also relates to novel uses of the following proteins, nucleic acids, or compounds:

- (1) HGFs, or proteins or compounds that are functionally equivalent thereto;
- (2) nucleic acids encoding HGFs or proteins functionally equivalent thereto;
- (3) nucleic acids encoding HGFs or proteins functionally equivalent thereto, which are inserted into mammalian expression vectors; and
- (4) naked nucleic acids encoding HGFs or proteins functionally equivalent thereto.

**[0163]** Specifically, the present invention relates to the following:

**[0164]** lymphangiogenesis-promoting agents that comprise as active ingredients the proteins, nucleic acids, or compounds listed in (1) to (4) above;

**[0165]** methods for preventing or treating lymphedema, which comprise the step of administering the proteins, nucleic acids, or compounds listed in (1) to (4) above to subjects; and

**[0166]** uses of the proteins, nucleic acids, and compounds listed in (1) to (4) above for producing lymphangiogenesis-promoting agents or agents used to prevent or treat lymphedema.

**[0167]** In this context, HGFs are preferably human HGFs.

**[0168]** The lymphangiogenesis-promoting agents of the present invention comprise as active ingredients the HGFs obtained as described above. The term "comprising an HGF as an ingredient" means comprising an HGF as at least one active ingredient, and content need not be limited. Furthermore, the lymphangiogenesis-promoting agents of the present invention may comprise other active ingredients that promote lymphangiogenesis, in addition to HGFs.

**[0169]** HGFs of the present invention can be formulated according to standard methods (see, for example, Remington's Pharmaceutical Science, latest edition, Mark Publishing Company, Easton, USA), and may comprise pharmaceutically acceptable carriers and/or additives. For example, the following can be comprised: detergents (for example, PEG and Tween), excipients, antioxidants (for example, ascorbic acid), coloring agents, flavoring agents, preservatives, stabilizers, buffering agents (for example, phosphoric acid, citric acid, and other organic acids), chelating agents (for example, EDTA), suspending agents, isotonicizing agents, binders, disintegrators, lubricants, fluidity promoters, and corrigents. However, the lymphangiogenesis-promoting agents of the present invention are not limited thereto, and may comprise other appropriate conventional carriers. Specific examples of such carriers include light anhydrous silicic acid, lactose, crystalline cellulose, mannitol, starch, carmellose calcium, carmellose sodium, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylacetaldihethylaminoacetate, polyvinylpyrrolidone, gelatin, medium chain fatty acid triglyceride, polyoxyethylene hydrogenated castor oil 60, sucrose, carboxymethylcellulose, corn starch, and inorganic salt. The agents may also comprise other low-molecular-weight polypeptides; proteins such as serum albumin, gelatin, and immunoglobulin; and amino acids such as glycine, glutamine, asparagine, arginine, and lysine. When the agents are prepared as aqueous solutions for injection, HGFs are dissolved in isotonic solutions containing, for example,

physiological saline, dextrose, and other adjuvants including, for example, D-sorbitol, D-mannose, D-mannitol, and sodium chloride. In addition, appropriate solubilizing agents such as alcohols (for example, ethanol), polyalcohols (for example, propylene glycol and PEGs), and non-ionic detergents (polysorbate 80 and HCO-50).

**[0170]** If necessary, HGFs may be encapsulated in microcapsules (e.g., microcapsules made of hydroxymethylcellulose, gelatin, and polymethylmethacrylate), and made into components of colloidal drug delivery systems (e.g. liposomes, albumin microspheres, microemulsions, nanoparticles, and nanocapsules) (for example, see "Remington's Pharmaceutical Science 16th edition", Oslo Ed. (1980)). Moreover, methods for preparing sustained-release drugs are known, and these can be applied to HGFs (Langer et al., J. Biomed. Mater. Res. (1981) 15, 167-277; Langer, Chem. Tech. (1982) 12, 98-105; U.S. Pat. No. 3,773,919; European Patent Application (EP) No. 58,481; Sidman et al., Biopolymers (1983) 22, 547-56; EP No. 133,988).

**[0171]** The lymphangiogenesis-promoting agents of the present invention can be administered to any mammal, including humans, rats, and dogs. These agents can be administered either orally or parenterally, but are preferably administered parenterally. Specifically, the agents can be administered to patients percutaneously or by injection. For example, injections can be locally administered by intravenous injection, intramuscular injection, or subcutaneous injection; however, they are preferably injected locally, particularly intramuscularly, in and around the areas where promotion of lymphangiogenesis is desired. Furthermore, the method of administration can be appropriately selected according to the age and symptoms of the patient

**[0172]** When an active ingredient is a protein, a single dose effective to promote lymphangiogenesis or to prevent or treat lymphedema can be selected from between 0.0001 to 10 mg per kg of body weight. Alternatively, when administered to human patients, the dose can be selected from between 0.001 to 100 mg/body, and the single dose preferably includes about 0.01 to 5 mg/body of HGF protein. However, the doses of the lymphangiogenesis-promoting agents of the present invention are not limited to these doses.

**[0173]** The present invention also relates to lymphangiogenesis-promoting agents comprising as active ingredients nucleic acids (hereinafter sometimes referred to as "the HGF genes") encoding HGFs or proteins functionally equivalent thereto. The terms "HGF" and "protein functionally equivalent to an HGF" of the present invention are as described above. Also in this context, an HGF is preferably a human HOE

**[0174]** Nucleic acids encoding HGFs of the present invention include cDNAs, genomic DNAs, and chemically synthesized DNAs. Moreover, nucleic acids encoding HGFs may include arbitrary sequences based on degeneracy of the genetic code, as long as they encode HGFs. Furthermore, nucleic acids encoding HGFs include, in addition to the DNAs described above, derivatives thereof and artificially modified nucleic acids. Such artificially modified nucleic acids include, for example, DNAs in which their sugar chain structures have been modified, cDNAs, genomic DNAs, chemically synthesized DNAs, and derivatives thereof, but are not limited thereto.

**[0175]** Genomic DNAs and cDNAs can be prepared by conventional methods known to those skilled in the art. Genomic DNAs can be prepared, for example, by extracting

genomic DNAs from human-derived cells, preparing genomic libraries (plasmids, phages, cosmids, BAC, PAC, and the like can be used as vectors), developing them, and carrying out colony or plaque hybridization using probes prepared based on nucleic acids encoding HGFs (for example, the nucleic acid of SEQ ID NO: 1). Alternatively, the DNAs can also be prepared by preparing primers specific to HGF-encoding DNAs and performing PCR using the primers. In addition, cDNAs can be prepared, for example, by synthesizing cDNAs based on mRNAs extracted from human-derived cells, preparing cDNA libraries in which the cDNAs are inserted into vectors such as  $\lambda$ ZAP, developing them, and carrying out colony or plaque hybridization as described above, or performing PCR.

**[0176]** Nucleic acids encoding HGFs or proteins functionally equivalent thereto that are comprised as active ingredients in the lymphangiogenesis-promoting agents of the present invention may be naked nucleic acids, or may be included in viral envelopes, liposomes or the like. The term "naked nucleic acid" refers to a nucleic acid in a state where the nucleic acid molecule is not present in an inclusion body, such as a viral envelope or liposome, but is present in an aqueous solution in a naked form without being coated.

**[0177]** Nucleic acids encoding HGFs or proteins functionally equivalent thereto are preferably incorporated into nucleic acid vectors. Any nucleic acid vector can be used in the present invention, as long as the nucleic acid encoding an HGF of the present invention that is inserted into the vector can be expressed in mammals. Such vectors include, for example, plasmids; viral vectors such as retroviral vectors, adenoviral vectors, adeno-associated virus vectors, vaccinia virus vectors, lentiviral vectors, herpes virus vectors, alphavirus vectors, EB virus vectors, papilloma virus vectors, and foamy virus vectors; and non-viral vectors (Niitsu Y. et al., Molecular Medicine 35: 1385-1395 (1998)), but are not limited thereto. Vectors in the present invention are suitable for use as vectors for gene therapy.

**[0178]** Those skilled in the art can appropriately design and use desired vectors. Vectors used in the present invention may include polynucleotide sequences that allow more efficient expression of HGFs, such as a transcription initiation region and transcription termination region that function in expression hosts, in addition to an arbitrary polynucleotide to be introduced.

**[0179]** Preferred forms of nucleic acids encoding HGFs used in the present invention include, for example, a nucleic acid encoding an HGF that has been inserted into a plasmid vector, such as pVAX1HGF/MGB1 described in SEQ ID NO: 5 or pcDNA3.1(-)HGF described in SEQ ID NO: 6.

**[0180]** Nucleic acids of the present invention that encode HGFs or proteins functionally equivalent to HGFs may be incorporated into vectors, such as cationic liposomes, ligand-DNA complexes, and gene guns. The nucleic acids may be, for example, in a form where they are packaged in HVJ-E vectors derived from Sendai virus envelopes (Kaneda, Y. et al., Mol. Ther. 6, 219-226 (2002)). Alternatively, the nucleic acids may be in a naked form.

**[0181]** In the present invention, the HGF genes may be used alone or in combination with other genes and/or other pharmaceutical agents as needed. In the present invention, the HGF gene and the genes used in combination as needed are incorporated into appropriate vectors that ensure the in vivo expression of the genes, and then administered to affected areas of patients.

**[0182]** When lymphangiogenesis-promoting agents comprising as active ingredients nucleic acids encoding HGFs or proteins functionally equivalent thereto are administered, those skilled in the art can readily select an adequate administration route and dose. However, the agents are preferably administered by injection to and/or around the areas where promotion of lymphangiogenesis is desired, in particular by intramuscular injection to the muscles in and/or around the areas. Other administration methods may be used, as long as HGFs or proteins functionally equivalent thereto can be expressed from the administered nucleic acids in and around the areas where lymphangiogenesis is desired to be promoted.

**[0183]** When an active ingredient is a nucleic acid described above, the single dose of the nucleic acid that is effective to promote lymphangiogenesis or to prevent or treat lymphedema can be selected, for example, from the range of 0.001 to 10 mg per kg body weight. Alternatively, when administered to human patients, the dose of the nucleic acid can be selected, for example, from the range of 0.001 to 50 mg/body, and the single dose of the nucleic acid is preferably about 0.5 to 50 mg/body. However, the dose is not limited to those described above.

**[0184]** In a preferred embodiment of the lymphangiogenesis-promoting agents comprising as active ingredients nucleic acids encoding HGFs or proteins functionally equivalent thereto, for example, when the agent is administered to a human, a plasmid including a human HGF cDNA (for example, pVAX1HGF/MGB1 described in SEQ ID NO: 5 or pcDNA3.1(-)HGF described in SEQ ID NO: 6) dissolved in a physiological buffer or pharmaceutically acceptable carrier is intramuscularly injected to the muscle in and/or around an area where lymphangiogenesis is desired to be promoted in a human patient at a single dose of 1 mg to 20 mg, preferably 1 mg to 10 mg, and more preferably 2 mg to 10 mg (for example, 3 mg to 6 mg, or 2 mg to 4 mg). The agent may be administered twice or more, preferably three times or more, at intervals of about 1 to 10 weeks, preferably about 1 to 8 weeks (for example, 2 to 6 weeks, or 1 to 4 weeks). When administered to humans, the agents may be administered, for example, according to the method described by Morishita R. et al., *Hypertension* 44 (2):203-9 (2004), or an appropriately modified method thereof.

**[0185]** The lymphangiogenesis-promoting agents of the present invention can be used as pharmaceutical agents for preventing or treating lymphedema. Lymphedema refers to a condition where occlusion of lymphatic vessels causes abnormal congestion of tissue fluid, resulting in swelling, chronic inflammation, and/or fibrosis. Lymphedema that is a target for prevention or treatment by the lymphangiogenesis-promoting agents of the present invention include all conditions that have such symptoms as described above, regardless of their names. The lymphangiogenesis-promoting agents of the present invention are useful for preventing or treating diseases involving such symptoms.

**[0186]** In subjects at high risk for lymphedema (for example, patients whose lymph nodes have been extirpated along with malignant tumors by surgery), lymphedema can be prevented by administering the lymphangiogenesis-promoting agents of the present invention.

**[0187]** The present invention also relates to methods for preventing or treating lymphedema, which comprise the step of administering to subjects, HGFs, proteins or compounds functionally equivalent to HGFs, or nucleic acids encoding HGFs or proteins functionally equivalent thereto. Herein,

HGFs are preferably human HGFs. The administration site may be any site where lymphangiogenesis is desired to be promoted. Lymphedema can also be treated, for example, by applying the above-described methods to patients with lymphedema. More specifically, lymphangiogenesis can be promoted by administering the lymphangiogenesis-promoting agents, as described above.

**[0188]** Those skilled in the art can administer human HGFs or the human HGF genes to affected areas of patients with lymphedema, appropriately considering the purpose. The administration to the affected areas of patients can be achieved by methods known to those skilled in the art.

**[0189]** In the present invention, the proteins, nucleic acids, and compounds described above in (1) to (4) may be administered alone or in combination with genes encoding other lymphangiogenic factors, or as compositions in combination with pharmaceutically acceptable carriers and/or additives. When administered in a form of composition, specific embodiments of genes encoding other angiogenic factors, and pharmaceutically acceptable carriers and/or additives to be used in combination are as described above.

**[0190]** Affected areas to which the proteins, nucleic acids, and compounds described above in (1) to (4) are not particularly limited, so long as they are areas in which the symptoms of lymphedema have appeared or are predicted to appear. For example, local administration to the upper arm region can be considered for lymphedema of upper limb, which is often observed after breast cancer surgery and such; alternatively, local administration to the femoral region can be considered for lymphedema of lower limb, which is often observed after uterine cancer surgery and such.

**[0191]** The dose is as described above; however, it varies depending on the type of disease, patient's weight, age, sex, and symptoms, administration purpose, form of the gene to be introduced and such. Those skilled in the art can appropriately determine the dose.

**[0192]** The subjects to which compositions, including proteins, nucleic acids, or compounds of the present invention, are administered include any mammal, such as a monkey, dog, and cat, in addition to human.

**[0193]** Furthermore, in an embodiment of the present invention, the methods for promoting lymphangiogenesis also include methods in which activation of HGF receptors is induced and lymphangiogenesis is promoted through such activation.

**[0194]** HGF receptors are membrane proteins named c-met, and have a tyrosine kinase in their cytoplasmic domain. When an HGF binds to the HGF binding site in the extracellular domain of c-met, the above-mentioned cytoplasmic tyrosine kinase is activated and phosphorylates tyrosine residues (Y<sup>1349</sup> and Y<sup>1356</sup>) in the c-met molecule. This triggers activation of the intracellular signaling pathway and then the HGF exerts its biological function. This mechanism is well-known in the art (see, for example, Graziani, A. et al., *J. Biol. Chem.* 266, 22087-22090 (1991); Graziani, A. et al., *J. Biol. Chem.* 268, 9165-9168 (1993); and Nakagami, H. et al., *Hypertension*, 37 [part2]: 581-586 (2001)). Thus, herein, the activation of an HGF receptor refers to the activation of tyrosine kinase in the c-met molecule.

**[0195]** In the present invention, lymphangiogenesis can be promoted by activating c-met expressed in lymphatic endothelial cells. The c-met activation is induced by reacting HGFs, or proteins or compounds functionally equivalent thereto with c-met on lymphatic endothelial cells. Thus, the

lymphangiogenesis-promoting agents described above can also be utilized in this embodiment.

**[0196]** The present invention relates to methods of screening for compounds having lymphangiogenesis-promoting activity or compounds having an effect of preventing or treating lymphedema.

**[0197]** In an embodiment of the present invention, the screening methods are methods of screening for compounds having lymphangiogenesis-promoting activity, which comprise the following steps of (a) to (c):

- (a) contacting test compounds with HGF receptors or proteins functionally equivalent thereto;
- (b) detecting the binding of test compounds to HGF receptors or proteins functionally equivalent thereto; and
- (c) selecting test compounds that bind to HGF receptors or proteins functionally equivalent thereto.

**[0198]** In the first embodiment, HGF receptors or proteins functionally equivalent thereto are first contacted with test compounds. The HGF receptors and proteins functionally equivalent thereto in the screening methods of the present invention include c-met and proteins functionally equivalent thereto. Specifically, such proteins include the following (i) to (iv):

- (i) proteins having the amino acid sequence of SEQ ID NO: 4;
- (ii) proteins encoded by nucleic acids that include the coding region of the nucleotide sequence of SEQ ID NO: 3;
- (iii) proteins which have an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 4, and which are functionally equivalent to proteins having the amino acid sequence of SEQ ID NO: 4; and
- (iv) proteins encoded by nucleic acids that hybridize under stringent conditions to nucleic acids having the nucleotide sequence of SEQ ID NO: 3, which are functionally equivalent to proteins having the amino acid sequence of SEQ ID NO: 4.

**[0199]** The “test compounds” in the methods of the present invention are not particularly limited, and include, for example, single compounds such as natural compounds, organic compounds, inorganic compounds, proteins, and peptides; as well as compound libraries, expression products of gene libraries, cell extracts, cell culture supernatants, products of fermentation microorganisms, marine organism extracts, plant extracts, prokaryotic cell extracts, unicellular eukaryote extracts, and animal cell extracts. If needed, the above test compounds can be appropriately labeled before use. Labels include, for example, radiolabels and fluorescent labels.

**[0200]** In the present invention, “contacting” is achieved by the procedure described below. For example, “contacting” can be achieved by adding a test compound to a culture medium of cells expressing an HGF receptor or a protein functionally equivalent thereto or an extract of cells expressing an HGF receptor or a protein functionally equivalent thereto. When the test compound is a protein, “contacting” can be achieved, for example, by introducing a vector carrying a DNA encoding the protein into cells expressing an HGF receptor or a protein functionally equivalent thereto, or by adding the vector to an extract of cells expressing an HGF receptor or a protein functionally equivalent thereto. Alternatively, for example, the two hybrid method with yeast, animal cells or the like can be used.

**[0201]** In the first embodiment, the binding between the test compounds and HGF receptors or proteins functionally equivalent thereto is subsequently detected. Detection or

measurement of the binding between proteins can be carried out by using, for example, labels attached to the proteins. The types of labels include, fluorescent labels and radiolabels, for example. The binding can also be measured by known methods such as the enzyme two hybrid method and the measurement method using BIACORE. In the present methods, the test compounds bound to the above-mentioned HGF receptors are then selected. The selected test compounds include compounds having lymphangiogenesis-promoting activity or compounds having an effect of preventing or treating lymphedema. In addition, the selected test compounds may be used as test compounds in the following screening.

**[0202]** Furthermore, in the second embodiment of the screening methods, the present invention provides methods of screening for compounds having lymphangiogenesis-promoting activity or compounds having an effect of preventing or treating lymphedema, which comprise the following steps (a) to (c):

- (a) contacting test compounds with cells expressing HGF receptors or proteins functionally equivalent thereto;
- (b) measuring the growth capacity or migratory activity of the cells, or phosphorylation of signaling molecules;
- (c) selecting test compounds that increase the growth capacity or migratory activity of the cells, or that cause phosphorylation of signaling molecules, as compared to when the test compounds are not contacted.

**[0203]** In the second embodiment, test compounds are first contacted with cells expressing HGF receptors or proteins functionally equivalent thereto. The “cells expressing HGF receptors or proteins functionally equivalent thereto” include isolated cells expressing HGF receptors or proteins functionally equivalent thereto and transformed cells expressing recombinant HGF receptors, but are not limited thereto. These cells include, for example, lymphatic endothelial cells derived from thoracic duct. HGF receptors or proteins functionally equivalent thereto used in the instant methods include the HGF receptors described above and proteins functionally equivalent thereto.

**[0204]** In the second embodiment, then, cells expressing HGF receptors or proteins functionally equivalent thereto are measured for their growth capacity or migratory activity, or phosphorylation of signaling molecules such as MAPK, Akt, c-met, and Ras, which are known to be phosphorylated by the c-met signaling.

**[0205]** Those skilled in the art can readily measure the degree of an increase in growth capacity and migratory activity, and the phosphorylation of MAPK or Akt, for example, by the methods described below in Examples.

**[0206]** In the screening methods of the present invention, the final step is selection of compounds that increase the growth capacity or migratory activity of the cells expressing HGF receptors or proteins functionally equivalent thereto, or that cause phosphorylation of molecules such as MAPK, Akt, c-met, and Ras, which are known to be phosphorylated by the c-met signaling, as compared to when the test compounds are not contacted. The compounds thus selected can be used as active ingredients of the lymphangiogenesis-promoting agents of the present invention or the pharmaceutical agents of the present invention for preventing or treating lymphedema.

**[0207]** All prior art documents cited herein are incorporated herein by reference.

## EXAMPLES

[0208] Herein below, the present invention will be specifically described with reference to Examples, but is not to be construed as being limited thereto.

## Example 1

## Effect of HGFs on Primary Cultured Lymphatic Endothelial Cells Derived from Canine Thoracic Duct

[0209] (1) Preparation of Primary Cultured Lymphatic Endothelial Cells Derived from the Canine Thoracic Duct

[0210] Primary cultured lymphatic endothelial cells derived from the canine thoracic duct were prepared by a known method (Microcirculation 6, 75-78 (1999)). Adult mongrel dogs (including both male and female; 6 to 12 kg) were sacrificed by bleeding from the femoral artery under anesthesia. Then, the thoracic ducts (10 to 15 cm) were isolated and placed in cold (4° C.) Hanks' balanced salt solution (HBSS). Connective tissues and adipocytes were removed. Branches of the thoracic ducts were ligated with sterilized silk thread, and the lymphatic vessels were washed with cold HBSS. Next, the thoracic ducts were incubated in a collagenase solution (250 U/ml HBSS) at 37° C. for 10 minutes. The ducts were washed with MEM containing 10% fetal bovine serum (FBS) and the cells were collected. After collagenase was removed by centrifugation, the cells were suspended in complete MEM supplemented with 20% FBS and an antibiotic, and then cultured in 35-mm dishes coated with type I collagen. The cells were cultured under a wet condition with 5% CO<sub>2</sub> at 37° C. When the cells reached confluence, they were detached with trypsin/EDTA and collected. Lymphatic endothelial cells were subcloned to prepare lymphatic endothelial cell clones. The clonal cells were able to be passaged at least 10 or more times.

[0211] The lymphatic endothelial cells thus obtained were confirmed to be immunostained with different types of lymphatic endothelial cell-specific antibodies (including anti-VEGFR-3 antibody and anti-Prox1 antibody). Furthermore, the expression of c-met, an HGF receptor, in these cells was confirmed by immunostaining. This novel finding is attributable to the present invention (FIG. 1).

(2) Enhancement of Growth Capacity by a Recombinant Human HGF

[0212] A recombinant human HGF was added at a concentration of 0, 2, 10, or 50 ng/ml to medium containing the lymphatic endothelial cells derived from the canine thoracic duct (70% or more confluent in 96-well plates) prepared above in (1). After three days, the MTS assay was carried out using the CellTiter 96 One Solution Reagent (Promega). The assay was independently performed on nine wells for each concentration. The results are shown in Table 1 and FIG. 2.

TABLE 1

	HGF 0 ng/ml	HGF 2 ng/ml	HGF 10 ng/ml	HGF 50 ng/ml
No1	0.22	0.257	0.339	0.285
No2	0.203	0.26	0.291	0.299
No3	0.203	0.279	0.344	0.341
No4	0.183	0.248	0.251	0.25
No5	0.208	0.262	0.3	0.301

TABLE 1-continued

	HGF 0 ng/ml	HGF 2 ng/ml	HGF 10 ng/ml	HGF 50 ng/ml
No6	0.193	0.247	0.338	0.406
No7	0.26	0.367	0.403	0.477
No8	0.288	0.417	0.499	0.503
No9	0.289	0.387	0.549	0.437
Mean	0.227	0.303	0.368	0.367
Standard deviation (SD)	0.041	0.068	0.099	0.092
Standard error (SE)	0.014	0.023	0.033	0.031

[0213] Even in the presence of 2 ng/ml of HGF, the measured values significantly increased. These results suggest that the HGF strongly promotes the growth of lymphatic endothelial cells.

(3) Enhancement of Migratory Activity by a Recombinant Human HGF

(i) Enhancement of Migratory Activity by a Recombinant HGF

[0214] The migratory activity of cells was measured by a previously reported method using the Boyden chamber (Arterioscler Thromb Vasc Biol. 22:108-114 (2002)). A polyvinylpyrrolidone (PVP)-free polycarbonate membrane with a pore size of 8 µm (Neuro Probe Inc., Gaithersburg, Md.) was coated with 0.1% gelatin and excess gelatin was removed using phosphate buffered saline (PBS). The above-described membrane was placed onto the lower chamber of the Boyden chamber containing 28 µl of EBM2 medium supplemented with 1% FBS, and 50 µl of medium containing 10<sup>6</sup> of the lymphatic endothelial cells derived from the canine thoracic duct prepared above in (1) was added to the upper chamber. A recombinant human HGF was added to the medium at a concentration of 0, 10, or 50 ng/ml. The cells were cultured under a wet condition with 5% CO<sub>2</sub> at 37° C. for four hours. The membrane was removed, and the cells on the upper surface of the membrane were detached. The cells on the lower surface of the membrane were stained with the Diff-Quick (Sysmex, Hyogo, Japan) and counted. This assay was independently performed in three wells for each HGF concentration. The results are shown in Table 2 and FIG. 3

TABLE 2

	HGF (—)	HGF 10 ng/ml	HGF 50 ng/ml
No1	3	224	271
No2	4	238	307
No3	6	222	254
Mean	4.33	228.00	277.33
SD	1.53	8.72	27.06
SE	0.0882	5.033	15.624

[0215] Table 2 shown above and FIG. 3 suggest that even in the presence of 10 ng/ml of HGF, the measured values significantly increases and thus the HGF promotes migration of lymphatic endothelial cells.

[0216] The above results suggest that HGFs promote lymphangiogenesis by promoting the growth and migration of lymphatic endothelial cells.



## (ii) Enhancement of Migratory Activity by a Paracrine Recombinant HGF

[0217] An expression plasmid for human HGF cDNA, pVAX1HGF/MGB1 (SEQ ID NO: 5), was packaged in the HVJ-E vector by a previously reported method (Kaneda, Y. et al., Mol. Ther. 6, 219-226 (2002)).

[0218] BHK cells were cultured until they reached confluence in 100-mm dishes. 50 HAU of the vector containing the plasmid was added to the culture medium of the BHK cells and then introduced into the cells. After 24 hours of culture, the cells were transferred into inserts for 24-well plates and cultured until they were almost confluent.

[0219] Meanwhile, the lymphatic endothelial cells derived from the canine thoracic duct described above in (1) were cultured until 70% or more confluent in 24-well plates, and the inserts containing the above-described BHK cells introduced with the expression plasmid for human HGF cDNA pVAX1HGF/MGB1 were placed in the wells. After 48 hours of culture, in the same way as described above, the cells were subjected to the MTS assay. The assay was independently performed on six wells for each sample. The cells treated in the same way using a GFP expression plasmid instead of the expression plasmid for human HGF cDNA pVAX1HGF/MGB1, were used as a negative control. The results are shown in Table 3 and FIG. 4.

TABLE 3

	GFP	HGF
No1	0.260	0.497
No2	0.320	0.451
No3	0.338	0.397
No4	0.272	0.414
No5	0.306	0.440
No6	0.307	0.400
Mean	0.301	0.433
SD	0.029	0.038
SE	0.012	0.016

[0220] When the cells were co-cultured with the cells introduced with the HGF cDNA, the measured values in the MTS assay were significantly increased as compared to the negative control. These results demonstrate that the introduction of the HGF gene is effective to promote the growth of lymphatic endothelial cells.

[0221] To further confirm the promotion of lymphatic endothelial cell growth by the HGF, c-fos promoter activity was measured by a previously reported method (Hypertension 37 (2); 581-586 (2001)).

[0222] The assay was independently performed on six wells for each sample. The cells treated using a GFP expression plasmid in the same way as described above, were used as a negative control. The results are shown in Table 4 and FIG. 5.

TABLE 4

	GFP	HGF
No1	231101	317234
No2	210645	344423
No3	143101	303133
No4	141224	319125
No5	195948	384934
No6	189940	270410
Mean	185326.50	323209.83

TABLE 4-continued

	GFP	HGF
SD	36327.55	38738.02
SE	14830.66	15814.73

[0223] The c-fos promoter activity is known to be positively correlated with the cell growth capacity. Thus, given the fact that the HGF increases the c-fos promoter activity, it is apparent that the introduction of the HGF gene is effective to promote the lymphatic endothelial cell growth.

(5) Comparison of Growth Promoting Effects Between a Human HGF cDNA and VEGF cDNA in an Autocrine System

[0224] Using Lipofectamine Plus (GIBCO-BRL), an expression plasmid for a naked human HGF cDNA, pVAX1HGF/MGB1, and a c-fos luciferase reporter gene plasmid (p2FTL) for c-fos promoter assay were introduced into the above-described lymphatic endothelial cells cultured until subconfluent in 6-well plates (J. Hypertens. 16: 993-1000 (1998)). After 24 hours of culture, the cells were cultured in serum-free culture medium for 24 hours. The luciferase activity was then determined by a conventional method, and found to be increased. Next, using the human VEGF cDNA instead of the human HGF cDNA, the same experiments as described above in (4) were carried out to compare the effects of human HGF cDNA and VEGF cDNA on the c-fos promoter activity. The assay was independently performed on four wells for each sample. The results are shown in Table 5 and FIG. 6.

TABLE 5

	GFP	VEGF	HGF
No1	28135	9213	120035
No2	19431	14273	80231
No3	6789	3728	46335
No4	9085	2936	39688
Mean	15860.00	7537.50	71572.25
SD	9859.20	5287.10	36865.23
SE	4929.599	2643.551	18432.617

[0225] These results suggest that introduction of the expression plasmid for HGF cDNA significantly increases the c-fos promoter activity through the autocrine of HGF, while the expression plasmid for VEGF cDNA does not increase the c-fos promoter activity and thus does not promote the lymphatic endothelial cell growth.

[0226] The results of experiments using the primary cell culture system described above suggest that lymphangiogenesis is promoted by HGFs, that lymphangiogenesis can be promoted by introducing the HGF gene, and that the introduction of an expression plasmid for a naked HGF cDNA, namely the introduction of a nucleic acid encoding an HGF, is effective for the promotion.

[0227] Furthermore, lymphatic endothelial cells isolated from adult dogs were used in the present Example, and thus HGFs were demonstrated to have the above-described activity on lymphatic endothelial cells of adult animals. This suggests that HGFs are useful as therapeutic agents for lymphedema.

[0228] Moreover, in order to confirm that the lymphangiogenesis-promoting activity of human HGF was not specific to

canine lymphatic endothelial cells, the growth-promoting activity of HGF on primary cultured aortic and venous endothelial cells isolated from adult mongrel dogs in the same way as described above was investigated. As a result, the human HGF was found to have growth-promoting activity on these cells. Furthermore, this growth-promoting activity was comparable to that on equivalent human cells (data not shown). Consequently, human HGFs also act on canine vascular endothelial cells in the same manner as on human vascular endothelial cells. These findings suggest that the lymphangiogenesis-promoting activity of human HGFs is not specific to canine lymphatic endothelial cells.

#### (6) Confirmation of HGF Activity on Human Lymphatic Endothelial Cells

**[0229]** It was postulated that human lymphatic endothelial cells would give the same result as that obtained using the lymphatic endothelial cells isolated from adult dogs. To confirm this prediction, the effect of HGF on the growth and migratory capacities of human lymphatic endothelial cells was assayed (AngioBio Co. (Del Mar, Calif.)).

**[0230]** The cells at passage 5 to 8 were used in the experiments. By immunostaining, it was confirmed that von Willebrand factor, VEGF receptor-3, Prox1, and c-Met, which are lymphatic endothelial cell markers, were also expressed in the cells (data not shown). The MTS assay and migration assay were carried out in the same way as the experiments using canine lymphatic endothelial cells.

**[0231]** The results of MTS assay and migration assay are shown in FIGS. 7A and 7B, respectively. Both cell growth capacity and migratory capacity were increased by adding a recombinant human HGF. They were significantly increased at 10 ng/ml or higher concentrations as compared to the cells in the absence of HGF.

**[0232]** The results confirm the initial prediction that an HGF would promote lymphangiogenesis of human lymphatic endothelial cells as well as the lymphatic endothelial cells from adult dogs described above in (1) to (5).

**[0233]** The above results suggest that HGFs have lymphangiogenesis-promoting activity not only on dogs but also on other mammals (including humans).

### Example 2

#### Effect of HGFs on Rat Models of Lymphedema

**[0234]** Based on the findings described above, the effect of an HGF on lymphedema was tested using rat models to demonstrate the in vivo effect.

#### (1) Preparation of Rat Models of Lymphedema

**[0235]** Rat models of Lymphedema were prepared according to Slavin S A et al., (Anals of Surgery 229, 421-427 (1999)). To confirm the presence of lymphedema in the tail of rat models, physiological saline containing 0.5% Patent Blue dye was injected at a position several centimeters distant from the base of tail immediately before surgery. The tail region was dissected on the day following the surgery, and the blue dye was observed in the lymphatic vessels (data not shown). Furthermore, the base of tail was evidently thicker as compared to control rats, and thus the above rats were demonstrated to be useful as lymphedema models.

#### (2) Lymphedema-Improving Effect of an Expression Plasmid for HGF cDNA on Rat Models of Lymphedema

**[0236]** The effect of introduction of the human HGF gene on lymphedema was investigated using the rat models prepared as described above.

**[0237]** 200 µg (/100 µl) of an expression plasmid for a naked human HGF cDNA, pVAX1HGF/MGB1, and 200 µg (/100 µl) of an expression plasmid for a naked VEGF cDNA, and as a control, 200 µg (/100 µl) of a naked GFP expression plasmid (Venus plasmid) were intramuscularly administered one, seven, and 14 days after surgery. In the group with surgery alone, only surgery was carried out without injection. In the group without surgery, no surgery was carried out. In the physiological saline group, 100 µl of physiological saline was intramuscularly injected one, seven, and 14 days after surgery. The tail thickness in each group was measured every seven days up to day 35 after surgery. Five rats in each group were tested and the mean was determined, which is shown in FIG. 8.

**[0238]** In all groups, the tail thickness transiently increased after surgery. However, the tail thickness was more rapidly decreased and the degree was greater only in the group introduced with HGF cDNA as compared to the other groups. This difference was significant after day 21. Thus, lymphedema was found to be improved by introducing the naked HGF cDNA plasmid.

**[0239]** The areas under the curves shown in FIG. 8 were determined. The result is shown in FIG. 9. There was no difference between rats introduced with the VEGF gene and rats in the control Venus group; however, the tail thickness was found to be significantly decreased in rats introduced with the HGF gene. This decrease in the thickness of tail affected with lymphedema implies the relief or cure of lymphedema. Thus, lymphedema was clearly demonstrated to be relieved or cured by introducing a nucleic acid encoding an HGF (the HGF gene).

**[0240]** Tissue samples were collected from the above-described surgical site in the rat tails on days 4, 10, and 17 after surgery, and human HGF mRNA level was determined by real-time RT-PCR using a conventional method. As a result, human HGF expression was confirmed up to day 17 after surgery only in the group introduced with the human HGF gene (data not shown).

**[0241]** Furthermore, in the rats introduced with the HGF gene, rats introduced with the VEGF gene, and control rats injected with physiological saline, expression of endothelial cell marker (PECAM-1), lymphatic endothelial cell markers (LYVE-1 and Prox1), and c-met was detected by immunostaining at the injection sites on day 35 after surgery. Typical staining images are shown in FIG. 10 (FIG. 10A). In addition, the expression level of each marker was determined by counting the number of immunostaining-positive vessels in microscopic fields randomly selected using a known method (Yoon Y S, et al., J. Clin. Invest. 111: 717-725 (2003)) (FIG. 10B). This result showed that there was no difference in the number of vessels positive for an endothelial cell marker, PECAM-1, in both rats introduced with the HGF gene and rats introduced with the VEGF gene as compared to the control. On the other hand, the numbers of vessels positive for lymphatic endothelial cell markers, LYVE-1 and Prox1, were both significantly increased in the rats introduced with the HGF gene, while there was no difference between the control and rats introduced with the VEGF gene. Furthermore, the number of vessels positive for c-met also tended to increase only in the

rats introduced with the HGF gene as compared to the control. These results confirm that the HGF gene promotes lymphangiogenesis but the VEGF gene does not.

[0242] The results obtained with the rat models of lymphedema demonstrate that when a nucleic acid encoding an HGF is injected and expressed near sites affected with lymphedema, lymphatic vessels are newly generated around the injection sites. The rat models can reflect any type of lymphedema. Thus, the findings obtained herein with the rat models, that the symptom of lymphedema is relieved or cured in vivo by administering the HGF gene, suggest that HGFs and their genes are useful as therapeutic agents for lymphedema in mammals including humans.

### Example 3

#### Confirmation of HGF Signaling in Lymphatic Endothelial Cells

[0243] To confirm the above-described mechanism of action of HGFs, it was demonstrated that phosphorylation of MAPK and Akt, which is known to be induced by HGFs in vascular endothelial cells, was also induced in lymphatic endothelial cells by HGF stimulation. The phosphorylation of MAPK and Akt is known to be essential for vascular endothelial cell growth promoted by HGFs (Nakagami H., Hypertension 37 [part 2]: 581-586 (2001)).

[0244] The culture medium of the above-described lymphatic endothelial cells derived from canine thoracic duct was changed with MEM containing 0.5% FCS or FCS-free MEM 12 hours or more before addition of an HGF. A recombinant human HGF was added at a concentration of 100 ng/ml, and after 0 to 15 minutes the medium was removed and the cells were lysed with lysis buffer (50 mM Tris-Cl, 2.5 mM EGTA, 1 mM EDTA, 10 mM NaF, 1% deoxycorticosterone, 1% Triton X-100, 1 mM PMSF, and 2 mM sodium vanadate (pH 7.5)). The genome molecules were disrupted by sonication. Samples containing 20 µg proteins were subjected to 10% SDS-PAGE according to a conventional method. After transfer to nitrocellulose membrane, Western blotting was carried out using an anti-MAPK/ERK antibody, antibody specific to phosphorylated MAPK/ERK (phosphospecific; Tyr705 or Ser727), anti-Akt antibody, and antibody specific to phosphorylated Akt. The used primary antibodies were available from Cell Signaling Technology and others. Detection was achieved using the ECL kit (Amersham).

[0245] The results are shown in FIG. 11. FIG. 11A shows a result of Western blotting for MAPK. The result demonstrate that the phosphorylation of both p44 and p42 MAPKs is enhanced within five minutes after HGF stimulation. The result of Akt is shown in FIG. 11B. Likewise, the phosphorylation of Akt is enhanced within five minutes after HGF stimulation. Each bottom panel depicts detection of p44 and p42 MAPKs, or Akt as an internal control.

[0246] The phosphorylation of MAPK and Akt was also enhanced in lymphatic endothelial cells in response to HGF stimulation, as described above. This suggests that, in lymphatic endothelial cells, HGFs also induce, via c-met, the same phosphorylation cascade as in vascular endothelial cells. Thus, the phosphorylation of MAPK and Akt is presumed to be an essential signal for lymphatic endothelial cell growth.

[0247] Furthermore, the lymphatic endothelial cell growth-promoting activity of HGFs was found to be reduced by the MEK inhibitors U0126 (50 µM) and PD9805 (30 µM) and the PI3 kinase inhibitors Ly294002 (50 µM) and wortmannin (100 nM) (data not shown). This finding also suggests that the lymphangiogenic activity of HGFs is based on induction via c-met of the same signal cascade as in vascular endothelial cells by HGFs.

### INDUSTRIAL APPLICABILITY

[0248] The present invention provides novel lymphangiogenesis-promoting agents. The lymphangiogenesis-promoting agents provided by the present invention comprise human HGFs as active ingredients.

[0249] VEGF-C is a known peptidic factor that promotes lymphangiogenesis. However, other VEGF members belonging to the VEGF family have no lymphangiogenesis-promoting activity. Only VEGF-C has been known to have lymphangiogenesis-promoting activity.

[0250] Meanwhile, HGFs, which were discovered herein to have lymphangiogenesis-promoting activity, are known as angiogenic factors, like VEGF. Since VEGF has no lymphangiogenesis-promoting activity, those skilled in the art have presumed that HGFs also have no lymphangiogenesis-promoting activity. Accordingly, the findings of the present invention, that HGFs have lymphangiogenic activity, is considered as a remarkable fact.

[0251] In addition, VEGF-C may induce edema via cross-linking with VEGFR2; however, HGFs do not have such a risk. This is also a remarkable feature of the lymphangiogenesis-promoting agents of the present invention. The lymphangiogenesis-promoting agents are effective for preventing or treating lymphedema.

[0252] Furthermore, HGFs activate the growth and migration of lymphatic endothelial cells isolated not from fetal or neonatal systems but from adult animals, and thereby promote lymphangiogenesis. Most of patients suffering from lymphedema after surgical removal of cancer tissues and/or lymph nodes for cancer treatment are adult. Thus, the agents of the present invention are particularly useful as therapeutic agents for adult suffering from lymphedema after surgical removal of cancer tissues and/or lymph nodes for cancer treatment.

---

### SEQUENCE LISTING

```
<160> NUMBER OF SEQ ID NOS: 6
<210> SEQ ID NO 1
<211> LENGTH: 2187
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

-continued

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: hepatocyte growth factor (HGF)

&lt;400&gt; SEQUENCE: 1

atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc	60
ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat	120
gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa	180
acaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt	240
ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc	300
ttcaatagca tgtcaagtgg agtgaaaaaa gaatttgccc atgaatttga cctctatgaa	360
aacaagact acattagaaa ctgcatcatt ggtaaaggac gcagctacaa gggaacagta	420
tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac	480
agctttttgc cttcgagcta tcggggtaaa gacctacagg aaaactactg tcgaaatcct	540
cgaggggaag aagggggacc ctggtgtttc acaagcaatc cagaggtacg ctacgaagtc	600
tgtgacattc ctcagtgttc agaagttgaa tgcatgacct gcaatgggga gagttatcga	660
ggctctatgg atcatacaga atcaggcaag atttgtcagc gctgggatca tcagacacca	720
caccggcaca aattcttgcc tgaagatat cccgacaagg gctttgatga taattattgc	780
cgcaatcccc atggccagcc gaggccatgg tgctatactc ttgaccctca caccgctgg	840
gagtactgtg caattaaaac atgcgctgac aatactatga atgacactga tgttcctttg	900
gaaacaactg aatgcatcca aggtcaagga gaaggctaca ggggcactgt caataccatt	960
tggaatggaa ttccatgtca gcgttgggat tctcagatc ctcacgagca tgacatgact	1020
cctgaaaatt tcaagtgcga ggacctaca gaaaattact gccgaaatcc agatgggtct	1080
gaatcaccct ggtgttttac cactgatcca aacatccgag ttggctactg ctcccaaatt	1140
ccaaactgtg atatgtcaca tggacaagat tgttatcgtg ggaatggcaa aaattatatg	1200
ggcaacttat cccaacaag atctggacta acatgttcaa tgtgggacaa gaacatggaa	1260
gacttacatc gtcatatctt ctgggaacca gatgcaagta agctgaatga gaattactgc	1320
cgaaatccag atgatgatgc tcatggaccc tgggtgtaca cgggaaatcc actcattcct	1380
tgggattatt gccctatttc tcgttgtgaa ggtgatacca caccataaat agtcaattta	1440
gaccatcccc taatatcttg tgccaaaacg aaacaattgc gagttgtaaa tgggattcca	1500
acacgaacaa acataggatg gatggttagt ttgagataga gaaataaaca tatctgcgga	1560
ggatcattga taaaggagag ttgggttctt actgcacgac agtgtttccc ttctcgagac	1620
ttgaaagatt atgaagcttg gcttggaatt catgatgtcc acggaagagg agatgagaaa	1680
tgcaaacagg ttctcaatgt ttcccagctg gtatatggcc ctgaaggatc agatctggtt	1740
ttaatgaagc ttgccaggcc tgctgtcctg gatgattttg ttagtacgat tgattacct	1800
aattatggat gcacaattcc tgaagagacc agttgcagtg tttatggctg gggctacact	1860
ggattgatca actatgatgg cctattacga gtggcacatc tctatataat gggaaatgag	1920
aaatgcagcc agcatcatcg agggaaggty actctgaatg agtctgaaat atgtgctggg	1980
gctgaaaaga ttggatcagg accatgtgag ggggattatg gtggccact tgtttgtgag	2040
caacataaaa tgagaatggt tcttggtgtc attgttctcg gtcgtggatg tgccattcca	2100
aatcgtcctg gtatttttgt ccgagtagca tattatgcaa aatggataga caaaattatt	2160

-continued

ttaacatata aggtaccaca gtcatag

2187

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 728

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: hepatocyte growth factor (HGF)

&lt;400&gt; SEQUENCE: 2

Met Trp Val Thr Lys Leu Leu Pro Ala Leu Leu Leu Gln His Val Leu  
 1 5 10 15

Leu His Leu Leu Leu Leu Pro Ile Ala Ile Pro Tyr Ala Glu Gly Gln  
 20 25 30

Arg Lys Arg Arg Asn Thr Ile His Glu Phe Lys Lys Ser Ala Lys Thr  
 35 40 45

Thr Leu Ile Lys Ile Asp Pro Ala Leu Lys Ile Lys Thr Lys Lys Val  
 50 55 60

Asn Thr Ala Asp Gln Cys Ala Asn Arg Cys Thr Arg Asn Lys Gly Leu  
 65 70 75 80

Pro Phe Thr Cys Lys Ala Phe Val Phe Asp Lys Ala Arg Lys Gln Cys  
 85 90 95

Leu Trp Phe Pro Phe Asn Ser Met Ser Ser Gly Val Lys Lys Glu Phe  
 100 105 110

Gly His Glu Phe Asp Leu Tyr Glu Asn Lys Asp Tyr Ile Arg Asn Cys  
 115 120 125

Ile Ile Gly Lys Gly Arg Ser Tyr Lys Gly Thr Val Ser Ile Thr Lys  
 130 135 140

Ser Gly Ile Lys Cys Gln Pro Trp Ser Ser Met Ile Pro His Glu His  
 145 150 155 160

Ser Phe Leu Pro Ser Ser Tyr Arg Gly Lys Asp Leu Gln Glu Asn Tyr  
 165 170 175

Cys Arg Asn Pro Arg Gly Glu Glu Gly Gly Pro Trp Cys Phe Thr Ser  
 180 185 190

Asn Pro Glu Val Arg Tyr Glu Val Cys Asp Ile Pro Gln Cys Ser Glu  
 195 200 205

Val Glu Cys Met Thr Cys Asn Gly Glu Ser Tyr Arg Gly Leu Met Asp  
 210 215 220

His Thr Glu Ser Gly Lys Ile Cys Gln Arg Trp Asp His Gln Thr Pro  
 225 230 235 240

His Arg His Lys Phe Leu Pro Glu Arg Tyr Pro Asp Lys Gly Phe Asp  
 245 250 255

Asp Asn Tyr Cys Arg Asn Pro Asp Gly Gln Pro Arg Pro Trp Cys Tyr  
 260 265 270

Thr Leu Asp Pro His Thr Arg Trp Glu Tyr Cys Ala Ile Lys Thr Cys  
 275 280 285

Ala Asp Asn Thr Met Asn Asp Thr Asp Val Pro Leu Glu Thr Thr Glu  
 290 295 300

Cys Ile Gln Gly Gln Gly Glu Gly Tyr Arg Gly Thr Val Asn Thr Ile  
 305 310 315 320

Trp Asn Gly Ile Pro Cys Gln Arg Trp Asp Ser Gln Tyr Pro His Glu  
 325 330 335

-continued

---

His Asp Met Thr Pro Glu Asn Phe Lys Cys Lys Asp Leu Arg Glu Asn	340	345	350
Tyr Cys Arg Asn Pro Asp Gly Ser Glu Ser Pro Trp Cys Phe Thr Thr	355	360	365
Asp Pro Asn Ile Arg Val Gly Tyr Cys Ser Gln Ile Pro Asn Cys Asp	370	375	380
Met Ser His Gly Gln Asp Cys Tyr Arg Gly Asn Gly Lys Asn Tyr Met	385	390	395
Gly Asn Leu Ser Gln Thr Arg Ser Gly Leu Thr Cys Ser Met Trp Asp	405	410	415
Lys Asn Met Glu Asp Leu His Arg His Ile Phe Trp Glu Pro Asp Ala	420	425	430
Ser Lys Leu Asn Glu Asn Tyr Cys Arg Asn Pro Asp Asp Ala His	435	440	445
Gly Pro Trp Cys Tyr Thr Gly Asn Pro Leu Ile Pro Trp Asp Tyr Cys	450	455	460
Pro Ile Ser Arg Cys Glu Gly Asp Thr Thr Pro Thr Ile Val Asn Leu	465	470	475
Asp His Pro Val Ile Ser Cys Ala Lys Thr Lys Gln Leu Arg Val Val	485	490	495
Asn Gly Ile Pro Thr Arg Thr Asn Ile Gly Trp Met Val Ser Leu Arg	500	505	510
Tyr Arg Asn Lys His Ile Cys Gly Gly Ser Leu Ile Lys Glu Ser Trp	515	520	525
Val Leu Thr Ala Arg Gln Cys Phe Pro Ser Arg Asp Leu Lys Asp Tyr	530	535	540
Glu Ala Trp Leu Gly Ile His Asp Val His Gly Arg Gly Asp Glu Lys	545	550	555
Cys Lys Gln Val Leu Asn Val Ser Gln Leu Val Tyr Gly Pro Glu Gly	565	570	575
Ser Asp Leu Val Leu Met Lys Leu Ala Arg Pro Ala Val Leu Asp Asp	580	585	590
Phe Val Ser Thr Ile Asp Leu Pro Asn Tyr Gly Cys Thr Ile Pro Glu	595	600	605
Lys Thr Ser Cys Ser Val Tyr Gly Trp Gly Tyr Thr Gly Leu Ile Asn	610	615	620
Tyr Asp Gly Leu Leu Arg Val Ala His Leu Tyr Ile Met Gly Asn Glu	625	630	635
Lys Cys Ser Gln His Arg Gly Lys Val Thr Leu Asn Glu Ser Glu	645	650	655
Ile Cys Ala Gly Ala Glu Lys Ile Gly Ser Gly Pro Cys Glu Gly Asp	660	665	670
Tyr Gly Gly Pro Leu Val Cys Glu Gln His Lys Met Arg Met Val Leu	675	680	685
Gly Val Ile Val Pro Gly Arg Gly Cys Ala Ile Pro Asn Arg Pro Gly	690	695	700
Ile Phe Val Arg Val Ala Tyr Tyr Ala Lys Trp Ile His Lys Ile Ile	705	710	715
Leu Thr Tyr Lys Val Pro Gln Ser	725		

-continued

---

```

<210> SEQ ID NO 3
<211> LENGTH: 4173
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: hepatocyte growth factor (HGF) receptor

<400> SEQUENCE: 3

atgaaggccc ccgctgtgct tgcacctggc atcctcgtgc tcctgtttac cttggtgcag    60
aggagcaatg gggagtgtaa agaggcacta gcaaagtcg agatgaatgt gaatatgaag    120
tatcagcttc ccaacttcac cgcggaaca cccatccaga atgtcattct acatgagcat    180
cacattttcc ttggtgccac taactacatt tatgttttaa atgaggaaga ccttcagaag    240
gttgctgagt acaagactgg gcctgtgctg gaacaccagc attgtttccc atgtcaggac    300
tgcagcagca aagccaattt atcaggaggt gtttgaaaag ataacatcaa catggctcta    360
gttgctgaca cctactatga tgatcaactc attagctgtg gcagcgtcaa cagagggacc    420
tgccagcgac atgtctttcc ccacaatcat actgctgaca tacagtcgga ggttcactgc    480
atattctccc cacagataga agagcccagc cagtgtcctg actgtgtggt gagcgccctg    540
ggagccaaa gtcctttcatc tgtaaaggac cggttcatca acttctttgt aggcaatacc    600
ataaattctt cttattttcc agatcatcca ttgcattcga tatcagttag aaggctaaag    660
gaaacgaaag atggttttat gtttttgacg gaccagtcct acattgatgt tttacctgag    720
ttcagagatt cttaccccat taagtatgtc catgcctttg aaagcaacaa ttttatttac    780
ttcttgacgg tccaaaggga aactctagat gctcagactt ttcacacaag aataatcagg    840
ttctgttcca taaactctgg attgcattcc tacatggaaa tgcctctgga gtgtattctc    900
acagaaaaga gaaaaaagag atccacaaa gaggaaagt ttaataactc tcaggctgag    960
tatgtcagca agcctggggc ccagcttgct agacaaatag gagccagcct gaatgatgac   1020
attcttttgc ggggtgttcg acaaagcaag ccagattctg ccgaaccaat ggatcgatct   1080
gccatgtgtg cattccctat caaatatgtc aacgaacttc tcaacaagat cgtcaacaaa   1140
aacaatgtga gatgtctcca gcatttttac ggaccaatc atgagcactg ctttaatagg   1200
acacttctga gaaattcatc aggtctgtga gcgcgcgctg atgaatatcg aacagagttt   1260
accacagctt tgcagcgcgt tgacttatcc atgggtcaat tcagcgaagt cctcttaaca   1320
tctatatcca ccttcattaa aggagaccct accatagcta atcttgggac atcagagggg   1380
cgcttcatgc aggttgtggt ttctcgatca ggaccatcaa cccctcatgt gaattttctc   1440
ctggactccc atccagtgtc tccagaagtg attgtggagc atacattaaa ccaaaatggc   1500
tacacactgg ttatcactgg gaagaagatc acgaagatcc cattgaatgg cttgggctgc   1560
agacatttcc agtcctgcag tcaatgcctc tctgccccac ctttgttca gtgtggctgg   1620
tgccacgaca aatgtgtgag atcggaggaa tgctgagcgc ggacatggac tcaacagatc   1680
tgtctgcctg caatctacaa ggttttccca aatagtgcac ccctgaagg agggacaagg   1740
ctgaccatat gtggtggtga ctttgattt cgagggaata ataaatttga tttaaagaaa   1800
actagagttc tccttggaag tgagagctgc accttgactt taagttagag cacgatgaat   1860
acattgaaat gcacagtgtg tcctgccatg aataagcatt tcaatatgtc cataattatt   1920
tcaaattggc acgggacaac acaatacagt acattctcct atgtggatcc tgtaataaca   1980
agtatttctc cgaaatacgg tcctatggct ggtggcactt tacttacttt aactggaaat   2040

```

-continued

---

tacctaaca	gtgggaattc	tagacacatt	tcaattggtg	gaaaaacatg	tactttaaaa	2100
agtggtgcaa	acagtattct	tgaatgttat	acccagccc	aaaccatttc	aactgagttt	2160
gctgttaaat	tgaaaattga	cttagccaac	cgagagacaa	gcattctcag	ttaccgtgaa	2220
gatcccatg	tctatgaaat	tcattccaac	aaatctttta	ttagtggtgg	gagcacaata	2280
acaggtgtg	ggaaaaacct	gaattcagtt	agtgccccga	gaatggcat	aaatgtgcat	2340
gaagcaggaa	ggaactttac	agtggcatgt	caacatcgct	ctaattcaga	gataatctgt	2400
tgtaccactc	cttcctgca	acagctgaat	ctgcaactcc	cctgaaaac	caaagccttt	2460
ttcatgttag	atgggacctc	ttccaaatac	tttgatctca	tttatgtaca	taatcctgtg	2520
tttaagcctt	ttgaaaagcc	agtgatgac	tcaatgggca	atgaaaatgt	actggaaatt	2580
aagggaatg	atattgacct	tgaagcagtt	aaaggtgaag	tgtaaaaagt	tggaaataag	2640
agctgtgaga	atatacactt	acattctgaa	gccgttttat	gcacggccc	caatgacctg	2700
ctgaaattga	acagcgagct	aaatatagag	tggaagcaag	caattctctc	aaccgtcctt	2760
ggaaaagtaa	tagttcaacc	agatcagaat	ttcacaggat	tgattgctgg	tgttgtctca	2820
atatcaacag	cactgttatt	actactggg	ttttctctgt	ggctgaaaa	gagaaagcaa	2880
attaaagatc	tgggcagtg	attagtctgc	tacgatgcaa	gagtacacac	tcctcatttg	2940
gataggcttg	taagtgccg	aagtgtaac	ccaactacag	aaatggtttc	aatgaatct	3000
gtagactacc	gagctacttt	tccagaagat	cagtttctca	attcatctca	gaacggttca	3060
tgccgacaag	tgcatgatcc	tctgacagac	atgtccccc	tcctaactag	tggggactct	3120
gatatatcca	gtccattact	gcaaaatact	gtccacattg	acctcagtgc	tctaaatcca	3180
gagctggtcc	aggcagtgca	gcattgtatg	attgggcccc	gtagcctgat	tgtgcatttc	3240
aatgaagtca	taggaagagg	gcattttggt	tgtgtatata	atgggacttt	gttggaacaat	3300
gatggcaaga	aaattcactg	tgtgtgaaa	tccttgaaac	gaatcactga	cataggagaa	3360
gtttcccaat	ttctgaccga	gggaatcatc	atgaaagatt	ttagtcatcc	caatgtcctc	3420
tcgtctctgg	gaatctgctc	gcgaagtga	gggtctccgc	tgggtgctct	accatacatg	3480
aaacatggag	atcttcgaaa	tttcattcga	aatgagactc	ataatccaac	tgtaaaagat	3540
cttattggct	ttggtcttca	agtagccaaa	ggcatgaaat	atcttgcaag	caaaaagttt	3600
gtccacagag	acttggtctc	aagaaactgt	atgctggatg	aaaaattcac	agtcaagggt	3660
gctgattttg	gtcttgccag	agacatgtat	gataaagaat	actatagtgt	acacaacaaa	3720
acaggtgcaa	agctgccagt	gaagtggatg	gctttgaaa	gtctgcaaac	tcaaaaagttt	3780
accaccaagt	cagatgtgtg	gtcctttggc	gtgctcctct	gggagctgat	gacaagagga	3840
gccccacctt	atcctgacgt	aaacaccttt	gatataactg	tttacttggt	gcaagggaga	3900
agactcctac	aacccgaata	ctgccagac	cccttatatg	aagtaatgct	aaaatgctgg	3960
caccctaaag	cgaatgctg	cccatccttt	tctgaactgg	tgtcccgat	atcagcgatc	4020
ttctctactt	tcattgggga	gcactatgtc	catgtgaacg	ctacttatgt	gaacgtaaaa	4080
tgtgtcgtc	cgtatccttc	tctgtgtgca	tcagaagata	acgctgatga	tgaggtggac	4140
acacgaccag	cctccttctg	ggagacatca	tag			4173

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 1390



-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (0)...(0)
<223> OTHER INFORMATION: hepatocyte growth factor (HGF) receptor

<400> SEQUENCE: 4

Met Lys Ala Pro Ala Val Leu Ala Pro Gly Ile Leu Val Leu Leu Phe
 1             5             10             15

Thr Leu Val Gln Arg Ser Asn Gly Glu Cys Lys Glu Ala Leu Ala Lys
 20             25             30

Ser Glu Met Asn Val Asn Met Lys Tyr Gln Leu Pro Asn Phe Thr Ala
 35             40             45

Glu Thr Pro Ile Gln Asn Val Ile Leu His Glu His His Ile Phe Leu
 50             55             60

Gly Ala Thr Asn Tyr Ile Tyr Val Leu Asn Glu Glu Asp Leu Gln Lys
 65             70             75             80

Val Ala Glu Tyr Lys Thr Gly Pro Val Leu Glu His Pro Asp Cys Phe
 85             90             95

Pro Cys Gln Asp Cys Ser Ser Lys Ala Asn Leu Ser Gly Gly Val Trp
 100            105            110

Lys Asp Asn Ile Asn Met Ala Leu Val Val Asp Thr Tyr Tyr Asp Asp
 115            120            125

Gln Leu Ile Ser Cys Gly Ser Val Asn Arg Gly Thr Cys Gln Arg His
 130            135            140

Val Phe Pro His Asn His Thr Ala Asp Ile Gln Ser Glu Val His Cys
 145            150            155            160

Ile Phe Ser Pro Gln Ile Glu Glu Pro Ser Gln Cys Pro Asp Cys Val
 165            170            175

Val Ser Ala Leu Gly Ala Lys Val Leu Ser Ser Val Lys Asp Arg Phe
 180            185            190

Ile Asn Phe Phe Val Gly Asn Thr Ile Asn Ser Ser Tyr Phe Pro Asp
 195            200            205

His Pro Leu His Ser Ile Ser Val Arg Arg Leu Lys Glu Thr Lys Asp
 210            215            220

Gly Phe Met Phe Leu Thr Asp Gln Ser Tyr Ile Asp Val Leu Pro Glu
 225            230            235            240

Phe Arg Asp Ser Tyr Pro Ile Lys Tyr Val His Ala Phe Glu Ser Asn
 245            250            255

Asn Phe Ile Tyr Phe Leu Thr Val Gln Arg Glu Thr Leu Asp Ala Gln
 260            265            270

Thr Phe His Thr Arg Ile Ile Arg Phe Cys Ser Ile Asn Ser Gly Leu
 275            280            285

His Ser Tyr Met Glu Met Pro Leu Glu Cys Ile Leu Thr Glu Lys Arg
 290            295            300

Lys Lys Arg Ser Thr Lys Lys Glu Val Phe Asn Ile Leu Gln Ala Ala
 305            310            315            320

Tyr Val Ser Lys Pro Gly Ala Gln Leu Ala Arg Gln Ile Gly Ala Ser
 325            330            335

Leu Asn Asp Asp Ile Leu Phe Gly Val Phe Ala Gln Ser Lys Pro Asp
 340            345            350

Ser Ala Glu Pro Met Asp Arg Ser Ala Met Cys Ala Phe Pro Ile Lys

```

-continued

355					360					365								
Tyr	Val	Asn	Asp	Phe	Phe	Asn	Lys	Ile	Val	Asn	Lys	Asn	Asn	Val	Arg			
370					375					380								
Cys	Leu	Gln	His	Phe	Tyr	Gly	Pro	Asn	His	Glu	His	Cys	Phe	Asn	Arg			
385					390					395					400			
Thr	Leu	Leu	Arg	Asn	Ser	Ser	Gly	Cys	Glu	Ala	Arg	Arg	Asp	Glu	Tyr			
					405					410					415			
Arg	Thr	Glu	Phe	Thr	Thr	Ala	Leu	Gln	Arg	Val	Asp	Leu	Phe	Met	Gly			
					420					425					430			
Gln	Phe	Ser	Glu	Val	Leu	Leu	Thr	Ser	Ile	Ser	Thr	Phe	Ile	Lys	Gly			
					435					440					445			
Asp	Leu	Thr	Ile	Ala	Asn	Leu	Gly	Thr	Ser	Glu	Gly	Arg	Phe	Met	Gln			
					450					455					460			
Val	Val	Val	Ser	Arg	Ser	Gly	Pro	Ser	Thr	Pro	His	Val	Asn	Phe	Leu			
					465					470					475		480	
Leu	Asp	Ser	His	Pro	Val	Ser	Pro	Glu	Val	Ile	Val	Glu	His	Thr	Leu			
					485					490					495			
Asn	Gln	Asn	Gly	Tyr	Thr	Leu	Val	Ile	Thr	Gly	Lys	Lys	Ile	Thr	Lys			
					500					505					510			
Ile	Pro	Leu	Asn	Gly	Leu	Gly	Cys	Arg	His	Phe	Gln	Ser	Cys	Ser	Gln			
					515					520					525			
Cys	Leu	Ser	Ala	Pro	Pro	Phe	Val	Gln	Cys	Gly	Trp	Cys	His	Asp	Lys			
					530					535					540			
Cys	Val	Arg	Ser	Glu	Glu	Cys	Leu	Ser	Gly	Thr	Trp	Thr	Gln	Gln	Ile			
					545					550					555		560	
Cys	Leu	Pro	Ala	Ile	Tyr	Lys	Val	Phe	Pro	Asn	Ser	Ala	Pro	Leu	Glu			
					565					570					575			
Gly	Gly	Thr	Arg	Leu	Thr	Ile	Cys	Gly	Trp	Asp	Phe	Gly	Phe	Arg	Arg			
					580					585					590			
Asn	Asn	Lys	Phe	Asp	Leu	Lys	Lys	Thr	Arg	Val	Leu	Leu	Gly	Asn	Glu			
					595					600					605			
Ser	Cys	Thr	Leu	Thr	Leu	Ser	Glu	Ser	Thr	Met	Asn	Thr	Leu	Lys	Cys			
					610					615					620			
Thr	Val	Gly	Pro	Ala	Met	Asn	Lys	His	Phe	Asn	Met	Ser	Ile	Ile	Ile			
					625					630					635		640	
Ser	Asn	Gly	His	Gly	Thr	Thr	Gln	Tyr	Ser	Thr	Phe	Ser	Tyr	Val	Asp			
					645					650					655			
Pro	Val	Ile	Thr	Ser	Ile	Ser	Pro	Lys	Tyr	Gly	Pro	Met	Ala	Gly	Gly			
					660					665					670			
Thr	Leu	Leu	Thr	Leu	Thr	Gly	Asn	Tyr	Leu	Asn	Ser	Gly	Asn	Ser	Arg			
					675					680					685			
His	Ile	Ser	Ile	Gly	Gly	Lys	Thr	Cys	Thr	Leu	Lys	Ser	Val	Ser	Asn			
					690					695					700			
Ser	Ile	Leu	Glu	Cys	Tyr	Thr	Pro	Ala	Gln	Thr	Ile	Ser	Thr	Glu	Phe			
					705					710					715		720	
Ala	Val	Lys	Leu	Lys	Ile	Asp	Leu	Ala	Asn	Arg	Glu	Thr	Ser	Ile	Phe			
					725					730					735			
Ser	Tyr	Arg	Glu	Asp	Pro	Ile	Val	Tyr	Glu	Ile	His	Pro	Thr	Lys	Ser			
					740					745					750			
Phe	Ile	Ser	Gly	Gly	Ser	Thr	Ile	Thr	Gly	Val	Gly	Lys	Asn	Leu	Asn			
					755					760					765			

-continued

---

Ser	Val	Ser	Val	Pro	Arg	Met	Val	Ile	Asn	Val	His	Glu	Ala	Gly	Arg	770	775	780
Asn	Phe	Thr	Val	Ala	Cys	Gln	His	Arg	Ser	Asn	Ser	Glu	Ile	Ile	Cys	785	790	795
Cys	Thr	Thr	Pro	Ser	Leu	Gln	Gln	Leu	Asn	Leu	Gln	Leu	Pro	Leu	Lys	805	810	815
Thr	Lys	Ala	Phe	Phe	Met	Leu	Asp	Gly	Ile	Leu	Ser	Lys	Tyr	Phe	Asp	820	825	830
Leu	Ile	Tyr	Val	His	Asn	Pro	Val	Phe	Lys	Pro	Phe	Glu	Lys	Pro	Val	835	840	845
Met	Ile	Ser	Met	Gly	Asn	Glu	Asn	Val	Leu	Glu	Ile	Lys	Gly	Asn	Asp	850	855	860
Ile	Asp	Pro	Glu	Ala	Val	Lys	Gly	Glu	Val	Leu	Lys	Val	Gly	Asn	Lys	865	870	875
Ser	Cys	Glu	Asn	Ile	His	Leu	His	Ser	Glu	Ala	Val	Leu	Cys	Thr	Val	885	890	895
Pro	Asn	Asp	Leu	Leu	Lys	Leu	Asn	Ser	Glu	Leu	Asn	Ile	Glu	Trp	Lys	900	905	910
Gln	Ala	Ile	Ser	Ser	Thr	Val	Leu	Gly	Lys	Val	Ile	Val	Gln	Pro	Asp	915	920	925
Gln	Asn	Phe	Thr	Gly	Leu	Ile	Ala	Gly	Val	Val	Ser	Ile	Ser	Thr	Ala	930	935	940
Leu	Leu	Leu	Leu	Leu	Gly	Phe	Phe	Leu	Trp	Leu	Lys	Lys	Arg	Lys	Gln	945	950	955
Ile	Lys	Asp	Leu	Gly	Ser	Glu	Leu	Val	Arg	Tyr	Asp	Ala	Arg	Val	His	965	970	975
Thr	Pro	His	Leu	Asp	Arg	Leu	Val	Ser	Ala	Arg	Ser	Val	Ser	Pro	Thr	980	985	990
Thr	Glu	Met	Val	Ser	Asn	Glu	Ser	Val	Asp	Tyr	Arg	Ala	Thr	Phe	Pro	995	1000	1005
Glu	Asp	Gln	Phe	Pro	Asn	Ser	Ser	Gln	Asn	Gly	Ser	Cys	Arg	Gln	Val	1010	1015	1020
Gln	Tyr	Pro	Leu	Thr	Asp	Met	Ser	Pro	Ile	Leu	Thr	Ser	Gly	Asp	Ser	1025	1030	1035
Asp	Ile	Ser	Ser	Pro	Leu	Leu	Gln	Asn	Thr	Val	His	Ile	Asp	Leu	Ser	1045	1050	1055
Ala	Leu	Asn	Pro	Glu	Leu	Val	Gln	Ala	Val	Gln	His	Val	Val	Ile	Gly	1060	1065	1070
Pro	Ser	Ser	Leu	Ile	Val	His	Phe	Asn	Glu	Val	Ile	Gly	Arg	Gly	His	1075	1080	1085
Phe	Gly	Cys	Val	Tyr	His	Gly	Thr	Leu	Leu	Asp	Asn	Asp	Gly	Lys	Lys	1090	1095	1100
Ile	His	Cys	Ala	Val	Lys	Ser	Leu	Asn	Arg	Ile	Thr	Asp	Ile	Gly	Glu	1105	1110	1115
Val	Ser	Gln	Phe	Leu	Thr	Glu	Gly	Ile	Ile	Met	Lys	Asp	Phe	Ser	His	1125	1130	1135
Pro	Asn	Val	Leu	Ser	Leu	Leu	Gly	Ile	Cys	Leu	Arg	Ser	Glu	Gly	Ser	1140	1145	1150
Pro	Leu	Val	Val	Leu	Pro	Tyr	Met	Lys	His	Gly	Asp	Leu	Arg	Asn	Phe	1155	1160	1165

-continued

Ile	Arg	Asn	Glu	Thr	His	Asn	Pro	Thr	Val	Lys	Asp	Leu	Ile	Gly	Phe
1170						1175						1180			
Gly	Leu	Gln	Val	Ala	Lys	Gly	Met	Lys	Tyr	Leu	Ala	Ser	Lys	Lys	Phe
1185					1190					1195					1200
Val	His	Arg	Asp	Leu	Ala	Ala	Arg	Asn	Cys	Met	Leu	Asp	Glu	Lys	Phe
				1205					1210					1215	
Thr	Val	Lys	Val	Ala	Asp	Phe	Gly	Leu	Ala	Arg	Asp	Met	Tyr	Asp	Lys
				1220					1225					1230	
Glu	Tyr	Tyr	Ser	Val	His	Asn	Lys	Thr	Gly	Ala	Lys	Leu	Pro	Val	Lys
				1235				1240					1245		
Trp	Met	Ala	Leu	Glu	Ser	Leu	Gln	Thr	Gln	Lys	Phe	Thr	Thr	Lys	Ser
	1250					1255					1260				
Asp	Val	Trp	Ser	Phe	Gly	Val	Leu	Leu	Trp	Glu	Leu	Met	Thr	Arg	Gly
1265					1270					1275					1280
Ala	Pro	Pro	Tyr	Pro	Asp	Val	Asn	Thr	Phe	Asp	Ile	Thr	Val	Tyr	Leu
				1285					1290						1295
Leu	Gln	Gly	Arg	Arg	Leu	Leu	Gln	Pro	Glu	Tyr	Cys	Pro	Asp	Pro	Leu
				1300					1305					1310	
Tyr	Glu	Val	Met	Leu	Lys	Cys	Trp	His	Pro	Lys	Ala	Glu	Met	Arg	Pro
				1315				1320					1325		
Ser	Phe	Ser	Glu	Leu	Val	Ser	Arg	Ile	Ser	Ala	Ile	Phe	Ser	Thr	Phe
	1330					1335						1340			
Ile	Gly	Glu	His	Tyr	Val	His	Val	Asn	Ala	Thr	Tyr	Val	Asn	Val	Lys
1345					1350					1355					1360
Cys	Val	Ala	Pro	Tyr	Pro	Ser	Leu	Leu	Ser	Ser	Glu	Asp	Asn	Ala	Asp
				1365					1370					1375	
Asp	Glu	Val	Asp	Thr	Arg	Pro	Ala	Ser	Phe	Trp	Glu	Thr	Ser		
				1380					1385				1390		

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 5181

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: mammalian expression plasmid vector  
pVAX1HGF/MGB1 including human HGF cDNA packaged in HVJ-E vector

&lt;400&gt; SEQUENCE: 5

```

gctgcttcgc gatgtacggg ccagatatac gcgttgacat tgattattga ctagttatta      60
atagtaatca attacggggg cattagttca tagcccatat atggagttcc gcgttacata      120
acttacggta aatggccgcg ctggctgacc gcccaacgac ccccgcccat tgacgtcaat      180
aatgacgtat gttcccatag taacgccaat agggactttc cattgacgtc aatgggtgga      240
gtatttacgg taaactgccc acttggcagt acatcaagtg tatcatatgc caagtacgcc      300
ccctattgac gtcaatgacg gtaaatggcc cgcttgccat tatgccagt acatgacctt      360
atgggacttt cctacttggc agtacatcta cgtattagtc atcgctatta ccatggtgat      420
gcggttttgg cagtacatca atgggcgtgg atagcggttt gactcacggg gatttccaag      480
tctccacccc attgacgtca atgggagttt gttttggcac caaatcaac gggactttcc      540
aaaatgtcgt aacaactccg ccccatggac gcaaatgggc ggtaggcgtg tacggtggga      600
gggttatata agcagagctc tctggctaac tagagaaccc actgcttact ggcttatcga      660
aattaatacg actcactata gggagacca agctggctag cgtttaaact taagcttggt      720

```

-continued

---

accgagctcg gatccgccag cccgtccagc agcaccatgt gggtgaccaa actcctgcc	780
gccctgtctg tgcagcatgt cctcctgcat ctctcctgc tccccatgc catccctat	840
gcagagggac aaaggaag aagaaataca attcatgaat tcaaaaaatc agcaaagact	900
accctaataca aaatagatcc agcactgaag ataaaaacca aaaaagtga tactgcagac	960
caatgtgcta atagatgtac taggaataaa ggacttccat tcaattgcaa ggcttttgtt	1020
tttgataaag caagaaaaca atgcctctgg ttccccttca atagcatgtc aagtggagt	1080
aaaaaagaat ttggccatga atttgacctc tatgaaaaca aagactacat tagaaactgc	1140
atcattggta aaggacgcag ctacaaggga acagtatcta tcaactaagag tggcatcaaa	1200
tgtagccctt ggagttccat gataccacac gaacacagct ttttgccttc gagctatcgg	1260
ggtaagacc tacaggaaaa ctactgtcga aatcctcgag gggaagaagg gggaccctgg	1320
tggttcacaa gcaatccaga ggtacgctac gaagtctgtg acattcctca gtgttcagaa	1380
gttgaatgca tgacctgcaa tggggagagt tatcgaggtc tcatggatca tacagaatca	1440
ggcaagattt gtcagcgtg ggatcatcag acaccacacc ggcacaaatt cttgcctgaa	1500
agatatcccc acaagggctt tgatgataat tattgccgca atcccgatgg ccagccgagg	1560
ccatgggtgct atactcttga ccttcacacc cgctgggagt actgtgcaat taaaacatgc	1620
gctgacaata ctatgaatga cactgatgtt cctttggaaa caactgaatg catccaaggt	1680
caaggagaag gctacagggg cactgtcaat accatttga atggaattcc atgtcagcgt	1740
tgggattctc agtatcctca cgagcatgac atgactcctg aaaatttcaa gtgcaaggac	1800
ctacgagaaa attactgccg aaatccagat gggcttgaat caccctggtg ttttaccact	1860
gatccaaaca tccgagtgtg ctactgctcc caaattccaa actgtgatat gtcacatgga	1920
caagattggt atcgtgggaa tggcaaaaat tatatgggca acttatccca aacaagatct	1980
ggactaacat gttcaatgtg ggacaagaac atggaagact tacatcgta tatcttctgg	2040
gaaccagatg caagtaagct gaatgagaat tactgccgaa atccagatga tgatgctcat	2100
ggaccctggt gctacacggg aaatccactc attccttggg attattgcc tatttctcgt	2160
tgtgaagggt ataccacacc tacaatagtc aatttagacc atcccgtaat atcttgtgcc	2220
aaaacgaac aattgcgagt tgtaaatggg attccaacac gaacaaacat aggatggatg	2280
gttagtttga gatacagaaa taacatatac tgcggaggat cattgataaa ggagagttgg	2340
gttcttactg cagcagctg ttcccttct cgagacttga aagattatga agcttggctt	2400
ggaattcatg atgtccacgg aagaggagat gagaaatgca aacaggttct caatgtttcc	2460
cagctgggtat atggccctga aggatcagat ctggttttaa tgaagcttgc caggcctgct	2520
gtcctggatg attttgttag tacgattgat ttacctaatt atggatgcac aattcctgaa	2580
aagaccagtt gcagtgttta tggctggggc tactctggat tgatcaacta tgatggccta	2640
ttacgagtgg cacatctcta tataatggga aatgagaaat gcagccagca tcatcgaggg	2700
aagtgactc tgaatgagtc tgaatatgt gctggggctg aaaagattgg atcaggacca	2760
tgtgaggggg attatggtg cccacttgtt tgtgagcaac ataaaatgag aatggttctt	2820
gggtgcattg ttcttggtcg tggatgtgcc attccaaatc gtcttggtat ttttgtccga	2880
gtagcatatt atgcaaaatg gatacacaaa attattttaa catataaggt accacagtca	2940
tagctgttaa cccgggtcga agcggccgct cgagtctaga gggcccggtt aaaccgcgtg	3000

-continued

---

atcagcctcg actgtgcctt ctagtgtcca gccatctgtt gtttgcccct cccccgtgcc	3060
ttccttgacc ctggaagggtg ccactcccac tgtcctttcc taataaaatg aggaaattgc	3120
atcgcatgtg ctgagtaggt gtcattctat tctggggggt ggggtggggc aggacagcaa	3180
gggggaggtat tgggaagaca atagcaggca tgctggggat gcggtgggct ctatggcttc	3240
tactgggcgg ttttatggac agcaagcgaa ccggaattgc cagctggggc gccctctggt	3300
aaggttggga agccctgcaa agtaaaactgg atggccttct tgccccaag gatctgatgg	3360
cgcaggggat caagctctga tcaagagaca ggatgaggat cgtttcgcat gattgaacaa	3420
gatggattgc acgcagggtc tccggccgct tgggtggaga ggctattcgg ctatgactgg	3480
gcacaacaga caatcggtcg ctctgatgcc gccgtgttcc ggctgtcagc gcaggggccc	3540
ccggttcttt ttgtcaagac cgacctgtcc ggtgccctga atgaactgca agacgaggca	3600
gcgcggctat cgtggctggc cagcaggggc gttccttgcg cagctgtgct cgacgttgtc	3660
actgaagcgg gaagggactg gctgctattg ggcaagtgc cggggcagga tctcctgtca	3720
tctcaccttg ctctgcgga gaaagtatcc atcatggctg atgcaatgcg gcggctgcat	3780
acgcttgatc cggctacctg cccattcgac caccaagcga aacatcgcat cgagcgagca	3840
cgtactcgga tggaaagccgg tcttgtcgat caggatgac tggaacgaaga gcatcagggg	3900
ctcgcgccag ccgaactggt cgccaggctc aaggcgagca tgcccgacgg cgaggatctc	3960
gtcgtgaccc atggcgatgc ctgcttgcg aatatcatgg tggaaaatgg ccgcttttct	4020
ggattcatcg actgtggcgg gctgggtgtg gcggaccgct atcaggacat agcgttggtc	4080
accgctgata ttgctgaaga gcttggcggc gaatgggctg accgcttctc cgtgctttac	4140
gggtatcgccg ctcccgatc gcagcgcac gccttctatc gccttcttga cgagttcttc	4200
tgaattatta acgcttacaa tttcctgatg cggatatttc tcttacgca tctgtgcggg	4260
atttcacacc gcatcagggt gcaacttttc gggaaatgtg cgcggaaccc ctatttgttt	4320
atTTTTctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct	4380
tcaataatag cacgtgctaa aacttcattt ttaatttaaa aggatctagg tgaagatcct	4440
ttttgataat ctcatgacca aaatccctta acgtgagttt tcgttccact gagcgtcaga	4500
ccccgtagaa aagatcaaag gatcttcttg agatcctttt tttctgcgcg taatctgctg	4560
cttgcaacaa aaaaaaccac cgctaccagc ggtggtttgt ttgccggatc aagagctacc	4620
aactcttttt ccgaaggtaa ctggcttcag cagagcgagc ataccaaata ctgttcttct	4680
agtgtagccg tagttaggcc accacttcaa gaactctgta gcaccgcta catacctcgc	4740
tctgctaate ctgttaccag tggetgtgc cagtggcgat aagtcgtgtc ttaccgggtt	4800
ggactcaaga cgatagttag cgataaggc gcagcggtcg ggctgaacgg ggggttcgtg	4860
cacacagccc agcttgagc gaacgacctc caccgaactg agatacctac agcgtgagct	4920
atgagaaagc gccacgcttc ccgaaggagg aaaggcggac aggtatccgg taagcggcag	4980
ggtcggaaca ggagagcgca cgaggagct tccaggggga aacgcctggt atctttatag	5040
tctgtcggg tttccgacc tctgacttga gcgtcgattt ttgtgatgct cgtcaggggg	5100
gcggagccta tggaaaaacg ccagcaacgc ggccttttta cggttcctgg ccttttctg	5160
gccttttctg cacatgttct t	5181

-continued

---

```

<210> SEQ ID NO 6
<211> LENGTH: 7681
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: mammalian expression plasmid vector
      pcDNA3.1(-)HGF

<400> SEQUENCE: 6
gacggatcgg gagatctccc gatccctat ggtgcactct cagtacaatc tgcctctgatg      60
ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggtcgct gagtagtgcg      120
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc      180
ttaggggttag gcgtttttcg ctgcttcgcg atgtacgggc cagatatacg cgttgacatt      240
gattattgac tagttattaa tagtaatcaa ttacgggggc attagttcat agcccatata      300
tggagttccg cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc      360
cccgccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc      420
attgacgtca atgggtggag tatttacggg aaactgcccc cttggcagta catcaagtgt      480
atcatatgcc aagtagcgcc cctattgacg tcaatgacgg taaatggccc gcctggcatt      540
atgcccagta catgacctta tgggactttc ctacttgcca gtacatctac gtattagtca      600
tcgtatttac catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggtttg      660
actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc      720
aaaatcaacg ggactttcca aaatgtcgta acaactccgc ccattgacg caaatgggcg      780
gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact agagaaccca      840
ctgcttactg gcttatcgaa attaatacga ctactatag ggagacccaa gctggctagc      900
gtttaaacgg gccctctaga ctcgagcggc cgctctagaa ctagctggat cctgccagcc      960
cgteccagcag caccatgtgg gtgaccaaac tcctgccagc cctgctgctg cagcatgtcc      1020
tcctgcactc cctctgctc cccatcgcca tcccctatgc agagggacaa aggaaaagaa      1080
gaaatacaat tcataaattc aaaaaatcag caaagactac ctaaatcaaa atagatccag      1140
cactgaagat aaaaacccaa aaagtgaata ctgcagacca atgtgctaata agatgtacta      1200
ggaataaagg acttccattc acttgcaagg cttttgtttt tgataaagca agaaaacaat      1260
gcctctgggt ccccttcaat agcatgtcaa gtggagtga aaaaagaattt ggccatgaat      1320
ttgacctcta tgaaaacaaa gactacatta gaaactgcat cattggtaaa ggacgcagct      1380
acaagggaac agtatctatc actaagagtg gcatcaaag tcagccctgg agttccatga      1440
taccacacga acacagcttt ttgccttcga gctatcgggg taaagaccta caggaaaact      1500
actgtcgaaa tcctcgaggg gaagaagggg gacctggtg tttcacaagc aatccagagg      1560
tacgctacga agtctgtgac attcctcagt gttcagaagt tgaatgcatg acctgcaatg      1620
gggagagtta tcgaggtctc atggatcata cagaatcagg caagatttgt cagcgctggg      1680
atcatcagac accacaccgg cacaaattct tgcctgaaag atatcccgac aagggttttg      1740
atgataatta ttgccgcaat cccgatggcc agccgaggcc atggtgctat actcttgacc      1800
ctcacacccg ctgggagtac tgtgcaatta aaacatgcgc tgacaatact atgaatgaca      1860
ctgatgttcc ttggaaaca actgaatgca tccaaggtca aggagaaggc tacaggggca      1920
ctgtcaatac catttggaat ggaattccat gtcagcgttg ggattctcag tatectcacg      1980

```

-continued

agcatgacat gactcctgaa aatttcaagt gcaaggacct acgagaaaaat tactgccgaa	2040
atccagatgg gtctgaatca ccttggtgtt ttaccactga tccaaacatc cgagttggct	2100
actgctcccc aattccaaac tgtgatatgt cacatggaca agattgttat cgtgggaatg	2160
gcaaaaatta tatgggcaac ttatcccaaa caagatctgg actaacatgt tcaatgtggg	2220
acaagaacat ggaagactta catcgtcata tcttctggga accagatgca agtaagctga	2280
atgagaatta ctgccgaaat ccagatgatg atgctcatgg accctggtgc tacacgggaa	2340
atccactcat tccttgggat tattgccta tttctcgttg tgaaggtgat accacaccta	2400
caatagtcaa tttagacat cccgtaatat cttgtgcaa aacgaaacaa ttgcgagttg	2460
taaatgggat tccaacacga acaaacatag gatggatggt tagtttgaga tacagaaata	2520
aacatatctg cggaggatca ttgataaagg agagttgggt tcttactgca cgacagtgtt	2580
tccttctctg agacttgaaa gattatgaag cttggcttgg aattcatgat gtccacggaa	2640
gaggagatga gaaatgcaaa caggttctca atgttttcca gctggtatat ggcctgaag	2700
gatcagatct ggttttaatg aagcttgcca ggctgctgt cdtggatgat tttgttagta	2760
cgattgattt acctaatat ggatgcacaa ttctgaaaa gaccagtgc agtgtttatg	2820
gctggggcta cactggattg atcaactatg atggcctatt acgagtggca catctctata	2880
taatgggaaa tgagaaatgc agccagcatc atcgagggaa ggtgactctg aatgagtctg	2940
aaatatgtgc tggggctgaa aagattggat caggaccatg tgagggggat tatggtggcc	3000
cacttgtttg tgagcaacat aaaatgagaa tggttcttgg tgtcattgtt cctggctctg	3060
gatgtgccat tccaaatcgt cctggtattt ttgtccgagt agcatattat gcaaaatgga	3120
tacacaaaat tattttaaca tataaggtag cacagtcata gctgttaacc cgggtcgaag	3180
cggccgcca cgtgctggat atctgcagaa ttccaccaca ctggactagt ggatccgagc	3240
tcggtacca gcttaagttt aaaccgctga tcagcctcga ctgtgccttc tagttgccag	3300
ccatctgttg ttgcccctc ccccggtgct tccttgacct tggaagggtgc cactccact	3360
gtccttctct aataaaaatga ggaattgca tcgcattgtc tgagtagggt tcattctatt	3420
ctgggggggt ggggtgggca ggacagcaag ggggaggatt gggaagacaa tagcaggcat	3480
gctgggggat cgggtgggtc tatggttctt caggcggaac gaaccagctg gggctctagg	3540
gggtatcccc acgcgccttg tagcggcgca ttaagcgcgg cgggtgtggt ggttacgcgc	3600
agcgtgaccg ctacacttgc cagcgcctta gcgcgcctc ctttcgcttt cttcccttc	3660
tttctcgcca cgttcgcggg ctttccccgt caagctctaa atcgggggct ccctttaggg	3720
ttccgattta gtgctttacg gcacctcgac ccaaaaaac ttgattaggg tgatggttca	3780
cgtagtgggc catcgccctg atagacggtt tttcgccctt tgacgttggg gtccacgttc	3840
tttaatatgt gactcttgtt ccaaaactga acaacactca accctatctc ggtctattct	3900
tttgatttat aagggtttt gccgatttgc gcctattggt taaaaaatga gctgatttaa	3960
caaaaaatta acgcgaatta attctgtgga atgtgtgtca gttagggtgt ggaagtccc	4020
caggctcccc agcaggcaga agtatgcaaa gcatgcatct caattagtca gcaaccaggt	4080
gtggaaagtc cccaggctcc ccagcaggca gaagtatgca aagcatgcat ctcaattagt	4140
cagcaaacat agtccccccc ctaactccgc ccaccccgcc cctaactccg cccagtccg	4200
cccattctcc gcccctggc tgactaatth tttttattta tgcagaggcc gaggcgcct	4260



-continued

ctgcctctga gctattccag aagtagtgag gaggcctttt tggaggccta ggcttttgca	4320
aaaagctccc gggagcttgt atatccattt tcggatctga tcaagagaca ggatgaggat	4380
cgtttcgcat gattgaacaa gatggattgc acgcagggtc tccggccgct tgggtggaga	4440
ggctattcgg ctatgactgg gcacaacaga caatcggctg ctctgatgcc gccgtgttcc	4500
ggctgtcagc gcaggggcgc ccgggttctt ttgtcaagac cgacctgtcc ggtgccctga	4560
atgaactgca ggacgaggca gcgcggctat cgtggctggc cacgacgggc gttccttgcg	4620
cagctgtgct cgacgttgtc actgaagcgg gaagggactg gctgctattg ggcgaagtgc	4680
cggggcagga tctcctgca tctcaccttg ctctgcccga gaaagtatcc atcatggctg	4740
atgcaatgcg gcggctgcat acgcttgatc cggctacctg cccattcgac caccaagcga	4800
aacatcgcat cgagcgagca cgtactcgga tggaagccgg tcttgtcgat caggatgatc	4860
tggacgaaga gcatcagggg ctccgccag ccgaactgtt cgccaggctc aaggcgcgca	4920
tgcccgacgg cgaggatctc gtcgtgaccc atggcgatgc ctgcttgccg aatatcatgg	4980
tggaaaatgg ccgcttttct ggattcatcg actgtggccg gctgggtgtg gcggaccgct	5040
atcaggacat agcgttggtt acccgtgata ttgctgaaga gcttggcggc gaatgggctg	5100
accgcttctc cgtgctttac ggtatcgccg ctcccgatcc gcagcgcacg gccttctatc	5160
gccttcttga cgagttcttc tgagcggggc tctgggggtc gaaatgaccg accaagcgac	5220
gcccacctg ccatcacgag atttcgatcc caccgcgcgc ttctatgaaa ggttgggctt	5280
cggaaatcgtt ttcgggacg ccggctggat gatcctccag cgcggggatc tcatgctgga	5340
gttcttcgcc caccccaact tgtttattgc agcttataat ggttacaagt aaagcaatag	5400
catcacaaat ttcacaaata aagcattttt ttactgcat tctagttgtg gtttgtccaa	5460
actcatcaat gtatcttata atgtctgtat accgtcgacc tctagctaga gcttggcgta	5520
atcatggta tagctgttcc ctgtgtgaaa ttgttatccg ctcaaatcc cacaacaat	5580
acgagccgga agcataaagt gtaaacctg gggcgctaa tgagtgcgtt aactcacatt	5640
aattgcgttg cgtcactgc ccgctttcca gtcgggaaac ctgctgtgcc agctgcatta	5700
atgaatcggc caacgcgcgg ggagaggcgg tttgcgtatt gggcgctctt ccgcttctc	5760
gctcactgac tcgctgcgct cggctgttcc gctgcggcga gcggtatcag ctcactcaaa	5820
ggcggttaata cggttatcca cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa	5880
aggccagcaa aaggccagga accgtaaaaa ggccgcgttg ctggcgcttt tccataggt	5940
ccgccccctc gacgagcatc acaaaaatcg acgctcaagt cagaggtggc gaaaccgac	6000
aggactataa agataccagg cgtttcccc ccggaagctcc ctctgcgct ctctgttcc	6060
gacctgcgcg cttaccggat acctgtccgc ctttctccct tcgggaagcg tggcgcttcc	6120
tcatagctca cgtgttaggt atctcagttc ggtgtaggtc gttcgctcca agctgggctg	6180
tgtgcacgaa cccccgctc agcccgaccg ctgcgcctta tccggtaact atcgtcttga	6240
gtccaacccg gtaagacacg acttatcgcc actggcagca gccactggtg acaggattag	6300
cagagcgagg tatgtaggcg gtgctacaga gttcttgaag tggtagccta actacggcta	6360
cactagaaga acagtatttg gtatctgcgc tctgctgaag ccagttacct tcggaaaaag	6420
agttggtagc tcttgatccg gcaaaaaaac caccgctggt agcggttttt ttgtttgcaa	6480
gcagcagatt acgcgcagaa aaaaaggatc tcaagaagat cctttgatct tttctacggg	6540

-continued

gtctgacgct cagtggaaacg aaaactcacg ttaagggatt ttggatcatga gattatcaaa	6600
aaggatcttc acctagatcc ttttaaatca aaaatgaagt ttttaaatcaa tctaaagtat	6660
atatgagtaa acttggtctg acagttacca atgcttaatc agtgaggcac ctatctcagc	6720
gatctgtcta tttcgttcat ccatagttgc ctgactcccc gtcgtgtaga taactacgat	6780
acgggagggc ttaccatctg gccccagtcg tgcaatgata ccgcgagacc cagctcacc	6840
ggctccagat ttatcagcaa taaaccagcc agccggaagg gccgagcgca gaagtgggtcc	6900
tgcaacttta tccgcctcca tccagtttat taattgttgc cggaagcta gagtaagtag	6960
ttcgccagtt aatagtttgc gcaacgttgc tgccattgct acaggcatcg tgggtgcacg	7020
ctcgtcgttt ggtatggctt cattcagctc cggttcccaa cgatcaaggc gagttacatg	7080
atccccatg ttgtgcaaaa aagcggttag ctcttcggt cctccgatcg ttgtcagaag	7140
taagtgtggc gcagtgttat cactcatggt tatggcagca ctgcataatt ctcttactgt	7200
catgccatcc gtaagatgct tttctgtgac tggtagtagc tcaaccaagt cattctgaga	7260
atagtgtatg cggcgaccga gttgtctctg cccggcgctc ataccgggata ataccgcgcc	7320
acatagcaga actttaaaag tgctcatcat tggaaaacgt tcttcggggc gaaaactctc	7380
aaggatctta ccgctgttga gatccagttc gatgtaacct actcgtgcac ccaactgatc	7440
ttcagcatct tttactttca ccagcgtttc tgggtgagca aaaacaggaa ggcaaatgc	7500
cgcaaaaaag ggaataaggg cgacacggaa atgttgaata ctcatactct tcctttttca	7560
atattattga agcattttatc aggggttattg tctcatgagc ggatacatat ttgaatgtat	7620
ttagaaaaat aaacaaatag gggttccgag cacatttccc cgaaaagtgc cactgacgt	7680
c	7681

1. A lymphangiogenesis-promoting agent comprising as an active ingredient an HGF or a protein or compound functionally equivalent to an HGF.

2. The agent of claim 1, wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):

- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
- (b) a protein encoded by a nucleic acid comprising the coding sequence in the nucleotide sequence of SEQ ID NO: 1;
- (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
- (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.

3. A lymphangiogenesis-promoting agent, which comprises as an active ingredient a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF.

4. The agent of claim 3, wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
- (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
- (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
- (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.

5. The agent of claim 3 or 4, wherein the nucleic acid has been inserted into a mammalian expression vector.

6. The agent of any one of claims 3 to 5, wherein the nucleic acid is a naked nucleic acid.

7. The agent of any one of claims 1 to 6, which is used as a pharmaceutical agent for preventing or treating lymphedema.

8. A method for promoting lymphangiogenesis, which comprises the step of administering to a subject an HGF or a protein or compound functionally equivalent to an HGF.

9. The method of claim 8, wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):

- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
  - (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.
- 10.** A method for promoting lymphangiogenesis, which comprises the step of administering to a subject a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF.
- 11.** The method of claim **10**, wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):
- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
  - (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.
- 12.** The method of claim **10** or **11**, wherein the nucleic acid has been inserted into a mammalian expression vector.
- 13.** The method of any one of claims **10** to **12**, wherein the nucleic acid is a naked nucleic acid.
- 14.** A method for inducing activation of an HGF receptor and promoting lymphangiogenesis through the activation.
- 15.** A method for preventing or treating lymphedema, which comprises the step of administering to a subject an HGF or a protein or compound functionally equivalent to an HGF.
- 16.** The method of claim **15**, wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):
- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
  - (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.
- 17.** A method for preventing or treating lymphedema, which comprises the step of administering to a subject a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF.
- 18.** The method of claim **17**, wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):
- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
  - (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.
- 19.** The method of claim **17** or **18**, wherein the nucleic acid has been inserted into a mammalian expression vector.
- 20.** The method of any one of claims **17** to **19**, wherein the nucleic acid is a naked nucleic acid.
- 21.** Use of an HGF or a protein or compound functionally equivalent to an HGF for producing a lymphangiogenesis-promoting agent or a pharmaceutical agent to be used to prevent or treat lymphedema.
- 22.** The use of claim **21**, wherein the HGF or protein functionally equivalent to an HGF is any one of the following (a) to (d):
- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
  - (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.
- 23.** Use of a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF for producing a lymphangiogenesis-promoting agent or a pharmaceutical agent to be used to prevent or treat lymphedema.
- 24.** The use of claim **23**, wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
  - (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.
- 25.** The use of claim **23** or **24**, wherein the nucleic acid has been inserted into a mammalian expression vector.
- 26.** The use of any one of claims **23** to **25**, wherein the nucleic acid is a naked nucleic acid.
- 27.** A method of screening for a compound having lymphangiogenesis-promoting activity or a compound having an effect of preventing or treating lymphedema, wherein the method comprises the following steps:
- (a) contacting a test compound with an HGF receptor or a protein functionally equivalent to an HGF receptor;
  - (b) detecting the binding between the protein and test compound; and
  - (c) selecting a test compound that binds to the protein.
- 28.** The method of claim **27**, wherein the HGF receptor or protein functionally equivalent to an HGF receptor is selected from the following (i) to (iv):
- (i) a protein comprising the amino acid sequence of SEQ ID NO: 4;
  - (ii) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 3;
  - (iii) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 4, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4; and
  - (iv) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 3, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4.
- 29.** A method of screening for a compound having lymphangiogenesis-promoting activity or a compound having an effect of preventing or treating lymphedema, wherein the method comprises the following steps:
- (a) contacting a test compound with a cell expressing an HGF receptor or a protein functionally equivalent to an HGF receptor;
  - (b) measuring the growth capacity or migratory activity of the cell, or phosphorylation of a signaling molecule; and
  - (c) selecting a test compound that increases the growth capacity or migratory activity of the cell, or causes phosphorylation of the signaling molecule, as compared to when the test compound is not contacted.
- 30.** The method of claim **29**, wherein the HGF receptor or protein functionally equivalent to an HGF receptor is selected from the following (i) to (iv):
- (i) a protein comprising the amino acid sequence of SEQ ID NO: 4;
  - (ii) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 3;
  - (iii) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 4, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4; and
  - (iv) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 3, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 4.
- 31.** A vector for preventing or treating lymphedema, which is a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted.
- 32.** The vector of claim **31**, wherein the nucleic acid encoding an HGF or the protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):
- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
  - (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.
- 33.** The vector of claim **31** or **32**, which comprises the nucleotide sequence of SEQ ID NO: 5 or 6.
- 34.** The vector of any one of claims **31** to **33**, which is administered in a naked state.
- 35.** The vector of any one of claims **31** to **34**, which is administered by intramuscular injection to or around an affected area in a subject.
- 36.** A pharmaceutical agent for preventing or treating lymphedema, which comprises as an active ingredient a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted, wherein the vector is administered in a naked state by intramuscular injection to or around an affected area in a subject.
- 37.** The pharmaceutical agent of claim **36**, wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):
- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
  - (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
  - (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ

ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and

- (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.

**38.** The pharmaceutical agent of claim **36** or **37**, wherein the vector comprises the nucleotide sequence of SEQ ID NO: 5 or 6.

**39.** A method for preventing or treating lymphedema, which comprises the step of administering, in a naked state, a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted, to or around an affected area in a subject by intramuscular injection.

**40.** The method of claim **39**, wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
- (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
- (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
- (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising

the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.

**41.** The method of claim **40**, wherein the vector comprises the nucleotide sequence of SEQ ID NO: 5 or 6.

**42.** Use of a mammalian expression vector into which a nucleic acid encoding an HGF or a protein functionally equivalent to an HGF has been inserted, for producing a pharmaceutical agent for preventing or treating lymphedema, wherein the vector is administered in a naked state to or around an affected area in a subject by intramuscular injection.

**43.** The use of claim **42**, wherein the nucleic acid encoding an HGF or a protein functionally equivalent to an HGF is a nucleic acid encoding any one of the following (a) to (d):

- (a) a protein comprising the amino acid sequence of SEQ ID NO: 2;
- (b) a protein encoded by a nucleic acid comprising the coding region of the nucleotide sequence of SEQ ID NO: 1;
- (c) a protein comprising an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of SEQ ID NO: 2, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2; and
- (d) a protein encoded by a nucleic acid that hybridizes under stringent conditions to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, which is functionally equivalent to a protein comprising the amino acid sequence of SEQ ID NO: 2.

**44.** The use of claim **42** or **43**, wherein the vector comprises the nucleotide sequence of SEQ ID NO: 5 or 6.

\* \* \* \* \*