



US 20090076097A1

(19) **United States**

(12) **Patent Application Publication**

Czarnik

(10) **Pub. No.: US 2009/0076097 A1**

(43) **Pub. Date: Mar. 19, 2009**

(54) **DEUTERIUM-ENRICHED ATAZANAVIR**

Related U.S. Application Data

(75) Inventor: **Anthony W. Czarnik**, Reno, NV
(US)

(60) Provisional application No. 60/972,690, filed on Sep.
14, 2007.

Publication Classification

Correspondence Address:

VANCE INTELLECTUAL PROPERTY, PC

5467 HILL TOP STREET

CROZET, VA 22932-3167 (US)

(51) **Int. Cl.**
A61K 31/44 (2006.01)
C07D 213/04 (2006.01)
A61P 31/12 (2006.01)

(73) Assignee: **PROTIA, LLC**, Reno, NV (US)

(52) **U.S. Cl.** **514/357**; 546/332

(21) Appl. No.: **12/196,255**

(57) **ABSTRACT**

(22) Filed: **Aug. 21, 2008**

The present application describes deuterium-enriched atazanavir, pharmaceutically acceptable salt forms thereof, and methods of treating using the same.

DEUTERIUM-ENRICHED ATAZANAVIR**CROSS-REFERENCE TO RELATED APPLICATIONS**


[0001] The present application claims priority benefit under 35 U.S.C. § 119(e) of U.S. Provisional Patent Application Ser. No. 60/972,690 filed 14 Sep. 2007. The disclosure of this application is incorporated herein by reference.

FIELD OF THE INVENTION

[0002] This invention relates generally to deuterium-enriched atazanavir, pharmaceutical compositions containing the same, and methods of using the same.

BACKGROUND OF THE INVENTION

[0003] Atazanavir, shown below, is a well known protease inhibitor.



Since atazanavir is a known and useful pharmaceutical, it is desirable to discover novel derivatives thereof. Atazanavir is described in U.S. Pat. Nos. 5,849,911, 6,300,519, and 6,166,004; the contents of which are incorporated herein by reference.

SUMMARY OF THE INVENTION

[0004] Accordingly, one object of the present invention is to provide deuterium-enriched atazanavir or a pharmaceutically acceptable salt thereof.

[0005] It is another object of the present invention to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the deuterium-enriched compounds of the present invention or a pharmaceutically acceptable salt thereof.

[0006] It is another object of the present invention to provide a method for treating human immunodeficiency virus, comprising administering to a host in need of such treatment

a therapeutically effective amount of at least one of the deuterium-enriched compounds of the present invention or a pharmaceutically acceptable salt thereof.

[0007] It is another object of the present invention to provide a novel deuterium-enriched atazanavir or a pharmaceutically acceptable salt thereof for use in therapy.

[0008] It is another object of the present invention to provide the use of a novel deuterium-enriched atazanavir or a pharmaceutically acceptable salt thereof for the manufacture of a medicament (e.g., for the treatment of human immunodeficiency virus).

[0009] These and other objects, which will become apparent during the following detailed description, have been achieved by the inventor's discovery of the presently claimed deuterium-enriched atazanavir.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS


[0010] Deuterium (D or ^2H) is a stable, non-radioactive isotope of hydrogen and has an atomic weight of 2.0144.

Hydrogen naturally occurs as a mixture of the isotopes ^1H (hydrogen or protium), D (^2H or deuterium), and T (^3H or tritium). The natural abundance of deuterium is 0.015%. One of ordinary skill in the art recognizes that in all chemical compounds with a H atom, the H atom actually represents a mixture of H and D, with about 0.015% being D. Thus, compounds with a level of deuterium that has been enriched to be greater than its natural abundance of 0.015%, should be considered unnatural and, as a result, novel over their non-enriched counterparts.

[0011] All percentages given for the amount of deuterium present are mole percentages.

[0012] It can be quite difficult in the laboratory to achieve 100% deuteration at any one site of a lab scale amount of compound (e.g., milligram or greater). When 100% deuteration is recited or a deuterium atom is specifically shown in a structure, it is assumed that a small percentage of hydrogen may still be present. Deuterium-enriched can be achieved by either exchanging protons with deuterium or by synthesizing the molecule with enriched starting materials.

[0013] The present invention provides deuterium-enriched atazanavir or a pharmaceutically acceptable salt thereof. There are fifty-two hydrogen atoms in the atazanavir portion of atazanavir as show by variables R_1 - R_{52} in formula I below.



[0014] The hydrogens present on atazanavir have different capacities for exchange with deuterium. Hydrogen atoms R_1 - R_5 are easily exchangeable under physiological conditions and, if replaced by deuterium atoms, it is expected that they will readily exchange for protons after administration to a patient. The remaining hydrogen atoms are not easily exchangeable for deuterium atoms. However, deuterium atoms at the remaining positions may be incorporated by the use of deuterated starting materials or intermediates during the construction of atazanavir.

[0015] The present invention is based on increasing the amount of deuterium present in atazanavir above its natural abundance. This increasing is called enrichment or deuterium-enrichment. If not specifically noted, the percentage of enrichment refers to the percentage of deuterium present in the compound, mixture of compounds, or composition. Examples of the amount of enrichment include from about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 16, 21, 25, 29, 33, 37, 42, 46, 50, 54, 58, 63, 67, 71, 75, 79, 84, 88, 92, 96, to about 100 mol %. Since there are 52 hydrogens in atazanavir, replacement of a single hydrogen atom with deuterium would result in a molecule with about 2% deuterium enrichment. In order to achieve enrichment less than about 2%, but above the natural abundance, only partial deuteration of one site is required. Thus, less than about 2% enrichment would still refer to deuterium-enriched atazanavir.

[0016] With the natural abundance of deuterium being 0.015%, one would expect that for approximately every 6,667 molecules of atazanavir ($1/0.00015=6,667$), there is one naturally occurring molecule with one deuterium present. Since atazanavir has 52 positions, one would roughly expect that for approximately every 346,684 molecules of atazanavir ($52 \times 6,667$), all 52 different, naturally occurring, mono-deuterated

atazanavirs would be present. This approximation is a rough estimate as it doesn't take into account the different exchange rates of the hydrogen atoms on atazanavir. For naturally occurring molecules with more than one deuterium, the num-

bers become vastly larger. In view of this natural abundance, the present invention, in an embodiment, relates to an amount of a deuterium enriched compound, whereby the enrichment recited will be more than naturally occurring deuterated molecules.

[0017] In view of the natural abundance of deuterium-enriched atazanavir, the present invention also relates to isolated or purified deuterium-enriched atazanavir. The isolated or purified deuterium-enriched atazanavir is a group of molecules whose deuterium levels are above the naturally occurring levels (e.g., 2%). The isolated or purified deuterium-enriched atazanavir can be obtained by techniques known to those of skill in the art (e.g., see the syntheses described below).

[0018] The present invention also relates to compositions comprising deuterium-enriched atazanavir. The compositions require the presence of deuterium-enriched atazanavir which is greater than its natural abundance. For example, the compositions of the present invention can comprise (a) a μg of a deuterium-enriched atazanavir; (b) a mg of a deuterium-enriched atazanavir; and, (c) a gram of a deuterium-enriched atazanavir.

[0019] In an embodiment, the present invention provides an amount of a novel deuterium-enriched atazanavir.


[0020] Examples of amounts include, but are not limited to (a) at least 0.01, 0.02, 0.03, 0.04, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, to 1 mole, (b) at least 0.1 moles, and (c) at least 1 mole of the compound. The present amounts also cover lab-scale (e.g., gram scale), kilo-lab scale (e.g., kilogram scale), and industrial or commercial scale (e.g., multi-kilogram or above scale) quantities as these will be more useful in the actual manufacture of a pharmaceutical. Industrial/commercial scale refers

to the amount of product that would be produced in a batch that was designed for clinical testing, formulation, sale/distribution to the public, etc.

[0021] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof.

is at least 13%. The abundance can also be (a) at least 25%, (b) at least 38%, (c) at least 50%, (d) at least 63%, (e) at least 75%, (f) at least 88%, and (g) 100%.

[0027] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abun-



[0022] wherein R₁-R₅₂ are independently selected from H and D; and the abundance of deuterium in R₁-R₅₂ is at least 2%. The abundance can also be (a) at least 4%, (b) at least 10%, (c) at least 15%, (d) at least 21%, (e) at least 27%, (f) at least 33%, (g) at least 38%, (h) at least 44%, (i) at least 50%, (j) at least 56%, (k) at least 62%, (l) at least 67%, (m) at least 73%, (n) at least 79%, (o) at least 85%, (p) at least 90%, (q) at least 96%, and (r) 100%.

[0023] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₁-R₅ is at least 20%. The abundance can also be (a) at least 40%, (b) at least 60%, (c) at least 80%, and (d) 100%.

[0024] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₆-R₈ is at least 20%. The abundance can also be (a) at least 40%, (b) at least 60%, (c) at least 80%, and (d) 100%.

[0025] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₉-R₁₈ is at least 10%. The abundance can also be (a) at least 20%, (b) at least 30%, (c) at least 40%, (d) at least 50%, (e) at least 60%, (f) at least 70%, (g) at least 80%, (h) at least 90%, and (i) 100%.

[0026] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I, wherein the abundance of deuterium in R₁₉-R₂₁ and R₂₇-R₃₁

dance of deuterium in R₂₂-R₂₆ is at least 25%. The abundance can also be (a) at least 50%, (b) at least 75%, and (c) 100%.

[0028] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₃₂-R₃₅ is at least 25%. The abundance can also be (a) at least 50%, (b) at least 75%, and (c) 100%.


[0029] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₃₆-R₃₉ is at least 5%. The abundance can also be (a) at least 9%, (b) at least 12%, (c) at least 18%, (d) at least 23%, (e) at least 27%, (f) at least 32%, (g) at least 36%, (h) at least 41%, (i) at least 45%, (j) at least 50%, (k) at least 55%, (l) at least 59%, (m) at least 62%, (n) at least 68%, (o) at least 73%, (p) at least 77%, (q) at least 82%, (r) at least 86%, (s) at least 91%, (t) at least 95%, and (u) 100%.

[0030] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₄₀-R₄₉ is at least 10%. The abundance can also be (a) at least 20%, (b) at least 30%, (c) at least 40%, (d) at least 50%, (e) at least 60%, (f) at least 70%, (g) at least 80%, (h) at least 90%, and (i) 100%.

[0031] In another embodiment, the present invention provides a novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₅₀-R₅₂ is at least 33%. The abundance can also be (a) at least 67%, and (b) 100%.

[0032] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof.

[0038] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof,



[0033] wherein R₁-R₅₂ are independently selected from H and D; and the abundance of deuterium in R₁-R₅₂ is at least 2%. The abundance can also be (a) at least 4%, (b) at least 10%, (c) at least 15%, (d) at least 21%, (e) at least 27%, (f) at least 33%, (g) at least 38%, (h) at least 44%, (i) at least 50%, (j) at least 56%, (k) at least 62%, (l) at least 67%, (m) at least 73%, (n) at least 79%, (o) at least 85%, (p) at least 90%, (q) at least 96%, and (r) 100%.

[0034] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₁-R₅ is at least 20%. The abundance can also be (a) at least 40%, (b) at least 60%, (c) at least 80%, and (d) 100%.

[0035] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₆-R₉ is at least 20%. The abundance can also be (a) at least 40%, (b) at least 60%, (c) at least 80%, and (d) 100%.

[0036] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₉-R₁₈ is at least 10%. The abundance can also be (a) at least 20%, (b) at least 30%, (c) at least 40%, (d) at least 50%, (e) at least 60%, (f) at least 70%, (g) at least 80%, (h) at least 90%, and (i) 100%.

[0037] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I, wherein the abundance of deuterium in R₁₉-R₂₁ and R₂₇-R₃₁ is at least 13%. The abundance can also be (a) at least 25%, (b) at least 38%, (c) at least 50%, (d) at least 63%, (e) at least 75%, (f) at least 88%, and (g) 100%.

wherein the abundance of deuterium in R₂₂-R₂₆ is at least 25%. The abundance can also be (a) at least 50%, (b) at least 75%, and (c) 100%.


[0039] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₃₂-R₃₅ is at least 25%. The abundance can also be (a) at least 50%, (b) at least 75%, and (c) 100%.

[0040] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₃₆-R₃₉ is at least 5%. The abundance can also be (a) at least 9%, (b) at least 12%, (c) at least 18%, (d) at least 23%, (e) at least 27%, (f) at least 32%, (g) at least 36%, (h) at least 41%, (i) at least 45%, (j) at least 50%, (k) at least 55%, (l) at least 59%, (m) at least 62%, (n) at least 68%, (o) at least 73%, (p) at least 77%, (q) at least 82%, (r) at least 86%, (s) at least 91%, (t) at least 95%, and (u) 100%.

[0041] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₄₀-R₄₉ is at least 10%. The abundance can also be (a) at least 20%, (b) at least 30%, (c) at least 40%, (d) at least 50%, (e) at least 60%, (f) at least 70%, (g) at least 80%, (h) at least 90%, and (i) 100%.

[0042] In another embodiment, the present invention provides an isolated novel, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R₅₀-R₅₂ is at least 33%. The abundance can also be (a) at least 67%, and (b) 100%.

[0043] In another embodiment, the present invention provides novel mixture of deuterium enriched compounds of formula I or a pharmaceutically acceptable salt thereof.



[0044] wherein R_1 - R_{52} are independently selected from H and D; and the abundance of deuterium in R_1 - R_{52} is at least 2%. The abundance can also be (a) at least 4%, (b) at least 10%, (c) at least 15%, (d) at least 21%, (e) at least 27%, (f) at least 33%, (g) at least 38%, (h) at least 44%, (i) at least 50%, (j) at least 56%, (k) at least 62%, (l) at least 67%, (m) at least 73%, (n) at least 79%, (o) at least 85%, (p) at least 90%, (q) at least 96%, and (r) 100%.

[0045] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_1 - R_5 is at least 20%. The abundance can also be (a) at least 40%, (b) at least 60%, (c) at least 80%, and (d) 100%.

[0046] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_6 - R_8 is at least 20%. The abundance can also be (a) at least 40%, (b) at least 60%, (c) at least 80%, and (d) 100%.

[0047] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_9 - R_{18} is at least 10%. The abundance can also be (a) at least 20%, (b) at least 30%, (c) at least 40%, (d) at least 50%, (e) at least 60%, (f) at least 70%, (g) at least 80%, (h) at least 90%, and (i) 100%.

[0048] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I, wherein the abundance of deuterium in R_{19} - R_{21} and R_{27} - R_{31} is at least 13%. The abundance can also be (a) at least 25%, (b) at least 38%, (c) at least 50%, (d) at least 63%, (e) at least 75%, (f) at least 88%, and (g) 100%.

[0049] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_{22} - R_{26} is at least 25%. The abundance can also be (a) at least 50%, (b) at least 75%, and (c) 100%.

[0050] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_{32} - R_{35} is at least 25%. The abundance can also be (a) at least 50%, (b) at least 75%, and (c) 100%.

[0051] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_{36} - R_{39} is at least 5%. The abundance can also be (a) at least 9%, (b) at least 12%, (c) at least 18%, (d) at least 23%, (e) at least 27%, (f) at least 32%, (g) at least 36%, (h) at least 41%, (i) at least 45%, (j) at least 50%, (k) at least 55%, (l) at least 59%, (m) at least 62%, (n) at least 68%, (o) at least 73%, (p) at least 77%, (q) at least 82%, (r) at least 86%, (s) at least 91%, (t) at least 95%, and (u) 100%.

[0052] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_{40} - R_{49} is at least 10%. The abundance can also be (a) at least 20%, (b) at least 30%, (c) at least 40%, (d) at least 50%, (e) at least 60%, (f) at least 70%, (g) at least 80%, (h) at least 90%, and (i) 100%.

[0053] In another embodiment, the present invention provides a novel mixture of, deuterium enriched compound of formula I or a pharmaceutically acceptable salt thereof, wherein the abundance of deuterium in R_{50} - R_{52} is at least 33%. The abundance can also be (a) at least 67%, and (b) 100%.

[0054] In another embodiment, the present invention provides novel pharmaceutical compositions, comprising: a pharmaceutically acceptable carrier and a therapeutically effective amount of a deuterium-enriched compound of the present invention.

[0055] In another embodiment, the present invention provides a novel method for treating human immunodeficiency virus comprising: administering to a patient in need thereof a therapeutically effective amount of a deuterium-enriched compound of the present invention.

[0056] In another embodiment, the present invention provides an amount of a deuterium-enriched compound of the present invention as described above for use in therapy.

[0057] In another embodiment, the present invention provides the use of an amount of a deuterium-enriched compound of the present invention for the manufacture of a medicament (e.g., for the treatment of human immunodeficiency virus).

[0058] The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. This invention encompasses all combinations of preferred aspects of the invention noted herein. It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment or embodiments to describe additional more preferred embodiments. It is also to be understood that each individual element of the preferred embodiments is intended to be taken individually as its own independent preferred embodiment. Furthermore, any element of an embodiment is meant to be combined with any and all other elements from any embodiment to describe an additional embodiment.

DEFINITIONS

[0059] The examples provided in the definitions present in this application are non-inclusive unless otherwise stated. They include but are not limited to the recited examples.

[0060] The compounds of the present invention may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. All processes used to prepare compounds of the present invention and intermediates made therein are considered to be part of the present invention. All tautomers of shown or described compounds are also considered to be part of the present invention.

[0061] "Host" preferably refers to a human. It also includes other mammals including the equine, porcine, bovine, feline, and canine families.

[0062] "Treating" or "treatment" covers the treatment of a disease-state in a mammal, and includes: (a) preventing the disease-state from occurring in a mammal, in particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it; (b) inhibiting the disease-state, e.g., arresting its development; and/or (c) relieving the disease-state, e.g., causing regression of the disease state until

a desired endpoint is reached. Treating also includes the amelioration of a symptom of a disease (e.g., lessen the pain or discomfort), wherein such amelioration may or may not be directly affecting the disease (e.g., cause, transmission, expression, etc.).


[0063] "Therapeutically effective amount" includes an amount of a compound of the present invention that is effective when administered alone or in combination to treat the desired condition or disorder. "Therapeutically effective amount" includes an amount of the combination of compounds claimed that is effective to treat the desired condition or disorder. The combination of compounds is preferably a synergistic combination. Synergy, as described, for example, by Chou and Talalay, *Adv. Enzyme Regul.* 1984, 22:27-55, occurs when the effect of the compounds when administered in combination is greater than the additive effect of the compounds when administered alone as a single agent. In general, a synergistic effect is most clearly demonstrated at sub-optimal concentrations of the compounds. Synergy can be in terms of lower cytotoxicity, increased antiviral effect, or some other beneficial effect of the combination compared with the individual components.

[0064] "Pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of the basic residues. The pharmaceutically acceptable salts include the conventional quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include, but are not limited to, those derived from inorganic and organic acids selected from 1, 2-ethanedithiolonic, 2-acetoxybenzoic, 2-hydroxyethanesulfonic, acetic, ascorbic, benzenesulfonic, benzoic, bicarbonic, carbonic, citric, edetic, ethane disulfonic, ethane sulfonic, fumaric, glucoheptonic, gluconic, glutamic, glycolic, glycolylarsanilic, hexylresorcinic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxymaleic, hydroxynaphthoic, isethionic, lactic, lactobionic, lauryl sulfonic, maleic, malic, mandelic, methane-sulfonic, napsylic, nitric, oxalic, pamoic, pantothenic, phenylacetic, phosphoric, polygalacturonic, propionic, salicylic, stearic, subacetic, succinic, sulfamic, sulfanilic, sulfuric, tartaric, and toluenesulfonic.


Synthesis

[0065]


Scheme 1 shows an example of how to prepare atazanavir.



-continued




-continued



[0066] Using combinations of various deuterated starting materials and intermediates shown in Scheme 1, a person skilled in the art of organic chemistry should be able to prepare a wide variety of deuterated atazanavir analogs.


EXAMPLES

[0067] Table 1 provides compounds that are representative examples of the present invention. When one of R₁-R₅₂ is present, it is selected from H or D.




-continued

2




3




-continued

4




5




-continued

6




7




-continued

8




9




-continued

10




[0068] Table 2 provides compounds that are representative examples of the present invention. Where H is shown, it represents naturally abundant hydrogen.

11




-continued

12




13




-continued

14




15




-continued

16




17




-continued

18




19



-continued


20



[0069] Numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

What is claimed is:

1. A deuterium-enriched compound of formula I or a pharmaceutically acceptable salt thereof:



wherein R_1 - R_{52} are independently selected from H and D;
and

the abundance of deuterium in R_1 - R_{52} is at least 2%.

2. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_1 - R_{52} is selected from at least 2%, at least 4%, at least 10%, at least 15%, at least 21%, at least 27%, at least 33%, at least 38%, at least 44%, at least 50%, at least 56%, at least 62%, at least 67%, at least 73%, at least 79%, at least 85%, at least 90%, at least 96%, and 100%.

3. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_1 - R_5 is selected from at least 20%, at least 40%, at least 60%, at least 80%, and 100%.

4. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_6 - R_8 is selected from at least 33%, at least 67%, and 100%.

5. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_9 - R_{18} is selected from at least

8. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_{32} - R_{35} is selected from at least 25%, at least 50%, at least 75%, and 100%.

9. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_{36} - R_{39} is selected from at least 25%, at least 50%, at least 75%, and 100%.


10. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_{40} - R_{49} is selected from at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, and 100%.

11. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_{50} - R_{52} is selected from at least 33%, at least 67%, and 100%.

12. A deuterium-enriched compound of claim 1, wherein the compound is selected from compounds 1-10 of Table 1:

13. A deuterium-enriched compound of claim 1, wherein the compound is selected from compounds 11-20 of Table 2:

14. An isolated deuterium-enriched compound of formula I or a pharmaceutically acceptable salt thereof:



10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, and 100%.

6. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_{19} - R_{21} and R_{27} - R_{31} is selected from at least 13%, at least 25%, at least 38%, at least 50%, at least 63%, at least 75%, at least 88%, and 100%.

7. A deuterium-enriched compound of claim 1, wherein the abundance of deuterium in R_{22} - R_{26} is selected from at least 20%, at least 40%, at least 60%, at least 80%, and 100%.


wherein R_1 - R_{52} are independently selected from H and D;
and

the abundance of deuterium in R_1 - R_{52} is at least 2%.

15. An isolated deuterium-enriched compound of claim 14, wherein the compound is selected from compounds 1-10 of Table 1.

16. An isolated deuterium-enriched compound of claim 14, wherein the compound is selected from compounds 11-20 of Table 2.

17. A mixture of deuterium-enriched compounds of formula I or a pharmaceutically acceptable salt thereof:



wherein R_1 - R_{52} are independently selected from H and D;

and

the abundance of deuterium in R_1 - R_{52} is at least 2%.

18. A mixture of deuterium-enriched compound of claim 27, wherein the compound is selected from compounds 1-10 of Table 1.

19. A mixture of deuterium-enriched compound of claim 27, wherein the compound is selected from compounds 11-20 of Table 2.

20. A pharmaceutical composition, comprising: a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 1 or a pharmaceutically acceptable salt form thereof.

21. A method for treating human immunodeficiency virus comprising: administering, to a patient in need thereof, a therapeutically effective amount of a compound of claim 1 or a pharmaceutically acceptable salt form thereof.

* * * * *