



US006870031B2

(12) **United States Patent**
Hilton et al.

(10) **Patent No.:** **US 6,870,031 B2**
(45) **Date of Patent:** **Mar. 22, 2005**

(54) **HAEMOPOIETIN RECEPTOR AND
GENETIC SEQUENCES ENCODING SAME**

(75) Inventors: **Douglas James Hilton**, Warrandyte
(AU); **Tracy Willson**, North Balwyn
(AU); **Timothy Gainsford**, North
Carlton (AU); **Warren S Alexander**,
Moonee Ponds (AU); **Donald Metcalf**,
Balwyn (AU); **Ashley Ng**, Balwyn
(AU); **Nicos A Nicola**, Mont Albert
(AU)

(73) Assignee: **Amrad Operations Pty. Ltd.**, Victoria
(AU)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 204 days.

(21) Appl. No.: **10/014,156**

(22) Filed: **Dec. 7, 2001**

(65) **Prior Publication Data**

US 2005/0004015 A1 Jan. 6, 2005

Related U.S. Application Data

(62) Division of application No. 09/043,816, filed on Sep. 17,
1998, now Pat. No. 6,414,128.

(30) **Foreign Application Priority Data**

Sep. 26, 1995 (AU) PN 5641
Sep. 26, 1996 (AU) PCT/AU96/00607

(51) **Int. Cl.⁷** **C07K 14/705**

(52) **U.S. Cl.** **530/350; 514/2**

(58) **Field of Search** **530/350; 514/2**

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,869,610 A * 2/1999 Snodgrass et al. 530/350

OTHER PUBLICATIONS

X. X. Du, et al. (1994), "Interleukin-11: A Multifunctional
Growth Factor Derived From the Hematopoietic Microen-
vironment", *Blood*, 83: pp. 2023-2030.

Y.C. Yang, et al. (1992), "Interleukin-11 and Its Receptor",
Biofactors, 4: pp. 15-21.

S.R. Paul, et al. (1990), "Molecular Cloning of a cDNA
Encoding Interleukin 11, a Stromal Cell-Derived Lympho-
poietic and Hematopoietic Cytokine", *Proc. Natl. Acad. Sci.
USA*, 87: pp. 7512-7516.

M. Musashi, et al. (1991), "Synergistic Interactions Between
Interleukin-11 and Interleukin-4 in Support of Proliferation
of Primitive Hematopoietic Progenitors of Mice", *Blood*, 78:
pp. 1448-1451.

K.R. Schibler, et al. (1992), "Effect of Interleukin-11 on
Cycling Status and Clonogenic Maturation of Fetal and
Adult Hematopoietic Progenitors", *Blood*, 4: pp. 900-903.

K. Tsuji, et al. (1992), "Enhancement of Murine Hematopoie-
sis by Synergistic Interactions Between Steel Factor (ligand
for c-kit), Interleukin-11, and Other Early Acting Factors in
Culture", *Blood*, 79: pp. 2855-2860.

Samuel A. Burstein, et al. (1992), "Leukemia Inhibitory
Factor and Interleukin-11 Promote Maturation of Murine
and Human Megakaryocytes in Vitro", *J. Cell. Physiol*, 153:
pp. 305-312.

Giao Hangoc, et al. (1993), "In Vivo Effects of Recombinant
Interleukin-11 on Myelopoiesis in Mice", *Blood*, 81: pp.
965-972.

Masanao Teramura, et al. (1992), "Interleukin-11 Enhances
Human Megakaryocytopoiesis in Vitro", *Blood*, 79: pp.
327-331.

Yuji Yonemura, et al., (1992), "Synergistic Effects of Inter-
leukin 3 and Interleukin-11 on Murine Megakaryopoiesis in
Serum-free Culture", *Exp. Hematol*, 20: pp. 1011-1016.

Heinz Baumann, et al., (1991), "Interleukin-11 Regulates
the Hepatic Expression of the Same Plasma Protein Genes as
Interleukin-6", *J. Biol. Chem.*, 266: pp. 20424-20427.

Ichiro Kawashima, et al. (1991), "Molecular Cloning of
cDNA Encoding Adipogenesis Inhibitory Factor and Identi-
tity with Interleukin-11", *Febs. Lett.*, 283: pp. 199-202.

D.C. Keller, et al. (1993), "Interleukin-11 Inhibits Adipo-
genesis and Stimulates Myelopoiesis in Human Long-Term
Marrow Cultures", *Blood*, 82: pp. 1428-1435.

J. Sambrook, et al., (1989), "Molecular Cloning", *A Labo-
ratory Manual Cold Spring Harbor Laboratory*, Cold
Spring Harbor, NY.

Douglas J. Hilton, et al. (1994), "Cloning of a Murine IL-11
Receptor α -chain; Requirement for gp130 for High Affinity
Binding and Signal Transduction", *EMBO Journal*, 13: pp.
4765-4777.

Da-Wei Gong, et al. (1996), "Genomic Structure and Pro-
moter Analysis of the Human obese Gene", *J. Biol. Chem.*,
271: pp. 3971-3974.

Seiichi Mizushima, et al. (1990), "pEF-BOS, a Powerful
Mammalian Expression Vector", *Nucleic Acids Research*,
18: pp. 5322.

Nicos A. Nicola (1996), "Molecular Cloning of Two Novel
Transmembrane Ligands for EphRelated Kinases (LERKS)
that are Related to LERK-2", *Growth Factors*, 13: pp.
141-149.

(List continued on next page.)

Primary Examiner—John Ulm

(74) *Attorney, Agent, or Firm*—Scully, Scott, Murphy &
Presser

(57) **ABSTRACT**

The present invention is directed to a novel haemopoietin
receptor or a derivative thereof and to genetic sequences
encoding same. The receptor molecule and its derivatives
and the genetic sequences encoding same of the present
invention are useful in the development of a wide range of
agonists, antagonists, therapeutics and diagnostic reagents
based on ligand interaction with its receptor. The present
invention particularly relates to a receptor for leptin.

4 Claims, 20 Drawing Sheets

OTHER PUBLICATIONS

Douglas J. Hilton, et al. (1992), "Kinetic Analyses of the Binding of Leukemia Inhibitory Factor to Receptors on Cells and Membranes and in Detergent Solution", *J. Biol. Chem.*, 267: 10238–10247.

George Scatchard (1949), "The Attractions of Proteins for Small Molecules and Ions", *Ann. N.Y. Acad. Sciences*, 51: 660–672.

Unit 3.3.1 (1995), "Mapping by Partial Endonuclease Digestions", *Current Protocols in Molecular Biology*.

Marvin B. Shapiro, et al. (1987), "RNA Splice Junctions of Different Classes of Eukaryotes: Sequence Statistics and Functional Implications in Gene Expression", *Nucleic Acids Research*, 15: 7155–7174.

* cited by examiner




Figure 1

-255 cgaattcgcgggcgcg
-240 gtcgaccgcggncccagctcgggagacatggggggcggttaaagctctcgtggnattatcc
-180 ttcagtggggstattggactgactttttcttatgctgggatgtgccttagaggattatgga
-120 tttggcagttcacccctgaccatcttgaaaataagttatctctgatctctgtctgtatggt
-60 acttctctccccctcaccaacggagaacaaatgtgggcaaagtgtacttctctgaagtaag
1 ATGATTTGTCAAAAATTCTGTGTGGTTTTGTTACATTGGGAATTTATTTATGTGATAACT
1 M I C Q K F C V V L L H W E F I Y V I T
61 GCGTTTAACTTGTCATATCCAATTACTCCTTGGAGATTTAAGTTGTCTTGCATGCCACCA
21 A F N L S Y P I T P W R F K L S C M P P
121 AATTCAACCTATGACTACTTCCTTTTGCCTGCTGGACTCTCAAAGAATACTTCAAATTCTG
41 N S T Y D Y F L L P A G L S K N T S N S

A — — — — — **A**

Figure 2A

**Figure 2B**

**Figure 2C**

**Figure 2D**

D — — — — — **D**

1081 GTTCCCTCAAAAGAGATTGTTTGGTGGATGAATTTAGCTGAGAAAATTCCTCAAAGCCAG
361 V P S K E I V W W M N L A E K I P Q S Q

1141 TATGATGTTGTGAGTGATCATGTTAGCAAAGTTACTTTTTTCAATCTGAATGAAACCAAA
381 Y D V V S D H V S K V T F F N L N E T K

1201 CCTCGAGGAAAGTTTACCTATGATGCAGTGTACTGCTGCAATGAACATGAATGCCATCAT
401 P R G K F T Y D A V Y C C N E H E C H H

1261 CGCTATGCTGAATTATATGTGATTGATGTCAATATCAATATCTCATGTGAAACTGATGGG
421 R Y A E L Y V I D V N I N I S C E T D G

1321 TACTTAACTAAAATGACTTGCAGATGGTCAACCAGTACAATCCAGTCACTTGCGGAAAGC
441 Y L T K M T C R W S T S T I Q S L A E S

E — — — — — **E**

Figure 2E

E — — — — — **E**

1381 ACTTTGCAATTGAGGTATCATAGGAGCAGCCTTTACTGTTCTGATATTCCATCTATTCAT
461 T L Q L R Y H R S S L Y C S D I P S I H

1441 CCCATATCTGAGCCCAAAGATTGCTATTTGCAGAGTGATGGTTTTTATGAATGCATTTTC
481 P I S E P K D C Y L Q S D G F Y E C I F

1501 CAGCCAATCTTCCTATTATCTGGCTACACAATGTGGATTAGGATCAATCACTCTCTAGGT
501 Q P I F L L S G Y T M W I R I N H S L G

1561 TCACTTGACTCTCCACCAACATGTGTCCTTCCTGATTCTGTGGTGAAGcCACTGcCTCCA
521 S L D S P P T C V L P D S V V K P L P P

1621 TCCAGTGTGAAAGCAGAAATTACTATAAACATTGGATTATTGAAAATATCTTGGGAAAAG
541 S S V K A E I T I N I G L L K I S W E K

F — — — — — **F**

Figure 2F

**Figure 2G**

G — — — — — **G**

1981 ACTTTACTTTGGAAGCCCCTGATGAAAAATGACTCATTGTGCAGTGTTTCAGAGATATGTG
 661 T L L W K P L M K N D S L C S V Q R Y V

2041 ATAAACCATCATACTTCCTsCAATGGAACATGGTCAGAAGATGTGGGAAATCACACGAAA
 681 I N H H T S X N G T W S E D V G N H T K

2101 TTCACTTTCCTGTGGACAGAGCAAGCACATACTGTTACGGTTCTGGCCATCAATTCAATT
 701 F T F L W T E Q A H T V T V L A I N S I

2161 GGTGCTTCTGTTgCaAATtTTAATTTAACCTTTTTCATGGCCtATGAGCAAAGTAAATATC
 721 G A S V A N F N L T F S W P M S K V N I

2221 GTGCAGTCACTCAGTGCTTATCCTTTAAACAGCAGTTGTGTGATTGTTTCCTGGATACTA
 741 V Q S L S A Y P L N S S C V I V S W I L

H — — — — — **H**

Figure 2H

H — — — — — **H**

2281 TCACCCAGTGATTACAAGCTAATGTATTTTATTATTGAGTGGAAAAATCTTAATGAAGAT
761 S P S D Y K L M Y F I I E W K N L N E D

2341 GGTGAAATAAAATGGCTTAGAATCTCTTCATCTGTTAAGAAGTATTATATCCATGATCAT
781 G E I K W L R I S S S V K K Y Y I H D H

2401 TTTATCCCCATTGAGAAGTACCAGTTCAGTCTTTACCCAA_tATTTATGGAAGGAGTGGGA
801 F I P I E K Y Q F S L Y P I F M E G V G

2461 AAACCAAAGATAATTAATAGTTTCACTCAAGATGATATTGAAAAACACCAGAGTGATGCA
821 K P K I I N S F T Q D D I E K H Q S D A

2521 GGTTTATATGTAATTGTGCCAGTAATTATTTCTCTTCCATCTTATTGCTTGGAACATTA
841 G L Y V I V P V I I S S S I L L L G T L

I — — — — — **I**

Figure 21

2581 TTAATATCACACCAAAGAATGAAAAAGCTATTTTGGGAAGATGTTCCGAACCCCAAGAAT
861 L I S H Q R M K K L F W E D V P N P K N

2641 TGTTCTTGGGCACAAGGACTTAATTTTCAGAAGAGAACGGACATTCTTtgaagtctaattc
881 C S W A Q G L N F Q K R T D I L *





2701 atgatcactacagatgaacccaatgtgccaacttcccaacagtctatagagtattagaag
3761 atttttacatttttgaagaaggggagcaaatctaaaaaaaaattcagttgaacttctgagag
2821 ttaacatatggtggattatgttgatttagaacttaaaaatagatgtcatttaaaccaagt
2881 ttacatctaaactcaggtcaaacctacacactaattaaaagtttagtagatttcaaatt
2941 ttcatacataagtaactaaagaccgaaaactaaacagtataaggaccagtattttgtaattc
3001 ttttaataaccgacaacgacagtaatgtatagataatttacagtagtttatacatcatctg
3061 ttaggacattaatccacttgagattttgacgttgtagactgtttatcgaaatttttatgt
3121 tactaatattcataccttagtcacttttataaatcaaacataaaaaatacagggtttgaaaa


Figure 2J


J. — — — — — J


3181 ggtaaaatctaaggaaatatctgtgcagtcggatTTTTtagtcggataagcccacaagaaa
3241 acttatagaggaccgtaaaaaacatagattgaaacaagttagacccttaaagtcaaaagtt
3301 ataggaaactTTTaccgaattcactattgaaggcaaagtcaatTTTccttcgggcttcaac
3361 acaaacacgacgggtgtcctgtcacccctcaatgtcaagtatagtcctactgggatgtatg
3421 ggtccagtcctaactgccctgggtcttcccttgtagctgaagattacaggtgcgaaagaaca
3481 aattaatactggatTTtagattaaatgaagggtgacttggtaggttctggagaccgtccgtc
3541 cctttacccgctcactasgtTTTTTccctctgagaaacctcgaaaatacttatcaagtacc
3601 actcctgtcttgaaaagatgaaagtctgtctgacgaacgatcaaaatacttaag


Figure 2K


**Figure 3A****Figure 3B****Figure 3C****Figure 3D**

**Figure 4**

**Figure 5A**

**Figure 5B**

**Figure 6**



Cross-species conservation of the NR-2 gene

Figure 7





Figure 8



Expression of the Leptin Receptor (NR2) in murine tissues

↑TOP

Figure 9

HAEMOPOIETIN RECEPTOR AND GENETIC SEQUENCES ENCODING SAME

CROSS REFERENCE TO RELATED APPLICATION

This application is a divisional of Ser. No. 09/043,816, filed on Sept. 17, 1998, now U.S. Pat. No. 6,414,128.

BACKGROUND OF THE INVENTION

The preferred haemopoietin receptor of the present invention is referred to herein as "NR2". The NR2 receptor interacts with leptin and is referred to as a "leptin receptor". The terms "haemopoietin receptor", "NR2" and "leptin receptor" are used interchangeably throughout the subject specification. The species from which a particular NR2 is desired is given in single letter abbreviation in lower case before NR2. For example, murine NR2 is "mNR2" and human NR2 is "hNR2". A recombinant form may have the prefix "r".

The rapidly increasing sophistication of recombinant DNA techniques is greatly facilitating research into the medical and allied health fields. Cytokine research is of particular importance, especially as these molecules regulate the proliferation, differentiation and function of a wide variety of cells. Administration of recombinant cytokines or regulating cytokine function and/or synthesis is becoming increasingly the focus of medical research into the treatment of a range of disease conditions.

Despite the discovery of a range of cytokines and other secreted regulators of cell function, comparatively few cytokines are directly used or targeted in therapeutic regimens. One reason for this is the pleiotropic nature of many cytokines. For example, interleukin (IL)-11 is a functionally pleiotropic molecule (1,2), initially characterized by its ability to stimulate proliferation of the IL-6-dependent plasmacytoma cell line, T11 65 (3). Other biological actions of IL-11 include induction of multipotential haemopoietin progenitor cell proliferation (4,5,6), enhancement of megakaryocyte and platelet formation (7,8,9,10), stimulation of acute phase protein synthesis (11) and inhibition of adipocyte lipoprotein lipase activity (12, 13). The diverse and pleiotropic function of IL-11 and other haemopoietin cytokines makes these molecules an important group to study, especially at the level of interaction of the cytokines with their receptors. Manipulation and control of cytokine receptors and of cytokine-receptor interaction is potentially very important in many therapeutic situations, especially where the target cytokine is functionally pleiotropic and it is desired to block certain functions of a target cytokine but not all functions.

Another important aspect of cytokine receptors is in the search for new cytokines. In this regard, the inventors have used a procedure for cloning haemopoietin receptors without prior knowledge of their ligands. Identification of receptors then provides a screening procedure for potentially new cytokines and for previously characterised cytokines. In addition, identification of new haemopoietin receptors allows for selective blocking of pleiotropic cytokine function.

In accordance with the present invention, the inventors identified a novel haemopoietin receptor which interacts with leptin, a hormone which regulates adipose tissue mass.

BRIEF SUMMARY OF THE INVENTION

The present invention is directed to a novel haemopoietin receptor or a derivative thereof and to genetic sequences

encoding same. The receptor molecule and its derivatives and the genetic sequences encoding same of the present invention are useful in the development of a wide range of agonists, antagonists, therapeutics and diagnostic reagents based on ligand interaction with its receptor. The present invention particularly relates to a receptor for leptin.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic representation showing size of NR2 cDNA clones isolated and schematic structure of the predicted NR2 protein.

FIG. 2 is a representation of the nucleotide sequence and corresponding amino acid sequence of the haemopoietin receptor.

FIG. 3 is a representation of a FACS analysis of NR2 expression by BA/F₃ cells.

FIG. 4 is a photographic representation of a silver-stained gel of NR2 expression by BA/F₃ cells.

FIG. 5 is a graphical representation showing specific binding of ¹²⁵I human leptin to Ba/F₃ cells stably transfected to express hNR2 on their cell surface.

(a) Saturation binding curve for ¹²⁵I h leptin binding to Ba/F₃/hNR2 cells at 23° C.

(b) Scatchard transformation of the data in (a). The slope of the curve indicates an equilibrium dissociation constant (K_D) of 120 pM.

FIG. 6 is a graphical representation showing specific binding of ¹²⁵I human leptin to COS-7 cells transiently transfected to express hNR2 on their cell surface (a) or to purified soluble human NR2 (b). Saturation binding curves at 23° C. are shown.

FIG. 7 is a photographic representation showing cross species conservation of the NR2 gene. Southern blot of genomic DNA probed with a specific cDNA probe for NR2.

FIG. 8 is a diagrammatic representation of the NR2 locus. A map of the NR2 locus, showing positioning of the clones isolated from genomic libraries. The results of the restriction enzyme mapping using NcoI and the positioning of the exons on these fragments are also shown.

FIG. 9 is a photographic representation showing expression of leptin receptor (NR2) in murine tissues.

DETAILED DESCRIPTION OF THE INVENTION

Bibliographic details of the publications numerically referred to in this specification are collected at the end of the description. Sequence Identity Numbers (SEQ ID NOs.) for the nucleotide and amino acid sequences referred to in the specification are defined following the bibliography.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

One aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a haemopoietin receptor or a derivative thereof wherein said sequence of nucleotides or a complementary form thereof is capable of hybridising under medium stringent conditions to the oligonucleotide:

5'-(A/G)CTCCA(A/G)TC(A/G)CTCCA-3' [SEQ ID NO: 1].

In a preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence or a complementary form thereof which hybridises under medium stringent conditions to the oligonucleotides:

5'-(A/G)CTCCA(A/G)TC(A/G)CTCCA-3' [SEQ ID NO: 1]

5'-ACTAGCAGGGATGTAGCTGAG-3' [SEQ ID NO:4]

5'-CTGCTCCTATGATACCT-3' [SEQ ID NO:6]

5'-CCTCTTCCATCTTATTGCTTGG-3' [SEQ ID NO:7]

5'-ATCGGTCGTGACATACAAGG-3' [SEQ ID NO:8]

In an even more preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence or a complementary form thereof which hybridises under medium stringent conditions to one or more of the following oligonucleotides:

5'-(A/G)CTCCA(A/G)TC(A/G)CTCCA-3' [SEQ ID NO: 1].

5'-ACTAGCAGGGATGTAGCTGAG-3' [SEQ ID NO:4]

5'-CTCAGCTACATCCCTGCTAGT-3' [SEQ ID NO:5]

5'-CTGCTCCTATGATACCT-3' [SEQ ID NO:6]

5'-CCTCTTCCATCTTATTGCTTGG-3' [SEQ ID NO:7]

5'-ATCGGTCGTGACATACAAGG-3' [SEQ ID NO:8]

5'-AGCTAAGCTTTCTAGATATCCAATTACTCCTTGGAGAT-3' [SEQ ID NO:9]

5'-AGCTTCTAGATCAATCACTCTGGTGTGTTTTCAAT-3' [SEQ ID NO:10]

5'-AGCTTCTAGATCAAACCTTTTATATCCATGACAAC-3' [SEQ ID NO:11].

In a still more preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence or complementary form thereof which is capable of hybridising separately under medium stringent conditions to each of oligonucleotide SEQ ID NO:1 and SEQ ID NO:4 to SEQ ID NO:11.

In a most preferred embodiment, the present invention provides a nucleic acid molecule comprising a sequence of nucleotides or a complementary form thereof substantially as set forth in FIG. 2 [SEQ ID NO:12] or a sequence of nucleotides capable of hybridising to all or part thereof under medium stringent conditions.

Accordingly, a preferred embodiment of the present invention is also directed to a nucleic acid molecule encoding a haemopoietin receptor or a derivative thereof and comprising a nucleotide sequence as set forth in SEQ ID NO:12 or is capable of hybridising to all or part thereof under medium stringent conditions.

The haemopoietin receptor of the present invention is referred to herein as "NR2". In accordance with the present invention, NR2 is capable of interacting with leptin and, hence, is also referred to as a "leptin receptor".

The term "derivative" includes any or all parts, fragments, portions, homologues or analogues to the nucleotide sequence set forth in SEQ ID NO:12 as well as hybrid molecules between the NR2 receptor and other receptors or other molecules. Derivatives include single or multiple nucleotide substitutions, deletions and/or additions to the nucleotide sequence set forth in SEQ ID NO:12.

Another aspect of the present invention contemplates a recombinant haemopoietin receptor encoded by the nucleic acid molecules as hereinbefore described.

According to one aspect of this embodiment, there is provided recombinant haemopoietin receptor encoded by a nucleic acid molecule which comprises a nucleotide sequence or a complementary form thereof which is capable of hybridising to SEQ ID NO:1 under medium stringent conditions.

In a preferred embodiment, the recombinant haemopoietin receptor is encoded by a nucleic acid molecule which comprises a nucleotide sequence or a complementary form thereof which is capable of hybridising to SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7 and SEQ ID NO:8 under medium stringent conditions.

In an even more preferred embodiment, the recombinant haemopoietin receptor is encoded by a nucleic acid molecule which comprises a nucleotide sequence or complementary form thereof which hybridises under medium stringency conditions to one or more of SEQ ID NO:1 and SEQ ID NO:4 to SEQ ID NO:11.

In still an even more preferred embodiment, the recombinant haemopoietin receptor is encoded by a nucleic acid molecule which comprises a nucleotide sequence or a complementary form thereof which hybridises under medium stringent conditions to each of oligonucleotides SEQ ID NO:1 and SEQ ID NO:4 to SEQ ID NO:11.

In a most preferred embodiment, the present invention is directed to a recombinant NR2 encoded by a nucleic acid molecule comprising a nucleotide sequence or complementary form thereof substantially as set forth in SEQ ID NO:12 or a sequence capable of hybridising to all or part thereof under medium stringent conditions.

According to this latter aspect of the present invention, there is provided a recombinant NR2 having an amino acid sequence substantially as set forth in FIG. 2 [SEQ ID NO:13] or having at least about 60% similarity to all or part thereof, more preferably at least about 70%, still more preferably at least about 80% and still more preferably at least about 90–95% or above (e.g. 96%, 97%, 98% or greater than or equal to 99%) similarly to all or part of the amino acid sequence set forth in SEQ ID NO:13.

The recombinant NR2 or a genetic sequence encoding same is preferably in isolated form meaning that a composition of matter comprises at least about 10%, more preferably at least about 20%, still more preferably at least about 30–40%, even more preferably at least about 50–60%, still even more preferably at least about 70–80% or greater (e.g. 85%, 90% or 95%) of the recombinant receptor or genetic sequence encoding same relative to other components in the composition as determined by, for example, molecular weight, activity, nucleic acid content or composition or other convenient means.

Reference herein to "recombinant haemopoietin receptor", "NR2" or "leptin receptor" includes reference to derivatives thereof such as parts, fragments, portions, homologues, hybrids or analogues thereof. The derivatives may be functional or not or may be non-functional but immunologically interactive with antibodies to all or part of the receptor. Derivatives of the receptor also cover agonists or antagonists of receptor-ligand interaction. Function is conveniently defined by an ability of NR2 to interact with leptin or for soluble NR2 to compete with leptin-induced activities of certain cells.

For the purposes of defining the level of stringency, reference can conveniently be made to Sambrook et al (14) which is herein incorporated by reference where the washing steps disclosed at pages 952–957 are considered high stringency. A low stringency is defined herein as being in 4–6×SSC/0.1–0.5% w/v SDS at 37–45° C. for 2–3 hours. Depending on the source and concentration of nucleic acid involved in the hybridisation, alternative conditions of stringency may be employed such as medium stringent conditions which are considered herein to be 1–4×SSC/0.25–0.5% w/v SDS at ≥45° C. for 2–3 hours or high stringent conditions considered herein to be 0.1–1×SSC/0.1% w/v SDS at ≥60° C. for 1–3 hours.

The nucleic acid molecule is preferably derivable from the human genome but genomes and nucleotide sequences from non-human animals are also encompassed by the present invention. Non-human animals contemplated by the present invention include livestock animals (e.g. sheep, cows, pigs, goats, horses, donkeys), laboratory test animals (e.g. mice, rats, guinea pigs, hamsters, rabbits), domestic companion animals (e.g. dogs, cats), birds (e.g. chickens, geese, ducks and other poultry birds, game birds, emus, ostriches) and captive wild or tamed animals (e.g. foxes, kangaroos, dingoes).

Preferred human genetic sequences encoding NR2 include sequences from cells of bone marrow, brain, liver, kidney, heart, testis, stomach, lymph nodes, colon, spleen and ovary, neonatal tissue, embryonic tissue, cancer or tumour-derived tissues.

The nucleic acid molecule of the present invention may be single or double stranded, linear or closed circle DNA (e.g. genomic DNA), cDNA or mRNA or combinations thereof such as in the form of DNA:RNA hybrids. The nucleic acid molecule may also include a vector such as an expression vector component to facilitate expression of the haemopoietin receptor or its components or parts.

As stated above, the present invention further contemplates a range of derivatives of NR2. Derivatives include fragments, parts, portions, mutants, homologues and analogues of the NR2 polypeptide and corresponding genetic sequence. Derivatives also include single or multiple amino acid substitutions, deletions and/or additions to NR2 or single or multiple nucleotide substitutions, deletions and/or additions to the genetic sequence encoding NR2. "Additions" to amino acid sequences or nucleotide sequences include fusions with other peptides, polypeptides or proteins or fusions to nucleotide sequences. Reference herein to "NR2" includes reference to all derivatives thereof including functional derivatives or "NR2" immunologically interactive derivatives.

Analogues of NR2 contemplated herein include, but are not limited to, modification to side chains, incorporating of unnatural amino acids and/or their derivatives during peptide, polypeptide or protein synthesis and the use of crosslinkers and other methods which impose conformational constraints on the proteinaceous molecule or their analogues.

Examples of side chain modifications contemplated by the present invention include modifications of amino groups such as by reductive alkylation by reaction with an aldehyde followed by reduction with NaBH_4 ; amidination with methylacetimidate; acylation with acetic anhydride; carbamoylation of amino groups with cyanate; trinitrobenzylation of amino groups with 2, 4, 6-trinitrobenzene sulphonic acid (TNBS); acylation of amino groups with succinic anhydride and tetrahydrophthalic anhydride; and pyridoxylation of lysine with pyridoxal-5-phosphate followed by reduction with NaBH_4 .

The guanidine group of arginine residues may be modified by the formation of heterocyclic condensation products with reagents such as 2,3-butanedione, phenylglyoxal and glyoxal.

The carboxyl group may be modified by carbodiimide activation via O-acylisourea formation followed by subsequent derivatisation, for example, to a corresponding amide.

Sulphydryl groups may be modified by methods such as carboxymethylation with iodoacetic acid or iodoacetamide; performic acid oxidation to cysteic acid; formation of a mixed disulphides with other thiol compounds; reaction with maleimide, maleic anhydride or other substituted male-

imide; formation of mercurial derivatives using 4-chloromercuribenzoate, 4-chloromercuriphenylsulphonic acid, phenylmercury chloride, 2-chloromercuri-4-nitrophenol and other mercurials; carbamoylation with cyanate at alkaline pH.

Tryptophan residues may be modified by, for example, oxidation with N-bromosuccinimide or alkylation of the indole ring with 2-hydroxy-5-nitrobenzyl bromide or sulphenyl halides. Tyrosine residues on the other hand, may be altered by nitration with tetranitromethane to form a 3-nitrotyrosine derivative.

Modification of the imidazole ring of a histidine residue may be accomplished by alkylation with iodoacetic acid derivatives or N-carbethoxylation with diethylpyrocarbonate.

Examples of incorporating unnatural amino acids and derivatives during peptide synthesis include, but are not limited to, use of norleucine, 4-amino butyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, ornithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid, 2-thienyl alanine and/or D-isomers of amino acids. A list of unnatural amino acid, contemplated herein is shown in Table 1.

Crosslinkers can be used, for example, to stabilise 3D conformations, using homo-bifunctional crosslinkers such as the bifunctional imido esters having $(\text{CH}_2)_n$ spacer groups with $n=1$ to $n=6$, glutaraldehyde, N-hydroxysuccinimide esters and hetero-bifunctional reagents which usually contain an amino-reactive moiety such as N-hydroxysuccinimide and another group specific-reactive moiety such as maleimido or dithio moiety (SH) or carbodiimide (COOH). In addition, peptides can be conformationally constrained by, for example, incorporation of C_α and N_α -methylamino acids, introduction of double bonds between C_α and C_β atoms of amino acids and the formation of cyclic peptides or analogues by introducing covalent bonds such as forming an amide bond between the N and C termini, between two side chains or between a side chain and the N or C terminus.

These types of modifications may be important to stabilise NR2 if administered to an individual or for use as a diagnostic reagent.

The present invention further contemplates chemical analogues of NR2 capable of acting as antagonists or agonists of NR2 or which can act as functional analogues of NR2. Chemical analogues may not necessarily be derived from NR2 but may share certain conformational similarities. Alternatively, chemical analogues may be specifically designed to mimic certain physiochemical properties of NR2. Chemical analogues may be chemically synthesised or may be detected following, for example, natural product screening.

The identification of NR2 permits the generation of a range of therapeutic molecules capable of modulating expression of NR2 or modulating the activity of NR2. Modulators contemplated by the present invention includes agonists and antagonists of NR2 expression. Antagonists of NR2 expression include antisense molecules, ribozymes and co-suppression molecules. Agonists include molecules which increase promoter ability or interfere with negative regulatory mechanisms. Agonists of NR2 include molecules which overcome any negative regulatory mechanism. Antagonists of NR2 include antibodies and inhibitor peptide fragments.

TABLE 1

Non-conventional amino acid	Code	Non-conventional amino acid	Code
α -aminobutyric acid	Abu	L-N-methylalanine	Nmala
α -amino- α -methylbutyrate	Mgab	L-N-methylarginine	Nmarg
aminocyclopropane-carboxylate	Cpro	L-N-methylasparagine	Nmasn
aminoisobutyric acid	Aib	L-N-methylaspartic acid	Nmasp
aminonorbornyl-carboxylate	Norb	L-N-methylcysteine	Nmcys
cyclohexylalanine	Chexa	L-N-methylglutamine	Nmgln
cyclopentylalanine	Cpen	L-N-methylglutamic acid	Nmglu
D-alanine	Dal	L-N-methylhistidine	Nmhis
D-arginine	Darg	L-N-methylisoleucine	Nmile
D-aspartic acid	Dasp	L-N-methylleucine	Nmleu
D-cysteine	Dcys	L-N-methyllysine	Nmlys
D-glutamine	Dgln	L-N-methylmethionine	Nmmet
D-glutamic acid	Dglu	L-N-methylnorleucine	Nnmle
D-histidine	Dhis	L-N-methylnorvaline	Nnmva
D-isoleucine	Dile	L-N-methylornithine	Nmorn
D-leucine	Dleu	L-N-methylphenylalanine	Nmphe
D-lysine	Dlys	L-N-methylproline	Nmpro
D-methionine	Dmet	L-N-methylserine	Nmser
D-ornithine	Dorn	L-N-methylthreonine	Nmthr
D-phenylalanine	Dphe	L-N-methyltryptophan	Nmtrp
D-proline	Dpro	L-N-methyltyrosine	Nmtyr
D-serine	Dser	L-N-methylvaline	Nmval
D-threonine	Dthr	L-N-methylethylglycine	Nmetg
D-tryptophan	Dtrp	L-N-methyl-t-butylglycine	Nmtbug
D-tyrosine	Dtyr	L-norleucine	Nle
D-valine	Dval	L-norvaline	Nva
D- α -methylalanine	Dmala	α -methyl-aminoisobutyrate	Maib
D- α -methylarginine	Dmarg	α -methyl- γ -amino-butyrate	Mgab
D- α -methylasparagine	Dmasn	α -methylcyclohexylalanine	Mchexa
D- α -methylaspartate	Dmasp	α -methylcyclopentylalanine	Mcpen
D- α -methylcysteine	Dmcys	α -methyl- α -naphthylalanine	Manap
D- α -methylglutamine	Dmgln	α -methylpenicillamine	Mpen
D- α -methylhistidine	Dmhis	N-(4-aminobutyl)glycine	Nglu
D- α -methylisoleucine	Dmile	N-(2-aminoethyl)glycine	Nacg
D- α -methylleucine	Dmleu	N-(3-aminopropyl)glycine	Norn
D- α -methyllysine	Dmlys	N-amino- α -methylbutyrate	Nmaabu
D- α -methylmethionine	Dmmet	α -naphthylalanine	Anap
D- α -methylornithine	Dmorn	N-benzylglycine	Nphe
D- α -methylphenylalanine	Dmphe	N-(2-carbamylethyl)glycine	Ngln
D- α -methylproline	Dmpro	N-(carbamylmethyl)glycine	Nasn
D- α -methylserine	Dmser	N-(2-carboxyethyl)glycine	Nglu
D- α -methylthreonine	Dmthr	N-(carboxymethyl)glycine	Nasp
D- α -methyltryptophan	Dmtrp	N-cyclobutylglycine	Ncbut
D- α -methyltyrosine	Dmtyr	N-cycloheptylglycine	Nchep
D- α -methylvaline	Dmval	N-cyclohexylglycine	Nchex
D-N-methylalanine	Dnmala	N-cyclodecylglycine	Ncdcc
D-N-methylarginine	Dnmarg	N-cyclododecylglycine	Ncdod
D-N-methylasparagine	Dnmasn	N-cyclooctylglycine	Ncoct
D-N-methylaspartate	Dnmasp	N-cyclopropylglycine	Ncprio
D-N-methylcysteine	Dnmcys	N-cycloundecylglycine	Ncund
D-N-methylglutamine	Dnmgln	N-(2,2-diphenylethyl)glycine	Nbhm
D-N-methylglutamate	Dnmglu	N-(3,3-diphenylpropyl)glycine	Nbhe
D-N-methylhistidine	Dnmhis	N-(3-guanidinopropyl)glycine	Narg
D-N-methylisoleucine	Dnmile	N-(1-hydroxyethyl)glycine	Nthr
D-N-methylleucine	Dnmleu	N-(hydroxyethyl)glycine	Nser
D-N-methyllysine	Dnmlys	N-(imidazolethyl)glycine	Nhis
N-methylcyclohexylalanine	Nmchexa	N-(3-indolylethyl)glycine	Nhtrp
D-N-methylornithine	Dnmorn	N-methyl- γ -aminobuty-	Nmgabu
N-methylglycine	Nala		
N-methylaminoisobutyrate	Nmaib		
N-(1-methylpropyl)glycine	Nile		
N-(2-methylpropyl)glycine	Nleu		
D-N-methyltryptophan	Dnmtrp		
D-N-methyltyrosine	Dnmtyr		
D-N-methylvaline	Dnmval		
γ -aminobutyric acid	Gabu		
L-t-butylglycine	Tbug		
L-ethylglycine	Etg		

TABLE 1-continued

Non-conventional amino acid	Code	Non-conventional amino acid	Code
L-homophenylalanine	Hphe	rate	
L- α -methylarginine	Marg	D-N-methylmethionine	Dnmmet
L- α -methylaspartate	Masp	N-methylcyclopentylalanine	Nmcpen
L- α -methylcysteine	Mcys	D-N-methylphenylalanine	Dnmphe
L- α -methylglutamine	Mgln	D-N-methylproline	Dnmpro
L- α -methylhistidine	Mhis	D-N-methylserine	Dnmser
L- α -methylisoleucine	Mile	D-N-methylthreonine	Dnmthr
L- α -methylleucine	Mleu	N-(1-methylethyl)glycine	Nval
L- α -methylmethionine	Mmet	N-methyl-naphthylalanine	Nmanap
L- α -methylnorvaline	Mnva	N-methylpenicillamine	Nmpen
L- α -methylphenylalanine	Mphe	N-(p-hydroxyphenyl)glycine	Nhtyr
L- α -methylserine	Mser	N-(thiomethyl)glycine	Ncys
L- α -methyltryptophan	Mtrp	penicillamine	Pen
L- α -methylvaline	Mval	L- α -methylalanine	Mala
N-(N-(2,2-diphenylethyl)-carbamylmethyl)glycine	Nnbhm	L- α -methylasparagine	Masn
1-carboxy-1-(2,2-diphenylethylamino)-cyclopropane	Nmbc	L- α -methyl-t-butylglycine	Mtbug
		L-methylethylglycine	Metg
		L- α -methylglutamate	Mglu
		L- α -methylhomophenylalanine	Mhphe
		N-(2-methylthioethyl)glycine	Nmet
		L- α -methyllysine	Mlys
		L- α -methylnorleucine	Mnle
		L- α -methylornithine	Morn
		L- α -methylproline	Mpro
		L- α -methylthreonine	Mthr
		L- α -methyltyrosine	Mtyr
		L-N-methylhomophenylalanine	Nmhph
		N-(N-(3,3-diphenylpropyl)-carbamylmethyl)glycine	Nnbhe

Other derivatives contemplated by the present invention include a range of glycosylation variants from a completely unglycosylated molecule to a modified glycosylated molecule. Altered glycosylation patterns may result from expression of recombinant molecules in different host cells.

Another embodiment of the present invention contemplates a method for modulating expression of NR2 in a human, said method comprising contacting the NR2 gene encoding NR2 with an effective amount of a modulator of NR2 expression for a time and under conditions sufficient to up-regulate or down-regulate or otherwise modulate expression of NR2. For example, a nucleic acid molecule encoding NR2 or a derivative thereof may be introduced into a cell to enhance NR2 related activities of that cell. Conversely, NR2 antisense sequences (or sense sequences for co-suppression) such as oligonucleotides may be introduced to decrease NR2-related activities of any cell expressing the endogenous NR2 gene. Ribozymes may also be used.

Another aspect of the present invention contemplates a method of modulating activity of NR2 in a human, said method comprising administering to said mammal a modulating effective amount of a molecule for a time and under conditions sufficient to increase or decrease NR2 activity. The molecule may be a proteinaceous molecule or a chemical entity and may also be a derivative of NR2 or its receptor or a chemical analogue or truncation mutant of NR2 or its receptor.

Accordingly, the present invention contemplates a pharmaceutical composition comprising NR2 or a derivative thereof or a modulator of NR2 expression or NR2 activity

and one or more pharmaceutically acceptable carriers and/or diluents. These components are referred to as the "active ingredients".

In this regard there is provided a pharmaceutical composition comprising a recombinant haemopoietin receptor as hereinbefore described or a ligand (e.g. leptin) binding portion thereof and one or more pharmaceutically acceptable carriers and/or diluents.

In another embodiment, there is provided a pharmaceutical composition comprising a ligand (e.g. leptin) to the recombinant haemopoietin receptor as hereinbefore described and one or more pharmaceutically acceptable carriers and/or diluents.

Still a further aspect of the present invention contemplates a method of treatment of an animal comprising administering to said animal a treatment effective amount of a recombinant haemopoietin receptor as hereinbefore described or a ligand binding portion thereof or a ligand (e.g. leptin) to said haemopoietic receptor for a time and under conditions sufficient for said treatment to be substantially effected or the conditions to be substantially ameliorated.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion or may be in the form of a cream or other form suitable for topical application. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

When the active ingredients are suitably protected they may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsule, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets,

troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 1% by weight of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 5 to about 80% of the weight of the unit. The amount of active compound in such therapeutically useful compositions in such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 0.1 μ g and 2000 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the components as listed hereafter: A binder such as gum, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound(s) may be incorporated into sustained-release preparations and formulations.

The present invention also extends to forms suitable for topical application such as creams, lotions and gels.

Pharmaceutically acceptable carriers and/or diluents include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active material for the treatment of disease in living subjects having a diseased condition in which bodily health is impaired.

The principal active ingredient is compounded for convenient and effective administration in effective amounts with a suitable pharmaceutically acceptable carrier in dosage unit form as hereinbefore disclosed. A unit dosage form can, for example, contain the principal active compound in amounts ranging from 0.5 μ g to about 2000 mg. Expressed in proportions, the active compound is generally present in from about 0.5 μ g to about 2000 mg/ml of carrier. In the case

of compositions containing supplementary active ingredients, the dosages are determined by reference to the usual dose and manner of administration of the said ingredients.

The pharmaceutical composition may also comprise genetic molecules such as a vector capable of transfecting target cells where the vector carries a nucleic acid molecule capable of modulating NR2 expression or NR2 activity. The vector may, for example, be a viral vector.

Still another aspect of the present invention is directed to antibodies to NR2 and its derivatives or its ligands (e.g. leptin). Such antibodies may be monoclonal or polyclonal and may be selected from naturally occurring antibodies to NR2 or may be specifically raised to NR2 or derivatives thereof. In the case of the latter, NR2 or its derivatives may first need to be associated with a carrier molecule. The antibodies and/or recombinant NR2 or its derivatives of the present invention are particularly useful as therapeutic or diagnostic agents.

For example, NR2 and its derivatives can be used to screen for naturally occurring antibodies to NR2. These may occur, for example in some autoimmune diseases. Alternatively, specific antibodies can be used to screen for NR2. Techniques for such assays are well known in the art and include, for example, sandwich assays and ELISA. Knowledge of NR2 levels may be important for diagnosis of certain cancers or a predisposition to cancers or for monitoring certain therapeutic protocols.

Antibodies to NR2 of the present invention may be monoclonal or polyclonal. Alternatively, fragments of antibodies may be used such as Fab fragments. Furthermore, the present invention extends to recombinant and synthetic antibodies and to antibody hybrids. A "synthetic antibody" is considered herein to include fragments and hybrids of antibodies. The antibodies of this aspect of the present invention are particularly useful for immunotherapy and may also be used as a diagnostic tool for assessing the receptor or receptor-ligand interaction or monitoring the program of a therapeutic regimen.

For example, specific antibodies can be used to screen for NR2 proteins. The latter would be important, for example, as a means for screening for levels of NR2 in a cell extract or other biological fluid or purifying NR2 made by recombinant means from culture supernatant fluid. Techniques for the assays contemplated herein are known in the art and include, for example, sandwich assays and ELISA.

It is within the scope of this invention to include any second antibodies (monoclonal, polyclonal or fragments of antibodies or synthetic antibodies) directed to the first mentioned antibodies discussed above. Both the first and second antibodies may be used in detection assays or a first antibody may be used with a commercially available anti-immunoglobulin antibody. An antibody as contemplated herein includes any antibody specific to any region of NR2.

Both polyclonal and monoclonal antibodies are obtainable by immunization with the enzyme or protein and either type is utilizable for immunoassays. The methods of obtaining both types of sera are well known in the art. Polyclonal sera are less preferred but are relatively easily prepared by injection of a suitable laboratory animal with an effective amount of NR2, or antigenic parts thereof, collecting serum from the animal, and isolating specific sera by any of the known immunoadsorbent techniques. Although antibodies produced by this method are utilizable in virtually any type of immunoassay, they are generally less favoured because of the potential heterogeneity of the product.

The use of monoclonal antibodies in an immunoassay is particularly preferred because of the ability to produce them

in large quantities and the homogeneity of the product. The preparation of hybridoma cell lines for monoclonal antibody production derived by fusing an immortal cell line and lymphocytes sensitized against the immunogenic preparation can be done by techniques which are well known to those who are skilled in the art.

Another aspect of the present invention contemplates a method for detecting NR2 in a biological sample from a subject said method comprising contacting said biological sample with an antibody specific for NR2 or its derivatives or homologues for a time and under conditions sufficient for an antibody-NR2 complex to form, and then detecting said complex.

The presence of NR2 may be accomplished in a number of ways such as by Western blotting and ELISA procedures. A wide range of immunoassay techniques are available as can be seen by reference to U.S. Pat. Nos. 4,016,043, 4,424,279 and 4,018,653. These, of course, includes both single-site and two-site or "sandwich" assays of the non-competitive types, as well as in the traditional competitive binding assays. These assays also include direct binding of a labelled antibody to a target.

Sandwich assays are among the most useful and commonly used assays and are favoured for use in the present invention. A number of variations of the sandwich assay technique exist, and all are intended to be encompassed by the present invention. Briefly, in a typical forward assay, an unlabelled antibody is immobilized on a solid substrate and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen complex, a second antibody specific to the antigen, labelled with a reporter molecule capable of producing a detectable signal is then added and incubated, allowing time sufficient for the formation of another complex of antibody-antigen-labelled antibody. Any unreacted material is washed away, and the presence of the antigen is determined by observation of a signal produced by the reporter molecule. The results may either be qualitative, by simple observation of the visible signal, or may be quantitated by comparing with a control sample containing known amounts of hapten. Variations on the forward assay include a simultaneous assay, in which both sample and labelled antibody are added simultaneously to the bound antibody. These techniques are well known to those skilled in the art, including any minor variations as will be readily apparent. In accordance with the present invention the sample is one which might contain NR2 including cell extract, tissue biopsy or possibly serum, saliva, mucosal secretions, lymph, tissue fluid and respiratory fluid. The sample is, therefore, generally a biological sample comprising biological fluid, cell extract, bone marrow or lymph, tissue extract (e.g. from kidney, liver, spleen, etc), fermentation fluid and supernatant fluid such as from a cell culture and cell conditioned medium.

In the typical forward sandwich assay, a first antibody having specificity for the NR2 or antigenic parts thereof, is either covalently or passively bound to a solid surface. The solid surface is typically glass or a polymer, the most commonly used polymers being cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene. The solid supports may be in the form of tubes, beads, discs of microplates, or any other surface suitable for conducting an immunoassay. The binding processes are well-known in the art and generally consist of cross-linking covalently binding or physically adsorbing, the polymer-antibody complex is washed in preparation for the test sample. An aliquot of the sample to be tested is then added to the solid phase

complex and incubated for a period of time sufficient (e.g. 2–40 minutes) and under suitable conditions (e.g. 25° C.) to allow binding of any subunit present in the antibody. Following the incubation period, the antibody subunit solid phase is washed and dried and incubated with a second antibody specific for a portion of the hapten. The second antibody is linked to a reporter molecule which is used to indicate the binding of the second antibody to the hapten.

An alternative method involves immobilizing the target molecules in the biological sample and then exposing the immobilized target to specific antibody which may or may not be labelled with a reporter molecule. Depending on the amount of target and the strength of the reporter molecule signal, a bound target may be detectable by direct labelling with the antibody. Alternatively, a second labelled antibody, specific to the first antibody is exposed to the target-first antibody complex to form a target-first antibody-second antibody tertiary complex. The complex is detected by the signal emitted by the reporter molecule.

By “reporter molecule” as used in the present specification, is meant a molecule which, by its chemical nature, provides an analytically identifiable signal which allows the detection of antigen-bound antibody. Detection may be either qualitative or quantitative. The most commonly used reporter molecules in this type of assay are either enzymes, fluorophores or radionuclide containing molecules (i.e. radioisotopes) and chemiluminescent molecules. In the case of an enzyme immunoassay, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. As will be readily recognized, however, a wide variety of different conjugation techniques exist, which are readily available to the skilled artisan. Commonly used enzymes include horseradish peroxidase, glucose oxidase, beta-galactosidase and alkaline phosphatase, amongst others. The substrates to be used with the specific enzymes are generally chosen for the production, upon hydrolysis by the corresponding enzyme, of a detectable colour change. Examples of suitable enzymes include alkaline phosphatase and peroxidase. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates noted above. In all cases, the enzyme-labelled antibody is added to the first antibody hapten complex, allowed to bind, and then the excess reagent is washed away. A solution containing the appropriate substrate is then added to the complex of antibody-antigen-antibody. The substrate will react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of hapten which was present in the sample. “Reporter molecule” also extends to use of cell agglutination or inhibition of agglutination such as red blood cells on latex beads, and the like.

Alternately, fluorescent compounds, such as fluorescein and rhodamine, may be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labelled antibody adsorbs the light energy, inducing a state to excitability in the molecule, followed by emission of the light at a characteristic colour visually detectable with a light microscope. As in the EIA, the fluorescent labelled antibody is allowed to bind to the first antibody-hapten complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to the light of the appropriate wavelength the fluorescence observed indicates the presence of the hapten of interest. Immunofluorescence and EIA techniques are both very well established in the art and are particularly preferred for the present method. However, other reporter molecules, such as radioisotope, chemiluminescent or bioluminescent molecules, may also be employed.

The present invention also contemplates genetic assays such as involving PCR analysis to detect NR2 gene or its derivatives. Alternative methods or methods used in conjunction include direct nucleotide sequencing or mutation scanning such as single stranded conformation polymorphisms analysis (SSCP) as specific oligonucleotide hybridisation, as methods such as direct protein truncation tests. Such genetic tests may be important, for example, in genetic screening of animals (e.g. humans) for non-expression or substantial absence of expression or expression of mutant forms of NR2 leading to conditions such as obesity or other effects of leptin-receptor interaction.

The nucleic acid molecules of the present invention may be DNA or RNA. When the nucleic acid molecule is in DNA form, it may be genomic DNA or cDNA. RNA forms of the nucleic acid molecules of the present invention are generally mRNA.

Although the nucleic acid molecules of the present invention are generally in isolated form, they may be integrated into or ligated to or otherwise fused or associated with other genetic molecules such as vector molecules and in particular expression vector molecules. Vectors and expression vectors are generally capable of replication and, if applicable, expression in one or both of a prokaryotic cell or a eukaryotic cell. Preferably, prokaryotic cells include *E. coli*, *Bacillus* sp and *Pseudomonas* sp. Preferred eukaryotic cells include yeast, fungal, mammalian and insect cells.

Accordingly, another aspect of the present invention contemplates a genetic construct comprising a vector portion and a mammalian and more particularly a human NR2 gene portion, which NR2 gene portion is capable of encoding an NR2 polypeptide or a functional or immunologically interactive derivative thereof.

Preferably, the NR2 gene portion of the genetic construct is operably linked to a promoter on the vector such that said promoter is capable of directing expression of said NR2 gene portion in an appropriate cell.

In addition, the NR2 gene portion of the genetic construct may comprise all or part of the gene fused to another genetic sequence such as a nucleotide sequence encoding glutathione-S-transferase or part thereof or a cytokine or another haemopoietic receptor. Hybrid receptor molecules are particularly useful in the development of multi functional therapeutic and diagnostic agents.

The present invention extends to such genetic constructs and to prokaryotic or eukaryotic cells comprising same.

The present invention also extends to any or all derivatives of NR2 including mutants, part, fragments, portions, homologues and analogues or their encoding genetic sequence including single or multiple nucleotide or amino acid substitutions, additions and/or deletions to the naturally occurring nucleotide or amino acid sequence.

The NR2 and its genetic sequence of the present invention will be useful in the generation of a range of therapeutic and diagnostic reagents and will be especially useful in the detection of a corresponding ligand. For example, recombinant NR2 may be bound or fused to a reporter molecule capable of producing an identifiable signal, contacted with a biological sample putatively containing a ligand and screening for binding of the labelled NR2 to the ligand. Alternatively, labelled NR2 may be used to screen expression libraries of putative ligand genes or functional parts thereof.

In another embodiment, the NR2 is first immobilised. According to this embodiment, there is provided a method comprising contacting a biological sample containing a putative ligand with said haemopoietic receptor or a ligand binding portion thereof immobilised to a solid support for a time and under conditions sufficient for a complex to form between said receptor and said ligand if said ligand is present in said biological sample, eluting bound ligand and isolating same.

Soluble NR2 polypeptides are also contemplated to be useful in the treatment of disease, injury or abnormality in the nervous system, e.g. in relation to central or peripheral nervous system to treat Cerebral Palsy, trauma induced paralysis, vascular ischaemia associated with stroke, neuronal tumours, motoneurone disease, Parkinson's disease, Huntington's disease, Alzheimer's disease, Multiple Sclerosis, peripheral neuropathies associated with diabetes, heavy metal or alcohol toxicity, renal failure and infectious diseases such as herpes, rubella, measles, chicken pox, HIV

-continued

Amino Acid	Three-letter Abbreviation	One-letter Symbol
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V
Any residue	Xaa	X

TABLE 2

SEQUENCE OF OLIGONUCLEOTIDES		
OLIGONUCLEOTIDE	SEQUENCE	SEQ ID NOs
HYB2	5'-(A/G)CTCCA(A/G)TC(A/G)CTCCA-3'	1
T3	5'-TAATACGACTCACTATAGGGAGA-3'	2
T7	5'-ATTAACCTCACTAAAGGGA-3'	3
721	5'-ACTAGCAGGGATGTAGCTGAG-3'	4
722	5'-CTCAGCTACATCCCTGCTAGT-3'	5
761	5'-CTGCTCTATGATACCT-3'	6
875	5'-CCTCTTCCATCTTATTGCTTGG-3'	7
939	5'-ATCGGTGCTGACATACAAGG-3'	8
1056	5'-AGCTAAGCTTTCTAGATATCCAATTACTCCTTGGAGA-3'	9
1092	5'-AGCTTCTAGATCAATCACTCTGGTGTCTTTTCAAT-3'	10
1094	5'-AGCTTCTAGATCAAACCTTTATATCCATGACAAAC-3'	11

or HTLV-1. The NR2 polypeptides may also be important for regulating cytokine activity such as leptin activity, modulating haemopoiesis and/or regulating or modulating adipose tissue.

As stated above, the NR2 or its ligand of the present invention or their functional derivatives may be provided in a pharmaceutical composition together with one or more pharmaceutically acceptable carriers and/or diluents. In addition, the present invention contemplates a method of treatment comprising the administration of an effective amount of NR2 of the present invention. The present invention also extends to antagonists and agonists of NR2 and/or its ligand and their use in therapeutic compositions and methodologies.

A further aspect of the present invention contemplates the use of NR2 or its functional derivatives in the manufacture of a medicament for the treatment of NR2 mediated conditions defective or deficient.

The present invention is further described with reference to the following non-limiting Examples.

The following single and three letter abbreviations for amino acid residues are used in the specification:

Amino Acid	Three-letter Abbreviation	One-letter Symbol
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartic acid	Asp	D
Cysteine	Cys	C
Glutamine	Gln	Q
Glutamic acid	Glu	E
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S

EXAMPLE 1

Cloning of a Human NR2 (Haemopoietin Receptor) cDNA

A cDNA library constructed from mRNA from a the bone marrow of a patient recovering from chemotherapy was constructed by C. G. Begley, Cancer Research Unit, WEHI in IZAP (Stratagene, Calif., USA) were used to infect *Escherichia coli* of the strain LE392. Infected bacteria were grown on twenty 150 mm agar plates, to give approximately 50,000 plaques per plate. Plaques were then transferred to duplicate 150 mm diameter nylon membranes (Colony/Plaque Screen™, NEN Research Products, MA, USA), bacteria were lysed and the DNA was denatured fixed by autoclaving at 100° C. for 1 min with dry exhaust. The filters were rinsed twice in 0.1% (w/v) sodium dodecyl sulfate (SDS), 0.1×SSC (SSC is 150 mM sodium chloride, 15 mM sodium citrate dihydrate) at room temperature and pre-hybridised overnight at 42° C. in 6×SSC containing 2 mg/ml bovine serum albumin, 2 mg/ml Ficoll, 2 mg/ml polyvinylpyrrolidone, 100 mM ATP, 10 mg/ml tRNA, 2 mM sodium pyrophosphate, 2 mg/ml salmon sperm DNA, 0.1% SDS and 200 mg/ml sodium azide. The pre-hybridisation buffer was removed. 1.2 mg of the degenerate oligonucleotides for hybridisation (HYB2; Table 2 above) were phosphorylated with T4 polynucleotide kinase using 960 mCi of $g^{32}P$ -ATP (Bresatec, S. A., Australia). Unincorporated ATP was separated from the labelled oligonucleotide using a pre-packed gel filtration column (NAP-5; Pharmacia, Uppsala, Sweden). Filters were hybridised overnight at 37° C. in 80 ml of the prehybridisation buffer containing and 10^6 – 10^7 cpm/ml of labelled oligonucleotide. Filters were briefly rinsed twice at room temperature in 6×SSC, 0.1% (v/v) SDS, twice for 30 min at 45° C. in a shaking waterbath containing 1.51 of the same buffer and then briefly in 6×SSC at room temperature. Filters were then blotted dry and exposed to autoradiographic film at –70° C. using intensifying screens, for 7–14 days prior to development.

Plaques that appeared to hybridise to the probe on duplicate filters were picked and eluted for 2 days at 4° C. in 0.5 ml of 100 mM NaCl, 10 mM MgCl₂, 10 mM Tris.HCl pH7.4

containing 0.5% (w/v) gelatin and 0.5% (v/v) chloroform. 5 ml aliquots of each eluate was used as the substrate for two PCR reactions containing 5 ml of 10× concentrated PCR buffer (Boehringer Mannheim GmbH, Mannheim, Germany), 1 ml of 10 mM dATP, dCTP, dGTP and dTTP, 2.5 ml of the oligonucleotides HYB2 and either T3 or T7 at a concentration of 100 mg/ml, 0.5 ml of Taq polymerase (Boehringer Mannheim GmbH) and water to a final volume of 50 ml. PCR was carried out in a Perkin-Elmer 9600 by heating the reactions to 96° C. for 2 min and then for 25 cycles at 96° C. for 30 sec, 55° C. for 30 sec and 72° C. for 2 min. The reactions were then electrophoresed on a 1% (w/v) low melting point agarose gel in TAE. Any positive products were excised, the gel slice was melted and 2 ml was used as the substrate for a second PCR reaction using conditions identical to the first. The product from the second reaction was purified using an ultrafree-MC centrifugal filtration unit (Millipore Corp.) by centrifugation for 15 min at 2000 g in an eppendorf centrifuge, adding 0.5 ml of 10 mM Tris.HCl, 1 mM EDTA pH8 and recentrifuging. This procedure was repeated three times and the DNA was recovered in 50 ml of 10 mM Tris.HCl, 1 mM EDTA pH8.

Approximately 500 ng of DNA from each PCR reaction was sequenced using a fmol sequencing kit (Promega Corporation, WI, USA), according to the manufacturer's instructions with the ³³P-labelled oligonucleotide primer HYB2. The products were resolved on a 6% w/v polyacrylamide gel and the sequence of each clone was analysed using the Blast database comparison programs and the translation function of the Wisconsin suite of DNA programs. The sequence of the PCR product derived from the primary plaque eluate number CF.32 appeared to be novel since it had no homologues in the databases of DNA sequences that were searched, and upon inspection of the sequence of the conceptually translated products appeared also to be a member of the haemopoietin receptor family. This clone was given the name of new receptor 2 or NR2.

The positively hybridising bacteriophage present in the eluate from the primary plug NR2-CF-32-1 was purified using a second round of screening performed in a manner identical to the first, except that plaques were grown on smaller, 82 mm, plates of agar. Once purified DNA, the positive cDNA cloned into the plasmid pBluescript was excised from the λ-ZAP II bacteriophage according to the manufacturer's instructions (Statagene). A CsCl purified preparation of the DNA was made and this was sequenced on both strands. Sequencing was performed using an Applied Biosystems automated DNA sequencer, with fluorescent dideoxynucleotide analogues according to the manufacturer's instructions. The DNA sequence was analysed using software supplied by Applied Biosystems.

EXAMPLE 2

Isolation of Additional NR2 cDNAs

NR2-CF.32 did not appear to contain the entire coding region of the novel receptor. In order to identify other cDNA libraries containing cDNA clones for NR2 we performed PCR upon 1 ml aliquots of λ-bacteriophage cDNA libraries made from mRNA from various human tissues and using oligonucleotides 722 and 761, designed from NR2-CF-32-1, as primers. The oligonucleotides are defined in Table 2, above. Reactions contained the same elements as described

above and were performed in an identical manner. In addition to the original library, five other cDNA libraries appeared to contain NR2 cDNAs. These were screened using a ³²P-labelled oligonucleotide 721 and 761 designed from the 5'-end and the 3' end of the sequence derived from NR2-CF.32, using conditions identical to those described in section (i) except that filters were washed at 55° C. rather than 45° C. Again, as described in section (i), positively hybridising plaques were purified, the cDNAs were recovered and cloned into plasmids pBluescript II or pUC19. Ten independent cDNA clones were sequenced on both strands. Further clones were isolated in a similar manner by screening libraries with oligonucleotide 875 and 939.

The extent of each clone is illustrated in FIG. 1 and a composite sequence is shown in FIG. 2. NR2 clearly has all the features of a member of the haemopoietin receptor family.

EXAMPLE 3

Analysis of the Expression Pattern of NR2 mRNA

Northern blots of mRNA from various human tissues and cell lines were hybridised with a random-primed human NR2 cDNA fragment from the internal EcoR I site to the Hpa I site (FIG. 1). Using the protocol described previously by Hilton et al. (15), two human NR2 mRNA species were observed to be expressed at a low level in a range of adult tissues, and at higher levels in foetal tissues such as the lung and liver. FIG. 9 shows expression of NR2 in various mouse tissues using human NR2 cDNA as probe. Interestingly among a series of human haemopoietin cell lines the megakaryocytic cell line MEG01 expressed high levels of NR2 mRNA suggesting that NR2 and its cognate ligand may play a role in the regulation of the megakaryocyte proliferation, differentiation and/or function.

EXAMPLE 4

Generation of Plasmids Directing the Expression of Full-Length and Secreted Forms Human NR2

Since antibodies to NR2 were not available to monitor expression, constructs were engineered to express full length and two soluble versions of NR2 with an N-terminal "FLAG" epitope (International Biotechnologies/Eastman Kodak, New Haven Conn.). First, a derivative of the mammalian expression vector pEF-BOS was generated so that it contained DNA encoding the signal sequence of murine IL-3 (MVLASSTSIHTMLLLLLMLFHLGLQASIS [SEQ ID NO. 14]) and the FLAG epitope (DYKDDDDK [SEQ ID NO. 15]) followed by a unique Xba I cloning site. This vector was named pEF/IL3SIG/FLAG.

The 5' end of the mature NR2 coding region was generated by PCR using primers 1056 and 721 on clone 60-58-7 (FIG. 1). The EcoR I/Hpa I fragment of clone 60-55-7-6 containing the 3' end of the NR2 coding region and a portion of the 3'-untranslated region was cloned into the EcoR I/SmaI digested pBluescript (FIG. 1). This construct was digested with Hind III and EcoR I and into it was cloned the 5'-NR2 PCR product digested with the same enzymes. The resulting construct was digested with Xba I to yield a fragment which contained the coding region of human NR2

from Y26 to the natural last amino acid L897 (FIG. 1) and a segment of 3'-untranslated region and was cloned into the Xba I site of pEF/IL3SIG/FLAG to give pEF/IL3SIG/FLAG/NR2/897. A soluble derivative of human NR2 was also engineered. PCR was carried out either using primers 1056 and 1092 to amplify the predicted mature coding region of the extracellular portion of human NR2 (Y26 to D839; FIG. 1). The PCR products were digested with Xba I and subcloned into Xba I digested pEF/IL3SIG/FLAG to give pEF/IL3SIG/FLAG/NR2/839. The identity of each construct was confirmed by dideoxy sequencing.

EXAMPLE 5

Transient Expression of Full Length and Secreted Forms of Human NR2 IN COS Cells

In order to confirm that full length and soluble NR2 could be produced using the expression vectors pEF/IL3SIG/FLAG/NR2/897 and pEF/IL3SIG/FLAG/NR2/839, COS cells were transiently transfected with these constructs. Briefly, COS cells from a confluent 175 cm² tissue culture flask were resuspended in PBS and electroporated (BioRad Gene pulser; 500 mF, 300 V) with 20 mg of uncut pEF/IL3SIG/FLAG/NR2/897 or pEF/IL3SIG/FLAG/NR2/839 in a 0.4 cm cuvette (BioRad). After 2 to 3 days at 37° C. in a fully humidified incubator containing 10% v/v CO₂ in air cells were used for analyses of protein expression. Conditioned medium was collected by centrifugation and stored sterile at 4° C. Cells were also harvested and lysed for 5 min in 500 ml of 50 mM Tris.HCl pH7.4 containing 150 mM NaCl, 2 mM EDTA and 1% v/v Triton X-100. The intact nuclei were removed by centrifugation at 10,000 g for 5 min. 500 ml of 50 mM Tris.HCl pH7.4 containing 150 mM NaCl, 2 mM EDTA, 1% v/v Triton X-100, 1% w/v sodium deoxycholate and 0.2% w/v SDS. 15 ml of anti-FLAG M2 affinity gel (International Biotechnologies/Eastman Kodak, New Haven Conn.) was then added to the cell extract or to 1 ml of conditioned medium and precipitation was carried out overnight at 4° C. The affinity gel was then washed three times in cold PBS and the precipitated protein was eluted by resuspending the gel in 80 ml of 100 mM sodium phosphate pH7.2, 10 mM EDTA, 0.1% w/v SDS and 1% 2-mercaptoethanol and boiling for 5 min. The supernatant was removed and 8 ml of 10% b-octyl glucoside was added. One half of each sample was incubated for 16 hours with 0.6 U of N-Glycanase-F (Boehringer-Mannheim), while the remainder was left untreated. An equal volume of 2× SDS-PAGE sample buffer was added to the samples which were then boiled and electrophoresed on pre-cast 4–15% w/v polyacrylamide gels (BioRad). The resolved proteins were then electroblotted onto Immobolon membranes, which were then blocked with 5% w/v skim milk, 0.1% v/v Tween 20 in PBS, rinsed and incubated with 5 ml of anti-FLAG M2 antibody in 2.5 ml of PBS containing 0.1% v/v Tween 20, rinsed and incubated with peroxidase-conjugated human anti-mouse Ig in 5% w/v skim milk, 0.1% v/v Tween 20 in PBS, rinsed and incubated with ECL reagent for 1 min. Filters were then blotted dry and exposed to autoradiographic film for 1 min.

COS cells that were mock transfected contained no reactive protein, while COS cells transfected with pEF/IL3SIG/

FLAG/NR2/897 expressed an immunoreactive protein of between 120,000 and 140,000 molecular weight. Deglycosylation with N-Glycanase-F resulted in a reduction in the apparent molecular weight to approximately 110,000 close to that predicted from the cDNA sequence of NR2. The immunoreactivity observed was completely inhibited by inclusion of an excess of the FLAG peptide during the immunoprecipitation step. No specific immunoreactive proteins could be detected in the medium conditioned by COS cells transfected with pEF/IL3SIG/FLAG/NR2/897. In contrast immunoreactive proteins were found in the medium and the cell pellet of COS cells transfected with DNA encoding the secreted form of NR2—pEF/IL3SIG/FLAG/NR2/839. The secreted form of NR2, as predicted, exhibited a lower apparent molecular weight than full length NR2, 10,000 to 120,000. This again decreased upon deglycosylation, to approximately 100,000.

COS cells transfected with pEF/IL3SIG/FLAG/NR2/897 were also examined for cell surface expression of NR2 by immunofluorescence staining. 5×10⁵ COS cells were resuspended in 100 ml of PBS containing 5% fetal calf serum and incubated with FITC-conjugated anti-FLAG M2 antibody for 45 min on ice, the cells were fixed and examined using a fluorescence microscope. No positive cells were observed in mock transfected samples, while approximately 10% of COS cells transfected with pEF/IL3SIG/FLAG/NR2/897 stained brightly positive. This data was consistent with the expected transient transfection efficiency of COS cells using electroporation.

EXAMPLE 6

Stable Expression of Full Length Human NR2

As described below certain routes to the identification of the NR2 ligand require stable expression of full-length NR2 in haemopoietic cell lines and the production and purification of large (mg) amounts of secreted NR2. Stable transfection of the pEF/IL3SIG/FLAG/NR2/897 and pEF/IL3SIG/FLAG/NR2/839 plasmids was achieved by electroporation. Briefly, the plasmids were linearised by digestion with the restriction enzyme Aat II. 20 mg of the linearised pEF/IL3SIG/FLAG/NR2/897 plasmid and 2 mg of pPGKpuropA, pPGKneopA or pPGKhydropA (plasmids directing the expression of the puromycin, neomycin and hygromycin resistance genes) were electroporated into 4×10⁶ parental Ba/F3 cells, Ba/F3 cells engineered to express human gp130 with or without coexpression of the human LIF receptor, Ba/F3 cells expressing the human b-chain common to the IL-3, IL-5 and GM-CSF receptors, Ba/F3 cells expressing the human IL-2 receptor b- and g-chains, CTLL cells or CHO cells. Briefly, cells were washed twice in ice-cold PBS and resuspended in PBS at 5×10⁶ per ml. 4×10⁶ cells were aliquoted into 0.4 mm electroporation cuvettes with the DNA. DNA and cells were incubated for 10 min on ice and electroporated at 270 V and 960 mF in a Bio-Rad Gene-Pulser (Bio-Rad Laboratories, Calif., USA). The cells were mixed with 1 ml of culture medium, centrifuged through 3 ml of FCS and resuspended in 100 ml of culture medium. Cells were then aliquoted into four 24 well dishes. After two days, selection was commenced by the addition puromycin to a concentration of 20

mg/ml, G418 to a concentration of 1.2 mg/ml or hygromycin to a concentration of 1 mg/ml. After 10–14 days, clones of proliferating cells were transferred to flasks and after expansion were tested for receptor expression.

FACS analysis using the anti-FLAG M2 antibody (FIG. 3) illustrates that Ba/F3 cells transfected with the pEF/IL3SIG/FLAG/NR2/897 express NR2 on the cell surface. Similar results have been obtained with other cell lines. As with COS cells, CHO cells transfected with pEF/IL3SIG/FLAG/NR2/839 secrete the NR2 extracellular domain. The extracellular domain of NR2 has been purified on an anti-FLAG M2 antibody affinity column using the FLAG peptide as the means of elution. This results in a high degree of purification of the NR2 extracellular domain as seen in the silver-stained poly-acrylamide gel illustrated in FIG. 4.

EXAMPLE 7

Strategies for Isolation of the Ligand for NR2

The stable expression of full-length and secreted NR2 enables steps to be taken to generate specific monoclonal antibodies to NR2 and allows a number of strategies to be employed to identify the cognate ligands of NR2.

(a) Expression of NR2 in Factor Dependent Cell Lines;

A variety of haemopoietin cell lines have been described which are dependent on the presence of exogenous growth factor for survival and proliferation in vitro. Among these are the murine cell lines Ba/F3, FDCP-1, 32D, CTLL, NFS-60, B6SutA, DA-1 and DA-3 and the human cell lines M07 and TF-1. FLAG-tagged murine and human NR2 may be stably expressed in each of these cell lines. The capacity of medium conditioned by a variety of murine and human cell lines and tissues to stimulate the survival and division of factor dependent cell lines expressing NR2 will be compared to the ability of the same medium to stimulate parental cell lines that do not express NR2. Medium that shows a greater ability to stimulate the proliferation cells expressing NR2 will be considered as a potential source of NR2.

NR2 has also been co-expressed in Ba/F3 cells with the LIF receptor α -chain and gp130, with the IL-2 receptor β - and γ -chains of the IL-2 receptor and with the common β -chain of the IL-3, IL-5 and GM-CSF receptors. Again conditioned medium will be tested for their ability to stimulate the proliferation of these cell lines.

(b) Identification of the NR2 Ligand Using the Cytosensor;

The haemopoietin cell lines expressing NR2 described above and additional non-haemopoietin cell lines engineered to express NR2 will be used in conjunction with the Cytosensor to screen conditioned medium for the presence of a ligand capable of altering cellular ion fluxes. Positive conditioned medium will be considered as a potential source of NR2 ligands.

(c) Selection of Ba/F3 Cells Expressing the NR2 Ligand;

Ba/F3 cells expressing NR2 with or without additional receptor components will be mutated with EMS or with a retrovirus. Mutants that are capable of proliferation in the absence of added growth factor will be selected. The medium from such clones will then be tested for their ability to stimulate the proliferation of Ba/F3 cells expressing NR2 with or without additional receptor components compared with the corresponding Ba/F3 cells that do not express NR2. Positive conditioned medium will be considered as a potential source of the NR2 ligand.

(d) Expression of NR2 in Cell Lines that may be Induced to Differentiate;

Similar experiments may be performed by expressing FLAF-tagged NR2 in cells that may be induced to differentiate by cytokines. Such cells include the murine lines M1 and WEHI-3BD+ and the human lines HL-60 and U937. The capacity of medium conditioned by a variety of murine and human cell lines and tissues to induce the differentiation of such cell lines expressing NR2 will be compared to the ability of the same medium to stimulate parental cell lines that do not express NR2. Medium that shows a greater ability to stimulate the differentiation of cells expressing NR2 will be considered as a potential source of NR2 ligand.

(e) Use of Secreted NR2 Extracellular Domain as a Probe on the Biosensor;

Purified extracellular domain of NR2 has been obtained and is being immobilized on the surface of a Biosensor chip. Medium conditioned by a variety of murine and human cell lines and tissues will be passed across the chip and specific changes in the surface plasmon resonance will be noted. Positive medium will be considered as a potential source of NR2 ligand.

(f) Use of Secreted NR2 Extracellular Domain as the Basis of an Affinity Column;

Purified extracellular domain of NR2 has been obtained and is being immobilized using a variety of chemistries. Affinity columns will be constructed and medium conditioned by a variety of murine and human cell lines and tissues will be passed through. Proteins that bind to the column will be considered to be candidate NR2 ligands and will be further characterised.

EXAMPLE 8

Human Leptin

A human leptin cDNA (16) was cloned into the pFBOS expression vector (17) in frame with the interleukin-3 leader sequence followed by the FLAGTM epitope sequence (18). CHO cells were transfected with this vector by electroporation and supernatant harvested from exponentially growing cultures. The supernatant was concentrated over a YM-10 membrane (10-fold) and then applied to an affinity column containing immobilised anti-FLAGTM antibody M2. The column was eluted with FLAGTM peptide according to the manufacturers instructions (Eastman Kodak, Rochester, N.Y.). The monomeric form of human leptin was purified by gel filtration chromatography on a Superose 12 column (Pharmacia, Uppsala, Sweden) and exchanged into 20 mM phosphate buffered (pH7.4) saline (0.15 M) containing 0.02% v/v Tween 20 and 0.02% w/v sodium azide (PBS) by gel filtration on Sephadex G-25 M (PD-10) columns (Pharmacia). Human leptin was iodinated with ¹²⁵I using a modified iodine monochloride method (19) to a specific radioactivity of approximately 10⁷ cpm/pmol and exchanged into PBS as above.

EXAMPLE 9

Binding of ¹²⁵I Human Leptin to Cells Expressing NR2 or to Soluble NR2

Cos-hNR2 are COS-7 cells electroporated with pFBOS-hNR2 and harvested at 3. days (5×10⁴ cells used per point).

23

Ba/F3-hNR2 are Ba/F3 cells stably transfected with peFBOS-hNR2 (9×10^5 cells used per point).

Solh NR2 is a soluble form of human NR2 purified by anti-FLAGTM affinity chromatography from the supernatant (48 hr) of COS cells transfected with peFBOS-solh NR2 (approx. 0.1 $\mu\text{g/ml}$ final concentration in binding assay).

For cells, the total reaction volume was 100 μl in RPMI-medium containing 10 mM Hepes pH7.4 and 10% v/v foetal calf serum (RHF). The reaction mixture also contained ^{125}I h leptin $0-6 \times 10^5$ cpm as indicated with or without unlabelled h leptin (approx. 1 $\mu\text{g/ml}$).

The mixture was incubated for 1–1.5 hr at 23° C. and then layered over 200 μl cold foetal calf serum in small, tapered centrifuge tubes (Elkay, Melbourne) and centrifuged at 12000 g for 10 sec. The cell pellet was removed by cutting the tubes with a scalpel blade and the cell bound (pellet) radioactivity and the unbound radioactivity (the rest of the tube) were separately counted in a Packard γ -counter. Specifically bound ^{125}I h leptin was determined as the difference in counts between otherwise identical tubes that contained or did not contain the unlabelled excess h leptin. The data were plotted as saturation curves (specifically bound versus added ^{125}I h leptin) and as Scatchard transformations (specific bound/free radioactivity versus specific bound radioactivity to determine the equilibrium dissociation constants [K_d]) (20).

For soluble receptors (sol hNR2) incubations were as above but after 1 hr at 23° C., 20 μl of concavalin A-sepharose 4B beads ($\frac{1}{4}$ suspension in 0.1 M acetate pH5) were added and incubation continued for a further 30 min. Subsequently, the beads were centrifuged and processed as above. The results are shown in FIGS. 5 and 6. Human leptin binds to Ba/F3/COS cells transfected with hNR2 cDNA and to soluble hNR2.

EXAMPLE 9

Expression of NR2 in Animal Species

Genomic DNA from various sources was digested with EcoRI. This was then blotted onto a nylon membrane (GeneScreen Plus®, NEN Research Products, USA). The filter was then probed using a 1.1 kb cDNA fragment of NR2. The fragment covers the 3' half of the first haemopoietin domain and extends to cover the whole of the second haemopoietin domain, terminating the type III fibronectin domain. The filter was prehybridised and hybridised in 0.5M sodium phosphate, 7% w/v SDS and 1 mM EDTA at 50° C. overnight. The filter was then washed in 40 mM sodium phosphate and 1% w/v SDS at 50° C. The results are shown in FIG. 7.

EXAMPLE 10

Cloning of the Human NR2 Locus

In order to obtain genomic clones of the human NR2 locus, various genomic libraries were screened. These libraries were screened with either oligonucleotide or cDNA probes. Oligonucleotide screening conditions: 1×10^6 clones

24

were fixed to nylon filters (Colony/Plaque ScreenTM, NEN Research Products, USA). These filters were then prehybridised in a 6 \times SSC buffer containing 0.2% Ficoll, 0.2% w/v bovine serum albumin, 0.2% polyvinylpyrrolidone, 0.1M ATP, 50 $\mu\text{g/ml}$ transfer RNA, 2 mM tetra-sodium pyrophosphate, 50 $\mu\text{g/ml}$ herring sperm DNA and 0.1% w/v sodium azide at 37° C. for at least 2 hours. They were hybridised overnight under the same conditions, with at least 2×10^6 cpm/mL of ^{32}P -labelled oligonucleotide probe. The filters were then washed in 6 \times SSC/0.1% w/v SDS at 50–55° C. depending on the sequence of the specific oligonucleotide (Melting Temp –10C).

cDNA screening conditions: 10×10^6 clones were fixed to nylon filters. These filters were then prehybridised in a 2 \times SSC buffer containing 0.2% Ficoll, 0.2% w/v bovine serum albumin, 0.2% polyvinylpyrrolidone, 0.1M ATP, 50 $\mu\text{g/ml}$ transfer RNA, 2 mM tetra-sodium pyrophosphate, 50 $\mu\text{g/ml}$ herring sperm DNA and 0.1% w/v sodium azide at 37° C. for at least 2 hours at 65° C. They were hybridised overnight under the same conditions, with at least 2×10^6 cpm/mL of ^{32}P -labelled cDNA fragment. The filters were then washed in 2 \times SSC/0.1% w/v SDS at 65° C.

EXAMPLE 11

Restriction Enzyme Mapping

The clones obtained were characterised by mapping with partial endonuclease digestion (21).

In order to determine on which fragments the various exons were present, specific oligonucleotide probes were used. The various clones were digested with a range of restriction enzymes. These were then blotted to a nylon membrane (GeneScreen Plus®), NEN Research Products, USA). Oligonucleotides derived from the cDNA sequence (and known to be specific for a particular exon), were then hybridised to the digested fragments. These hybridisations were done under the same conditions as mentioned above for oligonucleotides. Exons could then be mapped to specific fragments by a positive hybridisation after overnight exposure.

Intron/exon boundary sequences were determined by sequencing across the intron/exon boundaries. Primers specific for sequence on either side of the boundary were used in a sequencing PCR reaction. Sequencing was performed on an ABI 373 sequencer using the Taq cycle sequencing kit (Applied Biosystems). These sequences were then compared to the consensus intron/exon boundary sequence (22). The results are shown in FIG. 8 and in Table 3.

EXAMPLE 12

Determination of Amino Acid Sequence of hNR2

The N-terminal amino acid sequence of hNR2 was determined. The results are shown below. The actual sequence starts at amino acid 16. The sequence is as follows:

Asp Ser Ile Ser Ser Ser Asp Tyr Lys Asp Asp Asp Glu Ser Arg [SEQ ID NO:44]
 5 10 15
 Tyr Pro Ile Thr Pro Trp Arg Phe Lys Leu Ser Xaa Met Pro Pro
 20 25 30
 Xaa Ser Thr Tyr Asp
 35

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

- 10
 10. Yonemura, Y., Kawakita, M., Masuda, T., Fujimoto, K., Kato, K. and Takatsuki, K. (1992) *Exp. Hematol.* 20: 1011–6.
 11. Baumann, H. and Schendel, P. (1991) *J. Biol. Chem.* 266: 20424–7.
 12. Kawashima, I., Ohsumi, J., Mita-Honjo, K., Shimoda-Takano, K., Ishikawa, H., Sakakibara, S., Miyadai, K. and Takiguchi, Y. (1991) *Febs. Lett.* 283: 199–202.

TABLE 3

Intron-Exon junctions of the human NR2 gene								
Exon	Exon size (bp)	DONOR			Intron size	ACCEPTOR		
ss	60	AITGGG	gtaagttatt	[SEQ ID NO. 16]		ccttttcag	GTGTAT	[SEQ ID NO. 29]
lg	330	AAATAG	gtaagcatta	[SEQ ID NO. 17]		tcctaacag	AAATTA	[SEQ ID NO. 30]
SD100A	124	TGTTCT	gtaagtacca	[SEQ ID NO. 18]		ttaaatcag	ATGCAA	[SEQ ID NO. 31]
SD100B								
SD100B	145	CACAAG	gtaggttatg	[SEQ ID NO. 19]		tatttaacag	GCTGAC	[SEQ ID NO. 32]
Ig	291	TGATTG	gtaagaaca	[SEQ ID NO. 20]	0.16	ctcattacag	ATGTCA	[SEQ ID NO. 33]
SD100A'	118	AITGAG	gtatcatagg	[SEQ ID NO. 21]	2.3	tttcaaatag	ATGTGA	[SEQ ID NO. 34]
SD100A'	200	CTGTGG	gtatgtcaag	[SEQ ID NO. 22]	2.4	tcttttaag	GAGCAG	[SEQ ID NO. 35]
SD100B'	149	TGGAAG	gtaccttta	[SEQ ID NO. 23]		aaatttcag	TGAAGC	[SEQ ID NO. 36]
SD100B'	160	TAAAAG	gtctgcagag	[SEQ ID NO. 24]	0.2	tattttacag	ATGTAT	[SEQ ID NO. 37]
FnIII	83	TGGAGG	gtatnccaat	[SEQ ID NO. 25]	>7 kbp	catttgccag	TTCCTA	[SEQ ID NO. 38]
FnIII	161	CAATTC	aattggtgct	[SEQ ID NO. 26]		tttactacag	CCCCTG	[SEQ ID NO. 39]
FnIII'								
FnIII'								
Tm	106	CCAAAG	gtattgtact	[SEQ ID NO. 27]	1.4	tcttttcag	ATGATA	[SEQ ID NO. 40]
Cyt (Box I)	76	CATAAG	gttgettttt	[SEQ ID NO. 28]	3	ccctttgtag	AATGAA	[SEQ ID NO. 41]
Cyt' (NR2.2)	212				3	cccttttcag	AAAATG	[SEQ ID NO. 42]
3'UTR	>1085				3	atctaacag	AGAACG	[SEQ ID NO. 43]
Consensus		AG	gt ^a /gagt			tc rich-cag	G	

REFERENCES

- 45
 1. Du, X. X. and Williams, D. A. (1994) *Blood* 83: 2023–2030.
 2. Yang, Y. C. and Yin, T. (1992) *Biofactors* 4: 15–21.
 3. Paul, S. R., Bennett, F., Calvetti, J. A., Kelleher, K., Wood, C. R., O'Hara, R. J. J., Leary, A. C., Sibley, B., Clark, S. C., Williams, D. A. and Yang, Y. -C. (1990) *Proc. Natl. Acad. Sci. USA* 87: 7512.
 4. Musashi, M., Clark, S. C., Sudo, T., Urdal, D. L., and Ogawa, M. (1991) *Blood* 78: 1448–1451.
 5. Schibler, K. R., Yang, Y. C. and Christensen, R. D. (1992) *Blood* 80: 900–3.
 6. Tsuji, K., Lyman, S. D., Sudo, T., Clark, S. C., and Ogawa, M. (1992) *Blood* 79: 2855–60.
 7. Burstein, S. A., Mei, R. L., Henthorn, J., Friese, P. and turner, K. (1992) *J. Cell. Physiol.* 153: 305–12.
 8. Hangoc, G., Yin, T., Cooper, S., Schendel, P., Yang, Y. C. and Broxmeyer, H. E. (1993) *Blood* 81: 965–72.
 9. Teramura, M., Kobayashi, S., Hoshino, S., Oshimi, K. and Mizoguchi, H. (1992) *Blood* 79: 327–31.
 10. Keller, D. C., Du, X. X., Srouf, E. f., Hoffman, R. and Williams, D. A. (1993) *Blood* 82:1428–35.
 11. Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989) *Molecular cloning: A Laboratory Manual* Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
 12. Hilton, D., et al (1994) *EMBO Journal* 13: 4765–4777.
 13. Gong et al (1996) *J. Biol. Chem.* 271: 3971–3974.
 14. Mizushima and Nagata (1990) *Nucleic Acids Research* 18: 5322.
 15. Nicola et al (1996) *Growth Factors* 13: 141–149.
 16. Hilton and Nicola (1992) *J. Biol. Chem.* 267: 10238–10247.
 17. Scatchard (1949) *Ann. N.Y. Acad. Sciences* 51: 660–672.
 18. Unit 3.3.1 (1995) *Current Protocols in Molecular Biology*. Editors Ausubel, F. M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A. and Strul, K.
 19. Shapiro, M. B. and Senapathy, P. (1987) *Nucleic Acids Research* 15: 7155–7174.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 1

rctccartcr ctcca 15

<210> SEQ ID NO 2
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 2

taatacgact cactataggg aga 23

<210> SEQ ID NO 3
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 3

attaccctca ctaaaggga 19

<210> SEQ ID NO 4
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 4

actagcaggg atgtagctga g 21

<210> SEQ ID NO 5
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 5

ctcagctaca tccctgctag t 21

<210> SEQ ID NO 6
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 6

ctgctcctat gatacct 17

<210> SEQ ID NO 7
 <211> LENGTH: 22

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 7

cctcttccat cttattgctt gg                                     22

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 8

atcggtcgtg acatacaagg                                         20

<210> SEQ ID NO 9
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 9

agctaagctt tctagatatc caattactcc ttggaga                     37

<210> SEQ ID NO 10
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 10

agcttctaga tcaatcactc tgggtgtttt caat                         34

<210> SEQ ID NO 11
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 11

acgttctaga tcaaactttt atatccatga caac                        34

<210> SEQ ID NO 12
<211> LENGTH: 3909
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (27)..(68)
<223> OTHER INFORMATION: N is a or g or c or t
<221> NAME/KEY: unsure
<222> LOCATION: (923)
<223> OTHER INFORMATION: R is g or a
<221> NAME/KEY: unsure
<222> LOCATION: (2315)
<223> OTHER INFORMATION: S is g or c

<400> SEQUENCE: 12

cgaattcgcg ggcgcgtcga ccgcggnccc agctcgggag acatgggggg cgttaaagct   60
ctcgtggnat tatccttcag tggggstatt ggactgactt ttcttatgct gggatgtgcc   120

```

-continued

ttagaggatt atggatttgg cagttcaccc tgaccatctt gaaaataagt tatctctgat	180
ctctgtctgt atgttacttc tctcccctca ccaacggaga acaaatgtgg gcaaagtgtg	240
cttctctgaa gtaagatgat ttgtcaaaaa ttctgtgtgg ttttgttaca ttgggaattt	300
atztatgtga taactgcgtt taacttgtca tatccaatta ctcttgag atttaagttg	360
tcttgcatgc caccaaatc aacctatgac tacttccttt tgcctgctgg actctcaaag	420
aatacttcaa attcgaatgg acattatgag acagctgttg aacctaatg taattcaagt	480
ggtactcact tttctaact atccaaaaca actttccact gttgctttcg gagtgagcaa	540
gatagaaact gctccttatg tgacagacaac attgaaggaa ggacatttgt ttcaacagta	600
aattccttag tttttcaaca aatagatgca aactggaaca tacagtgtg gctaaaagga	660
gacttaaaat tattcatctg ttatgtggag tcattattta agaactctatt caggaaattat	720
aactataagg tccatctttt atatgttctg cctgaagtgt tagaagattc acctctggtt	780
ccccaaaag gcagttttca gatggttcac tgcaattgca gtgttcata atgttgtgaa	840
tgtcttgtgc ctgtgccaac agccaaactc aacgacactc tccttatgtg tttgaaatc	900
acatctgggt gagtaatttt corgtcacct ctaatgtcag ttcagcccat aaatatgggtg	960
aagcctgac caccattagg ttgcatatg gaaatcacag atgatggtaa tttaaagatt	1020
tcttggtcca gccaccatt ggtaccattt ccacttcaat atcaagtga atattcagag	1080
aattctaca cagttatcag agaagctgac aagattgtct cagctacatc cctgctagta	1140
gacagtatac ttcctgggtc ttcgtatgag gttcagtgta ggggcaagag actggatggc	1200
ccaggaatct ggagtactg gagtactcct cgtgtcttta ccacacaaga tgtcatatac	1260
tttccacct aaattctgac aagtgttggg tctaattgtt cttttcactg catctataag	1320
aaggaaaaca agattgttcc ctcaaagag attgtttggt ggatgaattt agctgagaaa	1380
attcctcaa gccagatga tgttgtgagt gatcatgta gcaaagtac tttttcaat	1440
ctgaatgaaa ccaaacctcg aggaaagttt acctatgatg cagtgtactg ctgcaatgaa	1500
catgaatgcc atcatcgcta tgcgaatta tatgtgattg atgtcaatat caatatctca	1560
tgtgaaactg atgggtactt aactaaaatg acttgcagat ggtcaaccag tacaatccag	1620
tcacttgctg aaagcacttt gcaattgagg tatcatagga gcagccttta ctgttctgat	1680
attccatcta ttcaccccat atctgagccc aaagattgct atttgagag tgatggtttt	1740
tatgaatgca ttttccagcc aatcttcta ttatctggtc acacaatgtg gattaggatc	1800
aatcactctc taggttact tgactctcca ccaacatgtg tccttcctga ttctgtggtg	1860
aagccactgc ctccatccag tgtgaaagca gaaattacta taaacattgg attattgaaa	1920
atatcttggg aaaagccagt ctttccagag aataaccttc aattccagat tcgctatggt	1980
ttaagtggaa aagaagtaca atggaagatg tatgaggttt atgatccaaa accaaaatct	2040
gtcagctctc cagttccaga cttgtgtgca gtctatgctg ttcaggtggc gtttaagagg	2100
ctagatggac tgggatattg gagtaattgg agcaatccag cctacacagt tgtcatggat	2160
ataaaagtct ctatgagagg acctgaattt tggagaataa ttaatggaga tactatgaaa	2220
aaggagaaaa atgtcacttt actttggaag ccctgatga aaaatgactc attgtgcagt	2280
gttcagagat atgtgataaa ccatcatact tcctscatg gaacatggc agaagatgtg	2340
ggaaatcaca cgaaattcac tttctgtgg acagagcaag cacatactgt tacggttctg	2400
gccatcaatt caattgtgct tctgttgca aattttaatt taaccttttc atggcctatg	2460
agcaaagtaa atatcgtgca gtcactcagt gcttatcctt taaacagcag ttgtgtgatt	2520

-continued

```

gtttcctgga tactatcacc cagtgattac aagctaagt atttattat tgagtggaaa 2580
aatcttaatg aagatgggta aataaaatgg cttagaatct cttcatctgt taagaagtat 2640
tatatccatg atcattttat ccccatgtgag aagtaccagt tcagtcctta cccaatattt 2700
atggaaggag tgggaaaacc aaagataatt aatagtttca ctcaagatga tattgaaaaa 2760
caccagagtg atgcagggtt atatgtaatt gtgccagtaa ttatttcctc ttccatctta 2820
ttgcttgga cattaataat atcacaccaa agaagaaaa agctattttg ggaagatgtt 2880
ccgaacccca agaattgttc ctgggcacaa ggacttaatt ttcagaagag aacggacatt 2940
ctttgaagtc taatcatgat cactacagat gaaccaatg tgccaacttc ccaacagtct 3000
atagagtatt agaagatttt tacattttga agaaggggag caaatctaaa aaaaattcag 3060
ttgaacttct gagagttaac atatggtgga ttatgttgat ttagaactta aaatagatgt 3120
catttaaacc caagttttac atctaaactc aggtcaaacc tacacactaa ttaaagtttt 3180
agtagatttc aaattttcat cataagtact aaagaccgaa aactaaacag tataaggacc 3240
agtattttgt aattctttta ataccgacaa cgacagtaat gtatagataa tttacagtag 3300
tttatacatc atctgttagg acattaatcc acttgagatt ttgacgttgt agactgttta 3360
tcgaaatfff tatgttacta atattcatac cttagtcact tttataaatc aaacataaaa 3420
atacaggttt gaaaaggtaa aatctaagga aatatctgtg cagtcggatt tttagtcgga 3480
taagcccaca agaaaactta tagaggaccg taaaaacata gattgaaaca agttagaccc 3540
ttaaagtcaa aagttatagg aacttttacc gaattcacta ttgaaggcaa agtcaatfff 3600
ccttcgggct tcaacacaaa caccgacggg gtctgtcac cctcaatgac aagtatagtc 3660
ctactgggat gtatgggtcc agtctaactg ccctggtctt ccctgttagc tgaagattac 3720
aggtgcgaaa gaacaaatta atactggatt tagattaaat gaaggtgact tggtaggttc 3780
tggagaccgt ccgtcccttt acccgctact asgttttttc cctctgagaa acctcgaaaa 3840
tacttatcaa gtaccactcc tgtcttgaag agatgaaagt ctgtctgacg aacgatcaaa 3900
atacttaag 3909

```

```

<210> SEQ ID NO 13
<211> LENGTH: 896
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: UNSURE
<222> LOCATION: (223)
<223> OTHER INFORMATION: Xaa is unknown or other.
<221> NAME/KEY: UNSURE
<222> LOCATION: (687)
<223> OTHER INFORMATION: Xaa is unknown or other.

```

```

<400> SEQUENCE: 13

```

```

Met Ile Cys Gly Lys Phe Cys Val Val Leu Leu His Trp Gln Phe Ile
 1             5             10            15

Tyr Val Ile Thr Ala Phe Asn Leu Ser Tyr Pro Ile Thr Pro Trp Arg
      20            25            30

Phe Lys Leu Ser Cys Met Pro Pro Asn Ser Thr Thr Asn Tyr Phe Leu
      35            40            45

Leu Pro Ala Gly Leu Ser Lys Asn Thr Ser Asn Ser Asn Gly His Tyr
      50            55            60

Glu Thr Ala Val Glu Pro Lys Phe Asn Ser Ser Gly Thr His Phe Ser
      65            70            75            80

```


-continued

500					505					510					
Ile	Arg	Ile	Asn	His	Ser	Leu	Gly	Ser	Leu	Asn	Ser	Pro	Pro	Thr	Cys
		515					520					525			
Val	Leu	Pro	Asp	Ser	Val	Val	Lys	Pro	Leu	Pro	Pro	Ser	Ser	Val	Lys
	530					535					540				
Ala	Glu	Ile	Thr	Ile	Asn	Ile	Gly	Leu	Leu	Lys	Ile	Ser	Trp	Glu	Lys
545					550					555					560
Pro	Val	Phe	Pro	Glu	Asn	Asn	Leu	Gln	Phe	Gln	Ile	Arg	Thr	Gly	Leu
				565					570					575	
Ser	Gly	Lys	Glu	Val	Gln	Trp	Lys	Met	Tyr	Glu	Val	Thr	Asn	Pro	Lys
		580						585					590		
Pro	Lys	Ser	Val	Ser	Leu	Pro	Val	Pro	Asp	Leu	Cys	Ala	Val	Tyr	Ala
		595				600						605			
Val	Gln	Val	Arg	Phe	Lys	Arg	Leu	Asp	Gly	Leu	Gly	Tyr	Trp	Ser	Asn
	610					615					620				
Trp	Ser	Asn	Pro	Ala	Tyr	Thr	Val	Val	Met	Asp	Ile	Lys	Val	Pro	Met
625					630					635					640
Arg	Gly	Pro	Glu	Phe	Trp	Arg	Ile	Ile	Asn	Gly	Asp	Thr	Met	Lys	Lys
				645					650					655	
Glu	Lys	Asn	Val	Tyr	Leu	Leu	Trp	Lys	Pro	Leu	Met	Lys	Asn	Asp	Ser
			660					665					670		
Leu	Cys	Ser	Val	Gln	Arg	Tyr	Val	Ile	Asn	His	His	Thr	Ser	Xaa	Asn
		675					680					685			
Gly	Thr	Trp	Ser	Glu	Asn	Val	Gly	Asn	His	Thr	Lys	Phe	Thr	Phe	Leu
	690					695					700				
Trp	Thr	Glu	Gln	Ala	His	Thr	Val	Thr	Val	Leu	Ala	Ile	Asn	Ser	Ile
705					710					715					720
Gly	Ala	Ser	Val	Ala	Asn	Phe	Asn	Leu	Thr	Phe	Ser	Trp	Pro	Met	Ser
			725						730					735	
Lys	Val	Asn	Ile	Val	Gln	Ser	Leu	Ser	Ala	Tyr	Pro	Leu	Asn	Ser	Ser
			740					745					750		
Cys	Val	Ile	Val	Ser	Trp	Ile	Leu	Ser	Pro	Ser	Asp	Val	Lys	Leu	Met
	755						760					765			
Tyr	Pro	Ile	Ile	Glu	Trp	Lys	Asn	Leu	Asn	Glu	Asp	Gly	Glu	Ile	Lys
	770					775					780				
Trp	Leu	Arg	Ile	Ser	Ser	Ser	Val	Lys	Lys	Tyr	Tyr	Ile	His	Asp	His
785				790						795					800
Phe	Ile	Pro	Ile	Glu	Lys	Tyr	Gln	Phe	Ser	Leu	Tyr	Pro	Ile	Phe	Met
			805						810					815	
Glu	Gly	Val	Gly	Lys	Pro	Lys	Ile	Ile	Asn	Ser	Phe	Thr	Gln	Asn	Asn
		820						825					830		
Ile	Glu	Lys	His	Gln	Ser	Asp	Ala	Gly	Leu	Tyr	Val	Ile	Val	Pro	Val
	835						840					845			
Ile	Ile	Ser	Ser	Ser	Ile	Leu	Leu	Leu	Gly	Thr	Leu	Leu	Ile	Ser	His
	850					855					860				
Gln	Arg	Met	Lys	Lys	Leu	Phe	Trp	Glu	Asp	Val	Pro	Asn	Pro	Lys	Asn
865					870					875					880
Cys	Ser	Trp	Ala	Gln	Gly	Leu	Asn	Phe	Gln	Lys	Arg	Thr	Asn	Ile	Leu
			885						890					895	

<210> SEQ ID NO 14

<211> LENGTH: 30

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 14

Met Val Leu Ala Ser Ser Thr Thr Ser Ile His Thr Met Leu Leu Leu
 1             5             10             15

Leu Leu Met Leu Pro His Leu Gly Leu Gly Ala Ser Ile Ser
      20             25             30

<210> SEQ ID NO 15
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 15

Ala Thr Leu Ala Ala Ala Ala Leu
 1             5

<210> SEQ ID NO 16
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 16

attggggtaa gttatt                                     16

<210> SEQ ID NO 17
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:sSynthetic

<400> SEQUENCE: 17

aaataggtaa gcatta                                     16

<210> SEQ ID NO 18
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 18

tgttctgtaa gtacca                                     16

<210> SEQ ID NO 19
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 19

cacaaggtag gttatg                                     16

<210> SEQ ID NO 20
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

```

-continued

<400> SEQUENCE: 20

tgattggtaa gaaaca

16

<210> SEQ ID NO 21

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 21

attgaggtat catagg

16

<210> SEQ ID NO 22

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 22

ctgtgggtat gtcaag

16

<210> SEQ ID NO 23

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 23

tggaaggtac ctttta

16

<210> SEQ ID NO 24

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 24

taaaaggtct gcagag

16

<210> SEQ ID NO 25

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<221> NAME/KEY: unsure

<222> LOCATION: (11)

<223> OTHER INFORMATION: N is a or g or c or t

<400> SEQUENCE: 25

tggaggggtat nccaat

16

<210> SEQ ID NO 26

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 26

caattcaatt ggtgct

16

-continued

<210> SEQ ID NO 27
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 27

ccaaaggtat tgtact 16

<210> SEQ ID NO 28
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 28

cataaggttg cttttt 16

<210> SEQ ID NO 29
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 29

ccttttccag gtgtat 16

<210> SEQ ID NO 30
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 30

tcctaacaga attta 15

<210> SEQ ID NO 31
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 31

ttaaattcag atgcaa 16

<210> SEQ ID NO 32
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 32

tatttaacag gctgac 16

<210> SEQ ID NO 33
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 33

ctcattacag atgtca 16

<210> SEQ ID NO 34
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 34

tttcaaataag atgtga 16

<210> SEQ ID NO 35
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 35

tccttttaaag gagcag 16

<210> SEQ ID NO 36
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 36

aaattttctag tgaagc 16

<210> SEQ ID NO 37
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 37

tatttttacag atgtat 16

<210> SEQ ID NO 38
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 38

catttggcag ttccta 16

<210> SEQ ID NO 39
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 39

tttactacag cccctg 16

-continued

```

<210> SEQ ID NO 40
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 40

tctttttcag atgata                                     16

<210> SEQ ID NO 41
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 41

ccctttgtag aatgaa                                     16

<210> SEQ ID NO 42
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 42

ccttttccag aaaatg                                     16

<210> SEQ ID NO 43
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Synthetic

<400> SEQUENCE: 43

atctaaacag agaacg                                     16

<210> SEQ ID NO 44
<211> LENGTH: 35
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: UNSURE
<222> LOCATION: (27)
<223> OTHER INFORMATION: Xaa is unknown or other.
<221> NAME/KEY: UNSURE
<222> LOCATION: (31)
<223> OTHER INFORMATION: Xaa is unknown or other.

<400> SEQUENCE: 44

Ala Ser Ile Ser Ser Ser Ala Thr Leu Ala Ala Ala Gly Ser Ala Thr
 1           5           10           15

Pro Ile Thr Pro Thr Ala Pro Leu Leu Ser Xaa Met Pro Pro Xaa Ser
          20           25           30

Thr Tyr Asp
          35

```

49

- What is claimed is:
1. A recombinant haemopoietin receptor comprising the amino acid sequence as set forth in SEQ ID NO: 13.
 2. An isolated polypeptide comprising the amino acid sequence as represented by Y26 to D839 of SEQ ID NO: 13. 5
 3. A pharmaceutical composition comprising a recombinant haemopoietin receptor and at least one of a pharmaceutically acceptable carrier or a diluent, wherein said

50

- recombinant haemopoietin receptor comprises the amino acid sequence as set forth in SEQ ID NO:13.
4. A pharmaceutical composition comprising an isolated polypeptide and at least one of a pharmaceutically acceptable carrier or a diluent, wherein said isolated polypeptide comprises the amino acid sequence as represented by Y26 to D839 of SEQ ID NO:13.

* * * * *