

Orbit Propagator Theory

Cooper Chang Chien
Taiwan Space Agency (TASA), System Engineer

Abstract

This is a collection of general theories and equations that establishes the algorithm as the building block of the orbit propagation simulator. This note includes the basics of orbital mechanics starting from the six orbital elements, introduction on GCRF, ITRF, and LVLH reference frames, towards Keplerian orbital equations. In later sections, a detailed discussion on non-keplerian motions were conducted, this section mainly focuses on the derivation of the gravitational potential function due to Earth's oblateness, and to arrive to the acceleration used in propagation.

NOTE: Some of the concepts here require extensive literature reading, serve this either as a starting point or collection of knowledge, as I'll do my best to include materials that are helpful during my studies.

Index Terms — Orbit propagation, perturbation, Legendre polynomials, gravitational potential, satellite

I INTRODUCTORY ON ASTRODYNAMICS

This section will be focusing on the fundamental theory and concept of astrodynamics, starting from coordinate reference frames, orbital elements, to relating time with satellite orbit position.

I-A Reference Frame - Earth Based System

One of the fundamental requirements of describing an orbit is to select a suitable reference frame. Beginning January 1, 1998, the IAU adopted the **International Celestial Reference System** (ICRS) as the standard reference system, introducing the ICRF frame where the origin is at the barycentre of the solar system. In 1991, the IAU introduced the Barycentric (BCRS) and Geocentric (GCRS) Celestial Reference Systems. The IAU-2000 Resolutions use both the celestial (GCRF) and terrestrial (ITRF) frames, where the origin references the Earth's centre.

I-A1 GCRF

I-A2 ITRF

I-A3 PQW

I-A4 LVLH

I-B Defining an Orbit - Orbit Elements

I-C Position and Velocity

II SPECIAL PERTURBATIONS

The oblateness perturbation is one of the principal perturbing accelerations acting on Earth satellites from LEO to GEO, although its strength decays relatively rapidly with the orbital radius. Departures of the gravitational field from the spherical symmetry due to finer variations in the Earth's mass distribution are captured through a perturbing potential. Perturbations due to small longitudinal asymmetries (i.e., along the parallels) are particularly important for geostationary orbit, as they affect the stability of satellites on the geostationary ring. This section we look into the derivation and theory behind the gravitational potential function, how to interpret it and arrive to the acceleration term that are use in propagations.

II-A Gravitational Potential Model

Given that the gradient of the potential for a spherical central body will yield the acceleration, we must examine how to form a **potential function** that includes the perturbing accelerations due to nonspherical central body.

II-A1 Deriving the Aspherical-Potential Function

If we examine an infinite number of masses, m_Q , at point Q , the potential per unit mass at point P is the summation of the potential due to each of these points, and the acceleration is ∇U . Suppose ρ is the radius from the point mass m_Q to the spacecraft. The change in potential due to an infinitesimal element of mass, dm_\oplus , is

$$U = G \sum_{Q=1}^{\infty} \frac{m_Q}{\rho_Q}, \quad dU = g \frac{dm_\oplus}{\rho_Q} \quad (1)$$

Considering Earth as being composed by a set of such infinitesimal masses (see Fig. (1)), each giving a contribution of dU to the total Gravitational potential, thus, the summation of each mass approaches an integral and can be express as

$$U = G \int_{body} \frac{1}{\rho_Q} dm_\oplus \quad (2)$$

To relate the potential to the position vector of the spacecraft with respect to the Earth's centre of mass \mathbf{r} , we can express ρ using the law of cosines,

$$\rho_Q = \mathbf{r} - \mathbf{r}_Q = \sqrt{r^2 + r_Q^2 - 2rr_Q \cos(\Lambda)} \quad (3)$$

where the angle $\cos(\Lambda)$ can be determined using dot product, $\cos(\Lambda) = (\mathbf{r} \cdot \mathbf{r}_Q)/(rr_Q)$. Substitute the above equation into Eq. (2), the potential of the body then becomes

$$\begin{aligned} U &= \int_{body} \frac{dm_\oplus}{\sqrt{r^2 + r_Q^2 - 2rr_Q \cos(\Lambda)}} \\ &= \int_{body} \frac{dm_\oplus}{r \sqrt{1 + \alpha^2 - 2\alpha \cos(\Lambda)}} \end{aligned} \quad (4)$$

Where $\alpha = r_Q/r$. The parameter (α) will always be $\alpha < 1$ because the spacecraft (point) is outside of the central body ($r > R_\oplus$). Thus, we can use the binomial theorem to expand the denominator of the potential in a series.

$$\frac{1}{\sqrt{1 - 2\alpha \cos(\Lambda) + \alpha^2}} = \frac{1}{\sqrt{1 + x}} = \sum_{l=0}^{\infty} \alpha^l P_l[\cos(\Lambda)] \quad (5)$$

This expression is a series of *Legendre polynomials*¹, P_l . Setting $\gamma = \cos(\Lambda)$ for convenience, Rodriguez's formula gives the *conventional* Legendre polynomials,

$$\begin{aligned} P_l[\gamma] &= \frac{1}{2^l l!} \frac{d^l(\gamma^2 - 1)^l}{d\gamma^l} \\ P_l[\gamma] &= \frac{1}{2^l} \sum_{j=0}^{\frac{l}{2}} \frac{(-1)^j (2l - 2j)!}{j!(l - j)!(l - 2j)!} \gamma^{l-2j} \end{aligned} \quad (6)$$

The first six Legendre polynomials are,

$$\begin{aligned} P_0[\gamma] &= 1 \\ P_1[\gamma] &= \gamma \\ P_2[\gamma] &= \frac{1}{2}(3\gamma^2 - 1) \\ P_3[\gamma] &= \frac{1}{2}(5\gamma^3 - 3\gamma) \\ P_4[\gamma] &= \frac{1}{8}(35\gamma^4 - 30\gamma^2 + 3) \\ P_5[\gamma] &= \frac{1}{8}(63\gamma^5 - 70\gamma^3 + 15\gamma) \end{aligned} \quad (7)$$

Incorporating the concept of Legendre polynomials, we can write the potential as

$$U = \frac{G}{r} \int_{body} \sum_{l=0}^{\frac{l}{2}} \alpha^l P_l[\cos(\Lambda)] dm_\oplus \quad (8)$$

The following section provides a detailed look into the implication of the potential equation. By combining the *MacCullagh's formula* and the *addition theorem* of spherical harmonics, we can thus derive a more comprehensive potential function to determine our acceleration for numerical propagations.

¹The Legendre polynomials here is defined as the coefficients in a formal expression in powers of α of the *generating function*. Generating functions is a way of encoding an infinite sequence of numbers by treating them as the coefficients of a formal power series.




Figure 1: **Deriving the Gravitational Potential.** Schematic of Earth being composed with point masses, m_Q . Note that the frame is in ITRF().

II-B MacCullagh Geometric Method

II-B1 MacCullagh's formula

Using *MacCullagh's* approach, we express the potential in Eq. (8) as below. This permits us to evaluate each gravitational component and to exploit the implication of each gravitational potential term.

$$U = \underbrace{U_0}_{\text{two body}} + \underbrace{U_1}_{\text{centre of mass}} + \underbrace{U_2}_{\text{moments of inertia}} + \dots \quad (9)$$

1) U_0 , First Potential Term

Using the first Legendre polynomial terms for the first term P_0 ,

$$U_0 = \frac{G}{r} \int dm_{\oplus} = \frac{Gm_{\oplus}}{r} = \frac{\mu}{r} \quad (10)$$

Conveniently, what the the first term, U_0 , assumed is that the central body is spherically symmetric and homogeneous, which is what we often referred as the *two-body* potential.

2) U_1 , Second Potential Term

using the Legendre polynomial term by letting $l = 1$ and the expression of $\cos(\Lambda)$ from dot product manipulation,

$$\begin{aligned} U_1 &= \frac{G}{r} \int \cos(\Lambda) \alpha dm_{\oplus} = \frac{G}{r} \int \frac{x\xi + y\eta + z\zeta}{r^2} dm_{\oplus} \\ &= \frac{G}{r^3} \left(x \int \xi dm_{\oplus} + y \int \eta dm_{\oplus} + z \int \zeta dm_{\oplus} \right) \end{aligned} \quad (11)$$

Each integral (for ξ , η , and ζ) represents the component of the potential to account for a displacement of the centre of mass from the origin of the coordinate frame. Even if the origin coincides with the geometric centre of the body, it may not coincide with the centre

of mass. However, if the origin coincides with the centre of mass,

$$\int \xi dm_{\oplus} = \int \eta dm_{\oplus} = \int \zeta dm_{\oplus} = 0 \quad (12)$$

Thus, $U_1 = 0$ because the origin we chose is at the centre of mass. The expressions above are definitions of the centre of mass.

3) U_2 , Third Potential Term

The third term in the expansion is

$$U_2 = \frac{1}{2} \frac{G}{r^3} \int (3\gamma^2 - 1) r_Q^2 dm_{\oplus} \quad (13)$$

and since we defined $\gamma = \cos(\Lambda)$, we can substitute it in

$$U_2 = \frac{G}{2r^3} \underbrace{\int 2r_Q^2 dm_{\oplus}}_{A+B+C} - \frac{G}{2r^3} \underbrace{\int 3r_Q^2 \sin^2(\Lambda) dm_{\oplus}}_I \quad (14)$$

where A, B and C represents the *moments of inertia* about the three coordinate axes

$$\begin{aligned} A &= \int (\eta^2 + \zeta^2) dm_{\oplus} \\ B &= \int (\xi^2 + \zeta^2) dm_{\oplus} \\ C &= \int (\xi^2 + \eta^2) dm_{\oplus} \end{aligned} \quad (15)$$

And the remaining term is the *polar moment of inertia*, I , that refers to any point r_Q

$$I = \int r_Q^2 \sin^2(\Lambda) dm_{\oplus} \quad (16)$$

Thus, we can rewrite the second potential term as

$$U_2 = \frac{G}{2r^3} (A + B + C - 3I) \quad (17)$$

By substituting all the terms (U_0-U_2) and assuming the coordinate frame's origin is at the centre of mass, gives us the aspherical potential

$$U = \frac{Gm_{\oplus}}{r} + \frac{G}{2r^3}(A + B + C - 3I) \quad (18)$$

This is the *MacCullagh's formula*. Note that evaluating the integrals defining A, B, C and I presents practical problem requiring detailed knowledge on Earth's mass distribution, which is often quite unrealistic to assume uniform density. Therefore, we need to incorporate an alternative by introducing a geometric approach for Eq. (8).

II-B2 Geometric Approach (Spherical Harmonics)

The *geometric approach* examines the potential function and incorporate spherical trigonometry to develop an equation to find the angle, Λ , directly. Note that a satellite's latitude will always be a *geocentric* value ($\phi_{gc_{sat}}$). If we substitute $\phi_{gc_{sat}}$ with Λ , and using cosine law of spherical trigonometry,

$$\begin{aligned} \cos(\Lambda) &= \cos(90^\circ - \phi_{gc_Q})\cos(90^\circ - \phi_{gc_{sat}}) \\ &+ \sin(90^\circ - \phi_{gc_Q})\sin(90^\circ - \phi_{gc_{sat}})\cos(\lambda_Q - \lambda_{sat}) \end{aligned} \quad (19)$$

Reduction gives us

$$\begin{aligned} \cos(\Lambda) &= \sin(\phi_{gc_Q})\sin(\phi_{gc_{sat}}) \\ &+ \cos(\phi_{gc_Q})\cos(\phi_{gc_{sat}})\cos(\lambda_Q - \lambda_{sat}) \end{aligned} \quad (20)$$

The *addition theorem* of spherical harmonics provides a way to expand the expression for Λ into Eq. (8)

$$\begin{aligned} P_l[\cos(\Lambda)] &= P_l[\sin(\phi_{gc_Q})]P_l[\sin(\phi_{gc_{sat}})] \\ &+ 2 \sum_{m=1}^l \frac{(l-m)!}{(l+m)!} \{A_{l,m}A'_{l,m} + B_{l,m}B'_{l,m}\} \end{aligned} \quad (21)$$

$$\begin{aligned} A_{l,m} &= P_{l,m}[\sin(\phi_{gc_Q})]\cos(m\lambda_Q) \\ A'_{l,m} &= P_{l,m}[\sin(\phi_{gc_{sat}})]\cos(m\lambda_{sat}) \\ B_{l,m} &= P_{l,m}[\sin(\phi_{gc_Q})]\sin(m\lambda_Q) \\ B'_{l,m} &= P_{l,m}[\sin(\phi_{gc_{sat}})]\sin(m\lambda_{sat}) \end{aligned}$$

where the summation introduces " l " and " m " indices as *degree* and *order*, respectively. The term $P_{l,m}$ are called the

associated Legendre functions, $P_{l,m}$. Notice that for zero order ($m = 0$), the associated Legendre functions are simply the conventional Legendre polynomials. The general form were given by Lambeck as the following

$$P_{l,m}[\gamma] = \frac{1}{2^l l!} (1 - \gamma^2)^{m/2} \frac{d^{l+m}}{d\gamma^{l+m}} (\gamma^2 - 1)^l$$

l	m	$P_{l,m}(\sin\phi)$
0	0	1
1	0	$\sin(\phi_{gc_{sat}})$
1	1	$\cos(\phi_{gc_{sat}})$
2	0	$\frac{1}{2}\{3\sin^2(\phi_{gc_{sat}}) - 1\}$
2	1	$3\sin(\phi_{gc_{sat}})\cos(\phi_{gc_{sat}})$
2	2	$3\cos^2(\phi_{gc_{sat}})$
3	0	$\frac{1}{2}\{5\sin^3(\phi_{gc_{sat}}) - 3\sin(\phi_{gc_{sat}})\}$
3	1	$\frac{1}{2}\cos(\phi_{gc_{sat}})\{15\sin^2(\phi_{gc_{sat}}) - 3\}$
3	2	$15\cos^2(\phi_{gc_{sat}})\sin(\phi_{gc_{sat}})$
3	3	$15\cos^3(\phi_{gc_{sat}})$

By separating all the terms that are independent of the satellite's location in Eq. (8), we can arrive at a solution that isolates terms which depends only on the central body and those which relate that satellite's position. First we define two new variables, $C'_{l,m}$ and $S'_{l,m}$, as follow

$$\begin{aligned} C'_{l,m} &= \int_{body} r_Q^l \frac{(l-m)!}{(l+m)!} P_{l,m}[\sin(\phi_{gc_{sat}})] \cos(m\lambda_Q) dm_{\oplus} \\ S'_{l,m} &= \int_{body} r_Q^l \frac{(l-m)!}{(l+m)!} P_{l,m}[\sin(\phi_{gc_{sat}})] \sin(m\lambda_Q) dm_{\oplus} \end{aligned} \quad (22)$$

The two coefficients, $C'_{l,m}$ and $S'_{l,m}$, represent the mathematical modelling for the Earth's shape using spherical harmonics. The special case for the *zonal harmonics* is

$$C'_{l,0} = \int_{body} r_Q^l P_l[\sin(\phi_{gc_Q})] dm_{\oplus} \quad (23)$$

which uses the conventional Legendre polynomials. Note that $S'_{l,0}$ is zero. Substitute these values into the potential in Eq. (8) we can get the following expression for our potential.

$$\begin{aligned} U &= \frac{G}{r} \sum_{l=0}^{\infty} \frac{P_l[\sin(\phi_{gc_{sat}})]}{r^l} C'_{l,0} + \frac{G}{r} \sum_{l=1}^{\infty} \sum_{m=1}^l \frac{P_{l,m}[\sin(\phi_{gc_{sat}})]}{r^l} \{C'_{l,m} \cos(m\lambda_{sat}) + S'_{l,m} \sin(m\lambda_{sat})\} \\ &= \frac{\mu}{r} \sum_{l=0}^{\infty} P_l[\sin(\phi_{gc_{sat}})] \left(\frac{R_{\oplus}}{r}\right)^l C'_{l,0} + \frac{\mu}{r} \sum_{l=1}^{\infty} \sum_{m=1}^l P_{l,m}[\sin(\phi_{gc_{sat}})] \left(\frac{R_{\oplus}}{r}\right)^l \{C'_{l,m} \cos(m\lambda_{sat}) + S'_{l,m} \sin(m\lambda_{sat})\} \end{aligned} \quad (24)$$

A unit analysis suggest removing units from the above result. This leads to the C and S coefficients. We can undimensionalise the gravitational coefficients.

$$C'_{l,m} = C_{l,m} R_{\oplus}^l m_{\oplus}, \quad S'_{l,m} = S_{l,m} R_{\oplus}^l m_{\oplus} \quad (25)$$

The International Astronomical Union (1961) has adopted a form for the aspherical potential. First, a double summation encompasses the associated Legendre polynomials, $P_{l,m}$, and the harmonic coefficients, and $S_{l,0} = 0$. Notice that both summation indices begin at zero. Recall, as part of the derivation, if the centre of the coordinate system coincides with the attracting body's centre of mass, the coefficients

where R_{\oplus} and m_{\oplus} are radius and mass of the earth, respectively. Using the undimensionalise gravitational parameter, we can express gravitational potential as

$C_{1,0}$, $C_{1,1}$, and $S_{1,1}$ are all zero ($S_{1,0}$ is also zero by definition). This result corresponds to $l = 1$ in MacCullagh's approach. It leads to another common form of this relation, which separates the spherical potential and therefore requires us to adjust the summation limits. The 0^{th} term is within the μ/r term, and the first-degree terms are zero.

$$\begin{aligned} U &= \frac{\mu}{r} \sum_{l=0}^{\infty} \sum_{m=0}^l \left(\frac{R_{\oplus}}{r} \right)^l P_{l,m}[\sin(\phi_{gc_{sat}})] \{C_{l,m} \cos(m\lambda_{sat}) + S_{l,m} \sin(m\lambda_{sat})\} \\ &= \frac{\mu}{r} \left[1 + \sum_{l=2}^{\infty} \sum_{m=0}^l \left(\frac{R_{\oplus}}{r} \right)^l P_{l,m}[\sin(\phi_{gc_{sat}})] \{C_{l,m} \cos(m\lambda_{sat}) + S_{l,m} \sin(m\lambda_{sat})\} \right] \end{aligned} \quad (26)$$

This expression describes the gravitational attraction resulting from the irregular distribution of the Earth's mass using a potential function.

indices l and m , *degree* and *order*, determine lines on the sphere along which the functions vanish. The harmonic coefficients $C_{l,m}$, $S_{l,m}$ represent the relative importance of these mass distributions.

II-C Spherical Harmonics

The trigonometric argument of the Legendre polynomials constitutes surface's **spherical harmonics**, for they are periodic on the surface of a unit sphere. When the surface's spherical harmonics are divided by r^{l+1} , they're usually called solid spherical harmonics. The Sturm-Liouville theorem states that the solid spherical harmonics are eigenfunctions that constitute an independent basis for the gravitational model. In essence, they are Fourier series. The

II-C1 Zonal Harmonics

Zonal harmonics are defined by zeroth order ($m = 0$), where the dependence of the potential on longitude vanished and the field is symmetrical about the polar axis, and are simply **bands of latitude**.

Zonal harmonic coefficients are commonly defined in literature with the notation

$$J_l = C_{l,0} \quad (27)$$




Figure 2: **Zonal Harmonics**. The band reflects the Earth's oblateness, the shading region indicates regions of additional mass and the numbers link regions between the views().

II-C2 Sectoral Harmonics

Sectoral Harmonics occur when $l = m$ and represent bands of longitude (shown in Fig. (3)). The associated Legendre functions, $P_{l,l}[\sin(\phi_{gc_{sat}})]$ only have zeros at the pole, $\phi = \pm 90^\circ$, therefore, the only changes in sign are due

to the trigonometric functions $\cos(m\lambda)$ and $\sin(m\lambda)$. On the other hand the terms $[\cos(l\lambda)$ and $\sin(l\lambda)]$ are zero for $2l$ different values of λ . Hence the lines along which the functions $[\cos(l\lambda)$ and $\sin(l\lambda)]$ times $P_{l,l}[\sin(\phi_{gc_{sat}})]$ vanish are meridians of longitude. Each sector represents l positive (shaded) and l negative (white) mass concentrations.




Figure 3: **Sectoral Harmonics**. The band reflects the Earth's oblateness, the shading region indicates regions of additional mass and the numbers link regions between the views().

II-C3 Tesseral Harmonics

Tesseral harmonics refer to the cases for $l \neq m \neq 0$, the physical depiction is a specific region (tile) of the earth, for the sphere is divided up to a checkerboard array (shown

in Fig. (4)). The number of circles of latitude along which $P_{l,m}[\sin(\phi_{gc_{sat}})]$ is zero is equal to $(l - m)$, whereas the terms $[\cos(l\lambda)$ and $\sin(l\lambda)]$ vanish along $2m$ meridians of longitude. These zero lines represent the centre of the latitude/longitude bands.




Figure 4: **Tesseral Harmonics**. The band reflects the Earth's oblateness, the shading region indicates regions of additional mass and the numbers link regions between the views().

II-D Acceleration From Gravitational Potential

Since we've developed our gravitational potential that describes the uneven distribution of mass for Earth, we then proceed to take the gradient (partial derivative) of the potential and obtain the acceleration term and apply it during orbit propagation. Using matrix differentiation with $\vec{r} = r_I \hat{I} + r_J \hat{J} + r_K \hat{K}$ in the ITRF² frame, we can directly

determine the partial derivative of the aspherical potential function. Legendre functions are often differentiated in spherical coordinate, therefore, chain rule is required to find acceleration in cartesian coordinate, as shown below.

$$\vec{a} = \frac{\partial U}{\partial r} \left(\frac{\partial r}{\partial \vec{r}} \right)^T + \frac{\partial U}{\partial \phi_{gc_{sat}}} \left(\frac{\partial \phi_{gc_{sat}}}{\partial \vec{r}} \right)^T + \frac{\partial u}{\partial \lambda_{sat}} \left(\frac{\partial \lambda_{sat}}{\partial \vec{r}} \right)^T \quad (28)$$

²Note that accelerations are usually calculated in body-fixed frame (ECEF, or ITRF), whereas numerical integration are done in an inertial frame (ECI, or GCRF)

The derivatives of the position vector also proceed directly. Notice that the three partial derivatives below are unit vectors.

Then using the nonspherical portion of Eq. (26), we can calculate the respective partial derivatives we need for our acceleration

$$\begin{aligned}\frac{\partial \mathbf{r}}{\partial \vec{\mathbf{r}}} &= \frac{\vec{\mathbf{r}}^T}{r} \\ \frac{\partial \phi_{gc_{sat}}}{\partial \vec{\mathbf{r}}} &= \frac{1}{\sqrt{r_I^2 + r_J^2}} \left(-\frac{\vec{\mathbf{r}}^T \mathbf{r}_K}{r^2} + \frac{\partial r_K}{\partial \vec{\mathbf{r}}} \right) \\ \frac{\partial \lambda_{sat}}{\partial \vec{\mathbf{r}}} &= \frac{1}{r_I^2 + r_J^2} \left(r_I \frac{\partial r_J}{\partial \vec{\mathbf{r}}} - r_J \frac{\partial r_I}{\partial \vec{\mathbf{r}}} \right)\end{aligned}\quad (29)$$

$$\begin{aligned}\frac{\partial U}{\partial r} &= -\frac{\mu}{r} \sum_{l=2}^{\infty} \sum_{m=0}^l \left(\frac{R_{\oplus}}{r} \right)^l (l+1) P_{l,m}[\text{SIN}(\phi_{gc_{sat}})] \left\{ C_{l,m} \text{COS}(m\lambda_{sat}) + S_{l,m} \text{SIN}(m\lambda_{sat}) \right\} \\ \frac{\partial U}{\partial \phi_{gc_{sat}}} &= \frac{\mu}{r} \sum_{l=2}^{\infty} \sum_{m=0}^l \left(\frac{R_{\oplus}}{r} \right)^l \left\{ P_{l,m+1}[\text{SIN}(\phi_{gc_{sat}})] - m \text{TAN}(\phi_{gc_{sat}}) P_{l,m}[\text{SIN}(\phi_{gc_{sat}})] \right\} \left\{ C_{l,m} \text{COS}(m\lambda_{sat}) + S_{l,m} \text{SIN}(m\lambda_{sat}) \right\} \\ \frac{\partial U}{\partial \lambda_{sat}} &= \frac{\mu}{r} \sum_{l=2}^{\infty} \sum_{m=0}^l \left(\frac{R_{\oplus}}{r} \right)^l P_{l,m}[\text{SIN}(\phi_{gc_{sat}})] \left\{ S_{l,m} \text{COS}(m\lambda_{sat}) - C_{l,m} \text{SIN}(m\lambda_{sat}) \right\} \\ &\Downarrow\end{aligned}\quad (30)$$

$$\begin{aligned}a_I &= \left\{ \frac{1}{r} \frac{\partial U}{\partial r} - \frac{r_K}{r^2 \sqrt{r_I^2 + r_J^2}} \frac{\partial U}{\partial \phi_{gc_{sat}}} \right\} r_I - \left\{ \frac{1}{r_I^2 + r_J^2} \frac{\partial U}{\partial \lambda_{sat}} \right\} r_J - \frac{\mu r}{r^3} \\ a_J &= \left\{ \frac{1}{r} \frac{\partial U}{\partial r} - \frac{r_K}{r^2 \sqrt{r_I^2 + r_J^2}} \frac{\partial U}{\partial \phi_{gc_{sat}}} \right\} r_J + \left\{ \frac{1}{r_I^2 + r_J^2} \frac{\partial U}{\partial \lambda_{sat}} \right\} r_I - \frac{\mu r}{r^3} \\ a_K &= \frac{1}{r} \frac{\partial U}{\partial r} r_K + \frac{\sqrt{r_I^2 + r_J^2}}{r^2} \frac{\partial U}{\partial \phi_{gc_{sat}}} - \frac{\mu r}{r^3}\end{aligned}\quad (31)$$
