

# **FRST302: Forest Genetics**

## **Lecture 1.1: Classical Genetics and its Molecular Mechanisms**

# Outline for Today

- Short history of genetics
- Mendel's laws
- Chromosomes

**What is genetics?**


**What is genetics?**

**Genetics is the study of genes**, of variation and heredity across all branches of the tree of life

# *What are the major questions in genetics?*



# How can we apply a knowledge of genetics?



<sup>1</sup>Modified from Figure 1 - Zuidhof et al. 2014

# History of Genetics

Humans have probably pondered inheritance for all history:

- For much of history, the mechanisms of inheritance were basically unknown
- The inheritance of acquired characteristics was widely accepted for much of history (from Hippocrates to Aristotal to Lamarck)

# History of Genetics

Humans have probably pondered inheritance for all history:

- For much of history, the mechanisms of inheritance were basically unknown
- The inheritance of acquired characteristics was widely accepted for much of history (from Hippocrates to Aristotal to Lamarck)
- *Early microscopists thought that they had seen small humans inhabiting sperm cells!*






# History of Genetics

At the time Darwin came around, the dominant theory was **blending inheritance**

- The notion that an offspring's traits are simply the average of the parents' traits.
- This is intuitively appealing - continuously varying traits are often intermediate between their parents
- There is one big problem with blending inheritance!



*What's the big problem with  
blending inheritance?*

Blending inheritance only really makes sense when you are thinking about continuously varying traits

But different modes of variation are common:

Blending inheritance only really makes sense when you are thinking about continuously varying traits

But different modes of variation are common:

- Continuous - traits measured on a numerical scale (e.g. height, diameter, chlorophyll fluorescence)

Blending inheritance only really makes sense when you are thinking about continuously varying traits

But different modes of variation are common:

- Continuous - traits measured on a numerical scale (e.g. height, diameter, chlorophyll fluorescence)
- Discrete - traits that exhibit categorical differences (e.g. different leaf forms, distinct flower colour)


Blending inheritance only really makes sense when you are thinking about continuously varying traits

But different modes of variation are common:

- Continuous - traits measured on a numerical scale (e.g. height, diameter, chlorophyll fluorescence)
- Discrete - traits that exhibit categorical differences (e.g. different leaf forms, distinct flower colour)
- Ordinal - discrete traits with some informative order (e.g. high, medium and low shade tolerance)

# Particulate Inheritance


Through careful experimentation analysing discrete traits in peas, Franciscan Friar Gregor Mendel found evidence supporting a model of particulate inheritance



Mmmmm...  
Peas Peas Peas Peas Peas

# Mendel's Crosses

Mendel examined variation and inheritance of several discrete characteristics of pea plants






# Mendel's Crosses


Garden peas are capable of self-fertilization, so Mendel was able to generate "true" lines of peas that exhibit a particular trait/phenotype

- Crossing lines produces an F<sub>1</sub> generation
- The patterns of variation among the F<sub>2</sub> generations led Mendel to develop his notions of particulate inheritance



# Particulate Inheritance


- Proposed in 1865 and 1866
- 6-7 years after Darwin's Theory of Evolution
- Represents the foundation of modern genetics
- ASD



More peas please

# Reconciling the Mendelians and the Biometricians

**Distribution of Trait Values Assuming  
100 Genes Segregating According to Mendelian Inheritance**



# Particulate Inheritance

**Below is the R code to make the figures on the infinitesimal model - feel free to play around with it**

```
# Demonstrate the distribution of trait values for a quantitative trait  
# Under Mendelian segregation for an arbitrary number of genes  
# Assumes random mating, constant effect sizes, constant allele frequencies
```

```
nGenes = 100  
alleleFrequency = 0.2  
popSize = 5000  
effectSize = 1
```

```
hist(  
  replicate(popSize ,  
    sum( 1 * rbinom(nGenes, 2, alleleFrequency) ) ),  
  col = "#e69b99",  
  xlab= " Trait - Value",  
  main= paste(" Distribution - of - Trait - Values - Assuming\n", nGenes ,  
    " Genes - Segregating - According - to - Mendelian - Inheritance" ),  
  breaks = 40)
```