# Physics 5150

### Homework Set # 3

# Due 5 pm Thursday 2/8/2018

## **SOLUTIONS**

#### Problem 1: Thermonuclear fusion in stars

A 5 billion year old main-sequence star called Sol, belonging to the spectral type G2, has a mass  $M_{\odot} = 2 \times 10^{33}$  g and a total luminosity  $L_{\odot} = 4 \times 10^{33}$  erg/s. The star is powered by the pp chain thermonuclear reaction, the net result of which is that 4 protons are turned into a He<sup>4</sup> nucleus (an  $\alpha$ -particle) releasing 26.7 MeV of energy in the process.

- (a) Assuming that the star's parameters have not changed appreciably since it was born, how much hydrogen (by mass) has the star consumed over its lifetime?
- (b) In addition to producing helium, each pp chain reaction also produces 2 neutrinos that fly out of the star freely, without interacting with anything (they carry only 2% of the total fusion energy, so the corresponding energy losses can be neglected). How many neutrinos fly each second through a 1 cm<sup>2</sup> thumbnail of John Bahcall standing on a seashore on a small rocky planet called Terra, 150 mln km from the star?

#### **Solution:**

The number of pp-chain reactions happening in the Sun each second is

$$\dot{N} = \frac{L_{\odot}}{\epsilon} = 0.93 \times 10^{38} \,\text{sec}^{-1} \,.$$
 (1)

(a) In each reaction 4 hydrogen nuclei (protons) are consumed. Therefore, the total hydrogen consumption rate is

$$\dot{M}_H = 4m_p \dot{N} \simeq 6.3 \times 10^{11} \,\text{kg/sec} \,.$$
 (2)

Then, the amount of hydrogen consumed over the star's lifetime is

$$\Delta M_H = \dot{M}_H \times 5 \cdot 10^9 \times 3.15 \times 10^7 \,\text{sec} \simeq 10^{29} \,\text{kg}$$
. (3)

(b) The total neutrino production rate is

$$\dot{N}_{\nu} = 2\dot{N} \simeq 1.9 \times 10^{38} \,\text{sec}^{-1}$$
. (4)

The neutrino flux at R = 1 AU then is:

$$\Gamma_{\nu} = \frac{\dot{N}_{\nu}}{4\pi R^2} \simeq 6.6 \times 10^{10} \,\mathrm{cm}^{-2} \,\mathrm{sec}^{-1}$$
 (5)

### Problem 2: Fusion power plant

What is the annual tritium mass consumption of a continuously  $(24 \times 7)$  operating D-T thermonuclear fusion reactor with the total ( $\alpha$ -particles plus neutrons) fusion power output of 1 GW?

#### **Solution:**

Fusion energy released in 1 reaction is  $\epsilon \simeq 17\,\mathrm{MeV} \simeq 2.7 \times 10^{-12}\,\mathrm{J}$ . The number of reactions per unit time is then:

$$\dot{N} = \frac{P}{\epsilon} \simeq 3.7 \times 10^{20} \,\text{sec}^{-1} \,.$$
 (6)

The number of tritium nuclei consumed each year is

$$N_T = \dot{N} \times 3.16 \cdot 10^7 \,\text{sec/yr} \simeq 1.16 \times 10^{28} \,,$$
 (7)

and hence the annual tritium mass consumption is

$$M_T = N_T \times 3m_p \simeq 58 \,\mathrm{kg} \,. \tag{8}$$

### <u>Problem 3:</u> Tokamak plasma density:

Equilibrium and stability consideration limit the total (electrons + ions) plasma pressure inside a tokamak to be less than about 10% of the pressure of the confining magnetic field. What is the maximum plasma electron density in a tokamak with a magnetic field of 5 Tesla and temperature  $T_i = T_e = 10 \text{ keV}$ ?

#### **Solution:**

The maximum plasma pressure is  $P_{\text{max}} = 0.1 P_{\text{mag}} = 0.1 B^2 / 8\pi$ . The corresponding plasma density, related to the plasma pressure via  $P_{\text{max}} = n_{\text{max}}(T_e + T_i)$ , is

$$n_{\rm max} = 0.1 \, \frac{B^2}{8\pi (T_c + T_i)} \simeq 3 \times 10^{14} \, {\rm cm}^{-3} \, .$$