

Topological Energy Condensate Theory (TECT): 30

Jusang Lee*
Independent Researcher, Incheon, Korea
(Dated: March, 2026)

I. ACTUAL PDE COMPLETION PROTOCOL

A. Honest status

The present goal is not to rebuild the theorem chain from scratch. It is to populate the already-closed carrier-level certificate with actual branch data extracted from the full microscopic TECT linearization:

proof complete, certificate incomplete.

Equivalently, the remaining frontier is:

finite microscopic audit, not new theorem-building.

B. Actual PDE meaning

The phrase “complete the theory from the actual PDE” should be understood as follows.
One starts from the actual Bloch–Floquet linearized operator around the intended BCC background:

$$L(k).$$

Then one extracts the finite reduced carrier data by the chain

$L(k) \longrightarrow K_i := \partial_{k_i} L|_{G_*} \longrightarrow P_* K_i P_* \longrightarrow M_{3i} \longrightarrow \lambda_{\parallel},$

together with the transverse derivative data at the same shell point.

Thus the remaining work is not “solve the whole nonlinear PDE in one shot”, but rather:

extract the branch-dependent finite coefficients from the actual microscopic operator.

C. Gate 1: remote-gap numerical insertion

The first certificate slot is

$\Delta_{\text{bench},\rho}^{\text{fin}} > \eta_{R,\rho}.$

Hence the first numerical task is:

$$\Delta_{\text{bench},\rho}^{\text{fin}} = \min(\text{finite benchmark channel gaps}), \quad \eta_{R,\rho} = \text{remote-sector correction bound},$$

and then evaluate the signed margin

$\mathfrak{G}_1 := \Delta_{\text{bench},\rho}^{\text{fin}} - \eta_{R,\rho}.$

Gate 1 passes iff

$$\mathfrak{G}_1 > 0.$$

* jusanglee@kakao.com

D. Gate 2: actual longitudinal extraction

The exact intervalley witness is

$$\mathcal{M}_{ab}^{(JJ)} = \int_{\text{cell}} d^3x \, e^{-iG_* \cdot x} \overline{u_a^+(x)} \left(\hat{n}^i K_i^{(\Pi, J)}(x) \right) u_b^-(x).$$

The reduced longitudinal coefficient is

$$m_3 = -\frac{1}{2\Gamma_J} \left(\mathcal{M}_{11}^{(JJ)} - \mathcal{M}_{22}^{(JJ)} \right).$$

At the refined locked working level this is compressed to

$$\Delta_{00} = \eta_+^\dagger \Sigma_3 \eta_-, \quad L_* = m_{\parallel} u_{\parallel} = \frac{m_{\parallel}}{2} (u_{11} - u_{22}).$$

A practical sufficient condition is

$$|\Delta_{00} \alpha_X L_* A_{G_*}| > |R_N| \implies \lambda_{\parallel} \neq 0.$$

Thus the numerical longitudinal margin is

$$\mathfrak{G}_2 := |\Delta_{00} \alpha_X L_* A_{G_*}| - |R_N|.$$

Gate 2 passes iff

$$\mathfrak{G}_2 > 0.$$

E. Gate 3: derivative-resolved transverse audit

The honest transport-side transverse witnesses are

$$c_E = \frac{1}{4} \text{Tr}(\Sigma_1 \mathcal{T}_1 + \Sigma_2 \mathcal{T}_2), \quad c_O = \frac{1}{4} \text{Tr}(\Sigma_2 \mathcal{T}_1 - \Sigma_1 \mathcal{T}_2),$$

with

$$\alpha = c_E I_{\theta}, \quad \beta = c_O I_{\theta}.$$

Hence the transport-side pass condition is

$$(c_E, c_O) \neq (0, 0).$$

On the explicit folded-Bragg working branch, the strongest practical sufficient test is

$$A_{G_*}(C_1 \cdot G_*) \neq 0, \quad A_{G_*}(C_2 \cdot G_*) \neq 0.$$

The calibration layer is

$$(C_E, C_O) \neq (0, 0).$$

Thus a practical transverse numerical audit computes

$$\mathfrak{G}_{3,\text{tr}} := |c_E| + |c_O|, \quad \mathfrak{G}_{3,\text{cal}} := |C_E| + |C_O|.$$

Gate 3 passes iff

$$\mathfrak{G}_{3,\text{tr}} > 0, \quad \mathfrak{G}_{3,\text{cal}} > 0.$$

F. Gate 4: constant-sector commutator audit

The residual generator is

$$Q_* = \tau_3 \otimes s_3.$$

The full corrected microscopic constant-sector test is

$$[Q_*, H_{\Psi\Psi}^{\text{const}}] = 0, \quad [Q_*, \delta H_{\text{Class II}}^{\text{const}}] = 0, \quad [Q_*, \Sigma(0,0)] = 0.$$

In the present working truncation,

$$\mathcal{H}_{\text{extra}} = \emptyset \implies V(0) = 0 \implies \Sigma(0,0) = 0,$$

so Gate 4 is largely reduced to a finite low-block constant audit.

G. Final actual certificate

The fully populated actual branch certificate is

$$\Delta_{\text{bench},\rho}^{\text{fin}} > \eta_{R,\rho},$$

$$\lambda_{\parallel} \neq 0,$$

$$(c_E, c_O) \neq (0,0), \quad (C_E, C_O) \neq (0,0),$$

$$[Q_*, H_{\Psi\Psi}^{\text{const}}] = 0, \quad [Q_*, \delta H_{\text{Class II}}^{\text{const}}] = 0, \quad [Q_*, \Sigma(0,0)] = 0.$$

Therefore

$$\text{Gate 1} \wedge \text{Gate 2} \wedge \text{Gate 3} \wedge \text{Gate 4} \implies \text{isolated protected anisotropic massless Dirac carrier.}$$

II. FINAL SUMMARY

Yes: the next correct target is full completion from the actual microscopic PDE data.

But “actual PDE completion” now means:

$$\text{extract finite branch-dependent coefficients from the full microscopic linearized operator}$$

rather than

$$\text{re-solve the whole nonlinear PDE from scratch.}$$

Numerical extraction is not optional. It is exactly the remaining task.

The practical computational order is:

$$\text{Gate 1} \rightarrow \text{Gate 2} \rightarrow \text{Gate 3} \rightarrow \text{Gate 4.}$$

Among these, Gate 4 is already nearly closed in the present working truncation, so the main numerical burden is the actual evaluation of:

$$\Delta_{\text{bench},\rho}^{\text{fin}}, \quad \eta_{R,\rho}, \quad \Delta_{00}, \quad L_*, \quad A_{G_*}, \quad (c_E, c_O), \quad (C_E, C_O).$$

III. GATE 1 ACTUAL PDE CLOSURE ON THE EXPLICIT CANONICAL PURE-EVEN BRANCH

IV. SHORT EXPLANATION

At the present stage, Gate 1 is no longer a conceptual theorem problem. For the explicit canonical pure-even branch, it is reduced to one explicit finite inequality.

The exact support condition remains

$$\Delta_{\text{bench},\rho}^{\text{fin}} > \eta_{R,\rho}.$$

On the explicit branch

$$T_\star = T^1, \quad \eta = z_0, \quad \Lambda_O^{(0)} = 0,$$

the sharpest currently available practical form is obtained by inserting the same-shell non-enriched scalar lower bound and the fully inserted residual budget.

V. 1. MASTER GATE-1 CONDITION

The benchmark side is

$$\Delta_{\text{bench},\rho}^{\text{fin}} = \min \left\{ 4K - |r_\star|, \sqrt{M_\sigma^2 + c_\sigma^2(q_\star - \rho)^2}, q_\star^2 \left| (\lambda + \mu) + \frac{g^2}{M_\sigma^2 + c_\sigma^2 q_\star^2} \right|, \Xi_{\text{non}}^{(1)}(\rho) \right\}.$$

The residual side satisfies

$$\eta_{\text{rem},\rho} \leq 2 \left[\kappa \rho (|C_E| + |C_O|) + \frac{1}{M_X^2} (\alpha_X^2 + |\alpha_X \beta_X| + \beta_X^2) \right] + 2C_{\text{high},\text{non}}\rho + \sum_{a \in \mathcal{A}_{\text{act}}} K_a \rho^{r_a} + \sum_{a \in \mathcal{A}_{\text{act}}^{(H)}} \frac{\tilde{K}_a}{M_a^{\sigma_a}} \rho^{r_a} + \eta_{\text{diag},\rho}.$$

Hence a sufficient actual branch Gate-1 condition is

$$\min \left\{ 4K - |r_\star|, \sqrt{M_\sigma^2 + c_\sigma^2(q_\star - \rho)^2}, q_\star^2 \left| (\lambda + \mu) + \frac{g^2}{M_\sigma^2 + c_\sigma^2 q_\star^2} \right|, \Xi_{\text{non}}^{(1)}(\rho) \right\} > \eta_{\text{rem},\rho}.$$

VI. 2. SAME-SHELL NON-ENRICHED SCALAR INSERTION

The usable same-shell benchmark entry is reduced to

$$\Delta_{\text{non-enr},\rho}^{(1)} - \varepsilon_{\text{non-enr},\rho}^{(1)} \geq \Xi_{\text{non}}^{(1)}(\rho),$$

with

$$\Xi_{\text{non}}^{(1)}(\rho) := \left| -\mu^2 + \frac{Zq_\star^2}{4} - \frac{Yq_\star^4}{16} + 12uA^2 \right| - 4uA^2 - 2a_K \mathcal{S}_{\text{diag}}^{EE} - \varepsilon_{\text{small-}p,\rho}^{(1)} - \varepsilon_{\text{non-enr},\rho}^{(1)},$$

and

$$a_K = \frac{\beta_X^2}{M_X^2} (\Lambda_E^{(0)})^2.$$

Therefore the non-enriched benchmark is positive once

$$\left| -\mu^2 + \frac{Zq_\star^2}{4} - \frac{Yq_\star^4}{16} + 12uA^2 \right| - 4uA^2 > 2a_K \mathcal{S}_{\text{diag}}^{EE} + \varepsilon_{\text{small-}p,\rho}^{(1)} + \varepsilon_{\text{non-enr},\rho}^{(1)}.$$

VII. 3. SMALL- p DRIFT BUDGET

For the minimal same-shell truncation $N = 1$,

$$H_{\text{non-enr}}^{(1)}(p) = \begin{pmatrix} d_+(p) & m(p) \\ m(p) & d_-(p) \end{pmatrix},$$

with expansions

$$d_{\pm}(p) = d_{\pm}^{(0)} + p_i a_{\pm,i} + R_{\pm}^{(2)}(p), \quad m(p) = m_0 + p_i b_i + R_m^{(2)}(p).$$

Define

$$A_+ = \left(\sum_i |a_{+,i}|^2 \right)^{1/2}, \quad A_- = \left(\sum_i |a_{-,i}|^2 \right)^{1/2}, \quad B = \left(\sum_i |b_i|^2 \right)^{1/2}.$$

Then

$$\varepsilon_{\text{small-}p,\rho}^{(1)} \leq \left[(\rho A_+ + C_+ \rho^2)^2 + (\rho A_- + C_- \rho^2)^2 + 2(\rho B + C_m \rho^2)^2 \right]^{1/2}.$$

At leading order,

$$\varepsilon_{\text{small-}p,\rho}^{(1)} = \rho L_{\text{non}}^{(1)} + O(\rho^2), \quad L_{\text{non}}^{(1)} = \sqrt{A_+^2 + A_-^2 + 2B^2}.$$

VIII. 4. ACTUAL DERIVATIVE COEFFICIENTS

On the explicit pure-even branch,

$$a_{+,i} = \delta_{i\parallel} q_* \left(Z - \frac{Y q_*^2}{2} \right) - a_K \ell_{+,i}^{EE},$$

$$a_{-,i} = -\delta_{i\parallel} q_* \left(Z - \frac{Y q_*^2}{2} \right) - a_K \ell_{-,i}^{EE},$$

$$b_i = -a_K \ell_{0,i}^{EE}.$$

Hence

$$A_+ \leq \left| q_* \left(Z - \frac{Y q_*^2}{2} \right) \right| + a_K L_+, \quad A_- \leq \left| q_* \left(Z - \frac{Y q_*^2}{2} \right) \right| + a_K L_-, \quad B \leq a_K L_0.$$

Therefore the small- p drift is itself reduced to the finite overlap-derivative vectors

$$\ell_{+,i}^{EE}, \quad \ell_{-,i}^{EE}, \quad \ell_{0,i}^{EE}.$$

IX. 5. PURE-EVEN CANONICAL SHARPENING

If the residual transport side is also taken pure-even and one adopts the canonical normalization

$$C_{\parallel} = C_E = C_O = 1,$$

then

$$|C_E| + |C_O| \longrightarrow |C_E| = 1,$$

so the non-wrong estimate sharpens to

$$\eta_{\text{non,wrong},\rho} \leq \kappa\rho + \frac{1}{M_X^2} (\alpha_X^2 + |\alpha_X\beta_X| + \beta_X^2).$$

Thus the sharpest explicit sufficient Gate-1 inequality becomes

$$\min \left\{ 4K - |r_*|, \sqrt{M_\sigma^2 + c_\sigma^2(q_* - \rho)^2}, q_*^2 \left| (\lambda + \mu) + \frac{g^2}{M_\sigma^2 + c_\sigma^2 q_*^2} \right|, \Xi_{\text{non}}^{(1)}(\rho) \right\}$$

$$> 2 \left[\kappa\rho + \frac{1}{M_X^2} (\alpha_X^2 + |\alpha_X\beta_X| + \beta_X^2) \right] + 2C_{\text{high,non}}\rho + \sum_{a \in \mathcal{A}_{\text{act}}} K_a \rho^{r_a} + \sum_{a \in \mathcal{A}_{\text{act}}^{(H)}} \frac{\tilde{K}_a}{M_a^{\sigma_a}} \rho^{r_a} + \eta_{\text{diag},\rho}.$$

X. 6. WHAT IS ACTUALLY LEFT

So Gate 1 is now reduced to the finite insertion of the branch constants

$$\mathcal{S}_{\text{diag}}^{EE}, \ell_{+,i}^{EE}, \ell_{-,i}^{EE}, \ell_{0,i}^{EE}, \varepsilon_{\text{non-enr},\rho}^{(1)},$$

$$C_{\text{high,non}}, K_a, \tilde{K}_a, r_a, \sigma_a, \eta_{\text{diag},\rho},$$

together with the already explicit bulk-shell parameters

$$K, r_*, M_\sigma, c_\sigma, q_*, \lambda + \mu, g, \mu, Z, Y, u, A, \Lambda_E^{(0)}, \alpha_X, \beta_X, M_X.$$

Therefore:

$$\text{Gate 1 is structurally closed and executable,}$$

but

$$\text{its PASS/FAIL stamp still requires the actual branch constants above.}$$

XI. FINAL SUMMARY

At the actual PDE stage, the correct next move is Gate 1 finite insertion.

For the explicit canonical pure-even branch, the sharpest available Gate-1 sufficient inequality is

$$\min \left\{ 4K - |r_*|, \sqrt{M_\sigma^2 + c_\sigma^2(q_* - \rho)^2}, q_*^2 \left| (\lambda + \mu) + \frac{g^2}{M_\sigma^2 + c_\sigma^2 q_*^2} \right|, \Xi_{\text{non}}^{(1)}(\rho) \right\} > \eta_{\text{rem},\rho},$$

with

$$\Xi_{\text{non}}^{(1)}(\rho) = \left| -\mu^2 + \frac{Zq_*^2}{4} - \frac{Yq_*^4}{16} + 12uA^2 \right| - 4uA^2 - 2a_K \mathcal{S}_{\text{diag}}^{EE} - \varepsilon_{\text{small-}p,\rho}^{(1)} - \varepsilon_{\text{non-enr},\rho}^{(1)},$$

$$a_K = \frac{\beta_X^2}{M_X^2} (\Lambda_E^{(0)})^2.$$

The residual side is

$$\eta_{\text{rem},\rho} \leq 2 \left[\kappa\rho + \frac{1}{M_X^2} (\alpha_X^2 + |\alpha_X \beta_X| + \beta_X^2) \right] + 2C_{\text{high,non}}\rho + \sum_{a \in \mathcal{A}_{\text{act}}} K_a \rho^{r_a} + \sum_{a \in \mathcal{A}_{\text{act}}^{(H)}} \frac{\tilde{K}_a}{M_a^{\sigma_a}} \rho^{r_a} + \eta_{\text{diag},\rho},$$

after the pure-even canonical sharpening.

Thus Gate 1 is fully reduced to finite branch constants. No new theorem is needed.

What still blocks the final numerical PASS/FAIL stamp is only the missing insertion of

$$\mathcal{S}_{\text{diag}}^{EE}, \quad \ell_{+,i}^{EE}, \ell_{-,i}^{EE}, \ell_{0,i}^{EE}, \quad \varepsilon_{\text{non-enr},\rho}^{(1)}, \quad C_{\text{high,non}}, \quad K_a, \tilde{K}_a, r_a, \sigma_a, \quad \eta_{\text{diag},\rho}.$$

XII. ACTUAL PDE COMPLETION ON THE EXPLICIT CANONICAL PURE-EVEN BRANCH

XIII. SHORT EXPLANATION

For the current explicit canonical pure-even branch

$$T_\star = T^1, \quad \eta = z_0, \quad \Lambda_O^{(0)} = 0,$$

the reduced infrared PDE is already completed at the practical working-branch level.

What is still missing is not the PDE form itself, but the final finite audit needed for the full unconditional microscopic certificate.

XIV. 1. EXACT COMPLETED INFRARED PDE

The reduced infrared symbol is

$$A_1(p) = \lambda_{\parallel} p_{\parallel} \sigma_3 + \alpha(p_1 \sigma_1 + p_2 \sigma_2) + \beta(-p_2 \sigma_1 + p_1 \sigma_2). \quad (1)$$

On the explicit pure-even working branch, one has

$$\beta = 0, \quad (2)$$

so the PDE simplifies to

$$\left[i\partial_t - \lambda_{\parallel} \sigma_3 (-i\partial_{\parallel}) - \alpha(\sigma_1 (-i\partial_1) + \sigma_2 (-i\partial_2)) \right] \chi = 0. \quad (3)$$

XV. 2. EXACT COEFFICIENT DICTIONARY

In Branch B, the exact overlap dictionary is

$$\lambda_{\parallel} = \frac{\lambda_c}{2} I_{\theta'^3}, \quad \alpha = I_{\theta} \Lambda_E, \quad \beta = I_{\theta} \Lambda_O. \quad (4)$$

Using the explicit defect overlaps,

$$I_{\theta'^3} = -\frac{3\sqrt{3}\pi}{256}\theta_0^3\kappa^3, \quad I_\theta = -\frac{\sqrt{3}\pi}{16}\theta_0\kappa, \quad (5)$$

the pure-even working realization gives

$$\lambda_{\parallel} = \frac{3\sqrt{3}\pi}{512} \frac{\alpha_X \beta_X}{M_X^2} \theta_0^3 \kappa^3, \quad (6)$$

$$\alpha = \frac{\sqrt{3}\pi}{16} \frac{\beta_X}{M_X^2} \theta_0 \kappa, \quad \beta = 0. \quad (7)$$

Therefore the explicit actual PDE is

$$\left[i\partial_t - \frac{3\sqrt{3}\pi}{512} \frac{\alpha_X \beta_X}{M_X^2} \theta_0^3 \kappa^3 \sigma_3(-i\partial_{\parallel}) - \frac{\sqrt{3}\pi}{16} \frac{\beta_X}{M_X^2} \theta_0 \kappa (\sigma_1(-i\partial_1) + \sigma_2(-i\partial_2)) \right] \chi = 0. \quad (8)$$

XVI. 3. NUMERICAL COEFFICIENT SHEET

Numerically,

$$\lambda_{\parallel} \approx 0.03188319195 \frac{\alpha_X \beta_X}{M_X^2} (\theta_0 \kappa)^3, \quad (9)$$

$$\alpha \approx 0.34008738079 \frac{\beta_X}{M_X^2} (\theta_0 \kappa), \quad \beta = 0. \quad (10)$$

Since

$$\det M = \lambda_{\parallel} (\alpha^2 + \beta^2),$$

the explicit pure-even branch gives

$$\det M = \frac{9\sqrt{3}\pi^3}{131072} \frac{\alpha_X \beta_X^3}{M_X^6} \theta_0^5 \kappa^5, \quad (11)$$

hence numerically

$$\det M \approx 0.003687591698 \frac{\alpha_X \beta_X^3}{M_X^6} (\theta_0 \kappa)^5. \quad (12)$$

Therefore a sufficient nondegeneracy condition is

$$\alpha_X \neq 0, \quad \beta_X \neq 0, \quad \theta_0 \kappa \neq 0. \quad (13)$$

XVII. 4. GATE 2 AND GATE 3 FINITE PASS SHEET

The current carrier-side longitudinal witness is already reduced to

$$\Delta_{00} = \eta_+^\dagger \Sigma_3 \eta_-, \quad L_* = m_{\parallel} u_{\parallel}. \quad (14)$$

The practical sufficient condition is

$$|\Delta_{00} \alpha_X L_* A_{G_*}| > |R_N| \implies \lambda_{\parallel} \neq 0. \quad (15)$$

In the ultra-minimal truncation $R_N = 0$, it is enough to verify

$$\Delta_{00} \neq 0, \quad L_* \neq 0, \quad \alpha_X \neq 0, \quad A_{G_*} \neq 0. \quad (16)$$

For the transverse sector, the pass condition remains

$$(c_E, c_O) \neq (0, 0), \quad (17)$$

equivalently, on the explicit folded-Bragg branch,

$$A_{G_*}(C_1 \cdot G_*) \neq 0, \quad A_{G_*}(C_2 \cdot G_*) \neq 0. \quad (18)$$

XVIII. 5. GATE 1: WHAT STILL BLOCKS FULL UNCONDITIONAL CLOSURE

The sharpest currently available Gate-1 inequality on this branch is

$$\Delta_{\text{bench}, \rho}^{\text{fin}} = \min \left\{ 4K - |r_*|, \sqrt{M_\sigma^2 + c_\sigma^2 (q_* - \rho)^2}, q_*^2 \left| (\lambda + \mu) + \frac{g^2}{M_\sigma^2 + c_\sigma^2 q_*^2} \right|, \Xi_{\text{non}}^{(1)}(\rho) \right\}, \quad (19)$$

with

$$\Xi_{\text{non}}^{(1)}(\rho) = \left| -\mu^2 + \frac{Zq_*^2}{4} - \frac{Yq_*^4}{16} + 12uA^2 \right| - 4uA^2 - 2a_K \mathcal{S}_{\text{diag}}^{EE} - \varepsilon_{\text{small-}p, \rho}^{(1)} - \varepsilon_{\text{non-enr}, \rho}^{(1)}, \quad (20)$$

and residual side

$$\eta_{\text{rem}, \rho} \leq 2 \left[\kappa \rho (|C_E| + |C_O|) + \frac{1}{M_X^2} (\alpha_X^2 + |\alpha_X \beta_X| + \beta_X^2) \right] + 2C_{\text{high}, \text{non}} \rho + \sum_{a \in \mathcal{A}_{\text{act}}} K_a \rho^{r_a} + \sum_{a \in \mathcal{A}_{\text{act}}^{(H)}} \frac{\tilde{K}_a}{M_a^{\sigma_a}} \rho^{r_a} + \eta_{\text{diag}, \rho}. \quad (21)$$

Thus the full unconditional Gate-1 pass condition is

$$\Delta_{\text{bench}, \rho}^{\text{fin}} > \eta_{\text{rem}, \rho}. \quad (22)$$

At the present stage, the only genuinely missing part is the finite insertion of

$$\mathcal{S}_{\text{diag}}^{EE}, \quad \varepsilon_{\text{small-}p, \rho}^{(1)}, \quad \varepsilon_{\text{non-enr}, \rho}^{(1)}, \quad C_{\text{high}, \text{non}}, \quad K_a, \tilde{K}_a, r_a, \sigma_a, \quad \eta_{\text{diag}, \rho}. \quad (23)$$

XIX. FINAL SUMMARY

Therefore:

$$\text{the actual reduced infrared PDE is already completed on the present working branch,} \quad (24)$$

with explicit coefficients

$$\lambda_{\parallel}, \alpha, \beta$$

and explicit numerical prefactors.

However,

$$\boxed{\text{the full unconditional microscopic certificate is not yet numerically complete,}} \quad (25)$$

because Gate 1 still requires the insertion of finitely many branch-specific constants.

Equivalently,

$$\boxed{\text{working-branch PDE: closed,} \quad \text{full unconditional audit: still awaiting finite insertion.}} \quad (26)$$

XX. HOW TO EXTRACT ALL REMAINING FINITE INSERTION CONSTANTS

XXI. SHORT EXPLANATION

At the present stage, all unresolved quantities are already reduced to finite explicit audits. No new structural theorem is needed.

The remaining constants fall into four computational classes:

$$\boxed{\text{finite overlap sums,} \quad \text{derivatives at } p = 0, \quad \text{finite residual norms/tails,} \quad \text{constant-block commutators.}}$$

Therefore the right question is no longer

“Can these constants be defined?”

but rather

“Which finite operator or overlap must be evaluated for each constant?”

XXII. 1. GLOBAL MAP OF THE REMAINING UNRESOLVED SLOTS

The genuinely remaining finite audit slots are:

$$\boxed{\Delta_{\text{non-enr}, \rho}^{(N)} - \varepsilon_{\text{non-enr}, \rho}^{(N)}, \quad \eta_{R, \rho},}$$

$$\boxed{\Xi_{JJ} \quad \text{or equivalently} \quad L_* = m_{\parallel} u_{\parallel},}$$

$$\boxed{(c_E, c_O), \quad (C_E, C_O),}$$

$$\boxed{[Q_*, H_{\Psi\Psi}^{\text{const}}], \quad [Q_*, \delta H_{\text{Class II}}^{\text{const}}], \quad [Q_*, \Sigma(0, 0)].}$$

So every remaining object is already a finite insertion target.

XXIII. 2. GATE 1: SAME-SHELL NON-ENRICHED BENCHMARK

A. 2.1. Core strategy

The same-shell non-enriched slot is already reduced to the finite Hermitian block

$$\boxed{H_{\text{non-enr}}^{(N)}(p).}$$

For $N = 1$,

$$\boxed{H_{\text{non-enr}}^{(1)}(p) = \begin{pmatrix} d_+(p) & m(p) \\ m(p) & d_-(p) \end{pmatrix}.}$$

Thus the extraction protocol is:

1. compute $H_{\text{non-enr}}^{(N)}(0)$;
2. compute its spectral distance from zero;
3. bound the perturbation on $|p| \leq \rho$.

This yields

$$\Delta_{\text{non-enr},\rho}^{(N)} \geq \Delta_0^{(N)} - \varepsilon_{\text{small-}p,\rho}^{(N)},$$

and after truncation control,

$$\Delta_{\text{non-enr},\rho}^{(N)} - \varepsilon_{\text{non-enr},\rho}^{(N)}.$$

B. 2.2. How to extract $\mathcal{S}_{\text{diag}}^{EE}$

At $p = 0$, the explicit pure-even branch gives

$$d_+^{(0)} = E_{z_0}^{\text{dir}}(0) - a_K \mathcal{S}_{++}^{EE}, \quad d_-^{(0)} = E_{z_0}^{\text{dir}}(0) - a_K \mathcal{S}_{--}^{EE},$$

$$m_0 = B_{z_0}(0) - a_K \mathcal{S}_{+-}^{EE}, \quad a_K = \frac{\beta_X^2}{M_X^2} (\Lambda_E^{(0)})^2.$$

If the residual is valley-symmetric,

$$\mathcal{S}_{++}^{EE} = \mathcal{S}_{--}^{EE} =: \mathcal{S}_{\text{diag}}^{EE}.$$

Therefore $\mathcal{S}_{\text{diag}}^{EE}$ is extracted by:

1. build the finite EE -overlap matrix of the non-enriched same-shell sector at $p = 0$;
2. evaluate its diagonal valley entries $\mathcal{S}_{++}^{EE}, \mathcal{S}_{--}^{EE}$;
3. in the symmetric branch, identify their common value as $\mathcal{S}_{\text{diag}}^{EE}$;
4. otherwise keep the asymmetric pair and use the larger one in the bound.

Operationally, this is a finite cell-integral / finite Fourier-sum computation.

C. 2.3. How to extract $\ell_{+,i}^{EE}, \ell_{-,i}^{EE}, \ell_{0,i}^{EE}$

The derivative overlap vectors are defined by

$$\ell_{+,i}^{EE} = \partial_{p_i} \mathcal{S}_{++}^{EE}(p)|_{p=0}, \quad \ell_{-,i}^{EE} = \partial_{p_i} \mathcal{S}_{--}^{EE}(p)|_{p=0},$$

$$\ell_{0,i}^{EE} = \partial_{p_i} \mathcal{S}_{+-}^{EE}(p)|_{p=0}.$$

So these are obtained by:

1. construct the p -dependent overlap sums $\mathcal{S}_{++}^{EE}(p), \mathcal{S}_{--}^{EE}(p), \mathcal{S}_{+-}^{EE}(p)$;
2. differentiate them with respect to p_i ;
3. evaluate at $p = 0$.

These are genuine first-derivative data, not new unknown parameters. Their norms are

$$L_+ = \left(\sum_i |\ell_{+,i}^{EE}|^2 \right)^{1/2}, \quad L_- = \left(\sum_i |\ell_{-,i}^{EE}|^2 \right)^{1/2}, \quad L_0 = \left(\sum_i |\ell_{0,i}^{EE}|^2 \right)^{1/2}.$$

D. 2.4. How to extract C_+, C_-, C_m and $\varepsilon_{\text{small-}p, \rho}^{(1)}$

Near $p = 0$,

$$d_{\pm}(p) = d_{\pm}^{(0)} + p_i a_{\pm, i} + R_{\pm}^{(2)}(p), \quad m(p) = m_0 + p_i b_i + R_m^{(2)}(p),$$

with

$$|R_+^{(2)}(p)| \leq C_+ \rho^2, \quad |R_-^{(2)}(p)| \leq C_- \rho^2, \quad |R_m^{(2)}(p)| \leq C_m \rho^2.$$

Hence C_+, C_-, C_m are second-derivative remainder bounds. They are extracted by:

1. compute or bound the Hessians $\partial_{p_i} \partial_{p_j} d_{\pm}(p)$, $\partial_{p_i} \partial_{p_j} m(p)$ on $|p| \leq \rho$;
2. take uniform sup norms;
3. convert them into quadratic Taylor constants C_+, C_-, C_m .

Then

$$\boxed{\varepsilon_{\text{small-}p, \rho}^{(1)} \leq \left[(\rho A_+ + C_+ \rho^2)^2 + (\rho A_- + C_- \rho^2)^2 + 2(\rho B + C_m \rho^2)^2 \right]^{1/2}},$$

where

$$A_{\pm} = \left(\sum_i |a_{\pm, i}|^2 \right)^{1/2}, \quad B = \left(\sum_i |b_i|^2 \right)^{1/2}.$$

Thus $\varepsilon_{\text{small-}p, \rho}^{(1)}$ is not an extra free constant: it is produced from first- and second-derivative data.

E. 2.5. How to extract $\varepsilon_{\text{non-enr}, \rho}^{(N)}$

The truncation error enters through the upgrade

$$\varepsilon_{\text{small-}p, \rho}^{(N)} + \varepsilon_{\text{non-enr}, \rho}^{(N)} < \Delta_{\text{non-enr}, 0}^{(N)}.$$

The same-shell notes reduce this to:

1. a finite heavy-overlap sum $\mathcal{S}_{\text{diag}}^{EE, (N)}$,
2. a finite tail bound $\mathcal{R}_{\text{diag}}^{EE, (N)}$,
3. a finite small- p plus truncation budget.

Therefore $\varepsilon_{\text{non-enr}, \rho}^{(N)}$ is extracted by one of two equivalent routes:

a. Route A: rigorous tail route. Estimate the omitted shell/basis tail

$$\mathcal{R}_{\text{diag}}^{EE, (N)} = \sum_{\text{omitted modes}} |\cdots|,$$

using decay/power-counting of the overlap kernel, and turn it into a norm bound.

b. Route B: monotone numerical stabilization route. Compute the same-shell benchmark at $N, N+1, N+2, \dots$, verify convergence, and use the last increment as a rigorous-enough posteriori upper bound only if monotone decay is proven. Without monotone control, this route is heuristic, not theorem-level.

So the fully honest theorem-level route is Route A.

XXIV. 3. GATE 1: RESIDUAL REMOTE BUDGET

A. 3.1. Master decomposition

The residual side is already reduced to

$$\eta_{R,\rho} \leq \|E_\rho\|_2 + \eta_{\text{tail},\rho} + \eta_{\text{diag},\rho},$$

or more weakly

$$\eta_{R,\rho} \leq \max\{S_{\text{vol}}, S_{\text{high}}, S_{\text{non}}, S_{\text{wrong}}\} + \eta_{\text{tail},\rho} + \eta_{\text{diag},\rho}.$$

Thus there are three extraction tasks.

B. 3.2. How to extract the pair budgets and $C_{\text{high},\text{non}}$

The remaining six pair budgets are

$$\eta_{\text{vol},\text{high},\rho}, \eta_{\text{vol},\text{non},\rho}, \eta_{\text{vol},\text{wrong},\rho}, \eta_{\text{high},\text{non},\rho}, \eta_{\text{high},\text{wrong},\rho}, \eta_{\text{non},\text{wrong},\rho}.$$

For the dominant explicit branch, the inserted notes already give

$$\eta_{\text{non},\text{wrong},\rho} \leq \kappa\rho(|C_E| + |C_O|) + \frac{1}{M_X^2} (\alpha_X^2 + |\alpha_X\beta_X| + \beta_X^2),$$

and

$$\eta_{\text{high},\text{non},\rho} \leq C_{\text{high},\text{non}}\rho.$$

Therefore $C_{\text{high},\text{non}}$ is extracted as:

$$C_{\text{high},\text{non}} = \sup_{0 < |p| \leq \rho_0} \frac{\|E_{\text{high},\text{non}}(p)\|}{|p|},$$

or via a corresponding row-sum norm bound on that pair block.

So the method is:

1. isolate the high/non block of the remote mixing matrix;
2. differentiate or directly bound it for small p ;
3. divide by $|p|$ and take the uniform supremum.

C. 3.3. How to extract $K_a, \tilde{K}_a, r_a, \sigma_a$

The tail theorem already has the form

$$\eta_{\text{tail},\rho} \leq \sum_{a \in \mathcal{A}_{\text{act}}} K_a \rho^{r_a} + \sum_{a \in \mathcal{A}_{\text{act}}^{(H)}} \frac{\tilde{K}_a}{M_a^{\sigma_a}} \rho^{r_a}, \quad r_a \geq 1.$$

Hence these constants are extracted by:

1. enumerate the active omitted channel families $a \in \mathcal{A}_{\text{act}}, \mathcal{A}_{\text{act}}^{(H)}$;
2. determine the low- p scaling exponent r_a of each family;
3. determine the heavy-scale suppression exponent σ_a where applicable;
4. compute the corresponding prefactors $K_a, \tilde{K}_a$ from the channelwise norm bounds.

This is a pure power-counting + family-enumeration problem.

D. 3.4. How to extract $\eta_{\text{diag},\rho}$

The diagonal correction budget is

$$\eta_{\text{diag},\rho} = \sup_{|p| \leq \rho} \|R_R^{\text{diag}}(p)\|.$$

So the method is literal: construct the diagonal remainder $R_R^{\text{diag}}(p)$ after the chosen truncation/splitting, then take the operator-norm supremum on the ball $|p| \leq \rho$.

This is usually the cleanest object to compute numerically after the symbolic subtraction has been fixed.

XXV. 4. GATE 2: LONGITUDINAL FINITE DATA

A. 4.1. How to extract Ξ_{JJ} and L_*

The exact longitudinal witness is already reduced to

$$\mathcal{M}_{ab}^{(JJ)} = \int_{\text{cell}} d^3x e^{-iG_* \cdot x} \overline{u_a^+(x)} (\hat{n}^i K_i^{(\Pi,J)}(x)) u_b^-(x).$$

Then

$$m_3 = -\frac{1}{2\Gamma_J} (\mathcal{M}_{11}^{(JJ)} - \mathcal{M}_{22}^{(JJ)}).$$

At the refined locked working level this is compressed to

$$L_* = m_{\parallel} u_{\parallel}.$$

Therefore the extraction protocol is:

1. build the pure- JJ intervalley block $K_i^{(\Pi,J)}$;
2. compute the cell overlap matrix $\mathcal{M}_{ab}^{(JJ)}$;
3. take the diagonal difference $\mathcal{M}_{11}^{(JJ)} - \mathcal{M}_{22}^{(JJ)}$;
4. identify Ξ_{JJ} or equivalently $L_* = m_{\parallel} u_{\parallel}$.

B. 4.2. How to extract Δ_{00}

The working-level scalar is

$$\Delta_{00} = \eta_+^\dagger \Sigma_3 \eta_-.$$

So one must:

1. solve the low-slot eigenproblem at $p = 0$;
2. obtain the normalized valley basis vectors η_+, η_- ;
3. evaluate the single 2×2 Pauli overlap $\eta_+^\dagger \Sigma_3 \eta_-$.

This is a finite eigensystem computation, not a PDE problem anymore.

C. 4.3. How to extract A_{G_*}

On the explicit folded-Bragg branch, A_{G_*} is the Fourier amplitude of the relevant G_* -harmonic. Hence it is extracted by:

$$A_{G_*} = \hat{A}(G_*) = \int_{\text{cell}} d^3x e^{-iG_* \cdot x} A(x),$$

or by the equivalent discrete Fourier coefficient in the Bloch/Fourier basis.

XXVI. 5. GATE 3: TRANSVERSE WITNESSES AND CALIBRATION

A. 5.1. How to extract (c_E, c_O)

The honest transverse frontier is

$$\partial_{p_\nu} \hat{\Sigma}_{JK}(G_*; p)|_{p=0}, \quad \partial_{p_\nu} \hat{\Sigma}_{KK}(G_*; p)|_{p=0}, \quad \nu = 1, 2.$$

The Pauli witnesses are

$$c_E = \frac{1}{4} \text{Tr}(\Sigma_1 \mathcal{T}_1 + \Sigma_2 \mathcal{T}_2), \quad c_O = \frac{1}{4} \text{Tr}(\Sigma_2 \mathcal{T}_1 - \Sigma_1 \mathcal{T}_2).$$

Hence the extraction protocol is:

1. compute the derivative-resolved shell blocks at $Q = G_*$;
2. assemble the transverse shell tensors $\mathcal{T}_1, \mathcal{T}_2$;
3. project onto the Σ_1, Σ_2 Pauli basis;
4. read off c_E, c_O .

Inside the explicit folded-Bragg Class II branch, this compresses to

$$\hat{T}_{(II)}^\alpha(G_*) = -\frac{\alpha_X \beta_X}{M_X^2} \sum_A \sum_{Q+Q'=G_*} \hat{J}_i^A(Q) \hat{K}_{i,\alpha}^A(Q'),$$

and at first-shell dominant level to

$$\hat{T}_{(II)}^\alpha(G_*) \propto i A_{G_*}(C_\alpha \cdot G_*).$$

So on the working branch, a fast sufficient test is just to evaluate

$$A_{G_*}(C_1 \cdot G_*), \quad A_{G_*}(C_2 \cdot G_*).$$

B. 5.2. How to extract (C_E, C_O)

The Bloch calibration layer is exact:

$$\alpha = C_E g_E, \quad \beta = C_O g_O.$$

The overlap formulas are

$$C_E = \frac{1}{4} \text{Tr}(\sigma_1 M_{(E)}^1 + \sigma_2 M_{(E)}^2),$$

$$C_O = \frac{1}{4} \text{Tr}(\sigma_2 M_{(O)}^1 - \sigma_1 M_{(O)}^2).$$

So the extraction method is:

1. build the normalized E - and O -unit inputs;
2. compute the folded-Bragg overlap matrices $M_{(E)}^1, M_{(E)}^2, M_{(O)}^1, M_{(O)}^2$;
3. evaluate the two trace formulas above.

This is a pure finite calibration computation.

XXVII. 6. GATE 4: CONSTANT-SECTOR COMMUTATOR AUDIT

The remaining Gate-4 test is

$$\boxed{[Q_*, H_{\Psi\Psi}^{\text{const}}] = 0, \quad [Q_*, \delta H_{\text{Class II}}^{\text{const}}] = 0, \quad [Q_*, \Sigma(0, 0)] = 0,} \quad Q_* = \tau_3 \otimes s_3.$$

Thus the extraction protocol is:

1. project the constant block of the microscopic operator to the 4×4 carrier space;
2. write each constant correction in the explicit $\tau_a \otimes s_b$ basis;
3. compute the three commutators with Q_* ;
4. verify that all coefficients outside the Q_* -commuting algebra vanish.

So Gate 4 is not an unknown-constant problem at all: it is a finite matrix-algebra check.

XXVIII. 7. WHAT IS RIGOROUS, WHAT IS ONLY PRACTICAL

The following routes are theorem-level rigorous:

1. finite overlap/cell-integral evaluation;
2. $p = 0$ derivative evaluation;
3. operator-norm or row-sum supremum bounds;
4. explicit tail power-counting bounds;
5. direct commutator computation.

The following can be useful numerically but are not theorem-level by themselves:

1. plain convergence-by-eye in N ;
2. fitting unknown exponents from data without a prior bound;
3. assuming valley symmetry or canonical normalization without verifying them.

XXIX. FINAL SUMMARY

Yes: in principle all remaining insertion constants can be extracted.

More precisely, they reduce to:

$$\boxed{\mathcal{S}_{\text{diag}}^{EE}, \ell_{\pm, i}^{EE}, \ell_{0, i}^{EE}, C_{\pm}, C_m \longleftarrow \text{same-shell overlap sums and their first/second } p\text{-derivatives;}} \quad (27)$$

$$\boxed{\varepsilon_{\text{non-enr}, \rho}^{(N)}, K_a, \tilde{K}_a, r_a, \sigma_a, C_{\text{high, non}}, \eta_{\text{diag}, \rho} \longleftarrow \text{finite residual norms} + \text{tail power counting;}} \quad (28)$$

$$\boxed{\Delta_{00}, \Xi_{JJ}, L_*, A_{G_*} \longleftarrow \text{finite low-slot eigensystem} + \text{cell overlaps} + \text{Fourier coefficient};} \quad (29)$$

$$\boxed{(c_E, c_O), (C_E, C_O) \longleftarrow \text{derivative-resolved shell audit} + \text{Bloch calibration traces};} \quad (30)$$

$$\boxed{[Q_*, H_{\Psi\Psi}^{\text{const}}], [Q_*, \delta H_{\text{Class II}}^{\text{const}}], [Q_*, \Sigma(0, 0)] \longleftarrow \text{finite } 4 \times 4 \text{ commutator audit.}} \quad (31)$$

So the remaining project is algorithmic, not conceptual.

XXX. GATE 1 SAME-SHELL EXTRACTION LAYER: EXACT FORMULAS FOR

$$\mathcal{S}_{\text{diag}}^{EE}, \ell_{\pm, i}^{EE}, \ell_{0, i}^{EE}, C_{\pm}, C_m$$

XXXI. SHORT EXPLANATION

All six unresolved same-shell constants are generated by one common heavy-resolvent object:

$$\boxed{\mathcal{S}_{ab}^{EE}(p)}.$$

Once $\mathcal{S}_{ab}^{EE}(p)$ is available, we obtain

$$\mathcal{S}_{\text{diag}}^{EE} = \mathcal{S}_{++}^{EE}(0) = \mathcal{S}_{--}^{EE}(0),$$

$$\ell_{+, i}^{EE} = \partial_{p_i} \mathcal{S}_{++}^{EE}(0), \quad \ell_{-, i}^{EE} = \partial_{p_i} \mathcal{S}_{--}^{EE}(0), \quad \ell_{0, i}^{EE} = \partial_{p_i} \mathcal{S}_{+-}^{EE}(0),$$

and the quadratic drift constants C_+, C_-, C_m from second derivatives.

Thus the whole problem reduces to one source-resolvent construction.

XXXII. 1. LOW-SLOT STATES AND HEAVY BLOCK

Let

$$|\phi_+(p)\rangle, \quad |\phi_-(p)\rangle$$

denote the same-shell non-enriched low states in the common Bloch sector $k_* + p$. Let

$$\Pi_H$$

be the projector onto the heavy complementary space, and let

$$\mathbb{O}_H(p)$$

be the heavy block of the reference microscopic operator restricted to $\Pi_H \mathcal{H}$.

Assume

$$\boxed{\mathbb{O}_H(p) \geq \Delta_H > 0}$$

for $|p| \leq \rho$, so that the heavy resolvent exists:

$$\boxed{R_H(p) := \mathbb{O}_H(p)^{-1} \quad \text{on } \Pi_H \mathcal{H}.}$$

XXXIII. 2. EVEN SOURCE VECTORS

Let $\mathcal{O}_E(p)$ denote the even heavy source operator induced by the pure-even residual KK -sector. Define the heavy source vectors

$$|v_a^E(p)\rangle := \Pi_H \mathcal{O}_E(p) |\phi_a(p)\rangle, \quad a \in \{+, -\}. \quad (32)$$

Then the exact heavy-overlap tensor is

$$\mathcal{S}_{ab}^{EE}(p) := \langle v_a^E(p), R_H(p) v_b^E(p) \rangle. \quad (33)$$

This is the common generating object for all remaining same-shell constants.

XXXIV. 3. SPECTRAL-SUM REPRESENTATION

If one diagonalizes the heavy block,

$$\mathbb{O}_H(p) |\chi_\alpha(p)\rangle = E_\alpha(p) |\chi_\alpha(p)\rangle, \quad E_\alpha(p) \geq \Delta_H,$$

and defines

$$\mathcal{E}_{a\alpha}(p) := \langle \chi_\alpha(p) | v_a^E(p) \rangle = \langle \chi_\alpha(p) | \mathcal{O}_E(p) | \phi_a(p) \rangle,$$

then

$$\mathcal{S}_{ab}^{EE}(p) = \sum_{\alpha} \frac{\mathcal{E}_{a\alpha}(p) \overline{\mathcal{E}_{b\alpha}(p)}}{E_{\alpha}(p)}. \quad (34)$$

Thus the already-used Schur form

$$-\sum_{\alpha} \frac{K_{a\alpha} \overline{K_{b\alpha}}}{E_{\alpha}}$$

is exactly the spectral form of $\mathcal{S}_{ab}^{EE}(p)$.

XXXV. 4. COORDINATE-SPACE CELL-INTEGRAL FORMULA

Work in the Bloch gauge

$$\phi_a(p, x) = e^{i(k_*+p)\cdot x} u_a(p, x), \quad \chi_\alpha(p, x) = e^{i(k_*+p)\cdot x} w_\alpha(p, x),$$

with u_a, w_α periodic on the primitive cell.

Define the Bloch-conjugated even source operator

$$\mathcal{O}_E^{(k_*+p)} := e^{-i(k_*+p)\cdot x} \mathcal{O}_E(p) e^{i(k_*+p)\cdot x}.$$

Then

$$\mathcal{E}_{a\alpha}(p) = \int_{\text{cell}} d^3x \overline{w_\alpha(p, x)} (\mathcal{O}_E^{(k_*+p)} u_a(p, \cdot))(x). \quad (35)$$

Hence

$$\mathcal{S}_{ab}^{EE}(p) = \sum_{\alpha} \frac{1}{E_{\alpha}(p)} \left(\int_{\text{cell}} d^3x \overline{w_\alpha} \mathcal{O}_E^{(k_*+p)} u_a \right) \overline{\left(\int_{\text{cell}} d^3x \overline{w_\alpha} \mathcal{O}_E^{(k_*+p)} u_b \right)}. \quad (36)$$

This is the cleanest theorem-level coordinate-space formula.

XXXVI. 5. RECIPROCAL-SPACE FINITE FOURIER-SUM FORMULA

Expand the periodic parts in reciprocal lattice modes:

$$u_a(p, x) = \sum_{Q \in \mathcal{G}} u_{a,Q}(p) e^{iQ \cdot x}, \quad w_\alpha(p, x) = \sum_{Q \in \mathcal{G}} c_{\alpha,Q}(p) e^{iQ \cdot x}.$$

Likewise write the periodic operator kernel as

$$\mathcal{O}_E^{(k_*+p)}(x) = \sum_{G \in \mathcal{G}} \hat{\mathcal{O}}_E^{(k_*+p)}(G) e^{iG \cdot x}.$$

Then the overlap amplitudes become the finite/discretized Fourier sums

$$\mathcal{E}_{a\alpha}(p) = \sum_{Q, P \in \mathcal{G}} \overline{c_{\alpha,Q}(p)} \hat{\mathcal{O}}_E^{(k_*+p)}(Q - P) u_{a,P}(p). \quad (37)$$

Therefore

$$\mathcal{S}_{ab}^{EE}(p) = \sum_{\alpha} \frac{1}{E_{\alpha}(p)} \left(\sum_{Q, P} \overline{c_{\alpha,Q}} \hat{\mathcal{O}}_E(Q - P) u_{a,P} \right) \overline{\left(\sum_{Q', P'} \overline{c_{\alpha,Q'}} \hat{\mathcal{O}}_E(Q' - P') u_{b,P'} \right)}. \quad (38)$$

This is the formula to implement directly in a shell-Fourier code.

XXXVII. 6. EXTRACTION OF $\mathcal{S}_{\text{diag}}^{EE}$

At $p = 0$,

$$\mathcal{S}_{++}^{EE}(0), \quad \mathcal{S}_{--}^{EE}(0), \quad \mathcal{S}_{+-}^{EE}(0) \quad (39)$$

are computed from (33) or (34).

If valley symmetry holds,

$$\mathcal{S}_{++}^{EE}(0) = \mathcal{S}_{--}^{EE}(0) =: \mathcal{S}_{\text{diag}}^{EE}. \quad (40)$$

Thus

$$\mathcal{S}_{\text{diag}}^{EE} = \langle v_+^E(0), R_H(0) v_+^E(0) \rangle = \langle v_-^E(0), R_H(0) v_-^E(0) \rangle. \quad (41)$$

XXXVIII. 7. FIRST DERIVATIVES AND EXTRACTION OF

$$\ell_{+,i}^{EE}, \ell_{-,i}^{EE}, \ell_{0,i}^{EE}$$

Differentiate (33). Since

$$\partial_{p_i} R_H = -R_H(\partial_{p_i} \mathbb{O}_H) R_H,$$

we obtain

$$\partial_{p_i} \mathcal{S}_{ab}^{EE}(p) = \langle \partial_i v_a^E, R_H v_b^E \rangle + \langle v_a^E, R_H \partial_i v_b^E \rangle - \langle v_a^E, R_H(\partial_i \mathbb{O}_H) R_H v_b^E \rangle. \quad (42)$$

Evaluating at $p = 0$,

$$\ell_{+,i}^{EE} = \partial_{p_i} \mathcal{S}_{++}^{EE}(0), \quad \ell_{-,i}^{EE} = \partial_{p_i} \mathcal{S}_{--}^{EE}(0), \quad \ell_{0,i}^{EE} = \partial_{p_i} \mathcal{S}_{+-}^{EE}(0). \quad (43)$$

Equivalently, in source-resolvent form,

$$\ell_{+,i}^{EE} = \langle \partial_i v_+^E, R_H v_+^E \rangle + \langle v_+^E, R_H \partial_i v_+^E \rangle - \langle v_+^E, R_H(\partial_i \mathbb{O}_H) R_H v_+^E \rangle, \quad (44)$$

$$\ell_{-,i}^{EE} = \langle \partial_i v_-^E, R_H v_-^E \rangle + \langle v_-^E, R_H \partial_i v_-^E \rangle - \langle v_-^E, R_H(\partial_i \mathbb{O}_H) R_H v_-^E \rangle, \quad (45)$$

$$\ell_{0,i}^{EE} = \langle \partial_i v_+^E, R_H v_-^E \rangle + \langle v_+^E, R_H \partial_i v_-^E \rangle - \langle v_+^E, R_H(\partial_i \mathbb{O}_H) R_H v_-^E \rangle, \quad (46)$$

all evaluated at $p = 0$.

XXXIX. 8. LINEAR-SOLVE REALIZATION WITHOUT HEAVY DIAGONALIZATION

Define

$$|y_b(p)\rangle := R_H(p)|v_b^E(p)\rangle, \quad \text{so} \quad \mathbb{O}_H(p)|y_b(p)\rangle = |v_b^E(p)\rangle.$$

Then

$$\mathcal{S}_{ab}^{EE}(p) = \langle v_a^E(p), y_b(p) \rangle.$$

Differentiate once:

$$\mathbb{O}_H|y_{b,i}^{(1)}\rangle = |\partial_i v_b^E\rangle - (\partial_i \mathbb{O}_H)|y_b\rangle,$$

where

$$|y_{b,i}^{(1)}\rangle := \partial_{p_i}|y_b\rangle.$$

Therefore

$$\partial_{p_i} \mathcal{S}_{ab}^{EE} = \langle \partial_i v_a^E, y_b \rangle + \langle v_a^E, y_{b,i}^{(1)} \rangle. \quad (47)$$

This is the most implementation-friendly route: it avoids summing over the full heavy spectrum.

XL. 9. SECOND DERIVATIVES AND EXTRACTION OF

C_+, C_-, C_m

From the same-shell block,

$$d_+(p) = d_+^{(0)} + p_i a_{+,i} + R_+^{(2)}(p), \quad d_-(p) = d_-^{(0)} + p_i a_{-,i} + R_-^{(2)}(p),$$

$$m(p) = m_0 + p_i b_i + R_m^{(2)}(p),$$

with

$$a_{+,i} = \delta_{i\parallel} q_* \left(Z - \frac{Y q_*^2}{2} \right) - a_K \ell_{+,i}^{EE},$$

$$a_{-,i} = -\delta_{i\parallel} q_* \left(Z - \frac{Y q_*^2}{2} \right) - a_K \ell_{-,i}^{EE}, \quad b_i = -a_K \ell_{0,i}^{EE}.$$

Now define the Hessians

$$H_{ij}^{(+)}(p) := \partial_{p_i} \partial_{p_j} d_+(p), \quad H_{ij}^{(-)}(p) := \partial_{p_i} \partial_{p_j} d_-(p), \quad H_{ij}^{(m)}(p) := \partial_{p_i} \partial_{p_j} m(p).$$

A rigorous Taylor bound gives

$$C_+ := \frac{1}{2} \sup_{|p| \leq \rho} \|H^{(+)}(p)\|_{\text{op}}, \quad C_- := \frac{1}{2} \sup_{|p| \leq \rho} \|H^{(-)}(p)\|_{\text{op}}, \quad C_m := \frac{1}{2} \sup_{|p| \leq \rho} \|H^{(m)}(p)\|_{\text{op}}. \quad (48)$$

Equivalently,

$$|R_+^{(2)}(p)| \leq C_+ \rho^2, \quad |R_-^{(2)}(p)| \leq C_- \rho^2, \quad |R_m^{(2)}(p)| \leq C_m \rho^2.$$

Since

$$d_+(p) = E_{z_0}^{\text{dir}}(p) - a_K \mathcal{S}_{++}^{EE}(p), \quad d_-(p) = E_{z_0}^{\text{dir}}(p) - a_K \mathcal{S}_{--}^{EE}(p),$$

$$m(p) = B_{z_0}(p) - a_K \mathcal{S}_{+-}^{EE}(p),$$

we get

$$H_{ij}^{(+)} = \partial_{ij} E_{z_0}^{\text{dir}} - a_K \partial_{ij} \mathcal{S}_{++}^{EE}, \quad (49)$$

$$H_{ij}^{(-)} = \partial_{ij} E_{z_0}^{\text{dir}} - a_K \partial_{ij} \mathcal{S}_{--}^{EE}, \quad (50)$$

$$H_{ij}^{(m)} = \partial_{ij} B_{z_0} - a_K \partial_{ij} \mathcal{S}_{+-}^{EE}. \quad (51)$$

Thus the quadratic constants are entirely determined by the second p -derivatives of the same overlap tensor.

XLI. 10. SECOND-DERIVATIVE LINEAR-SOLVE ROUTE

Differentiate the heavy solve a second time. Let

$$|y_{b,ij}^{(2)}\rangle := \partial_{p_i} \partial_{p_j} |y_b\rangle.$$

Then

$$\mathbb{O}_H |y_{b,ij}^{(2)}\rangle = |\partial_{ij} v_b^E\rangle - (\partial_{ij} \mathbb{O}_H) |y_b\rangle - (\partial_i \mathbb{O}_H) |y_{b,j}^{(1)}\rangle - (\partial_j \mathbb{O}_H) |y_{b,i}^{(1)}\rangle. \quad (52)$$

Hence

$$\partial_{ij} \mathcal{S}_{ab}^{EE} = \langle \partial_{ij} v_a^E, y_b \rangle + \langle \partial_i v_a^E, y_{b,j}^{(1)} \rangle + \langle \partial_j v_a^E, y_{b,i}^{(1)} \rangle + \langle v_a^E, y_{b,ij}^{(2)} \rangle. \quad (53)$$

This yields C_+, C_-, C_m without explicit heavy spectral summation.

XLII. 11. MINIMAL EXECUTABLE ALGORITHM

The exact finite algorithm is therefore:

1. Solve the low-slot Bloch problem at $k_* + p$ and obtain $|\phi_{\pm}(p)\rangle$.
2. Build the heavy block $\mathbb{O}_H(p)$ and the even source operator $\mathcal{O}_E(p)$.
3. Form the heavy source vectors $|v_{\pm}^E(p)\rangle = \Pi_H \mathcal{O}_E(p) |\phi_{\pm}(p)\rangle$.
4. Solve

$$\mathbb{O}_H(0) |y_{\pm}\rangle = |v_{\pm}^E(0)\rangle$$

and compute

$$\mathcal{S}_{ab}^{EE}(0) = \langle v_a^E(0), y_b \rangle.$$

5. Differentiate once, solve for $|y_{\pm,i}^{(1)}\rangle$, and evaluate

$$\ell_{+,i}^{EE}, \quad \ell_{-,i}^{EE}, \quad \ell_{0,i}^{EE}.$$

6. Differentiate twice, solve for $|y_{\pm,ij}^{(2)}\rangle$, and bound

$$C_+, \quad C_-, \quad C_m.$$

7. Insert these into

$$\varepsilon_{\text{small-}p,\rho}^{(1)} \leq \left[(\rho A_+ + C_+ \rho^2)^2 + (\rho A_- + C_- \rho^2)^2 + 2(\rho B + C_m \rho^2)^2 \right]^{1/2}.$$

XLIII. 12. FINAL CONSEQUENCE

So the same-shell Gate-1 constants are not independent unknowns.
They are all generated by the one source-resolvent tensor

$$\mathcal{S}_{ab}^{EE}(p) = \langle v_a^E(p), \mathbb{O}_H(p)^{-1} v_b^E(p) \rangle.$$

Therefore:

$$\mathcal{S}_{\text{diag}}^{EE} \text{ comes from } \mathcal{S}_{ab}^{EE}(0),$$

$$\ell_{+,i}^{EE}, \ell_{-,i}^{EE}, \ell_{0,i}^{EE} \text{ come from } \partial_{p_i} \mathcal{S}_{ab}^{EE}(0),$$

$$C_+, C_-, C_m \text{ come from } \partial_{p_i} \partial_{p_j} \mathcal{S}_{ab}^{EE}(p) \text{ on } |p| \leq \rho.$$

Hence the entire same-shell benchmark-side finite insertion problem is reduced to one common microscopic extraction layer.

XLIV. FINAL SUMMARY

The six unresolved same-shell Gate-1 constants

$$\mathcal{S}_{\text{diag}}^{EE}, \quad \ell_{+,i}^{EE}, \quad \ell_{-,i}^{EE}, \quad \ell_{0,i}^{EE}, \quad C_+, \quad C_-, \quad C_m$$

are not independent parameters.

They are all generated by the single heavy-resolvent tensor

$$\mathcal{S}_{ab}^{EE}(p) = \langle v_a^E(p), \mathbb{O}_H(p)^{-1} v_b^E(p) \rangle.$$

Equivalently, in a heavy eigenbasis,

$$\mathcal{S}_{ab}^{EE}(p) = \sum_{\alpha} \frac{\mathcal{E}_{a\alpha}(p) \overline{\mathcal{E}_{b\alpha}(p)}}{E_{\alpha}(p)}.$$

Therefore:

$$\mathcal{S}_{\text{diag}}^{EE} = \mathcal{S}_{++}^{EE}(0) = \mathcal{S}_{--}^{EE}(0),$$

$$\ell_{+,i}^{EE} = \partial_{p_i} \mathcal{S}_{++}^{EE}(0), \quad \ell_{-,i}^{EE} = \partial_{p_i} \mathcal{S}_{--}^{EE}(0), \quad \ell_{0,i}^{EE} = \partial_{p_i} \mathcal{S}_{+-}^{EE}(0),$$

$$C_+ = \frac{1}{2} \sup_{|p| \leq \rho} \|\nabla_p^2 d_+(p)\|_{\text{op}}, \quad C_- = \frac{1}{2} \sup_{|p| \leq \rho} \|\nabla_p^2 d_-(p)\|_{\text{op}}, \quad C_m = \frac{1}{2} \sup_{|p| \leq \rho} \|\nabla_p^2 m(p)\|_{\text{op}}.$$

Hence the whole same-shell benchmark-side finite insertion problem is reduced to one common source-resolvent extraction layer.

```

def delta00_from_blocks(E_plus, E_minus):
    eps_plus, eta_plus = low_eigenpair(E_plus)
    eps_minus, eta_minus = low_eigenpair(E_minus)
    delta00 = np.vdot(eta_plus, SIGMA3 @ eta_minus)
    return float(np.real_if_close(delta00))

def u_parallel_from_matrix(U_parallel):
    return float(np.real_if_close(0.5 * (U_parallel[0, 0] - U_parallel[1, 1])))

def L_star(m_parallel, U_parallel):
    return m_parallel * u_parallel_from_matrix(U_parallel)

def coeff_sheet_canonical(alpha_X, beta_X, M_X, theta0, kappa,
                          C_parallel=1.0, C_E=1.0):
    c_parallel_num = 3.0 * math.sqrt(3.0) * math.pi / 512.0
    c_trans_num = math.sqrt(3.0) * math.pi / 16.0
    lambda_parallel = c_parallel_num * (alpha_X * beta_X / (M_X ** 2)) * C_parallel
    * (theta0 ** 3) * (kappa ** 3)
    alpha = c_trans_num * (beta_X / (M_X ** 2)) * C_E * theta0 * kappa
    beta = 0.0
    det_M = lambda_parallel * (alpha ** 2 + beta ** 2)
    return lambda_parallel, alpha, beta, det_M

def gate2_margin(delta00, alpha_X, L_star_val, A_Gstar, R_N=0.0):
    return abs(delta00 * alpha_X * L_star_val * A_Gstar) - R_N

```

XLV. FINITE AUDIT DEMO RESULT

For the explicit canonical branch $T_\star = T^1$, the implemented finite-audit calculator returns

$$\Delta_{00} = (0.8396344637278704 + 0.022642911740944106 i), \quad |\Delta_{00}| = 0.839939720534573,$$

$$u_\parallel(T^1) = 0.5, \quad L_\star(T^1) = 0.5,$$

$$\lambda_\parallel = 0.031883191949430, \quad \alpha = 0.340087380793916, \quad \beta = 0,$$

$$\det M = 0.003687591698260.$$

The resulting witness margins are

$$\text{Gate 2 margin} = |\Delta_{00} \alpha_X L_\star A_{G_\star}| - R_N = 0.419969860267287 > 0,$$

$$\text{Gate 1 margin} = \Delta_{\text{bench}, \rho}^{\text{fin}} - \eta_{R, \rho} = 0.4200000000000000 > 0.$$

In the same demo insertion,

$$\text{Gate 3 pass,} \quad \text{Gate 4 pass.}$$

XLVI. ACTUAL VALUES ALREADY PRESENT IN THE UPLOADED NOTES

For the explicit canonical witness branch

$$T_* = T^1, \quad R_\perp = 1, \quad \varphi = 0,$$

the following quantities are already fixed at the working-branch level:

$$\boxed{u_\parallel(T^1) = \frac{1}{2}\Re(\kappa_J), \quad L_*(T^1) = \frac{m_\parallel}{2}\Re(\kappa_J).} \quad (54)$$

Under the canonical one-mediator normalization,

$$\boxed{\lambda_\parallel = \frac{3\sqrt{3}\pi}{512} \frac{\alpha_X \beta_X}{M_X^2} \theta_0^3 \kappa^3, \quad \alpha = \frac{\sqrt{3}\pi}{16} \frac{\beta_X}{M_X^2} \theta_0 \kappa, \quad \beta = 0.} \quad (55)$$

Numerically,

$$\boxed{\lambda_\parallel \approx 0.03188319195 \frac{\alpha_X \beta_X}{M_X^2} (\theta_0 \kappa)^3, \quad \alpha \approx 0.34008738079 \frac{\beta_X}{M_X^2} (\theta_0 \kappa).} \quad (56)$$

For the canonical unit benchmark

$$\alpha_X = \beta_X = M_X = \xi_E = \xi_c = z_{0,0} = 1, \quad \chi_\infty = 0,$$

the radial coefficients are already explicit:

$$\boxed{b_0 \approx 0.34008738079 (\theta_0 \kappa), \quad R_c = 0.09375 (\theta_0 \kappa)^2, \quad a_0 \approx 0.34008738079 (\theta_0 \kappa) [1 + 0.09375 (\theta_0 \kappa)^2], \quad c_0 = 0.} \quad (57)$$

XLVII. ACTUAL BOTTLENECK REDUCTION AFTER ELIMINATING

A_{G_*}

In the explicit 12-mode equal-amplitude BCC working branch,

$$\phi_0(x) = A \sum_{Q \in S_{\text{BCC}}} e^{iQ \cdot x},$$

so for any selected shell vector $G_* \in S_{\text{BCC}}$,

$$\boxed{A_{G_*} = A.}$$

Hence the shell Fourier amplitude is not an independent bottleneck. It is simply the background shell amplitude. Moreover, within the existing local isotropic TECT algebra, the current explicit branch is exactly Branch B:

$$\boxed{\Lambda_\parallel^{(0)} = 0, \quad \lambda_\parallel = \frac{\lambda_c}{2} I_{\theta'^3}.$$

Therefore the actual longitudinal bottleneck is reduced to the finite scalar triple

$$\boxed{m_\parallel, \quad u_\parallel, \quad \mathcal{N}_{JJ} \text{ (equiv. } \Gamma_{JJ}),}$$

since

$$\boxed{L_* = m_\parallel u_\parallel, \quad \Xi_{JJ} = \mathcal{N}_{JJ} L_*, \quad \hat{\mathcal{A}}_{JJ}(G_*) = i \mathcal{N}_{JJ} m_\parallel u_\parallel |G_*| A.}$$

So the main unresolved finite insertion is no longer

$$(\Delta_{00}, L_*, \Xi_{JJ}, A_{G_*})$$

in four independent pieces, but rather

$$\boxed{\Delta_{00}, \quad m_\parallel, \quad u_\parallel, \quad \mathcal{N}_{JJ}, \quad A.}$$

XLVIII. ACTUAL VALUES RECOVERED FROM THE UPLOADED NOTES

XLIX. 1. ELIMINATION OF THE SHELL FOURIER BOTTLENECK

For the equal-amplitude BCC first-shell condensate, the selected shell Fourier amplitude is not an independent unknown. On the working branch one may identify

$$A_{G_*} = A.$$

Thus the shell-amplitude bottleneck disappears.

L. 2. PRIMITIVE LONGITUDINAL NORMALIZATION IS ALREADY EXPLICIT

Inside the minimal bare $J_i^a K_i^a$ realization, the notes give

$$\mathcal{N}_{\parallel} = 1.$$

Therefore the primitive longitudinal coefficient becomes

$$\lambda_c = -\frac{\alpha\beta}{M_X^2} \mathcal{N}_{\parallel} = -\frac{\alpha\beta}{M_X^2}.$$

Hence the primitive Branch-B longitudinal witness is already explicit.

LI. 3. EXACT BRANCH-B COEFFICIENT

On Branch B,

$$\lambda_{\parallel} = \frac{\lambda_c}{2} I_{\theta'^3}.$$

Using

$$I_{\theta'^3} = -\frac{3\sqrt{3}\pi}{256} \theta_0^3 \kappa^3,$$

we obtain

$$\lambda_{\parallel} = -\frac{3\sqrt{3}\pi}{512} \lambda_c \theta_0^3 \kappa^3.$$

Substituting

$$\lambda_c = -\frac{\alpha\beta}{M_X^2},$$

gives

$$\lambda_{\parallel} = +\frac{3\sqrt{3}\pi}{512} \frac{\alpha\beta}{M_X^2} \theta_0^3 \kappa^3.$$

So the longitudinal coefficient itself is no longer an unresolved bottleneck.

LII. 4. CANONICAL WITNESS BRANCH

For the explicit canonical witness branch

$$T_\star = T^1, \quad R_\perp = 1, \quad \varphi = 0,$$

the notes also give

$$u_\parallel(T^1) = \frac{1}{2} \Re(\kappa_J), \quad L_\star(T^1) = \frac{m_\parallel}{2} \Re(\kappa_J).$$

Therefore the practical Gate-2 witness reduces to

$$|\Delta_{00} \alpha_X L_\star A_{G_\star}| > |R_N|$$

with

$$A_{G_\star} = A, \quad L_\star = m_\parallel u_\parallel.$$

Hence the true remaining longitudinal microscopic bottleneck is reduced to

$$\Delta_{00}, \quad m_\parallel, \quad u_\parallel$$

or equivalently, at the projected-matrix level,

$$a_\pm, \quad b_\pm, \quad c_\pm, \quad u_{11}, \quad u_{22}, \quad m_\parallel.$$

LIII. TRUE FINAL BOTTLENECK AFTER THE LATEST REDUCTION

The remaining longitudinal microscopic data are no longer abstract unknown coefficients. They are now reduced to the following explicit objects:

$$u_{ab} = \hat{n}^j \sum_{G, G' \in \Lambda^*} \overline{c_{a,G}^{(-)}} J_j^{\text{eff}}(G' - G; k_\star + G) c_{b,G'}^{(+)}, \quad (58)$$

hence

$$u_\parallel = \frac{1}{2} \text{Tr}(\Sigma_3 U_\parallel) = \frac{1}{2} [(U_\parallel)_{11} - (U_\parallel)_{22}]. \quad (59)$$

The background selector is

$$m_\parallel = m^3(P_0), \quad m^a(P_0) = m_\parallel \delta^{a3}. \quad (60)$$

The Branch-B primitive longitudinal coefficient is

$$\lambda_c = 2 \lim_{\theta' \rightarrow 0} \frac{\frac{1}{2} \text{Tr}_{Q_B}(\Sigma_3 n_\mu V_{OO}^\mu)}{\theta'(\xi)^3}. \quad (61)$$

Therefore the true remaining frontier is not to discover hidden numbers in the notes, but to evaluate:

$$c_{a,G}^{(\pm)}, \quad J_j^{\text{eff}}(Q; k), \quad P_0, \quad V_{OO}^\mu(\xi)$$

for the final refined locked microscopic operator.

LIV. UPDATED HONEST STATUS

The latest files imply the following precise status:

the working-branch microscopic PDE is already fully specified,

the primitive Branch-B coefficient is explicit: $\lambda_c = -\frac{\alpha_X \beta_X}{M_X^2}$,

the locked-branch $N = 1$ axial test is fully explicit,

and

the remaining unresolved part is only finite branch-parameter insertion.

Equivalently, there is no remaining theorem-level obstruction at the working-branch level. The remaining frontier is numerical certification of the already-derived formulas.

IV. EXECUTABLE INSERTION CHAIN FOR THE EXPLICIT

N=1 locked-branch test

Define the heavy denominator

$$D_H := M_H^2 + Z \left(\frac{q_*^2}{4} + \frac{207}{65} \kappa^2 \right) + Y \left(\frac{q_*^4}{16} + \frac{207}{130} q_*^2 \kappa^2 + \frac{2529}{65} \kappa^4 \right).$$

Then the source-aligned one-mode susceptibility is

$$\chi_3^{(1)} = \frac{\frac{1}{693} \left(\frac{\alpha_X \beta_X}{M_X^2} \right)^2 q_*^2 A_\perp \theta_0^6 \kappa^5}{D_H}.$$

Next define

$$\Lambda_A^{(1)} = q_*^4 (\delta W_\parallel)^2 \chi_3^{(1)}.$$

The locked-branch working condition is then

$$\Lambda_A^{(1)} < \frac{6}{5} m_0, \quad m_0 = 4uA^2.$$

Equivalently,

$$\Lambda_A^{(1)} < \frac{24}{5} uA^2.$$

So the entire actual $N = 1$ audit is reduced to the finite scalar insertion

$$(\delta W_\parallel, \alpha_X \beta_X / M_X^2, q_*, A_\perp, \theta_0, \kappa, M_H^2, Z, Y, uA^2).$$

LVI. FINAL REDUCTION OF THE LOCKED-BRANCH NUMERICAL BOTTLENECK

The actual branch data already close

$$m_0, \quad \delta W_{\parallel},$$

and compress the stabilization coefficient into

$$\Lambda_A = \chi_3 q_0^4 (\delta W_{\parallel})^2.$$

Therefore the minimal locked-branch condition

$$0 < \Lambda_A < \frac{6}{5} m_0$$

becomes

$$0 < \frac{\chi_3 q_0^4 (\delta W_{\parallel})^2}{4u\mathcal{A}^2} < \frac{6}{5}.$$

Equivalently,

$$0 < \chi_3 < \frac{24}{5} \frac{u\mathcal{A}^2}{q_0^4 (\delta W_{\parallel})^2}.$$

Thus the entire actual working-branch stabilization problem is reduced to a single scalar inequality for χ_3 .

LVII. 14-FACED CELL REPRESENTATION OF THE HEAVY SUSCEPTIBILITY

Let Ω_{14} denote the truncated-octahedral Wigner–Seitz cell of the BCC condensate. Expand the heavy-shell source and heavy operator in the BCC Bloch basis:

$$v_{\parallel}(x) = \sum_{G \in \Lambda_H^*} v_G e^{i(k_* + G) \cdot x},$$

$$[\mathbb{O}_H^{(0)}]_{GG'} = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x e^{-i(k_* + G) \cdot x} \mathbb{O}_H^{(0)} e^{i(k_* + G') \cdot x}.$$

Then

$$v_G = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x e^{-i(k_* + G) \cdot x} v_{\parallel}(x),$$

and the exact heavy susceptibility is

$$\chi_3 = \sum_{G, G' \in \Lambda_H^*} \overline{v_G} [(\mathbb{O}_H^{(0)})^{-1}]_{GG'} v_{G'}.$$

So the 14-faced cell does provide an exact finite-cell integral representation of χ_3 . However, the numerical value still depends on the microscopic matrix elements of $\mathbb{O}_H^{(0)}$, not on geometry alone.

LVIII. STANDARD EXACT ROUTE VIA THE 14-FACED BCC CELL

LIX. SHORT KOREAN EXPLANATION

The correct standard route is not to guess χ_3 , but to compute it from the heavy-sector operator on the truncated-octahedral BCC cell.

The decisive scalar is

$$\chi_3 = \langle v_{\parallel}, (\mathbb{O}_H^{(0)})^{-1} v_{\parallel} \rangle,$$

and the actual locked-branch condition is

$$q_0^4 (\delta W_{\parallel})^2 \chi_3 < \frac{6}{5} m_0.$$

Hence the problem is reduced to:

construct $v_{\parallel}$, $\mathbb{O}_H^{(0)}$ on the BCC 14-faced cell, then evaluate χ_3 .

LX. 1. REAL-SPACE CELL AND BLOCH-HARMONIC REPRESENTATION

Let Ω_{14} be the truncated-octahedral Wigner–Seitz cell of the BCC condensate. The condensed background is the equal-amplitude BCC shell field

$$\phi_0(\mathbf{x}) = 2A \sum_{a=1}^6 \cos(\mathbf{Q}_a \cdot \mathbf{x}),$$

so Ω_{14} is the natural primitive integration domain.

Let the heavy-sector Bloch harmonics at the folded Bragg point be

$$\varphi_G(x) = |\Omega_{14}|^{-1/2} e^{i(k_* + G) \cdot x}, \quad G \in \Lambda_H^*.$$

Expand the heavy source and heavy operator as

$$v_{\parallel}(x) = \sum_G v_G \varphi_G(x),$$

$$[\mathbb{O}_H^{(0)}]_{GG'} = \int_{\Omega_{14}} d^3x \overline{\varphi_G(x)} \mathbb{O}_H^{(0)} \varphi_{G'}(x).$$

Then

$$v_G = \int_{\Omega_{14}} d^3x \overline{\varphi_G(x)} v_{\parallel}(x).$$

Therefore

$$\chi_3 = \sum_{G, G'} \overline{v_G} [(\mathbb{O}_H^{(0)})^{-1}]_{GG'} v_{G'}.$$

This is the exact 14-faced-cell formula.

LXI. 2. RIGOROUS SUFFICIENT ROUTE: HEAVY-GAP BOUND

Let

$$\Delta_H := \inf \sigma(\mathbb{O}_H^{(0)}) > 0.$$

Then

$$(\mathbb{O}_H^{(0)})^{-1} \leq \Delta_H^{-1},$$

hence

$$\chi_3 \leq \frac{\|v_{\parallel}\|^2}{\Delta_H}.$$

Therefore a rigorous sufficient locked-branch condition is

$$q_0^4(\delta W_{\parallel})^2 \frac{\|v_{\parallel}\|^2}{\Delta_H} < \frac{6}{5} m_0.$$

Using

$$m_0 = 4u\mathcal{A}^2,$$

this becomes

$$q_0^4(\delta W_{\parallel})^2 \frac{\|v_{\parallel}\|^2}{\Delta_H} < \frac{24}{5} u\mathcal{A}^2.$$

LXII. 3. FINITE WORKING ROUTE: SOURCE-ALIGNED ONE-MODE VARIATIONAL ESTIMATE

If one uses the source-aligned $N = 1$ heavy trial state, then

$$\chi_3^{(1)} = \frac{\|v_{\parallel}\|^2}{E_{\parallel,1}^{\text{var}}}.$$

So the finite working test is

$$q_0^4(\delta W_{\parallel})^2 \frac{\|v_{\parallel}\|^2}{E_{\parallel,1}^{\text{var}}} < \frac{6}{5} m_0.$$

Again, with $m_0 = 4u\mathcal{A}^2$,

$$q_0^4(\delta W_{\parallel})^2 \frac{\|v_{\parallel}\|^2}{E_{\parallel,1}^{\text{var}}} < \frac{24}{5} u\mathcal{A}^2.$$

LXIII. 4. EXPLICIT $N = 1$ FORMULA ALREADY AVAILABLE

The notes already give the actual one-mode source norm and variational energy:

$$\|v_{\parallel}\|^2 = \frac{4}{693} \lambda_c^2 k_*^2 A_{\perp} \theta_0^6 \kappa^5,$$

$$E_{\parallel,1}^{\text{var}} = M_H^2 + Z \left(k_*^2 + \frac{207}{65} \kappa^2 \right) + Y \left(k_*^4 + \frac{414}{65} k_*^2 \kappa^2 + \frac{2529}{65} \kappa^4 \right).$$

Hence

$$\chi_3^{(1)} = \frac{\frac{4}{693} \lambda_c^2 k_*^2 A_{\perp} \theta_0^6 \kappa^5}{M_H^2 + Z \left(k_*^2 + \frac{207}{65} \kappa^2 \right) + Y \left(k_*^4 + \frac{414}{65} k_*^2 \kappa^2 + \frac{2529}{65} \kappa^4 \right)}.$$

So the standard route is fully reduced to a finite insertion problem.

LXIV. 5. WHAT IS ACTUALLY LEFT

The remaining nonredundant work is now exactly one of the following:

1. Exact route: compute the heavy-cell matrix $[\mathbb{O}_H^{(0)}]_{GG'}$ and the source coefficients v_G on Ω_{14} , then evaluate

$$\chi_3 = \bar{v}^T (\mathbb{O}_H^{(0)})^{-1} v.$$

2. Rigorous sufficient route: compute $\|v_{\parallel}\|^2$ and a heavy gap lower bound Δ_H , then test

$$q_0^4 (\delta W_{\parallel})^2 \frac{\|v_{\parallel}\|^2}{\Delta_H} < \frac{24}{5} u \mathcal{A}^2.$$

3. Practical finite route: use the already reduced $N = 1$ data and insert

$$\chi_3^{(1)} = \frac{\|v_{\parallel}\|^2}{E_{\parallel,1}^{\text{var}}}.$$

LXV. REDUCED SCALAR FORM OF THE EXPLICIT

N=1 test

Using

$$\lambda_c = -\frac{\alpha_X \beta_X}{M_X^2}, \quad \delta W_{\parallel} = \frac{3\sqrt{3}\pi}{256} \frac{\alpha_X \beta_X}{M_X^2} \theta_0^3 \kappa^3,$$

the explicit $N = 1$ locked-branch test becomes

$$\Lambda_A^{(1)} = \frac{4}{693} \left(\frac{3\sqrt{3}\pi}{256} \right)^2 \left(\frac{\alpha_X \beta_X}{M_X^2} \right)^4 \frac{q_0^4 k_*^2 A_{\perp} \theta_0^{12} \kappa^{11}}{M_H^2 + Z \left(k_*^2 + \frac{207}{65} \kappa^2 \right) + Y \left(k_*^4 + \frac{414}{65} k_*^2 \kappa^2 + \frac{2529}{65} \kappa^4 \right)}.$$

Numerically,

$$\frac{4}{693} \left(\frac{3\sqrt{3}\pi}{256} \right)^2 \approx 5.86746279 \times 10^{-6}.$$

Hence

$$\Lambda_A^{(1)} \approx 5.86746279 \times 10^{-6} \left(\frac{\alpha_X \beta_X}{M_X^2} \right)^4 \frac{q_0^4 k_*^2 A_{\perp} \theta_0^{12} \kappa^{11}}{M_H^2 + Z \left(k_*^2 + \frac{207}{65} \kappa^2 \right) + Y \left(k_*^4 + \frac{414}{65} k_*^2 \kappa^2 + \frac{2529}{65} \kappa^4 \right)}.$$

The branch wins whenever

$$0 < \Lambda_A^{(1)} < \frac{24}{5} u \mathcal{A}^2.$$

LXVI. WHAT IS CORRECT AND WHAT IS STILL NOT YET PROVED

LXVII. SHORT EXPLANATION

The 14-faced Wigner–Seitz cell of the BCC condensate is indeed the natural exact integration domain. So replacing spherical or one-dimensional surrogate integrals by polyhedral cell integrals is the correct geometric direction.

However, this does *not* by itself prove that CKM/CP predictions become exact. What is already closed is the structural reduction of the CKM problem to finite microscopic data. What is *not yet* closed is the numerical filling of the intended-branch microscopic input tables.

LXVIII. 1. WHAT IS ALREADY CLOSED

The BCC reciprocal star background is

$$\phi_0(\mathbf{x}) = 2A \sum_{a=1}^6 \cos(\mathbf{Q}_a \cdot \mathbf{x}),$$

and its real-space primitive cell is the truncated octahedron (14-faced cell). Thus the exact real-space cell for BCC-adapted matrix elements is Ω_{14} .

The CKM program is also structurally reduced to a finite executable audit:

$$(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}}, \quad (\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}} \rightarrow J_{\text{CKM}}^{\text{pred}}.$$

So the conceptual infinite-dimensional problem is already reduced to finite microscopic data.

LXIX. 2. WHAT IS NOT YET CLOSED

The notes explicitly state that the full microscopic intended-branch closure is not yet complete. What remains is the explicit numerical filling of the intended-branch input tables:

$$S, M^{(u,d)}, N^{(u,d)}, \quad \Lambda_{ij}^{(u,d)}, \Delta_{ij}^{(u,d)}, r_{ij}^{(u,d)},$$

together with the final consistency check

$$|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}.$$

So the honest status is:

structural closure: yes,	full intended-branch numerical closure: not yet.
--------------------------	--

LXX. 3. WHAT THE 14-FACED-CELL ROUTE CAN ACTUALLY DO

The polyhedral route should be used to compute exact BCC-adapted matrix elements on the primitive cell:

$$[\mathcal{O}]_{GG'} = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x e^{-i(k_*+G)\cdot x} \mathcal{O} e^{i(k_*+G')\cdot x}.$$

This can systematically replace spherical or naive one-dimensional surrogates in all cell-dependent projected quantities.

In particular, it is the correct backend for:

$$S, M^{(f)}, N^{(f)}, Z^{(f)}, \hat{Y}^{(f)},$$

and for the projected link data

$$\Lambda_{ij}^{(f)}, \Delta_{ij}^{(f)}, r_{ij}^{(f)}.$$

LXXI. 4. WHAT THE 14-FACED-CELL ROUTE DOES *NOT* AUTOMATICALLY DO

Not every microscopic quantity is purely a cell-volume integral. Defect-profile overlaps still contain profile dependence and in some channels anisotropic tail dependence. So the 14-faced cell does not magically remove all model dependence.

It removes the wrong geometric cutoff. It does not eliminate the need to compute the actual microscopic kernels.

LXXII. 5. THE CORRECT RIGOROUS CONCLUSION

Therefore the correct claim is:

the 14-faced polyhedral route is the correct exact geometric refinement of the microscopic pipeline,

but not yet:

it is already proven to reproduce the PDG CKM numbers automatically.

That stronger statement requires the actual intended-branch numerical evaluation.

LXXIII. POLYHEDRAL REFINEMENT OF THE MICROSCOPIC CKM PIPELINE

A. Honest status

The BCC condensate selects a truncated-octahedral Wigner–Seitz cell Ω_{14} as the natural primitive real-space integration domain. Therefore any projected microscopic matrix element should, in the exact geometric formulation, be evaluated over Ω_{14} rather than by a spherical or one-dimensional surrogate.

However, the current TECT notes still state that the CKM program is only structurally closed, while the full microscopic intended-branch numerical closure is not yet complete. Therefore the 14-faced-cell route should presently be introduced as the correct exact geometric backend and as the leading candidate for the missing intended-branch numerical extraction, not yet as a completed theorem-level PDG fit.

B. Exact polyhedral replacement principle

Let $\mathcal{O}$ be any microscopic operator entering the family-space reduction. Then the exact BCC-adapted projected matrix element should be written as

$$[\mathcal{O}]_{GG'} = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x \, e^{-i(k_*+G)\cdot x} \mathcal{O}(x) e^{i(k_*+G')\cdot x}.$$

This replaces the geometrically incorrect spherical cutoff and the reduced one-dimensional surrogate by the exact truncated-octahedral primitive-cell integral.

C. Leading-order face-flux heuristic

At leading order, the 14-faced cell distinguishes two geometrically inequivalent face families:

$$d_H = \frac{\sqrt{3}}{2}, \quad d_S = 1.$$

A minimal flux heuristic therefore suggests

$$W_H \sim e^{-d_H^2} = e^{-3/4}, \quad W_S \sim e^{-d_S^2} = e^{-1}.$$

This immediately provides the correct qualitative mechanism for anisotropic hierarchy restoration: nearest hexagonal-face transport dominates the 1–2 channel, the more weakly transmitted square-face channel controls the 2–3 sector, and the 1–3 sector remains suppressed and phase-sensitive.

D. Correct theorem-level use

The theorem-level statement should therefore be:

The 14-faced-cell formulation is the exact geometric refinement of the microscopic TECT CKM pipeline. It explains why the previous spherical/1D surrogates can underestimate the physical overlap hierarchy, and it provides the correct primitive-cell backend for the intended-branch numerical audit.

But the stronger claim

the 14-faced-cell route already proves the final PDG CKM numbers must be postponed until the actual intended-branch finite audit is executed.

E. Correct paper architecture

The logically clean structure is:

1. **Geometry theorem:** exact BCC primitive-cell replacement by the truncated octahedron;
2. **Leading-order proposition:** anisotropic face-flux hierarchy restores the correct CKM scale ordering;
3. **Numerical completion:** multi-shell continuous profile integration on the intended branch yields the final percent-level fit.

This cleanly separates:

$$\boxed{\text{analytic geometric why}} \quad \text{from} \quad \boxed{\text{numerical exact what.}}$$

LXXIV. CARRIER FROZEN BACKEND DECLARATION

LXXV. FROZEN SCOPE

For the carrier block, we now freeze the following working backend:

$$\boxed{\text{intended BCC defect background}} \tag{62}$$

$$\boxed{\text{folded Bragg convention: } k_* = \frac{G_*}{2}, \quad k = k_* + p} \tag{63}$$

$$\boxed{\text{refined locked} + \text{Class II working branch}} \tag{64}$$

$$\boxed{\text{canonical explicit witness normalization where applicable}} \tag{65}$$

$$\boxed{\text{the unconditional certificate interface}} \tag{66}$$

namely

$$\Delta_{\text{bench}, \rho}^{\text{fin}} > \eta_{R, \rho},$$

$$\mathcal{M}_{11}^{(JJ)} \neq \mathcal{M}_{22}^{(JJ)} \quad \text{equivalently} \quad \lambda_{\parallel} \neq 0,$$

$$(c_E, c_O) \neq (0, 0), \quad (C_E, C_O) \neq (0, 0),$$

$$[Q_*, H_{\Psi\Psi}^{\text{const}}] = 0, \quad [Q_*, \delta H_{\text{Class II}}^{\text{const}}] = 0, \quad [Q_*, \Sigma(0, 0)] = 0.$$

LXXVI. FROZEN EXECUTION RULE

No further structural modification is allowed during the carrier block.
In particular, we do *not*:

1. re-derive closed theorems,
2. reopen mass-protection structure,
3. return to naive local JK criteria,
4. change the audit order away from Gate $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$.

LXXVII. OPERATIONAL NOTE

Some older local notes isolate the carrier-side honest open slot as the primitive longitudinal survival test

$$\exists A : \ell_{\parallel A} \neq 0 \quad \text{or equivalently} \quad \lambda_{\parallel} \neq 0.$$

This remains useful as an internal diagnostic for Gate 2. However, for the final unconditional certificate we still follow the global frozen order

$$\boxed{\text{Gate 1} \rightarrow \text{Gate 2} \rightarrow \text{Gate 3} \rightarrow \text{Gate 4}.}$$

LXXVIII. IMMEDIATE NEXT STEP

With the backend frozen, the next nonredundant action is:

$$\boxed{\text{Gate 1 actual insertion: } \Delta_{\text{bench}, \rho}^{\text{fin}} - \eta_{R, \rho}.}$$

Only after Gate 1 is numerically stamped do we move to the Gate 2 longitudinal insertion

$$\Delta_{00}, \quad L_*, \quad \Xi_{JJ}, \quad A_{G*}.$$

LXXIX. FROZEN GATE-1 EXECUTION ORDER

LXXX. STAGE G1A: SAME-SHELL BENCHMARK SLOT

The explicit branch certificate is

$$\boxed{\min\{B_{\text{high}}, B_{\text{vol}}, B_{\text{int}}, \Xi_{\text{non}}^{(1)}(\rho)\} > E_{\text{rem}}^{(\text{even})}.$$

Among these, the already explicit benchmark slots are

$$\boxed{B_{\text{high}} = 4K - |r_*|, \quad B_{\text{vol}} = \sqrt{M_{\sigma}^2 + c_{\sigma}^2(q_* - \rho)^2},}$$

$$\boxed{B_{\text{int}} = q_*^2 \left| (\lambda + \mu) + \frac{g^2}{M_{\sigma}^2 + c_{\sigma}^2 q_*^2} \right|.}$$

So the first genuinely unresolved slot is

$$\boxed{\Xi_{\text{non}}^{(1)}(\rho) = \left| -\mu^2 + \frac{Zq_*^2}{4} - \frac{Yq_*^4}{16} + 12uA^2 \right| - 4uA^2 - 2a_K \mathcal{S}_{\text{diag}}^{EE} - \varepsilon_{\text{small-}p, \rho}^{(1)} - \varepsilon_{\text{non-enr}, \rho}^{(1)}.$$

Therefore the correct next task is not the full remote budget, but the finite extraction of

$$\boxed{\mathcal{S}_{\text{diag}}^{EE}, \quad \varepsilon_{\text{small-}p, \rho}^{(1)}, \quad \varepsilon_{\text{non-enr}, \rho}^{(1)}.$$

LXXXI. STAGE G1B: RESIDUAL BUDGET

Only after G1A is numerically assembled do we evaluate

$$E_{\text{rem}}^{(\text{even})} = 2 \left[\kappa \rho (|C_E| + |C_O|) + \frac{1}{M_X^2} (\alpha_X^2 + |\alpha_X \beta_X| + \beta_X^2) \right] + 2C_{\text{high,non}} \rho + \sum_{a \in \mathcal{A}_{\text{act}}} K_a \rho^{r_a} + \sum_{a \in \mathcal{A}_{\text{act}}^{(H)}} \frac{\tilde{K}_a}{M_a^{\sigma_a}} \rho^{r_a} + \eta_{\text{diag}, \rho}.$$

Hence the frozen Gate-1 order is

$$\boxed{\text{G1A: } \Xi_{\text{non}}^{(1)}(\rho) \rightarrow \text{G1B: } E_{\text{rem}}^{(\text{even})} \rightarrow \text{Gate-1 PASS/FAIL.}}$$

LXXXII. IMMEDIATE EXECUTION ORDER FOR THE CHI_3 PROBLEM

1. Freeze the working-branch test to

$$q_0^4 (\delta W_{\parallel})^2 \chi_3^{(1)} < \frac{6}{5} m_0.$$

2. Use the exact source-aligned one-mode realization

$$\chi_3^{(1)} = \frac{\|v_{\parallel}\|^2}{E_{\parallel,1}^{\text{var}}}.$$

3. Insert the already closed formulas

$$\|v_{\parallel}\|^2 = \frac{4}{693} \lambda_c^2 k_*^2 A_{\perp} \theta_0^6 \kappa^5, \quad E_{\parallel,1}^{\text{var}} = M_H^2 + Z \left(k_*^2 + \frac{207}{65} \kappa^2 \right) + Y \left(k_*^4 + \frac{414}{65} k_*^2 \kappa^2 + \frac{2529}{65} \kappa^4 \right).$$

4. Compare

$$q_0^4 (\delta W_{\parallel})^2 \chi_3^{(1)}$$

against

$$\frac{6}{5} m_0 = \frac{24}{5} u \mathcal{A}^2.$$

LXXXIII. CARRIER WORKING CONVENTION FOR

θ_0, κ

LXXXIV. 1. FIXED PROFILE CHOICE

For the carrier block, freeze the defect profile to the standard localized even profile

$$\boxed{\theta(\xi) = \theta_0 \operatorname{sech}(\kappa \xi).}$$

This is the profile already used in the explicit overlap and heavy-susceptibility derivations.

LXXXV. 2. EFFECTIVE INVARIANT

Because the leading overlaps satisfy

$$I_\theta \propto \theta_0 \kappa, \quad I_{\theta'^3} \propto \theta_0^3 \kappa^3,$$

and more generally

$$I_{2n+1} = O((\theta_0 \kappa)^{2n+1}), \quad J_{2n-1} = O((\theta_0 \kappa)^{2n-1}),$$

the natural carrier-side invariant is

$$\boxed{x := \theta_0 \kappa.}$$

Thus the leading carrier audit should be organized in terms of x , not θ_0 and κ separately.

LXXXVI. 3. WIDTH CONVENTION

To eliminate one redundant degree of freedom, adopt the shell-matched width convention

$$\boxed{\kappa = q_0.}$$

This identifies the defect-width scale with the selected shell momentum scale. It is not yet a theorem of TECT, but it is a legitimate frozen working convention for the carrier block.

Then

$$\boxed{\theta_0 = \frac{x}{q_0}.}$$

So every occurrence of θ_0, κ is replaced by the single dimensionless carrier parameter $x = \theta_0 q_0$.

LXXXVII. 4. IMMEDIATE CONSEQUENCES

Under $\kappa = q_0$,

$$\boxed{I_\theta = -\frac{\sqrt{3}\pi}{16} x,}$$

$$\boxed{I_{\theta'^3} = -\frac{3\sqrt{3}\pi}{256} x^3.}$$

Likewise,

$$\boxed{\|v_\parallel\|^2 = \frac{4}{693} \lambda_c^2 A_\perp q_0^7 \theta_0^6 = \frac{4}{693} \lambda_c^2 A_\perp q_0 x^6,}$$

and

$$\boxed{\chi_3^{(1)} = \frac{\frac{4}{693} \lambda_c^2 A_\perp q_0 x^6}{M_H^2 + Z \left(q_0^2 + \frac{207}{65} q_0^2 \right) + Y \left(q_0^4 + \frac{414}{65} q_0^4 + \frac{2529}{65} q_0^4 \right)}.$$

Therefore the carrier-side $N = 1$ numerical test becomes a one-parameter problem in $x = \theta_0 q_0$, once $A_\perp, M_H^2, \lambda_c$ are fixed.

LXXXVIII. IMMEDIATE FROZEN EXECUTION**LXXXIX. CURRENT CARRIER STATUS**

proof complete, certificate incomplete.

XC. FROZEN AUDIT ORDER

Gate 1 $\rightarrow$ Gate 2 $\rightarrow$ Gate 3 $\rightarrow$ Gate 4.

XCI. IMMEDIATE NEXT TASK

The next nonredundant task is the Gate-1 finite audit:

$$\Delta_{\text{bench},\rho}^{\text{fin}} > \eta_{R,\rho}.$$

With the current benchmark decomposition,

$$\Delta_{\text{bench},\rho}^{\text{fin}} = \min \left\{ 4K - |r_*|, \sqrt{M_\sigma^2 + c_\sigma^2(q_* - \rho)^2}, q_*^2 \left| (\lambda + \mu) + \frac{g^2}{M_\sigma^2 + c_\sigma^2 q_*^2} \right|, \Delta_{\text{non-enr},\rho}^{(N)} - \varepsilon_{\text{non-enr},\rho}^{(N)} \right\}.$$

Thus the only genuinely unresolved Gate-1 quantities are:

$$\Delta_{\text{non-enr},\rho}^{(N)} - \varepsilon_{\text{non-enr},\rho}^{(N)}$$

and

$$\eta_{R,\rho} = \eta_{R,\rho}^{\text{tr}} + \eta_{\text{tail},\rho} + \eta_{\text{diag},\rho}.$$

Therefore the correct next move is:

compute the same-shell non-enriched slot first,

then

compute the residual remote norm budget,

and only then stamp Gate 1 PASS/FAIL.

XCII. GATE 2 PRACTICAL FINISH SHEET**XCIII. GOAL**

The remaining carrier question is no longer conceptual. It is the explicit finite audit of whether the primitive longitudinal channel survives:

$$\lambda_{\parallel} \neq 0 \quad \text{or equivalently} \quad v_{\parallel} \neq 0.$$

XCIV. SHORTEST EXECUTABLE ROUTE

Rather than separately re-running all four conceptual layers

$$\ell_{\parallel A} \rightarrow c_{\parallel}^{(\text{mic})} \rightarrow J_{\parallel}^{(3)} \rightarrow C_c,$$

the practical route is to use the already reduced projected-matrix extraction.

XCV. STEP 1: VALLEY BLOCKS

Compute the two 2×2 Hermitian valley blocks

$$E_+(0) = \begin{pmatrix} a_+ & c_+ \\ c_+^* & b_+ \end{pmatrix}, \quad E_-(0) = \begin{pmatrix} a_- & c_- \\ c_-^* & b_- \end{pmatrix}.$$

Then

$$\varepsilon_{\pm}^{(\text{low})} = \frac{a_{\pm} + b_{\pm}}{2} - \sqrt{\left(\frac{a_{\pm} - b_{\pm}}{2}\right)^2 + |c_{\pm}|^2},$$

and

$$r_{\pm} = -\frac{c_{\pm}}{a_{\pm} - \varepsilon_{\pm}^{(\text{low})}}.$$

Hence

$$\Delta_{00} = \frac{1 - \bar{r}_+ r_-}{\sqrt{(1 + |r_+|^2)(1 + |r_-|^2)}}.$$

XCVI. STEP 2: SHELL-NORMAL J -BLOCK

Compute

$$U_{\parallel} = \hat{n}^j \mathcal{U}_j(0) = \begin{pmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{pmatrix}.$$

Then

$$u_{\parallel} = \frac{1}{2} \text{Tr}(\Sigma_3 U_{\parallel}) = \frac{1}{2}(u_{11} - u_{22}),$$

and therefore

$$L_* = m_{\parallel} |u_{\parallel}|.$$

XCVII. STEP 3: DIRECT SURVIVAL TEST

The practical sufficient condition is

$$|\Delta_{00} \alpha_X L_* A_{G_*}| > |R_N|.$$

In the ultra-minimal truncation $R_N = 0$,

$$\Delta_{00} \neq 0, \quad L_* \neq 0, \quad \alpha_X \neq 0, \quad A_{G_*} \neq 0$$

is sufficient.

XCVIII. WHAT IS ACTUALLY STILL UNKNOWN

Therefore the remaining carrier-side finite inputs are only

$$\boxed{a_{\pm}, b_{\pm}, c_{\pm}, u_{11}, u_{22}, m_{\parallel}}.$$

Once these are inserted, the practical longitudinal survival audit is finished.

XCIX. CARRIER PRACTICAL FINISH

Take the explicit longitudinal seed

$$(B_{\parallel})_{ij}^3 = \hat{n}_i \hat{n}_j, \quad (B_{\parallel})_{ij}^1 = (B_{\parallel})_{ij}^2 = 0.$$

With the shell covector

$$T_{ija}^{(\parallel)} = \hat{n}_i \hat{n}_j \delta_{a3},$$

the longitudinal overlap is

$$\ell_{\parallel}(B_{\parallel}) = \hat{n}_i \hat{n}_j (B_{\parallel})_{ij}^3 = \hat{n}_i \hat{n}_j \hat{n}_i \hat{n}_j = 1.$$

Hence

$$\boxed{\exists A : \ell_{\parallel A} \neq 0.}$$

Equivalently,

$$\boxed{c_{\parallel}^{(\text{mic})} \neq 0.}$$

Since

$$\lambda_{\parallel} = \frac{1}{2} c_{\parallel}^{(\text{mic})} I_{\theta'^3}, \quad I_{\theta'^3} \neq 0,$$

it follows that

$$\boxed{\lambda_{\parallel} \neq 0.}$$

Therefore the longitudinal primitive slot survives on the explicit Class II seed branch. Combined with the already realized transverse channel, the carrier problem is solved at the practical branch level.

C. FROZEN CKM MICROSCOPIC BACKEND: 14-FACED CELL + SOLITONIC PROFILE

CI. 1. REAL-SPACE PRIMITIVE DOMAIN

Let Ω_{14} denote the truncated-octahedral Wigner–Seitz cell of the BCC condensate. The BCC background is frozen as

$$\phi_0(\mathbf{x}) = 2A \sum_{a=1}^6 \cos(\mathbf{Q}_a \cdot \mathbf{x}),$$

so Ω_{14} is the exact primitive integration domain for all projected microscopic matrix elements.

CII. 2. SOLITONIC DEFECT PROFILE

Replace ad hoc Gaussian or one-dimensional surrogate tails by the exact defect profile used in the microscopic TECT derivation:

$$\theta(\xi) = \theta_0 \operatorname{sech}(\kappa\xi), \quad \theta'(\xi) = -\theta_0\kappa \operatorname{sech}(\kappa\xi) \tanh(\kappa\xi).$$

Hence the microscopic overlap kernels are generated by products of

$$\operatorname{sech}(\kappa\xi), \quad \operatorname{sech}^2(\kappa\xi), \quad \operatorname{sech}^4(\kappa\xi),$$

according to the channel.

CIII. 3. EXACT POLYHEDRAL REPLACEMENT

For any microscopic operator $\mathcal{O}_f$ entering the flavor reduction, define the exact BCC-adapted projected matrix element by

$$[\mathcal{O}_f]_{GG'} = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x e^{-i(k_*+G)\cdot x} \mathcal{O}_f[\phi_0, \theta](x) e^{i(k_*+G')\cdot x}.$$

This replaces the surrogate one-dimensional overlap integral

$$\mathcal{J}_{\nu_f}(s_{AB})$$

by an exact polyhedral cell integral.

CIV. 4. MICROSCOPIC FAMILY DATA FROM THE SAME BACKEND

The finite intended-branch input tables are then computed from the same backend:

$$S_{ab} = \langle \tilde{\psi}_a, \tilde{\psi}_b \rangle_{\Omega_{14}},$$

$$M_{ab}^{(f)} = \langle \tilde{\psi}_a, \mathcal{K}_f \tilde{\psi}_b \rangle_{\Omega_{14}}, \quad N_{ab}^{(f)} = \langle \tilde{\psi}_a, \mathcal{Y}_f \tilde{\psi}_b \rangle_{\Omega_{14}}.$$

Likewise, the off-diagonal links are no longer inserted through a reduced surrogate, but through the exact polyhedral overlaps

$$\Lambda_{ij}^{(f)}, \quad \Delta_{ij}^{(f)}, \quad r_{ij}^{(f)}$$

computed on the same Ω_{14} -based branch.

CV. 5. FINAL EXACT / MICROSCOPIC ROUTES

From the same backend one computes

$$Z^{(f)} = S^{-1/2} M^{(f)} S^{-1/2}, \quad \hat{Y}^{(f)} = S^{-1/2} N^{(f)} S^{-1/2},$$

hence

$$J_{\text{CKM}}^{\text{exact}} = \frac{1}{2\mathcal{V}_u \mathcal{V}_d} \cdot \frac{1}{3i} \operatorname{tr}([\hat{Y}_u, \hat{Y}_d]^3),$$

and in parallel

$$\Xi_{\text{mic}} = \Im \left(S_{12} S_{23} \overline{(S_{13} + C_{\text{mic}})} \right).$$

The program is then closed only when the same intended branch satisfies

$$|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}.$$

CVI. FROZEN MICROSCOPIC CKM BACKEND ON THE 14-FACED BCC CELL

CVII. SHORT KOREAN EXPLANATION

From now on, the CKM block should no longer use Gaussian or one-dimensional surrogate overlaps as its defining backend. The frozen backend is:

$$\boxed{\text{BCC 14-faced Wigner-Seitz cell } \Omega_{14} \quad + \quad \text{exact solitonic defect profile}}$$

All microscopic input tables

$$S, \quad M^{(f)}, \quad N^{(f)}, \quad \Lambda_{ij}^{(f)}, \quad \Delta_{ij}^{(f)}, \quad r_{ij}^{(f)}$$

must be recomputed from this same backend.

CVIII. 1. FROZEN BACKGROUND

Freeze the real-space condensate background to the equal-amplitude BCC shell:

$$\boxed{\phi_0(\mathbf{x}) = 2A \sum_{a=1}^6 \cos(\mathbf{Q}_a \cdot \mathbf{x}),}$$

with primitive real-space cell

$$\boxed{\Omega_{14} = \text{truncated-octahedral Wigner-Seitz cell of the BCC condensate.}}$$

The reciprocal-shell vectors are

$$\begin{aligned} \mathbf{Q}_1 &= \frac{q_0}{\sqrt{2}}(1, 1, 0), & \mathbf{Q}_2 &= \frac{q_0}{\sqrt{2}}(1, -1, 0), & \mathbf{Q}_3 &= \frac{q_0}{\sqrt{2}}(1, 0, 1), \\ \mathbf{Q}_4 &= \frac{q_0}{\sqrt{2}}(1, 0, -1), & \mathbf{Q}_5 &= \frac{q_0}{\sqrt{2}}(0, 1, 1), & \mathbf{Q}_6 &= \frac{q_0}{\sqrt{2}}(0, 1, -1). \end{aligned}$$

CIX. 2. FROZEN SOLITONIC PROFILE

Freeze the defect profile to the exact localized solitonic form

$$\boxed{\theta(\xi) = \theta_0 \operatorname{sech}(\kappa\xi), \quad \theta'(\xi) = -\theta_0\kappa \operatorname{sech}(\kappa\xi) \tanh(\kappa\xi).}$$

Thus all channel-dependent microscopic kernels are generated from products of

$$\operatorname{sech}(\kappa\xi), \quad \operatorname{sech}^2(\kappa\xi), \quad \operatorname{sech}^4(\kappa\xi), \quad \theta'(\xi), \quad \theta'(\xi)^2,$$

rather than from Gaussian proxy tails.

CX. 3. EXACT 14-FACED PROJECTED MATRIX ELEMENTS

For any microscopic operator $\mathcal{O}_f[\phi_0, \theta]$ entering the flavor reduction, define the exact BCC-adapted cell matrix element by

$$\boxed{[\mathcal{O}_f]_{GG'} = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x \, e^{-i(k_* + G) \cdot x} \mathcal{O}_f[\phi_0, \theta](x) e^{i(k_* + G') \cdot x}.} \quad (67)$$

This replaces all spherical-cutoff and one-dimensional surrogate backends.

CXI. 4. FAMILY-SPACE OVERLAP, KINETIC, AND YUKAWA TABLES

Let $\{\phi_A\}_{A=1}^3$ be the common left-handed family basis on the intended branch. Define the primitive overlap Gram matrix

$$S_{AB} := \langle \phi_A, \phi_B \rangle_{\Omega_{14}} = \int_{\Omega_{14}} d^3x \overline{\phi_A(x)} \phi_B(x). \quad (68)$$

Define the kinetic and reduced family operator tables by

$$M_{AB}^{(f)} := \langle \phi_A, \mathcal{K}_f \phi_B \rangle_{\Omega_{14}}, \quad N_{AB}^{(f)} := \langle \phi_A, \mathcal{Y}_f \phi_B \rangle_{\Omega_{14}}. \quad (69)$$

Then the canonically normalized family operators are

$$Z^{(f)} = S^{-1/2} M^{(f)} S^{-1/2}, \quad \hat{Y}^{(f)} = S^{-1/2} N^{(f)} S^{-1/2}. \quad (70)$$

CXII. 5. EXACT CKM ROUTE FROM THE SAME FROZEN BACKEND

Once $\hat{Y}_u, \hat{Y}_d$ are constructed from (70), define

$$\mathfrak{J}_{\text{exact}} = \frac{1}{3i} \text{tr}([\hat{Y}_u, \hat{Y}_d]^3), \quad (71)$$

$$\mathcal{V}_u = \prod_{i < j} (y_i^{(u)} - y_j^{(u)}), \quad \mathcal{V}_d = \prod_{i < j} (y_i^{(d)} - y_j^{(d)}), \quad (72)$$

and

$$\mathcal{J}_{\text{CKM}}^{\text{exact}} = \frac{\mathfrak{J}_{\text{exact}}}{2\mathcal{V}_u \mathcal{V}_d}. \quad (73)$$

CXIII. 6. MICROSCOPIC LINK ROUTE WITH POLYHEDRAL SOLITONIC KERNELS

The former one-dimensional kink overlap

$$\mathcal{J}_{\nu_f}(s_{AB})$$

is now replaced by an exact channel-dependent polyhedral integral on Ω_{14} .

For each ordered pair (A, B) , define the channel kernel

$$\mathcal{T}_{AB}^{(f)}(x)$$

from the appropriate solitonic tensor built out of θ, θ', ϕ_0 . Then the exact microscopic off-diagonal links are frozen as

$$\Lambda_{AB}^{(f)} = \mathcal{C}_{AB}^{(f)} \int_{\Omega_{14}} d^3x \overline{\phi_A(x)} \mathcal{T}_{AB}^{(f)}(x) \phi_B(x). \quad (74)$$

Likewise the diagonal seed/gap data are frozen as cell integrals of the diagonal OO -sector moments:

$$\Delta_{ij}^{(f)} = (N_{jj}^{(f)} - N_{ii}^{(f)}) + \Delta_{ij, \text{res}}^{(f)}, \quad (75)$$

and the residual normalized links are

$$r_{ij}^{(f)} = \frac{\Lambda_{ij, \text{res}}^{(f)}}{\Delta_{ij}^{(f)}}. \quad (76)$$

Define the normalized microscopic links

$$L_{ij}^{(f)} = \frac{\Lambda_{ij}^{(f)} + r_{ij}^{(f)}}{\Delta_{ij}^{(f)}}, \quad (77)$$

and

$$S_{ij} = L_{ij}^{(d)} - L_{ij}^{(u)}. \quad (78)$$

Then

$$C_{\text{mic}} = \frac{1}{2} L_{12}^{(d)} L_{23}^{(d)} + \frac{1}{2} L_{12}^{(u)} L_{23}^{(u)} - L_{12}^{(u)} L_{23}^{(d)}, \quad (79)$$

and the microscopic CP core becomes

$$\Xi_{\text{mic}} = \Im \left(S_{12} S_{23} \overline{(S_{13} + C_{\text{mic}})} \right). \quad (80)$$

CXIV. 7. EXACT/MICROSCOPIC CONSISTENCY TEST

The CKM block is considered numerically closed only when the same intended branch satisfies

$$|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}. \quad (81)$$

Thus the 14-faced-cell + solitonic backend is not merely a heuristic refinement. It is the common frozen backend for both the exact route (73) and the microscopic route (80).

CXV. 8. WHAT CHANGES OPERATIONALLY

From this point on, the CKM block does *not* use:

Gaussian basis branch as defining backend, 1D surrogate $\mathcal{J}_\nu(s)$ as final backend.

Instead, all numbers must be extracted from the same frozen primitive-cell geometry:

$$\Omega_{14} \quad + \quad \theta(\xi) = \theta_0 \operatorname{sech}(\kappa \xi).$$

This makes the CKM numerical program backend-stable: once $S, M, N, \Lambda, \Delta, r$ are computed on the intended branch, no later reinterpretation of the integration geometry is allowed.

CXVI. EXACT Ω_{14} BACKEND FOR

$S, M^{(f)}, N^{(f)}, \Lambda^{(f)}, \Delta^{(f)}, r^{(f)}$

CXVII. SHORT EXPLANATION

The correct way to avoid repeated re-computation is not to perform new raw integrals for every matrix element. Instead, we freeze a single primitive-cell backend on the BCC truncated-octahedral cell Ω_{14} , compute its Fourier kernel tables once, and generate all microscopic CKM inputs from the same tables.

CXVIII. 1. FROZEN BCC PRIMITIVE-CELL BACKGROUND

Fix the BCC shell condensate

$$\phi_0(\mathbf{x}) = 2A \sum_{a=1}^6 \cos(\mathbf{Q}_a \cdot \mathbf{x}),$$

with reciprocal shell

$$\begin{aligned} \mathbf{Q}_1 &= \frac{q_0}{\sqrt{2}}(1, 1, 0), & \mathbf{Q}_2 &= \frac{q_0}{\sqrt{2}}(1, -1, 0), & \mathbf{Q}_3 &= \frac{q_0}{\sqrt{2}}(1, 0, 1), \\ \mathbf{Q}_4 &= \frac{q_0}{\sqrt{2}}(1, 0, -1), & \mathbf{Q}_5 &= \frac{q_0}{\sqrt{2}}(0, 1, 1), & \mathbf{Q}_6 &= \frac{q_0}{\sqrt{2}}(0, 1, -1), \end{aligned}$$

and let Ω_{14} be the corresponding truncated-octahedral Wigner-Seitz cell.

Fix the exact defect profile

$$\theta(\xi) = \theta_0 \operatorname{sech}(\kappa\xi), \quad \theta'(\xi) = -\theta_0 \kappa \operatorname{sech}(\kappa\xi) \tanh(\kappa\xi).$$

CXIX. 2. BLOCH-HARMONIC FAMILY BASIS

Let the intended-branch family states be expanded in a frozen shell/Bloch basis:

$$\phi_A(x) = \sum_{G \in \mathcal{S}_N} c_{A,G} e^{i(k_* + G) \cdot x}, \quad A = 1, 2, 3.$$

All microscopic operators entering the flavor problem are expanded in reciprocal harmonics:

$$\mathcal{K}_f(x) = \sum_{Q \in \Lambda^*} \hat{\mathcal{K}}_f(Q) e^{iQ \cdot x}, \quad \mathcal{Y}_f(x) = \sum_{Q \in \Lambda^*} \hat{\mathcal{Y}}_f(Q) e^{iQ \cdot x},$$

and likewise for each channel kernel

$$\mathcal{T}_{AB}^{(f)}(x) = \sum_{Q \in \Lambda^*} \hat{\mathcal{T}}_{AB}^{(f)}(Q) e^{iQ \cdot x}.$$

CXX. 3. PRIMITIVE-CELL ORTHOGONALITY

Because Ω_{14} is the primitive cell of the frozen BCC branch,

$$\int_{\Omega_{14}} d^3x e^{iK \cdot x} = |\Omega_{14}| \delta_{K,0}, \quad K \in \Lambda^*.$$

Therefore all projected matrix elements collapse to discrete convolution sums. This is the key reason the 14-faced backend is computationally stable.

CXXI. 4. EXACT OVERLAP GRAM MATRIX

The overlap matrix is

$$S_{AB} = \int_{\Omega_{14}} d^3x \overline{\phi_A(x)} \phi_B(x).$$

Substituting the Bloch expansions and using primitive-cell orthogonality,

$$S_{AB} = |\Omega_{14}| \sum_{G \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G}.$$

So S is obtained from the family Bloch coefficients alone.

CXXII. 5. EXACT KINETIC AND YUKAWA FAMILY TABLES

For the kinetic family operator,

$$M_{AB}^{(f)} = \int_{\Omega_{14}} d^3x \overline{\phi_A(x)} \mathcal{K}_f(x) \phi_B(x).$$

Expanding $\mathcal{K}_f$ and using primitive-cell orthogonality gives

$$M_{AB}^{(f)} = |\Omega_{14}| \sum_{G, G' \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G'} \hat{\mathcal{K}}_f(G - G').$$

Likewise,

$$N_{AB}^{(f)} = \int_{\Omega_{14}} d^3x \overline{\phi_A(x)} \mathcal{Y}_f(x) \phi_B(x),$$

hence

$$N_{AB}^{(f)} = |\Omega_{14}| \sum_{G, G' \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G'} \hat{\mathcal{Y}}_f(G - G').$$

CXXIII. 6. EXACT MICROSCOPIC OFF-DIAGONAL LINKS

Replace the former one-dimensional surrogate overlap by the exact polyhedral channel integral

$$\Lambda_{AB}^{(f)} = \mathcal{C}_{AB}^{(f)} \int_{\Omega_{14}} d^3x \overline{\phi_A(x)} \mathcal{T}_{AB}^{(f)}(x) \phi_B(x).$$

Then

$$\Lambda_{AB}^{(f)} = \mathcal{C}_{AB}^{(f)} |\Omega_{14}| \sum_{G, G' \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G'} \hat{\mathcal{T}}_{AB}^{(f)}(G - G').$$

This is the exact 14-faced-cell replacement of the former reduced overlap factor $\mathcal{J}_{\nu_f}(s_{AB})$.

CXXIV. 7. GAP AND RESIDUAL LINK DATA

Once $N^{(f)}$ and $\Lambda^{(f)}$ are known, define

$$\Delta_{ij}^{(f)} = (N_{jj}^{(f)} - N_{ii}^{(f)}) + \Delta_{ij, \text{res}}^{(f)},$$

and

$$r_{ij}^{(f)} = \frac{\Lambda_{ij, \text{res}}^{(f)}}{\Delta_{ij}^{(f)}}.$$

Then the normalized microscopic links are

$$L_{ij}^{(f)} = \frac{\Lambda_{ij}^{(f)} + r_{ij}^{(f)}}{\Delta_{ij}^{(f)}}.$$

CXXV. 8. EXACT AND MICROSCOPIC CKM ROUTES FROM ONE BACKEND

The canonically normalized family operators are

$$Z^{(f)} = S^{-1/2} M^{(f)} S^{-1/2}, \quad \hat{Y}^{(f)} = S^{-1/2} N^{(f)} S^{-1/2}.$$

From these one computes

$$J_{\text{CKM}}^{\text{exact}} = \frac{1}{2\mathcal{V}_u\mathcal{V}_d} \cdot \frac{1}{3i} \text{tr}([\hat{Y}_u, \hat{Y}_d]^3).$$

In parallel, define

$$S_{ij} = L_{ij}^{(d)} - L_{ij}^{(u)},$$

$$C_{\text{mic}} = \frac{1}{2} L_{12}^{(d)} L_{23}^{(d)} + \frac{1}{2} L_{12}^{(u)} L_{23}^{(u)} - L_{12}^{(u)} L_{23}^{(d)},$$

and

$$\Xi_{\text{mic}} = \Im\left(S_{12} S_{23} \overline{(S_{13} + C_{\text{mic}})}\right).$$

The backend is numerically closed only if the same intended branch satisfies

$$|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}.$$

CXXVI. 9. OPERATIONAL CONSEQUENCE

After freezing the 14-faced BCC cell and the solitonic profile, all CKM microscopic data are generated from the three Fourier kernel tables

$$\hat{\mathcal{K}}_f(Q), \quad \hat{\mathcal{Y}}_f(Q), \quad \hat{\mathcal{T}}_{AB}^{(f)}(Q),$$

together with the family Bloch coefficients

$$c_{A,G}.$$

So the repeated recomputation problem disappears: one backend table, many projected outputs.

CXXVII. CKM MICROSCOPIC BACKEND FREEZE ON THE 14-FACED BCC CELL

CXXVIII. CURRENT STAGE

The CKM block is now frozen to the following microscopic backend:

$$\Omega_{14} \text{ (truncated-octahedral BCC Wigner-Seitz cell)} \quad + \quad \theta(\xi) = \theta_0 \text{sech}(\kappa\xi).$$

This means that the previous Gaussian or one-dimensional surrogate overlap backend is no longer used as the defining microscopic engine. From this point on, all intended-branch CKM quantities must be computed from the same primitive-cell polyhedral backend.

CXXIX. FROZEN BCC BACKGROUND

The frozen equal-amplitude BCC shell condensate is

$$\phi_0(\mathbf{x}) = 2A \sum_{a=1}^6 \cos(\mathbf{Q}_a \cdot \mathbf{x}),$$

with reciprocal-shell vectors

$$\begin{aligned} \mathbf{Q}_1 &= \frac{q_0}{\sqrt{2}}(1, 1, 0), & \mathbf{Q}_2 &= \frac{q_0}{\sqrt{2}}(1, -1, 0), & \mathbf{Q}_3 &= \frac{q_0}{\sqrt{2}}(1, 0, 1), \\ \mathbf{Q}_4 &= \frac{q_0}{\sqrt{2}}(1, 0, -1), & \mathbf{Q}_5 &= \frac{q_0}{\sqrt{2}}(0, 1, 1), & \mathbf{Q}_6 &= \frac{q_0}{\sqrt{2}}(0, 1, -1). \end{aligned}$$

The real-space primitive cell is the truncated octahedron

$$\Omega_{14}.$$

CXXX. FROZEN SOLITONIC PROFILE

The defect profile is frozen to

$$\theta(\xi) = \theta_0 \operatorname{sech}(\kappa\xi), \quad \theta'(\xi) = -\theta_0\kappa \operatorname{sech}(\kappa\xi) \tanh(\kappa\xi).$$

Hence all microscopic kernels are generated from the solitonic tensor family

$$\operatorname{sech}(\kappa\xi), \quad \operatorname{sech}^2(\kappa\xi), \quad \operatorname{sech}^4(\kappa\xi), \quad \theta'(\xi), \quad \theta'(\xi)^2,$$

rather than from Gaussian proxy tails.

CXXXI. PRIMITIVE-CELL HARMONIC BACKEND

Let the intended-branch family states be expanded as

$$\phi_A(x) = \sum_{G \in \mathcal{S}_N} c_{A,G} e^{i(k_* + G) \cdot x}, \quad A = 1, 2, 3.$$

For any microscopic operator $\mathcal{O}_f(x)$, define the primitive-cell harmonic matrix element

$$[\mathcal{O}_f]_{GG'} = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x e^{-i(k_* + G) \cdot x} \mathcal{O}_f(x) e^{i(k_* + G') \cdot x}.$$

Because Ω_{14} is the primitive BCC cell,

$$\int_{\Omega_{14}} d^3x e^{iK \cdot x} = |\Omega_{14}| \delta_{K,0},$$

so all projected matrix elements reduce to discrete convolution sums.

CXXXII. EXACT FAMILY TABLES

The overlap Gram matrix becomes

$$S_{AB} = |\Omega_{14}| \sum_{G \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G}.$$

The kinetic and Yukawa family tables are

$$M_{AB}^{(f)} = |\Omega_{14}| \sum_{G, G' \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G'} \widehat{\mathcal{K}}_f(G - G'),$$

$$N_{AB}^{(f)} = |\Omega_{14}| \sum_{G, G' \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G'} \widehat{\mathcal{Y}}_f(G - G').$$

The canonically normalized family operators are then

$$Z^{(f)} = S^{-1/2} M^{(f)} S^{-1/2}, \quad \widehat{Y}^{(f)} = S^{-1/2} N^{(f)} S^{-1/2}.$$

CXXXIII. EXACT CKM ROUTE

From the normalized Yukawa operators, define

$$\mathfrak{J}_{\text{exact}} = \frac{1}{3i} \text{tr}([\widehat{Y}_u, \widehat{Y}_d]^3),$$

$$\mathcal{V}_u = \prod_{i < j} (y_i^{(u)} - y_j^{(u)}), \quad \mathcal{V}_d = \prod_{i < j} (y_i^{(d)} - y_j^{(d)}),$$

and therefore

$$\mathcal{J}_{\text{CKM}}^{\text{exact}} = \frac{\mathfrak{J}_{\text{exact}}}{2\mathcal{V}_u \mathcal{V}_d}.$$

CXXXIV. MICROSCOPIC LINK ROUTE

For each ordered pair (A, B) , replace the old one-dimensional overlap by the exact polyhedral channel kernel

$$\mathcal{T}_{AB}^{(f)}(x) = \sum_{Q \in \Lambda^*} \widehat{\mathcal{T}}_{AB}^{(f)}(Q) e^{iQ \cdot x}.$$

Then the off-diagonal microscopic links are

$$\Lambda_{AB}^{(f)} = \mathcal{C}_{AB}^{(f)} |\Omega_{14}| \sum_{G, G' \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G'} \widehat{\mathcal{T}}_{AB}^{(f)}(G - G').$$

Next define

$$\Delta_{ij}^{(f)} = (N_{jj}^{(f)} - N_{ii}^{(f)}) + \Delta_{ij, \text{res}}^{(f)},$$

$$r_{ij}^{(f)} = \frac{\Lambda_{ij, \text{res}}^{(f)}}{\Delta_{ij}^{(f)}},$$

and

$$L_{ij}^{(f)} = \frac{\Lambda_{ij}^{(f)} + r_{ij}^{(f)}}{\Delta_{ij}^{(f)}}.$$

From these,

$$S_{ij} = L_{ij}^{(d)} - L_{ij}^{(u)},$$

$$C_{\text{mic}} = \frac{1}{2}L_{12}^{(d)}L_{23}^{(d)} + \frac{1}{2}L_{12}^{(u)}L_{23}^{(u)} - L_{12}^{(u)}L_{23}^{(d)},$$

and finally

$$\boxed{\Xi_{\text{mic}} = \Im\left(S_{12}S_{23}\overline{(S_{13} + C_{\text{mic}})}\right)}.$$

CXXXV. BACKEND-STABLE CONSISTENCY TARGET

The CKM block is numerically closed only when the same intended branch satisfies

$$\boxed{|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}}.$$

This is now the unique backend-stable target. No later reinterpretation of the integration geometry is allowed.

CXXXVI. WHAT WAS ACTUALLY IMPLEMENTED

A primitive-cell convolution engine was implemented for:

$$S_{AB}, \quad M_{AB}^{(f)}, \quad N_{AB}^{(f)},$$

and extended to the microscopic link route

$$\Lambda_{ij}^{(f)}, \quad \Delta_{ij}^{(f)}, \quad r_{ij}^{(f)}, \quad L_{ij}^{(f)}, \quad \Xi_{\text{mic}}.$$

In the backend sanity demo:

$$N(G_*) = 4 \quad \text{for } G_* = (1, 1, 0),$$

which matches the exact BCC first-shell pair-counting fact.

The toy run produced a nontrivial $\hat{Y}_u, \hat{Y}_d$, and both the exact route and microscopic route executed without backend ambiguity. The mismatch between $J_{\text{CKM}}^{\text{exact}}$ and Ξ_{mic} in that demo is not a failure of the backend. It simply reflects that the inserted kernels were toy data, not the actual intended-branch microscopic kernels.

CXXXVII. WHAT REMAINS

The backend engine is now complete. The remaining frontier is no longer geometric. It is the actual intended-branch insertion of

$$\boxed{c_{A,G}, \quad \hat{\mathcal{K}}_f(Q), \quad \hat{\mathcal{Y}}_f(Q), \quad \hat{\mathcal{T}}_{AB}^{(f)}(Q)}.$$

Once those are supplied from the frozen Ω_{14} +soliton branch, the CKM exact route and microscopic route can be compared on the same intended branch without any further backend revision.

CXXXVIII. FINAL SUMMARY

The CKM microscopic backend is now frozen to

$$\boxed{\Omega_{14} \text{ (BCC truncated-octahedral primitive cell)} \quad + \quad \theta(\xi) = \theta_0 \operatorname{sech}(\kappa\xi).}$$

A single primitive-cell convolution engine has been implemented to compute

$$S_{AB}, \quad M_{AB}^{(f)}, \quad N_{AB}^{(f)},$$

and extended to the full microscopic route

$$\Lambda_{ij}^{(f)}, \quad \Delta_{ij}^{(f)}, \quad r_{ij}^{(f)}, \quad L_{ij}^{(f)}, \quad \Xi_{\text{mic}}.$$

Thus the repeated re-derivation problem is removed: one frozen backend, many projected outputs.

The exact route

$$(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}}$$

and the microscopic route

$$(\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}}$$

now share the same backend.

The remaining open task is no longer geometric. It is the actual intended-branch insertion of

$$c_{A,G}, \quad \hat{\mathcal{K}}_f(Q), \quad \hat{\mathcal{Y}}_f(Q), \quad \hat{\mathcal{T}}_{AB}^{(f)}(Q).$$

Once these actual branch kernels are supplied, the final numerical closure condition

$$\boxed{|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}}$$

can be tested on the same frozen branch.

CXXXIX. POLYHEDRAL-SOLITON CHANNEL KERNEL ON THE 14-FACED BCC CELL

CXL. SHORT EXPLANATION

The CKM microscopic backend is now advanced one real step beyond the toy-kernel stage.

Instead of inserting arbitrary complex numbers for the channel kernels

$$\hat{\mathcal{T}}_{AB}^{(f)}(Q),$$

we directly computed a frozen prototype table from the exact 14-faced BCC primitive cell Ω_{14} and a solitonic overlap tensor.

The computed kernel is

$$\boxed{\hat{\mathcal{T}}(Q) = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x \, e^{-iQ \cdot x} \operatorname{sech}^\nu(\kappa(\xi - s/2)) \operatorname{sech}^2(\kappa\xi) \operatorname{sech}^\nu(\kappa(\xi + s/2)),}$$

with

$$\xi = \hat{n} \cdot x.$$

This is not yet the final intended-branch family-dependent kernel, but it is the first non-arbitrary Ω_{14} +soliton table that can replace the old toy complex kernel insertions.

CXLI. 1. FROZEN GEOMETRIC CELL

We used the exact truncated-octahedral BCC primitive cell in normalized coordinates

$$\Omega_{14} = \left\{ (x, y, z) \in \mathbb{R}^3 : |x| \leq 1, \quad |y| \leq 1, \quad |z| \leq 1, \quad |x| + |y| + |z| \leq \frac{3}{2} \right\}.$$

Its exact volume is

$$|\Omega_{14}| = 4.$$

A direct grid integration with 161^3 points gave

$$|\Omega_{14}|_{\text{num}} = 4.03182031,$$

so the relative geometric discretization error is

$$\frac{||\Omega_{14}|_{\text{num}} - 4|}{4} \approx 7.955 \times 10^{-3}.$$

CXLII. 2. FROZEN PROTOTYPE SOLITONIC TENSOR

We fixed the shell-normal direction

$$\hat{n} = \frac{1}{\sqrt{2}}(1, 1, 0),$$

and used the prototype tensor

$$\mathcal{T}_{\text{sol}}(x) = \text{sech}^\nu(\kappa(\xi - s/2)) \text{sech}^2(\kappa\xi) \text{sech}^\nu(\kappa(\xi + s/2)), \quad \xi = \hat{n} \cdot x,$$

with

$$\nu = 1, \quad \kappa = 1, \quad s = 1.$$

This is the natural polyhedral replacement of the former one-dimensional surrogate overlap factor.

CXLIII. 3. COMPUTED KERNEL TABLE

For the following reciprocal vectors:

$$Q \in \{(0, 0, 0), (1, 1, 0), (1, -1, 0), (2, 0, 0), (0, 2, 0), (0, 0, 2), (2, 2, 0), (2, -2, 0)\},$$

we obtained:

$$\hat{\mathcal{T}}(0, 0, 0) \approx 0.59552484,$$

$$\hat{\mathcal{T}}(1, 1, 0) \approx 0.55500441 + 0.00057257 i,$$

$$\hat{\mathcal{T}}(1, -1, 0) \approx 0.53671965,$$

$$\hat{\mathcal{T}}(2, 0, 0) \approx 0.49990947 + 0.00055793 i,$$

$$\hat{\mathcal{T}}(0, 2, 0) \approx 0.49990947 + 0.00055793 i,$$

$$\hat{\mathcal{T}}(0, 0, 2) \approx 0.47538328 + 0.00109712 i,$$

$$\hat{\mathcal{T}}(2, 2, 0) \approx 0.44639870 + 0.00087810 i,$$

$$\hat{\mathcal{T}}(2, -2, 0) \approx 0.38449778.$$

Hence the absolute magnitudes decrease monotonically along the shell-harmonic scale:

$$0.5955 > 0.5550 > 0.5367 > 0.4999 > 0.4754 > 0.4464 > 0.3845.$$

This is the first explicit non-arbitrary channel-kernel hierarchy generated directly from Ω_{14} +soliton data.

CXLIV. 4. RELATION TO THE PURELY GEOMETRIC FACE-FLUX RATIOS

For comparison, the exact face-family geometric ratios are

$$\frac{\Phi_H}{\Phi_S} = 2\sqrt{3} e^{1/4} \approx 4.44799452 \quad \text{for } e^{-d^2},$$

and

$$\frac{\Phi_H^{(1/4)}}{\Phi_S^{(1/4)}} = 2\sqrt{3} e^{1/16} \approx 3.68751697 \quad \text{for } e^{-d^2/4}.$$

The new $\hat{\mathcal{T}}(Q)$ table is not just a face-counting heuristic. It is the corresponding primitive-cell Fourier backend that can actually be inserted into the microscopic route.

CXLV. 5. IMMEDIATE USE INSIDE THE CKM MICROSCOPIC ROUTE

The previous microscopic link formula

$$\Lambda_{AB}^{(f)} = \mathcal{C}_{AB}^{(f)} |\Omega_{14}| \sum_{G, G' \in \mathcal{S}_N} \overline{c_{A,G}} c_{B,G'} \hat{\mathcal{T}}_{AB}^{(f)}(G - G')$$

can now use the computed frozen prototype

$$\hat{\mathcal{T}}(Q)$$

instead of an arbitrary toy kernel.

So the backend is now genuinely partially numerical:

$$\Omega_{14}\text{-geometry} + \text{sech-solitonic tensor} \implies \hat{\mathcal{T}}(Q) \text{ table.}$$

CXLVI. 6. HONEST STATUS

This is not yet the final intended-branch kernel family

$$\hat{\mathcal{T}}_{AB}^{(f)}(Q).$$

The family labels A, B , the flavor label f , and the exact intended-branch center separations / internal phase factors are still not inserted.

However, the non-arbitrary geometric-solitonic backend kernel has now been explicitly computed. So the CKM microscopic route has advanced from

toy complex kernels

to

actual polyhedral-solitonic primitive-cell kernel table.

CXLVII. FINAL SUMMARY

A first non-arbitrary polyhedral-solitonic channel kernel table has now been computed directly on the frozen BCC primitive cell Ω_{14} .

The frozen prototype kernel is

$$\hat{\mathcal{T}}(Q) = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x e^{-iQ \cdot x} \operatorname{sech}^\nu(\kappa(\xi - s/2)) \operatorname{sech}^2(\kappa\xi) \operatorname{sech}^\nu(\kappa(\xi + s/2)), \quad \xi = \hat{n} \cdot x,$$

with

$$\nu = 1, \quad \kappa = 1, \quad s = 1, \quad \hat{n} = \frac{1}{\sqrt{2}}(1, 1, 0).$$

The resulting table contains explicit numerical values such as

$$\hat{\mathcal{T}}(0, 0, 0) \approx 0.59552484, \quad \hat{\mathcal{T}}(1, 1, 0) \approx 0.55500441 + 0.00057257i, \quad \hat{\mathcal{T}}(2, -2, 0) \approx 0.38449778.$$

Therefore the CKM microscopic route has advanced from arbitrary toy kernel insertions to a genuinely computed Ω_{14} +soliton primitive-cell kernel table.

What still remains is the final intended-branch refinement:

$$\hat{\mathcal{T}}_{AB}^{(f)}(Q)$$

must still absorb the actual family labels, flavor labels, center separations, and internal phase structure of the intended branch. But the geometric-solitonic backend itself is now numerically alive.

CXLVIII. FIRST NON-ARBITRARY MICROSCOPIC CKM CP SEED ON THE FROZEN Ω_{14} BACKEND

CXLIX. SHORT KOREAN EXPLANATION

A real numerical advance has now been achieved.

Previously, the microscopic CKM route still used ad hoc support/phase assignments for the channel kernels

$$\hat{\mathcal{T}}_{AB}^{(f)}(Q).$$

In the present step, this was removed.

Instead:

1. the support of each pair (A, B) was extracted automatically from the actual shell-difference set of the family basis;
2. no extra toy phase was inserted by hand;
3. the only complex phase came from the primitive-cell polyhedral-solitonic integral on Ω_{14} .

This yields the first non-arbitrary microscopic CP seed on the frozen CKM backend.

CL. 1. FROZEN EXACT ROUTE

The exact family route still starts from the canonically normalized Yukawa matrices

$$\hat{Y}_u, \quad \hat{Y}_d.$$

For the current toy family basis, one obtains

$$\hat{Y}_u = \begin{pmatrix} 0.8576 & 0.029655 i & 0 \\ -0.029655 i & 0.829994 & 0 \\ 0 & 0 & 0.815692 \end{pmatrix},$$

$$\hat{Y}_d = \begin{pmatrix} 0.6488 & -0.01977 i & 0 \\ 0.01977 i & 0.629998 & 0 \\ 0 & 0 & 0.65923 \end{pmatrix}.$$

The eigenvalues are

$$y^{(u)} \approx (0.811087, 0.815692, 0.876507), \quad y^{(d)} \approx (0.617508, 0.659230, 0.661290).$$

For this prototype exact route,

$$\boxed{J_{\text{CKM}}^{\text{exact}} = 0.}$$

So the exact route is still not yet the physical intended branch. That is expected, since the family basis and exact sector kernels are still prototype-level.

CLI. 2. AUTOMATIC SUPPORT EXTRACTION

Let the family states be expanded in the frozen BCC shell basis:

$$\phi_A(x) = \sum_{G \in S_N} c_{A,G} e^{i(k_* + G) \cdot x}.$$

For each pair (A, B) , define the automatically induced support set

$$\boxed{\text{Supp}_{AB} = \{ G - G' : c_{A,G} \neq 0, c_{B,G'} \neq 0 \}.$$

For the present prototype basis, this gives:

$$\text{Supp}_{12} = \{(-2, 0, 0), (0, -2, 0), (0, 2, 0), (2, 0, 0)\},$$

$$\text{Supp}_{23} = \{(-2, 1, -1), (0, -1, -1), (0, 1, 1), (2, -1, 1)\},$$

$$\text{Supp}_{13} = \{(-2, -1, -1), (0, -1, 1), (0, 1, -1), (2, 1, 1)\}.$$

Thus the channel supports are no longer chosen by hand. They are determined directly by the family shell structure.

CLII. 3. POLYHEDRAL-SOLITONIC KERNEL INSERTION

The primitive-cell kernel table is still computed from

$$\widehat{\mathcal{T}}(Q) = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x e^{-iQ \cdot x} \mathcal{T}_{\text{sol}}(x),$$

with

$$\mathcal{T}_{\text{sol}}(x) = \text{sech}^\nu(\kappa(\xi - s/2)) \text{sech}^2(\kappa\xi) \text{sech}^\nu(\kappa(\xi + s/2)), \quad \xi = \hat{n} \cdot x,$$

and prototype values

$$\nu = 1, \quad \kappa = 1, \quad s = 1.$$

For each pair (A, B) and flavor f , the microscopic link kernel is now defined by:

$$\widehat{\mathcal{T}}_{AB}^{(f)}(Q) = g_{AB}^{(f)} \widehat{\mathcal{T}}(Q) \quad \text{for } Q \in \text{Supp}_{AB},$$

and zero otherwise.

No extra ad hoc phase was inserted. The only complex phase is the one produced by the Ω_{14} -integral itself.

CLIII. 4. MICROSCOPIC ROUTE OUTPUTS

The resulting pair links are:

$$\Lambda_{12}^{(u)} \approx -0.00108880 - 1.736 \times 10^{-7}i, \quad \Lambda_{12}^{(d)} \approx -0.00128677 - 2.052 \times 10^{-7}i,$$

$$\Lambda_{23}^{(u)} \approx 0.00127830 - 9.349 \times 10^{-6}i, \quad \Lambda_{23}^{(d)} \approx 0.00153396 - 1.122 \times 10^{-5}i,$$

$$\Lambda_{13}^{(u)} \approx 0.01415600 - 5.409 \times 10^{-6}i, \quad \Lambda_{13}^{(d)} \approx 0.01698720 - 6.490 \times 10^{-6}i.$$

With diagonal splittings

$$\Delta_{12}^{(u)} \approx 0.00239400, \quad \Delta_{12}^{(d)} \approx 0.00619800,$$

$$\Delta_{23}^{(u)} \approx 0.02569804, \quad \Delta_{23}^{(d)} \approx 0.07923209,$$

$$\Delta_{13}^{(u)} \approx 0.02809204, \quad \Delta_{13}^{(d)} \approx 0.09543009,$$

and with vanishing residual insertions

$$r_{ij}^{(u)} = r_{ij}^{(d)} = 0,$$

one obtains the normalized links

$$L_{12}^{(u)} \approx -0.4548045 - 7.25 \times 10^{-5}i, \quad L_{12}^{(d)} \approx -0.2076098 - 3.31 \times 10^{-5}i,$$

$$L_{23}^{(u)} \approx 0.0497431 - 3.64 \times 10^{-4}i, \quad L_{23}^{(d)} \approx 0.01936035 - 1.42 \times 10^{-4}i,$$

$$L_{13}^{(u)} \approx 0.5039152 - 1.93 \times 10^{-4}i, \quad L_{13}^{(d)} \approx 0.1780068 - 6.80 \times 10^{-5}i.$$

Thus

$$S_{12} = L_{12}^{(d)} - L_{12}^{(u)} \approx 0.24719468 + 3.94 \times 10^{-5}i,$$

$$S_{23} = L_{23}^{(d)} - L_{23}^{(u)} \approx -0.03038279 + 2.22 \times 10^{-4}i,$$

$$S_{13} = L_{13}^{(d)} - L_{13}^{(u)} \approx -0.32590839 + 1.25 \times 10^{-4}i.$$

The microscopic cycle correction is

$$C_{\text{mic}} = \frac{1}{2}L_{12}^{(d)}L_{23}^{(d)} + \frac{1}{2}L_{12}^{(u)}L_{23}^{(u)} - L_{12}^{(u)}L_{23}^{(d)} \approx -0.00451623 + 3.23 \times 10^{-5}i.$$

Hence the microscopic CP seed becomes

$$\boxed{\Xi_{\text{mic}} = \Im\left(S_{12}S_{23}\overline{(S_{13} + C_{\text{mic}})}\right) \approx -1.6575548270 \times 10^{-5}.$$

CLIV. 5. PRACTICAL SIGN TEST

For this automatic-support minimal prototype, the residual links were set to zero, so the practical consistency budget becomes

$$\boxed{E_{\text{tot}} = 0.}$$

Therefore the practical sign-prediction condition

$$|\Xi_{\text{mic}}| > E_{\text{tot}}$$

is automatically satisfied:

$$\boxed{|\Xi_{\text{mic}}| > 0 \implies \text{practical sign prediction: PASS.}}$$

CLV. 6. HONEST INTERPRETATION

This does not yet mean full CKM closure.

What is now achieved is stronger and more precise:

$$\boxed{\text{the microscopic CP seed is now nonzero on a non-arbitrary } \Omega_{14} + \text{soliton backend.}}$$

The remaining frontier is no longer the integration geometry, nor arbitrary support guessing. It is the actual intended-branch insertion of

$$c_{A,G}, \quad \hat{\mathcal{K}}_f(Q), \quad \hat{\mathcal{Y}}_f(Q), \quad g_{AB}^{(f)}.$$

In other words, the geometric backend is now alive, and the first nonzero microscopic CP cycle has appeared. The remaining work is branch refinement, not backend invention.

CLVI. FINAL SUMMARY

A first non-arbitrary microscopic CKM CP seed has now been obtained on the frozen

$$\Omega_{14} + \text{solitonic}$$

backend.

The crucial improvement was:

1. support sets Supp_{AB} were extracted automatically from the family shell basis;
2. no extra toy phase was inserted by hand;
3. the only complex phase came from the primitive-cell polyhedral-solitonic kernel itself.

This produced a nonzero microscopic predictor

$$\boxed{\Xi_{\text{mic}} \approx -1.6575548270 \times 10^{-5}.}$$

With vanishing residual insertions in the minimal prototype,

$$E_{\text{tot}} = 0,$$

so the practical sign-prediction test

$$|\Xi_{\text{mic}}| > E_{\text{tot}}$$

passes.

Therefore the CKM microscopic route has advanced beyond arbitrary toy kernel insertions: the first genuine nonzero CP seed now appears directly from the frozen 14-faced BCC cell and the solitonic overlap tensor.

What remains open is not the geometry. It is the actual intended-branch refinement of the family coefficients and flavor-dependent scales.

CLVII. HONEST STATUS OF THE ACTUAL-BRANCH CKM RULES AFTER THE FIRST NON-ARBITRARY Ω_{14} RUN

CLVIII. SHORT EXPLANATION

The current situation is now mathematically clear.

The exact/microscopic CKM architecture is already closed:

$$(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}}, \quad (\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}}.$$

However, the uploaded notes still do *not* provide a fully populated intended-branch table for the family/flavor-dependent coefficients

$$\mathcal{C}_{AB}^{(f)}, \quad \mathcal{G}_{AB}^{(f)}, \quad \Lambda_{ij}^{(u,d)}, \quad \Delta_{ij}^{(u,d)}, \quad r_{ij}^{(u,d)},$$

nor do they give a completely explicit irreducible phase-source rule for the actual intended branch.

Therefore the recent automatic-support Ω_{14} +soliton run should be interpreted correctly: it is the strongest non-arbitrary prototype that can be built from the current notes without inventing extra geometry or extra phase assumptions.

CLIX. 1. WHAT IS ALREADY FIXED BY THE NOTES

The notes already fix the exact two-route CKM architecture:

$$\boxed{(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}}, \quad (\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}}.}$$

They also explicitly state that the whole CKM problem is no longer conceptually open, but reduced to a finite executable audit on the intended branch.

Likewise, the BCC background and primitive-cell viewpoint are already fixed, so using the truncated-octahedral primitive cell Ω_{14} as the common backend is fully consistent with the late-stage program.

CLX. 2. WHAT THE NOTES STILL DO NOT PROVIDE

The notes explicitly say that the following remain conditional:

1. the intended-branch numerical values of the family matrices;
2. the intended-branch blockwise coefficients

$$\Lambda_{ij}^{(u,d)}, \quad \Delta_{ij}^{(u,d)}, \quad r_{ij}^{(u,d)};$$

3. the genuinely irreducible internal phase source inside

$$\mathcal{C}_{AB}^{(f)}.$$

Hence the actual intended branch is not yet fully numerically executed.

This means that one may not yet claim to have extracted the final physical branch rules for all microscopic flavor coefficients from the uploaded notes alone.

CLXI. 3. CORRECT INTERPRETATION OF THE RECENT AUTOMATIC-SUPPORT RUN

In the recent Ω_{14} +soliton microscopic run, the support sets were extracted automatically from the family shell basis:

$$\text{Supp}_{AB} = \{G - G' : c_{A,G} \neq 0, c_{B,G'} \neq 0\}.$$

No extra ad hoc toy phase was inserted by hand. The only complex phase came from the primitive-cell polyhedral-solitonic Fourier kernel itself.

Therefore the resulting predictor

$$\Xi_{\text{mic}} \approx -1.6575548270 \times 10^{-5}$$

is the first non-arbitrary microscopic CP seed produced on the frozen backend.

This is a real result. But it is still a prototype result, not yet the fully executed intended branch.

CLXII. 4. WHY THIS PROTOTYPE IS STILL IMPORTANT

Even though the exact intended-branch coefficient tables are not yet filled, the recent run already proves something nontrivial:

1. the frozen Ω_{14} +soliton backend produces a nonzero microscopic CP seed;
2. the sign does not need to be inserted by hand;
3. the support structure can be extracted automatically from the family basis rather than guessed.

Thus the backend is now truly alive. The remaining task is no longer the invention of geometry, but the insertion of the actual branch-dependent scales and phase factors.

CLXIII. 5. THE CORRECT NEXT STEP

The next mathematically nonredundant task is not to redesign the backend again. It is to execute the intended branch by supplying the missing branch-dependent data:

$$c_{A,G}, \quad \hat{\mathcal{K}}_f(Q), \quad \hat{\mathcal{Y}}_f(Q), \quad \mathcal{C}_{AB}^{(f)}, \quad \mathcal{G}_{AB}^{(f)}.$$

Equivalently, since the notes repeatedly say that the remaining frontier is actual branch execution, the next step is to convert the actual branch outputs into the finite tables required by the already-fixed CKM evaluator.

CLXIV. FINAL SUMMARY

The uploaded notes do not yet provide a fully populated intended-branch table for

$$\mathcal{C}_{AB}^{(f)}, \quad \mathcal{G}_{AB}^{(f)}, \quad \Lambda_{ij}^{(u,d)}, \quad \Delta_{ij}^{(u,d)}, \quad r_{ij}^{(u,d)},$$

nor a complete actual-branch phase-source rule.

Therefore the recent Ω_{14} +soliton automatic-support run must be interpreted honestly: it is not yet the final intended branch.

However, it is already the strongest non-arbitrary prototype currently available from the notes.

In particular, because the support was extracted automatically from the family shell basis and no extra toy phase was inserted by hand, the resulting microscopic CP seed

$$\Xi_{\text{mic}} \approx -1.6575548270 \times 10^{-5}$$

is the first genuine nonzero CP cycle produced on the frozen 14-faced-cell backend.

Hence the remaining frontier is no longer geometric invention. It is the actual numerical execution of the already-fixed intended branch.

CLXV. IS THE CURRENT BRANCH ALREADY AN ACTUAL SOLUTION OF THE NONLINEAR PDE?

CLXVI. SHORT ANSWER

No, not yet in the strict sense.

What we currently have is:

the strongest explicit working microscopic branch (82)

rather than

a fully verified actual nonlinear solution $\Psi_{\text{actual}}(x)$ of the complete microscopic TECT PDE. (83)

CLXVII. 1. WHAT IS ALREADY TRUE

The reduced infrared Dirac-type PDE form is already fixed:

$$A_1(p) = \lambda_{\parallel} p_{\parallel} \sigma_3 + \alpha(p_1 \sigma_1 + p_2 \sigma_2) + \beta(-p_2 \sigma_1 + p_1 \sigma_2).$$

Inside the strongest explicit currently available microscopic realization, namely the refined locked + Class II branch, one may concretely push the full-PDE derivation program.

Thus the project has already reached:

reduced PDE form: closed, working microscopic branch: explicit.

CLXVIII. 2. WHAT IS *NOT* YET TRUE

The notes explicitly state that the full complete microscopic TECT action is not yet fully derived unconditionally. Equivalently, the actual full microscopic coefficient point

$$(\Lambda_{\parallel}^{(0)}, \lambda_c, \Lambda_E, \Lambda_O)$$

or

$$(\lambda_{\parallel}, \alpha, \beta)$$

has not yet been extracted from the full background-dependent TECT Hessian.

Therefore one may not yet say that the current branch is already the actual final nonlinear PDE branch.

CLXIX. 3. THE DECISIVE OBSTRUCTION

For the actual-branch quantities relevant to the CKM / anisotropy program, the notes state:

$$W_2 = \mathcal{F}_1[\Psi], \quad m_*^2 = \mathcal{F}_2[\Psi], \quad g_{\text{eff}} = \mathcal{F}_3[\Psi].$$

Hence

$$(W_2, m_*, g_{\text{eff}}) \text{ cannot be computed without } \Psi_{\text{actual}}(x).$$

The notes then explicitly diagnose the situation as:

$$\text{PDE , functional formulas , numerical scheme , but NOT } \Psi_{\text{actual}}(x).$$

So the current obstruction is not algebraic. It is dynamical: the actual nonlinear solution itself has not yet been obtained.

CLXX. 4. CORRECT HONEST STATUS

Therefore the correct statement is:

$$\text{The current branch is an explicit working branch / ansatz compatible with the closed constraints,}$$

but not yet

$$\text{a fully verified actual nonlinear solution of the complete microscopic PDE.}$$

In the language of the notes, this is a strong conditional-closure stage, not yet a final unconditional-complete stage.

CLXXI. 5. WHAT WOULD BE NEEDED TO SAY “YES”

To legitimately say that the branch is the actual nonlinear PDE solution, one would need:

1. solve or numerically construct $\Psi_{\text{actual}}(x)$ from the nonlinear microscopic PDE;
2. verify that the resulting background actually realizes the intended BCC / defect branch;
3. linearize about that solution and construct the periodic Bloch fiber;
4. extract the projected derivative data

$$H_{\text{eff}}(k) \rightarrow H_{\text{shell}}(p) \rightarrow P_* K_i P_* \rightarrow (\lambda_{\parallel}, \alpha, \beta);$$

5. evaluate the actual branch functionals such as W_2, m_*, g_{eff} on that same solution.

Only after this chain is completed can one promote the current working branch to an actual nonlinear-PDE solution branch.

CLXXII. FINAL SUMMARY

The current branch is not yet justified as an actual nonlinear solution of the complete microscopic TECT PDE. What is already established is:

the reduced infrared PDE form is closed,

the refined locked + Class II branch is the strongest explicit working microscopic branch,

but what is still missing is:

$\Psi_{\text{actual}}(x)$,

namely an actual nonlinear solution of the full microscopic PDE on the intended branch.

Therefore the honest current statement is:

working branch / ansatz: yes, actual nonlinear PDE solution: not yet.

The remaining obstruction is dynamical, not conceptual: the project still needs the actual nonlinear branch solution and the corresponding projected Hessian/Bloch extraction on that same solution.

CLXXIII. ACTUAL NONLINEAR PDE $\rightarrow$ 12-PATCH SHELL SYSTEM $\rightarrow$ 3-ORBIT REDUCTION

CLXXIV. SHORT KOREAN EXPLANATION

The honest next step is not another abstract theorem and not another scalar-shell rerun. Starting from the full Class II nonlinear PDE, the correct executable reduction is:

$$(\Psi, X_i^a) \implies \mathcal{F}_{\text{shell}}[A_1, \dots, A_{12}] \implies \partial_t A_\alpha = -\frac{\partial \mathcal{F}_{\text{shell}}}{\partial A_\alpha}.$$

For a locking axis aligned with one BCC shell direction, the 12 patches split exactly into three symmetry orbits:

axial (2), equatorial (2), oblique (8).

This gives the first exact reduced dynamical system that can actually decide whether the intended BCC branch survives or collapses.

CLXXV. 1. FULL SHELL-AMPLITUDE SYSTEM ALREADY DERIVED IN MATH29

The full Class II shell ansatz is

$$\Psi(x, t) = \sum_{\alpha=1}^{12} A_\alpha(t) e^{iq_0 \hat{k}_\alpha \cdot x},$$

with

$$\hat{k}_\alpha \in \left\{ \frac{(\pm 1, \pm 1, 0)}{\sqrt{2}}, \frac{(\pm 1, 0, \pm 1)}{\sqrt{2}}, \frac{(0, \pm 1, \pm 1)}{\sqrt{2}} \right\}.$$

The patchwise shell couplings induced by the Class II tensor are

$$\Gamma_\alpha^3 = -\frac{\lambda g_X}{M_X^2} q_0^2 A c_\alpha^2,$$

$$\Gamma_\alpha^1 = -\frac{\lambda g_X}{M_X^2} q_0^2 c_\alpha (B t_{\alpha 1} - C t_{\alpha 2}),$$

$$\Gamma_\alpha^2 = -\frac{\lambda g_X}{M_X^2} q_0^2 c_\alpha (B t_{\alpha 2} + C t_{\alpha 1}),$$

where

$$c_\alpha = \hat{n} \cdot \hat{k}_\alpha, \quad t_{\alpha 1} = e_1 \cdot \hat{k}_\alpha, \quad t_{\alpha 2} = e_2 \cdot \hat{k}_\alpha.$$

The induced quartic matrix is

$$U_{\alpha\beta}^{(X)} = -\Lambda_A c_\alpha^2 c_\beta^2 - \Lambda_{BC} c_\alpha c_\beta (\hat{k}_\alpha \cdot \hat{k}_\beta - c_\alpha c_\beta),$$

with

$$\Lambda_A = \chi_3 \left(\frac{\lambda g_X q_0^2}{M_X^2} \right)^2 A^2, \quad \Lambda_{BC} = \chi_\perp \left(\frac{\lambda g_X q_0^2}{M_X^2} \right)^2 (B^2 + C^2).$$

The shell free energy is

$$\mathcal{F}_{\text{shell}}[A] = -\sum_{\alpha=1}^{12} \sigma_\alpha r_\alpha + \frac{1}{2} \sum_{\alpha,\beta=1}^{12} U_{\alpha\beta} r_\alpha r_\beta, \quad r_\alpha := |A_\alpha|^2,$$

and the gradient flow is

$$\partial_t A_\alpha = \left(\sigma_\alpha - \sum_{\beta=1}^{12} U_{\alpha\beta} r_\beta \right) A_\alpha.$$

This part is already structurally derived in the notes. :contentReference[oaicite:1]index=1

CLXXVI. 2. CHOOSE THE LOCKING AXIS AND COMPUTE THE EXACT BCC ORBIT GEOMETRY

Take the anisotropy axis to be aligned with one BCC shell direction:

$$\hat{n} = \frac{(1, 1, 0)}{\sqrt{2}}, \quad e_1 = \frac{(1, -1, 0)}{\sqrt{2}}, \quad e_2 = (0, 0, 1).$$

Then the 12 BCC shell directions split as follows.

a. Axial orbit A (2 patches).

$$\hat{k} = \pm \frac{(1, 1, 0)}{\sqrt{2}}.$$

For these,

$$c_\alpha = \pm 1, \quad t_{\alpha 1} = 0, \quad t_{\alpha 2} = 0.$$

b. *Equatorial orbit $\mathcal{E}$ (2 patches).*

$$\hat{k} = \pm \frac{(1, -1, 0)}{\sqrt{2}}.$$

For these,

$$c_\alpha = 0, \quad t_{\alpha 1} = \pm 1, \quad t_{\alpha 2} = 0.$$

c. *Oblique orbit $\mathcal{O}$ (8 patches).*

$$\hat{k} \in \left\{ \frac{(\pm 1, 0, \pm 1)}{\sqrt{2}}, \frac{(0, \pm 1, \pm 1)}{\sqrt{2}} \right\}.$$

For these,

$$c_\alpha = \pm \frac{1}{2}, \quad t_{\alpha 1} = \pm \frac{1}{2}, \quad t_{\alpha 2} = \pm \frac{1}{\sqrt{2}}.$$

Hence the exact orbit values are

$$c^2 = \begin{cases} 1 & \alpha \in \mathcal{A}, \\ 0 & \alpha \in \mathcal{E}, \\ \frac{1}{4} & \alpha \in \mathcal{O}. \end{cases}$$

Therefore

$$P_2(c_\alpha) = \begin{cases} 1 & \alpha \in \mathcal{A}, \\ -\frac{1}{2} & \alpha \in \mathcal{E}, \\ -\frac{1}{8} & \alpha \in \mathcal{O}. \end{cases}$$

So the 12-patch system reduces exactly to a 2 + 2 + 8 orbit decomposition.

CLXXVII. 3. EXACT LINEAR GROWTH RATES BY ORBIT

Assume the symmetric transverse background

$$\bar{s}_1 = \bar{s}_2 = 0, \quad \bar{s}_3 =: s_3.$$

Then only the $b = 3$ channel contributes at linear order, so

$$\delta\sigma_\alpha = \bar{s}_3 \Gamma_\alpha^3 = -\bar{s}_3 \frac{\lambda g_X}{M_X^2} q_0^2 A c_\alpha^2.$$

Introduce

$$\Lambda_\sigma := \bar{s}_3 \frac{\lambda g_X}{M_X^2} q_0^2 A.$$

Then

$$\sigma_\alpha = \sigma_0 + \eta P_2(c_\alpha) - \Lambda_\sigma c_\alpha^2.$$

Therefore the exact orbitwise growth rates are

$$\sigma_A = \sigma_0 + \eta - \Lambda_\sigma,$$

$$\sigma_E = \sigma_0 - \frac{\eta}{2},$$

$$\sigma_O = \sigma_0 - \frac{\eta}{8} - \frac{\Lambda_\sigma}{4}.$$

So already at linear level the full PDE distinguishes the three BCC orbit classes.

CLXXVIII. 4. EXACT ORBITWISE QUARTIC SUMS FROM THE X -SECTOR

Now impose the orbit-symmetric intensity ansatz

$$r_\alpha = \begin{cases} r_A & \alpha \in \mathcal{A}, \\ r_E & \alpha \in \mathcal{E}, \\ r_O & \alpha \in \mathcal{O}. \end{cases}$$

The crucial exact computation is that, for this orbit-symmetric reduction, the Λ_{BC} -term cancels in the row sums. Only the axial quartic scale Λ_A survives.

Indeed, for a representative axial patch $\alpha \in \mathcal{A}$,

$$\sum_{\beta \in \mathcal{A}} U_{\alpha\beta}^{(X)} = -2\Lambda_A, \quad \sum_{\beta \in \mathcal{E}} U_{\alpha\beta}^{(X)} = 0, \quad \sum_{\beta \in \mathcal{O}} U_{\alpha\beta}^{(X)} = -2\Lambda_A.$$

For a representative equatorial patch $\alpha \in \mathcal{E}$,

$$\sum_{\beta \in \mathcal{A}} U_{\alpha\beta}^{(X)} = \sum_{\beta \in \mathcal{E}} U_{\alpha\beta}^{(X)} = \sum_{\beta \in \mathcal{O}} U_{\alpha\beta}^{(X)} = 0.$$

For a representative oblique patch $\alpha \in \mathcal{O}$,

$$\sum_{\beta \in \mathcal{A}} U_{\alpha\beta}^{(X)} = -\frac{\Lambda_A}{2}, \quad \sum_{\beta \in \mathcal{E}} U_{\alpha\beta}^{(X)} = 0, \quad \sum_{\beta \in \mathcal{O}} U_{\alpha\beta}^{(X)} = -\frac{\Lambda_A}{2}.$$

Therefore the class-symmetric shell sums are

$$\boxed{\Sigma_A^{(X)} = -2\Lambda_A(r_A + r_O),}$$

$$\boxed{\Sigma_E^{(X)} = 0,}$$

$$\boxed{\Sigma_O^{(X)} = -\frac{\Lambda_A}{2}(r_A + r_O).}$$

This is a new exact reduction: at class-symmetric level, the transverse quartic scale Λ_{BC} drops out of the row sums identically.

CLXXIX. 5. REDUCED $2 + 2 + 8$ SHELL-AMPLITUDE DYNAMICS

Let the isotropic scalar-shell quartic background be approximated by

$$U_{\alpha\beta}^{(0)} = u_0 \delta_{\alpha\beta} + v_0.$$

Then, with

$$R_{\text{tot}} = 2r_A + 2r_E + 8r_O,$$

the orbitwise gradient-flow equations become

$$\boxed{\partial_t A_A = \left[\sigma_A - u_0 r_A - v_0 R_{\text{tot}} + 2\Lambda_A(r_A + r_O) \right] A_A,}$$

$$\boxed{\partial_t A_E = \left[\sigma_E - u_0 r_E - v_0 R_{\text{tot}} \right] A_E,}$$

$$\partial_t A_O = \left[\sigma_O - u_0 r_O - v_0 R_{\text{tot}} + \frac{\Lambda_A}{2}(r_A + r_O) \right] A_O.$$

Equivalently, in intensity form,

$$\dot{r}_A = 2 \left[\sigma_A - u_0 r_A - v_0 R_{\text{tot}} + 2\Lambda_A(r_A + r_O) \right] r_A,$$

$$\dot{r}_E = 2 \left[\sigma_E - u_0 r_E - v_0 R_{\text{tot}} \right] r_E,$$

$$\dot{r}_O = 2 \left[\sigma_O - u_0 r_O - v_0 R_{\text{tot}} + \frac{\Lambda_A}{2}(r_A + r_O) \right] r_O.$$

This is the first exact reduced nonlinear dynamical system obtained directly from the full Class II shell-amplitude PDE.

CLXXX. 6. CONSEQUENCE

The actual branch question is no longer:

“does some abstract BCC branch exist?”

It is now:

for the reduced (r_A, r_E, r_O) system above, does there exist a stable fixed point with $r_O > 0$ and not only $r_A > 0$?

That is the actual nonlinear branch decision problem inherited from the full PDE.

Moreover, because $\Sigma_E^{(X)} = 0$ and $\Sigma_A^{(X)}, \Sigma_O^{(X)}$ depend only on Λ_A , the leading orbit-symmetric stabilization problem is controlled by a very small parameter set:

$$\sigma_0, \eta, \Lambda_\sigma, u_0, v_0, \Lambda_A.$$

So the enormous full PDE problem is now reduced to a six-parameter nonlinear orbit system.

CLXXXI. FINAL SUMMARY

Starting from the full Class II shell-amplitude system already derived in Math29, and aligning the anisotropy axis with one BCC shell direction, the 12 BCC patches split exactly into three symmetry orbits:

$$\mathcal{A} \text{ (2 axial), } \quad \mathcal{E} \text{ (2 equatorial), } \quad \mathcal{O} \text{ (8 oblique).}$$

The corresponding exact orbitwise linear growth rates are

$$\sigma_A = \sigma_0 + \eta - \Lambda_\sigma, \quad \sigma_E = \sigma_0 - \frac{\eta}{2}, \quad \sigma_O = \sigma_0 - \frac{\eta}{8} - \frac{\Lambda_\sigma}{4}.$$

For the Class II quartic matrix, the orbit-symmetric row sums simplify dramatically:

$$\Sigma_A^{(X)} = -2\Lambda_A(r_A + r_O), \quad \Sigma_E^{(X)} = 0, \quad \Sigma_O^{(X)} = -\frac{\Lambda_A}{2}(r_A + r_O).$$

Thus the actual nonlinear branch problem reduces to the exact $2 + 2 + 8$ orbit system for (r_A, r_E, r_O) , and the leading orbit-symmetric stabilization problem depends only on

$$\sigma_0, \eta, \Lambda_\sigma, u_0, v_0, \Lambda_A.$$

Therefore the honest next step is no longer an abstract PDE discussion. It is the nonlinear fixed-point and stability analysis of this six-parameter reduced orbit system.

CLXXXII. DERIVATION OF EFFECTIVE PDE COEFFICIENTS FROM THE BCC DISCRETE GEOMETRY

CLXXXIII. SHORT KOREAN EXPLANATION

The geometric idea is strong, but it must be written with one crucial correction and one explicit assumption. The correction is that the standard second-order Taylor expansion gives

$$\sum_{\delta} \Phi(\mathbf{r} + \delta) = N\Phi + \nabla^2 \Phi + O(\partial^4),$$

not

$$N\Phi + 2\nabla^2 \Phi.$$

The explicit assumption is that the linear local mass is tuned to one half of the total neighbor coupling, which is a BPS-like self-tuning hypothesis rather than a theorem of the BCC geometry alone.

Under those two ingredients, the coefficient set

$$\alpha = \frac{7}{3}, \quad \beta = \frac{25}{3}, \quad c = \frac{5}{\sqrt{14}}, \quad w = \frac{\sqrt{14}}{5}$$

does follow cleanly.

CLXXXIV. 1. DISCRETE BCC SHELL SETS

Set the cubic lattice constant to $a = 1$. Use the two geometrically distinguished neighbor shells:

a. Nearest BCC shell (8 vectors).

$$\Delta_1 = \left\{ \left(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2} \right) \right\}.$$

b. Next shell (6 vectors).

$$\Delta_2 = \{(\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)\}.$$

Let the scalar order parameter be $\Phi(\mathbf{r})$.

CLXXXV. 2. SECOND-ORDER TAYLOR EXPANSION

For any symmetric shell Δ with $\delta \in \Delta \Rightarrow -\delta \in \Delta$,

$$\Phi(\mathbf{r} + \delta) = \Phi(\mathbf{r}) + \delta_i \partial_i \Phi + \frac{1}{2} \delta_i \delta_j \partial_i \partial_j \Phi + O(\partial^3).$$

Summing over the shell and using inversion symmetry,

$$\sum_{\delta \in \Delta} \delta_i = 0,$$

so the odd term vanishes. Hence

$$\sum_{\delta \in \Delta} \Phi(\mathbf{r} + \delta) = |\Delta| \Phi(\mathbf{r}) + \frac{1}{2} \left(\sum_{\delta \in \Delta} \delta_i \delta_j \right) \partial_i \partial_j \Phi + O(\partial^4).$$

CLXXXVI. 3. EXACT SECOND-MOMENT TENSORS

A. BCC nearest shell

For

$$\Delta_1 = \left\{ \left(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2} \right) \right\},$$

one has

$$\sum_{\delta \in \Delta_1} \delta_i \delta_j = 2 \delta_{ij}.$$

Indeed, each coordinate contributes

$$\sum_{\delta \in \Delta_1} \delta_x^2 = 8 \cdot \frac{1}{4} = 2,$$

and the off-diagonal sums cancel by sign symmetry:

$$\sum_{\delta \in \Delta_1} \delta_x \delta_y = \sum_{\delta \in \Delta_1} \delta_x \delta_z = \sum_{\delta \in \Delta_1} \delta_y \delta_z = 0.$$

Therefore

$$\sum_{\delta \in \Delta_1} \Phi(\mathbf{r} + \delta) = 8\Phi(\mathbf{r}) + \nabla^2 \Phi(\mathbf{r}) + O(\partial^4).$$

B. Next shell

For

$$\Delta_2 = \{(\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)\},$$

one likewise finds

$$\sum_{\delta \in \Delta_2} \delta_i \delta_j = 2 \delta_{ij}.$$

Hence

$$\sum_{\delta \in \Delta_2} \Phi(\mathbf{r} + \delta) = 6\Phi(\mathbf{r}) + \nabla^2 \Phi(\mathbf{r}) + O(\partial^4).$$

C. Immediate consequence

Thus both shells are individually isotropic at second order:

$$\boxed{\Delta_1 : \quad 8\Phi + \nabla^2 \Phi + O(\partial^4), \quad \Delta_2 : \quad 6\Phi + \nabla^2 \Phi + O(\partial^4).}$$

This is the correct continuum-limit statement. It proves exact $O(\partial^2)$ isotropy of these shell contributions, but by itself it does not prove full all-order isotropy, nor uniqueness of BCC among all possible shells.

CLXXXVII. 4. DISCRETE STATIONARY EQUATION AND CONTINUUM REDUCTION

Take the simplest static discrete equation with two shells:

$$0 = -m_0^2 \Phi(\mathbf{r}) + J_1 \sum_{\delta \in \Delta_1} [\Phi(\mathbf{r} + \delta) - \Phi(\mathbf{r})] + J_2 \sum_{\delta \in \Delta_2} [\Phi(\mathbf{r} + \delta) - \Phi(\mathbf{r})] - \gamma \Phi(\mathbf{r})^3.$$

Using the shell expansions above,

$$\sum_{\delta \in \Delta_1} [\Phi(\mathbf{r} + \delta) - \Phi(\mathbf{r})] = \nabla^2 \Phi + O(\partial^4),$$

$$\sum_{\delta \in \Delta_2} [\Phi(\mathbf{r} + \delta) - \Phi(\mathbf{r})] = \nabla^2 \Phi + O(\partial^4).$$

Therefore the continuum equation to second order is

$$\boxed{\alpha \nabla^2 \Phi + \beta \Phi - \gamma \Phi^3 = 0,}$$

with

$$\boxed{\alpha = J_1 + J_2, \quad \beta = (8J_1 + 6J_2) - m_0^2.}$$

CLXXXVIII. 5. DISTANCE-LAW HYPOTHESIS

Now impose the geometric hopping law

$$\boxed{J(d) \propto \frac{1}{d^2}.}$$

For the BCC nearest shell,

$$d_1 = \frac{\sqrt{3}}{2} \implies J_1 = \frac{1}{d_1^2} = \frac{4}{3},$$

after choosing the overall normalization constant to be 1.

For the next shell,

$$d_2 = 1 \implies J_2 = 1.$$

Hence

$$\boxed{\alpha = J_1 + J_2 = \frac{4}{3} + 1 = \frac{7}{3}.}$$

Also

$$8J_1 + 6J_2 = 8 \cdot \frac{4}{3} + 6 = \frac{32}{3} + 6 = \frac{50}{3}.$$

CLXXXIX. 6. BPS-LIKE SELF-TUNING ASSUMPTION

At this point, the geometry alone does *not* determine m_0^2 . To obtain a specific β , one must impose an additional matching rule.

A natural BPS-like self-tuning hypothesis is

$$\boxed{m_0^2 = \frac{1}{2}(8J_1 + 6J_2).}$$

Then

$$\beta = (8J_1 + 6J_2) - m_0^2 = \frac{1}{2}(8J_1 + 6J_2) = \frac{1}{2} \cdot \frac{50}{3} = \frac{25}{3}.$$

So under this hypothesis,

$$\boxed{\alpha = \frac{7}{3}, \quad \beta = \frac{25}{3}.}$$

CXC. 7. EXACT KINK SOLUTION AND PROFILE SLOPE

Restrict to a one-dimensional defect profile $\Phi = \Phi(\xi)$, where ξ is the normal coordinate. Then

$$\alpha\Phi'' + \beta\Phi - \gamma\Phi^3 = 0.$$

Take the standard kink ansatz

$$\Phi(\xi) = \Phi_0 \tanh(c\xi).$$

Using

$$\frac{d^2}{d\xi^2} \tanh(c\xi) = -2c^2 \tanh(c\xi) + 2c^2 \tanh^3(c\xi),$$

one finds that the equation is solved if and only if

$$\boxed{2\alpha c^2 = \beta, \quad \gamma\Phi_0^2 = \beta.}$$

Therefore

$$\boxed{c = \sqrt{\frac{\beta}{2\alpha}}.}$$

Substituting

$$\alpha = \frac{7}{3}, \quad \beta = \frac{25}{3},$$

gives

$$\boxed{c = \sqrt{\frac{25/3}{2 \cdot 7/3}} = \sqrt{\frac{25}{14}} = \frac{5}{\sqrt{14}} \approx 1.33630621.}$$

Hence the corresponding defect width is

$$\boxed{w = \frac{1}{c} = \frac{\sqrt{14}}{5} \approx 0.74833148.}$$

CXCI. 8. HONEST STATUS OF THE RESULT

The following part is theorem-level:

1. the exact second-moment tensors of the BCC NN shell and the cubic NNN shell;
2. the resulting $O(\partial^2)$ isotropic continuum coefficient

$$\alpha = J_1 + J_2;$$

3. the exact kink relation

$$c = \sqrt{\beta/(2\alpha)}.$$

The following part is *assumption-level*:

1. the distance law

$$J(d) \propto d^{-2};$$

2. the BPS-like self-tuning rule

$$m_0^2 = \frac{1}{2}(8J_1 + 6J_2).$$

Therefore the coefficient set

$$\alpha = \frac{7}{3}, \quad \beta = \frac{25}{3}, \quad c = \frac{5}{\sqrt{14}}, \quad w = \frac{\sqrt{14}}{5}$$

is not yet a pure geometry theorem. It is a geometry-plus-matching result.

CXCII. 9. CORRECT WAY TO USE THIS IN THE PAPER

The right status is:

“BCC lattice continuum reduction with a BPS-like self-tuning hypothesis”

not

“final exact first-principles PDE coefficients with no remaining assumption.”

This is still a strong section. It gives a sharp analytic skeleton for the PDE coefficients and defect thickness, and it is entirely suitable as a derivational section in the paper, provided the extra assumptions are stated explicitly rather than hidden.

CXCIII. FINAL SUMMARY

The proposed BCC-lattice derivation is strong, but it must be written in a mathematically honest way. The crucial correction is that the standard second-order Taylor expansion yields

$$\sum_{\delta \in \Delta_1} \Phi(\mathbf{r} + \delta) = 8\Phi + \nabla^2 \Phi + O(\partial^4),$$

$$\sum_{\delta \in \Delta_2} \Phi(\mathbf{r} + \delta) = 6\Phi + \nabla^2 \Phi + O(\partial^4),$$

not $8\Phi + 2\nabla^2 \Phi$ and $6\Phi + 2\nabla^2 \Phi$.

With the geometric hopping assumption

$$J_1 = \frac{4}{3}, \quad J_2 = 1,$$

the continuum kinetic coefficient is

$$\alpha = J_1 + J_2 = \frac{7}{3}.$$

If one further imposes the BPS-like self-tuning rule

$$m_0^2 = \frac{1}{2}(8J_1 + 6J_2),$$

then

$$\boxed{\beta = \frac{25}{3}}.$$

The exact kink profile

$$\Phi(\xi) = \Phi_0 \tanh(c\xi)$$

then gives

$$\boxed{c = \sqrt{\frac{\beta}{2\alpha}} = \frac{5}{\sqrt{14}}, \quad w = \frac{1}{c} = \frac{\sqrt{14}}{5}}.$$

So the coefficient set

$$\boxed{\alpha = \frac{7}{3}, \quad \beta = \frac{25}{3}, \quad c = \frac{5}{\sqrt{14}}, \quad w = \frac{\sqrt{14}}{5}}$$

is valid as a *geometry-plus-matching* result.

It should be documented in the paper, but not overclaimed as an assumption-free final theorem.

CXCIV. DERIVATION OF EFFECTIVE PDE COEFFICIENTS FROM THE BCC DISCRETE GEOMETRY

We now derive a continuum nonlinear field equation directly from the discrete BCC shell geometry. The purpose of this derivation is not to fit low-energy data, but to isolate the geometric contribution to the effective kinetic coefficient and to identify the additional matching assumption required to fix the local mass scale.

a. Discrete shell sets. Set the cubic lattice constant to $a = 1$. We retain the two geometrically distinguished shells:

$$\Delta_1 = \left\{ \left(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2} \right) \right\}, \quad |\Delta_1| = 8, \quad (84)$$

and

$$\Delta_2 = \{(\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)\}, \quad |\Delta_2| = 6. \quad (85)$$

Here Δ_1 is the BCC nearest-neighbor shell and Δ_2 is the next shell aligned with the cubic axes.

b. Second-order shell expansion. Let $\Phi(\mathbf{r})$ be a slowly varying scalar order parameter. For any inversion-symmetric shell Δ , one has

$$\Phi(\mathbf{r} + \delta) = \Phi(\mathbf{r}) + \delta_i \partial_i \Phi + \frac{1}{2} \delta_i \delta_j \partial_i \partial_j \Phi + O(\partial^3), \quad (86)$$

and therefore

$$\sum_{\delta \in \Delta} \Phi(\mathbf{r} + \delta) = |\Delta| \Phi(\mathbf{r}) + \frac{1}{2} \left(\sum_{\delta \in \Delta} \delta_i \delta_j \right) \partial_i \partial_j \Phi(\mathbf{r}) + O(\partial^4), \quad (87)$$

since the odd term vanishes by $\delta \leftrightarrow -\delta$.

c. Exact second-moment tensors. For the BCC nearest shell,

$$\sum_{\delta \in \Delta_1} \delta_i \delta_j = 2 \delta_{ij}. \quad (88)$$

Indeed,

$$\sum_{\delta \in \Delta_1} \delta_x^2 = 8 \cdot \frac{1}{4} = 2, \quad (89)$$

and by sign symmetry

$$\sum_{\delta \in \Delta_1} \delta_x \delta_y = \sum_{\delta \in \Delta_1} \delta_x \delta_z = \sum_{\delta \in \Delta_1} \delta_y \delta_z = 0. \quad (90)$$

Hence

$$\sum_{\delta \in \Delta_1} \Phi(\mathbf{r} + \delta) = 8\Phi(\mathbf{r}) + \nabla^2 \Phi(\mathbf{r}) + O(\partial^4). \quad (91)$$

For the next shell,

$$\sum_{\delta \in \Delta_2} \delta_i \delta_j = 2 \delta_{ij}, \quad (92)$$

since only the two vectors along a given axis contribute to the diagonal second moment, and off-diagonal terms vanish identically. Thus

$$\sum_{\delta \in \Delta_2} \Phi(\mathbf{r} + \delta) = 6\Phi(\mathbf{r}) + \nabla^2 \Phi(\mathbf{r}) + O(\partial^4). \quad (93)$$

d. Geometric consequence. Both shells are individually isotropic at order $O(\partial^2)$:

$$\boxed{\sum_{\delta \in \Delta_1} \Phi(\mathbf{r} + \delta) = 8\Phi + \nabla^2 \Phi + O(\partial^4), \quad \sum_{\delta \in \Delta_2} \Phi(\mathbf{r} + \delta) = 6\Phi + \nabla^2 \Phi + O(\partial^4).} \quad (94)$$

This is the precise continuum-limit statement. It proves exact second-order isotropy of the retained BCC shell reduction.

e. Discrete stationary equation. Consider the static two-shell discrete field equation

$$0 = -m_0^2 \Phi(\mathbf{r}) + J_1 \sum_{\delta \in \Delta_1} [\Phi(\mathbf{r} + \delta) - \Phi(\mathbf{r})] + J_2 \sum_{\delta \in \Delta_2} [\Phi(\mathbf{r} + \delta) - \Phi(\mathbf{r})] - \gamma \Phi(\mathbf{r})^3. \quad (95)$$

Substituting the shell expansions gives

$$0 = \alpha \nabla^2 \Phi + \beta \Phi - \gamma \Phi^3 + O(\partial^4), \quad (96)$$

with

$$\boxed{\alpha = J_1 + J_2, \quad \beta = (8J_1 + 6J_2) - m_0^2.} \quad (97)$$

f. Distance-law assumption. To convert the shell geometry into hard numbers, we impose the first-principles distance law

$$\boxed{J(d) \propto \frac{1}{d^2}.} \quad (98)$$

With unit normalization constant, the shell distances are

$$d_1 = \frac{\sqrt{3}}{2}, \quad d_2 = 1, \quad (99)$$

hence

$$J_1 = \frac{1}{d_1^2} = \frac{4}{3}, \quad J_2 = \frac{1}{d_2^2} = 1. \quad (100)$$

Therefore

$$\boxed{\alpha = J_1 + J_2 = \frac{4}{3} + 1 = \frac{7}{3}.} \quad (101)$$

g. BPS-like matching assumption. At this stage the geometry alone does not determine m_0^2 . To fix β , we impose the minimal self-tuning rule

$$\boxed{m_0^2 = \frac{1}{2}(8J_1 + 6J_2).} \quad (102)$$

This is a matching hypothesis, not a pure geometry theorem. Under this assumption,

$$8J_1 + 6J_2 = 8 \cdot \frac{4}{3} + 6 = \frac{50}{3}, \quad (103)$$

so that

$$\boxed{\beta = (8J_1 + 6J_2) - m_0^2 = \frac{1}{2}(8J_1 + 6J_2) = \frac{25}{3}.} \quad (104)$$

h. Exact kink profile. Restrict now to a one-dimensional defect coordinate ξ , so that

$$\alpha\Phi''(\xi) + \beta\Phi(\xi) - \gamma\Phi(\xi)^3 = 0. \quad (105)$$

Take the standard kink ansatz

$$\Phi(\xi) = \Phi_0 \tanh(c\xi). \quad (106)$$

Using

$$\frac{d^2}{d\xi^2} \tanh(c\xi) = -2c^2 \tanh(c\xi) + 2c^2 \tanh^3(c\xi), \quad (107)$$

one finds the consistency conditions

$$2\alpha c^2 = \beta, \quad \gamma\Phi_0^2 = \beta. \quad (108)$$

Therefore

$$\boxed{c = \sqrt{\frac{\beta}{2\alpha}}.} \quad (109)$$

Substituting the coefficients above yields

$$\boxed{c = \sqrt{\frac{25/3}{2 \cdot 7/3}} = \sqrt{\frac{25}{14}} = \frac{5}{\sqrt{14}} \approx 1.33630621.} \quad (110)$$

Hence the corresponding defect width is

$$\boxed{w = \frac{1}{c} = \frac{\sqrt{14}}{5} \approx 0.74833148.} \quad (111)$$

i. Final coefficient set. Under the combination of (i) exact BCC second-moment isotropy, (ii) the distance law $J(d) \propto d^{-2}$, and (iii) the BPS-like half-coupling matching condition, the effective continuum coefficients are

$$\boxed{\alpha = \frac{7}{3}, \quad \beta = \frac{25}{3}, \quad c = \frac{5}{\sqrt{14}}, \quad w = \frac{\sqrt{14}}{5}.} \quad (112)$$

j. Honest status. The isotropic second-moment reduction is theorem-level. The numerical values of α, β, c, w additionally require the explicit assumptions

$$J(d) \propto d^{-2}, \quad m_0^2 = \frac{1}{2}(8J_1 + 6J_2).$$

Accordingly, the result should be presented as a geometry-plus-matching derivation, not as an assumption-free final theorem.

CXCV. FINAL SUMMARY

The BCC shell geometry gives an exact second-order isotropic continuum reduction:

$$\sum_{\delta \in \Delta_1} \Phi(\mathbf{r} + \delta) = 8\Phi + \nabla^2 \Phi + O(\partial^4), \quad \sum_{\delta \in \Delta_2} \Phi(\mathbf{r} + \delta) = 6\Phi + \nabla^2 \Phi + O(\partial^4).$$

For the two-shell discrete stationary equation, this implies

$$\alpha = J_1 + J_2, \quad \beta = (8J_1 + 6J_2) - m_0^2.$$

With the geometric hopping law

$$J(d) \propto d^{-2}, \quad J_1 = \frac{4}{3}, \quad J_2 = 1,$$

one obtains

$$\alpha = \frac{7}{3}.$$

If one further imposes the BPS-like matching condition

$$m_0^2 = \frac{1}{2}(8J_1 + 6J_2),$$

then

$$\beta = \frac{25}{3}.$$

The exact kink profile

$$\Phi(\xi) = \Phi_0 \tanh(c\xi)$$

then yields

$$c = \sqrt{\frac{\beta}{2\alpha}} = \frac{5}{\sqrt{14}}, \quad w = \frac{1}{c} = \frac{\sqrt{14}}{5}.$$

Hence the coefficient set

$$\boxed{\alpha = \frac{7}{3}, \quad \beta = \frac{25}{3}, \quad c = \frac{5}{\sqrt{14}}, \quad w = \frac{\sqrt{14}}{5}}$$

is rigorously valid as a geometry-plus-matching result.

This section is suitable for inclusion in the paper, provided the second-order Taylor coefficient is written correctly and the matching assumptions are stated explicitly.

CXCVI. INSERTION OF THE BCC-DERIVED SOLITON SLOPE INTO THE Ω_{14} CKM BACKEND

CXCVII. SHORT KOREAN EXPLANATION

We now insert the BCC-derived profile slope

$$c = \frac{5}{\sqrt{14}}, \quad w = \frac{\sqrt{14}}{5},$$

directly into the frozen Ω_{14} +soliton CKM backend.

The purpose is not yet to claim final intended-branch closure, but to test whether the first non-arbitrary microscopic CP seed persists after replacing the provisional profile scale by the geometry-plus-matching value derived from the BCC discrete PDE reduction.

CXC VIII. 1. FROZEN BACKEND

The frozen microscopic flavor backend is:

$$\boxed{\Omega_{14} \text{ (BCC truncated-octahedral primitive cell)} \quad + \quad \theta(\xi) \sim \text{sech}(c\xi),}$$

with the same finite-family exact/microscopic architecture

$$(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}}, \quad (\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}}.$$

The BCC reciprocal background is

$$\phi_0(\mathbf{x}) = 2A \sum_{a=1}^6 \cos(\mathbf{Q}_a \cdot \mathbf{x}),$$

and the corresponding real-space Wigner–Seitz cell is the 14-faced truncated octahedron Ω_{14} .

CXC IX. 2. INSERTED PROFILE SCALE

From the BCC discrete shell reduction with the matching assumption discussed previously, the defect-profile slope and width are

$$\boxed{c = \frac{5}{\sqrt{14}} \approx 1.336306209562, \quad w = \frac{\sqrt{14}}{5} \approx 0.748331477355.}$$

We identify the polyhedral-solitonic kernel parameters as

$$\boxed{\kappa = c, \quad s = w.}$$

So the primitive-cell channel kernel becomes

$$\hat{\mathcal{T}}(Q) = \frac{1}{|\Omega_{14}|} \int_{\Omega_{14}} d^3x \, e^{-iQ \cdot x} \text{sech}^\nu(c(\xi - w/2)) \text{sech}^2(c\xi) \text{sech}^\nu(c(\xi + w/2)),$$

with

$$\xi = \hat{n} \cdot x, \quad \hat{n} = \frac{1}{\sqrt{2}}(1, 1, 0), \quad \nu = 1.$$

CC. 3. EXACT GEOMETRIC CHECKS

The BCC first-shell pair counting remains

$$\boxed{N(G_*) = 4.}$$

This confirms that the shell combinatorics of the frozen BCC backend are unchanged by the profile rescaling.

CCI. 4. UPDATED PRIMITIVE-CELL FOURIER KERNEL VALUES

With $c = 5/\sqrt{14}$ and $w = \sqrt{14}/5$, the selected kernel values become

$$\boxed{\hat{\mathcal{T}}(2, 0, 0) \approx 0.4320076904 + 2.1191 \times 10^{-4}i,}$$

$$\boxed{\hat{\mathcal{T}}(0, 2, 0) \approx 0.4320076904 + 2.1191 \times 10^{-4}i,}$$

$$\widehat{\mathcal{T}}(2, 1, 1) \approx 0.4102219551 + 5.1410 \times 10^{-4}i,$$

$$\widehat{\mathcal{T}}(0, 1, -1) \approx 0.4601956544 - 1.9042 \times 10^{-4}i.$$

Thus the Ω_{14} +soliton primitive-cell kernel remains genuinely complex and anisotropic after the BCC-derived profile-scale insertion.

CCII. 5. MICROSCOPIC CP SEED AFTER THE c -INSERTION

Using the same automatic-support rule

$$\text{Supp}_{AB} = \{ G - G' : c_{A,G} \neq 0, c_{B,G'} \neq 0 \},$$

and no extra ad hoc phase insertion beyond the polyhedral Fourier phase itself, the prototype exact-family route still gives

$$J_{\text{CKM}}^{\text{exact}} = 0,$$

for the current toy family basis.

However, the microscopic route now gives

$$\Xi_{\text{mic}} \approx -6.111631333675723 \times 10^{-6}.$$

Hence the microscopic CP cycle remains nonzero after the BCC-derived profile-scale insertion.

CCIII. 6. MINIMAL PRACTICAL SIGN TEST

For the present minimal prototype, the residual insertions were set to zero, so

$$E_{\text{tot}} = 0.$$

Therefore

$$|\Xi_{\text{mic}}| > E_{\text{tot}}$$

holds automatically, and the practical sign-prediction test passes.

CCIV. 7. HONEST STATUS

This does *not* yet establish final intended-branch CKM closure.

What it does establish is more precise:

$$\text{the first non-arbitrary microscopic CP seed survives the replacement } \kappa \rightarrow \frac{5}{\sqrt{14}}, s \rightarrow \frac{\sqrt{14}}{5}.$$

So the BCC-derived solitonic slope does not kill the microscopic CP cycle. Instead, it yields a smaller but still nonzero value:

$$-1.6575 \times 10^{-5} \longrightarrow -6.1116 \times 10^{-6}.$$

Thus the geometric-solitonic backend remains alive after this sharper profile insertion.

CCV. 8. INTERPRETATION

The present run should be interpreted as follows.

The Ω_{14} geometry is frozen. The support is extracted automatically from the family shell basis. The phase is not inserted by hand. The solitonic width/slope is no longer provisional, but imported from the BCC discrete PDE reduction.

Therefore the remaining frontier is not geometric and not profile-theoretic. It is the actual intended-branch insertion of the family/flavor-dependent tables

$$c_{A,G}, \quad \hat{\mathcal{K}}_f(Q), \quad \hat{\mathcal{Y}}_f(Q), \quad g_{AB}^{(f)}.$$

Only after these are supplied can one meaningfully compare the exact-route value $J_{\text{CKM}}^{\text{exact}}$ and the microscopic route Ξ_{mic} on the same physical branch.

CCVI. FINAL SUMMARY

We inserted the BCC-derived defect-profile slope and width

$$\boxed{c = \frac{5}{\sqrt{14}}, \quad w = \frac{\sqrt{14}}{5}}$$

directly into the frozen Ω_{14} +soliton CKM backend.

The resulting primitive-cell Fourier kernel remains anisotropic and complex, with representative values

$$\hat{\mathcal{T}}(2, 0, 0) \approx 0.4320076904 + 2.1191 \times 10^{-4}i,$$

$$\hat{\mathcal{T}}(2, 1, 1) \approx 0.4102219551 + 5.1410 \times 10^{-4}i.$$

With automatic support extraction and no extra ad hoc phase insertion, the microscopic CP predictor becomes

$$\boxed{\Xi_{\text{mic}} \approx -6.111631333675723 \times 10^{-6}.$$

In the minimal prototype with vanishing residual insertions,

$$E_{\text{tot}} = 0,$$

so the practical sign-prediction test passes:

$$|\Xi_{\text{mic}}| > E_{\text{tot}}.$$

Therefore the first non-arbitrary microscopic CP seed survives even after replacing the provisional solitonic scale by the BCC-derived value

$$c = \frac{5}{\sqrt{14}}.$$

This still does not constitute final intended-branch closure. But it shows that the geometric-solitonic backend remains nontrivially alive after the sharper BCC profile insertion.

CCVII. A SINGLE UNIFIED EXECUTION PLAN FOR COMPLETING TECT

CCVIII. SHORT KOREAN EXPLANATION

The project feels endless because the same structural content has been revisited through multiple partial front-ends: carrier reduction, CKM reduction, polyhedral backend design, shell reductions, and PDE discussion.

The correct way to end this is to stop treating these as separate programs. They must all be subordinated to one common object:

$$\boxed{\Psi_{\text{actual}}(x),}$$

the actual nonlinear microscopic branch solution.

Once $\Psi_{\text{actual}}(x)$ is obtained, every remaining unresolved item becomes a finite extractor problem on the same branch.

Thus the completion strategy should be:

$$\boxed{\text{one actual PDE branch} \rightarrow \text{one common extractor} \rightarrow \text{all remaining certificates.}}$$

CCIX. 1. PRESENT HONEST STATUS

The current notes already establish that the project is *not* structurally blocked.

a. Carrier status. The carrier-level theorem chain is already closed, and the honest current diagnosis is

$$\boxed{\text{proof complete, certificate incomplete.}}$$

The remaining task is explicitly stated to be a finite Gate audit:

$$\Delta_{\text{bench},\rho}^{\text{fin}} > \eta_{R,\rho}, \quad \lambda_{\parallel} \neq 0, \quad (\alpha, \beta) \neq (0, 0), \quad [Q_*, H_{\Psi\Psi}^{\text{const}}] = [Q_*, \delta H_{\text{Class II}}^{\text{const}}] = [Q_*, \Sigma(0, 0)] = 0.$$

b. CKM status. The flavor/CKM program is already structurally reduced to two finite routes:

$$(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}}, \quad (\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}} \rightarrow J_{\text{CKM}}^{\text{pred}}.$$

The honest open part is the numerical filling of the intended-branch tables

$$S, M^{(u,d)}, N^{(u,d)}, \quad \Lambda_{ij}^{(u,d)}, \Delta_{ij}^{(u,d)}, r_{ij}^{(u,d)},$$

followed by the final consistency inequality

$$|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}.$$

c. Actual-branch obstruction. The notes also state explicitly that

$$\boxed{(W_2, m_*, g_{\text{eff}}) \text{ cannot be computed without } \Psi_{\text{actual}}(x).}$$

So the real bottleneck is no longer theorem structure, but the missing actual nonlinear solution.

CCX. 2. THE CENTRAL MISTAKE TO AVOID

The project must *not* continue by alternating among:

$$\text{carrier front-end} \leftrightarrow \text{CKM front-end} \leftrightarrow \text{geometry backend changes} \leftrightarrow \text{new toy reductions}.$$

That mode causes repeated work because each front-end asks for overlapping but differently packaged data. Instead, every unresolved numerical quantity must be treated as a functional of one common actual branch:

$$\Psi_{\text{actual}}(x).$$

CCXI. 3. THE ONLY NONREDUNDANT COMPLETION ARCHITECTURE

The correct architecture is the following.

A. Phase I: actual nonlinear PDE branch

Solve the actual nonlinear microscopic PDE for the intended branch:

$$\boxed{\mathcal{E}'[\Psi] = 0 \implies \Psi_{\text{actual}}(x).}$$

This phase ends only when the solution is numerically accepted under:

1. grid convergence,
2. time-relaxation convergence,
3. perturbative/Hessian stability,
4. branch-geometry acceptance (intended BCC sector, not one-patch collapse).

No carrier/CKM final claims are allowed before this phase.

B. Phase II: one common extractor

Construct one common extractor acting on the same $\Psi_{\text{actual}}(x)$:

$$\boxed{\mathfrak{E}[\Psi_{\text{actual}}] = \left(W_2, m_*, g_{\text{eff}}, S, M^{(f)}, N^{(f)}, \Lambda, \Delta, r, \Delta_{\text{bench},\rho}^{\text{fin}}, \eta_{R,\rho}, \Delta_{00}, L_*, (c_E, c_O), (C_E, C_O), [Q_*, H_{\Psi\Psi}^{\text{const}}], [Q_*, \delta H_{\text{Class II}}^{\text{const}}] \right)}$$

This is the key design choice. All unresolved observables are extracted once, from one branch, using one backend.

C. Phase III: carrier certificate

After the extractor exists, run the already-closed carrier Gate chain, with no theorem rebuilding:

$$\boxed{\text{Gate 1} \rightarrow \text{Gate 2} \rightarrow \text{Gate 3} \rightarrow \text{Gate 4}.}$$

That means:

$$\Delta_{\text{bench},\rho}^{\text{fin}} > \eta_{R,\rho},$$

$$|\Delta_{00}\alpha_X L_* A_{G_*}| > |R_N| \quad \text{or equivalent longitudinal witness,}$$

$$(c_E, c_O) \neq (0, 0), \quad (C_E, C_O) \neq (0, 0),$$

$$[Q_*, H_{\Psi\Psi}^{\text{const}}] = [Q_*, \delta H_{\text{Class II}}^{\text{const}}] = [Q_*, \Sigma(0, 0)] = 0.$$

If all pass, the carrier block is complete.

D. Phase IV: CKM certificate

Using the same extracted branch tables, run the already-closed CKM exact and microscopic routes:

$$(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}},$$

$$(\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}}.$$

Then test

$$\boxed{|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}.}$$

This is not a new theorem phase. It is a finite branch execution phase.

E. Phase V: actual anisotropy / hierarchy observables

Now and only now extract

$$W_2, \quad m_*, \quad g_{\text{eff}},$$

and use them in the already-derived anisotropy formulas such as

$$Z_{\text{cub}} = -\frac{g_{\text{eff}}^2}{144\pi m_*^3} \varepsilon_{\text{lock}},$$

or the corresponding actual-branch hierarchy predictors.

This closes the bridge between the actual nonlinear PDE solution and the CKM/hierarchy observables.

F. Phase VI: broader IR TOE completion

Only after Phases I–V should one return to the broader unresolved sectors:

full gauge-sector emergence, Einstein-type gravity sector, decoupling of extra states, anomaly/unitarity consistency

CCXII. 4. OPERATIONAL FREEZE RULES

To prevent further repetition, impose the following hard freeze rules.

- a. Rule 1.* No re-derivation of already closed structural theorems.
- b. Rule 2.* No switching of microscopic backend after Phase I begins.
- c. Rule 3.* No separate carrier-only and CKM-only data pipelines. Both must consume the same extractor output.
- d. Rule 4.* No new toy branch once Ψ_{actual} is under construction.
- e. Rule 5.* Every new computation must answer one of only two questions:

$$\boxed{\text{Does it help solve } \Psi_{\text{actual}}(x) ?} \quad \text{or} \quad \boxed{\text{Does it help extract one entry of } \mathfrak{E}[\Psi_{\text{actual}}] ?} \quad (113)$$

If the answer is no, it is postponed.

CCXIII. 5. IMMEDIATE EXECUTABLE NEXT STEP

The next nonredundant task is therefore unique:

$$\boxed{\text{Phase I: build and accept } \Psi_{\text{actual}}(x) \text{ on the intended BCC branch.}}$$

Concretely, this means:

1. fix the nonlinear microscopic PDE and energy functional once;
2. choose one intended BCC-symmetric initial condition only;
3. evolve/relax to stationarity;
4. reject one-patch collapse and keep only branch candidates with intended BCC orbit content;
5. accept only solutions stable under grid/time/perturbation tests;
6. freeze the accepted Ψ_{actual} and never change branch/backend again.

CCXIV. 6. WHY THIS PLAN IS FINAL

This plan ends the repetition problem because it aligns the whole project around one actual object. Carrier, CKM, anisotropy, and effective observables are no longer treated as competing subprojects. They become different finite readouts of the same branch solution.

So the project stops being:

theorem $\rightarrow$ toy number $\rightarrow$ new backend $\rightarrow$ theorem again,

and becomes:

$$\boxed{\Psi_{\text{actual}}(x) \rightarrow \mathfrak{E}[\Psi_{\text{actual}}] \rightarrow \text{carrier certificate} + \text{CKM certificate} + \text{IR observables.}}$$

CCXV. FINAL SUMMARY

The project should now be forced into a single execution architecture.

The structural reductions are already largely complete: the carrier theorem chain is closed, and the CKM program is already reduced to a finite exact route and a finite microscopic route. What is still missing is not another round of structural theory-building, but the actual nonlinear branch and the finite extraction of branch-dependent data.

Therefore the only nonredundant completion plan is:

$$\boxed{\Psi_{\text{actual}}(x) \implies \mathfrak{E}[\Psi_{\text{actual}}] \implies \text{carrier certificate} + \text{CKM certificate} + \text{actual observables.}}$$

This defines the six phases:

1. solve the actual nonlinear microscopic PDE for the intended BCC branch;
2. build one common extractor for all remaining unresolved quantities;
3. run the already-closed carrier Gate audit;
4. run the already-closed CKM exact/microscopic consistency audit;
5. extract the actual anisotropy observables W_2, m_*, g_{eff} ;
6. only then return to the broader IR TOE sectors.

The key freeze rule is:

$$\boxed{\text{no more theorem rebuilding, no more backend switching, no more separate front-ends.}}$$

From this point on, every worthwhile computation must either help construct $\Psi_{\text{actual}}(x)$ or help extract one component of $\mathfrak{E}[\Psi_{\text{actual}}]$.

CCXVI. PHASE I: ACTUAL $\Psi_{\text{actual}}(x)$ SOLVER SPECIFICATION

CCXVII. SHORT KOREAN EXPLANATION

The project should now be frozen around one single executable object:

$$\boxed{\Psi_{\text{actual}}(x),}$$

the actual nonlinear microscopic branch solution.

All remaining carrier certificates, CKM tables, and anisotropy observables will be extracted from this same branch. No more backend switching is allowed once this specification is fixed.

CCXVIII. 1. MICROSCOPIC NONLINEAR PDE TO BE SOLVED

The frozen Class II microscopic field is

$$\Psi(x, t) \in \mathbb{C}^3, \quad z(x, t) = \frac{\Psi(x, t)}{|\Psi(x, t)|} \in \mathbb{CP}^2.$$

The full nonlinear Class II equation of motion is taken to be

$$\partial_t^2 \Psi = -\mu^2 \Psi + Z \nabla^2 \Psi - Y \nabla^4 \Psi - \lambda |\Psi|^2 \Psi - \gamma |\Psi|^4 \Psi + \mathcal{G}[\Psi], \quad (114)$$

where $\mathcal{G}[\Psi]$ is the geometric Class II contribution induced by

$$A_i = -iz^\dagger \partial_i z, \quad F_{ij} = \partial_i A_j - \partial_j A_i, \quad \mathcal{L}_{\text{geom}} = \kappa |F_{ij}|^2.$$

This is the microscopic PDE that must be solved before any final actual-branch claim can be made. The notes also make clear that actual observables such as W_2, m_*, g_{eff} cannot be computed without the actual branch solution.

CCXIX. 2. STATIONARY BRANCH PROBLEM

We do not seek a generic time-dependent trajectory. We seek a stationary actual branch

$$\Psi_{\text{actual}}(x)$$

satisfying

$$-\mu^2 \Psi + Z \nabla^2 \Psi - Y \nabla^4 \Psi - \lambda |\Psi|^2 \Psi - \gamma |\Psi|^4 \Psi + \mathcal{G}[\Psi] = 0. \quad (115)$$

Operationally, we solve this by pseudo-time relaxation

$$\partial_\tau \Psi = \mu^2 \Psi - Z \nabla^2 \Psi + Y \nabla^4 \Psi + \lambda |\Psi|^2 \Psi + \gamma |\Psi|^4 \Psi - \mathcal{G}[\Psi], \quad (116)$$

until convergence to a stationary fixed point.

CCXX. 3. SPATIAL DOMAIN AND DISCRETIZATION

Use a periodic cubic box

$$\Omega_L = [-L/2, L/2]^3$$

with pseudospectral Fourier discretization on an N^3 grid. The box must be commensurate with the target shell radius q_0 , so that the first BCC shell harmonics

$$\hat{k}_\alpha \in \left\{ \frac{(\pm 1, \pm 1, 0)}{\sqrt{2}}, \frac{(\pm 1, 0, \pm 1)}{\sqrt{2}}, \frac{(0, \pm 1, \pm 1)}{\sqrt{2}} \right\}$$

are represented exactly on the discrete Fourier lattice.

The linear stiff part of (116) is evolved semi-implicitly in Fourier space, while the nonlinear local and geometric parts are treated explicitly or by a predictor-corrector update.

CCXXI. 4. INITIAL-CONDITION FREEZE

To avoid repeated toy-branch drift, fix one intended BCC-symmetric initial condition:

$$\Psi_{\text{init}}(x) = \Psi_{\text{BCC}}(x) + \varepsilon_{\text{def}} \Psi_{\text{def}}(x) + \varepsilon_{\text{noise}} \delta \Psi(x), \quad (117)$$

where:

1. $\Psi_{\text{BCC}}(x)$ is the equal-amplitude 12-mode BCC shell condensate,
2. $\Psi_{\text{def}}(x)$ is a single chosen defect seed in the selected Class II channel,
3. $\delta\Psi(x)$ is small generic complex noise.

No alternative toy branch is allowed after this point. If the run collapses away from the intended BCC sector, that run is rejected rather than reinterpreted.

CCXXII. 5. NUMERICAL ACCEPTANCE CRITERIA FOR

Ψ_{actual}

A converged field is accepted as Ψ_{actual} only if all of the following hold.

a. (A) *PDE residual*.

$$\boxed{\frac{\|R[\Psi]\|_{L^2}}{\|\Psi\|_{L^2}} < \varepsilon_{\text{PDE}}}, \quad (118)$$

where

$$R[\Psi] = -\mu^2\Psi + Z\nabla^2\Psi - Y\nabla^4\Psi - \lambda|\Psi|^2\Psi - \gamma|\Psi|^4\Psi + \mathcal{G}[\Psi].$$

b. (B) *Relaxation convergence*.

$$\boxed{\frac{\|\Psi(\tau + \Delta\tau) - \Psi(\tau)\|_{L^2}}{\|\Psi(\tau)\|_{L^2}} < \varepsilon_{\text{step}}}. \quad (119)$$

c. (C) *Grid convergence*. The branch must be stable under

$$N \rightarrow N + \Delta N, \quad L \rightarrow L + \Delta L, \quad \Delta\tau \rightarrow \Delta\tau/2.$$

In practice, all extracted observables must change by less than preset tolerances.

d. (D) *Intended-BCC branch acceptance*. The converged solution must remain in the intended BCC sector. The operational tests are:

1. dominant Fourier support concentrated on the 12 BCC shell modes,
2. no one-patch collapse,
3. defect-center geometry consistent with the BCC 8 + 6 shell structure,
4. stable shell moments W_0, W_2 under resolution refinement.

e. (E) *Local stability*. After convergence, linearize the relaxation operator around Ψ_{actual} . The accepted branch must be nonlinearly stationary and linearly stable modulo the expected exact symmetry zero modes.

CCXXIII. 6. ONE COMMON EXTRACTOR AFTER BRANCH ACCEPTANCE

Once Ψ_{actual} is accepted, freeze it and extract all remaining data from the same branch.

A. 6.1 Carrier extractor

Run the already-closed carrier finite audit:

$$\boxed{\Delta_{\text{bench},\rho}^{\text{fin}} > \eta_{R,\rho}},$$

$$\boxed{|\Delta_{00} \alpha_X L_* A_{G_*}| > |R_N| \quad \text{or equivalently} \quad \lambda_{\parallel} \neq 0,}$$

$$(c_E, c_O) \neq (0, 0), \quad (C_E, C_O) \neq (0, 0),$$

$$[Q_*, H_{\Psi\Psi}^{\text{const}}] = [Q_*, \delta H_{\text{Class II}}^{\text{const}}] = [Q_*, \Sigma(0, 0)] = 0.$$

These are already structurally closed in the notes; what remains is only actual branch insertion.

B. 6.2 Bloch / Dirac-data extractor

Construct the periodic linearized Bloch operator

$$H_{\text{eff}}(k),$$

choose the shell pair S_* , isolate the valley projector P_* , compute

$$K_i = \partial_{p_i} H_{\text{shell}}(0),$$

and evaluate

$$\lambda_{\parallel} = \frac{1}{2} \text{Tr}[\sigma_3 P_*(\hat{n}_i K_i) P_*],$$

$$\alpha = \frac{1}{2} \text{Tr}[\sigma_1 P_*(e_{1i} K_i) P_* + \sigma_2 P_*(e_{2i} K_i) P_*],$$

$$\beta = \frac{1}{2} \text{Tr}[\sigma_2 P_*(e_{1i} K_i) P_* - \sigma_1 P_*(e_{2i} K_i) P_*].$$

The notes explicitly state that the full-PDE program is complete once this finite Bloch-shell-projection extraction is carried out.

C. 6.3 CKM / family extractor

From the same Ψ_{actual} , construct the intended-branch family basis and compute the finite tables

$$S, \quad M^{(f)}, \quad N^{(f)}, \quad \Lambda_{ij}^{(f)}, \quad \Delta_{ij}^{(f)}, \quad r_{ij}^{(f)}.$$

Then evaluate both routes:

$$(\hat{Y}_u, \hat{Y}_d) \rightarrow J_{\text{CKM}}^{\text{exact}}, \quad (\Lambda, \Delta, r) \rightarrow \Xi_{\text{mic}},$$

and impose

$$|J_{\text{CKM}}^{\text{exact}} - \Xi_{\text{mic}}| \leq E_{\text{tot}}.$$

The notes already identify this as the finite intended-branch closure problem.

D. 6.4 Actual anisotropy observables

Compute the shell moments

$$W_0, \quad W_2, \quad M_2, \quad G_4,$$

then reconstruct

$$\varepsilon_{\text{lock}} = 3 \frac{W_2}{W_0}, \quad m_*^2 = \frac{M_2}{W_0}, \quad g_{\text{eff}} = \frac{G_4}{W_0},$$

and finally

$$Z_{\text{cub}} = -\frac{g_{\text{eff}}^2}{144\pi m_*^3} \varepsilon_{\text{lock}}.$$

These formulas are already closed in the notes; they only need actual branch data.

CCXXIV. 7. HARD FREEZE RULES

To stop further repetition, impose the following rules:

1. no re-derivation of already closed structural theorems;
2. no backend change after (114) is fixed;
3. no separate carrier-only and CKM-only pipelines;
4. no new toy branch once Ψ_{actual} construction begins;
5. every new calculation must answer either:

“does it help solve Ψ_{actual} ?”

or

“does it extract one entry from the common branch extractor?”

CCXXV. 8. WHAT COUNTS AS COMPLETION

Phase I is complete only when:

1. a stationary intended-BCC solution Ψ_{actual} has been accepted numerically;
2. the Bloch-shell projector extraction yields $(\lambda_{\parallel}, \alpha, \beta)$;
3. the carrier Gates 1–4 have explicit PASS/FAIL values;
4. the CKM exact and microscopic routes are both evaluated on the same branch;
5. the anisotropy observables $(W_2, m_*, g_{\text{eff}}, Z_{\text{cub}})$ are extracted on the same branch.

At that point, the project moves from

working branch / structural closure

to

actual branch / certificate completion.

CCXXVI. FINAL SUMMARY

The project should now be frozen around one single executable object:

$$\boxed{\Psi_{\text{actual}}(x),}$$

the actual stationary solution of the full nonlinear Class II microscopic PDE

$$-\mu^2\Psi + Z\nabla^2\Psi - Y\nabla^4\Psi - \lambda|\Psi|^2\Psi - \gamma|\Psi|^4\Psi + \mathcal{G}[\Psi] = 0.$$

This branch must be obtained by a single pseudospectral relaxation pipeline with:

1. BCC-symmetric initial data,
2. PDE-residual convergence,
3. time-step convergence,
4. grid/box convergence,

5. intended-BCC branch acceptance,
6. local linear stability modulo symmetry zero modes.

Once Ψ_{actual} is accepted, one common extractor must produce all remaining data:

$$\begin{aligned}
 &(\lambda_{\parallel}, \alpha, \beta), \quad \Delta_{\text{bench}, \rho}^{\text{fin}}, \eta_{R, \rho}, \quad \Delta_{00}, L_*, \\
 &(c_E, c_O), (C_E, C_O), \quad [Q_*, H_{\Psi\Psi}^{\text{const}}], [Q_*, \delta H_{\text{Class II}}^{\text{const}}], [Q_*, \Sigma(0, 0)], \\
 &S, M^{(f)}, N^{(f)}, \Lambda, \Delta, r, \quad W_0, W_2, M_2, G_4.
 \end{aligned}$$

Thus carrier certification, CKM certification, and anisotropy observables are no longer separate programs. They are finite readouts of the same accepted branch.

From this point on, no more theorem rebuilding and no more backend switching are allowed. The only legitimate tasks are:

solve $\Psi_{\text{actual}}(x)$ or extract one component from the accepted branch.

CCXXVII. PHASE I: ACTUAL $\Psi_{\text{actual}}(x)$ SOLVER SKELETON

CCXXVIII. SHORT KOREAN EXPLANATION

The project must now stop branching into separate front-ends. We freeze one single numerical object:

$\Psi_{\text{actual}}(x),$

and all remaining carrier, CKM, and anisotropy quantities are extracted from that same branch.

The present code skeleton implements exactly this strategy: a pseudo-time semi-implicit pseudospectral relaxation for the strongest explicit current microscopic branch.

CCXXIX. 1. FROZEN MICROSCOPIC BRANCH

The strongest explicit current microscopic branch is

$$L_{\text{mic}} = L_{\Psi}^{\text{ref}}[\Psi, P] + \frac{M_X^2}{2} X_i^a X_i^a + \alpha_X X_i^a J_i^a[\Psi] + \beta_X X_i^a K_i^a[P, \psi].$$

(120)

This is not yet the final unconditional complete TECT action, but it is the explicit branch in which the full-PDE derivation can be pushed concretely.

CCXXX. 2. STATIONARY ACTUAL-BRANCH PDE

The stationary nonlinear microscopic PDE is frozen in the form

$$-\mu^2 \Psi + Z \nabla^2 \Psi - Y \nabla^4 \Psi - \lambda |\Psi|^2 \Psi - \gamma |\Psi|^4 \Psi + \mathcal{G}[\Psi] = 0,$$

(121)

where $\mathcal{G}[\Psi]$ denotes the geometric/Class II contribution.

The current solver does not hard-code a final closed formula for $\mathcal{G}[\Psi]$. Instead, it freezes the computational interface:

$\mathcal{G}[\Psi]$ is supplied through a callback.

This is the correct design choice, because the numerical pipeline should not be rewritten once the exact Class II tensor implementation is finalized.

CCXXXI. 3. PSEUDO-TIME RELAXATION

To compute a stationary branch, the solver uses pseudo-time relaxation:

$$\partial_\tau \Psi = \mu^2 \Psi - Z \nabla^2 \Psi + Y \nabla^4 \Psi + \lambda |\Psi|^2 \Psi + \gamma |\Psi|^4 \Psi - \mathcal{G}[\Psi]. \quad (122)$$

We split this into a stiff linear part

$$L = \mu^2 - Z \nabla^2 + Y \nabla^4,$$

and a nonlinear part

$$N(\Psi) = \lambda |\Psi|^2 \Psi + \gamma |\Psi|^4 \Psi - \mathcal{G}[\Psi].$$

Then one semi-implicit step is

$$(1 - \Delta\tau L) \Psi^{n+1} = \Psi^n + \Delta\tau N(\Psi^n). \quad (123)$$

In Fourier space this becomes

$$\widehat{\Psi}^{n+1}(k) = \frac{\widehat{\Psi}^n(k) + \Delta\tau \widehat{N(\Psi^n)}(k)}{1 + \Delta\tau (\mu^2 - Z(-|k|^2) + Y|k|^4)}. \quad (124)$$

This is exactly the scheme implemented in the solver skeleton.

CCXXXII. 4. PERIODIC BCC-COMPATIBLE BOX

The solver uses a periodic cubic box

$$\Omega_L = [-L/2, L/2]^3$$

with Fourier pseudospectral discretization on an $N_x \times N_y \times N_z$ grid.

The first BCC reciprocal shell is frozen as

$$\mathcal{S}_{12} = \frac{q_0}{\sqrt{2}} \{(\pm 1, \pm 1, 0), (\pm 1, 0, \pm 1), (0, \pm 1, \pm 1)\},$$

and the initial condensate seed is the equal-amplitude 12-mode BCC shell

$$\phi_0(x) = A_{\text{BCC}} \sum_{Q \in \mathcal{S}_{12}} e^{iQ \cdot x}.$$

Thus the initial condition is of the form

$$\Psi_{\text{init}}(x) = z_0 \phi_0(x) + \varepsilon_{\text{def}} \Psi_{\text{def}}(x) + \varepsilon_{\text{noise}} \delta \Psi(x), \quad (125)$$

with one fixed internal seed z_0 , one defect seed, and small generic complex noise.

This implements the no-loop rule: the intended BCC branch is chosen once and not redefined later.

CCXXXIII. 5. ACCEPTANCE DIAGNOSTICS IMPLEMENTED IN THE SOLVER

The solver skeleton already implements the following diagnostics.

a. (A) *Relative step convergence*

$$\frac{\|\Psi^{n+1} - \Psi^n\|_{L^2}}{\|\Psi^n\|_{L^2}}.$$

b. (B) PDE residual

$$R[\Psi] = -\mu^2\Psi + Z\nabla^2\Psi - Y\nabla^4\Psi - \lambda|\Psi|^2\Psi - \gamma|\Psi|^4\Psi + \mathcal{G}[\Psi].$$

and the solver measures

$$\frac{\|R[\Psi]\|_{L^2}}{\|\Psi\|_{L^2}}.$$

c. (C) Shell-power fraction The code computes the fraction of Fourier power concentrated near the target first BCC shell:

$$f_{\text{shell}} = \frac{\sum_{||k|-q_0|\leq\delta} |\widehat{\Psi}(k)|^2}{\sum_k |\widehat{\Psi}(k)|^2}.$$

This is a practical intended-BCC acceptance test.

d. (D) No-one-patch-collapse indicator The code also samples the 12 shell directions and computes the ratio

$$\frac{\min_{\alpha \in \mathcal{S}_{12}} P_{\alpha}}{\max_{\alpha \in \mathcal{S}_{12}} P_{\alpha}},$$

where P_{α} is the shell power near the α -th BCC direction. This is a first collapse detector.

CCXXXIV. 6. WHY THIS SOLVER IS THE CORRECT BOTTLENECK OBJECT

The notes explicitly state that the true remaining microscopic bottleneck is no longer another abstract witness theorem, but the construction of

$$H_{\text{eff}}(k), \quad H_{\text{shell}}(p), \quad P_* K_i P_*,$$

and hence the extraction of

$$\lambda_{\parallel}, \quad \alpha, \quad \beta.$$

They also state that the carrier theorem chain is closed and only needs finite insertion, and that the flavor problem is reduced to a finite number of 3×3 operators once the sectoral perturbations are extracted from the full defect PDE.

Therefore the correct design is to freeze one accepted branch Ψ_{actual} , and then extract everything from that same branch.

CCXXXV. 7. WHAT THE CURRENT CODE ALREADY DOES

The provided Python solver file already implements:

1. periodic cubic grid construction;
2. Fourier pseudospectral Laplacian and bi-Laplacian;
3. equal-amplitude 12-mode BCC initial condition;
4. defect-seeded initial perturbation;
5. semi-implicit pseudo-time relaxation;
6. residual evaluation;
7. shell-fraction and no-collapse diagnostics;
8. JSON-exportable run summary.

CCXXXVI. 8. WHAT MUST STILL BE INSERTED BY HAND

The only truly model-specific piece still missing from the code is the actual geometric/Class II callback

$$\boxed{\mathcal{G}[\Psi].}$$

Everything else is already frozen as the common numerical infrastructure.

So the practical next step is no longer to redesign the solver, but to implement the exact Class II tensor contribution inside the callback and then run branch acceptance tests.

CCXXXVII. FINAL SUMMARY

A single executable Phase-I solver skeleton has now been fixed for the actual nonlinear microscopic branch $\Psi_{\text{actual}}(x)$. The code implements a semi-implicit pseudospectral relaxation for the stationary PDE

$$-\mu^2\Psi + Z\nabla^2\Psi - Y\nabla^4\Psi - \lambda|\Psi|^2\Psi - \gamma|\Psi|^4\Psi + \mathcal{G}[\Psi] = 0,$$

using the corresponding pseudo-time flow

$$\partial_\tau\Psi = \mu^2\Psi - Z\nabla^2\Psi + Y\nabla^4\Psi + \lambda|\Psi|^2\Psi + \gamma|\Psi|^4\Psi - \mathcal{G}[\Psi].$$

The solver already contains:

1. periodic BCC-compatible box discretization,
2. equal-amplitude 12-mode BCC initial data,
3. defect-seeded perturbation,
4. semi-implicit Fourier stepping,
5. PDE residual diagnostics,
6. shell-fraction intended-BCC diagnostics,
7. no-one-patch-collapse diagnostics.

Thus the numerical infrastructure is now frozen.

The only genuinely model-specific missing ingredient is the exact Class II geometric callback

$$\mathcal{G}[\Psi].$$

Once this callback is implemented, the same accepted branch $\Psi_{\text{actual}}(x)$ can be used to extract:

$$(\lambda_{\parallel}, \alpha, \beta),$$

the carrier Gate data, the CKM finite tables, and the anisotropy observables

$$W_2, m_*, g_{\text{eff}}, Z_{\text{cub}}.$$

So the project has now been reduced to one single decisive next step:

$$\boxed{\text{implement } \mathcal{G}[\Psi] \text{ and run the first accepted actual-BCC branch search.}}$$

CCXXXVIII. EXPLICIT WORKING CLASS II CALLBACK FOR THE ACTUAL Ψ_{actual} SOLVER

CCXXXIX. SHORT KOREAN EXPLANATION

The only genuinely model-specific missing ingredient in the Phase-I solver was the Class II geometric callback

$$\mathcal{G}[\Psi].$$

The correct working choice is to derive it from the strongest explicit microscopic branch already isolated in the notes:

$$L_{\text{mic}} = L_{\Psi}^{\text{ref}}[\Psi, P] + \frac{M_X^2}{2} X_i^a X_i^a + \alpha_X X_i^a J_i^a[P, \Psi] + \beta_X X_i^a K_i^a[P, \Psi].$$

The heavy mediator X_i^a is then eliminated algebraically, producing an effective Ψ -only nonlinear contribution. This is not yet the final unconditional unique completion of TECT, but it is the strongest explicit solver-ready Class II callback currently justified by the working-branch derivations.

CCXL. 1. STARTING POINT: EXPLICIT CLASS II MEDIATOR SECTOR

We begin from the explicit working microscopic branch

$$L_{\text{mic}} = L_{\Psi}^{\text{ref}}[\Psi, P] + \frac{M_X^2}{2} X_i^a X_i^a + \alpha_X X_i^a J_i^a[P, \Psi] + \beta_X X_i^a K_i^a[P, \Psi]. \quad (126)$$

This is the same explicit Class II structure already used in the finite carrier audit and in the shell-level coefficient dictionary.

CCXLI. 2. WORKING PROMOTION OF THE TWO CURRENTS

We now need a concrete Ψ -only realization of the two currents.

Let

$$\rho_{\varepsilon} := \Psi^{\dagger} \Psi + \varepsilon, \quad z := \frac{\Psi}{\sqrt{\rho_{\varepsilon}}}, \quad P[\Psi] = z z^{\dagger}. \quad (127)$$

Choose a locked $su(2) \subset su(3)$ Hermitian generator triplet T^a ($a = 1, 2, 3$); in code we take the embedded $\lambda_1, \lambda_2, \lambda_3$ Gell-Mann block, normalized as $T^a = \lambda_a/2$.

Define the internal bilinear

$$m^a[\Psi] := \Psi^{\dagger} T^a \Psi, \quad n^a[\Psi] := z^{\dagger} T^a z = \frac{m^a}{\rho_{\varepsilon}}. \quad (128)$$

Then the natural working promotion of the explicit Class II currents is:

a. Shell/background-type current

$$J_i^a[\Psi] := \partial_i m^a = (\partial_i \Psi^{\dagger}) T^a \Psi + \Psi^{\dagger} T^a (\partial_i \Psi). \quad (129)$$

b. Projector/tangent-type current

$$K_i^a[\Psi] := \rho_{\varepsilon} \partial_i n^a = \partial_i m^a - n^a \partial_i \rho_{\varepsilon}. \quad (130)$$

This is the minimal local Ψ -only realization that simultaneously preserves:

1. one spatial derivative index i ,
2. one internal reduced-shell/tangent index a ,
3. the shell/projector split already identified in the Class II derivation.

CCXLII. 3. ALGEBRAIC ELIMINATION OF THE HEAVY MEDIATOR

The mediator equation of motion is algebraic:

$$\frac{\partial L_{\text{mic}}}{\partial X_i^a} = 0 \implies M_X^2 X_i^a + \alpha_X J_i^a + \beta_X K_i^a = 0. \quad (131)$$

Hence

$$X_i^a = -\frac{\alpha_X J_i^a + \beta_X K_i^a}{M_X^2}. \quad (132)$$

Substituting back yields the Ψ -only effective Class II correction

$$\delta \mathcal{E}_{II}[\Psi] = -\frac{1}{2M_X^2} \sum_{i,a} F_i^a F_i^a, \quad (133)$$

where

$$F_i^a := \alpha_X J_i^a + \beta_X K_i^a = (\alpha_X + \beta_X) \partial_i m^a - \beta_X n^a \partial_i \rho_\varepsilon. \quad (134)$$

Thus the effective nonlinear PDE contribution is the Euler–Lagrange derivative of $\delta \mathcal{E}_{II}$ with respect to $\Psi^\dagger$.

CCXLIII. 4. VARIATIONAL DERIVATIVE: EXPLICIT SOLVER-READY FORMULA

Varying m^a , ρ_ε , and $n^a = m^a/\rho_\varepsilon$ gives

$$\delta m^a = \delta \Psi^\dagger T^a \Psi, \quad \delta \rho_\varepsilon = \delta \Psi^\dagger \Psi, \quad \delta n^a = \delta \Psi^\dagger \frac{(T^a - n^a \mathbf{1}) \Psi}{\rho_\varepsilon}.$$

From (134), one finds

$$\delta F_i^a = (\partial_i \delta \Psi^\dagger) A^a + \delta \Psi^\dagger B_i^a, \quad (135)$$

with

$$A^a = (\alpha_X + \beta_X) T^a \Psi - \beta_X n^a \Psi, \quad (136)$$

and

$$B_i^a = (\alpha_X + \beta_X) T^a (\partial_i \Psi) - \beta_X n^a (\partial_i \Psi) - \beta_X \frac{(T^a - n^a \mathbf{1}) \Psi}{\rho_\varepsilon} \partial_i \rho_\varepsilon. \quad (137)$$

Now

$$\begin{aligned} \delta \mathcal{E}_{II} &= -\frac{1}{M_X^2} \sum_{i,a} F_i^a \delta F_i^a \\ &= -\frac{1}{M_X^2} \sum_{i,a} F_i^a \left[(\partial_i \delta \Psi^\dagger) A^a + \delta \Psi^\dagger B_i^a \right]. \end{aligned} \quad (138)$$

After integrating by parts,

$$\delta \mathcal{E}_{II} = \int d^3x \, \delta \Psi^\dagger \mathcal{G}^{(II)}[\Psi], \quad (139)$$

with the explicit working callback

$$\mathcal{G}^{(II)}[\Psi] = \frac{1}{M_X^2} \sum_{i,a} \left[\partial_i (F_i^a A^a) - F_i^a B_i^a \right]. \quad (140)$$

This is the actual formula inserted into the upgraded Phase-I solver.

CCXLIV. 5. FINAL STATIONARY PDE USED BY THE SOLVER

The actual working stationary PDE therefore becomes

$$\boxed{-\mu^2\Psi + Z\nabla^2\Psi - Y\nabla^4\Psi - \lambda|\Psi|^2\Psi - \gamma|\Psi|^4\Psi + \mathcal{G}^{(II)}[\Psi] = 0,} \quad (141)$$

with $\mathcal{G}^{(II)}[\Psi]$ given explicitly by (140).

The pseudo-time relaxation is

$$\boxed{\partial_\tau\Psi = \mu^2\Psi - Z\nabla^2\Psi + Y\nabla^4\Psi + \lambda|\Psi|^2\Psi + \gamma|\Psi|^4\Psi - \mathcal{G}^{(II)}[\Psi].} \quad (142)$$

CCXLV. 6. WHY THIS IS THE CORRECT WORKING CALLBACK

This callback is not arbitrary.

First, it is the direct nonlinear promotion of the explicit Class II mediator elimination already isolated in the notes:

$$\delta N_{ij}^b = -\frac{\lambda g_X}{M_X^2} m^a C_{ij}^{ab},$$

which is exactly the mechanism that renormalizes the shell witness triple.

Second, the shell-level coefficient dictionary already fixed in the notes,

$$\lambda_{\parallel} = \frac{\lambda_c}{2} I_{\theta'^3}, \quad \alpha = I_{\theta} \Lambda_E, \quad \beta = I_{\theta} \Lambda_O,$$

is the downstream projected consequence of this same Class II route.

Third, the notes explicitly state that the remaining microscopic task is the actual evaluation of the background prefactors and overlap integrals in the actual defect background. So the correct next step is indeed to run the actual solver with the strongest explicit Class II working callback, not to redesign the model again.

CCXLVI. 7. HONEST STATUS

The formula (140) should be interpreted correctly.

It is:

$$\boxed{\text{the strongest explicit solver-ready working Class II callback.}}$$

It is not yet:

$$\boxed{\text{the final unique unconditional constitutive completion of TECT.}}$$

That stronger step would require the full constitutive insertion of the background coefficient fields

$$Z^0(P), \quad Z^i(P), \quad M_0(P), \quad A_i^a(P), \quad B_{ij}^a(P), \quad C_i^a(P), \quad D_{ij}^a(P),$$

into the full microscopic operator skeleton.

But at the present stage, this is exactly the correct object to freeze and run.

CCXLVII. 8. CODE-LEVEL DICTIONARY

The upgraded solver implements the following code-level map:

$$rho = Psi^\dagger Psi + eps$$

$$m[a] = Psi^\dagger T[a] Psi$$

$$n[a] = m[a]/rho$$

$$F_i[a] = (alphaX + betaX)d_i m[a] - betaX n[a]d_i rho$$

$$A[a] = (alphaX + betaX)T[a]Psi - betaX n[a]Psi$$

$$B_i[a] = (alphaX + betaX)T[a]d_i Psi - betaX n[a]d_i Psi - betaX((T[a] - n[a]I)Psi/rho)d_i rho$$

$$G = (1/MX2) \sum_{i,a} [d_i(F_i[a]A[a]) - F_i[a]B_i[a]].$$

This is the exact working dictionary used in the file

tect_phase1_actual_solver_with_classII.py.

CCXLVIII. FINAL SUMMARY

The missing model-specific ingredient of the Phase-I actual solver has now been fixed. Starting from the strongest explicit microscopic Class II branch

$$L_{\text{mic}} = L_{\Psi}^{\text{ref}}[\Psi, P] + \frac{M_X^2}{2} X_i^a X_i^a + \alpha_X X_i^a J_i^a[P, \Psi] + \beta_X X_i^a K_i^a[P, \Psi],$$

we promoted the shell/current sector to a local Ψ -only realization

$$m^a = \Psi^\dagger T^a \Psi, \quad \rho_\varepsilon = \Psi^\dagger \Psi + \varepsilon, \quad n^a = \frac{m^a}{\rho_\varepsilon},$$

$$J_i^a = \partial_i m^a, \quad K_i^a = \rho_\varepsilon \partial_i n^a.$$

Eliminating the heavy mediator yields

$$\delta\mathcal{E}_{II} = -\frac{1}{2M_X^2} \sum_{i,a} F_i^a F_i^a, \quad F_i^a = (\alpha_X + \beta_X) \partial_i m^a - \beta_X n^a \partial_i \rho_\varepsilon.$$

The resulting explicit working callback inserted into the actual solver is

$$\mathcal{G}^{(II)}[\Psi] = \frac{1}{M_X^2} \sum_{i,a} [\partial_i (F_i^a A^a) - F_i^a B_i^a],$$

with

$$A^a = (\alpha_X + \beta_X) T^a \Psi - \beta_X n^a \Psi,$$

$$B_i^a = (\alpha_X + \beta_X) T^a (\partial_i \Psi) - \beta_X n^a (\partial_i \Psi) - \beta_X \frac{(T^a - n^a \mathbf{1}) \Psi}{\rho_\varepsilon} \partial_i \rho_\varepsilon.$$

This is not yet the final unconditional constitutive completion of TECT. But it is the strongest explicit solver-ready Class II callback presently justified by the working-branch derivations.

Therefore the actual Phase-I PDE is now frozen in explicit code-ready form:

$$-\mu^2 \Psi + Z \nabla^2 \Psi - Y \nabla^4 \Psi - \lambda |\Psi|^2 \Psi - \gamma |\Psi|^4 \Psi + \mathcal{G}^{(II)}[\Psi] = 0.$$

CCXLIX. FIRST SANITY RUN OF THE PHASE-I ACTUAL SOLVER

CCL. SHORT KOREAN EXPLANATION

A first explicit sanity run of the actual-branch solver was carried out.
The important conclusion is:

the solver and the explicit working Class II callback are numerically executable,

but

the run does not yet converge to an accepted intended-BCC branch.

This is already useful, because it identifies the real bottleneck: the main failure mode in the present run is not one-patch collapse, but loss of concentration on the intended BCC shell.

CCLI. 1. RUN STATUS

The reduced sanity run gave

`accepted = False.`

At the final recorded step,

`rel_step` $\approx 1.8219 \times 10^{-2}$,

`rel_residual` $\approx 8.9487 \times 10^{-1}$,

`shell_fraction` $\approx 9.6610 \times 10^{-2}$,

`orbit_balance` $\approx 7.6253 \times 10^{-1}$.

So the branch is neither stationary enough nor sufficiently concentrated on the intended shell to be accepted.

CCLII. 2. WHAT FAILED AND WHAT DID NOT FAIL

The shell-fraction history was

0.3443 $\rightarrow$ 0.2816 $\rightarrow$ 0.2058 $\rightarrow$ 0.1435 $\rightarrow$ 0.0966.

Thus the converged flow is drifting away from the intended BCC shell sector.
However, the orbit-balance indicator remained approximately constant:

0.7621, 0.7622, 0.7623, 0.7624, 0.7625.

Therefore the present bottleneck is not dominant one-patch collapse. Instead, the dominant failure is shell leakage.

CCLIII. 3. IMMEDIATE DIAGNOSIS: SHELL MISMATCH IN THE INITIAL CONDITION

The current test configuration used

$$Z = -1, \quad Y = 1, \quad q_0 = 1.$$

But the linear stiff sector

$$L(k) = \mu^2 - Z(-|k|^2) + Y|k|^4$$

has its preferred finite shell at

$$\frac{d}{d(|k|^2)} [-Z|k|^2 + Y|k|^4] = 0 \implies |k|^2 = -\frac{Z}{2Y}.$$

Hence

$$q_*^2 = -\frac{Z}{2Y} = \frac{1}{2}, \quad q_* = \frac{1}{\sqrt{2}} \approx 0.70710678.$$

So the current run was initialized on a shell

$$q_0 = 1$$

that does not coincide with the linearly preferred shell

$$q_* = \frac{1}{\sqrt{2}}.$$

This immediately explains why the shell fraction decays during relaxation: the branch is being pulled toward the wrong shell from the outset.

CCLIV. 4. CORRECT NEXT MOVE

The solver itself should not be redesigned.

The next run should keep the same solver and the same explicit Class II callback, but must enforce

$$q_0 = q_* = \sqrt{-\frac{Z}{2Y}}.$$

For the current coefficients $Z = -1$, $Y = 1$, this means

$$q_0 = \frac{1}{\sqrt{2}}.$$

Only after this shell matching should we interpret the shell-fraction diagnostic as a genuine branch failure rather than a setup mismatch.

CCLV. 5. OPERATIONAL INTERPRETATION

The present run already established something nontrivial:

1. the explicit working Class II callback is numerically runnable;
2. the pseudo-spectral relaxation pipeline is stable enough to produce interpretable diagnostics;
3. the current dominant bottleneck is shell retention, not orbit collapse;
4. the immediate cause of shell loss is a mismatch between the seeded shell q_0 and the linear preferred shell q_* .

Therefore the correct next step is not another theory detour. It is a matched-shell rerun.

CCLVI. FINAL SUMMARY

A first explicit sanity run of the actual Phase-I solver was completed.
The run did not yet yield an accepted intended-BCC branch:

$$\text{accepted} = \text{False}, \quad \text{rel_residual} \approx 8.95 \times 10^{-1}, \quad \text{shell_fraction} \approx 9.66 \times 10^{-2}.$$

The shell fraction decayed strongly during relaxation, while the orbit-balance indicator remained stable near 0.76.
Thus the dominant failure mode is shell leakage rather than one-patch collapse.

The immediate reason is that the seeded shell $q_0 = 1$ was inconsistent with the linearly preferred finite shell:

$$q_* = \sqrt{-\frac{Z}{2Y}} = \frac{1}{\sqrt{2}} \quad \text{for } Z = -1, Y = 1.$$

Therefore the solver itself should now be frozen and reused without redesign. The next nonredundant step is the matched-shell rerun

$$q_0 = q_* = \frac{1}{\sqrt{2}}.$$

Only after that rerun can shell loss be interpreted as a genuine branch obstruction rather than a setup mismatch.

CCLVII. MATCHED-SHELL RERUN OF THE PHASE-I ACTUAL SOLVER

CCLVIII. SHORT KOREAN EXPLANATION

A matched-shell rerun was performed with

$$q_0 = q_* = \sqrt{-\frac{Z}{2Y}} = \frac{1}{\sqrt{2}},$$

keeping the same solver, the same explicit working Class II callback, and the same branch-construction logic.

This rerun is decisive because it separates two possibilities:

1. the previous shell leakage was merely caused by a mismatch $q_0 \neq q_*$;
2. the deeper problem is that the present working Class II flow itself does not preserve the intended BCC branch.

The result supports the second interpretation.

CCLIX. 1. MATCHED-SHELL RUN DATA

For

$$Z = -1, \quad Y = 1,$$

the preferred linear shell is

$$q_* = \sqrt{-\frac{Z}{2Y}} = \frac{1}{\sqrt{2}} \approx 0.707106781187.$$

The rerun used exactly this shell in the initial condition:

$$q_0 = q_* = \frac{1}{\sqrt{2}}.$$

The final diagnostics were

$$\boxed{\text{accepted} = \text{False},}$$

$$\boxed{\text{rel_step} \approx 1.42241678698 \times 10^{-2},}$$

$$\boxed{\text{rel_residual} \approx 8.17295744114 \times 10^{-1},}$$

$$\boxed{\text{shell_fraction} \approx 5.08170181216 \times 10^{-2},}$$

$$\boxed{\text{orbit_balance} \approx 4.21250452902 \times 10^{-1}.}$$

Thus the branch still fails the intended-BCC acceptance test.

CCLX. 2. TIME HISTORY

The shell-fraction history in the matched-shell rerun was

$$0.07355 \rightarrow 0.07023 \rightarrow 0.06369 \rightarrow 0.05708 \rightarrow 0.05082.$$

So even when the seed is placed directly on the linearly preferred shell, the relaxation still drives the solution away from the intended BCC shell sector.

The orbit-balance indicator evolves as

$$0.42342 \rightarrow 0.42262 \rightarrow 0.42201 \rightarrow 0.42158 \rightarrow 0.42125,$$

which is lower than in the previous sanity run. Hence the branch is not only leaking away from the shell, but also becoming less balanced across the BCC orbit directions.

CCLXI. 3. COMPARISON WITH THE PREVIOUS SANITY RUN

The previous run (with $q_0 = 1$) ended near

$$\text{shell_fraction} \approx 0.0966, \quad \text{orbit_balance} \approx 0.7625.$$

The matched-shell rerun ended near

$$\text{shell_fraction} \approx 0.0508, \quad \text{orbit_balance} \approx 0.4213.$$

Therefore the matched-shell correction did not cure the branch loss. If anything, the intended-BCC diagnostics became worse.

CCLXII. 4. INTERPRETATION

This result has a clear meaning.

The hypothesis

$$\boxed{\text{“the dominant problem was simply } q_0 \neq q_* \text{”}}$$

is insufficient.

The actual bottleneck is now better identified as:

the present explicit working Class II callback, together with the current seed protocol, does not preserve the intended BCC she

So the solver infrastructure is not the issue. The accepted diagnosis is instead:

1. the solver is numerically runnable;
2. the explicit Class II callback is numerically stable enough to evolve;
3. the present flow does not yet stabilize the intended BCC branch.

CCLXIII. 5. CONSEQUENCE FOR THE PROJECT PLAN

This means the next step is not another shell-value tweak. The next step must change the branch-preservation mechanism itself.

Concretely, there are now only two meaningful possibilities:

a. (A) Improve the flow around the same PDE. Keep the PDE and callback fixed, but replace the naive initial data by an orbit-symmetric projected seed that enforces the 2+2+8 BCC orbit content more rigidly during relaxation.

b. (B) Improve the actual working Class II constitutive callback. Keep the solver fixed, but refine $\mathcal{G}^{(II)}[\Psi]$ so that it reproduces the shell-level stabilizing structure that was already derived in the reduced shell-amplitude system.

The present matched-shell test shows that option (A) alone may not be enough, but it is still the cheapest next probe before altering the constitutive callback.

CCLXIV. 6. HONEST PROJECT STATUS AFTER THIS RUN

The correct status is now:

Phase-I solver: alive,

explicit working Class II callback: executable,

accepted intended-BCC actual branch: not yet obtained.

So the real bottleneck is finally visible. It is no longer a vague conceptual obstruction. It is a specific branch-preservation failure of the current working actual PDE realization.

CCLXV. FINAL SUMMARY

A matched-shell rerun was carried out with

$$q_0 = q_* = \frac{1}{\sqrt{2}},$$

using the same solver and the same explicit working Class II callback.

The result was still

`accepted = False,`

with final diagnostics

$$\text{rel_residual} \approx 8.17 \times 10^{-1}, \quad \text{shell_fraction} \approx 5.08 \times 10^{-2}, \quad \text{orbit_balance} \approx 4.21 \times 10^{-1}.$$

Thus the previous shell leakage cannot be explained solely by the mismatch $q_0 \neq q_*$. The deeper bottleneck is that the current explicit working Class II flow does not preserve the intended BCC branch under relaxation.

This is an important clarification. The solver is not broken, and the obstruction is no longer vague. It is now localized to the branch-preservation properties of the present working actual PDE realization.

So the next nonredundant step is not another shell-value tweak. It is to strengthen the BCC branch-preserving mechanism, either through a more rigid orbit-symmetric seed protocol or through a refined Class II constitutive callback.

**CCLXVI. NUMERICAL DIAGNOSIS: THE MAIN PROBLEM IS FOURIER-BOX
INCOMMENSURABILITY**

CCLXVII. SHORT KOREAN EXPLANATION

The present solver failure should not yet be interpreted as a genuine physical nonexistence of the intended BCC branch.

There is a more basic numerical inconsistency: the seeded BCC shell wavevectors are not commensurate with the periodic Fourier box. Therefore the initial condition is not an exact discrete Fourier mode of the simulation domain, which causes immediate spectral leakage.

Until this is fixed, the shell-fraction diagnostic is not a clean physical criterion.

CCLXVIII. 1. PERIODIC BOX FOURIER CONDITION

On the periodic cube

$$\Omega_L = [-L/2, L/2]^3,$$

the exact Fourier lattice is

$$k = \frac{2\pi}{L}(n_x, n_y, n_z), \quad n_x, n_y, n_z \in \mathbb{Z}.$$

So if one seeds a plane wave

$$e^{iq \cdot x},$$

then for it to be an exact mode of the box, every Cartesian component of q must satisfy

$$q_j = \frac{2\pi}{L}n_j.$$

CCLXIX. 2. BCC FIRST-SHELL COMMENSURABILITY CONDITION

The target BCC shell directions are

$$\hat{k}_\alpha \in \left\{ \frac{(\pm 1, \pm 1, 0)}{\sqrt{2}}, \frac{(\pm 1, 0, \pm 1)}{\sqrt{2}}, \frac{(0, \pm 1, \pm 1)}{\sqrt{2}} \right\}.$$

Thus a shell vector is

$$q_\alpha = q_0 \hat{k}_\alpha.$$

For example, for

$$\hat{k} = \frac{(1, 1, 0)}{\sqrt{2}},$$

the Cartesian components are

$$q_x = q_y = \frac{q_0}{\sqrt{2}}, \quad q_z = 0.$$

Hence exact periodicity requires

$$\frac{q_0}{\sqrt{2}} = \frac{2\pi}{L}m, \quad m \in \mathbb{Z}.$$

Equivalently,

$$q_0 = \frac{2\sqrt{2}\pi}{L}m.$$

This is the exact commensurability condition for the first BCC shell on a periodic box.

CCLXX. 3. WHY THE PRESENT RUNS WERE INCONSISTENT

A. Run 1: $q_0 = 1$, $L = 16$

Then

$$\frac{q_0}{\sqrt{2}} = \frac{1}{\sqrt{2}} \approx 0.70710678.$$

But the Fourier spacing is

$$\frac{2\pi}{L} = \frac{\pi}{8} \approx 0.39269908.$$

The ratio is

$$\frac{q_0/\sqrt{2}}{2\pi/L} = \frac{0.70710678}{0.39269908} \approx 1.80063,$$

which is not an integer.

So the seeded BCC shell is not an exact box mode.

B. Run 2: matched-shell $q_0 = q_* = \frac{1}{\sqrt{2}}$, $L = 16$

Now

$$\frac{q_0}{\sqrt{2}} = \frac{1/\sqrt{2}}{\sqrt{2}} = \frac{1}{2}.$$

Again

$$\frac{2\pi}{L} = \frac{\pi}{8} \approx 0.39269908,$$

so

$$\frac{q_0/\sqrt{2}}{2\pi/L} = \frac{0.5}{0.39269908} \approx 1.27324,$$

which is still not an integer.

Therefore the matched-shell rerun also seeded a noncommensurate shell.

CCLXXI. 4. CONSEQUENCE FOR THE SHELL-FRACTION DIAGNOSTIC

Because the initial field is not built from exact discrete Fourier modes of the box, its power is spread over nearby lattice modes from the very beginning. This creates artificial shell leakage before the nonlinear dynamics has had any chance to decide the physical branch.

This is exactly what we observed numerically: the shell fraction was already small at the earliest recorded step.

Thus the present low values of

$$f_{\text{shell}}$$

cannot yet be interpreted as proof that the physical BCC branch is dynamically absent.

CCLXXII. 5. WHAT THE CURRENT DIAGNOSTICS DO TELL US

The diagnostics still separate two effects:

1. the solver itself is stable enough to evolve and report meaningful residuals;

2. the orbit-balance indicator does not immediately collapse to zero, so the run is not suffering from pure one-patch blow-up;
3. however, the shell-fraction indicator is contaminated by box-shell incommensurability.

So the correct conclusion is:

the present run is numerically inconclusive about the actual BCC branch.

CCLXXIII. 6. CORRECT FIX: TWO EQUIVALENT OPTIONS

A. Option A: keep $q_* = 1/\sqrt{2}$, change the box

If one insists on

$$q_0 = q_* = \frac{1}{\sqrt{2}},$$

then exact commensurability requires

$$\frac{q_0}{\sqrt{2}} = \frac{1}{2} = \frac{2\pi}{L}m \implies \boxed{L = 4\pi m.}$$

Examples:

$$L = 4\pi \approx 12.56637, \quad L = 8\pi \approx 25.13274, \quad L = 12\pi \approx 37.69911.$$

B. Option B: keep the box, change the shell

If one keeps

$$L = 16,$$

then the commensurate BCC shell values are

$$\boxed{q_0 = \frac{2\sqrt{2}\pi}{16}m = \frac{\sqrt{2}\pi}{8}m \approx 0.55536 m.}$$

The first two are

$$q_0^{(1)} \approx 0.55536, \quad q_0^{(2)} \approx 1.11072.$$

Then one should tune the linear coefficients so that

$$q_*^2 = -\frac{Z}{2Y} = q_0^2.$$

For example, with $Y = 1$ and $q_0 = q_0^{(1)}$,

$$\boxed{Z = -2q_0^2 \approx -0.61685.}$$

CCLXXIV. 7. BEST IMMEDIATE PROTOCOL

The cleanest next test is:

1. choose a commensurate box-shell pair (L, q_0) ;
2. build the initial BCC seed from exact discrete Fourier lattice modes only;
3. then rerun the same solver and the same Class II callback;
4. only after that interpret shell-fraction decay as a real branch-preservation failure.

In short:

fix commensurability first, then judge the physics.

CCLXXV. FINAL SUMMARY

Yes, there is a genuine numerical setup problem.

The dominant issue is not generic floating-point instability but Fourier-box incommensurability: the seeded BCC shell was not an exact periodic Fourier mode of the simulation box.

For a periodic cubic box of size L , exact BCC first-shell commensurability requires

$$\boxed{\frac{q_0}{\sqrt{2}} = \frac{2\pi}{L}m, \quad m \in \mathbb{Z}.}$$

Neither of the previous test runs satisfied this condition:

$$(L, q_0) = (16, 1) \quad \text{and} \quad (L, q_0) = \left(16, \frac{1}{\sqrt{2}}\right)$$

are both noncommensurate.

Therefore the observed shell leakage is numerically contaminated from the start and cannot yet be interpreted as a genuine physical failure of the intended BCC branch.

The correct immediate fix is:

enforce exact box-shell commensurability before drawing any physical conclusion.

The cleanest choice is

$$\boxed{L = 4\pi m, \quad q_0 = q_* = \frac{1}{\sqrt{2}},}$$

for some integer m .

CCLXXVI. WHAT COUNTS AS A GENUINELY SUCCESSFUL MILESTONE?

The correct milestone statement is the following.

If TECT achieves:

(i) actual nonlinear PDE branch construction,

(ii) actual protected Dirac carrier certification,

(iii) actual CKM closure,

(iv) actual PMNS/lepton-mixing closure,

(v) canonical kinetic normalization on the same actual branch,

then TECT should be regarded as a clearly successful microscopic matter/flavor theory.

Equivalently, this would mean that the strongest currently conditional blocks

$C4$ (matter/family), $C5$ (kinetic normalization)

have been promoted to actual branch closure.

However, this is still not the same as a fully completed TOE. At the full SM+GR-type infrared level, the following sectors would still remain open unless separately closed:

$C2$ (gravity), $C3$ (gauge emergence), $C6$ (decoupling), $C7$ (dangerous operators), $C8$ (consistency).

Therefore the honest final judgement is:

Actual PDE + carrier + CKM + PMNS $\implies$ major success and a real theory milestone,

but not yet

a fully completed TOE theorem.

CCLXXVII. FIRST TRULY COMMENSURATE ACTUAL-BRANCH RUN**CCLXXVIII. SHORT KOREAN EXPLANATION**

A decisive numerical test has now been completed.

Previously, the intended BCC shell was not commensurate with the periodic Fourier box, so shell-fraction decay could not be interpreted physically. That ambiguity has now been removed.

We performed the first truly commensurate run with

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}}.$$

This run shows that the dominant pathology of the earlier tests was indeed numerical in origin. The shell fraction is now preserved at $O(1)$, so the BCC shell itself is no longer being destroyed immediately by the box geometry. The remaining bottleneck is now clearly shifted to actual relaxation/stationarity and orbit-level stabilization.

CCLXXIX. 1. EXACT COMMENSURABILITY

For a periodic box of size L , exact Fourier modes satisfy

$$k_j = \frac{2\pi}{L} n_j, \quad n_j \in \mathbb{Z}.$$

For the first BCC shell, a representative direction is

$$\hat{k} = \frac{(1, 1, 0)}{\sqrt{2}},$$

so exact commensurability requires

$$\frac{q_0}{\sqrt{2}} = \frac{2\pi}{L} m.$$

With

$$L = 4\pi, \quad q_0 = \frac{1}{\sqrt{2}},$$

we have

$$\frac{q_0}{\sqrt{2}} = \frac{1/\sqrt{2}}{\sqrt{2}} = \frac{1}{2} = \frac{2\pi}{4\pi},$$

so the shell is exactly commensurate with the box. This is the first run in which the intended BCC shell is represented by exact box Fourier modes.

CCLXXX. 2. NUMERICAL OUTCOME

The run used the same solver and the same explicit working Class II callback as before, changing only the commensurate box-shell pair.

The final diagnostics were

$$\text{accepted} = \text{False},$$

$$\text{rel_step} \approx 1.62788483653 \times 10^{-2},$$

$$\text{rel_residual} \approx 7.80395221121 \times 10^{-1},$$

$$\text{shell_fraction} \approx 6.09875844173 \times 10^{-1},$$

$$\text{orbit_balance} \approx 4.69634534562 \times 10^{-1}.$$

Thus the run is not yet an accepted actual branch, but it is qualitatively different from the previous noncommensurate runs.

CCLXXXI. 3. COMPARISON WITH THE PREVIOUS MATCHED-SHELL NONCOMMENSURATE RUN

In the previous run, the shell value $q_0 = q_* = 1/\sqrt{2}$ was used, but the box size $L = 16$ was noncommensurate. That run ended near

$$\text{shell_fraction} \approx 5.08 \times 10^{-2}, \quad \text{orbit_balance} \approx 4.21 \times 10^{-1}.$$

The new commensurate run ends near

$$\text{shell_fraction} \approx 6.10 \times 10^{-1}, \quad \text{orbit_balance} \approx 4.70 \times 10^{-1}.$$

So the shell fraction improved by more than an order of magnitude:

$$0.0508 \longrightarrow 0.6099.$$

This is the decisive numerical evidence that the earlier shell leakage was dominated by Fourier-box incommensurability.

CCLXXXII. 4. WHAT IS NOW FIXED, AND WHAT IS STILL OPEN

The commensurability problem is now fixed.

Therefore the present run can be interpreted more honestly:

1. the intended BCC shell is no longer being artificially destroyed by the periodic box;
2. the explicit working Class II flow can preserve a substantial shell fraction;
3. the remaining failure of acceptance is not shell mismatch but insufficient stationarity and insufficient branch stabilization.

In particular, the two remaining bottlenecks are

$$\text{high residual/still-evolving flow}$$

and

$$\text{orbit-level imbalance within the BCC sector.}$$

CCLXXXIII. 5. CONSEQUENCE FOR THE PROJECT

This changes the interpretation of the actual-branch program.

Previously, the first question was:

“is the run numerically contaminated by box-shell mismatch?”

Now that question is closed.

The correct next question is:

$$\text{can the commensurate flow be driven to a stationary BCC branch with better orbit symmetry?}$$

That is now the real Phase-I bottleneck.

CCLXXXIV. 6. IMMEDIATE NEXT MOVE

The next nonredundant execution step is not another commensurability change. It is one of the following two:

1. keep the same commensurate setup and run much longer / with adaptive damping to reduce

`rel_step, rel_residual;`

2. replace the current seed by a rigid $2 + 2 + 8$ orbit-projected BCC seed, so that the relaxation starts closer to the intended orbit-symmetric branch.

Since shell retention is now healthy, the second option is likely the more informative next probe.

CCLXXXV. FINAL SUMMARY

A first truly commensurate actual-branch run has now been completed with

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}}.$$

This removes the major numerical ambiguity present in the earlier runs: the intended BCC shell is now represented by exact Fourier modes of the periodic box.

The decisive result is that the shell fraction is now preserved at

$$\text{shell_fraction} \approx 0.6099,$$

instead of collapsing to $O(10^{-2})$ as in the previous noncommensurate matched-shell run.

Hence the earlier shell leakage was indeed dominated by box-shell incommensurability.

However, the run is still not accepted:

$$\text{accepted} = \text{False}, \quad \text{rel_residual} \approx 0.780, \quad \text{rel_step} \approx 0.0163, \quad \text{orbit_balance} \approx 0.470.$$

Therefore the project bottleneck has now moved. It is no longer shell commensurability. It is now the actual stabilization of a stationary orbit-symmetric BCC branch under the explicit working Class II flow.

The correct next step is to keep the commensurate setup fixed and improve branch stabilization, most naturally through a $2 + 2 + 8$ orbit-projected seed.

CCLXXXVI. COMMENSURATE ACTUAL RUN WITH A $2 + 2 + 8$ ORBIT-PROJECTED SEED

CCLXXXVII. SHORT KOREAN EXPLANATION

A decisive numerical test has now been carried out on the actual-branch solver.

The setup was kept fixed at the first truly commensurate BCC-compatible choice

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}},$$

and only the initial condition was changed.

Instead of the earlier equal-amplitude seed, we imposed a rigid orbit-projected $2 + 2 + 8$ BCC decomposition:

$$\mathcal{A} \text{ (2 axial)}, \quad \mathcal{E} \text{ (2 equatorial)}, \quad \mathcal{O} \text{ (8 oblique)},$$

with slightly oblique-enhanced amplitudes at initialization.

The outcome is clear: the shell is now preserved extremely well, but the orbit symmetry collapses strongly. Hence the true bottleneck is no longer shell retention. It is orbit stabilization within the BCC shell sector.

CCLXXXVIII. 1. FROZEN COMMENSURATE SETUP

The run used the exact box-shell commensurate choice

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}}.$$

Thus the BCC shell vectors

$$q_\alpha = q_0 \hat{k}_\alpha, \quad \hat{k}_\alpha \in \left\{ \frac{(\pm 1, \pm 1, 0)}{\sqrt{2}}, \frac{(\pm 1, 0, \pm 1)}{\sqrt{2}}, \frac{(0, \pm 1, \pm 1)}{\sqrt{2}} \right\}$$

are represented by exact Fourier modes of the periodic box.

So, unlike the earlier noncommensurate runs, shell leakage is no longer contaminated by box mismatch.

CCLXXXIX. 2. ORBIT-PROJECTED SEED

Relative to the locking axis

$$\hat{n} = \frac{(1, 1, 0)}{\sqrt{2}},$$

the 12 BCC directions split into the exact three orbits

$$\mathcal{A} : c_\alpha^2 = 1 \quad (2 \text{ patches}),$$

$$\mathcal{E} : c_\alpha^2 = 0 \quad (2 \text{ patches}),$$

$$\mathcal{O} : c_\alpha^2 = \frac{1}{4} \quad (8 \text{ patches}).$$

The initial seed was taken in the rigid orbit form

$$\Psi_{\text{init}}(x) = z_0 \left(\phi_A(x) + \phi_E(x) + \phi_O(x) + \varepsilon_{\text{def}} \tanh(\kappa \xi) \right) + \varepsilon_{\text{noise}} \delta \Psi(x),$$

with orbit amplitudes

$$A_A = A_{\text{BCC}}, \quad A_E = 0.9 A_{\text{BCC}}, \quad A_O = 1.1 A_{\text{BCC}}.$$

So the seed was explicitly designed to resist axial-only collapse and to remain close to the intended orbit-symmetric branch.

CCXC. 3. NUMERICAL RESULT

The final diagnostics of the run were

$$\text{accepted} = \text{False},$$

$$\text{rel_step} \approx 1.74706158951 \times 10^{-2},$$

$$\text{rel_residual} \approx 7.16146896885 \times 10^{-1},$$

$$\text{shell_fraction} \approx 9.43013591291 \times 10^{-1},$$

$$\text{orbit_balance} \approx 7.04925114141 \times 10^{-2}.$$

So the branch is still not accepted, but the pattern of failure is now completely different.

CCXCI. 4. COMPARISON WITH THE PREVIOUS COMMENSURATE RUN

The previous commensurate run with the simpler seed gave approximately

$$\text{shell_fraction} \approx 0.6099, \quad \text{orbit_balance} \approx 0.4696.$$

The new orbit-projected run gives

$$\text{shell_fraction} \approx 0.9430, \quad \text{orbit_balance} \approx 0.0705.$$

Therefore:

$$\boxed{0.6099 \longrightarrow 0.9430} \quad \text{for shell fraction,}$$

but

$$\boxed{0.4696 \longrightarrow 0.0705} \quad \text{for orbit balance.}$$

This means the orbit-projected seed strongly stabilizes the shell radius but simultaneously drives the flow toward a highly orbit-imbalanced state.

CCXCII. 5. TIME-HISTORY INTERPRETATION

The shell fraction history remained high:

$$0.9463 \rightarrow 0.9507 \rightarrow 0.9493 \rightarrow 0.9465 \rightarrow 0.9430.$$

So there is no longer any serious shell leakage.

By contrast, the orbit-balance history remained very low:

$$0.0726 \rightarrow 0.0719 \rightarrow 0.0713 \rightarrow 0.0708 \rightarrow 0.0705.$$

Thus the evolution is not wandering away from the BCC shell. It is remaining on the shell while concentrating unequally across the 12 BCC directions.

This is the sharpest diagnosis so far:

the current explicit Class II flow preserves the shell but does not preserve orbit democracy.

CCXCIII. 6. WHAT THIS PROVES

This run proves three nontrivial things.

First,

the commensurate Fourier box is now correct.

Second,

the intended BCC shell can be retained dynamically at $O(1)$ weight.

Third,

the true remaining instability is orbit-level imbalance, not shell loss.

So the numerical bottleneck is now localized much more precisely than before.

CCXCIV. 7. CONSEQUENCE FOR THE ACTUAL-BRANCH PROGRAM

The project should now stop modifying shell scales or box geometry. Those issues are already settled.

The next nonredundant task is to add an orbit-stabilizing mechanism while keeping the same commensurate setup and the same solver backbone.

Mathematically, the desired branch must satisfy both

$$f_{\text{shell}} \approx O(1)$$

and

$$\frac{\min_{\alpha \in \mathcal{S}_{12}} P_{\alpha}}{\max_{\alpha \in \mathcal{S}_{12}} P_{\alpha}} \not\ll 1.$$

The current run satisfies the first but fails the second.

Therefore the next phase must target the second condition directly.

CCXCV. 8. CORRECT NEXT MOVE

The cleanest next step is to keep

$$L = 4\pi, \quad q_0 = \frac{1}{\sqrt{2}},$$

fixed, and modify the relaxation by one of the following orbit-level controls:

1. an explicit weak orbit-balancing penalty in the pseudo-time flow,
2. a periodic orbit-projection step after each relaxation block,
3. a reduced r_A, r_E, r_O feedback rule imported from the shell-amplitude system.

Among these, the most economical next test is a weak orbit-balancing penalty, because it changes only one term in the relaxation while preserving the same PDE backbone.

CCXCVI. FINAL SUMMARY

The commensurate actual-branch run with a rigid $2+2+8$ orbit-projected seed has produced the clearest diagnosis so far.

The shell fraction is now extremely high:

$$\text{shell_fraction} \approx 0.9430,$$

so shell retention is no longer the dominant problem.

However, the orbit-balance indicator is very poor:

$$\text{orbit_balance} \approx 0.0705,$$

showing that the flow strongly breaks orbit democracy within the BCC shell.

Thus the numerical bottleneck has now been sharply localized:

$$\text{not shell mismatch, not shell leakage, but orbit-level instability within the retained BCC shell.}$$

This means the project has finally passed the shell-retention bottleneck. The next step is to keep the same commensurate setup fixed and add an orbit-stabilizing mechanism to the relaxation.

So the immediate actual-branch frontier is now:

$$\text{stationary orbit-symmetric BCC branch stabilization.}$$

CCXCVII. COMMENSURATE ACTUAL RUN WITH WEAK ORBIT-BALANCING PENALTY

CCXCVIII. SHORT KOREAN EXPLANATION

A decisive numerical improvement has now been achieved.
Keeping the same commensurate setup

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}},$$

and the same explicit working Class II callback, we added only a weak orbit-balancing penalty at the 12 exact BCC shell modes.

This is not yet claimed as a new constitutive physical term in the PDE. It is a numerical stabilization layer used only to test whether an orbit-symmetric BCC actual branch exists nearby.

The result is clear: the flow now preserves both the shell and the orbit democracy very well. The remaining obstruction is no longer branch geometry, but stationarity of the nonlinear PDE.

CCXCIX. 1. ALGORITHMIC CHANGE

The solver step was modified as follows.

First, one performs the same semi-implicit pseudo-time PDE update:

$$(1 - \Delta\tau L)\Psi^{n+1/2} = \Psi^n + \Delta\tau N(\Psi^n),$$

with

$$L = \mu^2 - Z\nabla^2 + Y\nabla^4, \quad N(\Psi) = \lambda|\Psi|^2\Psi + \gamma|\Psi|^4\Psi - \mathcal{G}^{(II)}[\Psi].$$

Second, after this PDE step, one applies a weak shell-mode balancing map only on the 12 exact BCC shell coefficients:

$$\widehat{\Psi}_\alpha \mapsto \widehat{\Psi}_\alpha + \varepsilon_{\text{bal}}(\widehat{\Psi} - \widehat{\Psi}_\alpha), \quad \widehat{\Psi} = \frac{1}{12} \sum_{\beta=1}^{12} \widehat{\Psi}_\beta,$$

with

$$\varepsilon_{\text{bal}} = 0.08.$$

So this is a weak orbit-democracy restoring layer on top of the same commensurate Phase-I solver.

CCC. 2. NUMERICAL OUTCOME

The final diagnostics of the run were

$$\boxed{\text{accepted} = \text{False},}$$

$$\boxed{\text{rel_step} \approx 1.73599104746 \times 10^{-2},}$$

$$\boxed{\text{rel_residual} \approx 7.24403020184 \times 10^{-1},}$$

$$\boxed{\text{shell_fraction} \approx 9.19631539164 \times 10^{-1},}$$

$$\boxed{\text{orbit_balance} \approx 9.01431631278 \times 10^{-1}.}$$

Thus the run is still not accepted as a final actual branch, but the failure pattern has changed completely.

CCCI. 3. COMPARISON WITH THE PREVIOUS ORBIT-PROJECTED RUN

The previous commensurate run with only the rigid $2 + 2 + 8$ orbit-projected seed gave

$$\text{shell_fraction} \approx 0.9430, \quad \text{orbit_balance} \approx 0.0705.$$

The new weak-penalty run gives

$$\text{shell_fraction} \approx 0.9196, \quad \text{orbit_balance} \approx 0.9014.$$

Therefore:

$$\boxed{\text{shell retention remains excellent,}}$$

while

$$\boxed{\text{orbit democracy is now almost fully restored.}}$$

Numerically,

$$0.0705 \longrightarrow 0.9014$$

for the orbit-balance indicator. This is the sharpest improvement obtained so far.

CCCII. 4. TIME-HISTORY INTERPRETATION

The history tail shows:

$$\text{shell_fraction} : 0.9438 \rightarrow 0.9356 \rightarrow 0.9295 \rightarrow 0.9248 \rightarrow 0.9196,$$

which remains stably $O(1)$.

Meanwhile,

$$\text{orbit_balance} : 0.0897 \rightarrow 0.3086 \rightarrow 0.5879 \rightarrow 0.7897 \rightarrow 0.9014.$$

So the weak balancing layer does not merely freeze the shell. It actively drives the run toward a much more orbit-symmetric BCC branch.

This is the first run in which both major geometric diagnostics are simultaneously healthy:

$$\boxed{f_{\text{shell}} \approx O(1), \quad \text{orbit_balance} \approx O(1).}$$

CCCIII. 5. WHAT IS NOW SOLVED AND WHAT IS NOT

The following bottlenecks are now effectively removed:

1. box-shell incommensurability,
2. catastrophic shell leakage,
3. orbit-level collapse inside the retained BCC shell.

The remaining obstruction is almost purely dynamical:

$$\boxed{\text{rel_residual} \approx 0.724}$$

is still too large, and

$$\boxed{\text{rel_step} \approx 1.74 \times 10^{-2}}$$

shows that the flow is still evolving.

So the accepted diagnosis is now:

$$\boxed{\text{the branch geometry is under control, but the PDE has not yet relaxed to a stationary fixed point.}}$$

CCCIV. 6. HONEST STATUS OF THE PENALTY LAYER

This improvement must be described honestly.

The weak orbit-balancing step is currently an algorithmic stabilization device. It is not yet claimed to be part of the final constitutive microscopic PDE.

Therefore the correct interpretation is:

$$\boxed{\text{there exists a nearby orbit-symmetric BCC branch candidate under the present working Class II flow,}} \quad (143)$$

but not yet

$$\boxed{\text{a final accepted actual branch of the unmodified constitutive PDE.}} \quad (144)$$

Still, this is a major success, because it proves that the project is no longer blocked by branch-geometry instability.

CCCV. 7. CORRECT NEXT STEP

Since the shell and orbit diagnostics are now both healthy, the next nonredundant move is no longer geometric. The next step must target residual reduction and stationarity:

1. longer relaxation runs,
2. adaptive time stepping,
3. possibly stronger semi-implicit damping for the stiff nonlinear sector,
4. grid refinement once the coarse-grid stationary trend is confirmed.

In other words, the project has now entered the true fixed-point convergence phase of Phase I.

CCCVI. FINAL SUMMARY

A commensurate actual-branch run with a weak orbit-balancing penalty has now produced the clearest numerical status so far.

The run yields

$$\boxed{\text{shell_fraction} \approx 0.9196, \quad \text{orbit_balance} \approx 0.9014.}$$

So the BCC shell is retained and the orbit democracy is almost fully restored. This means that the project has effectively passed the branch-geometry bottlenecks:

commensurability, shell retention, orbit stabilization.

The run is still not accepted because

$$\text{rel_residual} \approx 0.724, \quad \text{rel_step} \approx 1.74 \times 10^{-2},$$

so the nonlinear PDE has not yet relaxed to a stationary fixed point.

Therefore the remaining bottleneck is no longer geometric. It is now purely dynamical:

$$\boxed{\text{actual stationary convergence of the nonlinear PDE.}}$$

The weak orbit-balancing layer should currently be interpreted as an algorithmic stabilization device, not yet as a final physical constitutive term. But its success shows that a nearby orbit-symmetric BCC actual branch candidate is numerically present.

CCCVII. STAGED-DAMPING RESULT: GEOMETRY IS UNDER CONTROL, RESIDUAL PLATEAU IS THE FINAL BOTTLENECK

CCCVIII. SHORT KOREAN EXPLANATION

The latest numerical run changes the status of the actual-branch program in an important way. Keeping fixed:

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}},$$

and using the same weak orbit-balancing penalty, we ran a longer staged-damping relaxation.

The outcome is decisive: both the shell retention and the orbit symmetry are now excellent, while the PDE residual remains stuck near a plateau of order 10^{-1} .

Therefore the project is no longer blocked by branch geometry. It is now blocked almost purely by stationary PDE convergence.

CCCIX. 1. SETUP

The run used the same commensurate BCC-compatible setup:

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}}.$$

The same $2 + 2 + 8$ orbit-projected seed and the same weak shell-mode orbit-balancing layer were kept. Only the time step was scheduled in stages:

$$\Delta\tau : 0.008 \rightarrow 0.006 \rightarrow 0.004 \rightarrow 0.003.$$

Thus the purpose of the run was not to alter branch geometry, but only to improve relaxation convergence.

CCCX. 2. FINAL DIAGNOSTICS

The final state at step 80 gave

$$\text{accepted} = \text{False},$$

$$\text{rel_step} \approx 5.256752 \times 10^{-3},$$

$$\text{rel_residual} \approx 7.265510 \times 10^{-1},$$

$$\text{shell_fraction} \approx 9.183939 \times 10^{-1},$$

$$\text{orbit_balance} \approx 9.962593 \times 10^{-1}.$$

Hence the branch is still not accepted, but the reason is now highly localized.

CCCXI. 3. WHAT HAS NOW BEEN EFFECTIVELY SOLVED

The run shows that the following difficulties are no longer dominant:

a. (i) Box-shell mismatch. This was already eliminated by the commensurate choice $L = 4\pi$, $q_0 = 1/\sqrt{2}$.

b. (ii) Shell leakage. The shell fraction remains stably large:

$$0.9355, 0.9307, 0.9280, 0.9255, 0.9236, 0.9216, 0.9200, 0.9184.$$

So the branch stays on the intended BCC shell.

c. (iii) Orbit collapse. The orbit-balance indicator improves monotonically:

$$0.3089 \rightarrow 0.5882 \rightarrow 0.7900 \rightarrow 0.9016 \rightarrow 0.9558 \rightarrow 0.9805 \rightarrow 0.9914 \rightarrow 0.9963.$$

Thus the branch also becomes nearly perfectly orbit-symmetric within the retained shell. Therefore the geometric part of the actual-branch search is now effectively under control.

CCCXII. 4. THE TRUE REMAINING BOTTLENECK

In sharp contrast, the residual does not decay toward zero:

$$0.7102, 0.6947, 0.7018, 0.7099, 0.7152, 0.7200, 0.7234, 0.7266.$$

So the flow approaches a shell-preserving, orbit-symmetric state, but not a true stationary solution of the present PDE. The relative step does decrease,

$$0.0151 \rightarrow 0.0145 \rightarrow 0.0108 \rightarrow 0.0106 \rightarrow 0.00708 \rightarrow 0.00704 \rightarrow 0.00527 \rightarrow 0.00526,$$

but this only indicates numerical slowing, not actual PDE satisfaction.

Hence the correct diagnosis is now:

the run is approaching a stabilized quasi-branch, but not yet a true stationary PDE fixed point.

CCCXIII. 5. INTERPRETATION

This result has two possible explanations.

a. Possibility A: constitutive mismatch. The current explicit working Class II callback

$$\mathcal{G}^{(II)}[\Psi]$$

may still differ in an essential way from the actual constitutive term required by the intended microscopic branch. Then the flow can preserve the right shell and orbit geometry without satisfying the true stationary Euler–Lagrange condition.

b. Possibility B: algorithmic projection artifact. The weak orbit-balancing penalty may stabilize a nearby orbit-symmetric branch candidate that is not itself a stationary solution of the unmodified PDE. Then the penalty helps reveal the nearby branch geometry, but not the true physical fixed point.

In either case, the obstruction is now no longer geometric. It is a stationary residual obstruction.

CCCXIV. 6. CORRECT PROJECT STATUS AFTER THIS RUN

The actual-branch program has now reached the following precise status:

shell-retaining BCC branch candidate: yes,

orbit-symmetric BCC branch candidate: yes,

accepted stationary PDE solution: not yet.

This is a major narrowing of the frontier.

CCCXV. 7. CORRECT NEXT STEP

Because the geometry is now under control, the next nonredundant task is no longer to tune shell or orbit mechanisms.

The next step must directly attack the residual plateau:

1. remove or soften the penalty after the orbit-symmetric branch has formed, and see whether the branch remains nearby;
2. compare the residual with and without the penalty on the same branch;
3. refine the explicit Class II callback so that the branch geometry and the stationary PDE condition become compatible;
4. only after that attempt longer / finer-grid acceptance runs.

The key point is that the frontier has shifted from branch discovery to constitutive/stationary compatibility.

CCCXVI. FINAL SUMMARY

The staged-damping run has now isolated the actual bottleneck of the Phase-I program.

With the commensurate setup

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}},$$

together with the weak orbit-balancing stabilization, the numerical branch satisfies

$$\text{shell_fraction} \approx 0.9184, \quad \text{orbit_balance} \approx 0.9963.$$

So the solver now successfully preserves both the BCC shell and the orbit democracy. This means the branch-geometry bottlenecks are effectively solved.

However, the residual remains stuck near

$$\text{rel_residual} \approx 0.7266,$$

while the relative step only decreases to

$$\text{rel_step} \approx 5.26 \times 10^{-3}.$$

Therefore the remaining obstruction is no longer geometric. It is the mismatch between the presently stabilized branch and a true stationary solution of the nonlinear PDE.

The correct current status is:

$$\text{orbit-symmetric BCC branch candidate: found,}$$

but

$$\text{accepted stationary actual branch: not yet found.}$$

Hence the next step must attack the residual plateau directly, rather than further modifying shell or orbit geometry.

CCCXVII. PENALTY CONTINUATION TEST: THE ORBIT-SYMMETRIC BCC BRANCH PERSISTS WITHOUT PENALTY

CCCXVIII. SHORT KOREAN EXPLANATION

A continuation test was performed in which the weak orbit-balancing penalty was gradually turned off while keeping the same commensurate setup

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}}.$$

The result is decisive:

1. the shell-retaining, orbit-symmetric BCC branch does not disappear when the penalty is removed;
2. however, the PDE residual does not relax toward zero and instead drifts upward slightly.

Therefore the penalty should now be interpreted as a basin-entry device, not as the source of the branch itself.

CCCXIX. 1. CONTINUATION SCHEDULE

The run used the schedule

$$(\Delta\tau, \varepsilon_{\text{pen}}) = (0.008, 0.08) \rightarrow (0.006, 0.05) \rightarrow (0.004, 0.02) \rightarrow (0.003, 0.005) \rightarrow (0.003, 0).$$

Thus the orbit-balancing layer was turned off continuously rather than removed abruptly.

CCCXX. 2. BLOCKWISE DIAGNOSTICS

The block summaries were:

$$\varepsilon_{\text{pen}} = 0.08 : \quad \text{rel_residual} \approx 0.6947, \quad \text{shell_fraction} \approx 0.9307, \quad \text{orbit_balance} \approx 0.5882.$$

$$\varepsilon_{\text{pen}} = 0.05 : \quad \text{rel_residual} \approx 0.7099, \quad \text{shell_fraction} \approx 0.9256, \quad \text{orbit_balance} \approx 0.8226.$$

$$\varepsilon_{\text{pen}} = 0.02 : \quad \text{rel_residual} \approx 0.7200, \quad \text{shell_fraction} \approx 0.9217, \quad \text{orbit_balance} \approx 0.8770.$$

$$\varepsilon_{\text{pen}} = 0.005 : \quad \text{rel_residual} \approx 0.7266, \quad \text{shell_fraction} \approx 0.9184, \quad \text{orbit_balance} \approx 0.8879.$$

$$\varepsilon_{\text{pen}} = 0 : \quad \text{rel_residual} \approx 0.7322, \quad \text{shell_fraction} \approx 0.9149, \quad \text{orbit_balance} \approx 0.8878.$$

CCCXXI. 3. MAIN INTERPRETATION

The decisive point is that after the penalty is turned fully off,

$$\boxed{\text{shell_fraction} \approx 0.9149}$$

and

$$\boxed{\text{orbit_balance} \approx 0.8878}$$

remain large.

Therefore the orbit-symmetric BCC branch candidate does not collapse when the penalty disappears. This means the penalty is not merely forcing an artificial branch. Instead, it helps the flow enter the basin of a nearby unpenalized branch candidate.

So the correct statement is:

the near-BCC branch appears to survive continuation to the unpenalized flow.

CCCXXII. 4. WHAT STILL FAILS

The final diagnostics were

$$\text{rel_step} \approx 5.229 \times 10^{-3}, \quad \text{rel_residual} \approx 7.322 \times 10^{-1}.$$

Thus the flow still does not approach a true stationary PDE solution. Moreover, the residual slightly increases as the penalty is reduced:

$$0.6947 \rightarrow 0.7099 \rightarrow 0.7200 \rightarrow 0.7266 \rightarrow 0.7322.$$

Hence the remaining obstruction is not shell geometry and not orbit symmetry. It is the stationary residual compatibility of the current explicit working Class II callback.

CCCXXIII. 5. UPDATED PROJECT STATUS

The actual Phase-I status is now:

commensurate shell-retaining branch candidate: yes,

orbit-symmetric branch candidate: yes,

branch survives continuation to zero penalty: yes,

accepted stationary solution of the present PDE: not yet.

This is a major narrowing of the frontier.

CCCXXIV. 6. CORRECT NEXT MOVE

Since the branch now survives removal of the stabilization layer, the next nonredundant step is no longer another continuation test.

The correct next step is to decompose the residual:

$$R[\Psi] = R_{\text{lin}} + R_{\text{loc}} + R_{\text{Class II}},$$

with

$$R_{\text{lin}} = -\mu^2 \Psi + Z \nabla^2 \Psi - Y \nabla^4 \Psi,$$

$$R_{\text{loc}} = -\lambda |\Psi|^2 \Psi - \gamma |\Psi|^4 \Psi,$$

$$R_{\text{Class II}} = \mathcal{G}^{(II)}[\Psi].$$

Then one must diagnose which sector dominates the plateau:

$$\|R_{\text{lin}}\|, \quad \|R_{\text{loc}}\|, \quad \|R_{\text{Class II}}\|.$$

This will tell us whether the plateau comes mainly from:

1. the linear shell choice,
2. the local nonlinear potential,
3. or the current explicit Class II constitutive callback.

That is now the sharpest next step.

CCCXXV. FINAL SUMMARY

The continuation test shows that the shell-retaining, orbit-symmetric BCC branch candidate survives even after the weak orbit-balancing penalty is reduced all the way to zero.

At the final unpenalized stage one still finds

$$\text{shell_fraction} \approx 0.9149, \quad \text{orbit_balance} \approx 0.8878.$$

Therefore the penalty is best interpreted as a basin-entry numerical device, not as the source of the branch itself. However, the PDE residual remains large:

$$\text{rel_residual} \approx 0.7322.$$

So the true remaining bottleneck is now sharply identified:

$$\text{not branch geometry, but stationary residual compatibility of the current explicit Class II callback.}$$

The next nonredundant task is therefore to decompose the residual into its linear, local-nonlinear, and Class-II pieces and identify which sector is responsible for the plateau.

CCCXXVI. RESIDUAL DECOMPOSITION: THE PLATEAU IS DOMINATED BY THE LINEAR SECTOR

CCCXXVII. SHORT KOREAN EXPLANATION

The residual decomposition has now isolated the true source of the plateau.

On the final unpenalized continuation branch candidate, the relative residual norms are:

$$\|R_{\text{tot}}\|_{\text{rel}} \approx 0.7322,$$

$$\|R_{\text{lin}}\|_{\text{rel}} \approx 0.7577, \quad \|R_{\text{loc}}\|_{\text{rel}} \approx 0.0341, \quad \|R_{\text{Class II}}\|_{\text{rel}} \approx 1.18 \times 10^{-5}.$$

Thus the plateau is not dominated by the Class II constitutive term. It is dominated almost entirely by the linear shell operator.

CCCXXVIII. 1. RESIDUAL DECOMPOSITION

The stationary PDE residual was decomposed as

$$R[\Psi] = R_{\text{lin}} + R_{\text{loc}} + R_{\text{Class II}},$$

with

$$R_{\text{lin}} = -\mu^2 \Psi + Z \nabla^2 \Psi - Y \nabla^4 \Psi,$$

$$R_{\text{loc}} = -\lambda |\Psi|^2 \Psi - \gamma |\Psi|^4 \Psi,$$

$$R_{\text{Class II}} = \mathcal{G}^{(II)}[\Psi].$$

At the final unpenalized continuation state, we found

$$\|R_{\text{lin}}\|_{\text{rel}} \approx 0.7577,$$

$$\|R_{\text{loc}}\|_{\text{rel}} \approx 0.0341,$$

$$\|R_{\text{Class II}}\|_{\text{rel}} \approx 1.18 \times 10^{-5},$$

while

$$\|R_{\text{tot}}\|_{\text{rel}} \approx 0.7322.$$

Therefore the residual plateau is overwhelmingly linear-sector dominated.

CCCXXIX. 2. SHELL-MARGINALITY CHECK

For a shell mode

$$\Psi \sim e^{iq \cdot x},$$

the linear operator contributes

$$\nabla^2 \Psi = -q^2 \Psi, \quad \nabla^4 \Psi = q^4 \Psi,$$

so

$$R_{\text{lin}} = \left(-\mu^2 - Zq^2 - Yq^4\right)\Psi.$$

The present commensurate actual-branch setup uses

$$Z = -1, \quad Y = 1, \quad q_* = \sqrt{-\frac{Z}{2Y}} = \frac{1}{\sqrt{2}}.$$

Hence

$$q_*^2 = \frac{1}{2}, \quad q_*^4 = \frac{1}{4},$$

and therefore

$$\boxed{\mathcal{L}_{\text{lin}}(q_*) = -\mu^2 + \frac{1}{4}.$$

With the current skeleton choice

$$\mu^2 = 1,$$

this gives

$$\boxed{\mathcal{L}_{\text{lin}}(q_*) = -\frac{3}{4} = -0.75.$$

This agrees strikingly with the observed residual scale

$$\|R_{\text{lin}}\|_{\text{rel}} \approx 0.7577.$$

So the plateau is not mysterious. It is the direct consequence of a shell-mass mismatch in the linear sector.

CCCXXX. 3. IMMEDIATE CONSEQUENCE

To make the preferred shell marginal at linear order, one must impose

$$\mathcal{L}_{\text{lin}}(q_*) = 0.$$

This yields

$$-\mu^2 + \frac{1}{4} = 0 \quad \Longrightarrow \quad \boxed{\mu^2 = \frac{1}{4}}.$$

Equivalently, in the general (Z, Y, q_*) notation,

$$\boxed{\mu^2 = -Zq_*^2 - Yq_*^4}.$$

For the present fixed values $Z = -1$, $Y = 1$, $q_* = 1/\sqrt{2}$, this is exactly

$$\boxed{\mu_{\text{crit}}^2 = \frac{1}{4}}.$$

CCCXXXI. 4. INTERPRETATION

This changes the project diagnosis substantially.

Previously, the residual plateau seemed to suggest a possible incompatibility between the current explicit Class II callback and the stationary PDE.

But the actual decomposition shows:

$$\boxed{\text{the dominant obstruction is not } \mathcal{G}^{(II)}[\Psi],}$$

$$\boxed{\text{it is the arbitrary skeleton choice } \mu^2 = 1.}$$

Thus the next move is not to redesign the Class II constitutive callback. It is to retune the linear mass parameter so that the retained BCC shell is actually marginal at linear order.

CCCXXXII. 5. UPDATED PROJECT STATUS

The actual Phase-I status is now:

$$\boxed{\text{commensurate shell retention: solved,}}$$

$$\boxed{\text{orbit-symmetric branch survival: solved,}}$$

$$\boxed{\text{residual plateau source: identified,}}$$

$$\boxed{\text{remaining task: linear mass retuning } \mu^2 \rightarrow \mu_{\text{crit}}^2.}$$

So the bottleneck is no longer vague and no longer geometric. It is a single scalar mismatch in the linear sector.

CCCXXXIII. 6. CORRECT NEXT STEP

The immediate nonredundant next computation is therefore:

$$\boxed{\mu^2 : 1 \longrightarrow \frac{1}{4}}$$

with all other ingredients frozen:

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}}, \quad Z = -1, \quad Y = 1,$$

and the same continuation branch protocol.

Only after that returned run should we re-evaluate whether the stationary actual branch is reached.

CCCXXXIV. FINAL SUMMARY

The residual plateau has now been cleanly diagnosed.

On the final unpenalized continuation branch candidate,

$$\|R_{\text{tot}}\|_{\text{rel}} \approx 0.7322,$$

but the decomposition shows

$$\boxed{\|R_{\text{lin}}\|_{\text{rel}} \approx 0.7577, \quad \|R_{\text{loc}}\|_{\text{rel}} \approx 0.0341, \quad \|R_{\text{Class II}}\|_{\text{rel}} \approx 1.18 \times 10^{-5}.$$

Therefore the plateau is dominated almost entirely by the linear sector.

For the retained shell

$$q_* = \frac{1}{\sqrt{2}},$$

with

$$Z = -1, \quad Y = 1,$$

the linear shell coefficient is

$$\mathcal{L}_{\text{lin}}(q_*) = -\mu^2 + \frac{1}{4}.$$

The present skeleton choice $\mu^2 = 1$ therefore gives

$$\mathcal{L}_{\text{lin}}(q_*) = -\frac{3}{4},$$

which matches the observed plateau scale.

So the next step is now sharply determined:

$$\boxed{\mu^2 \rightarrow \mu_{\text{crit}}^2 = \frac{1}{4}.$$

The project is no longer blocked by shell geometry, orbit symmetry, or an obviously dominant Class II inconsistency. The remaining obstruction is a linear mass mismatch in the actual solver skeleton.

CCCXXXV. WHAT THE NUMBERS ACTUALLY MEAN

The computed quantities are not mere numerical outputs. Each of them is a coordinate of an underlying geometric or dynamical structure.

$$\boxed{\text{shell_fraction} \longleftrightarrow \text{retention of the BCC finite-}q_0 \text{ shell geometry,}} \quad (145)$$

$$\boxed{\text{orbit_balance} \longleftrightarrow \text{survival of the 12-direction BCC orbit democracy,}} \quad (146)$$

$$\boxed{N(G_*) = 4 \longleftrightarrow \text{the exact BCC shell pair combinatorics entering the longitudinal witness,}} \quad (147)$$

$$\boxed{R < 12U_d \longleftrightarrow \text{the exact energetic condition distinguishing true BCC stabilization from one-patch collapse.}} \quad (148)$$

So the numbers are not “just numbers.” They are the measurable coordinates of the geometric branch structure. The recent actual-branch computations already show that:

1. the BCC shell can be retained,
2. the orbit-symmetric branch can survive continuation to zero penalty,
3. the remaining obstruction is no longer branch geometry but stationary PDE compatibility.

Thus the program has moved from

“is there any structure at all?”

to

“why is the surviving structure not yet an exact stationary solution?”

which is a much sharper and more advanced stage.

CCCXXXVI. LINEAR-SECTOR RETUNING TEST IN THE ACTUAL BRANCH PROGRAM

CCCXXXVII. SHORT KOREAN EXPLANATION

A light but decisive continuation comparison was carried out on the actual-branch solver. The setup was kept fixed at the commensurate BCC-compatible values

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}},$$

together with the same penalty-continuation schedule. Only the linear mass parameter was changed:

$$\mu^2 : 1 \longrightarrow \frac{1}{4}.$$

The result confirms that the previously identified shell-marginality diagnosis was correct.

CCCCXXVIII. 1. WHY $\mu^2 = \frac{1}{4}$ IS THE CRITICAL VALUE

For a shell mode

$$\Psi \sim e^{iq \cdot x},$$

the linear stationary operator contributes

$$R_{\text{lin}} = -\mu^2 \Psi + Z \nabla^2 \Psi - Y \nabla^4 \Psi = (-\mu^2 - Zq^2 - Yq^4) \Psi.$$

With the frozen solver values

$$Z = -1, \quad Y = 1, \quad q_* = \sqrt{-\frac{Z}{2Y}} = \frac{1}{\sqrt{2}},$$

we have

$$q_*^2 = \frac{1}{2}, \quad q_*^4 = \frac{1}{4}.$$

Hence the shell-marginality condition

$$-\mu^2 - Zq_*^2 - Yq_*^4 = 0$$

becomes

$$-\mu^2 + \frac{1}{2} - \frac{1}{4} = 0,$$

so

$$\boxed{\mu_{\text{crit}}^2 = \frac{1}{4}}.$$

Thus $\mu^2 = \frac{1}{4}$ is not an arbitrary retune. It is the unique linear value that makes the retained shell marginal.

CCCCXXIX. 2. CONTINUATION COMPARISON

We compare two continuation runs:

$$\mu^2 = 1 \quad \text{versus} \quad \mu^2 = \frac{1}{4},$$

with the same schedule

$$(\Delta\tau, \varepsilon_{\text{pen}}) = (0.008, 0.08) \rightarrow (0.006, 0.05) \rightarrow (0.004, 0.02) \rightarrow (0.003, 0.005) \rightarrow (0.003, 0).$$

At the final unpenalized stage, the original choice $\mu^2 = 1$ gives

$$\boxed{\text{rel_residual} \approx 0.7056},$$

$$\boxed{\|R_{\text{lin}}\|_{\text{rel}} \approx 0.7608, \quad \|R_{\text{loc}}\|_{\text{rel}} \approx 0.0724, \quad \|R_{\text{Class II}}\|_{\text{rel}} \approx 2.74 \times 10^{-5}}.$$

After retuning to $\mu^2 = \frac{1}{4}$, the final unpenalized stage gives

$$\boxed{\text{rel_residual} \approx 0.1196},$$

$$\boxed{\|R_{\text{lin}}\|_{\text{rel}} \approx 0.0659, \quad \|R_{\text{loc}}\|_{\text{rel}} \approx 0.0959, \quad \|R_{\text{Class II}}\|_{\text{rel}} \approx 3.87 \times 10^{-5}}.$$

CCCXL. 3. MAIN CONCLUSION

The decisive changes are

$$0.7056 \rightarrow 0.1196$$

for the total residual, and

$$0.7608 \rightarrow 0.0659$$

for the linear residual.

Meanwhile the branch-geometry observables remain essentially unchanged:

$$\text{shell_fraction} : 0.9264 \rightarrow 0.9261, \quad \text{orbit_balance} : 0.5664 \rightarrow 0.5663.$$

Therefore the retuning

$$\mu^2 = 1 \rightarrow \mu^2 = \frac{1}{4}$$

fixes the dominant linear obstruction without damaging the BCC shell structure.

CCCXLI. 4. UPDATED BOTTLENECK DIAGNOSIS

After the retuning, the dominant residual is no longer the linear part. Instead,

$$\|R_{\text{loc}}\|_{\text{rel}} \approx 0.0959$$

is now the leading nontrivial contribution, while

$$\|R_{\text{Class II}}\|_{\text{rel}} \sim 10^{-5}$$

remains negligible at this scale.

So the Phase-I bottleneck has shifted from

linear shell mismatch

to

$$\text{local nonlinear-sector mismatch and full stationary convergence.}$$

CCCXLII. 5. IMMEDIATE NEXT STEP

The next nonredundant computation is to freeze

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}}, \quad \mu^2 = \frac{1}{4}$$

and rerun the full 16^3 continuation at this retuned value.

Only after that full-scale run should one decide whether additional tuning of the local nonlinear potential (λ, γ) is necessary.

CCCXLIII. FINAL SUMMARY

The shell-marginality retuning test confirms that the previous residual plateau was indeed dominated by the linear sector.

With the commensurate setup fixed,

$$L = 4\pi, \quad q_0 = q_* = \frac{1}{\sqrt{2}},$$

the retuning

$$\mu^2 = 1 \rightarrow \frac{1}{4}$$

reduces the final unpenalized continuation residual from

$$0.7056 \rightarrow 0.1196,$$

while the linear residual drops from

$$0.7608 \rightarrow 0.0659.$$

At the same time, the shell and orbit diagnostics remain essentially unchanged.

Therefore the retuning is validated. The dominant remaining obstruction is no longer the linear shell operator. It is now the local nonlinear sector:

$$\|R_{\text{loc}}\|_{\text{rel}} \approx 0.0959.$$

The correct next move is now fixed: freeze

$$L = 4\pi, \quad q_0 = \frac{1}{\sqrt{2}}, \quad \mu^2 = \frac{1}{4}$$

and run the full 16^3 continuation at this value.

CCCXLIV. EXACT REDUCED PROGRAM FOR THE RIGOROUS EXTRACTION OF THE BACKEND NUMBERS

CCCXLIV. SHORT KOREAN EXPLANATION

We now stop the broad numerical scan of the backend point

$$(\mu^2, \lambda, \gamma)$$

and replace it by a finite analytic extraction program.

The key idea is to move from a direct search in the PDE parameter space to an exact reduced shell-amplitude theory induced by the Class II X -sector. The remaining numerical task must then be only the finite evaluation of a small set of projected coefficients.

CCCXLVI. 1. MINIMAL HONEST SHELL-AMPLITUDE SYSTEM

The correct reduced starting point is the inter-patch-coupled shell-amplitude equation

$$\partial_t A_\alpha = \left(\sigma_0 + \eta P_2^\alpha\right) A_\alpha - u |A_\alpha|^2 A_\alpha - v \left(\sum_\beta |A_\beta|^2\right) A_\alpha - \kappa \left(\sum_\beta M_{\alpha\beta} |A_\beta|^2\right) A_\alpha. \quad (149)$$

Here:

$$P_2^\alpha = P_2(\hat{n} \cdot \hat{k}_\alpha), \quad M_{\alpha\beta} \sim (\hat{k}_\alpha \cdot \hat{k}_\beta)^2,$$

and the five coefficients

$$\boxed{\sigma_0, \quad \eta, \quad u, \quad v, \quad \kappa}$$

are the true reduced unknowns.

Thus the problem is no longer to scan directly over

$$(\mu^2, \lambda, \gamma),$$

but to extract a finite reduced coefficient set.

CCCXLVII. 2. STATIONARY INTENSITY EQUATIONS

Write

$$r_\alpha := |A_\alpha|^2.$$

For stationary amplitudes with fixed phases, (149) reduces to

$$\boxed{0 = \sigma_0 + \eta P_2^\alpha - u r_\alpha - v \sum_{\beta} r_\beta - \kappa \sum_{\beta} M_{\alpha\beta} r_\beta, \quad \alpha = 1, \dots, 12.} \quad (150)$$

This is the exact finite algebraic system whose critical points determine the actual BCC branch basin.

CCCXLVIII. 3. BCC-SYMMETRIC QUARTIC REDUCTION

By BCC symmetry, the quartic coupling matrix is specified by only four numbers:

$$\boxed{U_d, \quad U_{\text{opp}}, \quad U_{\perp}, \quad U_{\text{obl}}.} \quad (151)$$

The corresponding row sum is

$$\boxed{R = U_d + U_{\text{opp}} + 2U_{\perp} + 8U_{\text{obl}}.} \quad (152)$$

Then the one-patch branch has

$$\boxed{r_{\text{1patch}} = \frac{\sigma}{U_d}, \quad \mathcal{F}_{\text{1patch}} = -\frac{\sigma^2}{2U_d},} \quad (153)$$

while the equal-amplitude 12-patch BCC branch has

$$\boxed{r_{\text{BCC}} = \frac{\sigma}{R}, \quad \mathcal{F}_{\text{BCC}} = -6\frac{\sigma^2}{R}.} \quad (154)$$

Therefore the sharp energetic criterion is

$$\boxed{\mathcal{F}_{\text{BCC}} < \mathcal{F}_{\text{1patch}} \iff R < 12U_d.} \quad (155)$$

This converts the branch-selection problem into an exact algebraic inequality.

CCCXLIX. 4. ORBIT REDUCTION

Because the axis-locked branch naturally splits the shell into

$$2 + 2 + 8,$$

we may reduce the 12 amplitudes to three orbit variables:

$$A_A, \quad A_E, \quad A_O.$$

Equivalently,

$$r_A = |A_A|^2, \quad r_E = |A_E|^2, \quad r_O = |A_O|^2.$$

Then the full stationary system (150) reduces to a 3×3 algebraic critical-point problem

$$\boxed{\frac{\partial \mathcal{F}_{\text{orb}}}{\partial r_A} = 0, \quad \frac{\partial \mathcal{F}_{\text{orb}}}{\partial r_E} = 0, \quad \frac{\partial \mathcal{F}_{\text{orb}}}{\partial r_O} = 0.} \quad (156)$$

The backend numbers are then no longer selected by brute-force PDE scanning, but by the critical point of this reduced orbit free energy.

CCCL. 5. FINAL FINITE EXTRACTION TASK

Once the reduced formulas are fixed, the only remaining numerical task is the finite projection of the coefficients

$$\sigma_0, \quad \eta, \quad u, \quad v, \quad \kappa$$

or equivalently

$$U_d, \quad U_{\text{opp}}, \quad U_{\perp}, \quad U_{\text{obl}}$$

from the intended BCC branch of the microscopic TECT/Class II system.

At that point, the backend point

$$(\mu^2, \lambda, \gamma)$$

is not searched for directly. It is obtained as the induced stationary critical point of the reduced amplitude theory.

CCCLI. 6. IMMEDIATE ROADMAP

The rigorous backend program is therefore:

$$\boxed{(\Psi, X_i^a) \implies \mathcal{F}_{\text{shell}}[A_1, \dots, A_{12}] \implies \mathcal{F}_{\text{orb}}(r_A, r_E, r_O) \implies (r_A, r_E, r_O)_{\text{crit}} \implies (\mu^2, \lambda, \gamma)_{\text{backend}}.} \quad (157)$$

So the next step is not another parameter scan. It is the explicit derivation of $\mathcal{F}_{\text{orb}}$ and its stationary equations.

CCCLII. EXACT ORBIT FREE ENERGY FOR THE BACKEND-EXTRACTION PROGRAM

CCCLIII. SHORT KOREAN EXPLANATION

We now perform the second step of the rigorous backend program.

The direct numerical scan over

$$(\mu^2, \lambda, \gamma)$$

is replaced by an exact reduced orbit free energy on the retained BCC shell. The key point is that the 12-patch quartic interaction reduces to only four coupling numbers,

$$U_d, \quad U_{\text{opp}}, \quad U_{\perp}, \quad U_{\text{obl}},$$

and the axis-locked branch reduces further to the $2 + 2 + 8$ orbit variables

$$r_A, \quad r_E, \quad r_O.$$

This turns the backend-extraction problem into a finite algebraic critical-point problem.

CCCLIV. 1. STARTING POINT: THE HONEST REDUCED SHELL-AMPLITUDE SYSTEM

We start from the minimal inter-patch shell system

$$\partial_t A_\alpha = \left(\sigma_0 + \eta P_2^\alpha \right) A_\alpha - \sum_{\beta=1}^{12} U_{\alpha\beta} |A_\beta|^2 A_\alpha, \quad \alpha = 1, \dots, 12, \quad (158)$$

where

$$U_{\alpha\beta} = u \delta_{\alpha\beta} + v + \kappa M_{\alpha\beta}. \quad (159)$$

We assume the BCC shell

$$\hat{k}_\alpha \in \mathcal{S}_{12},$$

with locking axis

$$\hat{n} = \frac{(1, 1, 0)}{\sqrt{2}}.$$

The Legendre anisotropy values are then

$$P_2(\hat{n} \cdot \hat{k}_\alpha) = \begin{cases} 1, & \alpha \in A \quad (2 \text{ axial patches}), \\ -\frac{1}{2}, & \alpha \in E \quad (2 \text{ equatorial patches}), \\ -\frac{1}{8}, & \alpha \in O \quad (8 \text{ oblique patches}). \end{cases} \quad (160)$$

Hence the effective linear coefficients are

$$\boxed{\sigma_A = \sigma_0 + \eta, \quad \sigma_E = \sigma_0 - \frac{\eta}{2}, \quad \sigma_O = \sigma_0 - \frac{\eta}{8}.} \quad (161)$$

CCCLV. 2. EXACT FOUR-NUMBER REDUCTION OF THE QUARTIC MATRIX

By shell symmetry, the quartic kernel $U_{\alpha\beta}$ takes only four values, depending on the relation of β to α :

$$\boxed{U_d, \quad U_{\text{opp}}, \quad U_{\perp}, \quad U_{\text{obl}}.} \quad (162)$$

Here:

$$U_d : \beta = \alpha, \quad U_{\text{opp}} : \hat{k}_\beta = -\hat{k}_\alpha, \quad U_{\perp} : \hat{k}_\alpha \cdot \hat{k}_\beta = 0, \quad U_{\text{obl}} : |\hat{k}_\alpha \cdot \hat{k}_\beta| = \frac{1}{2}.$$

Thus the full 12×12 quartic matrix is specified by only the four numbers in (162).

CCCLVI. 3. ORBIT COUNTING FOR THE $2 + 2 + 8$ SPLIT

Let

$$N_A = 2, \quad N_E = 2, \quad N_O = 8.$$

For the orbit-symmetric ansatz

$$r_\alpha = \begin{cases} r_A, & \alpha \in A, \\ r_E, & \alpha \in E, \\ r_O, & \alpha \in O, \end{cases} \quad r_I := |A_I|^2, \quad (163)$$

the ordered pair counts are exactly:

$$\boxed{C_{AA} = 2U_d + 2U_{\text{opp}}, \quad C_{EE} = 2U_d + 2U_{\text{opp}},} \quad (164)$$

$$\boxed{C_{AE} = 4U_\perp, \quad C_{AO} = 16U_{\text{obl}}, \quad C_{EO} = 16U_{\text{obl}},} \quad (165)$$

$$\boxed{C_{OO} = 8U_d + 8U_{\text{opp}} + 16U_\perp + 32U_{\text{obl}}.} \quad (166)$$

These are exact consequences of the cuboctahedral BCC shell combinatorics.

CCCLVII. 4. EXACT ORBIT FREE ENERGY

The reduced free energy whose gradient reproduces the stationary version of (158) is

$$\mathcal{F} = - \sum_{\alpha=1}^{12} \sigma_\alpha r_\alpha + \frac{1}{2} \sum_{\alpha,\beta=1}^{12} U_{\alpha\beta} r_\alpha r_\beta. \quad (167)$$

Under the orbit ansatz (163), this becomes

$$\begin{aligned} \boxed{\mathcal{F}_{\text{orb}}(r_A, r_E, r_O)} &= -2\sigma_A r_A - 2\sigma_E r_E - 8\sigma_O r_O \\ &\quad + (U_d + U_{\text{opp}})(r_A^2 + r_E^2) + 4U_\perp r_A r_E \\ &\quad + 16U_{\text{obl}}(r_A + r_E)r_O + (4U_d + 4U_{\text{opp}} + 8U_\perp + 16U_{\text{obl}})r_O^2. \end{aligned} \quad (168)$$

This is the exact $2 + 2 + 8$ orbit free energy.

CCCLVIII. 5. EXACT STATIONARY EQUATIONS

Taking derivatives of (168), we obtain the stationary orbit system

$$\boxed{\sigma_A = (U_d + U_{\text{opp}})r_A + 2U_\perp r_E + 8U_{\text{obl}}r_O,} \quad (169)$$

$$\boxed{\sigma_E = 2U_\perp r_A + (U_d + U_{\text{opp}})r_E + 8U_{\text{obl}}r_O,} \quad (170)$$

$$\boxed{\sigma_O = 2U_{\text{obl}}(r_A + r_E) + (U_d + U_{\text{opp}} + 2U_\perp + 4U_{\text{obl}})r_O.} \quad (171)$$

Thus the full 12-patch stationary problem has been reduced exactly to a 3×3 algebraic system.

CCCLIX. 6. SUM/DIFFERENCE REDUCTION AND CLOSED-FORM SOLUTION

Define

$$S := r_A + r_E, \quad \Delta_r := r_A - r_E. \quad (172)$$

Subtracting (170) from (169) gives

$$\Delta_r = \frac{\sigma_A - \sigma_E}{U_d + U_{\text{opp}} - 2U_{\perp}} = \frac{3\eta/2}{U_d + U_{\text{opp}} - 2U_{\perp}}. \quad (173)$$

Adding (169) and (170), we obtain

$$(U_d + U_{\text{opp}} + 2U_{\perp})S + 16U_{\text{obl}}r_O = 2\sigma_0 + \frac{\eta}{2}. \quad (174)$$

Equation (171) becomes

$$2U_{\text{obl}}S + (U_d + U_{\text{opp}} + 2U_{\perp} + 4U_{\text{obl}})r_O = \sigma_0 - \frac{\eta}{8}. \quad (175)$$

Introduce

$$\mathcal{A} := U_d + U_{\text{opp}} + 2U_{\perp}, \quad \mathcal{B} := 16U_{\text{obl}}, \quad \mathcal{C} := 2U_{\text{obl}}, \quad \mathcal{D} := U_d + U_{\text{opp}} + 2U_{\perp} + 4U_{\text{obl}}. \quad (176)$$

Then (174)–(175) read

$$\mathcal{A}S + \mathcal{B}r_O = 2\sigma_0 + \frac{\eta}{2}, \quad \mathcal{C}S + \mathcal{D}r_O = \sigma_0 - \frac{\eta}{8}. \quad (177)$$

Hence

$$S = \frac{(2\sigma_0 + \frac{\eta}{2})\mathcal{D} - \mathcal{B}(\sigma_0 - \frac{\eta}{8})}{\mathcal{A}\mathcal{D} - \mathcal{B}\mathcal{C}}, \quad (178)$$

$$r_O = \frac{\mathcal{A}(\sigma_0 - \frac{\eta}{8}) - \mathcal{C}(2\sigma_0 + \frac{\eta}{2})}{\mathcal{A}\mathcal{D} - \mathcal{B}\mathcal{C}}. \quad (179)$$

Finally,

$$r_A = \frac{S + \Delta_r}{2}, \quad r_E = \frac{S - \Delta_r}{2}. \quad (180)$$

Thus the exact orbit critical point is obtained in closed algebraic form.

CCCLX. 7. SPECIALIZATION TO THE EXPLICIT KERNEL $M_{\alpha\beta} = (\hat{k}_{\alpha} \cdot \hat{k}_{\beta})^2$

For the explicit angular kernel

$$M_{\alpha\beta} = (\hat{k}_{\alpha} \cdot \hat{k}_{\beta})^2, \quad (181)$$

the four quartic numbers become

$$U_d = u + v + \kappa, \quad U_{\text{opp}} = v + \kappa, \quad U_{\perp} = v, \quad U_{\text{obl}} = v + \frac{\kappa}{4}. \quad (182)$$

Then the row sum

$$R := U_d + U_{\text{opp}} + 2U_{\perp} + 8U_{\text{obl}} \quad (183)$$

reduces to

$$\boxed{R = u + 12v + 4\kappa.} \quad (184)$$

The one-patch and equal-amplitude 12-patch BCC branches satisfy

$$r_{\text{1patch}} = \frac{\sigma}{U_d}, \quad \mathcal{F}_{\text{1patch}} = -\frac{\sigma^2}{2U_d}, \quad (185)$$

and

$$r_{\text{BCC}} = \frac{\sigma}{R}, \quad \mathcal{F}_{\text{BCC}} = -6\frac{\sigma^2}{R}. \quad (186)$$

Therefore the sharp BCC stability criterion

$$\mathcal{F}_{\text{BCC}} < \mathcal{F}_{\text{1patch}} \iff R < 12U_d$$

becomes

$$\boxed{11u + 8\kappa > 0.} \quad (187)$$

This is an exact closed criterion for avoiding one-patch collapse in the explicit reduced theory.

CCCLXI. 8. ORBIT FREE ENERGY DIRECTLY IN (u, v, κ) VARIABLES

Substituting (182) into (168), we obtain

$$\begin{aligned} \boxed{\mathcal{F}_{\text{orb}}(r_A, r_E, r_O)} &= -2\sigma_A r_A - 2\sigma_E r_E - 8\sigma_O r_O \\ &\quad + (u + 2v + 2\kappa)(r_A^2 + r_E^2) + 4v r_A r_E \\ &\quad + (16v + 4\kappa)(r_A + r_E)r_O + (4u + 32v + 12\kappa)r_O^2. \end{aligned} \quad (188)$$

The stationary equations are then

$$\boxed{\sigma_A = (u + 2v + 2\kappa)r_A + 2v r_E + (8v + 2\kappa)r_O,} \quad (189)$$

$$\boxed{\sigma_E = 2v r_A + (u + 2v + 2\kappa)r_E + (8v + 2\kappa)r_O,} \quad (190)$$

$$\boxed{\sigma_O = \left(2v + \frac{\kappa}{2}\right)(r_A + r_E) + (u + 8v + 3\kappa)r_O.} \quad (191)$$

The orbit splitting becomes especially simple:

$$\boxed{\Delta_r = \frac{3\eta/2}{u + 2\kappa}.} \quad (192)$$

Thus the axial-equatorial splitting is controlled only by u and κ , while the isotropic background v drops out completely.

CCCLXII. 9. CONSEQUENCE FOR THE BACKEND NUMBERS

At this point the backend-extraction problem is no longer a broad scan in

$$(\mu^2, \lambda, \gamma).$$

Instead, it has been reduced to the finite extraction of the reduced coefficients

$$u, \quad v, \quad \kappa, \quad \sigma_0, \quad \eta$$

from the intended microscopic BCC branch.

Once these are computed, the critical point

$$(r_A, r_E, r_O)$$

is fixed analytically by (178), (179), and (180), and the backend numbers follow from the induced stationary branch.

So the next step is not another brute-force search. It is the explicit microscopic projection of

$$u, \quad v, \quad \kappa, \quad \sigma_0, \quad \eta.$$

CCCLXIII. FINAL SUMMARY

The backend-extraction problem has now been reduced exactly to a finite orbit free-energy problem.

The quartic shell interaction is specified by only four numbers

$$U_d, \quad U_{\text{opp}}, \quad U_{\perp}, \quad U_{\text{obl}},$$

and the axis-locked 12-patch shell reduces to the $2 + 2 + 8$ orbit variables

$$r_A, \quad r_E, \quad r_O.$$

The exact orbit free energy is

$$\mathcal{F}_{\text{orb}}(r_A, r_E, r_O),$$

and its stationary equations reduce to a closed 3×3 algebraic system. In sum/difference variables,

$$S = r_A + r_E, \quad \Delta_r = r_A - r_E,$$

the critical point is obtained explicitly in closed form.

For the explicit kernel

$$M_{\alpha\beta} = (\hat{k}_{\alpha} \cdot \hat{k}_{\beta})^2,$$

the sharp BCC stability condition becomes

$$\boxed{11u + 8\kappa > 0.}$$

Moreover, the axial-equatorial orbit splitting is

$$\boxed{\Delta_r = \frac{3\eta/2}{u + 2\kappa},}$$

so the isotropic coefficient v drops out of the orbit splitting entirely.

Therefore the remaining backend problem is no longer a broad numerical scan. It is the finite microscopic extraction of

$$u, \quad v, \quad \kappa, \quad \sigma_0, \quad \eta,$$

after which the orbit critical point and the induced backend numbers are fixed analytically.

CCCLXIV. MICROSCOPIC PROJECTION FORMULAS FOR THE REDUCED BACKEND COEFFICIENTS

CCCLXV. SHORT KOREAN EXPLANATION

We now perform the third step of the rigorous backend program.
The numerical search over

$$(\mu^2, \lambda, \gamma)$$

must be replaced by explicit microscopic projection formulas on the intended BCC branch. The task is to project the microscopic quadratic and quartic operators onto the retained shell basis and thereby extract the reduced coefficients

$$\sigma_0, \quad \eta, \quad u, \quad v, \quad \kappa.$$

Once these coefficients are known, the orbit critical point

$$(r_A, r_E, r_O)$$

is fixed analytically by the exact reduced theory derived in the previous step.

CCCLXVI. 1. INTENDED BCC SHELL BASIS

Let the intended microscopic background be the actual BCC branch

$$\Psi_{\text{BCC}}(x),$$

with shell radius

$$|k| = q_*.$$

Choose the 12 shell directions

$$\hat{k}_\alpha \in S_{\text{BCC}}, \quad \alpha = 1, \dots, 12,$$

and an adapted shell basis

$$\boxed{\varphi_\alpha(x) = \frac{1}{\sqrt{\text{Vol}(\Omega)}} \xi_\alpha e^{iq_* \hat{k}_\alpha \cdot x}, \quad \xi_\alpha^\dagger \xi_\alpha = 1,} \quad (193)$$

where Ω is the unit cell and ξ_α denotes the internal polarization / Bloch factor of the retained branch.

All reduced coefficients will now be extracted by projection onto the finite basis $\{\varphi_\alpha\}$.

CCCLXVII. 2. QUADRATIC PROJECTION: EXTRACTING THE LINEAR SHELL DATA

Let

$$\boxed{\mathcal{H}_{\text{mic}}^{(2)} := \left. \frac{\delta^2 \mathcal{F}_{\text{mic}}}{\delta \Psi^\dagger \delta \Psi} \right|_{\Psi = \Psi_{\text{BCC}}}} \quad (194)$$

be the microscopic quadratic Hessian on the intended BCC branch.

Define the shell-projected quadratic matrix

$$\boxed{H_{\alpha\beta}^{(2)} = \int_{\Omega} d^3x \overline{\varphi_\alpha(x)} \mathcal{H}_{\text{mic}}^{(2)} \varphi_\beta(x).} \quad (195)$$

At the reduced orbit level, the first quantity needed is the shell-diagonal part

$$\boxed{h_\alpha := H_{\alpha\alpha}^{(2)}} \quad (196)$$

The reduced shell free energy begins as

$$\mathcal{F}_{\text{shell}} = - \sum_{\alpha=1}^{12} h_\alpha r_\alpha + \frac{1}{2} \sum_{\alpha,\beta=1}^{12} U_{\alpha\beta} r_\alpha r_\beta + \dots, \quad r_\alpha := |A_\alpha|^2. \quad (197)$$

CCCLXVIII. 3. EXACT PROJECTION FORMULAS FOR σ_0 AND η

Let

$$P_2^\alpha := P_2(\hat{n} \cdot \hat{k}_\alpha),$$

with the axis-locked orbit values

$$P_2^\alpha = \begin{cases} 1, & \alpha \in A, \\ -\frac{1}{2}, & \alpha \in E, \\ -\frac{1}{8}, & \alpha \in O. \end{cases} \quad (198)$$

The reduced linear coefficients are defined by the exact projection

$$\boxed{h_\alpha = \sigma_0 + \eta P_2^\alpha + h_\alpha^{\text{rem}},} \quad (199)$$

where h_α^{rem} is the part orthogonal to the $\{1, P_2\}$ sector.

The orthogonal projection onto $\{1, P_2\}$ gives

$$\boxed{\sigma_0 = \frac{1}{12} \sum_{\alpha=1}^{12} h_\alpha,} \quad (200)$$

because

$$\sum_{\alpha=1}^{12} P_2^\alpha = 0.$$

Next,

$$\sum_{\alpha=1}^{12} (P_2^\alpha)^2 = 2 \cdot 1^2 + 2 \cdot \left(\frac{1}{2}\right)^2 + 8 \cdot \left(\frac{1}{8}\right)^2 = \frac{21}{8}.$$

Therefore

$$\boxed{\eta = \frac{\sum_{\alpha=1}^{12} P_2^\alpha h_\alpha}{\sum_{\alpha=1}^{12} (P_2^\alpha)^2} = \frac{8}{21} \sum_{\alpha=1}^{12} P_2^\alpha h_\alpha.} \quad (201)$$

Thus σ_0 and η are obtained exactly by finite projection of the microscopic quadratic Hessian.

CCCLXIX. 4. ORBIT-AVERAGE FORM OF THE SAME FORMULAS

Let

$$\boxed{\sigma_A := \frac{1}{2} \sum_{\alpha \in A} h_\alpha, \quad \sigma_E := \frac{1}{2} \sum_{\alpha \in E} h_\alpha, \quad \sigma_O := \frac{1}{8} \sum_{\alpha \in O} h_\alpha.} \quad (202)$$

Then

$$\boxed{\sigma_A = \sigma_0 + \eta, \quad \sigma_E = \sigma_0 - \frac{\eta}{2}, \quad \sigma_O = \sigma_0 - \frac{\eta}{8}.} \quad (203)$$

Hence

$$\boxed{\eta = \frac{2}{3}(\sigma_A - \sigma_E), \quad \sigma_0 = \frac{\sigma_A + 2\sigma_E}{3},} \quad (204)$$

and consistency predicts

$$\boxed{\sigma_O = \frac{\sigma_A + 3\sigma_E}{4}.} \quad (205)$$

Equation (205) is the exact internal check that the linear shell data is exhausted by the (σ_0, η) truncation.

CCCLXX. 5. QUARTIC PROJECTION: EXTRACTING THE SHELL-COUPLING MATRIX

Let

$$\Gamma_{\text{mic}}^{(4)} := \frac{\delta^4 \mathcal{F}_{\text{mic}}}{\delta \Psi^\dagger \delta \Psi^\dagger \delta \Psi \delta \Psi} \Big|_{\Psi = \Psi_{\text{BCC}}} \quad (206)$$

denote the microscopic four-point vertex on the intended branch.

Instead of writing all tensor indices explicitly, define the reduced quartic shell free energy by

$$\mathcal{F}_{\text{shell}}^{(4)} = \frac{1}{2} \sum_{\alpha, \beta=1}^{12} U_{\alpha\beta} r_\alpha r_\beta. \quad (207)$$

Equivalently,

$$U_{\alpha\beta} = \frac{\partial^2 \mathcal{F}_{\text{shell}}^{(4)}}{\partial r_\alpha \partial r_\beta}. \quad (208)$$

This definition is exact and removes all normalization ambiguity. The remaining task is therefore to compute the finite matrix $U_{\alpha\beta}$ from the microscopic projected quartic vertex.

CCCLXXI. 6. EXACT CLASS AVERAGES $U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}$

The BCC shell geometry implies that each ordered pair (α, β) belongs to exactly one of four classes:

$$\beta = \alpha, \quad \beta = \bar{\alpha}, \quad \hat{k}_\alpha \cdot \hat{k}_\beta = 0, \quad |\hat{k}_\alpha \cdot \hat{k}_\beta| = \frac{1}{2}.$$

Define the corresponding ordered-pair sets

$$\mathcal{C}_d, \quad \mathcal{C}_{\text{opp}}, \quad \mathcal{C}_\perp, \quad \mathcal{C}_{\text{obl}}.$$

Their cardinalities are

$$|\mathcal{C}_d| = 12, \quad |\mathcal{C}_{\text{opp}}| = 12, \quad |\mathcal{C}_\perp| = 24, \quad |\mathcal{C}_{\text{obl}}| = 96. \quad (209)$$

The class-averaged quartic numbers are therefore

$$U_d = \frac{1}{12} \sum_{(\alpha, \beta) \in \mathcal{C}_d} U_{\alpha\beta}, \quad (210)$$

$$U_{\text{opp}} = \frac{1}{12} \sum_{(\alpha, \beta) \in \mathcal{C}_{\text{opp}}} U_{\alpha\beta}, \quad (211)$$

$$U_\perp = \frac{1}{24} \sum_{(\alpha, \beta) \in \mathcal{C}_\perp} U_{\alpha\beta}, \quad (212)$$

$$U_{\text{obl}} = \frac{1}{96} \sum_{(\alpha, \beta) \in \mathcal{C}_{\text{obl}}} U_{\alpha\beta}. \quad (213)$$

Thus the full 12×12 quartic matrix has been reduced to four finite projected numbers.

CCCLXXII. 7. EXACT INVERSION TO u, v, κ

For the reduced kernel ansatz

$$U_{\alpha\beta} = u \delta_{\alpha\beta} + v + \kappa (\hat{k}_\alpha \cdot \hat{k}_\beta)^2, \quad (214)$$

the four class values are

$$\boxed{U_d = u + v + \kappa, \quad U_{\text{opp}} = v + \kappa, \quad U_\perp = v, \quad U_{\text{obl}} = v + \frac{\kappa}{4}.} \quad (215)$$

These relations can be inverted exactly:

$$\boxed{v = U_\perp,} \quad (216)$$

$$\boxed{\kappa = 4(U_{\text{obl}} - U_\perp),} \quad (217)$$

$$\boxed{u = U_d - U_{\text{opp}} = U_d - 4U_{\text{obl}} + 3U_\perp.} \quad (218)$$

So u, v, κ are exact finite functions of the projected quartic class averages.

CCCLXXIII. 8. CONSISTENCY DEFECT OF THE (u, v, κ) TRUNCATION

The explicit kernel ansatz (214) predicts the relation

$$\boxed{U_{\text{opp}} = 4U_{\text{obl}} - 3U_\perp.} \quad (219)$$

Hence define the quartic truncation defect

$$\boxed{\Delta_{\text{ker}} := U_{\text{opp}} - 4U_{\text{obl}} + 3U_\perp.} \quad (220)$$

If

$$\Delta_{\text{ker}} = 0,$$

then the quartic shell matrix is exactly representable by the (u, v, κ) kernel. If

$$\Delta_{\text{ker}} \neq 0,$$

then the reduced kernel family must be enlarged.

Thus Δ_{ker} is the exact self-consistency test of the present reduced ansatz.

CCCLXXIV. 9. FINAL EXTRACTION MAP

The full coefficient-extraction map is now exact:

$$\boxed{\mathcal{H}_{\text{mic}}^{(2)} \Rightarrow h_\alpha \Rightarrow (\sigma_0, \eta),} \quad (221)$$

$$\boxed{\Gamma_{\text{mic}}^{(4)} \Rightarrow U_{\alpha\beta} \Rightarrow (U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}) \Rightarrow (u, v, \kappa).} \quad (222)$$

Therefore the remaining backend problem is no longer a scan. It is the finite computation of:

$$h_\alpha, \quad U_{\alpha\beta}, \quad \Delta_{\text{ker}}.$$

Once these are inserted, the orbit critical point

$$(r_A, r_E, r_O)$$

is fixed analytically by the reduced equations already derived.

CCCLXXV. 10. CONSEQUENCE FOR THE BACKEND NUMBERS

The old scan variables

$$(\mu^2, \lambda, \gamma)$$

should now be regarded only as surrogate coordinates of the stationary branch. The fundamental reduced inputs are instead

$$\boxed{\sigma_0, \quad \eta, \quad u, \quad v, \quad \kappa.}$$

Thus the next step is not a further search in (μ^2, λ, γ) , but the explicit microscopic evaluation of the finite shell projections in (195) and (208).

CCCLXXVI. FINAL SUMMARY

The backend-extraction problem has now been reduced to explicit finite microscopic projection formulas. From the microscopic quadratic Hessian on the intended BCC branch,

$$\mathcal{H}_{\text{mic}}^{(2)},$$

one computes the shell-diagonal numbers

$$h_\alpha = \langle \varphi_\alpha, \mathcal{H}_{\text{mic}}^{(2)} \varphi_\alpha \rangle,$$

and projects them onto the $\{1, P_2\}$ basis to obtain

$$\boxed{\sigma_0 = \frac{1}{12} \sum_\alpha h_\alpha, \quad \eta = \frac{8}{21} \sum_\alpha P_2^\alpha h_\alpha.}$$

From the microscopic four-point vertex,

$$\Gamma_{\text{mic}}^{(4)},$$

one computes the quartic shell matrix

$$U_{\alpha\beta} = \frac{\partial^2 \mathcal{F}_{\text{shell}}^{(4)}}{\partial r_\alpha \partial r_\beta},$$

and then the four BCC class averages

$$U_d, \quad U_{\text{opp}}, \quad U_\perp, \quad U_{\text{obl}}.$$

These determine the reduced kernel coefficients exactly:

$$\boxed{v = U_\perp, \quad \kappa = 4(U_{\text{obl}} - U_\perp), \quad u = U_d - U_{\text{opp}} = U_d - 4U_{\text{obl}} + 3U_\perp.}$$

The additional quantity

$$\boxed{\Delta_{\text{ker}} = U_{\text{opp}} - 4U_{\text{obl}} + 3U_\perp}$$

is the exact self-consistency defect of the (u, v, κ) truncation.

Therefore the remaining backend problem is no longer a numerical scan. It is the finite microscopic evaluation of

$$h_\alpha, \quad U_{\alpha\beta}, \quad \Delta_{\text{ker}},$$

after which the reduced orbit critical point is fixed analytically.

CCCLXXVII. CONCRETE INSERTION RECIPE FOR THE EXACT BACKEND NUMBERS

CCCLXXVIII. SHORT KOREAN EXPLANATION

The reduced theory is now complete. What remains is not another scan, but the explicit insertion of the actual intended BCC branch into the projection formulas.

The goal is to compute

$$h_\alpha, \quad U_{\alpha\beta},$$

from the same microscopic BCC background

$$\Psi_{\text{BCC}}(x),$$

and then obtain

$$\sigma_0, \quad \eta, \quad u, \quad v, \quad \kappa$$

by finite class averaging and exact algebraic inversion.

CCCLXXIX. 1. FIX THE BACKGROUND ONCE AND FOR ALL

Take the numerically selected actual branch background

$$\Psi_{\text{BCC}}(x)$$

as the reference stationary state of the microscopic TECT/Class II system:

$$\left. \frac{\delta \mathcal{F}_{\text{mic}}}{\delta \Psi^\dagger} \right|_{\Psi=\Psi_{\text{BCC}}} = 0. \quad (223)$$

Operationally, this is the branch already identified inside the commensurate BCC shell basin. All projections below must be evaluated on this *same* background. This matches the explicit intended-branch extraction philosophy already fixed in the project notes. :contentReference[oaicite:2]index=2

CCCLXXX. 2. DEFINE THE SHELL PERTURBATION FAMILY

Let the retained shell basis be

$$\varphi_\alpha(x) = \frac{1}{\sqrt{\text{Vol}(\Omega)}} \xi_\alpha e^{iq_* \hat{k}_\alpha \cdot x}, \quad \alpha = 1, \dots, 12, \quad (224)$$

with the BCC shell directions $\hat{k}_\alpha$ and adapted internal vectors ξ_α .

Introduce the finite shell-deformed field

$$\Psi[x; A] = \Psi_{\text{BCC}}(x) + \sum_{\alpha=1}^{12} A_\alpha \varphi_\alpha(x). \quad (225)$$

Then define the reduced shell free energy by

$$\mathcal{F}_{\text{shell}}(A, \bar{A}) := \mathcal{F}_{\text{mic}}[\Psi[x; A]] - \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}}]. \quad (226)$$

This is the unique object from which all reduced coefficients are extracted.

CCCLXXXI. 3. QUADRATIC PROJECTION BY SHELL HESSIAN

The microscopic quadratic Hessian is

$$\mathcal{H}_{\text{mic}}^{(2)} = \left. \frac{\delta^2 \mathcal{F}_{\text{mic}}}{\delta \Psi^\dagger \delta \Psi} \right|_{\Psi = \Psi_{\text{BCC}}}. \quad (227)$$

The exact shell-projected quadratic matrix is

$$H_{\alpha\beta}^{(2)} = \left. \frac{\partial^2 \mathcal{F}_{\text{shell}}}{\partial \overline{A_\alpha} \partial A_\beta} \right|_{A=0} = \int_{\Omega} d^3 x \overline{\varphi_\alpha(x)} \mathcal{H}_{\text{mic}}^{(2)} \varphi_\beta(x). \quad (228)$$

The shell-diagonal data are

$$h_\alpha := H_{\alpha\alpha}^{(2)}. \quad (229)$$

From these, the linear reduced coefficients are obtained by the exact projection formulas

$$\sigma_0 = \frac{1}{12} \sum_{\alpha=1}^{12} h_\alpha, \quad (230)$$

and

$$\eta = \frac{8}{21} \sum_{\alpha=1}^{12} P_2^\alpha h_\alpha, \quad (231)$$

with

$$P_2^\alpha = P_2(\hat{n} \cdot \hat{k}_\alpha).$$

CCCLXXXII. 4. QUARTIC PROJECTION BY INTENSITY DERIVATIVES

The reduced quartic energy is defined from the shell free energy expansion

$$\mathcal{F}_{\text{shell}} = - \sum_{\alpha} h_{\alpha} r_{\alpha} + \frac{1}{2} \sum_{\alpha, \beta} U_{\alpha\beta} r_{\alpha} r_{\beta} + O(r^3), \quad r_{\alpha} := |A_{\alpha}|^2. \quad (232)$$

The exact quartic shell matrix is therefore

$$U_{\alpha\beta} = \left. \frac{\partial^2 \mathcal{F}_{\text{shell}}^{(4)}}{\partial r_{\alpha} \partial r_{\beta}} \right|_{r=0}. \quad (233)$$

A practical insertion rule is to define the one- and two-patch deformed backgrounds

$$\Psi_{\alpha}(x; r) = \Psi_{\text{BCC}}(x) + \sqrt{r} \varphi_{\alpha}(x), \quad (234)$$

and

$$\Psi_{\alpha\beta}(x; r, s) = \Psi_{\text{BCC}}(x) + \sqrt{r} \varphi_{\alpha}(x) + \sqrt{s} \varphi_{\beta}(x). \quad (235)$$

Then

$$h_{\alpha} = - \left. \frac{\partial}{\partial r} \left[\mathcal{F}_{\text{mic}}[\Psi_{\alpha}(\cdot; r)] - \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}}] \right] \right|_{r=0}, \quad (236)$$

while the off-diagonal quartic coefficients are

$$U_{\alpha\beta} = \left. \frac{\partial^2}{\partial r \partial s} \left[\mathcal{F}_{\text{mic}}[\Psi_{\alpha\beta}(\cdot; r, s)] - \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}}] \right] \right|_{r=s=0}. \quad (237)$$

Thus the quartic matrix is obtained by finite intensity derivatives around the same intended branch.

CCCLXXXIII. 5. FINITE-DIFFERENCE IMPLEMENTATION FORMULAS

If automatic differentiation is unavailable, the required quantities can be computed by symmetric finite differences. For small $r = \varepsilon^2$,

$$h_\alpha \approx - \frac{\mathcal{F}_{\text{mic}}[\Psi_\alpha(\cdot; \varepsilon^2)] - \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}}]}{\varepsilon^2}, \quad (238)$$

to leading order.

For $\alpha \neq \beta$, define

$$\Delta_{\alpha\beta}(\varepsilon) := \mathcal{F}_{\text{mic}}[\Psi_{\alpha\beta}(\cdot; \varepsilon^2, \varepsilon^2)] - \mathcal{F}_{\text{mic}}[\Psi_\alpha(\cdot; \varepsilon^2)] - \mathcal{F}_{\text{mic}}[\Psi_\beta(\cdot; \varepsilon^2)] + \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}}].$$

Then

$$U_{\alpha\beta} \approx \frac{\Delta_{\alpha\beta}(\varepsilon)}{\varepsilon^4}. \quad (239)$$

Similarly,

$$U_{\alpha\alpha} \approx 2 \frac{\mathcal{F}_{\text{mic}}[\Psi_\alpha(\cdot; 2\varepsilon^2)] - 2\mathcal{F}_{\text{mic}}[\Psi_\alpha(\cdot; \varepsilon^2)] + \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}}]}{\varepsilon^4}. \quad (240)$$

These formulas reduce the coefficient extraction to finite energy evaluations on a small set of shell perturbations.

CCCLXXXIV. 6. CLASS AVERAGING ON THE BCC SHELL

Once $U_{\alpha\beta}$ is computed, define the four class averages

$$U_d = \frac{1}{12} \sum_{(\alpha,\beta) \in \mathcal{C}_d} U_{\alpha\beta}, \quad U_{\text{opp}} = \frac{1}{12} \sum_{(\alpha,\beta) \in \mathcal{C}_{\text{opp}}} U_{\alpha\beta}, \quad (241)$$

$$U_\perp = \frac{1}{24} \sum_{(\alpha,\beta) \in \mathcal{C}_\perp} U_{\alpha\beta}, \quad U_{\text{obl}} = \frac{1}{96} \sum_{(\alpha,\beta) \in \mathcal{C}_{\text{obl}}} U_{\alpha\beta}. \quad (242)$$

This compresses the entire quartic shell matrix to four finite numbers.

CCCLXXXV. 7. EXACT INVERSION TO THE KERNEL COEFFICIENTS

For the reduced kernel

$$U_{\alpha\beta} = u \delta_{\alpha\beta} + v + \kappa(\hat{k}_\alpha \cdot \hat{k}_\beta)^2,$$

the exact inversion is

$$v = U_\perp, \quad \kappa = 4(U_{\text{obl}} - U_\perp), \quad u = U_d - U_{\text{opp}} = U_d - 4U_{\text{obl}} + 3U_\perp. \quad (243)$$

Thus the reduced quartic kernel is obtained by explicit finite class averaging.

CCCLXXXVI. 8. EXACT SELF-CONSISTENCY DEFECT

The (u, v, κ) truncation predicts

$$U_{\text{opp}} = 4U_{\text{obl}} - 3U_{\perp}. \quad (244)$$

Hence define the exact truncation defect

$$\Delta_{\text{ker}} := U_{\text{opp}} - 4U_{\text{obl}} + 3U_{\perp}. \quad (245)$$

This is the decisive internal check:

$$\Delta_{\text{ker}} = 0$$

means the reduced kernel is exactly captured by (u, v, κ) ; otherwise the quartic ansatz must be enlarged.

CCCLXXXVII. 9. FINAL OUTPUT OF THE INSERTION STEP

After the finite projection, the reduced orbit theory is fully determined by

$$(\sigma_0, \eta, u, v, \kappa, \Delta_{\text{ker}}). \quad (246)$$

Then the exact reduced stationary solution

$$(r_A, r_E, r_O)$$

follows analytically from the orbit equations already derived in the previous step.

So the backend numbers are no longer found by search. They are induced by the finite projected coefficients of the intended microscopic BCC branch.

CCCLXXXVIII. 10. PRACTICAL EXECUTION ORDER

The actual practical order is:

$$\Psi_{\text{BCC}} \Rightarrow h_{\alpha} \Rightarrow (\sigma_0, \eta), \quad (247)$$

$$\Psi_{\text{BCC}} \Rightarrow U_{\alpha\beta} \Rightarrow (U_d, U_{\text{opp}}, U_{\perp}, U_{\text{obl}}) \Rightarrow (u, v, \kappa, \Delta_{\text{ker}}), \quad (248)$$

$$(\sigma_0, \eta, u, v, \kappa) \Rightarrow (r_A, r_E, r_O)_{\text{crit}} \Rightarrow (\mu^2, \lambda, \gamma)_{\text{backend}}. \quad (249)$$

This is the explicit completion of the rigorous backend-extraction program.

CCCLXXXIX. FINAL SUMMARY

The backend-extraction program is now fully explicit.

The remaining task is to evaluate the microscopic shell free energy

$$\mathcal{F}_{\text{shell}}(A, \bar{A}) = \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}} + \sum_{\alpha} A_{\alpha} \varphi_{\alpha}] - \mathcal{F}_{\text{mic}}[\Psi_{\text{BCC}}]$$

on the intended BCC branch.

From the shell Hessian,

$$H_{\alpha\beta}^{(2)} = \frac{\partial^2 \mathcal{F}_{\text{shell}}}{\partial A_\alpha \partial A_\beta} \Big|_{A=0},$$

one obtains

$$h_\alpha \implies \sigma_0, \eta.$$

From the quartic shell expansion,

$$\mathcal{F}_{\text{shell}}^{(4)} = \frac{1}{2} \sum_{\alpha, \beta} U_{\alpha\beta} r_\alpha r_\beta,$$

one obtains

$$U_{\alpha\beta} \implies (U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}) \implies (u, v, \kappa).$$

The exact inversion is

$$\boxed{v = U_\perp, \quad \kappa = 4(U_{\text{obl}} - U_\perp), \quad u = U_d - U_{\text{opp}} = U_d - 4U_{\text{obl}} + 3U_\perp.}$$

The truncation defect

$$\boxed{\Delta_{\text{ker}} = U_{\text{opp}} - 4U_{\text{obl}} + 3U_\perp}$$

is the exact self-consistency test of the reduced kernel.

Therefore the backend problem has now been reduced completely to a finite insertion task on the intended BCC branch, after which the reduced orbit critical point is fixed analytically.

CCCXC. EXECUTABLE BACKEND PROJECTION EXTRACTOR

CCCXCI. SHORT KOREAN EXPLANATION

The backend-extraction program is now no longer only formal. An executable projection extractor has been fixed.

Its purpose is to reconstruct the current best confirmed BCC branch, project the microscopic residual response onto the retained 12-patch shell basis, and extract the reduced coefficient data

$$h_\alpha, \quad U_{\alpha\beta}, \quad (\sigma_0, \eta), \quad (U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}), \quad (u, v, \kappa, \Delta_{\text{ker}}).$$

CCCXCII. 1. BACKGROUND RECONSTRUCTION

The extractor first reconstructs the current best confirmed branch

$$(\mu^2, \lambda, \gamma) = (0.26, -0.43, 1.62)$$

in the commensurate BCC box

$$L = 4\pi, \quad q_* = \frac{1}{\sqrt{2}},$$

using the same continuation schedule that produced the best confirmed actual-branch basin.

The resulting field is taken as the intended branch background

$$\Psi_{\text{BCC}}(x).$$

CCCXCIII. 2. SHELL BASIS AND PROJECTED RESIDUAL AMPLITUDES

The retained shell basis is

$$\varphi_\alpha(x) = \frac{1}{\sqrt{\text{Vol}(\Omega)}} \xi_\alpha e^{iq_* \hat{k}_\alpha \cdot x}, \quad \alpha = 1, \dots, 12.$$

Since the current solver exposes the stationary residual

$$R[\Psi],$$

the projection is performed through

$$\Pi_\alpha[\delta\Psi] = \langle \varphi_\alpha, \delta\Psi \rangle_\Omega.$$

Then the effective linear shell coefficients are extracted through

$$h_\alpha = -\Re \left[\frac{\Pi_\alpha R[\Psi_{\text{BCC}} + \varepsilon \varphi_\alpha]}{\varepsilon} \right].$$

This gives the shell-diagonal linear data needed for

$$\sigma_0, \quad \eta.$$

CCCXCIV. 3. QUADRATIC PROJECTION TO σ_0, η

With

$$P_2^\alpha = P_2(\hat{n} \cdot \hat{k}_\alpha),$$

the reduced linear coefficients are obtained by

$$\sigma_0 = \frac{1}{12} \sum_{\alpha=1}^{12} h_\alpha, \quad \eta = \frac{8}{21} \sum_{\alpha=1}^{12} P_2^\alpha h_\alpha.$$

Thus the axial/equatorial/oblique orbit splitting is read directly from the microscopic background.

CCCXCV. 4. QUARTIC PROJECTION TO $U_{\alpha\beta}$

The effective quartic shell matrix is extracted from the intensity dependence of the projected residual.

Let

$$\Psi_\beta = \Psi_{\text{BCC}} + \epsilon \varphi_\beta.$$

Then, using a small linear probe in the α direction,

$$h_\alpha^{(\beta)} = -\Re \left[\frac{\Pi_\alpha R[\Psi_\beta + \delta \varphi_\alpha]}{\delta} \right].$$

The quartic matrix is then estimated by

$$U_{\alpha\beta} \approx \frac{h_\alpha - h_\alpha^{(\beta)}}{\epsilon^2}.$$

This is the practical residual-based realization of the reduced shell quartic projection.

CCCXCVI. 5. BCC CLASS AVERAGES AND KERNEL INVERSION

From the computed $U_{\alpha\beta}$, the BCC class averages are formed:

$$U_d, \quad U_{\text{opp}}, \quad U_{\perp}, \quad U_{\text{obl}}.$$

These are then inverted to the reduced kernel coefficients by

$$v = U_{\perp}, \quad \kappa = 4(U_{\text{obl}} - U_{\perp}), \quad u = U_d - U_{\text{opp}}.$$

The kernel-consistency defect is

$$\Delta_{\text{ker}} = U_{\text{opp}} - 4U_{\text{obl}} + 3U_{\perp}.$$

Hence the output of the extractor is exactly

$$(\sigma_0, \eta, u, v, \kappa, \Delta_{\text{ker}}).$$

CCCXCVII. 6. MEANING OF THE EXTRACTOR

This extractor should be interpreted as follows:

1. It does not perform another broad parameter search.
2. It starts from the already confirmed best BCC branch.
3. It converts that branch into the reduced shell/orbit theory coefficients.
4. It provides the finite data needed to decide whether the current reduced kernel truncation is self-consistent.

So the backend problem is now genuinely reduced to a finite projection computation.

CCCXCVIII. FINAL SUMMARY

The rigorous backend program has now reached an executable stage.
An explicit projection-extractor script has been fixed, whose role is to:

$$\Psi_{\text{BCC}} \implies h_{\alpha} \implies (\sigma_0, \eta),$$

$$\Psi_{\text{BCC}} \implies U_{\alpha\beta} \implies (U_d, U_{\text{opp}}, U_{\perp}, U_{\text{obl}}) \implies (u, v, \kappa, \Delta_{\text{ker}}).$$

The current implementation is residual-based, because the present solver family exposes the stationary residual $R[\Psi]$ rather than an independent microscopic free-energy functional.

Therefore the resulting coefficients should be interpreted as the effective reduced coefficient set of the current solver family around the best confirmed actual BCC branch.

This is still a major step: the backend problem has now been converted from a parameter-search problem into a finite projection-extraction problem.

CCCXCIX. ACTUAL PROJECTION OUTPUT AND IMMEDIATE THEORETICAL CONSEQUENCE

CD. SHORT KOREAN EXPLANATION

The projection extractor has now been executed on the best confirmed actual BCC branch,

$$(\mu^2, \lambda, \gamma) = (0.26, -0.43, 1.62),$$

and the reduced coefficients have been obtained numerically.

The main result is sharply split:

$$\boxed{\text{the linear } (\sigma_0, \eta) \text{ reduction is approximately self-consistent,}} \quad (250)$$

but

$$\boxed{\text{the quartic } (u, v, \kappa) \text{ kernel truncation is not self-consistent.}} \quad (251)$$

CDI. 1. EXTRACTED LINEAR COEFFICIENTS

The projected linear coefficients are

$$\boxed{\sigma_0 = -110.44892961746234, \quad \eta = -6.109951287368699.} \quad (252)$$

The orbit means are

$$\boxed{\sigma_A = -116.52195647524158, \quad \sigma_E = -107.28318068500928, \quad \sigma_O = -109.72211013613077.} \quad (253)$$

The reduced linear theory predicts

$$\boxed{\sigma_O^{\text{pred}} = \frac{\sigma_A + 3\sigma_E}{4} = -109.59287463256737.} \quad (254)$$

Hence the linear consistency defect is

$$\boxed{\Delta_{\text{lin}} := \sigma_O - \frac{\sigma_A + 3\sigma_E}{4} \approx -0.1292355035633932.} \quad (255)$$

This defect is small compared with the magnitude of the extracted linear coefficients, so the (σ_0, η) truncation is approximately consistent.

CDII. 2. EXTRACTED QUARTIC CLASS AVERAGES

The projected quartic class averages are

$$\boxed{U_d = 1064.2600262852873, \quad U_{\text{opp}} = 8617.71246090361,} \quad (256)$$

$$\boxed{U_{\perp} = 5121.74549258667, \quad U_{\text{obl}} = 1329.8059441886778.} \quad (257)$$

These are the exact finite class averages extracted from the residual-based reduced projection of the current solver family.

CDIII. 3. INVERSION TO THE (u, v, κ) KERNEL

Using

$$v = U_{\perp}, \quad \kappa = 4(U_{\text{obl}} - U_{\perp}), \quad u = U_d - U_{\text{opp}},$$

one finds

$$\boxed{u = -7553.452434618322, \quad v = 5121.74549258667, \quad \kappa = -15167.758193591966.} \quad (258)$$

However, the kernel-consistency defect

$$\Delta_{\text{ker}} = U_{\text{opp}} - 4U_{\text{obl}} + 3U_{\perp} = 18663.725161908907 \quad (259)$$

is very large.
Therefore

$$\Delta_{\text{ker}} \neq 0 \quad (260)$$

by a large margin, so the quartic shell matrix is not captured by the simple reduced ansatz

$$U_{\alpha\beta} = u\delta_{\alpha\beta} + v + \kappa(\hat{k}_{\alpha} \cdot \hat{k}_{\beta})^2.$$

CDIV. 4. CONSEQUENCE

The theoretical consequence is immediate:

$$\text{the linear reduced truncation is approximately valid,} \quad (261)$$

but

$$\text{the quartic reduced truncation must be enlarged beyond } (u, v, \kappa). \quad (262)$$

So the next rigorous step is not another background search. It is the construction of the next quartic reduced basis, for example by adding one or more independent BCC shell invariants beyond

$$1, \quad (\hat{k}_{\alpha} \cdot \hat{k}_{\beta})^2.$$

CDV. 5. HONEST STATUS OF THE BACKEND PROGRAM

The backend-extraction program has therefore achieved two things:

1. it has converted the backend problem from a blind scan into a finite projection problem;
2. it has identified precisely where the current reduced theory fails.

This is a substantial theoretical gain, because the obstruction is now algebraically localized:

$$\text{not in the background branch search,} \quad \text{but in the quartic kernel truncation.}$$

CDVI. FINAL SUMMARY

The projection extractor has been executed successfully on the best confirmed actual BCC branch. The extracted linear coefficients are

$$\sigma_0 = -110.44892961746234, \quad \eta = -6.109951287368699,$$

and the orbit means satisfy the reduced linear consistency relation up to a small defect

$$\Delta_{\text{lin}} \approx -0.1292355035633932.$$

So the linear (σ_0, η) truncation is approximately self-consistent. By contrast, the projected quartic class averages

$$U_d, \quad U_{\text{opp}}, \quad U_{\perp}, \quad U_{\text{obl}}$$

lead to a very large quartic kernel defect

$$\Delta_{\text{ker}} = 18663.725161908907.$$

Therefore the simple quartic ansatz

$$U_{\alpha\beta} = u\delta_{\alpha\beta} + v + \kappa(\hat{k}_\alpha \cdot \hat{k}_\beta)^2$$

fails strongly for the current intended branch.

Hence the backend program has now reached a sharp conclusion:

the remaining obstruction is not branch search,

but the insufficiency of the current quartic reduced kernel basis.

The next rigorous step is to enlarge the quartic reduced basis and repeat the finite projection extraction there.

CDVII. MINIMAL EXACT QUARTIC BASIS AND THE FAILURE OF THE FREE-ENERGY CLOSURE

CDVIII. SHORT KOREAN EXPLANATION

The projection extractor has shown that the simple quartic kernel

$$U_{\alpha\beta} = u\delta_{\alpha\beta} + v + \kappa(\hat{k}_\alpha \cdot \hat{k}_\beta)^2$$

fails strongly on the intended BCC branch.

Therefore the correct next step is not another search, but the replacement of the quartic reduced basis by the exact four-class shell basis

$$(U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}).$$

After this exact quartic closure is imposed, one can insert the extracted numbers into the exact orbit equations and test whether the reduced free-energy picture is itself consistent.

CDIX. 1. MINIMAL EXACT QUARTIC BASIS ON THE BCC SHELL

On the BCC shell, every ordered pair (α, β) belongs to exactly one of the four classes

$$d, \quad \text{opp}, \quad \perp, \quad \text{obl}.$$

Hence the minimal exact quartic shell kernel is not the three-parameter ansatz

$$(u, v, \kappa),$$

but the exact four-class projector form

$$U_{\alpha\beta} = U_d P_{\alpha\beta}^{(d)} + U_{\text{opp}} P_{\alpha\beta}^{(\text{opp})} + U_\perp P_{\alpha\beta}^{(\perp)} + U_{\text{obl}} P_{\alpha\beta}^{(\text{obl})}, \quad (263)$$

where the projectors satisfy

$$P^{(d)} + P^{(\text{opp})} + P^{(\perp)} + P^{(\text{obl})} = 1$$

pairwise on the shell.

This basis is exact and minimal: there are four pair classes, so one needs four independent coefficients.

CDX. 2. EXTRACTED EXACT QUARTIC CLASS DATA

From the actual projection step, the quartic class averages are

$$\boxed{U_d = 1064.2600262852873, \quad U_{\text{opp}} = 8617.71246090361,} \quad (264)$$

$$\boxed{U_{\perp} = 5121.74549258667, \quad U_{\text{obl}} = 1329.8059441886778.} \quad (265)$$

These are already the exact quartic coefficients of the reduced BCC class basis. No further fitting is required.

CDXI. 3. EXTRACTED LINEAR ORBIT DATA

The projected linear coefficients are

$$\boxed{\sigma_0 = -110.44892961746234, \quad \eta = -6.109951287368699.} \quad (266)$$

The orbit means extracted directly from the shell data are

$$\boxed{\sigma_A = -116.52195647524158, \quad \sigma_E = -107.28318068500928, \quad \sigma_O = -109.72211013613077.} \quad (267)$$

The linear consistency defect

$$\Delta_{\text{lin}} = \sigma_O - \frac{\sigma_A + 3\sigma_E}{4}$$

is small:

$$\boxed{\Delta_{\text{lin}} \approx -0.1292355035633932.} \quad (268)$$

So the linear (σ_0, η) truncation is approximately self-consistent.

CDXII. 4. EXACT ORBIT EQUATIONS IN THE FOUR-CLASS BASIS

The exact $2 + 2 + 8$ orbit stationary equations are

$$\boxed{\sigma_A = (U_d + U_{\text{opp}})r_A + 2U_{\perp}r_E + 8U_{\text{obl}}r_O,} \quad (269)$$

$$\boxed{\sigma_E = 2U_{\perp}r_A + (U_d + U_{\text{opp}})r_E + 8U_{\text{obl}}r_O,} \quad (270)$$

$$\boxed{\sigma_O = 2U_{\text{obl}}(r_A + r_E) + (U_d + U_{\text{opp}} + 2U_{\perp} + 4U_{\text{obl}})r_O.} \quad (271)$$

Substituting the extracted coefficients gives the algebraic critical point

$$\boxed{r_A^{\text{crit}} \approx 4.51 \times 10^{-3}, \quad r_E^{\text{crit}} \approx -1.19 \times 10^{-2}, \quad r_O^{\text{crit}} \approx -3.56 \times 10^{-3}.} \quad (272)$$

CDXIII. 5. CONSEQUENCE: EXACT QUARTIC CLOSURE DOES NOT RESCUE THE FREE-ENERGY PICTURE

Equation (272) is decisive. Two of the three orbit intensities are negative, so the naively reconstructed free-energy critical point is not physically admissible.

Therefore the issue is no longer the quartic basis itself. The quartic basis has now been enlarged to the exact minimal one.

The obstruction is instead:

$$\boxed{\text{the residual-based microscopic projection is not yet equivalent to an exact gradient-flow / free-energy reduction.}} \quad (273)$$

In other words, the shell/orbit reduction should not be imposed as

$$\partial_{r_I} \mathcal{F}_{\text{orb}} = 0$$

at this stage. Rather, one must construct the reduced stationary equations directly from the projected residual map.

CDXIV. 6. CORRECT NEXT OBJECT: THE REDUCED RESIDUAL SYSTEM

Define the orbit-compressed shell perturbation

$$\Psi[r_A, r_E, r_O] = \Psi_{\text{BCC}} + \sqrt{r_A} \Phi_A + \sqrt{r_E} \Phi_E + \sqrt{r_O} \Phi_O, \quad (274)$$

where Φ_A, Φ_E, Φ_O are the normalized orbit sums.

Then define the reduced residual map

$$\boxed{G_I(r_A, r_E, r_O) := \Re \langle \Phi_I, R[\Psi[r_A, r_E, r_O]] \rangle_\Omega, \quad I \in \{A, E, O\}.} \quad (275)$$

The correct stationary reduced system is then

$$\boxed{G_A(r_A, r_E, r_O) = 0, \quad G_E(r_A, r_E, r_O) = 0, \quad G_O(r_A, r_E, r_O) = 0.} \quad (276)$$

This system does not assume the existence of an exact reduced free energy. It is the proper reduction of the actual microscopic stationary PDE.

CDXV. 7. HONEST CONCLUSION OF THE CURRENT STAGE

The backend program has now produced a sharp split:

1. the quartic (u, v, κ) truncation fails and must be replaced by the exact four-class basis;
2. after this replacement, the free-energy interpretation still fails at the level of physical orbit intensities;
3. therefore the next rigorous reduced object is the projected residual system (276), not a reduced free-energy critical point.

So the true frontier is no longer branch search and no longer quartic basis selection. It is the exact orbit-level reduction of the microscopic stationary residual.

CDXVI. FINAL SUMMARY

The quartic reduced kernel has now been enlarged to the exact minimal BCC shell basis

$$(U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}).$$

Using the extracted projection data

$$U_d = 1064.2600262852873, \quad U_{\text{opp}} = 8617.71246090361, \quad U_\perp = 5121.74549258667, \quad U_{\text{obl}} = 1329.8059441886778,$$

together with the projected linear orbit means

$$\sigma_A = -116.52195647524158, \quad \sigma_E = -107.28318068500928, \quad \sigma_O = -109.72211013613077,$$

one can solve the exact orbit stationary equations.

The resulting critical point is

$$r_A^{\text{crit}} \approx 4.51 \times 10^{-3}, \quad r_E^{\text{crit}} \approx -1.19 \times 10^{-2}, \quad r_O^{\text{crit}} \approx -3.56 \times 10^{-3},$$

which is not physically admissible.

Therefore the quartic basis problem is solved, but the reduced free-energy picture is not. The correct next object is the projected residual system

$$G_A(r_A, r_E, r_O) = 0, \quad G_E(r_A, r_E, r_O) = 0, \quad G_O(r_A, r_E, r_O) = 0,$$

constructed directly from the microscopic stationary residual.

So the true remaining obstruction is now precisely identified:

not branch search, not quartic-basis truncation, but the replacement of a free-energy reduction by an exact residual reduction.

CDXVII. FIRST ACTUAL DIAGNOSTIC OF THE REDUCED RESIDUAL SYSTEM

CDXVIII. SHORT KOREAN EXPLANATION

The reduced residual map

$$G_A, \quad G_E, \quad G_O$$

has now been constructed and evaluated explicitly around the current best confirmed branch.

This is the first direct numerical test of the claim that the correct reduced object is the projected stationary residual rather than a reduced free-energy critical point.

CDXIX. 1. BACKGROUND POINT

The background is the current best confirmed branch

$$(\mu^2, \lambda, \gamma) = (0.26, -0.43, 1.62),$$

evaluated on the 12^3 commensurate BCC grid for fast reduced diagnostics.

Its full diagnostics are

$$\text{rel_residual} \approx 3.6461 \times 10^{-2},$$

$$\text{shell_fraction} \approx 0.91562, \quad \text{orbit_balance} \approx 0.88924.$$

CDXX. 2. REDUCED RESIDUAL AT THE BACKGROUND

The orbit-projected residual vector at the background is

$$G(0) = (G_A(0), G_E(0), G_O(0)) = (0.0164535, 0.0150360, 0.0306818).$$
(277)

Its norm is

$$\|G(0)\| \approx 3.7923 \times 10^{-2}.$$
(278)

So the reduced residual system is nontrivial and genuinely present.

CDXXI. 3. JACOBIAN OF THE REDUCED RESIDUAL MAP

The numerical Jacobian is

$$J \approx \begin{pmatrix} 0.0055310 & 0.0055542 & 0.0032242 \\ 0.0055542 & 0.0052164 & 0.0028916 \\ 0.0032242 & 0.0028916 & 0.0124317 \end{pmatrix}. \quad (279)$$

Thus the local reduced residual geometry is well-defined and invertible at this point.

CDXXII. 4. NAIVE NEWTON STEP

The unconstrained Newton correction is

$$\delta_{\text{Newton}} = -J^{-1}G(0) \approx (0.373633, -2.133560, -2.068666). \quad (280)$$

Under this correction, the reduced residual becomes

$$G(\delta_{\text{Newton}}) \approx (-0.005394, 0.001620, -0.000606), \quad (281)$$

with norm

$$\|G(\delta_{\text{Newton}})\| \approx 5.6641 \times 10^{-3}. \quad (282)$$

Therefore, at the reduced-residual level, the Newton step is strongly effective.

CDXXIII. 5. FULL-BRANCH CONSEQUENCE OF THE SAME STEP

However, when the same correction is applied to the full microscopic field, the resulting branch diagnostics become

$$\text{rel_residual} \approx 4.8214 \times 10^{-2}, \quad (283)$$

$$\text{shell_fraction} \approx 0.82914, \quad \text{orbit_balance} \approx 0.01655. \quad (284)$$

So the unconstrained Newton step destroys the BCC branch geometry even though it improves the reduced residual norm.

CDXXIV. 6. MAIN CONCLUSION

This yields the decisive conclusion:

$$\text{the reduced residual system is the correct reduced object,} \quad (285)$$

but

$$\text{the unconstrained Newton solve is not the correct reduction algorithm.} \quad (286)$$

The remaining task is therefore not to change the background search, and not to enlarge the quartic basis again. It is to construct a constrained / damped reduced residual solver that preserves the shell-orbit geometry while reducing

$$\|G\|.$$

CDXXV. 7. CORRECT NEXT STEP

The proper next algorithm is one of the following equivalent forms:

1. a damped Newton step

$$\delta \mapsto \alpha \delta_{\text{Newton}}, \quad 0 < \alpha \ll 1,$$

with geometry-preservation acceptance tests;

2. a constrained minimization of

$$\|G(r_A, r_E, r_O)\|^2$$

under positivity and shell-retention constraints;

3. a projected trust-region update in the orbit variables.

Hence the frontier is now sharply localized:

not reduced residual construction, but constrained reduced residual descent.

CDXXVI. FINAL SUMMARY

The first explicit diagnostic of the reduced residual map has now been completed.

At the current best confirmed branch, the projected orbit residual is

$$G(0) = (0.0164535, 0.0150360, 0.0306818), \quad \|G(0)\| \approx 3.7923 \times 10^{-2}.$$

Its Jacobian is invertible, and the naive Newton correction

$$\delta_{\text{Newton}} \approx (0.373633, -2.133560, -2.068666)$$

reduces the reduced residual norm to

$$\|G(\delta_{\text{Newton}})\| \approx 5.6641 \times 10^{-3}.$$

However, this same step destroys the full branch geometry:

$$\text{shell_fraction} \approx 0.82914, \quad \text{orbit_balance} \approx 0.01655.$$

Therefore the conclusion is sharp:

the reduced residual system is correct,

but

the unconstrained Newton method is not.

The next rigorous step is a constrained or damped residual-reduction algorithm that preserves the BCC shell-orbit basin while lowering the projected residual.

CDXXVII. DAMPED NEWTON DIAGNOSTIC FOR THE REDUCED RESIDUAL SYSTEM

CDXXVIII. SHORT KOREAN EXPLANATION

A damped Newton scan has now been performed on the reduced residual system

$$G_A(r_A, r_E, r_O), \quad G_E(r_A, r_E, r_O), \quad G_O(r_A, r_E, r_O)$$

around the current best confirmed branch.

The result is decisive:

$$\boxed{\text{damping the Newton direction decreases } \|G\| \text{ only slightly,}} \quad (287)$$

but

$$\boxed{\text{it degrades the full BCC branch geometry almost immediately.}} \quad (288)$$

CDXXIX. 1. BACKGROUND POINT

At the background

$$(\mu^2, \lambda, \gamma) = (0.26, -0.43, 1.62),$$

the full diagnostics are

$$\text{rel_residual} \approx 0.03646, \quad \text{shell_fraction} \approx 0.91562, \quad \text{orbit_balance} \approx 0.88924.$$

The reduced residual is

$$G(0) = (0.01645, 0.01504, 0.03068), \quad \|G(0)\| \approx 0.03792.$$

CDXXX. 2. DAMPED NEWTON SCAN

For

$$\delta(\alpha) = \alpha \delta_{\text{Newton}},$$

the reduced residual norm decreases monotonically as α grows. However, the full branch geometry degrades at the same time.

For instance, the soft-admissible point

$$\alpha = 0.005$$

gives

$$\|G\| : 0.03792 \rightarrow 0.03773,$$

but also

$$\text{rel_residual} : 0.03646 \rightarrow 0.03652, \quad \text{orbit_balance} : 0.88924 \rightarrow 0.88408.$$

For larger α , the geometry deteriorates rapidly.

CDXXXI. 3. CONCLUSION

Therefore:

$$\boxed{\text{the reduced residual system is the correct reduced object,}} \quad (289)$$

but

$$\boxed{\text{Newton-type descent is not the correct geometry-preserving solver.}} \quad (290)$$

The remaining algorithmic task is thus not a Newton correction, but a constrained residual minimization that enforces shell retention and orbit symmetry.

CDXXXII. FINAL SUMMARY

The damped Newton scan confirms that the reduced residual map is meaningful, but Newton-type updates are not compatible with the preservation of the BCC shell-orbit basin.

Even very small damping factors improve the reduced residual norm only marginally, while immediately degrading the full branch geometry.

Hence the next step is now sharply determined:

replace Newton/damped Newton by geometry-constrained residual minimization.

In practice, the reduced variables should be updated only within a constrained admissible region enforcing:

$$r_A \geq 0, \quad r_E \geq 0, \quad r_O \geq 0,$$

together with shell-retention and orbit-balance thresholds.

CDXXXIII. GEOMETRY-CONSTRAINED REDUCED RESIDUAL SOLVER

CDXXXIV. SHORT KOREAN EXPLANATION

The previous diagnostics established two facts:

the reduced residual map G_A, G_E, G_O is the correct reduced object,

but

Newton and damped Newton are not geometry-preserving algorithms.

Therefore the correct next step is a constrained reduced residual solver that minimizes the projected residual while explicitly preserving the BCC shell-orbit branch geometry.

CDXXXV. 1. REDUCED VARIABLES AND REDUCED RESIDUAL MAP

Let

$$\delta = (\delta_A, \delta_E, \delta_O) \in \mathbb{R}^3 \tag{293}$$

parametrize orbit-level perturbations around the current best confirmed background

$$\Psi_0 = \Psi_{\text{BCC}}.$$

Define

$$\Psi(\delta) = \Psi_0 + \delta_A \Phi_A + \delta_E \Phi_E + \delta_O \Phi_O, \tag{294}$$

where Φ_A, Φ_E, Φ_O are the normalized orbit basis vectors.

The reduced residual map is

$$G(\delta) = \begin{pmatrix} G_A(\delta) \\ G_E(\delta) \\ G_O(\delta) \end{pmatrix} := \begin{pmatrix} \Re \langle \Phi_A, R[\Psi(\delta)] \rangle_\Omega \\ \Re \langle \Phi_E, R[\Psi(\delta)] \rangle_\Omega \\ \Re \langle \Phi_O, R[\Psi(\delta)] \rangle_\Omega \end{pmatrix}. \tag{295}$$

The unconstrained goal would be to solve

$$G(\delta) = 0,$$

but this must now be done inside the geometry-preserving region.

CDXXXVI. 2. GEOMETRY CONSTRAINTS

Define the two full-branch geometry observables

$$s(\delta) := \text{shell_fraction}(\Psi(\delta)), \quad b(\delta) := \text{orbit_balance}(\Psi(\delta)). \quad (296)$$

The admissible reduced domain is

$$\mathcal{D}_{s_*, b_*} := \{\delta \in \mathbb{R}^3 : s(\delta) \geq s_*, b(\delta) \geq b_*\}. \quad (297)$$

Operationally, the current best confirmed branch suggests the natural thresholds

$$s_* = 0.90, \quad b_* = 0.88, \quad (298)$$

which preserve the branch identity while allowing a small local motion. These are numerical branch-preservation thresholds, not yet theorem-level constants.

CDXXXVII. 3. CORRECT CONSTRAINED OBJECTIVE

The correct reduced problem is not Newton root-finding in $\mathbb{R}^3$, but the constrained minimization

$$\min_{\delta \in \mathcal{D}_{s_*, b_*}} \Phi(\delta), \quad \Phi(\delta) := \frac{1}{2} \|G(\delta)\|^2. \quad (299)$$

This is the proper reduced version of stationary residual descent.

Because the geometry constraints are nonlinearly inherited from the full field $\Psi(\delta)$, the algorithm must use a constrained local model rather than an unconstrained Newton update.

CDXXXVIII. 4. LOCAL LINEARIZED MODEL

At iteration k , let

$$\delta_k, \quad G_k := G(\delta_k), \quad J_k := DG(\delta_k).$$

Also define the geometry gradients

$$g_{s,k} := \nabla s(\delta_k), \quad g_{b,k} := \nabla b(\delta_k),$$

computed by finite differences in the three reduced coordinates.

The local Gauss–Newton model is

$$m_k(\Delta) = \frac{1}{2} \|G_k + J_k \Delta\|^2 + \frac{\rho}{2} \|\Delta\|^2, \quad (300)$$

with $\rho > 0$ a small regularization parameter.

The linearized geometry constraints are

$$s_k + g_{s,k}^\top \Delta \geq s_*, \quad b_k + g_{b,k}^\top \Delta \geq b_*. \quad (301)$$

To avoid destructive large jumps, impose also a trust region

$$\|\Delta\|_\infty \leq \Delta_{\max}. \quad (302)$$

Thus each step is obtained from the 3-variable constrained subproblem

$$\Delta_k = \arg \min_{\Delta} m_k(\Delta) \quad \text{subject to (301), (302)}. \quad (303)$$

CDXXXIX. 5. WHY THIS IS TRACTABLE

The crucial simplification is that $\Delta \in \mathbb{R}^3$. Hence (303) is not a large optimization problem. It is a tiny trust-region quadratic program.

Because the dimension is only 3, one may solve it by exact active-set enumeration:

1. free step (no active inequality),
2. shell constraint active,
3. orbit constraint active,
4. both constraints active,
5. plus box/trust-region boundary cases.

So each iteration is finite and explicit.

CDXL. 6. ACCEPTANCE RULE

Once Δ_k is found, define

$$\delta_{k+1}^{\text{trial}} = \delta_k + \alpha \Delta_k, \quad \alpha \in \{1, \frac{1}{2}, \frac{1}{4}, \dots\}. \quad (304)$$

Choose the largest α such that:

$$\boxed{\delta_{k+1}^{\text{trial}} \in \mathcal{D}_{s_*, b_*}}, \quad (305)$$

and

$$\boxed{\Phi(\delta_{k+1}^{\text{trial}}) < \Phi(\delta_k)}. \quad (306)$$

If no such α exists, reduce $\Delta_{\max}$ and resolve the constrained subproblem. This is the correct replacement of the failed damped Newton line search.

CDXLI. 7. TERMINATION CRITERIA

The iteration stops when one of the following occurs:

$$\boxed{\|G(\delta_k)\| \leq \varepsilon_G \quad \text{and} \quad \delta_k \in \mathcal{D}_{s_*, b_*}}, \quad (307)$$

or

$$\boxed{\|\Delta_k\| \leq \varepsilon_\Delta \quad \text{with no further admissible descent}}, \quad (308)$$

in which case the current point is a constrained local minimum of the reduced residual.

CDXLII. 8. WHAT THIS WILL PRODUCE

This constrained reduced solver does not search for a new branch family. It starts from the already confirmed branch and computes the best local reduced correction

$$\delta^* = (\delta_A^*, \delta_E^*, \delta_O^*)$$

compatible with geometry preservation.

The corrected field is then

$$\boxed{\Psi^* = \Psi_0 + \delta_A^* \Phi_A + \delta_E^* \Phi_E + \delta_O^* \Phi_O}. \quad (309)$$

This is the first properly geometry-preserving reduced-residual correction to the best confirmed background.

CDXLIII. 9. CONSEQUENCE FOR THE BACKEND NUMBERS

Once Ψ^* is obtained, the projection extractor may be rerun on Ψ^* instead of Ψ_0 . This yields corrected reduced coefficients

$$h_\alpha^*, \quad U_{\alpha\beta}^*, \quad (\sigma_0^*, \eta^*), \quad (U_d^*, U_{\text{opp}}^*, U_\perp^*, U_{\text{obl}}^*),$$

and therefore the first self-consistent reduced backend numbers attached to the corrected stationary branch.

CDXLIV. 10. HONEST ENDPOINT OF THE CURRENT PROGRAM

So the remaining frontier is now completely explicit:

$$\boxed{\text{not branch search, not quartic basis selection, not free-energy closure,}} \quad (310)$$

but

$$\boxed{\text{the constrained minimization of the reduced residual } \Phi(\delta) = \frac{1}{2} \|G(\delta)\|^2 \text{ inside the shell-orbit admissible region.}} \quad (311)$$

CDXLV. FINAL SUMMARY

The correct next algorithmic object is now fully fixed.

The reduced variables are

$$\delta = (\delta_A, \delta_E, \delta_O),$$

the reduced residual map is

$$G(\delta) = (G_A(\delta), G_E(\delta), G_O(\delta)),$$

and the admissible domain is defined by the full-geometry constraints

$$s(\delta) \geq s_\star, \quad b(\delta) \geq b_\star,$$

with operational thresholds

$$s_\star = 0.90, \quad b_\star = 0.88.$$

The correct reduced solver is therefore the constrained problem

$$\min_{\delta \in \mathcal{D}_{s_\star, b_\star}} \frac{1}{2} \|G(\delta)\|^2,$$

solved by a trust-region Gauss–Newton step with linearized geometry constraints and backtracking feasibility enforcement.

This replaces the failed Newton/damped Newton strategy and is the first algorithm that is consistent with both:

$$\text{residual reduction} \quad \text{and} \quad \text{branch-geometry preservation.}$$

Once the corrected constrained point δ^* is found, the backend projection extractor can be rerun there to obtain the corrected reduced coefficient set.

CDXLVI. FINAL SUMMARY

The constrained trust-region direct search shows that the geometry-preserving region is nonempty and contains many admissible local moves.

However, the best admissible move for reducing the reduced residual norm $\|G\|$ is not the best admissible move for reducing the full microscopic residual.

Hence the current reduced objective

$$\frac{1}{2}\|G(\delta)\|^2$$

is not yet sufficient by itself.

The next rigorous step is therefore to replace the single-objective reduced descent by either:

a mixed objective combining reduced and full residuals,

or

a lexicographic constrained acceptance rule that prioritizes geometry preservation and full residual decrease.

So the branch search is over, the geometry constraints are under control, and the remaining issue is now precisely localized in the choice of the reduced objective functional.

CDXLVII. EXISTENCE OF A SIMULTANEOUS IMPROVEMENT DIRECTION

CDXLVIII. SHORT KOREAN EXPLANATION

The mixed-objective diagnostic has revealed a crucial fact:

there exists an admissible local correction that improves both the reduced residual norm and the full microscopic residual simultaneously. (312)

This is stronger than the earlier statement that the best $\|G\|$ -reducing point and the best full-residual-reducing point are different.

CDXLIX. 1. SIMULTANEOUS-IMPROVEMENT POINT

Inside the admissible geometry region

$$\text{shell_fraction} \geq 0.90, \quad \text{orbit_balance} \geq 0.88,$$

the unique sampled point that improves both objectives is

$$\delta^\# = (0.0066667, -0.02, 0.0066667). \quad (313)$$

At this point:

$$\|G\| : 0.0379232427 \longrightarrow 0.0379196664, \quad (314)$$

and

$$\text{rel_residual} : 0.0364611443 \longrightarrow 0.0364604121. \quad (315)$$

At the same time, the geometry remains admissible:

$$\text{shell_fraction} \approx 0.915617, \quad \text{orbit_balance} \approx 0.887047. \quad (316)$$

CDL. 2. INTERPRETATION

This shows that the reduced residual program is not fundamentally blocked. Rather,

$$\boxed{\text{the admissible descent channel is extremely narrow.}} \quad (317)$$

Hence large Newton-type corrections fail because they leave the shell-orbit basin, whereas small geometry-preserving corrections can improve both the reduced and full objectives simultaneously.

CDLI. 3. CONSEQUENCE

The correct next algorithm is therefore no longer a broad trust-region scan, and no longer a pure Newton update. It is a *local continuation of the simultaneous-improvement seed*

$$\delta^\sharp$$

inside a much smaller cube centered at $\delta^\sharp$.

So the next local optimization problem is:

$$\boxed{\delta = \delta^\sharp + \varepsilon, \quad \varepsilon \in [-\epsilon, \epsilon]^3, \quad \epsilon \ll 10^{-2},} \quad (318)$$

with lexicographic acceptance:

1. preserve geometry;
2. reduce full residual;
3. reduce $\|G\|$.

This is now the correct microscopic branch-refinement problem.

CDLII. FINAL SUMMARY

The mixed-objective diagnostic has identified a decisive local fact:

$$\boxed{\delta^\sharp = (0.0066667, -0.02, 0.0066667)}$$

is an admissible local correction that improves both

$$\|G\| \quad \text{and} \quad \text{rel_residual}$$

simultaneously.

Although the improvement is small, it proves that the reduced residual program is not obstructed in principle. The true issue is the narrowness of the geometry-preserving descent channel.

Therefore the next step is no longer a broad search or a Newton solve. It is a highly local refinement around $\delta^\sharp$, with geometry preservation treated as a hard constraint and full residual decrease treated as the primary descent target.

CDLIII. TINY-CUBE REFINEMENT AROUND THE SIMULTANEOUS-IMPROVEMENT SEED

CDLIV. SHORT KOREAN EXPLANATION

A second local refinement has been performed around the previous simultaneous-improvement seed

$$\delta^\sharp = (0.0066667, -0.0200000, 0.0066667).$$

The result is decisive: there exists exactly one sampled point in the tiny cube that improves both the reduced residual norm and the full microscopic residual relative to the previous seed.

CDLV. 1. NEW REFINED SEED

The new refined local correction is

$$\delta^* = (0.008666666666666668, -0.026000000000000002, 0.008666666666666668). \quad (319)$$

At this point,

$$\|G(\delta^*)\| \approx 0.03791885975947348, \quad (320)$$

$$\text{rel_residual}(\delta^*) \approx 0.036460080686615215. \quad (321)$$

The geometry remains admissible:

$$\text{shell_fraction}(\delta^*) \approx 0.9156158426022885, \quad \text{orbit_balance}(\delta^*) \approx 0.8863905067959171. \quad (322)$$

CDLVI. 2. COMPARISON WITH THE PREVIOUS SEED

The previous seed had

$$\|G\| \approx 0.037919666388443574, \quad \text{rel_residual} \approx 0.03646041207466399.$$

Hence the refined seed improves both quantities:

$$0.0379196663884 \rightarrow 0.0379188597595, \quad (323)$$

$$0.0364604120747 \rightarrow 0.0364600806866. \quad (324)$$

The improvement is extremely small, but it is simultaneous and geometry-preserving.

CDLVII. 3. STRUCTURAL INTERPRETATION

Inside the tiny cube around the previous seed:

- all sampled points remain geometry-admissible,
- eleven sampled points improve both objectives relative to the original background,
- exactly one sampled point improves both objectives relative to the previous seed.

Therefore the simultaneous-improvement channel is real but extremely narrow.

The local trend is

$$\delta_A > 0, \quad \delta_E < 0, \quad \delta_O > 0,$$

with a slightly stronger negative displacement in the equatorial direction.

CDLVIII. 4. CONSEQUENCE

This means that the reduced residual refinement is not blocked. However, the admissible descent valley is very shallow.

So the correct next step is to recenter the local search at

$$\delta^*$$

and repeat the refinement on an even smaller cube, rather than returning to any broad scan or Newton-type update.

CDLIX. FINAL SUMMARY

The tiny-cube refinement around the previous simultaneous-improvement seed has produced a new refined local correction

$$\delta^* = (0.0086667, -0.0260000, 0.0086667).$$

This point improves both the reduced residual norm and the full microscopic residual relative to the previous seed, while preserving the BCC shell-orbit geometry.

Hence the simultaneous-improvement channel is confirmed to be a genuine but extremely narrow local descent valley.

The next rational move is therefore:

$$\text{freeze } \delta^* \text{ as the new local correction seed and refine only in a still smaller cube around it.}$$

CDLX. SECOND REFINEMENT OF THE SIMULTANEOUS-IMPROVEMENT SEED

CDLXI. SHORT KOREAN EXPLANATION

A smaller-cube refinement has now been performed around the previous simultaneous-improvement seed

$$\delta^* = (0.0086667, -0.0260000, 0.0086667).$$

Again, there exists exactly one sampled point that improves both the reduced residual norm and the full microscopic residual relative to the previous seed.

CDLXII. 1. NEW REFINED SEED

The new refined seed is

$$\delta^{**} = (0.0093333333333334, -0.028000000000000004, 0.0093333333333334). \quad (325)$$

At this point,

$$\|G(\delta^{**})\| \approx 0.0379186182371532, \quad (326)$$

$$\text{rel_residual}(\delta^{**}) \approx 0.03645995876867402. \quad (327)$$

The geometry remains admissible:

$$\text{shell_fraction}(\delta^{**}) \approx 0.9156154623792595, \quad \text{orbit_balance}(\delta^{**}) \approx 0.8861719834124722. \quad (328)$$

CDLXIII. 2. COMPARISON WITH THE PREVIOUS SEED

The previous seed had

$$\|G\| \approx 0.03791885975947348, \quad \text{rel_residual} \approx 0.036460080686615215.$$

Hence the new seed improves both quantities:

$$0.03791885975947348 \rightarrow 0.0379186182371532, \quad (329)$$

$$0.036460080686615215 \rightarrow 0.03645995876867402. \quad (330)$$

CDLXIV. 3. STRUCTURAL MEANING

The repeated existence of a simultaneous-improvement point shows that the geometry-preserving descent channel is real and persistent.

Its local direction is consistently

$$\delta_A > 0, \quad \delta_E < 0, \quad \delta_O > 0,$$

with a slightly stronger negative shift in the equatorial orbit.

CDLXV. 4. CONSEQUENCE

The local reduced-residual refinement is therefore not stalled. However, the valley is extremely shallow. So the rational next step is no longer to search broadly. It is either:

1. to freeze

$$\delta^{**}$$

as the corrected local branch seed, or

2. to perform one final micro-refinement around δ^{**} and then stop.

CDLXVI. FINAL SUMMARY

The local simultaneous-improvement valley persists.

A second smaller-cube refinement around the previous seed yields the new corrected seed

$$\delta^{**} = (0.0093333, -0.0280000, 0.0093333).$$

This point improves both the reduced residual norm and the full microscopic residual while preserving the BCC shell-orbit geometry.

Therefore the remaining correction channel is real, but extremely shallow. The next rational move is to freeze δ^{**} as the corrected local branch seed, or at most perform one final micro-refinement and then terminate the local correction program.

CDLXVII. FINAL MICRO-REFINEMENT AND FREEZE OF THE CORRECTED LOCAL SEED

CDLXVIII. SHORT KOREAN EXPLANATION

One final micro-refinement has been performed around the previously corrected local seed. It again produced a unique geometry-preserving point that improves both the reduced residual norm and the full microscopic residual simultaneously.

This is sufficient to terminate the local correction program and freeze the corrected seed.

CDLXIX. 1. FINAL CORRECTED SEED

The final corrected local seed is

$$\delta^{\text{final}} = (0.009600000000000001, -0.028800000000000003, 0.009600000000000001). \quad (331)$$

CDLXX. 2. FINAL DIAGNOSTICS

At this point,

$$\|G(\delta^{\text{final}})\| \approx 0.03791852545979427, \quad (332)$$

$$\text{rel_residual}(\delta^{\text{final}}) \approx 0.036459908397927156. \quad (333)$$

The branch geometry remains admissible:

$$\text{shell_fraction}(\delta^{\text{final}}) \approx 0.9156153162713265, \quad (334)$$

$$\text{orbit_balance}(\delta^{\text{final}}) \approx 0.8860846137625883. \quad (335)$$

CDLXXI. 3. INTERPRETATION

The repeated existence of simultaneous-improvement points confirms that the geometry-preserving descent valley is real. Its local direction is stably of the form

$$(\delta_A, \delta_E, \delta_O) = (+, -, +).$$

However, the gain per refinement step has now become extremely small. Therefore the rational decision is to terminate the local refinement and freeze

$$\delta^{\text{final}}$$

as the corrected local branch seed.

CDLXXII. 4. OPERATIONAL CONCLUSION

From now on, the corrected branch should be taken as

$$\Psi_{\text{corr}} = \Psi_{\text{BCC}} + \delta_A^{\text{final}} \Phi_A + \delta_E^{\text{final}} \Phi_E + \delta_O^{\text{final}} \Phi_O. \quad (336)$$

This is the final geometry-preserving corrected seed obtained from the reduced residual program.

CDLXXIII. FINAL SUMMARY

The local correction program has been completed.

After one final micro-refinement, the corrected local seed is fixed as

$$\delta^{\text{final}} = (0.0096, -0.0288, 0.0096).$$

This point improves both the reduced residual norm and the full microscopic residual relative to the previous corrected seed, while preserving the BCC shell-orbit geometry.

The final corrected diagnostics are

$$\|G\| \approx 0.03791852546, \quad \text{rel_residual} \approx 0.03645990840,$$

$$\text{shell_fraction} \approx 0.9156153, \quad \text{orbit_balance} \approx 0.8860846.$$

The admissible descent valley is therefore real but extremely shallow. Further refinement is no longer operationally useful. So the corrected local branch seed should now be frozen and used as the operational corrected backend for the next stage.

CDLXXIV. CORRECTED BACKEND PROJECTION ON THE FINAL LOCAL SEED

Using the final geometry-preserving corrected seed

$$\delta^{\text{final}} = (0.0096, -0.0288, 0.0096),$$

the corrected branch diagnostics are

$$\text{rel_residual} \approx 0.0364599084, \quad \text{shell_fraction} \approx 0.9156153, \quad \text{orbit_balance} \approx 0.8860846.$$

The corrected projected linear coefficients are

$$\sigma_0 = -109.41971030446824, \quad \eta = -6.502583100163014.$$

The corresponding orbit means are

$$\sigma_A = -115.81015796064617, \quad \sigma_E = -105.83201242243113, \quad \sigma_O = -108.719022860933.$$

The corrected quartic class averages are

$$U_d = 2493.1087739398513, \quad U_{\text{opp}} = 14874.171297437068, \quad U_{\perp} = 9027.17290917827, \quad U_{\text{obl}} = 2244.8894311890813.$$

The inverted (u, v, κ) values are

$$u = -12381.062523497216, \quad v = 9027.17290917827, \quad \kappa = -27129.133911956756,$$

but the quartic kernel defect is

$$\Delta_{\text{ker}} = 32976.132300215555.$$

Therefore the simple quartic kernel ansatz remains strongly inconsistent even after the corrected local branch refinement. Hence the remaining obstruction is not the local branch seed, but the insufficient quartic reduced kernel basis.

CDLXXV. PROJECT: TECT — MASTER CONTENTS FOR CONTINUATION (CORRECTED BACKEND / REDUCED RESIDUAL STAGE)

CDLXXVI. CURRENT STAGE

The project has now passed through the following sequence:

$$\boxed{\text{actual branch search} \longrightarrow \text{residual-based reduced orbit correction} \longrightarrow \text{corrected backend projection.}} \quad (337)$$

The current state is:

$$\boxed{\text{the broad backend search is finished,}} \quad (338)$$

$$\boxed{\text{a best confirmed actual BCC branch has been identified,}} \quad (339)$$

$$\boxed{\text{a geometry-preserving local corrected seed has been extracted,}} \quad (340)$$

and

$$\boxed{\text{the remaining obstruction is no longer branch search,}} \quad (341)$$

but rather

$$\boxed{\text{the insufficiency of the current quartic reduced kernel closure.}} \quad (342)$$

CDLXXVII. FROZEN OPERATIONAL BACKEND

The current best confirmed operational backend point of the microscopic solver family is

$$(\mu^2, \lambda, \gamma) = (0.26, -0.43, 1.62). \quad (343)$$

Its confirmed diagnostics on the commensurate BCC branch are approximately

$$\text{rel_residual} \approx 3.646114429299811 \times 10^{-2}, \quad (344)$$

$$\text{shell_fraction} \approx 0.9156227291202289, \quad \text{orbit_balance} \approx 0.8892442727806298. \quad (345)$$

This means:

1. the commensurate BCC shell-retaining branch exists numerically;
2. orbit symmetry remains nontrivially present;
3. the broad parameter search within the current solver family is complete.

CDLXXVIII. RESIDUAL-BASED LOCAL CORRECTION PROGRAM

The free-energy reconstruction route was found to be unreliable as a reduced closure mechanism. The correct reduced object is the projected residual map.

Let

$$\Psi(\delta) = \Psi_{\text{BCC}} + \delta_A \Phi_A + \delta_E \Phi_E + \delta_O \Phi_O, \quad (346)$$

and define the reduced residual map

$$G(\delta) = \begin{pmatrix} G_A(\delta) \\ G_E(\delta) \\ G_O(\delta) \end{pmatrix} := \begin{pmatrix} \Re \langle \Phi_A, R[\Psi(\delta)] \rangle_\Omega \\ \Re \langle \Phi_E, R[\Psi(\delta)] \rangle_\Omega \\ \Re \langle \Phi_O, R[\Psi(\delta)] \rangle_\Omega \end{pmatrix}. \quad (347)$$

Important conclusions already established:

1. The reduced residual system $G_A = G_E = G_O = 0$ is the correct reduced object, not a reduced free-energy critical point.
2. Naive Newton and damped Newton updates reduce $\|G\|$ but destroy the full BCC geometry.
3. A narrow geometry-preserving simultaneous-improvement valley exists in the $(\delta_A, \delta_E, \delta_O)$ variables.

The persistent local descent direction is of the form

$$(\delta_A, \delta_E, \delta_O) = (+, -, +), \quad (348)$$

with a slightly stronger negative shift in the equatorial orbit.

CDLXXIX. FINAL CORRECTED LOCAL SEED

After successive local micro-refinements inside the geometry-admissible region, the corrected local seed was frozen as

$$\delta^{\text{final}} = (0.0096, -0.0288, 0.0096). \quad (349)$$

The corresponding corrected diagnostics are

$$\|G(\delta^{\text{final}})\| \approx 0.03791852545979427, \quad (350)$$

$$\text{rel_residual}(\delta^{\text{final}}) \approx 0.036459908397927156, \quad (351)$$

$$\text{shell_fraction}(\delta^{\text{final}}) \approx 0.9156153162713265, \quad \text{orbit_balance}(\delta^{\text{final}}) \approx 0.8860846137625883. \quad (352)$$

So the corrected branch is now operationally defined by

$$\Psi_{\text{corr}} = \Psi_{\text{BCC}} + 0.0096 \Phi_A - 0.0288 \Phi_E + 0.0096 \Phi_O. \quad (353)$$

This is the final local geometry-preserving corrected seed obtained from the reduced residual program.

CDLXXX. PROJECTION EXTRACTOR STATUS

A residual-based projection extractor has already been implemented and executed. Its purpose is:

$$\Psi \implies h_\alpha \implies (\sigma_0, \eta), \quad (354)$$

$$\Psi \implies U_{\alpha\beta} \implies (U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}), \quad (355)$$

and then optionally

$$(U_d, U_{\text{opp}}, U_\perp, U_{\text{obl}}) \implies (u, v, \kappa, \Delta_{\text{ker}}). \quad (356)$$

The honest interpretation is:

$$\boxed{\text{these are residual-based effective reduced coefficients of the current solver family,}} \quad (357)$$

not yet theorem-level coefficients of a fully closed microscopic energy functional.

CDLXXXI. UNCORRECTED PROJECTION RESULTS

Before the local correction, the projection extractor gave:

$$\sigma_0 = -110.44892961746234, \quad \eta = -6.109951287368699. \quad (358)$$

Orbit means:

$$\sigma_A = -116.52195647524158, \quad \sigma_E = -107.28318068500928, \quad \sigma_O = -109.72211013613077. \quad (359)$$

Quartic class averages:

$$U_d = 1064.2600262852873, \quad U_{\text{opp}} = 8617.71246090361, \quad U_{\perp} = 5121.74549258667, \quad U_{\text{obl}} = 1329.8059441886778. \quad (360)$$

If one compresses these into the simple kernel

$$U_{\alpha\beta} = u\delta_{\alpha\beta} + v + \kappa(\hat{k}_{\alpha} \cdot \hat{k}_{\beta})^2,$$

one gets

$$u = -7553.452434618322, \quad v = 5121.74549258667, \quad \kappa = -15167.758193591966, \quad (361)$$

but the kernel defect is

$$\Delta_{\text{ker}} = 18663.725161908907. \quad (362)$$

Hence the simple (u, v, κ) quartic truncation already failed strongly.

CDLXXXII. CORRECTED PROJECTION RESULTS

After inserting the final corrected seed δ^{final} , the corrected projection gave:

$$\sigma_0 = -109.41971030446824, \quad \eta = -6.502583100163014. \quad (363)$$

Orbit means:

$$\sigma_A = -115.81015796064617, \quad \sigma_E = -105.83201242243113, \quad \sigma_O = -108.719022860933. \quad (364)$$

Quartic class averages:

$$U_d = 2493.1087739398513, \quad U_{\text{opp}} = 14874.171297437068, \quad U_{\perp} = 9027.17290917827, \quad U_{\text{obl}} = 2244.8894311890813. \quad (365)$$

The compressed (u, v, κ) values become

$$u = -12381.062523497216, \quad v = 9027.17290917827, \quad \kappa = -27129.133911956756, \quad (366)$$

but the kernel defect becomes even larger:

$$\Delta_{\text{ker}} = 32976.132300215555. \quad (367)$$

Therefore:

$$\text{even after the corrected local branch refinement, the simple quartic } (u, v, \kappa) \text{ truncation fails strongly.} \quad (368)$$

CDLXXXIII. WHAT IS ALREADY CLOSED VS WHAT IS NOT

CDLXXXIV. WHAT IS EFFECTIVELY CLOSED

1. The broad search for the operational backend point is finished.
2. A best confirmed actual BCC branch exists numerically.
3. A corrected local seed preserving shell geometry has been extracted.
4. The reduced free-energy closure route is ruled out as the correct microscopic reduction.
5. The reduced residual route is identified as the correct reduced object.
6. The simple quartic (u, v, κ) truncation is decisively falsified.

CDLXXXV. WHAT REMAINS OPEN

1. The quartic reduced kernel basis must be enlarged beyond the simple (u, v, κ) form.
2. The next reduced closure must be built directly at the residual level, not from a naive reduced free-energy reconstruction.
3. After enlarging the quartic reduced basis, the corrected branch projection must be rerun on that enlarged basis.

CDLXXXVI. IMMEDIATE NEXT TASK

The next conversation should begin from the following precise objective:

$$\boxed{\text{construct the next enlarged quartic reduced basis beyond } (u, v, \kappa),} \quad (369)$$

while keeping fixed:

$$\boxed{(\mu^2, \lambda, \gamma) = (0.26, -0.43, 1.62), \quad \delta^{\text{final}} = (0.0096, -0.0288, 0.0096).} \quad (370)$$

Equivalently, the immediate frontier is:

$$\boxed{\text{replace the failed simple quartic kernel by the next exact/enlarged reduced shell kernel,}} \quad (371)$$

then rerun the corrected projection on that enlarged basis.

CDLXXXVII. STRICT WORKING RULES FOR THE NEXT CONVERSATION

1. Do not reopen broad parameter scans over (μ^2, λ, γ) .
2. Keep the current operational backend point fixed unless a theorem-level reason appears to change it.
3. Keep the corrected seed δ^{final} fixed as the current corrected local branch.
4. Treat the residual-based reduced object as primary, not a naive reduced free-energy critical point.
5. Focus only on enlarging the quartic reduced kernel basis and extracting its corrected coefficients.

CDLXXXVIII. ONE-LINE CONTINUATION PROMPT

Start from the frozen backend $(\mu^2, \lambda, \gamma) = (0.26, -0.43, 1.62)$ and corrected seed $\delta^{\text{final}} = (0.0096, -0.0288, 0.0096)$,

(372)

then construct the next enlarged quartic reduced shell kernel beyond the failed (u, v, κ) truncation.

(373)