



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

Search For Displaced Photons Using Timing.

Tambe E. Norbert¹ Shih-Chuan Kao¹ Yuichi Kubota¹

¹University Of Minnesota

**Long-Lived Meeting,
September 26, 2014**



Outline



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

- 1 Motivation
- 2 Production and Decay
- 3 Dataset and Trigger
- 4 Event Selection
 - Pre-Event Selection
 - Event Cleaning
 - Final Event Selection
- 5 Background Estimation
- 6 Systematics
- 7 Limits
- 8 Remaining Completion

How are we measuring the neutralino Lifetime?

- 1 Calculate time from distance travelled by Neutralino before it decays

Definition (Distance Travelled)

$$L = c\tau \cdot \gamma\beta \quad \rightarrow \quad c\tau = \frac{|\Delta \vec{r}|}{\gamma\beta}$$

- 2 Extract time directly from MC

Definition (MC Time)

$$c\tau = \frac{ct}{\gamma}$$



Production and Decay



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

Production and Decay Diagrams



Dataset and Trigger



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

Datasets and HLT Triggers used.

Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion





Figure : $1/\text{Slope} = 3065.60$ mm Figure : $1/\text{Slope} = 3083.56$ mm

Sample is $c\tau = 6000$ mm but we measure $c\tau \approx 3000$ mm



Background Estimation



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

Have fun lets see!



Systematics



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion



Limits



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

Are we measuring the original $c\tau$ of the neutralino?

CMS Official GMSB Samples

Λ [TeV]	mass[GeV]	C_{grav}	$c\tau$ [mm]	Fit Value[mm]
120	169	93.5	1000	657.89
120	169	162	3000	1942.12
140	198	162	3000	1550.38
140	198	187	4000	2064.83
140	198	229	6000	3083.56
180	256	93.5	1000	378.64
180	256	132	2000	749.45
180	256	162	3000	1104.85
180	256	229	6000	2203.61

We seem to be measuring neutralino $c\tau$ by some factor off.



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

By how much are we off in neutralino $c\tau$ measurements?

CMS Official GMSB Samples			
Λ [TeV]	$c\tau$ [mm]	Fit Value[mm]	Factor Off
120	3000	1942.12	1.54
140	3000	1550.38	1.93
180	3000	1104.85	2.71
140	6000	3083.56	1.9
180	6000	2203..61	2.7

Factor is **The SAME** for different neutralino $c\tau$ with same Λ value. However, factor is **NOT THE SAME** for the same $c\tau$ with different Λ values.

Is this due to how sample is generated?




Figure : $1/\text{Slope} = 3083.56$ mm




Figure : $1/\text{Slope} = 3067.48$ mm




Figure : $1/\text{Slope} = 3065.60$ mm




Figure : $1/\text{Slope} = 3037.66$ mm

Private GMSB sample seems to show same offset measurements



To Be Completed



Search For
Displaced
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Motivation

Production
and Decay

Dataset
and
Trigger

Event
Selection

Background
Estimation

Systematics

Limits

Remaining
Completion

- Offset in neutralino $c\tau$ seems to have a more subtle origin than expected. Probably how mass enters into the lifetime definition and implementation at MC generation level.
- GMSB samples with the same sample $c\tau$, hence C_{grav} , but with different Λ values have different offset factor.
- The observation that the $c\tau$ value for a given sample with Λ is different from the measured value is very unclear, even without looking at samples with different Λ values.
- Our next step involves understanding cause of this offset.