



Search For Delayed Photons Using Timing.

Tambe F Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results

Summary

# **Search For Delayed Photons Using** Timing.

Shih-Chuan Kao<sup>1</sup>

Yuichi Kubota<sup>1</sup> Tambe E. Norbert<sup>1</sup> Roger Rusack<sup>1</sup>

<sup>1</sup>University Of Minnesota

Long-Lived Meeting, December 11, 2014



### Outline



- Search For Delayed Photons Using Timing.
- Tambe E. Norbert
- Outline
- Introduction
- Production
- and Decay
- Dataset and Trigger Event
- Selection

  Background
- Estimation Systematics
- Results
- Summary

- Introduction
- Production and Decay
- 3 Dataset and Trigger
- 4 Event Selection
- 5 Background Estimation
- 6 Systematics
- Results
- 8 Summary



### Where are we now?



#### Search For Delayed Photons Using Timing.

Tambe E.

Outline

Outline

Introduction

Production and Decay

Dataset and

Trigger Event

Selection Background

Estimation Systematics

System

Results

Summary


#### The Universe Set

The set 
$$S = \{ \cdots 0, \frac{1}{2}, 1, \frac{3}{2}, 2 \cdots \} \cdot \hbar$$

where s is the spin of a particle. represents our past, current and probably future understanding of the universe around us. As of the moment Currently we know:

- $s = \frac{1}{2}\hbar$  Describes all the matter in our universe.
- ullet  ${f s}={f 1}\hbar$  Describes gauge interactions.
- ullet  ${f s}={f 0}\hbar$  Responsible for giving mass.
- $\mathbf{s} = \mathbf{2}\hbar$  Describes gravity (gauged?).
- $\mathbf{s} = \frac{3}{2}\hbar$  ?? Dark Matter?

However, this magic set only describes  $\approx 4\%$  of our total



universe



### Introduction



Search For Delaved Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results


Summary

#### Long-Lived Particle Models

- ★ Gauge Mediated Supersymmetry Breaking (GMSB)
  - ▶ Next-to-lightest SUSY (NLSP) is Neutralino ( $\tilde{\chi}_1^0$ )
  - $\triangleright eV keV$  Lightest-SUSY particle (LSP) is Gravitino ( $\tilde{G}$ ).
  - Gravitino is a Dark Matter Candidate.
  - ★ General Gauge Mediation (GGM)
    - ▶ NLSP is a mixture of fermions (Bino, Wino, Higssino).
    - Several SUSY particles can be NLSP.

#### ECAL Resolution

- ECAL timing resolution  $\sigma_t < 500$  ps.
- Use timing to identify photons and electrons from long-lived decay.





### LHC Supersymmetry Production





Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger


Event Selection

Background Estimation

Systematics

Results

Summary



SUSY production mostly in strong interactions at LHC.



# Cascade Decay Chain



Search For Delayed Photons Using Timing.

Tambe E.

Norbert Outline

Introduction

Production and Decay

Dataset and

and Trigger


Event Selection

Background Estimation

Systematics

Results

Summary



Y. Kats et al: arXiv:1110.6444v2



### Delayed Photon Production



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results


Summary

### Double Photon



2 Photons, 2 Jets, Large MET

#### Single Photon





#### Tranverse Decay Distance



Search For Delayed Photons Using Timing.

Tambe E.

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results


Summary

#### Distance Travelled

$$L_T = c\tau \cdot (\gamma \beta_T) = c\tau \cdot \left(\frac{p_T}{m}\right)$$

Proper Decay Length

$$c au_{ exttt{NLSP}} = C_{ exttt{grav}}^2 rac{1}{\kappa} \left(rac{m_{ exttt{NLSP}}}{GeV}
ight)^{-5} \left(rac{\sqrt{ exttt{F}}}{TeV}
ight)^4$$



J. Ruderman, D. Shih arXiv:1103.6083



### **Datasets**



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset

and Trigger

Event Selection

Background Estimation

Systematics

Results Summary • Data  $(19.1fb^{-1})$ 

Dataset Name	Recorded Luminosity $[fb^{-1}]$
/Run2012B/SinglePhoton/EXODisplacedPhoton-PromptSkim-v3	5.1
/Run2012C/SinglePhoton/EXODisplacedPhoton-PromptSkim-v3	6.9
/Run2012D/SinglePhoton/EXODisplacedPhoton-PromptSkim-v3	7.1
/Run2012C/Cosmics/Run2012C-22Jan2013-v1/REC0	3130384(events)
/Run2012D/Cosmics/Run2012C-22Jan2013-v1/REC0	52430 (events)
/SingleElectron/Run2012A-22Jan2013-v1/AOD	5.2
/DoubleElectron/Run2012C-22Jan2013-v1/AOD	4.8

• Signal MC [GMSB (SPS8)]

$\Lambda$ [TeV]	100	120	140	160	180	300
$M_{\tilde{\chi}_1^0} [GeV/c^2]$	140	169	198	227	256	430
$c\tau$	215	325	130	245	185	
(mm)	425	645	515	490	365	495
	1700	1290	1030	975	730	
	3400	1935	2060	1945	1100	995
	5100	2955	2920	2930	2195	2960
	6000	3870	3985	3910	3950	
	9300	5985	6000	5875	5980	6000
		9825	10450	9815	10450	10450

•  $\gamma+$  Jets MC

/   3003 141	_	
$\hat{p}_T$ [GeV /c]	$\sigma_{LO}$ (pb)	Number of events
50 - 80	3322.3	1995062
80 - 120	558.3	1992627
120 - 170	108.0	2000043
170 - 300	30.1	2000069
300 - 470	2.1	2000130
470 - 800	0.212	1975231



### HLT Trigger



Search For Delaved Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics


Results

Summary

• HLT\_Photon50\_CaloIdVL\_IsoL (Study Trigger)









# **ECAL** Timing



Search For Delayed Photons Using

Timing. Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation Systematics

Results

Summary


#### Time Reconstruction


- 10 digitized samples used.
- Fit and Weighted methods used to extract time.
- Time Measurement

$$T_{MAX} = \frac{\sum_{i} \frac{T_{MAX,i}}{\sigma_i^2}}{\sum_{i} \frac{1}{\sigma_i^2}}$$

#### Time Performance

• Time resolution better than 200 ps for  $E > 30~{\rm GeV}$ 







# ECAL Timing(2)

Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation


Systematics

Results

Summary

#### Photon Timing

- $T_{\gamma} =$  Average Time of all Crystals.
- ullet  $T_{\gamma}=$  Seed (most energetic) Crystal Time.



- Similar behavior seen in Seed and Average Time.
- We use seed time as Photon Measured Time in this analysis.



### ECAL Timing(3): MC Vs Data




Figure: (LEFT): Before (RIGHT): After

- Timing corrections from data applied to  $\gamma+$  Jets MC.
- $\gamma+$  Jets MC timing aligns better with data after corrections are applied.

Search For Delayed Photons Using Timing.

Tambe E.

Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Selection

Background

Estimation Systematics

Results



### Long-Lived Decay



Search For Delayed Photons Using Timing.

Tambe E.

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation


Systematics

Results

Summary

Source of Delayed Photon?

- Slow moving particle;  $\beta << 1$ ,
- Non-nominal flight path,
- Stopped in subdetectors,





# Slow Vs Off-Pointing Decay



Search For Delayed Photons Using Timing. Tambe E.

Norbert

Outline

Outlir

Introduction

Production and Decay

Dataset and Trigger


Event Selection


Background Estimation Systematics


Systeme

Results

Summary







Delayed photons mostly from slow moving neutralino decays.



## **Event Selection**



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results

Summary

Object Selection Criteria

Variable	Selection Cuts
Photon $p_T(\gamma^{1(2)})$	> 80(45)  GeV
$ \eta_{\gamma} $ ,(EB only),	< 3.0 (< 1.5)
Semi-minor axis $(S_{Minor})$	$0.12 \le S_{Minor} \le 0.38$
H/E	< 0.05
Track Vito, $\Delta R(\gamma, track)$	> 0.6
HCAL, ECAL, Track, Isolation	< 4.0, < 4.5, < 0.2
Cone Size(Iso $\gamma$ ) $\Delta R(\gamma,SC)$	< 0.4
Spike Swiss-Cross	$1 - E_4/E_1) < 0.98$
Jets must satisfy	JetID Requirements
Leading Jet $p_T$	$>35~{\sf GeV}$
Number Of Constituents	> 1
$\Delta R(\gamma, jet) = \sqrt{(\phi_{\gamma} - \phi_{jet})^2 + (\eta_{\gamma} - \eta_{jet})^2}$	> 0.3
$E_T^{miss}$	$>25~{\rm GeV}$



#### Kinematics Distribution





Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger


Event Selection


Background Estimation

Systematics

Results

Summary





• Different  $\Lambda$  values with the same  $c\tau(10 \text{ m})$ . Photon  $p_T$  is harder with higher values of  $\Lambda$ .



### Signal Efficiency and Acceptance





#### Norbert Outline

Introduction

..... oddeti.

Production and Decay

Dataset and Trigger


Event Selection


Background Estimation

Systematics

Results

Summary





Sharp drop in efficiency immediately beyond ECAL radius for slow moving neutralino decay as source of delayed photon.



### Signal Efficiency and Acceptance(II)





Norbert Outline

Introduction

Production and Decay

Dataset and Trigger


Event Selection

Background Estimation

Systematics

Results

Summary






Figure: Off-Pointing

Acceptance peaks at transverse decay length 800 mm with delayed photons from off-pointing neutralino decays.



### Signal Efficiency and Acceptance(III)



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results




Figure: 2 Dim Efficiency



# Sources Of Background



Search For Delayed Photons Using Timing.

Norbert Outline

<u>Int</u>roduction

Tambe E.

meroduction

Production and Decay

Dataset and Trigger

Trigger Event


Selection

Background
Estimation

Systematics

\_ .

Results





# **Background Estimation**



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation


Systematics

Results

Summary

Main sources of background to delayed photons are:

- Photons of events produced from Non-collision,
- Photons of events produced from collision with mis-measured ECAL time.



Features around  $\phi=0,\pm\pi$  and  $\eta$ -dependence shows that background sources originate from both collision and non-collision events.



#### In-Time Vs Out-Of-Time Events



Delayed Photons Using Timing.

Search For

Norbert Outline

Introduction

Introductio

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics Results

Summary

We estimate these background by defining two Control samples.

In-time events Control Sample (IT-CS)
Out-of-time events Control Sample (OT-CS)

#### Control Sample (In-time Events)

**IT-CS**: > 2 Jets Events with photon ECAL time,  $t \in [-1,1]$  ns.

#### Control Sample (Out-Of-time Events)

 $\mbox{OT-CS: }0$  Jet Events with photon ECAL time, t<-3 ns or t>2 ns.

Events from above CSs provide a unique approach to estimate possible background contribution in signal.



#### Halo Photon (HP)



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics


Results


Summary

#### Beam Halo Muons

- Proton beam interacting with gas/air particles in the beam pipe,
- Proton beam colliding with the collimators upstream prior to entering the CMS detector.

will produce energetic muons traveling parallel with main proton beam and showering in the Calorimeters.







### Halo Photon (II)



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production

and Decay

and Trigger


Event Selection

Background Estimation


Systematics Results


Summary

Using Halo kinematics, We can tag and estimate halo photons produced from halo muons showering in ECAL as follows:



#### Halo Expected Time







### HP Tagging Efficiency/mis-Tag Rate



Using Timing. Tambe E. Norbert

Search For Delayed Photons

Outline

Introduction

Production and Decay

Dataset

and Trigger

Event Selection

Background Estimation

Systematics

Results



#### Cosmic Muons



Delayed Photons Using Timing.

Search For

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and

Trigger Event

Selection Background

Estimation Estimation

Systematics

Results



#### Anomalous ECAL Spike



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

.....oudceic

Production and Decay

Dataset

and Trigger

Event Selection

Background Estimation

Systematics

Results



### ABCD Technique: Non-Collision Background



Using Timing. Tambe E. Norbert

Search For Delayed Photons

Outline

Introduction

Production

and Decay

Dataset

and Trigger

Event Selection

Background Estimation

Systematics

Results



### ABCD Technique: Collision Background



Photons
Using
Timing.

Tambe E.
Norbert

Search For Delayed

Outline

Introduction

Production

and Decay

Dataset

and Trigger

Event Selection

Background Estimation

Systematics

Results



### ABCD Technique: Total Background Estimation

Search For Delayed Photons Using Timing.

Norbert Outline

Introduction

Tambe E.

Production

and Decay

Dataset

and Trigger

Event Selection

Background Estimation Systematics

Results

Summary

Equations and Results.

**Closure Test**: Events with 1 - jet.



### Background Estimation Cross-Check



Photons Using Timing. Tambe E.

Search For Delayed

Norbert Outline

Introduction

Production and Decay

Dataset

and Trigger

Event Selection

Background Estimation

Systematics

Results

Summary

Using  $Z \to ee$  events.



# Systematics



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results

Summary

Background estimation is Data driven. Thus, most of a systematics come from signal,including:

#### Experimental Systematics

- Definition of Absolute or Zero time,
- ECAL time Resolution,
- Unclustered Energy,
- Jet energy scale,
- Jets energy resolution,
- Photon energy scale,
- Luminosity. We use standard CMS luminosity uncertainty.

#### Theoretical Systematics

- Choice of PDF.
- Re-normalization group equations.



# Systematics(II



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay

Dataset and

Trigger Event

Selection Background Estimation

Systematics

Results Summary Systematic Uncertainties

Cystematic Sheertametes		
Source	Uncertainty(%)	
Absolute time(Zero time)	$10 \sim 6$	
Unclustered Energy	$10 \sim 4$	
Photon Energy Scale	$4 \sim 2$	
ECAL Time Resolution	$5\sim 2$	
Jet Energy Scale	$9 \sim 3$	
Jet Energy Resolution	$9 \sim 2$	
Luminosity	2.6	
Choice of PDF	< 1	

 Systematics is obtained by studying the effects of varying by a few amount of a particular source of systematic on the total number of objects passing object selection cuts.



#### Results



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production

and Decay

Dataset

and Trigger

Event Selection

Background Estimation Systematics

Results

Summary

Events Passing Final Selection

Sample	Lifetime( $c au$ )[mm]	Number Of Events
GMSB $\Lambda=180~{ m TeV}$	10500	
${\rm GMSB}~\Lambda=180~{\rm TeV}$	6000	
${\rm GMSB}~\Lambda=180~{\rm TeV}$	4000	
${\rm GMSB}~\Lambda=180~{\rm TeV}$	3000	
${\rm GMSB}~\Lambda=180~{\rm TeV}$	2000	
${\rm GMSB}~\Lambda=180~{\rm TeV}$	1000	
${\rm GMSB}~\Lambda=180~{\rm TeV}$	500	
Data	1.00	
Background Total	0.014	



# CLs Upper Limits



Search For Delaved Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production

and Decay Dataset

and Trigger


Event Selection


Background Estimation

Systematics

Results

Summary





sample is  $c\tau=12000$  mm but we measure  $c\tau\approx10500$  mm



### $c\tau$ -Mass Limits



Search For Delayed Photons Using Timing.

Tambe E. Norbert

Outline

Introduction

Production and Decay


Dataset and Trigger

Event Selection

Background Estimation

Systematics

Results





# Summary



Delayed Photons Using Timing. Tambe E.

Search For

Norbert

Outline

Introduction

Production and Decay

Dataset

and Trigger

Event Selection

Background Estimation

Systematics Results