



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

1 / 48

Search For Delayed Photons Using Timing.

Shih-Chuan Kao¹ Yuichi Kubota¹ Tambe E. Norbert¹
Roger Rusack¹

¹University Of Minnesota

**Long-Lived Meeting,
December 16, 2014**



Outline



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

2 / 48

- 1 Introduction
- 2 Production and Decay
- 3 Dataset and Trigger
- 4 Event Selection
- 5 Analysis Strategy
- 6 Background Estimation
- 7 Systematics
- 8 Results
- 9 Summary


The Universe Set

The set $S = \{\cdots 0, \frac{1}{2}, 1, \frac{3}{2}, 2 \cdots\} \cdot \hbar$

where s is the spin of a particle. represents our past, current and probably future understanding of the universe around us. As of the moment Currently we know:

- $s = \frac{1}{2}\hbar$ Describes all the matter in our universe.
- $s = 1\hbar$ Describes gauge interactions.
- $s = 0\hbar$ Responsible for giving mass.
- $s = 2\hbar$ Describes gravity (gauged?).
- $s = \frac{3}{2}\hbar$?? **Dark Matter?**

However, this magic set only describes $\approx 4\%$ of our total




universe.

• Long-Lived Particle Models

- ★ Gauge Mediated Supersymmetry Breaking (GMSB)
 - ▷ Next-to-lightest SUSY (NLSP) is **Neutralino** ($\tilde{\chi}_1^0$)
 - ▷ $eV - keV$ Lightest-SUSY particle (LSP) is **Gravitino** (\tilde{G}).
 - ▷ Gravitino is a Dark Matter Candidate.
- ★ General Gauge Mediation (GGM)
 - ▷ NLSP is a mixture of fermions (Bino, Wino, Higgsino).
 - ▷ Several SUSY particles can be NLSP.

• ECAL Resolution

- † ECAL timing resolution
 $\sigma_t < 500$ ps.
- † Use timing to identify
photons and electrons from
long-lived decay.



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy


Background
Estimation

Systematics

Results


Summary

5 / 48



<https://twiki.cern.ch/twiki/bin/view/LHCPhysics/SUSYCrossSections>

arXiv:1206.2892



SUSY production mostly in strong interactions at LHC.

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection


Analysis
Strategy

Background
Estimation

Systematics

Results

Summary
6 / 48



Y. Kats et al: [arXiv:1110.6444v2](https://arxiv.org/abs/1110.6444v2)

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation


Systematics

Results

Summary


7 / 48

Double Photon



2 Photons, Jets, Large MET

Single Photon





1 Photon, Jets, Large MET

Distance Travelled

$$L_T = c\tau \cdot (\gamma\beta_T) = c\tau \cdot \left(\frac{p_T}{m}\right)$$

Proper Decay Length

$$c\tau_{\text{NLSP}} = C_{\text{grav}}^2 \frac{1}{\kappa} \left(\frac{m_{\text{NLSP}}}{\text{GeV}}\right)^{-5} \left(\frac{\sqrt{F}}{\text{TeV}}\right)^4$$



J. Ruderman, D. Shih [arXiv:1103.6083](https://arxiv.org/abs/1103.6083)



Datasets



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

9 / 48

• Data ($19.1 fb^{-1}$)

Dataset Name	Recorded Luminosity [fb^{-1}]
/Run2012B/SinglePhoton/EX0DisplacedPhoton-PromptSkim-v3	5.1
/Run2012C/SinglePhoton/EX0DisplacedPhoton-PromptSkim-v3	6.9
/Run2012D/SinglePhoton/EX0DisplacedPhoton-PromptSkim-v3	7.1
/Run2012C/Cosmics/Run2012C-22Jan2013-v1/RECO	3130384(events)
/Run2012D/Cosmics/Run2012C-22Jan2013-v1/RECO	52430 (events)
/SingleElectron/Run2012A-22Jan2013-v1/AOD	5.2
/DoubleElectron/Run2012C-22Jan2013-v1/AOD	4.8

• Signal MC [GMSB (SPS8)]

Λ [TeV]	100	120	140	160	180	300
$M_{\tilde{\chi}_1^0}$ [GeV/c^2]	140	169	198	227	256	430
$c\tau$	215	325	130	245	185	
(mm)	425	645	515	490	365	495
	1700	1290	1030	975	730	
	3400	1935	2060	1945	1100	995
	5100	2955	2920	2930	2195	2960
	6000	3870	3985	3910	3950	
	9300	5985	6000	5875	5980	6000
		9825	10450	9815	10450	10450

• $\gamma +$ Jets MC

\hat{p}_T [GeV / c]	σ_{LO} (pb)	Number of events
50 – 80	3322.3	1995062
80 – 120	558.3	1992627
120 – 170	108.0	2000043
170 – 300	30.1	2000069
300 – 470	2.1	2000130
470 – 800	0.212	1975231

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy




Background
Estimation

Systematics

Results

Summary

- **HLT_DisplacedPhoton65_CaloldVL_IsoL_PFMET25**
- HLT_Photon50_CaloldVL_IsoL (Study Trigger)



• Time Reconstruction



- 10 digitized samples used.
- Fit and Weighted methods used to extract time.

• Time Measurement

$$T_{MAX} = \frac{\sum_i \frac{T_{MAX,i}}{\sigma_i^2}}{\sum_i \frac{1}{\sigma_i^2}}$$


• Time Performance

- Time resolution better than 200 ps for $E > 30$ GeV



• Photon Timing

- T_γ = Average Time of all Crystals.
- T_γ = Seed (most energetic) Crystal Time.



- Similar behavior seen in Seed and Average Time.
- We use seed time as Photon Measured Time in this analysis.

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation


Systematics

Results

Summary

13 / 48

Ecal Time from Seed Crystal



Ecal Time from Seed Crystal





Figure : (LEFT): Before (RIGHT): After

- Timing corrections from data applied to γ + Jets MC.
- γ + Jets MC timing aligns better with data after corrections are applied.

• Source of Delayed Photon?



- Slow moving particle; $\beta \ll 1$,
- Non-nominal flight path,
- Stopped in subdetectors,



Photon Arrival Time

$$\Delta t_1 = (L1/c\beta) - (L1/c)$$

$$\Delta t_2 = (L1 + L2 - L3)/c$$



Delayed photons mostly from slow moving neutralino decays.



Event Selection



Object Selection Criteria

Variable	Selection Cuts
Photon $p_T(\gamma^{1(2)})$	$> 80(45) \text{ GeV}$
$ \eta_\gamma , (\text{EB only}),$	$< 3.0(< 1.5)$
Semi-minor axis(S_{Minor})	$0.12 \leq S_{Minor} \leq 0.38$
H/E	< 0.05
Track Veto, $\Delta R(\gamma, track)$	> 0.6
HCAL, ECAL, Track, Isolation	$< 4.0, < 4.5, < 0.2$
Cone Size(Iso γ) $\Delta R(\gamma, SC)$	< 0.4
Spike Swiss-Cross	$1 - E_4/E_1 < 0.98$
Jets must satisfy	JetID Requirements
Leading Jet p_T	$> 35 \text{ GeV}$
Number Of Constituents	> 1
$\Delta R(\gamma, jet) = \sqrt{(\phi_\gamma - \phi_{jet})^2 + (\eta_\gamma - \eta_{jet})^2}$	> 0.3
E_T^{miss}	$> 25 \text{ GeV}$

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

17 / 48




Figure : Photon p_T




Figure : E_T^{miss}

- Different Λ values with the same $c\tau(10 \text{ m})$. Photon p_T is harder with higher values of Λ .

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection



Analysis
Strategy

Background
Estimation

Systematics

Results

Summary
18 / 48



Sharp drop in efficiency immediately beyond ECAL radius for slow moving neutralino decay as source of delayed photon.

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary
19 / 48




Figure : Slow Moving




Figure : Off-Pointing

Acceptance peaks at transverse decay length 800 mm with delayed photons from off-pointing neutralino decays.

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

20 / 48




Figure : 2 Dim Efficiency



Analysis Strategy



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

21 / 48

- **Background Source**

- **Collision:** Mis-measured time of $Z/W/top$ events.
- **Non-Collision:** Out-time events from LHC proton Beam/Cosmic/Anomalous Spikes.

- **Strategy**

- I Identify, tag and reject Non-Collision events.
- II Perform ABCD background estimation technique on residual non-collision events.
- III Perform ABCD background estimation technique on collision events.
- IV Performed a combined ABCD background estimation technique.

- **Clusure Test:** Verify background estimation methodology by performing a combined ABCD technique on a control sample.
- **Cross-Check:** Background estimation of collision events on another Control Sample.



Sources Of Background



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy


Background
Estimation

Systematics



Results

Summary

22 / 48



- ▶ Non-collision events like proton Beam Induced Background (BIM or Halos)/Cosmic/Anomalous spikes contribute towards delayed photons ECAL timing.
- ▶ Need to define a cleaning mechanism for identifying and rejecting non-collision events.



Features around $\phi = 0, \pm\pi$ and η -dependence shows that background sources originate from both collision and non-collision events.



In-Time Vs Out-Of-Time Events



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

24 / 48

We estimate these background by defining two Control samples.

In-time events Control Sample (IT-CS)

Out-of-time events Control Sample (OT-CS)

Control Sample (In-time Events)

IT-CS: > 2 Jets Events with photon ECAL time, $t \in [-1, 1]$ ns.

Control Sample (Out-Of-time Events)

OT-CS: 0 Jet Events with photon ECAL time, $t < -3$ ns
or $t > 2$ ns.

Events from above CSs provide a unique approach to estimate possible background contribution in signal.

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results



Summary

25 / 48


Beam Halo Muons

- Proton beam interacting with gas/air particles in the beam pipe,
- Proton beam colliding with the collimators upstream prior to entering the CMS detector.

will produce energetic muons traveling parallel with main proton beam and showering in the Calorimeters.



Using Halo kinematics, We can tag and estimate halo photons produced from halo muons showering in ECAL as follows:




Halo Expected ECAL Time


$$t_0 = \frac{\rho}{c} = \frac{R}{\sin \theta} \frac{1}{c}, \quad t_{halo} = \frac{Z}{c} = \frac{R}{\tan \theta} \frac{1}{c}$$

$$\Delta t_H^{exp} = t_{halo} - t_0 = -\frac{R}{2c} \exp^{-\eta}$$

eta vs Ecal time




Additionally, using halo muon hits from CSC segment matched in ϕ to Superclusters in ECAL, we can additionally identify, tag and remove halo photon events with large timing.



Halo Photon Matching


$$\Delta\phi(CSCSeg, \gamma) = |\phi_{CSCSeg} - \phi_{\gamma}|$$



Satellite/Ghost Beam Halos

- Fill empty RF buckets.
- Trail main bunches by ≈ 5 ns.
- 10^{-5} protons compared to main bunches.
- Can contribute to main collision photons.
- Show a 2.5 ns pattern in EE,
- Tagged using $\Delta\phi(CSCseg, \gamma)$.

LHC LDM Proton Beam Profile




Halo Photon Event Properties

- Halo photons populate around $\phi = 0, \pm\pi$
- ECAL time mostly < -3 ns but can also arrive late(ghosts).
- Halo events most contain no jets (0-jet events).
- Rare cases can be associated with "pile-up" events.

Halo Photon Tagging Criteria

- Use $\Delta\phi(CSCseg, \gamma) < 0.05$ radians.
- Shower shape(
 $0.8 < S_{Major} < 1.65$ and
 $S_{minor} < 0.2$)

Ghost/Satellite EE





Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

30 / 48

Halo Photon Tagging Efficiency

- Control Sample Selection,
 - $\Delta\phi(CSCseg, \gamma) < 0.05$ radians
 - Same $\Delta t_H^{exp} = -\frac{R}{2c} \exp^{-\eta}$ ECAL time Vs η dependence.
- Efficiency evaluated in 5η bins for S_{Major} η dependence.

Halo Photon mis-Tag Rate

- Control Sample Selection:
 - ≥ 2 -jets events with $E_T^{miss} < 60$ GeV
 - ECAL time, $|t| < 1$ ns.
- mis-tag rate evaluated in 5η bins for S_{major} η dependence.

Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation


Systematics

Results

Summary


31 / 48

Tagging Efficiency



- Tagging Efficiency $\approx 98\%$

mis-Tag Rate





- mis-tag rate $\approx 3\%$

Cosmic Muons

- Muons from cosmic rays in CMS detector.
- Hits in muon detectors (DT/CSC) and shower in ECAL.
- Produce energetic photons with out-of-time.
- Using DT segment matched to ECAL cluster position in $\delta\eta$ and $\delta\phi$ can eliminate cosmic events.

$DT(\delta\eta, \delta\phi)$ Cosmic Muon dataset(left) and Data(Right)



$DT(\delta\eta, \delta\phi)$ tagging of cosmic muons in data and a pure cosmic sample(without LHC proton beam) is comparable.



Anomalous ECAL Spike



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results



Summary

33 / 48

ECAL Spikes


- Energetic particles(neutrons) from proton collision directly hitting APDs/VPTs.
- Associated with hadronic activity.
- Observed as photons with early time due to no crystal scintillation.
- Can produced late ECAL timing photons with small shower shape.
- ID and rejected requiring $1 - \frac{E_4}{E_1} < 0.9$ of crystal energy deposit and χ^2 from pulse shape fitting.

Spike Identification and Rejection



After tagging and cleaning Halo/Cosmic/Spike events, We apply **ABCD** background estimation technique on residual Non-collision background events to estimate their contribution to possible signal.

Event Tagging and Cleaning Performance





PF- E_T^{miss} (PF-MET) Adjustment



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

35 / 48

$$E_T^{\text{miss}}(\gamma)$$

PF-MET calculation fails to take into consideration E_T from out-of-time photons. We make PF-MET adjustment by taking into account the E_T of out-of-time photons for E_T^{miss} measurements. This new PF-MET is called $E_T^{\text{miss}}(\gamma)$.

As a result our signal selection criteria is defined as:

Signal Selection Criteria

SIGNAL: $\geq 1\gamma + \geq 2\text{Jets} + E_T^{\text{miss}} > 60, E_T^{\text{miss}}(\gamma) > 60 \text{ GeV}$

Assume similar distribution in earlier and delayed ECAL time for untagged non-collision events.


$$E_T^{\text{miss}} > 60 \text{ GeV}$$

$$\frac{D}{C} = \frac{B}{A}, \Rightarrow$$

	$E_T^{\text{miss}}(\gamma) < 60$	$E_T^{\text{miss}}(\gamma) > 60$
$t < -3 \text{ ns}$	A	B
$t > 3 \text{ ns}$	C	D

$$D = \frac{B}{A} \cdot C$$

Picture Explaining non-collision untagged background





ABCD Technique: Collision Background



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

37 / 48

Assume similar distribution in earlier and in-time, in-time and delayed ECAL time for collision events.

$$E_T^{\text{miss}}(\gamma) < 60 \text{ GeV}$$

	$E_T^{\text{miss}} < 60$	$E_T^{\text{miss}} > 60$
$t < -3 \text{ ns}$	B'	B
$ t < 2 \text{ ns}$	F'	F
$t > 3 \text{ ns}$	D'	D

$$\frac{Q}{D'} = \frac{F}{F'}, \Rightarrow$$


$$Q_d = \frac{F}{F'} \cdot D'$$

Picture

Explaining

collision

background





Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

38 / 48

Equations and Results.

Closure Test Results: 0,1-Jet Events





Results Of Background Estimation



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

39 / 48



Background Estimation Cross-Check



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

40 / 48

Using $Z \rightarrow ee$ events.



Background estimation is Data driven. Thus, most of a systematics come from signal, including:

Experimental Systematics

- Definition of Absolute or Zero time,
- ECAL time Resolution,
- Unclustered Energy,
- Jet energy scale,
- Jets energy resolution,
- Photon energy scale,
- Luminosity. We use standard CMS luminosity uncertainty.

Theoretical Systematics

- Choice of PDF.
- Re-normalization group equations.

Systematic Uncertainties

Source	Uncertainty(%)
Absolute time(Zero time)	10 ~ 6
Unclustered Energy	10 ~ 4
Photon Energy Scale	4 ~ 2
ECAL Time Resolution	5 ~ 2
Jet Energy Scale	9 ~ 3
Jet Energy Resolution	9 ~ 2
Luminosity	2.6
Choice of PDF	< 1

- Systematics is obtained by studying the effects of varying by a few amount of a particular source of systematic on the total number of objects passing object selection cuts.



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

43 / 48

Events Passing Final Selection

Sample	Lifetime($c\tau$)[mm]	Number Of Events
GMSB $\Lambda = 180$ TeV	10500	
GMSB $\Lambda = 180$ TeV	6000	
GMSB $\Lambda = 180$ TeV	4000	
GMSB $\Lambda = 180$ TeV	3000	
GMSB $\Lambda = 180$ TeV	2000	
GMSB $\Lambda = 180$ TeV	1000	
GMSB $\Lambda = 180$ TeV	500	
Data	1.00	
Background Total	0.014	



Observed Event



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation


Systematics

Results



Summary

44 / 48

3-D View



Transverse (left) and $\rho - Z$ (Right) View





Exclusion Limits



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

45 / 48




Figure : $c\tau$ Limits




Figure : Mass Limit

sample is $c\tau = 12000$ mm but we measure $c\tau \approx 10500$ mm



CT-Mass Limits



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy


Background
Estimation

Systematics

Results

Summary

46 / 48





Summary



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary

47 / 48



Search For
Delayed
Photons
Using
Timing.

Tambe E.
Norbert

Outline

Introduction

Production
and Decay

Dataset
and
Trigger

Event
Selection

Analysis
Strategy

Background
Estimation

Systematics

Results

Summary