

Age determination in herring (*Clupea harengus*)

Using otoliths in the
Quebec region


Helene Dionne

Senior Technician Herring Stock
Assessment

Pelagic and Ecosystem Science Branch

Maurice Lamontagne Institute

helene.dionne@dfo-mpo.gc.ca



Plan of the presentation

- Preparation of otoliths
- Reading otoliths
- Challenges
- Future projects
- Conclusion



Preparation of otoliths




- Extraction of sagitta
- Cleaning, placing on plate and drying in the oven (24hrs at 55°C)
- Bonding with resin composed of 80% Cytoseal™ 60 and 20% toluene
- Drying for 24 hours under a hood and in an oven (24 hours at 55°C) to remove fumes

Reading of otoliths

Determination of the spawning group

- Spring



- According to the shape of the first ring
- Pararost and Postrostrum
- Excisure
- Concave or convex core
- General shape (narrow or wide)
- Serrated edge

- Fall



Ref: Messieh, S.N. et al. 1989


Otolith reading

Age determination

- Herring harvested in spring (April-June)
 - ❖ Count the dark hyaline areas and add the last opaque growth area (if present)

For herring harvested in July, the addition or not of the opaque growth zone is at the discretion of the reader.

- Herring harvested in the fall (August-December)
 - ❖ Count dark hyaline areas



Otolith reading

We read the otoliths blind that is to say that we do not know the length or the stage of maturity of the fish

Ref: Penttila, J. and L.M. Dery 1988.

We read the 2 otoliths and in several places on the same otolith



We only count the rings that are visible all around the otolith



Starting from the core, the distance between the rings should go from larger to smaller




The age on antirostre is often exact or underestimated (-1 year)

Otolith reading Validation

- Validation of intra-reader accuracy with a sub-sample (50-100) each year
- Draw of 20% of otoliths made for validation by a second reader (inter-reader)
- Verification of agreement with CV ($\leq 5\%$) and % agreement ($\geq 80\%$)
- Discussion between readers and correction (if agreed)

Sample Type/year		Species		Hareng			
Aged 2X by ____		Date		1-4-21			
		Age Reader		Hélène Dionne +Geneviève Perrin			
					Bowker's test		
N Aged	120	Total CV	2.59%	Chi-sq	19.00		
N Agreed	101			d.f.	11		
Disagreed	19	% Agreement	84.2%	P-value	0.06		
					n/s		



2022	69589	9	C9	P	5	P	5		
		23	D11	A	5	A	5		
		27	E3	A	5	A	5		
		29	E5	P	5	P	5		
		31	E7	P	5	P?	5		
		38	F2	P	5	P	5		
		42	F6	A	4	A	4		
		45	F9	A	5	A?	5		
		46	F10	P?	5	P	5		
		47	F11	A	4	P?	5	A	5
		48	F12	A	6	A?	5		5
		49	G1	P	5	P	5		
		52	G4	P	6	P	6		
		57	G9	P	5	P	5		
		59	G11	P	3	P?	4	A	3

Challenges



- Determining the spawning group of juvenile herring is rather difficult.
 - ❖ The characteristics of the otoliths discussed earlier are not always very obvious.
 - ❖ Sometimes the characteristics contradict each other (with angle but first ring shorter than the antirostrum, etc.)
 - ❖ There is always a certain subjectivity depending on the reader (visual acuity, contrast detection, etc.)

Challenges


- New readers
 - ❖ Training with a reference collection
 - ❖ Intra and inter reader accuracy
- The last ring on the outer border is not present on the whole otolith. A doubt remains whether the latter should be added or not.
- The date of July 1 to separate the spawning groups

Future projects

- Otolith microstructure
- Shape R



FA 67266_137




FA 67266_137



In conclusion

- Despite the challenges encountered and discussed previously, we should not be too far from the truth because we manage to follow the cohorts!



References

Messieh, S.N., C. MacDougal J, and R. Claytor. 1989. Separation of Atlantic herring stocks in the southern Gulf of St. Lawrence using digitized otolith morphometrics and discriminant function analysis. Can. Tech. Rep. 1647: iv + 22 p.

Penttila, J., and L. M. Dery. 1988. Age determination methods for northwest Atlantic species. NOAA Tech. Rep. NMFS-72; 135 p.
[Available from: <http://www.nefsc.noaa.gov/fbi/age-man.html>]

Campana, S.E., M.C. Annand and J.I. McMillan. 1995. Graphical and Statistical methods for determining the consistency of age determinations. Trans. Am. Fish. Soc. 124: 131-138.

Appendix 3b. Proposed Training Outline for New Age Reader, Inter-Régional ageing workshop, June 17-18, 1997.