## Assignment 2

## T.Guru Balaji

## Download all python codes from

https://github.com/TGURUBALAJI/INTERNSHIP -IITH/Assignment2/code

and latex-tikz codes from

https://github.com/TGURUBALAJI/INTERNSHIP -IITH/Assignment2/gbalaji.tex

**2.106** Find the values of k for which the line

$$(k-3 - (4-k^2))\mathbf{x} + k^2 - 7k + 6 = 0$$
 (1)

is

- a) Parallel to the x-axis
- b) Parallel to the y-axis
- c) Passing through the origin

Solution Given equation of the line,

$$(k-3 - (4-k^2))\mathbf{x} + k^2 - 7k + 6 = 0$$
 (2)

of a general line equation  $\mathbf{n}\mathbf{x} = c$ here  $\mathbf{n} = (k-3 - \{4-k^2\})$ and  $c = -k^2 + 7k - 6$ 

a) Parallel to x-axis

 $\mathbf{n} = \begin{pmatrix} 0 & 1 \end{pmatrix}$  if the line is parallel to x-axis Equation of x-axis is  $\begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{x} = 0$ 

$$(1 \quad 0) \binom{k-3}{-\{4-k^2\}} = 0$$
 (3)

$$k - 3 = 0 \tag{4}$$

$$k = 3 \tag{5}$$

Substituting k = 3 in (2) Equation of line is,

$$\begin{pmatrix} 0 & 5 \end{pmatrix} \mathbf{x} = 6 \tag{6}$$

b) Parallel to y-axis  $\mathbf{n} = \begin{pmatrix} 1 & 0 \end{pmatrix}$  if the line is parallel to y-axis

Equation of y-axis is  $(0 \ 1)\mathbf{x} = 0$ 

$$(0 \quad 1) \binom{k-3}{-(4-k^2)} = 0 \tag{7}$$

$$4 - k^2 = 0 (8)$$

$$k = \pm 2 \tag{9}$$

Substituting k = 2 in (2). Equation of line is,

$$\begin{pmatrix} -1 & 0 \end{pmatrix} \mathbf{x} = 12 \tag{10}$$

Substituting k = -2 in (2). Equation of line is,

$$\begin{pmatrix} -5 & 0 \end{pmatrix} \mathbf{x} = -24 \tag{11}$$

c) Passing through origin

c = 0 if the line passes through origin Equation of line when passing through origin is

$$\mathbf{n}^{\mathsf{T}}\mathbf{x} = 0 \tag{12}$$

Hence

$$-k^2 + 7k - 6 = 0 ag{13}$$

$$(k-1)(k-6) = 0 (14)$$

$$k = 1, k = 6$$
 (15)

Substituting k = 1 in (2). The equation of line is,

$$(-2 \quad -3)\mathbf{x} = 0 \tag{16}$$

Substituting k = 6 in (2). The equation of line is,

$$(3 \quad 32)\mathbf{x} = 0 \tag{17}$$




Fig. 3: Plot of line equations