

Assignment 2

T.Guru Balaji

Download all python codes from

<https://github.com/TGURUBALAJI/INTERNSHIP-IITH/Assignment2/code>

and latex-tikz codes from

<https://github.com/TGURUBALAJI/INTERNSHIP-IITH/Assignment2/gbalaji.tex>

1 QUESTION No. 2.106 - LINEAR FORMS

Find the values of k for which the line

$$(k-3 \quad -(4-k^2))\mathbf{x} + k^2 - 7k + 6 = 0 \quad (1.0.1)$$

is

- a) Parallel to the x-axis
- b) Parallel to the y-axis
- c) Passing through the origin

2 SOLUTION

Given equation of the line,

$$(k-3 \quad -(4-k^2))\mathbf{x} + k^2 - 7k + 6 = 0 \quad (2.0.1)$$

- 1) Parallel to x-axis Equation of x-axis is $(1 \quad 0)\mathbf{x} = 0$

$$(1 \quad 0)\begin{pmatrix} k-3 \\ -(4-k^2) \end{pmatrix} = 0 \quad (2.0.2)$$

$$k-3 = 0 \quad (2.0.3)$$

$$k = 3 \quad (2.0.4)$$

Substituting $k = 3$ in (2.0.1)

Equation of line is,

$$(0 \quad 5)\mathbf{x} = 6 \quad (2.0.5)$$

- 2) Parallel to y-axis Equation of y-axis is $(0 \quad 1)\mathbf{x} = 0$

$$(0 \quad 1)\begin{pmatrix} k-3 \\ -(4-k^2) \end{pmatrix} = 0 \quad (2.0.6)$$

$$4 - k^2 = 0 \quad (2.0.7)$$

$$k = \pm 2 \quad (2.0.8)$$

Substituting $k = 2$ in (2.0.1)

Equation of line is,

$$(-1 \quad 0)\mathbf{x} = 12 \quad (2.0.9)$$

Substituting $k = -2$ in (2.0.1)

Equation of line is,

$$(-5 \quad 0)\mathbf{x} = -16 \quad (2.0.10)$$

- 3) passing through origin Equation of line when passing through origin is

$$\mathbf{n}^T \mathbf{x} = 0 \quad (2.0.11)$$

Hence

$$-k^2 + 7k - 6 = 0 \quad (2.0.12)$$

$$(k-1)(k-6) = 0 \quad (2.0.13)$$

$$k = 1, k = 6 \quad (2.0.14)$$

Substituting $k = 1$ in (2.0.1)

The equation of line is,

$$(-2 \quad -3)\mathbf{x} = 0 \quad (2.0.15)$$

Substituting $k = 6$ in (2.0.1)

The equation of line is,

$$(3 \quad 32)\mathbf{x} = 0 \quad (2.0.16)$$




Fig. 3: a) Plot of line parallel to x-axis




Fig. 3: b) Plot of line parallel to y-axis




Fig. 3: c) passing through origin