

Math 310

Homework 6

Due: 10/9/2024

Name: Gianluca Crescenzo

Exercise 1. Let $(x_k)_k$ be a sequence of strictly positive numbers such that

$$(kx_k)_k \rightarrow L > 0/.$$

Show that the infinite series $\sum_k x_k$ diverges.

Exercise 2. Let $(x_k)_k$ be a sequence of strictly positive numbers.

(i) If $\limsup_{n \rightarrow \infty} \frac{x_{k+1}}{x_k} < 1$, then the infinite series $\sum_k x_k$ converges.

(ii) If $\liminf_{n \rightarrow \infty} \frac{x_{k+1}}{x_k} > 1$, then the infinite series $\sum_k x_k$ diverges.

Exercise 3. Consider the sequence of functions:

$$f_n : \mathbf{R} \rightarrow \mathbf{R}; \quad f_n(x) = \arctan(nx).$$

(i) Show that $(f_n)_n \rightarrow \frac{\pi}{2} \text{sgn}$ point-wise.

(ii) Show that the convergence in (i) is nonuniform on $(0, \infty)$.

(iii) Show that the convergence in (i) is uniform on $[a, \infty)$ for a fixed $a > 0$.

Proof. Note that:

$$\begin{aligned} (\arctan(n))_n &\rightarrow \frac{\pi}{2}, \\ (\arctan(-n))_n &\rightarrow -\frac{\pi}{2}. \end{aligned}$$

So given $x > 0$, there exists $N_x \in \mathbf{N}$ such that $n \geq N_x$ implies $|\arctan(nx) - \frac{\pi}{2}| < \epsilon$. Similarly, given $x < 0$, there exists $N_x \in \mathbf{N}$ such that $n \geq N_x$ implies $|\arctan(nx) + \frac{\pi}{2}| < \epsilon$. For $x = 0$, we have that $(\arctan(0))_n = (0)_n \rightarrow 0_{\mathcal{F}(\mathbf{R}, \mathbf{R})}$. Hence $(\arctan(nx))_n \rightarrow \text{sign} \frac{\pi}{2}$.

Let $(x_k)_k = \frac{1}{k}$ and $n_k = k$. Observe that:

$$\begin{aligned} |f_{n_k}(x_k) - f(x_k)| &= \left| \arctan\left(k \cdot \frac{1}{k}\right) - \text{sign}\left(\frac{1}{k}\right) \cdot \frac{\pi}{2} \right| \\ &= \arctan(1). \end{aligned}$$

Picking $\epsilon_0 = \arctan(1)$ gives that $(\arctan(nx))_n$ does not converge uniformly on $(0, \infty)$.

Fix $a > 0$. Since $(d_u(f_n, f))_n = \left(\sup_{x \in [a, \infty)} \left| \arctan(nx) - \text{sign}(x) \frac{\pi}{2} \right| \right)_n$, we have:

$$\begin{aligned} \left| \sup_{x \in [a, \infty)} \left| \arctan(nx) - \text{sign}(x) \frac{\pi}{2} \right| \right| &\leq \sup_{x \in [a, \infty)} \left| \arctan(nx) - \text{sign}(x) \frac{\pi}{2} \right| \\ &= \sup_{x \in [a, \infty)} \left| \arctan(nx) - \frac{\pi}{2} \right| \\ &= 0. \end{aligned}$$

Thus $(f_n)_n$ converges uniformly on $[a, \infty)$. □

Exercise 4. Consider the sequence of functions:

$$f_n : [0, \infty) \rightarrow \mathbf{R}; \quad f_n(x) = \frac{\sin(nx)}{1 + nx}.$$

(i) Show that $(f_n)_n \rightarrow 0$ point-wise.

(ii) Show that the convergence in (i) is nonuniform on $(0, \infty)$.

(iii) Show that the convergence in (i) is uniform on $[a, \infty)$ for a fixed $a > 0$.

Proof. For $x = 0$, we have that $(f_n(0))_n = 0_{\mathcal{F}([0, \infty), \mathbf{R})}$. For $x > 0$:

$$\left| \frac{\sin(nx)}{1 + nx} \right| \leq \frac{1}{1 + n|x|} \leq \frac{1}{n|x|}.$$

Since $\left(\frac{1}{n|x|}\right)_n \rightarrow 0$, we have that $\left(\frac{\sin(nx)}{1+nx}\right)_n \rightarrow 0$. Hence $(f_n)_n \rightarrow 0_{\mathcal{F}([0, \infty), \mathbf{R})}$ pointwise.

Consider $x_k = \frac{\pi}{2k}$ and $n_k = k$. We have:

$$\begin{aligned} |f_{n_k}(x_k) - f(x_k)| &= \left| \frac{\sin(kx_k)}{1 + kx_k} \right| \\ &= \frac{\sin\left(k \cdot \frac{\pi}{2k}\right)}{1 + k \cdot \frac{\pi}{2k}} \\ &= \frac{\sin\left(\frac{\pi}{2}\right)}{1 + \frac{\pi}{2}} \\ &= \frac{1}{1 + \frac{\pi}{2}}. \end{aligned}$$

Picking $\epsilon_0 = \frac{1}{1 + \frac{\pi}{2}}$ gives that $(f_n)_n$ does not converge uniformly on $(0, \infty)$.

Fix $a > 0$. Since $(d_u(f_n, f))_n = \left(\sup_{x \in [a, \infty)} \left| \frac{\sin(nx)}{1+nx} \right| \right)_n$, we have:

$$\begin{aligned} \left| \sup_{x \in [a, \infty)} \left| \frac{\sin(nx)}{1 + nx} \right| \right| &\leq \sup_{x \in [a, \infty)} \left| \frac{\sin(nx)}{1 + nx} \right| \\ &\leq \sup_{x \in [a, \infty)} \left| \frac{1}{1 + nx} \right| \\ &= \frac{1}{1 + na} \\ &\leq \frac{1}{na}. \end{aligned}$$

Since $\left(\frac{1}{na}\right)_n \rightarrow 0$, then $(d_u(f_n, f))_n \rightarrow 0$. Thus f_n converges uniformly on $[a, \infty)$. □

Exercise 5. Show that the sequence of functions:

$$f_n : [0, \infty) \rightarrow \mathbf{R}; \quad f_n(x) = x^2 e^{-nx}$$

converges uniformly to 0.

Proof. Note that $(f_n)_n \rightarrow 0_{\mathcal{F}([0,\infty),\mathbf{R})}$. We have that $(d_u(f_n, f))_n = \left(\sup_{x \in [0,\infty)} |x^2 e^{-nx}| \right)_n$. Observe that:

$$\begin{aligned} \left| \sup_{x \in [0,\infty)} |x^2 e^{-nx}| \right| &\leq \sup_{x \in [0,\infty)} |x^2 e^{-nx}| \\ &\leq \sup_{x \in [0,\infty)} \left| \frac{x^2}{1 + x + \frac{n^2 x^2}{2}} \right| \\ &\leq \sup_{x \in [0,\infty)} \left| \frac{x^2}{\frac{n^2}{2} x^2} \right| \\ &= \sup_{x \in [0,\infty)} \left| \frac{2}{n^2} \right| \\ &= \frac{2}{n^2}. \end{aligned}$$

Since $\left(\frac{2}{n^2} \right)_n \rightarrow 0$, $(d_u(f_n, f))_n \rightarrow 0$. Thus $(f_n)_n$ converges uniformly on $[0, \infty)$. □

Exercise 6. Let $f_n = \mathbf{1}_{[n, n+1]}$. Show that $(f_n)_n \rightarrow 0$ point-wise on \mathbf{R} . Is the convergence uniform?

Exercise 7. Let $(f_n)_n$ and $(g_n)_n$ be sequences in $\ell_\infty(\Omega)$ with $(f_n)_n \rightarrow f$ and $(g_n)_n \rightarrow g$ uniformly on Ω . Prove that $(f_n g_n)_n \rightarrow fg$ uniformly on Ω .

Exercise 8. Find a sequence of functions $(f_n)_n$ defined on $[0, \infty)$ such that $\|f_n\|_u \geq n$ but with $(f_n)_n \rightarrow 0$ point-wise.

Exercise 9. Show that the series $\sum_{k=0}^{\infty} \frac{x^k}{k!}$ converges absolutely and uniformly on any closed and bounded interval $[a, b]$.