

Preface

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Contents

Contents	i
1 Normed Linear Spaces	1
1.1 Basic Definitions and Results	1
1.2 Completeness and Compactness	3
Exercises	4
1.3 Linear Operators	5
1.4 Bounded Linear Functionals	7
Exercises	10
1.5 ℓ^p Spaces	13
Exercises	14
1.6 Consequences of The Baire Category Theorem	15
Exercises	21
1.7 Continuous Dual, Adjoint Maps, and Weak Convergence	22
Exercises	22
1.8 L^p Spaces	23
Exercises	23
2 Hilbert Spaces	24
2.1 Inner Product Spaces	24
Exercises	26
2.2 Hilbert Spaces	27
Exercises	33
2.3 Compact Operators	40

Last update: 2026 March 16

Chapter 1

Normed Linear Spaces

§ 1.1. Basic Definitions and Results

Definition 1.1.1. A seminorm on an $\mathbb{F}$ -vector space X is a map:

$$\|\cdot\| : X \rightarrow [0, \infty)$$

satisfying:

- (1) $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{F}$ and $x \in X$ (*homogeneity*);
- (2) $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$ (*triangle inequality*).

If the following condition is also satisfied:

- (3) $\|x\| \geq 0$ and $\|x\| = 0$ if and only if $x = 0$ (*positive definiteness*),

then we say $\|\cdot\|$ is a norm. The ordered pair $(X, \|\cdot\|)$ is called a normed linear space.

Remark. In most cases, one can take $\mathbb{F}$ to be $\mathbb{R}$ or $\mathbb{C}$ without any loss of generality.

Remark. A normed linear space induces a metric by the equation $d(x, y) := \|y - x\|$.

Remark. A normed linear space is finite dimensional if it is as a vector space; otherwise it is infinite dimensional.

Definition 1.1.2. Let $(X, \|\cdot\|)$ be a normed linear space.

- (1) The open ball centered around $x_0 \in X$ with radius $r > 0$ is denoted $B(x_0, r) := \{x \in X \mid \|x - x_0\| < r\}$.
- (2) The closed ball centered around $x_0 \in X$ with radius $r > 0$ is denoted $\overline{B}(x_0, r) := \{x \in X \mid \|x - x_0\| \leq r\}$.
- (3) The sphere centered around $x_0 \in X$ with radius $r > 0$ is denoted $S(x_0, r) := \{x \in X \mid \|x - x_0\| = r\}$.

Definition 1.1.3. Two norms $\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ on a normed linear space X are said to be equivalent if there exists a constant $C \geq 1$ such that, for all $x \in X$:

$$\frac{1}{C} \|x\|_\beta \leq \|x\|_\alpha \leq C \|x\|_\beta.$$

Example 1.1.1. Let $x \in \mathbb{R}^n$, where $x := (x_1, \dots, x_n)$. Then:

$$\begin{aligned}\|x\|_2 &:= \left(\sum_{i=1}^n |x_i|^2 \right)^{\frac{1}{2}}, \\ \|x\|_1 &:= \sum_{i=1}^n |x_i|, \\ \|x\|_\infty &:= \max_{i=1}^n |x_i|, \\ \|x\|_p &:= \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}, \quad p > 1\end{aligned}$$

are norms on $\mathbb{R}^n$.

Example 1.1.2. Let $a, b \in \mathbb{R}$ with $a < b$. The set of continuous real-valued functions on $[a, b]$ is denoted $C([a, b]) := \{f : [a, b] \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$. Then:

$$\begin{aligned}\|f\|_\infty &:= \sup\{|f(x)| \mid x \in [a, b]\}, \\ \|f\|_2 &:= \left(\int_a^b |f(x)|^2 dx \right)^{\frac{1}{2}}, \\ \|f\|_p &:= \left(\int_a^b |f(x)|^p dx \right)^{\frac{1}{p}},\end{aligned}$$

are all norms on $C([a, b])$.

Definition 1.1.4. Let $(X, \|\cdot\|)$ be a normed linear space. A subset $V \subset X$ is convex if for all $x, y \in V$ and for all $t \in [0, 1]$, $tx + (1-t)y \in V$.

Example 1.1.3. Any subspace of a normed linear space is convex. Moreover, by the triangle inequality any open ball is convex.

Remark. Note that the *supremum norm* $\|\cdot\|_\infty$ on $C([a, b])$ only exists (by the Extreme Value Theorem) because $[a, b]$ is compact. Hence the above example can be generalized to a compact metric space Y . In particular, for $C(Y) := \{f : Y \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$, the norm $\|f\|_\infty := \sup\{|f(x)| \mid x \in Y\}$ always exists and is finite.

Proposition 1.1.1. Let (X, d) and (Y, ρ) be metric spaces. Let $f : X \rightarrow Y$. Then f is continuous if and only if the preimage of an open set in Y is open in X .

Proof. Suppose f is continuous. Let $V \subseteq Y$ be open. If $f^{-1}(V) = \emptyset$, then there is nothing to show. If not, pick any $x \in f^{-1}(V)$. Then $f(x) \in V$. Since V is open, there exists $\epsilon > 0$ such that $B(f(x), \epsilon) \subseteq V$. Since f is continuous, there exists a $\delta > 0$ such that $f(B(x, \delta)) \subseteq B(f(x), \epsilon) \subseteq V$. Thus $B(x, \delta) \subseteq f^{-1}(V)$; i.e., $f^{-1}(V)$ is open.

Conversely, let $x \in X$ and $\epsilon > 0$. Since $B(f(x), \epsilon)$ is open in Y , $f^{-1}(B(f(x), \epsilon))$ is open in X . Clearly $x \in f^{-1}(B(f(x), \epsilon))$. Since this set is open, there exists $\delta > 0$ such that $x \in B(x, \delta) \subseteq f^{-1}(B(f(x), \epsilon))$. Hence $f(B(x, \delta)) \subseteq B(f(x), \epsilon)$. Since the set of all open balls is a basis for a topology, any open set can be written as the union of open balls. Thus f is continuous. $\square$

Remark. It follows that f is continuous if and only if the preimage of a closed set is closed.

Definition 1.1.5. Let $(X, \|\cdot\|)$ be a normed linear space and $\{x_n\}_{n=1}^{\infty}$ be a sequence in X .

- (1) The sequence $\{x_n\}_{n=1}^{\infty}$ converges to $x \in X$ in the $\|\cdot\|$ -norm if $\|x - x_n\| \rightarrow 0$.
- (2) A series in X is a formal sum $\sum_{n=1}^{\infty} x_n$. We say the series $\sum_{n=1}^{\infty} x_n$ converges to $x \in X$ if the sequence of partial sums $\{\sum_{i=1}^n x_i\}_{n=1}^{\infty}$ converges to x . We say $\sum_{n=1}^{\infty} x_n$ is absolutely convergent if $\sum_{n=1}^{\infty} \|x_n\|$ converges.

Proposition 1.1.2. Let X be a normed linear space. The following are equivalent:

- (1) X is a Banach space;
- (2) Every absolutely convergent series in X converges.

Proof.

□

§ 1.2. Completeness and Compactness

Lemma 1.2.1. In $(\mathbb{R}^n, \|\cdot\|_{\infty})$, every closed ball $\overline{B}(0, c)$ is compact.

Proof.

□

Theorem 1.2.1. Let X be a finite dimensional normed linear space. Every closed and bounded subset of X is compact.

Proof.

□

Corollary 1.2.1. Every finite dimensional normed linear space is Banach.

Proof. Let X be a finite dimensional normed linear space and $\{x_n\}_{n=1}^{\infty}$ a Cauchy sequence. Then $\{x_n\}_{n=1}^{\infty}$ is closed and bounded, hence it is compact. Thus $\{x_n\}_{n=1}^{\infty}$ admits a convergent subsequence. But since $\{x_n\}_{n=1}^{\infty}$ is a Cauchy sequence which admits a convergent subsequence, $\{x_n\}_{n=1}^{\infty}$ is convergent. Thus X is Banach. □

Corollary 1.2.2. Every finite dimensional subspace of a normed linear space is closed.

Proof. This follows from Corollary 1.2.1 and Exercise 1.2.4. □

Lemma 1.2.2 (Riesz's Lemma). *Let $(X, \|\cdot\|)$ be a normed linear space. Let Z be a closed and proper subspace of X . Let $0 < \lambda < 1$ be fixed. Then there exists some $x \in X \setminus Z$ such that $\|x\| = 1$ and $\text{dist}(x, Z) > \lambda$.*

Proof.

□

Theorem 1.2.2. *Let X be a normed linear space. Then X is finite dimensional if and only if $\overline{B}(0, 1)$ is compact.*

Proof.

□

Exercises

Exercise 1.2.1. *Let K be a compact metric space (fine to assume that $K = [0, 1]$). Show that $C(K)$ paired with the sup-norm $\|\cdot\|_\infty$ is Banach.*

Proof.

□

Exercise 1.2.2. *In any metric space, a closed subset of a compact set is compact.*

Proof.

□

Exercise 1.2.3. *In any metric space, a Cauchy sequence which has a convergent subsequence must converge.*

Proof.

□

Exercise 1.2.4. *A subspace of a Banach space is closed if and only if it is Banach.*

Proof. Let X be a Banach space and Y a subspace. Suppose that Y is closed. Let $\{y_n\}_{n=1}^\infty \subseteq Y$ be a Cauchy sequence. Then $\{y_n\}_{n=1}^\infty$ is Cauchy in X . Therefore $y_n \rightarrow x$ for some $x \in X$. But since Y is closed, it contains all of its limit points. Hence $x \in Y$. Therefore $\{y_n\}_{n=1}^\infty$ converges in Y , establishing that Y is Banach.

Conversely, suppose that Y is Banach. Let $\{y_n\}_{n=1}^\infty \subseteq Y$ be a sequence such that $y_n \rightarrow x \in X$. Note that $\{y_n\}_{n=1}^\infty$ is convergent, hence Cauchy. But since Y is Banach, $\{y_n\}_{n=1}^\infty$ converges to some $y \in Y$. It must be the case that $x = y \in Y$. Therefore Y contains all of its limit points; i.e., it is closed in X . □

Exercise 1.2.5. In any normed linear space X with closed subspace Z , if $x \in X \setminus Z$ then $\text{dist}(x, Z) > 0$.

Proof. Suppose towards contradiction $\text{dist}(x, Z) = \inf\{\|x - z\| \mid z \in Z\} = 0$. We can find a sequence $\{z_n\}_{n=1}^\infty \subseteq Z$ such that $\|x - z_n\| \rightarrow 0$. Hence $z_n \rightarrow x$. But since Z is closed, $x \in Z$. This is a contradiction, hence $\text{dist}(x, Z) > 0$. $\square$

Exercise 1.2.6. In any normed linear space X with subspace Z , if $x \in X$ and $z \in Z$ then $\text{dist}(x, Z) = \text{dist}(x + z, Z)$.

Proof. Note that $\text{dist}(x + z, Z) = \inf\{\|(x + z) - z_0\| \mid z_0 \in Z\} = \inf\{\|x - (z_0 - z)\| \mid z_0 \in Z\}$. Since $\{z_0 - z \mid z_0 \in Z\} = Z$, we have that $\inf\{\|x - z\| \mid z \in Z\} = \inf\{\|(x + z) - z_0\| \mid z_0 \in Z\}$; i.e., $\text{dist}(x, Z) = \text{dist}(x + z, Z)$. $\square$

Exercise 1.2.7. Let X and Y be normed linear spaces and let $\varphi : X \rightarrow Y$ be a linear operator. Show that φ is continuous if and only if it is continuous at 0.

Proof. $\square$

§ 1.3. Linear Operators

Definition 1.3.1. Let X and Y be normed linear spaces. A map $T : X \rightarrow Y$ is linear if $T(x + \alpha y) = T(x) + \alpha T(y)$ for all $x, y \in X$ and $\alpha \in \mathbb{F}$. T is called a linear transformation (or linear operator). If X is a normed linear space over $\mathbb{F}$, then a linear transformation $\varphi : X \rightarrow \mathbb{F}$ is called a linear functional.

Remark. The norm on $\mathbb{F}$ is the usual absolute value in $\mathbb{R}$ or modulus operator in $\mathbb{C}$.

Example 1.3.1. Linear operators between $\mathbb{R}^n$ and $\mathbb{R}^m$ correspond to $n \times m$ matrices.

Example 1.3.2. Consider the normed linear space $(C([0, 1]), \|\cdot\|_\infty)$. Evaluation at zero is a linear functional, as well as the map $x \mapsto \int_0^1 x(t) dt$. Moreover, $T : C([0, 1]) \rightarrow C([0, 1])$ defined by $x(t) \mapsto \int_0^1 K(s, t)x(s) ds$ is a linear transformation, where K is a continuous function on $[0, 1] \times [0, 1]$.

Definition 1.3.2. Let X and Y be normed linear spaces and $T : X \rightarrow Y$ a linear transformation. T is bounded if there exists $M > 0$ such that for all $x \in X$, $\|x\| \leq 1$ implies $\|Tx\| \leq M$.

Theorem 1.3.1. Let X and Y be normed linear spaces. A linear transformation $T : X \rightarrow Y$ is bounded if and only if it is continuous.

Proof. Suppose T is continuous. Then T is continuous at zero. Choose $\delta > 0$ such that $\|x\| < \delta$ implies $\|Tx\| < 1$. Now let $x \in X$ with $\|x\| \leq 1$. Then $\|Tx\| = \left\| T\left(\frac{x\delta}{\delta}\right) \right\| = \frac{1}{\delta} \|T(\delta x)\| \leq \frac{1}{\delta}$. Therefore T is bounded.

Conversely, let $M > 0$ satisfy $\|Tx\| \leq M$ for all $x \in \overline{B}(0, 1)$. Let $\epsilon > 0$. Choose $\delta = \frac{\epsilon}{M}$. Then $\|x\| \leq \delta$ implies $\|\frac{M}{\epsilon}x\| \leq 1$. Hence by our assumption $\|T(\frac{M}{\epsilon}x)\| \leq M$, which is equivalent to $\|Tx\| \leq \epsilon$. Thus T is continuous at 0, hence it is continuous everywhere. $\square$

Example 1.3.3. Consider $D : C^1([0, 1]) \rightarrow C([0, 1])$ given by $Df = f'$. Consider the sequence of functions $\{f_n\}_{n=1}^\infty \subseteq C^1([0, 1])$ given by $f_n(x) = \frac{1}{\sqrt{n}}x^n$. Then for all $n \in \mathbb{N}$, $Df_n(x) = \sqrt{n}x^{n-1}$. Hence D is unbounded.

Theorem 1.3.2. Let X be a normed linear space. A linear functional $\varphi : X \rightarrow \mathbb{F}$ is continuous if and only if its kernel is closed.

Proof. Suppose φ is continuous. Then $\ker \varphi = \varphi^{-1}(\{0\})$ is closed, as the continuous preimage of a closed set is closed.

Conversely, suppose towards contradiction φ is not continuous. Then φ is unbounded; i.e., there exists a sequence $\{x_n\}_{n=1}^\infty \subset X$ such that $\|x_n\| = 1$ for all $n \in \mathbb{N}$ and $|\varphi(x_n)| \rightarrow \infty$. If $\ker \varphi = X$ then there is nothing to show. Assume $\ker \varphi \neq X$ and pick some $x \in X \setminus \ker \varphi$. Define $y_n := x - \frac{\varphi(x)}{\varphi(x_n)}x_n$. Then $\varphi(y_n) = \varphi(x) - \frac{\varphi(x)}{\varphi(x_n)}\varphi(x_n) = 0$. Hence $y_n \in \ker \varphi$ for all $n \in \mathbb{N}$. But note that $\|y_n - x\| = \left\| -\frac{\varphi(x)}{\varphi(x_n)}x_n \right\| = \frac{|\varphi(x)|}{|\varphi(x_n)|}\|x_n\| \rightarrow 0$. Hence $y_n \rightarrow x$. Since $\ker \varphi$ is closed, $x \in \ker \varphi$. This is a contradiction, establishing the proof. $\square$

Theorem 1.3.3. Let X be a finite dimensional normed linear space and Y a normed linear space. If $T : X \rightarrow Y$ is linear, then T is continuous.

Proof. Pick a basis for X such that $X = \text{span}\{b_1, \dots, b_n\}$. Define $M := \max_{i=1}^n \{T(b_i)\}$. For each $j = 1, \dots, n$, define a linear functional $\varphi_j : X \rightarrow \mathbb{F}$ by:

$$\begin{aligned} \varphi_j(x) &= \varphi_j\left(\sum_{i=1}^n \alpha_i b_i\right) \\ &= \sum_{i=1}^n \alpha_i \varphi_j(b_i) \\ &:= \alpha_j \end{aligned}$$

Since all subspaces of X are closed by Corollary 1.2.2, surely $\ker \varphi_j$ is closed. Hence by Theorem 1.3.2 φ_j is continuous. Since φ_j is continuous, find $L > 0$ sufficiently large enough

such that for every $x \in X$ with $\|x\| \leq 1$, we have $|\varphi_i(x)| \leq L$. Since $T : X \rightarrow Y$ is linear:

$$\begin{aligned} \|Tx\| &= \left\| T \sum_{i=1}^n \alpha_i b_i \right\| \\ &= \left\| \sum_{i=1}^n \alpha_i T b_i \right\| \\ &= \left\| \sum_{i=1}^n \varphi_i(x) T b_i \right\| \\ &\leq \sum_{i=1}^n |\varphi_i(x)| \|T b_i\| \\ &\leq nLM \\ &< \infty. \end{aligned}$$

Thus T is continuous. □

Theorem 1.3.4. *Let X be a finite dimensional vector space. All norms on X are equivalent.*

Proof. Let $\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ be two norms. Consider $\text{id} : (X, \|\cdot\|_\alpha) \rightarrow (X, \|\cdot\|_\beta)$. This map is surely linear, hence by Theorem 1.3.3 id is continuous. Hence we can find some $A > 0$ sufficiently large such that, for all $x \in X$ with $\|x\|_\alpha \leq 1$, we have $\|x\|_\beta \leq A$. Thus:

$$\begin{aligned} \|x\|_\beta &= \left\| \|x\|_\alpha \frac{x}{\|x\|_\alpha} \right\|_\beta \\ &= \|x\|_\alpha \left\| \frac{x}{\|x\|_\alpha} \right\|_\beta \\ &\leq \|x\|_\alpha A. \end{aligned}$$

Applying the same argument to id^{-1} gives $\|x\|_\alpha \leq \|x\|_\beta B$, establishing the theorem. □

§ 1.4. Bounded Linear Functionals

Definition 1.4.1. Let X and Y be normed linear spaces.

- (1) The space of bounded linear operators between X and Y is denoted $\mathcal{L}(X, Y) := \{T : X \rightarrow Y \mid T \text{ is bounded}\}$.
- (2) The operator norm of a bounded linear operator $T \in \mathcal{L}(X, Y)$ is $\|T\|_{\mathcal{L}(X, Y)} := \sup\{\|Tx\|_Y \mid x \in \overline{B}_X(0, 1)\}$.
- (3) The continuous dual of X is defined as $X^* := \mathcal{L}(X, \mathbb{F})$.

Remark. We denote $\mathcal{L}(X, X)$ by $\mathcal{L}(X)$.

Proposition 1.4.1. *If $T \in \mathcal{L}(X, Y)$ and $x \in X$, then $\|Tx\| \leq \|T\|_{\mathcal{L}(X, Y)} \cdot \|x\|$.*

Proof. If $x = 0$, then there is nothing to show. If $x \neq 0$, then:

$$\begin{aligned}\|Tx\| &= \left\| \|x\| T\left(\frac{x}{\|x\|}\right) \right\| \\ &= \left\| T\left(\frac{x}{\|x\|}\right) \right\| \|x\| \\ &\leq \|T\|_{\mathcal{L}(X,Y)} \|x\|. \quad \square\end{aligned}$$

Proposition 1.4.2. *If $S, T \in \mathcal{L}(X)$, then $S \circ T \in \mathcal{L}(X)$.*

Proof. Let $x \in X$ such that $\|x\| \leq 1$. By Proposition 1.4.1:

$$\begin{aligned}\|(S \circ T)(x)\| &\leq \|S(T(x))\| \\ &\leq \|S\|_{\mathcal{L}(X)} \|Tx\| \\ &\leq \|S\|_{\mathcal{L}(X)} \|T\|_{\mathcal{L}(X)} \|x\| \\ &\leq \|S\|_{\mathcal{L}(X)} \|T\|_{\mathcal{L}(X)}.\end{aligned}$$

Taking the supremum over all $\|x\| \leq 1$ gives $\|S \circ T\|_{\mathcal{L}(X)} \leq \|S\|_{\mathcal{L}(X)} \|T\|_{\mathcal{L}(X)}$. $\square$

Theorem 1.4.1. *Let X be a normed linear space and Y a Banach space. Then $(\mathcal{L}(X, Y), \|\cdot\|_{\mathcal{L}(X,Y)})$ is Banach.*

Proof. Let $\{T_n\}_{n=1}^\infty \subseteq \mathcal{L}(X, Y)$ be Cauchy. Note that for any $x \in X$, $\{T_n(x)\}_{n=1}^\infty$ will be Cauchy in Y . Since Y is complete, this sequence has a limit. Define $Tx := \lim_{n \rightarrow \infty} T_n x$. We must show that T is linear, bounded, and $T_n \rightarrow T$.

Let $x, y \in X$, $\alpha \in \mathbb{F}$, and $n \in \mathbb{N}$. Then $T_n(x + \alpha y) = T_n x + \alpha T_n y$. By limit laws, $T(x + \alpha y) = Tx + \alpha Ty$. Thus T is linear.

Since $\{T_n\}_{n=1}^\infty$ is Cauchy in $\mathcal{L}(X, Y)$, this means $\{T_n\}_{n=1}^\infty$ is bounded. Hence we can find $M > 0$ sufficiently large such that $\|T_n\| \leq M$ for all $n \in \mathbb{N}$. For every $x \in X$ with $\|x\| \leq 1$, we have:

$$\begin{aligned}\|Tx\| &= \left\| \lim_{n \rightarrow \infty} T_n x \right\| \\ &= \lim_{n \rightarrow \infty} \|T_n x\| \\ &\leq \lim_{n \rightarrow \infty} \|T_n\|_{\mathcal{L}(X,Y)} \|x\| \\ &\leq \lim_{n \rightarrow \infty} \|T_n\|_{\mathcal{L}(X,Y)} \\ &\leq M.\end{aligned}$$

Thus T is bounded.

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ large such that $n, m \geq N$ implies $\|T_n - T_m\|_{\mathcal{L}(X,Y)} < \epsilon$. Let $x \in X$ such that $\|x\| \leq 1$. Then:

$$\begin{aligned}\|T_n x - T_m x\| &\leq \|(T_n - T_m)x\| \\ &\leq \|T_n - T_m\|_{\mathcal{L}(X,Y)} \|x\| \\ &\leq \|T_n - T_m\|_{\mathcal{L}(X,Y)} \\ &< \epsilon.\end{aligned}$$

Take $m \rightarrow \infty$. Then $\|T_n x - Tx\| \leq \epsilon$. Hence $\|(T_n - T)x\| \leq \epsilon$. Taking the supremum over all $x \in \overline{B}_X(0, 1)$ gives $\|T_n - T\|_{\mathcal{L}(X, Y)} \leq \epsilon$. Thus $T_n \rightarrow T$ in the operator norm. $\square$

Corollary 1.4.1 (Neumann's Theorem). *Let X be a Banach space and $T \in \mathcal{L}(X)$. If $\|T\| < 1$, then $I - T$ is invertible, and $(I - T)^{-1} = \sum_{n=0}^{\infty} T^n$.*

Proof. $\square$

Question. For any normed linear space X , is X^* nonempty? How can one construct an arbitrary bounded linear functional on X ?

Lemma 1.4.1 (Zorn's Lemma). *In a partially ordered set, if every chain has a maximal element, then the partially ordered set has a maximal element.*

Theorem 1.4.2 (Hahn-Banach). *Let X be a $\mathbb{R}$ -vector space and $p : X \rightarrow \mathbb{R}$ a map satisfying:*

$$\begin{aligned} p(x + y) &\leq p(x) + p(y), \\ p(\lambda x) &= \lambda p(x), \text{ provided } \lambda > 0. \end{aligned}$$

Let $Y \subseteq X$ be a subspace and $\varphi : Y \rightarrow \mathbb{R}$ a linear functional with:

$$\varphi(y) \leq p(y) \text{ for all } y \in Y.$$

Then there exists a linear functional $\Phi : X \rightarrow \mathbb{R}$ such that $\Phi|_Y = \varphi$ and:

$$\Phi(x) \leq p(x) \text{ for all } x \in X.$$

Proof. $\square$

Corollary 1.4.2. *Let X be a normed linear space and $Y \subseteq X$ a subspace. If $\varphi \in Y^*$, then there exists an extension $\Phi \in X^*$ such that $\|\varphi\|_{Y^*} = \|\Phi\|_{X^*}$.*

Proof. Define $p : X \rightarrow \mathbb{R}$ by $p(x) := \|\varphi\|_{Y^*} \|x\|$. Clearly $p(x + y) \leq p(x) + p(y)$ and $p(\lambda x) = \lambda p(x)$ provided $\lambda > 0$. Moreover:

$$\begin{aligned} p(y) &= \|\varphi\|_{Y^*} \|y\| \\ &\geq |\varphi(y)| \\ &\geq \varphi(y). \end{aligned}$$

By the Hahn-Banach Theorem, there exists an extension $\Phi : X \rightarrow \mathbb{R}$ such that $\Phi|_Y = \varphi$ and $\Phi(x) \leq p(x)$. From this, we have the following inclusion of sets:

$$\left\{ \left| \varphi\left(\frac{y}{\|y\|}\right) \right| \mid y \in Y, y \neq 0 \right\} \subseteq \left\{ \left| \Phi\left(\frac{x}{\|x\|}\right) \right| \mid x \in X, x \neq 0 \right\}.$$

Hence $\|\varphi\|_{Y^*} \leq \|\Phi\|_{X^*}$. Conversely, by our construction of p we have:

$$\begin{aligned} |\Phi(x)| &\leq |p(x)| \\ &\leq \|\varphi\|_{Y^*} \|x\|. \end{aligned}$$

Rearranging gives $\left| \Phi\left(\frac{x}{\|x\|}\right) \right| \leq \|\varphi\|_{Y^*}$. Taking the supremum over all $x \in X$, $x \neq 0$ gives $\|\Phi\|_{X^*} \leq \|\varphi\|_{Y^*}$. Having established equality, we've finally shown $\Phi \in X^*$. $\square$

Corollary 1.4.3. *Let X be a normed linear space and $Y \subseteq X$ a subspace. If $w \in X \setminus Y$ such that $\text{dist}(w, Y) > 0$, then there exists a bounded linear functional $\varphi \in X^*$ such that $\varphi(w) = 1$, $\varphi(y) = 0$ for all $y \in Y$, and $\|\varphi\|_{X^*} = \frac{1}{\text{dist}(w, Y)}$.*

Proof. Define $\varphi : \text{span}\{Y, \{w\}\} \rightarrow \mathbb{R}$ by $\varphi(y + \lambda w) = \lambda$. This is a linear functional, and:

$$\begin{aligned} \|\varphi\| &= \sup \left\{ \left| \varphi\left(\frac{y + \lambda w}{\|y + \lambda w\|}\right) \right| : y \in Y, \lambda \in \mathbb{R}, y + \lambda w \neq 0 \right\} \\ &= \sup \left\{ \frac{|\lambda|}{\|y + \lambda w\|} : y \in Y, \lambda \in \mathbb{R}, y + \lambda w \neq 0 \right\} \\ &= \sup \left\{ \frac{1}{\left\| \frac{y}{\lambda} + w \right\|} : y \in Y, \lambda \in \mathbb{R}, y + \lambda w \neq 0 \right\} \\ &= \sup \left\{ \frac{1}{\|w - y\|} : y \in Y, \lambda \in \mathbb{R}, y + \lambda w \neq 0 \right\}^1 \\ &= \frac{1}{\inf\{\|w - y\| \mid y \in Y\}}^2 \\ &= \frac{1}{\text{dist}(w, Y)}. \end{aligned}$$

Applying Corollary 1.4.2 establishes the proof. $\square$

Remark. The previous corollary establishes that if $X = \{0\}$, then $X^* \neq \{0\}$.

Exercises

Exercise 1.4.1. *Let X be a normed linear space. Prove the limit laws: if $x_n \rightarrow x$, $y_n \rightarrow y$, and $\lambda \in \mathbb{F}$, then $x_n + \lambda y_n \rightarrow x + \lambda y$.*

Proof. $\square$

Exercise 1.4.2. *Let X be a normed linear space with norm $\|\cdot\|$. Prove that the map $\|\cdot\| : X \rightarrow \mathbb{R}$ is continuous.*

¹Since $\{\frac{y}{\lambda} \mid y \in Y, \lambda \in \mathbb{R}\} = Y$.

²Let $x_n \rightarrow \inf A$. Then $\frac{1}{x_n} \rightarrow \frac{1}{\inf A}$. Since $\frac{1}{\inf A}$ is an upper-bound and an adherent point of $\frac{1}{A}$, we have $\frac{1}{\inf A} = \sup \frac{1}{A}$.

Proof. Let $x, y \in X$. Then $\|x\| \leq \|x - y\| + \|y\|$, giving $\|x\| - \|y\| \leq \|x - y\|$. A similar tactic gives $\|y\| - \|x\| \leq \|x - y\|$. Hence $|\|x\| - \|y\|| \leq \|x - y\|$. This means $\|\cdot\|$ is Lipschitz, implying that it is continuous. $\square$

Exercise 1.4.3. Prove that the closure of a linear subspace in a normed linear space is also a subspace.

Proof. Let X be a normed linear space and Y a subspace. Let $u, v \in \bar{Y}$ and $\alpha \in \mathbb{F}$. There exists sequences $\{u_n\}_{n=1}^\infty, \{v_n\}_{n=1}^\infty$ such that $u_n \rightarrow u$ and $v_n \rightarrow v$. Hence $u_n + \alpha v_n \rightarrow u + \alpha v \in \bar{Y}$. Thus $\bar{Y}$ is a linear subspace. $\square$

Exercise 1.4.4. Prove that the operator norm has the three properties required of a norm

Proof. □

Exercise 1.4.5. Prove directly that if T is an unbounded linear operator, then it is discontinuous at 0.

Proof. Let $T : X \rightarrow Y$ be an unbounded linear operator. Since T is unbounded:

$$\|T\|_{\mathcal{L}(X,Y)} = \sup\{\|Tx\|_Y \mid x \in \bar{B}_X(0,1)\} = \infty.$$

We can find a sequence $\{x_n\}_{n=1}^\infty$ such that $\|x_n\|_X \leq 1$ and $\|Tx_n\|_Y \rightarrow \infty$. Consider the sequence $\left\{\frac{x_n}{\|Tx_n\|_Y}\right\}_{n=1}^\infty$. Clearly $\frac{x_n}{\|Tx_n\|_Y} \rightarrow 0$. But:

$$\left\|T\left(\frac{x_n}{\|Tx_n\|_Y}\right)\right\| = \frac{\|Tx_n\|_Y}{\|Tx_n\|_Y} = 1.$$

Hence the sequence $\left\{T\left(\frac{x_n}{\|Tx_n\|_Y}\right)\right\}_{n=1}^\infty$ does not converge to 0. Thus T is discontinuous at 0. $\square$

Exercise 1.4.6. Prove that the norm of a linear transformation is the infimum of all numbers M that satisfy the inequality $\|Tx\| \leq M\|x\|$ for all x .

Proof. Let M be such that $\|Tx\| \leq M\|x\|$ for all $x \in X$. Then $\frac{\|Tx\|}{\|x\|} \leq M$ for all $x \in X, x \neq 0$. Taking the supremum over all $x \in X, x \neq 0$ gives $\|T\|_{\mathcal{L}(X,Y)} \leq M$. Hence $\|T\|_{\mathcal{L}(X,Y)} \leq \inf\{M \mid \|Tx\| \leq M\|x\| \text{ for all } x \in X\}$.

Conversely, let $x \in X, x \neq 0$. Then $\|Tx\| \leq \|T\|_{\mathcal{L}(X,Y)}\|x\|$. Thus $\|T\|_{\mathcal{L}(X,Y)} \in \{M \mid \|Tx\| \leq M\|x\| \text{ for all } x \in X\}$. Surely $\|T\|_{\mathcal{L}(X,Y)} \geq \inf\{M \mid \|Tx\| \leq M\|x\| \text{ for all } x \in X\}$, establishing equality. $\square$

Exercise 1.4.7. On the space $C([0,1])$ we define "point-evaluation functionals" by $t^*(f) = f(t)$. Here $t \in [0,1]$ and $f \in C([0,1])$. Prove that $\|t^*\| = 1$. Prove that if $\varphi = \sum_{i=1}^n \lambda_i t_i^*$, where $t_1, t_2, \dots, t_n$ are distinct points in $[0,1]$, then $\|\varphi\| = \sum_{i=1}^n |\lambda_i|$.

Proof.

□

Exercise 1.4.8. Prove that if a linear transformation maps some nonvoid open set of the domain space to a bounded set in the range space, then it is continuous.

Proof. Let $T : X \rightarrow Y$ be linear. Let $U \subseteq X$ be open and nonvoid. Then there exists some $x \in U$. Find $r > 0$ such that $B(x, r) \subseteq U$.

Let $x_0 \in B(0, 1)$. Then $x + rx_0 \in B(x, r) \subseteq U$. Hence $T(x + rx_0) \in T(U)$. We can write:

$$\|Tx_0\| \leq \frac{\|T(x + rx_0)\| + \|Tx\|}{r}.$$

Since $T(U)$ is bounded, it must be the case that $\|Tx_0\| \leq M$ for some $M > 0$. Thus $T(B(0, 1))$ is bounded.

Now let $x \in X$. We have that:

$$\begin{aligned} \|Tx\| &= \left\| T\left(\frac{x}{2\|x\|} \cdot 2\|x\|\right) \right\| \\ &= 2 \left\| T\left(\frac{x}{2\|x\|}\right) \right\| \|x\| \\ &\leq 2M \|x\|. \end{aligned}$$

Thus T is continuous. □

Exercise 1.4.9. Let Y be a linear subspace in a normed linear space X . For a fixed $x \in X$, prove that:

$$\text{dist}(x, Y) = \sup\{\varphi(x) \mid \varphi \in X^*, \varphi|_Y \equiv 0, \|\varphi\| = 1\}.$$

Proof. Let $\varphi \in X^*$ such that $\varphi|_Y = 0$ and $\|\varphi\| = 1$. Observe that:

$$\begin{aligned} \varphi(x) &= \varphi(x) - \varphi(y) \\ &= \varphi(x - y) \\ &\leq \|\varphi\| \|x - y\| \\ &= \|x - y\|. \end{aligned}$$

Taking the infimum over all $y \in Y$ gives $\varphi(x) \leq \text{dist}(x, Y)$. Taking the supremum over all $\varphi \in X^*$ such that $\varphi|_Y = 0$ and $\|\varphi\| = 1$ gives $\sup\{\varphi(x) \mid \varphi \in X^*, \varphi|_Y = 0, \|\varphi\| = 1\} \leq \text{dist}(x, Y)$.

For the converse direction, we proceed by cases on $\text{dist}(x, Y)$. If $\text{dist}(x, Y) = 0$, then by the above inequality we have:

$$\begin{aligned} 0 &= \text{dist}(x, Y) \\ &\geq \sup\{\varphi(x) \mid \varphi \in X^*, \varphi|_Y = 0, \|\varphi\| = 1\} \\ &\geq 0. \end{aligned}$$

Hence $\text{dist}(x, Y) = \sup\{\varphi(x) \mid \varphi \in X^*, \varphi|_Y = 0, \|\varphi\| = 1\}$. Now suppose $\text{dist}(x, Y) > 0$. By Corollary 1.4.3 there exists a linear functional $\psi \in X^*$ such that $\psi(x) = 1$, $\psi|_Y = 0$, and

$\|\psi\| = \frac{1}{\text{dist}(x, Y)}$. Define a linear functional $\Phi : X \rightarrow \mathbb{F}$ by $\Phi(x_0) = \text{dist}(x_0, Y)\psi(x_0)$. Note that $\Phi|_Y = 0$ and:

$$\|\Phi\| = \text{dist}(x, Y)$$

$$\|\Phi\| = \text{dist}(x, Y)$$

□

§ 1.5. ℓ^p Spaces

Definition 1.5.1.

(1) Let $1 \leq p < \infty$. We define $\ell^p := \{x : \mathbb{N} \rightarrow \mathbb{F} \mid \sum_{j=1}^{\infty} |x_j|^p < \infty\}$. Likewise, define

$$\|\cdot\|_{\ell^p} : \ell^p \rightarrow \mathbb{F} \text{ by } \|x\|_{\ell^p} = \left(\sum_{j=1}^{\infty} |x_j|^p \right)^{\frac{1}{p}}.$$

(2) Let $p = \infty$. We define $\ell^\infty := \{x : \mathbb{N} \rightarrow \mathbb{F} \mid \sup_{n \in \mathbb{N}} |x_n| < \infty\}$. Likewise, define

$$\|\cdot\|_{\ell^\infty} : \ell^\infty \rightarrow \mathbb{F} \text{ by } \|x\|_{\ell^\infty} = \sup_{n \in \mathbb{N}} |x_n|.$$

Lemma 1.5.1. Let $a, b \geq 0$ and $0 \leq t \leq 1$. Then $((1-t)a + tb)^p \leq (1-t)a^p + tb^p$.

Proof.

□

Lemma 1.5.2. Let $a, b \geq 0$ and $0 \leq t \leq 1$. Then $(a+b)^p \leq 2^{p-1}(a^p + b^p)$.

Proof.

□

Lemma 1.5.3 (Minkowski's Inequality). Let $1 \leq p \leq \infty$. For any $x, y \in \ell^p$, we have $\|x + y\|_{\ell^p} \leq \|x\|_{\ell^p} + \|y\|_{\ell^p}$.

Theorem 1.5.1. For any $1 \leq p < \infty$, $(\ell^p, \|\cdot\|_{\ell^p})$ is a Banach space.

Proof. Lemma 1.5.1 establishes that ℓ^p is a vector space. Lemma 1.5.3 established that $\|\cdot\|_{\ell^p}$ satisfies the triangle inequality; the remaining norm axioms are routine to check. Hence $(\ell^p, \|\cdot\|_{\ell^p})$ is a normed linear space.

Let $\{f_n\}_{n=1}^{\infty} \subseteq \ell^p$ be a Cauchy sequence. For any $j, n, m \in \mathbb{N}$, we have:

$$\begin{aligned} |f_n(j) - f_m(j)| &\leq (|f_n(j) - f_m(j)|^p)^{\frac{1}{p}} \\ &\leq \left(\sum_{k=1}^{\infty} |f_n(k) - f_m(k)|^p \right)^{\frac{1}{p}} \\ &= \|f_n - f_m\|_{\ell^p}. \end{aligned}$$

Hence $\{f_n(j)\}_{n=1}^{\infty} \subseteq \mathbb{F}$ is Cauchy. Since $\mathbb{F}$ is complete, the limit of this sequence exists. Define $f(j) := \lim_{n \rightarrow \infty} f_n(j)$. We will show that $f \in \ell^p$ and that $f_n \rightarrow f$ in ℓ^p .

Find $N \in \mathbb{N}$ sufficiently large such that for all $n \geq N$, we have $\|f_n - f_N\|_{\ell^p} \leq 1$. Taking p^{th} powers on both sides gives the following statement: for all $n \geq N$ we have $\sum_{j=1}^{\infty} |f_n(j) - f_N(j)|^p \leq 1$. Now fix $J \in \mathbb{N}$. For all $n \geq N$ we have $\sum_{j=1}^J |f_n(j) - f_N(j)|^p \leq 1$. Take $n \rightarrow \infty$. Then we have $\sum_{j=1}^J |f(j) - f_N(j)|^p \leq 1$. In particular, for all $J \in \mathbb{N}$ we have:

$$\begin{aligned} \sum_{j=1}^J |f(j)|^p &= \sum_{j=1}^J |f(j) - f_N(j) + f_N(j)|^p \\ &\leq \sum_{j=1}^J 2^{p-1} (|f(j) - f_N(j)|^p + |f_N(j)|^p) \quad \text{Lemma 1.5.2} \\ &= 2^{p-1} \left(\sum_{j=1}^J |f(j) - f_N(j)|^p + \sum_{j=1}^J |f_N(j)|^p \right) \\ &\leq 2^{p-1} (1 + \|f_N\|_{\ell^p}^p) \\ &< \infty. \end{aligned}$$

By the Monotone Convergence Theorem, $\sum_{j=1}^{\infty} |f(j)|^p < \infty$. Hence $f \in \ell^p$.

Now let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that $\|f_n - f_N\|_{\ell^p} \leq \epsilon$. Then for all $n \geq N$ and for all $j \in \mathbb{N}$, we have $|f_n(j) - f_N(j)|^p \leq \epsilon^p$. Take $n \rightarrow \infty$. Then for all $J \in \mathbb{N}$ we have $\sum_{j=1}^J |f(j) - f_N(j)|^p \leq \epsilon^p$. Using the Monotone Convergence Theorem, take $J \rightarrow \infty$. Then $\sum_{j=1}^{\infty} |f(j) - f_N(j)|^p \leq \epsilon^p$. Therefore $\|f - f_N\|_{\ell^p} \leq \epsilon$. $\square$

Definition 1.5.2. The subspace of ℓ^∞ consisting of all convergent sequences is denoted $c := \{x \in \ell^\infty \mid x \text{ converges}\}$. The subspace of c consisting of all sequences which converge to zero is denoted $c_0 := \{x \in c \mid x \text{ converges to } 0\}$. The subspace of ℓ^∞ consisting of all sequences with only finitely many nonzero terms is denoted $c_{00} := \{x \in \ell^\infty \mid \text{supp}(x) < \infty\}$.

Definition 1.5.3. The standard coordinate vector in an infinite dimensional vector space is denoted $e^n := (0, \dots, 0, 1, 0, \dots)$; i.e., it is a sequence which is "1" in the n^{th} coordinate and "0" elsewhere.

Proposition 1.5.1. Let $y \in \ell^\infty$. Define $\varphi_y : \ell^1 \rightarrow \mathbb{F}$ by $\varphi_y(x) = \sum_{k=1}^{\infty} x_k y_k$. Then $\varphi \in (\ell^1)^*$ and $\|\varphi_y\|_{\ell^1}^* = \|y\|_{\ell^\infty}$.

Exercises

Exercise 1.5.1. Show that $(\ell^\infty, \|\cdot\|_\infty)$ is complete.

Proof.

$\square$

Exercise 1.5.2. Show that c and c_0 are closed in ℓ^∞ . Show that c_{00} is not closed by exhibiting a sequence in c_{00} which converges to a point $x \in \ell^\infty$ but is not in c_{00} .

Proof.

□

Exercise 1.5.3. Let $y \in \ell^\infty$ and define the linear functional:

$$\varphi_y : \ell^1 \rightarrow \mathbb{F}, \quad \varphi_y(x) := \sum_{j=1}^{\infty} x_j y_j.$$

Show that $\|\varphi_y\|_{(\ell^1)^*} = \|y\|_{\ell^\infty}$.*Proof.* Note that:

$$\begin{aligned} \sum_{i=1}^{\infty} |x_i y_i| &= \sum_{i=1}^{\infty} |x_i| |y_i| \\ &\leq \|y\|_{\ell^\infty} \sum_{i=1}^{\infty} |x_i| \\ &\leq \|y\|_{\ell^\infty} \|x\|_{\ell^1}. \end{aligned}$$

Hence:

$$\begin{aligned} |\varphi_y(x)| &= \left| \sum_{i=1}^{\infty} x_i y_i \right| \\ &\leq \sum_{i=1}^{\infty} |x_i y_i| \\ &\leq \|y\|_{\ell^\infty} \|x\|_{\ell^1}. \end{aligned}$$

Dividing by $\|x\|_{\ell^1}$ and taking the supremum over all $x \in \ell^1$, $x \neq 0$ gives the inequality $\|\varphi_y\|_{(\ell^1)^*} \leq \|y\|_{\ell^\infty}$. Hence $\varphi_y \in (\ell^1)^*$.For the converse direction, note that $e_j \in B_{\ell^1}(0, 1)$. Hence for any $j \in \mathbb{N}$, we have $|y_j| \leq |\varphi_y(e_j)| \leq \|\varphi_y\|_{(\ell^1)^*}$. Taking the supremum over all $j \in \mathbb{N}$ gives $\|y\|_{\ell^\infty} \leq \|\varphi_y\|_{(\ell^1)^*}$. Thus $\|\varphi_y\|_{(\ell^1)^*} = \|y\|_{\ell^\infty}$. □**§ 1.6. Consequences of The Baire Category Theorem****Theorem 1.6.1** (Baire Category Theorem). Let X be a complete metric space. If $\{U_n\}_{n=1}^{\infty}$ is a family of open and dense subsets of X , then $\bigcap_{n=1}^{\infty} U_n$ is still dense.*Proof.*

□

Corollary 1.6.1. Let X be a complete metric space and $\{Z_n\}_{n=1}^{\infty}$ a family of closed sets. If $X = \bigcup_{n=1}^{\infty} Z_n$, then there exists some $i \in \mathbb{N}$ such that Z_i has non-empty interior.

Proof.

□

Definition 1.6.1. Let X be a metric space.

- (1) A subset Y of X is called nowhere dense if $\bar{Y}$ has empty interior.
- (2) A subset of X is called Category I if it is the countable union of nowhere dense sets.
- (3) A subset of X which is not Category I is called Category II.

Remark. Theorem 1.6.1 and Corollary 1.6.1 can be rephrased as the assertion that every complete metric space is Category II.

Theorem 1.6.2 (Banach-Steinhouse). *Let X be a Banach space and Y a normed linear space. Let $\{T_\alpha\}_{\alpha \in A} \subset \mathcal{L}(X, Y)$ be an arbitrary family of bounded linear operators. Then $\sup\{\|T_\alpha\| \mid \alpha \in A\} < \infty$ if and only if $\{x \mid \sup_{\alpha \in A} \|T_\alpha x\| < \infty\}$ is Category II*

Proof. Suppose $C := \sup\{\|T_\alpha\| \mid \alpha \in A\} < \infty$. Then for all $x \in X$ and $\alpha \in A$, we have $\|T_\alpha x\|_Y \leq \|T_\alpha\| \|x\|_X \leq C \|x\|$. Hence $\{x \mid \sup_{\alpha \in A} \|T_\alpha x\| < \infty\} = X$, which is Category II by the Baire Category Theorem.

Conversely, suppose $F := \{x \mid \sup_{\alpha \in A} \|T_\alpha x\| < \infty\}$ is Category II. Define $F_n := \{x \mid \sup_{\alpha \in A} \|T_\alpha x\| \leq n\}$. We will show that F_n is closed in X . Note that:

$$F_n = \bigcap_{\alpha \in A} \{x \mid \|T_\alpha x\| \leq n\}.$$

Note that $\|\cdot\|$ and T_α are continuous, hence the composition $g_\alpha := \|T_\alpha(\cdot)\|$ is continuous. The above expression then becomes:

$$\begin{aligned} F_n &= \bigcap_{\alpha \in A} \{x \mid \|T_\alpha x\| \leq n\} \\ &= \bigcap_{\alpha \in A} \{x \mid g_\alpha(x) \leq n\} \\ &= \bigcap_{\alpha \in A} \{x \mid g_\alpha(x) \in [0, n]\} \\ &= \bigcap_{\alpha \in A} g_\alpha^{-1}([0, n]). \end{aligned}$$

Hence F_n is closed.

Since $F = \bigcup_{n=1}^{\infty} F_n$, by Corollary 1.6.1 there exists some $n \in \mathbb{N}$ such that F_n has nonempty interior. Choose $x_0 \in X$ and $r > 0$ such that $\bar{B}(x_0, r) \subset F_n$. Let $x \in X$ such that $\|x\| \leq 1$. Fix $\alpha \in A$. Note that $x_0 + rx \in \bar{B}(x_0, r)$, hence:

$$\begin{aligned} \|T_\alpha x\|_Y &= \left\| T_\alpha \left(\frac{1}{r} (x_0 + rx - x_0) \right) \right\|_Y \\ &\leq \frac{1}{r} (\|T_\alpha(x_0 + rx)\|_Y + \|T_\alpha x_0\|_Y) \\ &\leq \frac{2n}{r} \\ &< \infty. \end{aligned}$$

Taking the supremum over all α and $\|x\| \leq 1$ gives $\sup\{\|T_\alpha\| \mid \alpha \in A\} < \infty$. $\square$

Corollary 1.6.2 (Uniform Boundedness Principle). *Let X be a Banach space and Y a normed linear space. Let $\{T_\alpha\}_{\alpha \in A} \subset \mathcal{L}(X, Y)$ be an arbitrary family of bounded linear operators. If $\sup_{\alpha \in A} \|T_\alpha x\|_Y < \infty$ for all $x \in X$, then $\sup_{\alpha \in A} \|T_\alpha\| < \infty$.*

Proof. The set $\{x \mid \sup_{\alpha \in A} \|T_\alpha x\| < \infty\} = X$ is Category II. By Theorem 1.6.2 we have that $\sup_{\alpha \in A} \|T_\alpha\| < \infty$. $\square$

Definition 1.6.2. Let X and Y be normed linear spaces and $T : X \rightarrow Y$ linear.

- (1) The graph of T is $\text{Gr}(T) := \{(x, Tx) \mid x \in X\}$.
- (2) The map T has a closed graph if $\text{Gr}(T)$ is closed in $X \times Y$.

Remark. Regardless if $X \times Y$ is given the product or box topology, the induced metrics are equivalent.

Proposition 1.6.1. *Let X, Y be normed spaces and $T : X \rightarrow Y$ be linear. The following are equivalent:*

- (1) T has a closed graph;
- (2) If $\{x_n\}_{n=1}^\infty$ is a sequence such that $x_n \rightarrow x$ and $Tx_n \rightarrow y$, then $y = Tx$.

Proof. Suppose T has a closed graph. Let $\{x_n\}_{n=1}^\infty$ be a sequence such that $x_n \rightarrow x$ and $Tx_n \rightarrow y$. Then $\{(x_n, Tx_n)\}_{n=1}^\infty \subseteq \text{Gr}(T)$ is a sequence such that $(x_n, Tx_n) \rightarrow (x, y)$. But since T has a closed graph, $(x, y) \in \text{Gr}(T)$. Hence $y = Tx$.

Conversely, if $\{(x_n, Tx_n)\}_{n=1}^\infty \subseteq \text{Gr}(T)$ is a sequence such that $(x_n, Tx_n) \rightarrow (x, y) \in X \times Y$, then $x_n \rightarrow x$ and $Tx_n \rightarrow y$. By hypothesis $y = Tx$, hence $(x, y) \in \text{Gr}(T)$. Thus T has a closed graph. $\square$

Remark. A linear map T having a closed graph is weaker than T being continuous.

Example 1.6.1. Recall the following standard result from an introductory analysis course: let $J \subseteq \mathbb{R}$ be a bounded interval and let $\{f_n\}_{n=1}^\infty \subseteq J^\mathbb{R}$ be a sequence of functions. Suppose that there exists $x_0 \in J$ such that $\{f_n(x_0)\}_{n=1}^\infty$ converges, and that the sequence $\{f'_n\}_{n=1}^\infty$ of derivatives exists on J and converges uniformly on J to a function g . Then the sequence $\{f_n\}_{n=1}^\infty$ converges uniformly on J to a function f that has a derivative at every point of J and $f' = g$.

Indeed, define $D : C^1([0, 1]) \rightarrow C([0, 1])$ by $Df = f'$. Let $\{f_n\}_{n=1}^\infty \subseteq C^1([0, 1])$ be a sequence of functions such that $f_n \rightarrow f$ and $Df_n \rightarrow g$. Note that the convergence of $\{Df_n\}_{n=1}^\infty$ will be uniform by the compactness of $[0, 1]$. Hence by the above result, we have that $g = Df$. By Proposition 1.6.1, D has a closed graph. However, we showed in Example 1.3.3 that D is not continuous.

Definition 1.6.3. Let X, Y be normed linear spaces. A map $T : X \rightarrow Y$ is open if $T(U)$ is open in Y provided U is open in X .

Theorem 1.6.3 (Open-Mapping Theorem). *Let $T : X \rightarrow Y$ be a continuous, surjective, and linear mapping between Banach spaces. Then T is open.*

Remark. Many times throughout the following proof we will be using the fact that $B(0, n) = nB(0, 1)$.

Proof. Note that $Y = \bigcup_{n=1}^{\infty} T(nB_X(0, 1))$, hence $Y = \bigcup_{n=1}^{\infty} \overline{T(nB_X(0, 1))}$. Since Y is Banach, the Baire Category Theorem says there exists some $n \in \mathbb{N}$ such that $\overline{nT(B_X(0, 1))}$ has nonempty interior. So there exists $y_0 \in Y$ and $r > 0$ such that:

$$\begin{aligned} B_Y(y_0, r) &\subset \overline{nT(B_X(0, 1))} \\ &= \overline{nT(B_X(0, 1))^3}. \end{aligned}$$

Let $\epsilon > 0$. Since $y_0 \in B_Y(y_0, r)$, surely $y_0 \in \overline{nT(B_X(0, 1))}$. Hence there exists some $x_1 \in B_X(0, 1)$ such that $\|y_0 - nTx_1\|_Y < \epsilon$. Now let $y \in B(0, 1)$ be arbitrary. Then $y_0 + ry \in B_Y(y_0, r) \subset \overline{nT(B_X(0, 1))}$. Hence there exists $x_2 \in B_X(0, 1)$ such that $\|y_0 + ry - nTx_2\|_Y < \epsilon$. We can write:

$$\begin{aligned} y &= \frac{1}{r}(ry + y_0) - \frac{1}{r}y_0 \\ &= \frac{1}{r}(ry + y_0 - nTx_2) + \frac{n}{r}Tx_2 - \frac{1}{r}(y_0 - nTx_1) - \frac{n}{r}Tx_1 \\ &= \frac{1}{r}((y_0 + ry) - nTx_2) - \frac{1}{r}(y_0 - nTx_1) + \frac{n}{r}T(x_2 - x_1). \end{aligned}$$

We can rearrange the above equation as:

$$y - \frac{n}{r}T(x_2 - x_1) = \frac{1}{r}((y_0 + ry) - nTx_2) - \frac{1}{r}(y_0 - nTx_1).$$

Hence:

$$\begin{aligned} \left\| y - \frac{n}{r}T(x_2 - x_1) \right\| &= \left\| \frac{1}{r}((y_0 + ry) - nTx_2) - \frac{1}{r}(y_0 - nTx_1) \right\| \\ &\leq \frac{1}{r} \|((y_0 + ry) - nTx_2)\| + \frac{1}{r} \|(y_0 - nTx_1)\| \\ &< \frac{2\epsilon}{r}. \end{aligned}$$

Take $x' := x_2 - x_1 \in B(0, 2)$. Then since $\left\| y - \frac{n}{r}T(x') \right\| < \frac{2\epsilon}{r}$ can be made arbitrarily small, we have that $y \in \overline{T(B_X(0, \frac{2n}{r}))}$. Take $\lambda := \frac{2n}{r}$. Thus we've established the inclusion $B_Y(0, 1) \subset \overline{TB_X(0, \lambda)}$.

³Follows from the fact that $\overline{B(x, r)} = \overline{B}(x, r)$ and the above remark.

Let $y \in B_Y(0, 1) \subset \overline{T(B_X(0, \lambda))}$. Find $x_1 \in B_X(0, \lambda)$ such that $\|y - Tx_1\|_Y < \frac{1}{2}$. Then:

$$\begin{aligned} y - Tx_1 &\in B_Y\left(0, \frac{1}{2}\right) \\ &= \frac{1}{2}B_Y(0, 1) \\ &\subset \frac{1}{2}\overline{T(B_X(0, \lambda))} \\ &= \overline{\frac{1}{2}T(B_X(0, \lambda))} \\ &= \overline{T\left(B_X\left(0, \frac{\lambda}{2}\right)\right)}. \end{aligned}$$

Now pick $x_2 \in B_X(0, \frac{\lambda}{2})$ such that $\|y - Tx_1 - Tx_2\| < \frac{1}{2^2}$. Then:

$$\begin{aligned} y - Tx_1 - Tx_2 &\in B_Y\left(0, \frac{1}{2^2}\right) \\ &= \frac{1}{2^2}B_Y(0, 1) \\ &\subset \frac{1}{2^2}\overline{T(B_X(0, \lambda))} \\ &= \overline{\frac{1}{2^2}T(B_X(0, 1))} \\ &= \overline{T\left(B_X\left(0, \frac{\lambda}{2^2}\right)\right)}. \end{aligned}$$

Inductively, pick $x_k \in B_X(0, \frac{\lambda}{2^{k-1}})$ such that $\left\|y - \sum_{j=1}^k Tx_j\right\| < \frac{1}{2^k}$. Note that for all $k \in \mathbb{N}$, we have $\|x_k\|_X < \frac{\lambda}{2^{k-1}}$. Therefore $\sum_{j=1}^{\infty} x_j$ is absolutely convergent. Since X is Banach, by Proposition 1.1.2 the series $\sum_{j=1}^{\infty} x_j$ is convergent. In particular:

$$\begin{aligned} \left\|\sum_{j=1}^{\infty} x_j\right\|_X &\leq \sum_{j=1}^{\infty} \|x_j\|_X \\ &< \sum_{j=1}^{\infty} \frac{\lambda}{2^{j-1}} \\ &= 2\lambda. \end{aligned}$$

Hence $\sum_{j=1}^{\infty} x_j \in B_X(0, 2\lambda)$. Now note that $\sum_{j=1}^k x_j \rightarrow \sum_{j=1}^{\infty} x_j$ in X . Also, $\left\|y - \sum_{j=1}^k Tx_j\right\| < \frac{1}{2^k}$ implies that $T\left(\sum_{j=1}^k x_j\right) \rightarrow y$ in Y . Since T is continuous, T has a closed graph. Hence $y = T\sum_{j=1}^{\infty} x_j \in T(B_X(0, 2\lambda))$. Since $y \in B_Y(0, 1)$ was arbitrary, we've established the inclusion $B_Y(0, 1) \subset T(B_X(0, 2\lambda))$.

Now let $U \subseteq X$ be open. Let $y_0 \in T(U)$. Since T is surjective, find $x_0 \in U$ such that

$Tx_0 = y_0$. Since U is open there exists an $\epsilon > 0$ such that $B_X(x_0, \epsilon) \subseteq U$. Therefore:

$$\begin{aligned} T(U) &\supset T(B_X(x_0, \epsilon)) \\ &= Tx_0 + \frac{\epsilon}{2\lambda} T(B_X(0, 2\lambda)) \\ &\supset y_0 + \frac{\epsilon}{2\lambda} B_Y(0, 1) \\ &= B_Y\left(y_0, \frac{\epsilon}{2\lambda}\right) \end{aligned}$$

Thus $T(U)$ is open, establishing T as an open map. $\square$

Corollary 1.6.3. *Let $T : X \rightarrow Y$ be a bijective, linear, and continuous map between Banach spaces. Then T^{-1} is continuous.*

Proof. Since T is continuous, T has a closed graph. By the Open-Mapping Theorem T is an open map. Hence $(T^{-1})^{-1}(U) = T(U)$ is open. $\square$

Corollary 1.6.4. *Let X be a vector space. Suppose $\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ are norms such that $(X, \|\cdot\|_\alpha)$ and $(X, \|\cdot\|_\beta)$ are Banach. If there exists a $C > 0$ such that $\|x\|_\alpha \leq C\|x\|_\beta$ for all $x \in X$, then there exists $C' > 0$ such that $\|x\|_\beta \leq C'\|x\|_\alpha$ for all $x \in X$.*

Proof. Certainly the map $\text{id} : (X, \|\cdot\|_\beta) \rightarrow (X, \|\cdot\|_\alpha)$ is bijective and linear. Moreover, $\|\text{id } x\|_\alpha = \|x\|_\alpha \leq C\|x\|_\beta$. Hence id is continuous. By Corollary 1.6.3 id^{-1} is continuous. Hence there exists $C' > 0$ such that $\|x\|_\beta = \|\text{id}^{-1} x\|_\beta \leq C'\|x\|_\alpha$. $\square$

Remark. The above corollary can be rephrased as: if $\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ are norms such that $(X, \|\cdot\|_\alpha)$ and $(X, \|\cdot\|_\beta)$ are Banach, then $\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ are equivalent.

Theorem 1.6.4 (Closed Graph Theorem). *Let X and Y be Banach spaces. If $T : X \rightarrow Y$ is linear and has a closed graph, then T is continuous.*

Proof. Define $\|\cdot\|_* : X \rightarrow \mathbb{R}$ by $\|x\|_* := \|x\|_X + \|Tx\|_Y$. It is routine to check that $(X, \|\cdot\|_*)$ is a normed linear space. We will check that $(X, \|\cdot\|_*)$ is Banach. Suppose $\{x_k\}_{k=1}^\infty$ is Cauchy with respect to $\|\cdot\|_*$. Then clearly $\{x_k\}_{k=1}^\infty$ is Cauchy with respect to $\|\cdot\|_X$ and $\{Tx_k\}_{k=1}^\infty$ is Cauchy with respect to $\|\cdot\|_Y$. Since X and Y are Banach, there exists $x \in X$ and $y \in Y$ such that $x_k \rightarrow x$ with respect to $\|\cdot\|_X$ and $Tx_k \rightarrow y$ with respect to $\|\cdot\|_Y$. Since T has a closed graph, $y = Tx$. Hence:

$$\begin{aligned} \|x - x_k\|_* &= \|x - x_k\|_X + \|Tx - Tx_k\|_Y \\ &\rightarrow 0 \text{ as } k \rightarrow \infty. \end{aligned}$$

Thus $x_k \rightarrow x$ with respect to $\|\cdot\|_*$, establishing that $(X, \|\cdot\|_*)$ is Banach. But since $(X, \|\cdot\|_X)$ and $(X, \|\cdot\|_*)$ are Banach, and since $\|x\|_X \leq \|x\|_*$, Corollary 1.6.4 says there exists $C' > 0$ such that $\|x\|_* \leq C'\|x\|_X$. Therefore $\|Tx\|_Y \leq \|x\|_* \leq C'\|x\|_X$ for all $x \in X$. Thus T is continuous. $\square$

Example 1.6.2. *jan 29th, bijective but inverse is unbounded*

Example 1.6.3. bounded but inverse is unbounded**Exercises**

Exercise 1.6.1. Let f be infinitely differentiable on $\mathbb{R}$. Suppose for any $x \in \mathbb{R}$ there is some n such that $f^{(n)}(x) = 0$ (note that n may depend on x). Show that f must be a polynomial. Hint: Consider the sets $Z_n = \{x \mid f^{(n)}(x) = 0\}$ and recall Baire's Category Theorem.

Proof.

□

Exercise 1.6.2. Let $T : X \rightarrow Y$ be a linear map between normed spaces. Prove the assertion made in class: The graph $\{(x, Tx) \mid x \in X\}$ of T is closed in $X \times Y$ if and only if:

$$\text{if } x_n \rightarrow x \text{ and } Tx_n \rightarrow y \text{ then } y = Tx.$$

Proof.

□

Exercise 1.6.3. Define $T : c_0 \rightarrow c_0$ by $T(\{x_n\}_{n=1}^\infty) = \{x_{n+1}\}_{n=1}^\infty$. Determine if T is: injective, surjective, open, closed, invertible

Proof.

- Let $(x_1, x_2, x_3, \dots)$ and $(y_1, y_2, y_3, \dots)$ be sequences in c_0 and $\alpha \in \mathbb{F}$. Observe that:

$$\begin{aligned} T((x_1, x_2, x_3, \dots) + \alpha(y_1, y_2, y_3, \dots)) &= T((x_1 + \alpha y_1, x_2 + \alpha y_2, x_3 + \alpha y_3, \dots)) \\ &= (x_2 + \alpha y_2, x_3 + \alpha y_3, x_4 + \alpha y_4, \dots) \\ &= (x_2, x_3, x_4, \dots) + \alpha(y_2, y_3, y_4, \dots) \\ &= T((x_1, x_2, x_3, \dots)) + \alpha T((y_1, y_2, y_3, \dots)) \end{aligned}$$

Thus T is linear.

- Consider the sequences $(1, 0, 0, \dots)$ and $(2, 0, 0, \dots)$. These are elements of c_0 which are not equal, but $T((1, 0, 0, \dots)) = (0, 0, 0, \dots) = T((2, 0, 0, \dots))$. Therefore T is not injective.
- Let $(x_1, x_2, x_3, \dots) \in c_0$. Then $T((0, x_1, x_2, x_3, \dots)) = (x_1, x_2, x_3, \dots)$. Thus T is surjective.
- Let $(x_1, x_2, x_3, \dots) \in c_0$. Observe that:

$$\begin{aligned} \|T((x_1, x_2, x_3, \dots))\|_{\ell^\infty} &= \|(x_2, x_3, x_4, \dots)\|_{\ell^\infty} \\ &= \sup\{|x_i| \mid i = 2, 3, 4, \dots\} \\ &\leq \sup\{|x_i| \mid i = 1, 2, 3, \dots\} \\ &= \|(x_1, x_2, x_3, \dots)\|_{\ell^\infty}. \end{aligned}$$

Thus T is bounded, hence continuous.

- Since T is surjective, continuous, and linear, by the Open-Mapping Theorem T is open.

- Since T is continuous, T has a closed graph.
- Since T is not bijective, T is not invertible.

□

Exercise 1.6.4. Let $L : X \rightarrow Y$ be a linear map between Banach spaces. Suppose that the conditions $x_n \rightarrow 0$ and $Lx_n \rightarrow y$ imply that $y = 0$. Prove that L is continuous.

Proof. Let $\{(x_n, Tx_n)\}_{n=1}^{\infty} \subseteq \text{Gr}(L)$ be a sequence such that $(x_n, Tx_n) \rightarrow (x, y) \in X \times Y$. Then $x_n \rightarrow x$ and $Tx_n \rightarrow y$. Define $u_n := x_n - x$. Then $u_n \rightarrow 0$ and $Tu_n \rightarrow y - Tx$. By hypothesis $y - Tx = 0$, giving $y = Tx$. Thus $(x, y) \in \text{Gr}(T)$, which means T has a closed graph. By the Closed Graph Theorem, T is continuous. □

Exercise 1.6.5. Prove that if a closed map has an inverse then the inverse is also closed.

Proof. Let $\{y_n\}_{n=1}^{\infty}$ be a sequence in Y such that $y_n \rightarrow y$ and $T^{-1}y_n \rightarrow x$. Since T admits an inverse, it is bijective. Hence for each $y_n \in Y$, there exists $x_n \in X$ such that $y_n = Tx_n$. The two sequences above can be rewritten as $Tx_n \rightarrow y$ and $T^{-1}(Tx_n) = x_n \rightarrow x$. Since T has a closed graph, $y = Tx$. Since T is bijective, $x = T^{-1}y$. Thus T^{-1} has a closed graph. □

Exercise 1.6.6. Let $L : X \rightarrow Y$ be a continuous linear surjection, where X and Y are Banach spaces. Let $y_n \rightarrow y$ in Y . Prove that there exists points $x_n \in X$ and a constant $c \in \mathbb{R}$ such that $Lx_n = y_n$, the sequence $\{x_n\}_{n=1}^{\infty}$ converges, and $\|x_n\| \leq c\|y_n\|$.

Proof.

□

§ 1.7. Continuous Dual, Adjoint Maps, and Weak Convergence

Definition 1.7.1. Let X and Y be normed linear spaces and $T \in \mathcal{L}(X, Y)$. The adjoint of T is a map $T^* : Y^* \rightarrow X^*$ given by $T^*(\varphi) = \varphi(Tx)$.

Proposition 1.7.1. Let X and Y be normed linear spaces and $T \in \mathcal{L}(X, Y)$. Then $T^* \in \mathcal{L}(Y^*, X^*)$, and $\|T^*\|_{\mathcal{L}(Y^*, X^*)} = \|T\|_{\mathcal{L}(X, Y)}$.

Proof.

□

Exercises

Exercise 1.7.1. Let X be a normed linear space, let $K \subset X$, and let $\Omega \subset X^*$. Show that $K^\perp$ is closed in X^* and that $\Omega_\perp$ is closed in X .

Proof. Let $\{\varphi_n\}_{n=1}^\infty \subset K^\perp$ be a sequence such that $\varphi_n \rightarrow \Phi \in X^*$. Let $k \in K$ be arbitrary. Then $\varphi_n(k) \rightarrow \Phi(k)$. Since $\varphi_n(k) = 0$ for all $n \in \mathbb{N}$, it must be the case that $\Phi(k) = 0$. Since k was arbitrary, we have $\Phi \in K^\perp$. Since $K^\perp$ contains all of its limit points, it is closed in X^* .

Let $\{x_n\}_{n=1}^\infty \subset \Omega_\perp$ be a sequence such that $x_n \rightarrow y \in X$. Let $\varphi \in \Omega$ be arbitrary. Then $\varphi(x_n) \rightarrow \varphi(y)$ since φ is continuous. Since $\varphi(x_n) = 0$ for all $n \in \mathbb{N}$, it must be the case that $\varphi(y) = 0$. Since φ was arbitrary, $y \in \Omega_\perp$. Since $\Omega_\perp$ contains all of its limit points, it is closed in X . $\square$

Exercise 1.7.2. Let X be a normed linear space and $K \subset X$. Show that $(K^\perp)_\perp$ is the closure of the linear space K in X .

Proof.

$\square$

§ 1.8. L^p Spaces

Exercises

Exercise 1.8.1. For which $r \in \mathbb{R}$ is the function $f(x) = \frac{1}{|x|^r}$ in $L^1([0, 1])$? What about $L^p([0, 1])$?

Proof. Since $x \in [0, 1]$, we have $|x| = x$. Note that:

$$\int_0^1 x^{-r} dx = -\frac{1}{(r-1)x^{r-1}} \Big|_0^1$$

The right-hand side of this equation can only be evaluated if $r-1 > 0$. Hence $\frac{1}{|x|^r} \in L^1([0, 1])$ provided $r > 1$. Similarly, note that:

$$\begin{aligned} \int_0^1 (x^{-r})^p dx &= \int_0^1 x^{-rp} dx \\ &= -\frac{1}{(rp-1)x^{rp-1}} \Big|_0^1 \end{aligned}$$

The right-hand side of this equation can only be evaluated if $rp-1 > 0$. Hence $\frac{1}{|x|^r} \in L^p([0, 1])$ provided $r > \frac{1}{p}$. $\square$

Chapter 2

Hilbert Spaces

§ 2.1. Inner Product Spaces

Definition 2.1.1. Let V be a vector space over $\mathbb{F}$. An *inner product* is a map:

$$(\cdot, \cdot) : V \rightarrow \mathbb{F}$$

satisfying for all $x, y \in V$:

- (1) $(x, x) \geq 0$ and $(x, x) = 0$ if and only if $x = 0$;
- (2) $(\cdot, \cdot)$ is linear in the first slot.
- (3) $(x, y) = \overline{(y, x)}$.

Remark. Note that an inner product over $\mathbb{R}$ (resp. $\mathbb{C}$) can be defined as a positive-definite and symmetric (resp. Hermitian) bilinear form (resp. sesquilinear form). From (2) and (3) we have:

$$\begin{aligned} (x, \lambda y) &= \overline{(\lambda y, x)} \\ &= \overline{\lambda(y, x)} \\ &= \bar{\lambda} \overline{(y, x)} \\ &= \bar{\lambda}(x, y). \end{aligned}$$

Definition 2.1.2. Let V be an inner product space. We say $x, y \in V$ are orthogonal and write $x \perp y$ if $(x, y) = 0$.

Example 2.1.1.

- (1) The space $\mathbb{C}^n$ with $(x, y) := \sum_{j=1}^n x_j \overline{y_j}$ is an inner product space.
- (2) The space ℓ^2 with $(x, y) := \sum_{j=1}^{\infty} x_j \overline{y_j}$ is an inner product space. Note that the normed induced by the inner product is precisely the familiar ℓ^2 -norm.
- (3) The space $L^2(\Omega)$ with $(f, g) := \int_{\Omega} f(x) \overline{g(x)} dx$ is an inner product space. The norm induced by the inner product is also the familiar L^2 -norm.

Proposition 2.1.1. Let X be an inner product space and $x, y \in X$, $y \neq 0$. There exists $c \in \mathbb{F}$ such that $(x - cy) \perp y$.

Proof. By definition $(x - cy) \perp y$ means $(x - cy, y) = 0$. Hence $(x, y) - c(y, y) = 0$. Solving for c gives $c = \frac{(x, y)}{\|y\|^2}$. □

Theorem 2.1.1. Let X be an inner product space. Let $x, y \in X$.

- (1) If $x \perp y$, then $\|x + y\|^2 = \|x\|^2 + \|y\|^2$ (Pythagorean Theorem).
 (2) $|(x, y)| \leq \|x\| \|y\|$ (Cauchy-Schwartz Inequality).
 (3) $\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2$ (Parallelogram law).

Proof. (1) Observe that:

$$\begin{aligned}
 \|x + y\|^2 &= (x + y, x + y) \\
 &= (x, x) + (x, y) + (y, x) + (y, y) \\
 &= (x, x) + (x, y) + \overline{(x, y)} + (y, y) \\
 &= (x, x) + 2\Re((x, y)) + (y, y) \\
 &= (x, x) + (y, y) \\
 &= \|x\|^2 + \|y\|^2.
 \end{aligned}$$

(2) Assume $y \neq 0$. Find c such that $(x - cy) \perp y$. Then:

$$\begin{aligned}
 \|x\|^2 &= \|(x - cy) + cy\|^2 \\
 &= \|(x - cy)\|^2 + \|cy\|^2 \\
 &\geq |c|^2 \|y\|^2.
 \end{aligned}$$

Note that $c = \frac{(x, y)}{\|y\|^2}$ by Proposition 2.1.1. Hence $\|x\|^2 \geq \frac{|(x, y)|^2}{\|y\|^4} \|y\|^2$. Solving for $|(x, y)|$ establishes the inequality.

(3) Exercise 2.1.1. □

Proposition 2.1.2. Let V be an inner product space. Define $\|\cdot\| : V \rightarrow \mathbb{F}$ by $\|x\| = \sqrt{(x, x)}$. Then $(V, \|\cdot\|)$ is a normed linear space.

Proof. Let $v, w \in V$ and $\alpha \in \mathbb{F}$. We have that:

$$\begin{aligned}
 \|\alpha v\| &= \sqrt{(\alpha v, \alpha v)} \\
 &= \sqrt{\alpha \bar{\alpha} (v, v)} \\
 &= \sqrt{|\alpha|^2 (v, v)} \\
 &= |\alpha| \|v\|.
 \end{aligned}$$

Thus $\|\cdot\|$ is homogeneous. By Cauchy-Schwartz:

$$\begin{aligned}
 \|v + w\|_2^2 &= \langle v + w, v + w \rangle \\
 &= \langle v, v \rangle + \langle v, w \rangle + \langle w, v \rangle + \langle w, w \rangle \\
 &= \|v\|_2^2 + \langle v, w \rangle + \overline{\langle v, w \rangle} + \|w\|_2^2 \\
 &= \|v\|_2^2 + 2\Re(\langle v, w \rangle) + \|w\|_2^2 \\
 &\leq \|v\|_2^2 + 2|\langle v, w \rangle| + \|w\|_2^2 \\
 &\leq \|v\|_2^2 + 2\|v\|_2 \|w\|_2 + \|w\|_2^2 \\
 &= (\|v\|_2 + \|w\|_2)^2,
 \end{aligned}$$

Squaring both sides establishes the triangle inequality. For positive-definiteness, suppose $\|v\| = 0$. Then $\sqrt{(v, v)} = 0$, so it must be the case that $v = 0$. $\square$

Definition 2.1.3. Let X be an inner product space and $A \subset X$ a subspace. The orthogonal complement of A is the set $A^\perp := \{x \in X \mid (x, a) = 0 \text{ for all } a \in A\}$.

Proposition 2.1.3. If X is an inner product space and $A \subset X$ is a subspace, then $A^\perp$ is closed in X .

Proof. Define $\varphi_a : X \rightarrow \mathbb{F}$ by $x \mapsto (x, a)$. This map is continuous. Then $A^\perp = \bigcap_{a \in A} \varphi_a^{-1}(\{0\})$. Since φ_a is continuous, $A^\perp$ is closed. $\square$

Exercises

Exercise 2.1.1. Prove the Parallelogram Law in any inner-product space.

Proof. Observe that:

$$\begin{aligned} \|x + y\|^2 + \|x - y\|^2 &= (x + y, x + y) + (x - y, x - y) \\ &= (x, x) + 2\Re((x, y)) + (y, y) + (x, x) - 2\Re((x, y)) + (y, y) \\ &= 2(x, x) + 2(y, y) \\ &= 2\|x\|^2 + 2\|y\|^2. \end{aligned} \quad \square$$

Exercise 2.1.2. Let X be an inner-product space over $\mathbb{C}$ and fix $y \in X$. Prove that the map $\varphi_y : X \rightarrow \mathbb{C}$ defined by $\varphi_y(x) = (x, y)$ is continuous.

Proof. Let $x_1, x_2 \in X$. Observe that:

$$\begin{aligned} |\varphi_y(x_1) - \varphi_y(x_2)| &= |(x_1, y) - (x_2, y)| \\ &= |(x_1 - x_2, y)| \\ &\leq \|x_1 - x_2\| \|y\|. \end{aligned}$$

Since φ_y is Lipschitz, φ_y is continuous. $\square$

Exercise 2.1.3. Prove that in an inner-product space X , if $\{x_n\}_{n=1}^\infty \subset X$ is a sequence and $y \in X$ satisfy:

$$\|x_n\| \rightarrow \|y\| \text{ and } (x_n, y) \rightarrow \|y\|^2,$$

then $x_n \rightarrow y$.

Proof. Observe that:

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \|x_n - y\|^2 &= \lim_{n \rightarrow \infty} (x_n - y, x_n - y) \\
 &= \lim_{n \rightarrow \infty} \left((x_n, x_n) - (y, x_n) - (x_n, y) + (y, y) \right) \\
 &= \lim_{n \rightarrow \infty} \left(\|x_n\|^2 - 2\Re((x_n, y)) + \|y\|^2 \right) \\
 &= \lim_{n \rightarrow \infty} \|x_n\|^2 - \lim_{n \rightarrow \infty} 2\Re((x_n, y)) + \lim_{n \rightarrow \infty} \|y\|^2 \\
 &= \lim_{n \rightarrow \infty} \|x_n\|^2 - 2\Re\left(\lim_{n \rightarrow \infty} (x_n, y)\right) + \lim_{n \rightarrow \infty} \|y\|^2 \\
 &= \|y\|^2 - 2\|y\|^2 + \|y\|^2 \\
 &= 0.
 \end{aligned}$$

Since $\|x_n - y\|^2 \rightarrow 0$, it must be the case that $\|x_n - y\| \rightarrow 0$. Thus $x_n \rightarrow y$. $\square$

§ 2.2. Hilbert Spaces

Definition 2.2.1. An inner product space which is complete is called a Hilbert space.

Remark. It is standard to assume that the base field of a Hilbert space is $\mathbb{C}$.

Theorem 2.2.1. Let $\mathcal{H}$ be a Hilbert space and $K \subset \mathcal{H}$ a nonempty subset which is convex and closed. For any $x \in \mathcal{H}$, there exists a unique $y \in K$ which is closed to x .

Proof. Let $x \in X$. Let $\alpha := \text{dist}(x, K)$. Find a sequence $\{y_n\}_{n=1}^\infty \subset K$ such that $\|x - y_n\| \rightarrow \alpha$ as $n \rightarrow \infty$. For any $n, m \in \mathbb{N}$:

$$\begin{aligned}
 \|y_n - y_m\|^2 &= \|(y_n - x) + (x - y_m)\|^2 \\
 &= 2\|y_n - x\|^2 + 2\|x - y_m\|^2 - \|y_n + y_m - 2x\|^2 && \text{Parallelogram Law} \\
 &= 2\|y_n - x\|^2 + 2\|x - y_m\|^2 - 4\left\|\frac{y_n + y_m}{2} - x\right\|^2 && \text{See footnote}^1 \\
 &\leq 2\|y_n - x\|^2 + 2\|y_m - x\|^2 - 4\alpha^2.
 \end{aligned}$$

Since $\|x - y_n\| \rightarrow \alpha^2$ and $\|x - y_m\| \rightarrow \alpha^2$, note that $\|y_n - y_m\|^2$ can be made arbitrary small. Hence $\{y_n\}_{n=1}^\infty$ is Cauchy. Since K is a closed subset of a Hilbert space, it is complete. Hence there exists some $y \in K$ such that $y_n \rightarrow y$. Finally:

$$\begin{aligned}
 \|x - y\| &= \left\| x - \lim_{n \rightarrow \infty} y_n \right\| \\
 &= \lim_{n \rightarrow \infty} \|x - y_n\| \\
 &= \alpha.
 \end{aligned}$$

¹Note $\frac{y_n + y_m}{2} \in K$ because K is convex.

Now if y' is any other element in K with $\|y' - x\| = \alpha$, by our previous calculation:

$$\begin{aligned}\|y - y'\|^2 &= 2\|y - x\|^2 + 2\|y' - x\|^2 - \|y + y' - 2x\|^2 \\ &\leq 2\alpha^2 + 2\alpha^2 - 4\alpha^2 \\ &= 0.\end{aligned}$$

Thus $y = y'$. □

Theorem 2.2.2. *Let $\mathcal{H}$ be a Hilbert space. Let $Z \subset \mathcal{H}$ be a closed subspace. Then $\mathcal{H} = Z \oplus Z^\perp$.*

Proof. If $z \in Z \cap Z^\perp$, then $z \perp z$; i.e., $(z, z) = 0$. Therefore $z = 0$.

Now let $x \in \mathcal{H}$. By Theorem 2.2.1, let $z \in Z$ be the unique element closest to x . We show that $x - z \in Z^\perp$. Let $z' \in Z$ be nonzero. By Proposition 2.1.1, choose $\lambda \in \mathbb{C}$ such that $\lambda z' \perp (x - z) - \lambda z'$. Write:

$$x - z = \lambda z' + (x - z) - \lambda z'.$$

Then the Pythagorean Theorem gives:

$$\begin{aligned}\|x - z\|^2 &= \|\lambda z'\|^2 + \|x - (z + \lambda z')\|^2 \\ &\geq \|\lambda z'\|^2 + \|x - z\|^2.\end{aligned}$$

Note that the inequality is a result from the fact that z is the closest element in Z to x . Subtracting $\|x - z\|^2$ from both sides yields:

$$0 \geq \|\lambda z'\|^2$$

Because $z' \neq 0$, this forces $\lambda = 0$. Since $\lambda = \frac{(x - z, z')}{\|z'\|^2}$, we obtain $(x - z, z') = 0$. As $z' \in Z$ was arbitrary, $x - z \in Z^\perp$. Consequently, $x = z + (x - z) \in Z + Z^\perp$, establishing $x \in Z \oplus Z^\perp$. □

Definition 2.2.2. Let X be an inner product space. Let $\mathcal{U} := \{u_\alpha \mid \alpha \in I\} \subset X$. We say $\mathcal{U}$ is orthogonal if for all $\alpha \neq \beta$, we have $(u_\alpha, u_\beta) = 0$. We say $\mathcal{U}$ is orthonormal if for all $\alpha, \beta \in I$, we have:

$$(u_\alpha, u_\beta) = \begin{cases} 1, & \alpha = \beta \\ 0, & \alpha \neq \beta. \end{cases}$$

Theorem 2.2.3. *Let X be a nonzero inner product space. There exists a maximal orthonormal set by inclusion*

Proof. Order all possible orthonormal sets by inclusion. Show that every chain admits an upper bound. Then apply Zorn's Lemma. □

Theorem 2.2.4. *Let X be a separable inner product space. Then any orthonormal subset is at most countable.*

Proof. Let $\{a_j\}_{j=1}^\infty$ be a countable dense subset of X . Suppose towards contradiction $\mathcal{U} := \{u_\alpha \mid \alpha \in I\}$ is an uncountable orthonormal set in X . Pick $\alpha, \beta \in I$ such that $\alpha \neq \beta$. Then:

$$\begin{aligned}\|u_\alpha - u_\beta\|^2 &= \|u_\alpha\|^2 + \|u_\beta\|^2 \\ &= 2.\end{aligned}$$

Therefore, any two elements of $\mathcal{U}$ are $\sqrt{2}$ apart. Now for all $\alpha \in I$, consider the open balls $B(u_\alpha, \frac{\sqrt{2}}{2})$. Since each open ball has diameter less than $\sqrt{2}$, each ball can contain at most one element of $\{a_j\}_{j=1}^\infty$. Moreover, the collection $\{B(u_\alpha, \frac{\sqrt{2}}{2}) \mid \alpha \in I\}$ is disjoint. Indeed, if $y \in B(u_\alpha, \frac{\sqrt{2}}{2}) \cap B(u_\beta, \frac{\sqrt{2}}{2})$, then $\|u_\alpha - u_\beta\| \leq \|u_\alpha - y\| + \|y - u_\beta\| < \sqrt{2}$, which is a contradiction. Since $\{a_j\}_{j=1}^\infty$ is a dense subset, each a_j must be contained in a ball. We can define a surjection from $\{a_j\}_{j=1}^\infty$ to $\{B(u_\alpha, \frac{\sqrt{2}}{2}) \mid \alpha \in I\}$. But since $\{B(u_\alpha, \frac{\sqrt{2}}{2}) \mid \alpha \in I\}$ is indexed by an uncountable set, it must be the case that $\{a_j\}_{j=1}^\infty$ is uncountable. This is a contradiction, whence any orthonormal subset must be at most countable. $\square$

Theorem 2.2.5. Let X be an inner product space. Let $Y \subset X$ be a subspace with $\dim Y = n$.

- (1) Y has an orthonormal basis $\{u_1, \dots, u_n\}$;
- (2) For each $y \in Y$, we can write $y = \sum_{j=1}^n (y, u_j) u_j$. Moreover, $\|y\|^2 = \sum_{j=1}^n |(y, u_j)|^2$.
- (3) There is an isometric isomorphism $Y \rightarrow \mathbb{C}^n$.
- (4) For any $x \in X$, we have $x - \sum_{j=1}^n (x, u_j) u_j \perp Y$. Moreover, $\sum_{j=1}^n (x, u_j) u_j$ is the unique element in Y closest to x ,

Proof. (1) Apply the Gram-Schmidt process to a basis of Y .

(2) Let $y \in Y$. We can uniquely write $y = \sum_{j=1}^n a_j u_j$ for some $a_1, \dots, a_n \in \mathbb{C}$. Then $(y, u_j) = (\sum_{k=1}^n a_k u_k, u_j) = \sum_{k=1}^n a_k (u_k, u_j) = a_j$. Hence $y = \sum_{j=1}^n (y, u_j) u_j$.

(3) For any $y \in Y$, we can write $y = \sum_{j=1}^n a_j u_j$. Then $Y \rightarrow \mathbb{C}^n$ given by $y \mapsto (a_1, \dots, a_n)$ is linear, bijective, and norm preserving.

(4) Let $x \in X$. Define $a_j := (x, u_j)$ for each $j = 1, \dots, n$. For each $k = 1, \dots, n$:

$$\begin{aligned}\left(x - \sum_{j=1}^n a_j u_j, u_k\right) &= (x, u_k) - \left(\sum_{j=1}^n a_j u_j, u_k\right) \\ &= (x, u_k) - \sum_{j=1}^n a_j (u_j, u_k) \\ &= (x, u_k) - a_k \\ &= (x, u_k) - (x, u_k) \\ &= 0.\end{aligned}$$

Since any $y \in Y$ can be uniquely written as a linear combination of $u_1, \dots, u_n$, the above string of equalities implies $x - \sum_{j=1}^n (x, u_j) u_j \perp Y$. Now let $\sum_{j=1}^n b_j u_j$ be any element in Y .

Then:

$$\begin{aligned}
 \left\| x - \sum_{j=1}^n b_j u_j \right\|^2 &= \left\| x - \sum_{j=1}^n a_j u_j + \sum_{j=1}^n (a_j - b_j) u_j \right\|^2 \\
 &= \left\| x - \sum_{j=1}^n a_j u_j \right\|^2 + \left\| \sum_{j=1}^n (a_j - b_j) u_j \right\|^2 \\
 &= \left\| x - \sum_{j=1}^n a_j u_j \right\|^2 + \sum_{j=1}^n |a_j - b_j|^2 \\
 &\geq \left\| x - \sum_{j=1}^n a_j u_j \right\|^2,
 \end{aligned}$$

where we used the fact that $x - \sum_{j=1}^n a_j u_j \perp \sum_{j=1}^n (a_j - b_j) u_j$. This establishes that $\sum_{j=1}^n a_j u_j$ is the unique element in Y closest to x , where we have equality only when $a_j = b_j$ for all $j = 1, \dots, n$. $\square$

Theorem 2.2.6. *Let $\mathcal{H}$ be a Hilbert space. Let $\{x_j\}_{j=1}^\infty$ be an orthogonal sequence in $\mathcal{H}$. Then $\sum_{j=1}^\infty x_j$ converges in $\mathcal{H}$ if and only if $\sum_{j=1}^\infty \|x_j\|^2 < \infty$.*

Proof. Define:

$$\begin{aligned}
 z_n &:= \sum_{j=1}^n x_j, \\
 s_n &:= \sum_{j=1}^n \|x_j\|^2.
 \end{aligned}$$

Now $\sum_{j=1}^\infty x_j$ converges if and only if $\{z_n\}_{n=1}^\infty$ is Cauchy. For $n > m$:

$$\begin{aligned}
 \|z_n - z_m\|^2 &= \left\| \sum_{j=m+1}^n x_j \right\|^2 \\
 &= \sum_{j=m+1}^n \|x_j\|^2 \\
 &= |s_n - s_m|.
 \end{aligned}$$

From this, $\{z_n\}_{n=1}^\infty$ is Cauchy in $\mathcal{H}$ if and only if $\{s_n\}_{n=1}^\infty$ is Cauchy in $\mathbb{C}$. $\square$

Theorem 2.2.7 (Bessel's Inequality). *Let X be an inner-product space. Let $\{u_j\}_{j=1}^\infty$ be an orthonormal sequence in X . For all $x \in X$, we have $\sum_{j=1}^\infty |(x, u_j)|^2 \leq \|x\|^2$.*

Proof. For each $n \in \mathbb{N}$, define $Y_n := \text{span}\{u_1, \dots, u_n\}$. Let P_{Y_n} denote the projection onto Y_n . Then:

$$\begin{aligned} \sum_{j=1}^n |(x, u_j)|^2 &= \sum_{j=1}^n \|(x, u_j)u_j\|^2 \\ &= \left\| \sum_{j=1}^n (x, u_j)u_j \right\|^2 \\ &= \|P_{Y_n}(x)\|^2 \\ &\leq \|P_{Y_n}\|^2 \|x\|^2 \\ &= \|x\|^2. \end{aligned}$$

Note that $\|P_{Y_n}\| = 1$ is established in Exercise 2.2.5. Since n was arbitrary, by the Monotone Convergence Theorem $\sum_{j=1}^{\infty} |(x, u_j)|^2 \leq \|x\|^2$. $\square$

Theorem 2.2.8. Let $\mathcal{H}$ be a Hilbert space. Let $\{u_j\}_{j=1}^{\infty}$ be an orthonormal sequence in $\mathcal{H}$. The following are equivalent:

- (1) $\{u_j\}_{j=1}^{\infty}$ is a maximal orthonormal set;
- (2) For all $x \in \mathcal{H}$, if $x \perp \{u_j\}_{j=1}^{\infty}$, then $x = 0$;
- (3) For all $x \in \mathcal{H}$, there exists a unique sequence $\{a_j\}_{j=1}^{\infty}$ such that $x = \sum_{j=1}^{\infty} a_j u_j$.

Proof. We will only show (2) if and only if (3), as (1) if and only if (2) follows immediately by applying definitions.

Suppose for all $x \in \mathcal{H}$ we have $x \perp \{u_j\}_{j=1}^{\infty}$ implies $x = 0$. Let $x \in \mathcal{H}$. Claim: $x - \sum_{j=1}^{\infty} (x, u_j)u_j \perp \{u_j\}_{j=1}^{\infty}$. Indeed, for any $u_k \in \{u_j\}_{j=1}^{\infty}$:

$$\begin{aligned} \left(x - \sum_{j=1}^{\infty} (x, u_j)u_j, u_k \right) &= \lim_{n \rightarrow \infty} \left(x - \sum_{j=1}^n (x, u_j)u_j, u_k \right) \\ &= \lim_{n \rightarrow \infty} \left((x, u_k) - \sum_{j=1}^n ((x, u_j)u_j, u_k) \right) \\ &= \lim_{n \rightarrow \infty} ((x, u_k) - (x, u_k)) \\ &= 0. \end{aligned}$$

Hence $x - \sum_{j=1}^{\infty} (x, u_j)u_j = 0$, and rearranging gives $x = \sum_{j=1}^{\infty} (x, u_j)u_j$. Now suppose there exists a sequence $\{b_n\}_{n=1}^{\infty}$ such that $x = \sum_{j=1}^{\infty} b_j u_j$. Then $0 = \sum_{j=1}^{\infty} ((x, u_j) - b_j)u_j$, which by Bessel's Inequality results in $\sum_{j=1}^{\infty} |(x, u_j) - b_j|^2 \leq 0$. It must be the case that $(x, u_j) = b_j$ for all $j \in \mathbb{N}$, establishing uniqueness.

Let $x \in \mathcal{H}$ and suppose $x \perp \{u_j\}_{j=1}^{\infty}$. Write $x = \sum_{j=1}^{\infty} (x, u_j)u_j$. Then for all $k \in \mathbb{N}$:

$$\begin{aligned} (x, u_k) &= \left(\sum_{j=1}^{\infty} (x, u_j)u_j, u_k \right) \\ &= 0. \end{aligned}$$

Thus $x = 0$. □

Definition 2.2.3. Let $\mathcal{H}$ be a Hilbert space and $\{u_j\}_{j=1}^{\infty}$ an orthonormal sequence in $\mathcal{H}$. If any of the three equivalent statements in Theorem 2.2.8 hold true, we say $\{u_j\}_{j=1}^{\infty}$ is a orthonormal basis (or Hilbert basis).

Definition 2.2.4. Let $\mathcal{H}$ be a Hilbert space and $\{u_j\}_{j=1}^{\infty}$ an orthonormal basis. Let $x \in \mathcal{H}$.

- (1) The sequence $\{(x, u_j)\}_{j=1}^{\infty}$ is referred to as the generalized Fourier coefficients of x .
- (2) The series $\sum_{j=1}^{\infty} (x, u_j) u_j$ is called the generalized Fourier series of x .

Example 2.2.1. The set $\{e^n\}_{n=1}^{\infty}$ is an orthonormal basis in ℓ^2 . For any separable infinite dimensional hilbert space, let $\{u_j\}_{j=1}^{\infty}$ be an orthonormal basis. Define $H \rightarrow \ell^2$ by $\sum_{j=1}^{\infty} a_j u_j \mapsto (a_1, a_2, \dots)$. This is an isometric isomorphism.

Feb 12, stuff on L^2

Theorem 2.2.9 (Riez Representation Theorem). Let $\mathcal{H}$ be a Hilbert space. Let $\varphi \in \mathcal{H}^*$. There exists a unique $y \in H$ such that for all $x \in \mathcal{H}$, $\varphi(x) = (x, y)$.

Proof. If $y, y' \in \mathcal{H}$ satisfy $\varphi(x) = (x, y) = (x, y')$ for all $x \in \mathcal{H}$, then $(x, y - y') = 0$ for all $x \in \mathcal{H}$. Note that $y - y' \in \mathcal{H}$, hence we must have that $(y - y', y - y') = 0$. Thus $\|y - y'\| = 0$, hence $y = y'$.

We now prove existence. Let $\varphi \in \mathcal{H}^*$. Let $z \in N(\varphi)^\perp$ such that $\|z\| = 1$. For any $x \in H$, we have $x - \frac{\varphi(x)}{\varphi(z)} z \in N(\varphi)$ and the equation:

$$x = \left(x - \frac{\varphi(x)}{\varphi(z)} z \right) + \left(\frac{\varphi(x)}{\varphi(z)} z \right).$$

Now:

$$\begin{aligned} (x, \lambda z) &= \left(x - \frac{\varphi(x)}{\varphi(z)} z + \frac{\varphi(x)}{\varphi(z)} z, \lambda z \right) \\ &= \left(x - \frac{\varphi(x)}{\varphi(z)} z, \lambda z \right) + \left(\frac{\varphi(x)}{\varphi(z)} z, \lambda z \right). \end{aligned}$$

Because $z \in N(\varphi)^\perp$ and $x - \frac{\varphi(x)}{\varphi(z)} z \in N(\varphi)$, then $\left(x - \frac{\varphi(x)}{\varphi(z)} z, \lambda z \right) = 0$. We now have:

$$\begin{aligned} (x, \lambda z) &= \left(x - \frac{\varphi(x)}{\varphi(z)} z, \lambda z \right) + \left(\frac{\varphi(x)}{\varphi(z)} z, \lambda z \right) \\ &= \left(\frac{\varphi(x)}{\varphi(z)} z, \lambda z \right) \\ &= \frac{\varphi(x)}{\varphi(z)} \bar{\lambda} (z, z) \\ &= \frac{\varphi(x)}{\varphi(z)} \bar{\lambda} \|z\|^2 \\ &= \frac{\varphi(x)}{\varphi(z)} \bar{\lambda}. \end{aligned}$$

Let $\lambda := \overline{\varphi(z)}$. Then $(x, \varphi(z)) = \varphi(x)$, establishing the theorem. $\square$

self-adjoint operators

Exercises

Exercise 2.2.1. Let $H = L^2([0, 1])$. Use the Gram-Schmidt orthogonalization process from linear algebra to produce an orthonormal set $\{u_1, u_2, u_3\}$ which spans the same space as $\text{span}\{1, x, x^2\}$.

Proof. Let $v_1 := 1$, $v_2 := x$, and $v_3 := x^2$. The Gram-Schmidt process gives:

$$\begin{aligned} r_1 &= v_1 = 1, \\ \|r_1\| &= \sqrt{(r_1, r_1)} = 1. \\ r_2 &= v_2 - \frac{(v_2, r_1)}{(r_1, r_1)} r_1 = x - \frac{1}{2}, \\ \|r_2\| &= \sqrt{(r_2, r_2)} = \sqrt{\int_0^1 \left(x - \frac{1}{2}\right)^2 dx} = \frac{1}{\sqrt{12}}, \\ r_3 &= v_3 - \frac{(v_3, r_1)}{(r_1, r_1)} r_1 - \frac{(v_3, r_2)}{(r_2, r_2)} r_2 = x^2 - x + \frac{1}{6}, \\ \|r_3\| &= \sqrt{(r_3, r_3)} = \sqrt{\int_0^1 \left(x^2 - x + \frac{1}{6}\right)^2 dx} = \frac{1}{\sqrt{180}}. \end{aligned}$$

Thus an orthonormal basis for $\text{span}\{1, x, x^2\}$ is:

$$\begin{aligned} u_1 &:= \frac{r_1}{\|r_1\|} = 1 \\ u_2 &:= \frac{r_2}{\|r_2\|} = \sqrt{12} \left(x - \frac{1}{2}\right) \\ u_3 &:= \frac{r_3}{\|r_3\|} = \sqrt{180} \left(x^2 - x + \frac{1}{6}\right). \end{aligned} \quad \square$$

Exercise 2.2.2. Show directly that the functions:

$$u_n(x) = e^{i2\pi n x}, \quad n \in \mathbb{Z}$$

form an orthonormal sequence in $L^1([0, 1])$. Then write the function $f(x) = x$ in terms of this basis, that is find a_n such that:

$$x \sim \sum_{n=-\infty}^{\infty} a_n u_n(x).$$

What does Parseval's identity say in this case? (After some algebra you should be able to use your summation to find a formula for $\sum_{n=1}^{\infty} \frac{1}{n^2}$.)

Proof. Observe that:

$$\begin{aligned}
 \int_0^1 e^{i2\pi nx} e^{-i2\pi mx} dx &= \int_0^1 1 dx = 1. \\
 \int_0^1 e^{i2\pi nx} e^{-i2\pi mx} dx &= \int_0^1 e^{i2\pi(n-m)x} dx \\
 &= \frac{e^{i2\pi(n-m)x}}{i2\pi(n-m)} \Big|_0^1 \\
 &= \frac{e^{i2\pi(n-m)} - 1}{i2\pi(n-m)} \\
 &= \frac{\cos(2\pi(n-m)) - i \sin(2\pi(n-m)) - 1}{i2\pi(n-m)} \\
 &= \frac{1 - 0 - 1}{i2\pi(n-m)} \\
 &= 0.
 \end{aligned}$$

Thus $\{u_n\}_{n=1}^\infty$ is an orthonormal sequence in $L^1([0, 1])$. We proceed by cases on computing the Fourier coefficients of x . For $n = 0$:

$$a_0 = \int_0^1 x dx = \frac{1}{2}.$$

For $n \neq 0$:

$$\begin{aligned}
 a_n &= (x, u_n) \\
 &= \int_0^1 x e^{-i2\pi nx} dx \\
 &= -\frac{e^{i2\pi n}(e^{i2\pi n} - i2\pi n - 1)}{4n^2\pi^2} \quad \text{From Mathematica} \\
 &= \frac{i}{2n\pi}. \quad \text{Also from Mathematica}
 \end{aligned}$$

Using Parseval's Identity:

$$\begin{aligned}
 \frac{1}{3} &= \int_0^1 x^2 dx \\
 &= \|x\|^2 \\
 &= \sum_{n=-\infty}^{\infty} |a_n|^2 \\
 &= |a_0|^2 + 2 \sum_{n=1}^{\infty} |a_n|^2 \\
 &= \frac{1}{4} + 2 \sum_{n=1}^{\infty} \left| \frac{i}{2\pi n} \right|^2 \\
 &= \frac{1}{4} + \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2}.
 \end{aligned}$$

Solving for $\sum_{n=1}^{\infty} \frac{1}{n^2}$ gives:

$$\begin{aligned}
 \sum_{n=1}^{\infty} \frac{1}{n^2} &= 2\pi^2 \left(\frac{1}{3} - \frac{1}{4} \right) \\
 &= \frac{2\pi^2}{12} \\
 &= \frac{\pi^2}{6}.
 \end{aligned}$$

□

Exercise 2.2.3. Show directly that the functions:

$$1, \cos x, \sin x, \cos 2x, \sin 2x, \cos 3x, \dots$$

are mutually orthogonal in $L^2([-\pi, \pi])$.

Proof. For $n, m \neq 0$:

$$\begin{aligned}
 \int_{-\pi}^{\pi} \cos(nx) dx &= \frac{1}{n} \sin(nx) \Big|_{-\pi}^{\pi} = \frac{1}{n} \sin(n\pi) - \frac{1}{n} \sin(-n\pi) = \frac{2}{n} \sin(n\pi) = 0, \\
 \int_{-\pi}^{\pi} \sin(nx) dx &= -\frac{1}{n} \cos(nx) \Big|_{-\pi}^{\pi} = -\frac{1}{n} \cos(n\pi) + \frac{1}{n} \cos(-n\pi) = 0, \\
 \int_{-\pi}^{\pi} \cos(nx) \cos(mx) dx &= \int_{-\pi}^{\pi} \frac{\cos((n+m)x) + \cos((n-m)x)}{2} dx = 0, \\
 \int_{-\pi}^{\pi} \sin(nx) \sin(mx) dx &= \int_{-\pi}^{\pi} \frac{\cos((n-m)x) - \cos((n+m)x)}{2} dx = 0, \\
 \int_{-\pi}^{\pi} \cos(nx) \sin(mx) dx &= \int_{-\pi}^{\pi} \frac{\sin((n+m)x) + \sin((n-m)x)}{2} dx = 0.
 \end{aligned}$$

□

Exercise 2.2.4. Let $\{u_n\}_{n=1}^\infty$ be an orthonormal basis for a Hilbert space H . Let $\{v_n\}_{n=1}^\infty$ be an orthonormal seq which satisfies $\sum_{n=1}^\infty \|u_n - v_n\|^2 < 1$. Show that $\{v_n\}_{n=1}^\infty$ is also a Hilbert basis for H .

Proof. Let $x \in \mathcal{H}$. Suppose $x \perp \{v_j\}_{j=1}^\infty$. By Parseval's Identity:

$$\begin{aligned} \|x\|^2 &= \sum_{j=1}^\infty |(x, u_j)|^2 \\ &= \sum_{j=1}^\infty |(x, u_j) - (x, v_j)|^2 \\ &= \sum_{j=1}^\infty |(x, u_j - v_j)|^2 \\ &\leq \sum_{j=1}^\infty \|x\|^2 \|u_j - v_j\|^2. \end{aligned}$$

Suppose towards contradiction $x \neq 0$. We can divide both sides of the above equation by $\|x\|^2$, giving:

$$1 \leq \sum_{j=1}^\infty \|u_j - v_j\|^2 < 1.$$

This is a contradiction, hence $x = 0$. Since we've shown $x \perp \{v_j\}_{j=1}^\infty$ implies $x = 0$, we've established $\{v_j\}_{j=1}^\infty$ as an orthonormal basis for $\mathcal{H}$. $\square$

Exercise 2.2.5. Let H be a Hilbert space and $Y \neq \{0\}$ be a closed subspace, and let P denote orthogonal projection onto Y . Show that:

- (1) P is linear;
- (2) P is bounded and $\|P\| = 1$;
- (3) For all $x \in H$ we have $(Px, x) = \|Px\|^2$;
- (4) $I - P$ is orthogonal projection onto $Y^\perp$.

Proof. (a) For any $x_1, x_2 \in H$ and $\alpha \in \mathbb{C}$ we have:

$$\begin{aligned} P(x_1 + \alpha x_2) &= P(y_1 + y_1^\perp + \alpha y_2 + \alpha y_2^\perp) \\ &= P(y_1 + \alpha y_2 + (y_1 + \alpha y_2)^\perp) \\ &= y_1 + \alpha y_2 \\ &= P(x_1) + \alpha P(x_2). \end{aligned}$$

Thus T is linear.

(b) Let $x \in H$. Observe that:

$$\begin{aligned}\|Px\|^2 &= \|y\|^2 \\ &\leq \|y\|^2 + \|y^\perp\|^2 \\ &= \|y + y^\perp\|^2 \\ &= \|x\|^2.\end{aligned}$$

Squaring both sides gives $\|Px\| \leq \|x\|$. Dividing by $\|x\|$ on both sides of this inequality and taking the supremum over all $x \in H, x \neq 0$ gives $\|P\|_{\mathcal{L}(H,Y)} \leq 1$. Now let $y \in B_Y(0,1)$. Then:

$$\|Py\| = \|y\| = 1.$$

Hence $1 \in \{\|Px\| \mid x \in B_H(0,1)\}$. It must be the case that $1 \leq \|P\|_{\mathcal{L}(H,Y)}$. Therefore $\|P\|_{\mathcal{L}(H,Y)} = 1$.

(c) We will first show that P is self-adjoint and idempotent. Let $x_1, x_2 \in H$. We have:

$$\begin{aligned}(Px_1, x_2) &= (y, y_2 + y_2^\perp) \\ &= (y_1, y_2) + (y_1, y_2^\perp) \\ &= (y_1, y_2).\end{aligned}$$

$$\begin{aligned}(x_1, Px_2) &= (y_1 + y_1^\perp, y_2) \\ &= (y_1, y_2) + (y_1^\perp, y_2) \\ &= (y_1, y_2).\end{aligned}$$

Therefore $P^* = P$; i.e., P is self-adjoint. Now for any $x \in H$, notice that:

$$\begin{aligned}P(P(x)) &= P(y) \\ &= y \\ &= P(x).\end{aligned}$$

Thus $P^2 = P$. Using these two facts:

$$\begin{aligned}\|Px\|^2 &= (Px, Px) \\ &= (P^*Px, x) \\ &= (P^2x, x) \\ &= (Px, x).\end{aligned}$$

□

(d) For any $x \in H$:

$$\begin{aligned}(I - P)(x) &= Ix - Px \\ &= y + y^\perp - P(y + y^\perp) \\ &= y + y^\perp - y \\ &= y^\perp\end{aligned}$$

Exercise 2.2.6. Let H be a Hilbert Space with orthonormal basis $\{u_n\}_{n=1}^\infty$. We try to define a map:

$$T : H \rightarrow H \text{ by } Tx = \sum_{n=1}^{\infty} \lambda_n (x, u_n) u_n$$

for a sequence $(\lambda_n) \subset \mathbb{C}$. First show that this map is a bounded linear operator if and only if (λ_n) is a bounded sequence. Then, under what condition is T self-adjoint?

Proof. Suppose $T \in \mathcal{L}(H)$. Then:

$$\begin{aligned} |\lambda_n| &= |\lambda_n| \|u_n\| \\ &= \|\lambda_n u_n\| \\ &= \|Tu_n\| \\ &\leq \|T\| \|u_n\| \\ &= \|T\|. \end{aligned}$$

Taking the supremum over all $n \in \mathbb{N}$ gives $\sup_{n \in \mathbb{N}} |\lambda_n| \leq \|T\|_{\mathcal{L}(H)} < \infty$. Therefore (λ_n) is bounded. Conversely, suppose $\sup_{n \in \mathbb{N}} |\lambda_n| := M < \infty$. We must first show that T is linear. Let $x, y \in H$ and $\alpha \in \mathbb{C}$. Define $a_n := (x, u_n)$ and $b_n := (y, u_n)$. We can write:

$$\begin{aligned} x + \alpha y &= \sum_{n=1}^{\infty} a_n u_n + \alpha \sum_{n=1}^{\infty} b_n u_n \\ &= \sum_{n=1}^{\infty} (a_n + \alpha b_n) u_n. \end{aligned}$$

Applying T to $x + \alpha y$ gives:

$$\begin{aligned} T(x + \alpha y) &= \sum_{n=1}^{\infty} \lambda_n (a_n + \alpha b_n) u_n \\ &= \sum_{n=1}^{\infty} \lambda_n a_n u_n + \alpha \sum_{n=1}^{\infty} \lambda_n b_n u_n \\ &= Tx + \alpha Ty. \end{aligned}$$

Thus T is linear. Now let $x \in X$ and again define $a_n := (x, u_n)$. By Parseval's identity:

$$\begin{aligned}
 \|Tx\|^2 &= \sum_{n=1}^{\infty} |(Tx, u_n)|^2 \\
 &= \sum_{n=1}^{\infty} \left| \left(\sum_{j=1}^{\infty} \lambda_j a_n u_j, u_n \right) \right|^2 \\
 &= \sum_{n=1}^{\infty} \left| \sum_{j=1}^{\infty} (\lambda_j a_j u_j, u_n) \right|^2 \\
 &= \sum_{n=1}^{\infty} |\lambda_n a_n|^2 \\
 &\leq \sum_{n=1}^{\infty} |M|^2 |a_n|^2 \\
 &= |M|^2 \sum_{n=1}^{\infty} |a_n|^2 \\
 &= |M|^2 \|x\|^2.
 \end{aligned}$$

Rearranging and taking the square-root of both sides yields:

$$\frac{\|Tx\|}{\|x\|} \leq |M|.$$

Taking the supremum over all $x \in X$, $x \neq 0$ gives $\|T\|_{\mathcal{L}(H)} \leq |M| < \infty$. Thus $T \in \mathcal{L}(H)$.

Let $x, y \in H$. Observe that:

$$\begin{aligned}
 (Tx, y) &= \left(\sum_{n=1}^{\infty} \lambda_n (x, u_n) u_n, y \right) \\
 &= \sum_{n=1}^{\infty} (\lambda_n (x, u_n) u_n, y) \\
 &= \sum_{n=1}^{\infty} \lambda_n (x, u_n) (u_n, y). \\
 (x, Ty) &= \left(x, \sum_{n=1}^{\infty} \lambda_n (y, u_n) u_n \right) \\
 &= \sum_{n=1}^{\infty} (x, \lambda_n (y, u_n) u_n) \\
 &= \sum_{n=1}^{\infty} \overline{\lambda_n (y, u_n)} (x, u_n) \\
 &= \sum_{n=1}^{\infty} \overline{\lambda_n} (x, u_n) (y, u_n).
 \end{aligned}$$

From this, T is self-adjoint if and only if $\lambda_n = \overline{\lambda_n}$; i.e., whenever $(\lambda_n) \subset \mathbb{R}$. $\square$

§ 2.3. Compact Operators

Definition 2.3.1. Let X and Y be normed linear spaces and $T : X \rightarrow Y$ linear. We say T is compact if for every bounded sequence $\{x_n\}_{n=1}^\infty \subset X$, the sequence $\{Tx_n\}_{n=1}^\infty$ has a convergent subsequence. Equivalently, the operator T is said to be compact if $\overline{T(B_X(0,1))}$ is compact in Y .

Proposition 2.3.1. Let X and Y be normed linear spaces and $T : X \rightarrow Y$ linear.

- (1) If T is compact, then T is bounded.
- (2) If $T \in \mathcal{L}(X, Y)$ and $R(T)$ is finite-dimensional, then T is compact.

Proof. (1) Let $x \in X$, $\|x\| = 1$. Then $Tx \in T(B_X(0,1)) \subseteq \overline{T(B_X(0,1))}$. Since T is compact, $\overline{T(B_X(0,1))}$ is compact in Y . Since $\overline{T(B_X(0,1))}$ is compact, it is bounded. Hence $\|Tx\| < \infty$.

(2) Since $T \in \mathcal{L}(X, Y)$, the subspace $T(B_X(0,1))$ is bounded. Since $R(T)$ is finite-dimensional, by Corollary 1.2.2 $\overline{T(B_X(0,1))}$ is closed. Hence $\overline{T(B_X(0,1))} = T(B_X(0,1))$ is closed and bounded. By Theorem 1.2.1, $\overline{T(B_X(0,1))}$ is compact. Thus T is compact. $\square$

Example 2.3.1. Let X be a normed linear space. Then $\text{id} : X \rightarrow X$ is bounded, but not compact.

Define $T : \ell^2 \rightarrow \ell^2$ by $e^n \mapsto \frac{1}{\sqrt{n}}e^n$ and extend linearly. This map is compact, but $R(T)$ is not finite dimensional.

Theorem 2.3.1. Let X be a normed linear space and Y a Banach space. The set:

$$\{T \in \mathcal{L}(X, Y) \mid T \text{ is compact}\}$$

is closed in $\mathcal{L}(X, Y)$.

Proof. Let $\{T_n\}_{n=1}^\infty$ be a sequence of compact operators converging to $T \in \mathcal{L}(X, Y)$. Let $\{x_n\}_{n=1}^\infty$ be a bounded sequence. Choose a subsequence $\{x_{1,j}\}_{j=1}^\infty \subset \{x_n\}_{n=1}^\infty$ such that $\{T_1(x_{1,j})\}_{j=1}^\infty$ is convergent in Y . Note that $\{x_{1,j}\}_{j=1}^\infty$ is still bounded. Choose a subsequence $\{x_{2,j}\}_{j=1}^\infty \subset \{x_{1,j}\}_{j=1}^\infty \subset \{x_n\}_{n=1}^\infty$ such that $\{T_2(x_{2,j})\}_{j=1}^\infty$ is convergent in Y . In general, choose $\{x_{k,j}\}_{j=1}^\infty \subset \{x_{k-1,j}\}_{j=1}^\infty \subset \dots \subset \{x_n\}_{n=1}^\infty$ such that $\{T_k(x_{k,j})\}_{j=1}^\infty$ is convergent. Define $\{y_n\}_{n=1}^\infty$ by $y_n = x_{n,n}$. Then $\{y_n\}_{n=1}^\infty \subset \{x_n\}_{n=1}^\infty$. Then for any $k < n$, we have that $\{T_k y_n\}_{n=1}^\infty$ is convergent. $\square$