

§ 1. The Product Topology and the Tychonoff Theorem

Problem 19.3. *If each space X_α is a Hausdorff space, then $\prod X_\alpha$ is a Hausdorff space in both the box and product topologies.*

Proof. Given $\prod X_\alpha$ with the product topology, let $f, g \in \prod X_\alpha$ be distinct elements. Then there exists some α such that $f(\alpha), g(\alpha) \in X_\alpha$ with $f(\alpha) \neq g(\alpha)$. We can find open sets U_α containing $f(\alpha)$ and V_α containing $g(\alpha)$ such that $U_\alpha \cap V_\alpha = \emptyset$. Quickly note that:

$$\begin{aligned}\pi_\alpha^{-1}(U_\alpha) \cap \pi_\alpha^{-1}(V_\alpha) &= \pi_\alpha^{-1}(U_\alpha \cap V_\alpha) \\ &= \pi_\alpha^{-1}(\emptyset) \\ &= \emptyset.\end{aligned}$$

Both $\pi_\alpha^{-1}(U_\alpha)$ and $\pi_\alpha^{-1}(V_\alpha)$ are open in the product space $\prod X_\alpha$, and likewise $f \in \pi_\alpha^{-1}(U_\alpha)$ and $g \in \pi_\alpha^{-1}(V_\alpha)$. Thus the product space $\prod X_\alpha$ is Hausdorff. Since the box topology is finer than the product topology, $\prod X_\alpha$ is also Hausdorff with the box topology. ■

Problem 19.5. *One of the implications stated in Theorem 19.6 holds for the box topology. Which one?*

Proof. Example 2 which immediately follows Theorem 19.6 gives a counter example for the backwards implication. It must be that the forward implication holds for the box topology. ■

Problem 19.7. *Let $\mathbb{R}^\omega$ be the subset of $\mathbb{R}^\omega$ consisting of all sequences that are "eventually zero," that is, all sequences $(x_1, x_2, \dots)$ such that $x_i \neq 0$ for only finitely many values of i . What is the closure of $\mathbb{R}^\omega$ in $\mathbb{R}^\omega$ in the box and product topologies? Justify your answer.*

Proof. ■

§ 2. The Urysohn Lemma

Problem 33.2(a). *Show that a connected normal space having more than one point is uncountable.*

Proof. Let X be connected and normal. Let x_0 and x_1 be distinct points in X . Since X is normal, $\{x_0\}$ and $\{x_1\}$ are closed in X . Moreover, $\{x_0\} \cap \{x_1\} = \emptyset$. By Urysohn's Lemma, there exists a continuous function $f : X \rightarrow [a, b]$ such that $f(x_0) = a$ and $f(x_1) = b$. Since f is continuous and X is connected, $f(X)$ is also connected. It must be the case that $f(X)$ is an interval —otherwise $f(X)$ contains only a single element, which is immediately a contradiction as x_0 and x_1 are distinct points which map to two distinct elements in $[a, b]$. Since $f(X)$ is an interval, it is uncountable, which implies X is uncountable. ■

Problem 33.3. *Give a direct proof of the Urysohn lemma for a metric space (X, d) by setting:*

$$f(x) = \frac{d(x, A)}{d(x, A) + d(x, B)}.$$

Proof. Let X be a metric space. Let A and B be disjoint closed subsets of X . Since A and B are closed, $d(x, A) > 0$ for any $x \notin A$ and $d(x, B) > 0$ for any $x \notin B$. Moreover, $d(\cdot, A)$ and $d(\cdot, B)$ are continuous functions. Thus $f : X \rightarrow [0, 1]$ as given above is continuous, and $f(x) = 0$ for all $x \in A$ and $f(x) = 1$ for all $x \in B$. ■

Problem 33.8. Let X be completely regular, let K and C be disjoint closed subsets of X . Show that if K is compact, there is a continuous function $f : X \rightarrow [0, 1]$ such that $f(K) = \{0\}$ and $f(C) = \{1\}$.

Proof. Let $k \in K$. Then $k \notin C$. By the complete regularity of X , there exists $f'_k : X \rightarrow [0, 1]$ such that $f'_k(k) = 1$ and $f'_k(C) = \{0\}$. Define $f_k := 1 - f'_k$. Then $f_k(k) = 0$ and $f_k(C) = \{1\}$. Note that $[0, \frac{1}{2})$ is open in $[0, 1]$, hence $f_k^{-1}([0, \frac{1}{2}))$ is open in X . Moreover, it is clear that $k \in f_k^{-1}([0, \frac{1}{2}))$. We've obtained a cover $K \subseteq \bigcup_{k \in K} f_k^{-1}([0, \frac{1}{2}))$. Since K is compact, there exists points $\{k_1, \dots, k_n\}$ such that $K \subseteq \bigcup_{i=1}^n f_{k_i}^{-1}([0, \frac{1}{2}))$. Define $f := \min\{f_{k_1}, \dots, f_{k_n}\}$. Then $f(K) = [0, \frac{1}{2})$ and $f(C) = \{1\}$. Define $\varphi : [0, 1] \rightarrow [0, 1]$ by:

$$\varphi(t) := \begin{cases} 0, & 0 \leq t < \frac{1}{2} \\ 1, & t = 1 \end{cases}.$$

Since φ is continuous, the composition $\varphi \circ f$ is continuous. In particular, we have $(\varphi \circ f)(K) = \varphi([0, \frac{1}{2})) = \{0\}$ and $(\varphi \circ f)(C) = \varphi(\{1\}) = \{1\}$. This establishes the claim. ■

§ 3. The Baire Category Theorem

Problem 3.1. Show that if $\bigcup B_n$ is a nonempty Baire space, at least one of the sets $\overline{B_n}$ has a nonempty interior.

Proof. Suppose towards contradiction $\overline{B_n}^\circ = \emptyset$ for each n . Consider the collection of closed sets $\{\overline{B_n}\}_n$. Since $\bigcup B_n$ is a Baire space, it must be that $\left(\bigcup \overline{B_n}\right)^\circ = \emptyset$. But this means:

$$\begin{aligned} \bigcup B_n &\subseteq \bigcup \overline{B_n}^\circ \\ &\subseteq \left(\bigcup \overline{B_n}\right)^\circ \\ &= \emptyset. \end{aligned}$$

Thus $\bigcup B_n = \emptyset$, which is a contradiction. Therefore at least one of the sets $\overline{B_n}$ has a nonempty interior. ■

Problem 3.2. The Baire Category Theorem implies that $\mathbb{R}$ cannot be written as the countable union of closed subsets having empty interiors. Show this fails if the sets are not required to be closed.

Proof. ■

Problem 3.3. Show that the irrationals are a Baire space.

Proof. We will first show that any subset of a complete metric space which is a G_δ set is a Baire space. ■

Problem 3.4. Let (M, d) be a complete metric space with no isolated points. Prove that M is uncountable. (A point x in a topological space X is called a isolated point if the one point set $\{x\}$ is open in X).

Proof. Suppose towards contradiction M is countable. Since M has no isolated points, for each $x \in M$ the set $\{x\}^c$ is open. Moreover, $\overline{\{x\}^c} = (\{x\}^\circ)^c = \emptyset^c = M$. Hence $\{x\}$ is open and dense in M . But:

$$\begin{aligned} \bigcap_{i=1}^{\infty} \{x_i\}^c &= \left(\bigcup_{i=1}^{\infty} \{x_i\} \right)^c \\ &= M^c \\ &= \emptyset. \end{aligned}$$

Since we've found a countable collection of open and dense sets whose countable intersection is *not* dense, M is not a Baire Space. This contradicts the Baire Category Theorem, as M being a complete metric spaces implies that it is a Baire space. It must be the case that M is uncountable. ■

Problem 3.5. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be an infinitely differentiable function, and suppose that for each $x \in \mathbb{R}$, there exists a natural number n such that $f^{(n)}(x) = 0$. For each fixed $n \in \mathbb{N}$, consider the set

$$F_n := \{x \in \mathbb{R} \mid f^{(n)}(x) = 0\}$$

and let $I := [a, b]$.

(a) Show that $I = \bigcup_{n=0}^{\infty} (I \cap F_n)$.

(b) Show that there is a nonempty open interval $A \subseteq I$ where f is a polynomial.

Proof. ■