

Preface

Test

Contents

Contents	i
1 Complex Variables	1
1.1 Basic Definitions and Results	1
1.2 Winding Numbers and Cauchy’s Integral Formula	3
1.3 The Jordan Curve Theorem	4
1.4 Banach-Valued Integration	4
2 Banach Algebras	7
2.1 Basic Definitions and Results	7
2.2 Ideals and Quotients	8
2.3 The Spectrum	12
2.4 The Riesz Functional Calculus	12

Chapter 1

Complex Variables

§ 1.1. Basic Definitions and Results

Definition 1.1.1. Let $G \subset \mathbb{C}$ be open and $f : G \rightarrow \mathbb{C}$.

- (1) The function f is differentiable at a point $a \in G$ if

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

exists. The value of this limit is denoted $f'(a)$. If f is differentiable for each $a \in G$, then we simply say f is differentiable.

- (2) The function f is analytic if f is continuously differentiable.

Definition 1.1.2. A path is a continuous function $\gamma : [a, b] \rightarrow \mathbb{C}$.

Definition 1.1.3. Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a path.

- (1) If $\gamma'(t)$ exists for each $t \in [a, b]$ and $\gamma' : [a, b] \rightarrow \mathbb{C}$ is continuous, then γ is a smooth path.
- (2) If there is a partition of $[a, b]$ such that γ is smooth along each subinterval, then γ is piecewise smooth.
- (3) If γ is of bounded variation, then it is called a rectifiable path.
- (4) The set $\{\gamma(t) : a \leq t \leq b\}$ is called the trace of γ and is denoted by $\{\gamma\}$.
- (5) The path γ is said to be closed if $\gamma(a) = \gamma(b)$.

Theorem 1.1.1. Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a rectifiable path, and let $f : [a, b] \rightarrow \mathbb{C}$ be continuous. There exists a unique $I \in \mathbb{C}$ such that, for every $\epsilon > 0$, there exists $\delta > 0$ with the following property: if $\{t_0 < t_1, \dots, < t_m\}$ is a partition of $[a, b]$ with $\max\{t_k - t_{k-1} : 1 \leq k \leq m\} < \delta$, then for every choice of points $x_k \in [t_{k-1}, t_k]$ we have:

$$\left| I - \sum_{k=1}^m f(x_k)(\gamma(t_k) - \gamma(t_{k-1})) \right| < \epsilon.$$

Proof.



Definition 1.1.4. The number $I \in \mathbb{C}$ established in Theorem 1.1.1 is called the Riemann-Stieltjes integral of f with respect to γ over $[a, b]$ and is denoted by:

$$\int_a^b f d\gamma.$$

Theorem 1.1.2. If γ is a piecewise smooth path and $f : [a, b] \rightarrow \mathbb{C}$ is continuous then:

$$\int_a^b f d\gamma = \int_a^b f(t)\gamma'(t) dt.$$

Proof. ■

Definition 1.1.5. If $\gamma : [a, b] \rightarrow \mathbb{C}$ is a rectifiable path and $f : \{\gamma\} \rightarrow \mathbb{C}$ is continuous, then the line integral of f along γ is:

$$\int_a^b f(\gamma(t)) d\gamma(t).$$

This line integral is also denoted by $\int_\gamma f = \int_\gamma f(z) dz$.

Example 1.1.1. Let $a \in \mathbb{C}$ be arbitrary and consider the path $\gamma : [0, 1] \rightarrow \mathbb{C}$ given by $\gamma(t) := a + e^{2\pi i n t}$. Let $f : \mathbb{C} \setminus \{a\} \rightarrow \mathbb{C}$ be given by $f(z) := (z - a)^{-1}$. The line integral of f along γ is:

$$\begin{aligned} \int_\gamma f(z) dz &= \int_0^1 f(\gamma(t)) d\gamma(t) \\ &= \int_0^1 e^{-2\pi i n t} d\gamma(t). \end{aligned}$$

Since $e^{-2\pi i n t}$ is smooth, Theorem ?? gives:

$$\begin{aligned} \int_0^1 e^{-2\pi i n t} d\gamma(t) &= \int_0^1 e^{-2\pi i n t} \gamma'(t) dt \\ &= \int_0^1 e^{-2\pi i n t} (2\pi i n e^{2\pi i n t}) dt \\ &= 2\pi i n. \end{aligned}$$

Lemma 1.1.3. If $\gamma : [a, b] \rightarrow \mathbb{C}$ is a rectifiable path and $\varphi : [c, d] \rightarrow [a, b]$ is continuous, non-decreasing, and surjective, then $\gamma \circ \varphi : [c, d] \rightarrow \mathbb{C}$ is a rectifiable path and $\{\gamma\} = \{\gamma \circ \varphi\}$.

Proof. ■

Proposition 1.1.4. Let $\gamma : [a, b] \rightarrow \mathbb{C}$ and $\sigma : [c, d] \rightarrow \mathbb{C}$ be rectifiable paths. Define $\sigma \sim \gamma$ if there exists a function $\varphi : [c, d] \rightarrow [a, b]$ such that φ is continuous, strictly increasing, $\varphi(c) = a$ and $\varphi(d) = b$, and $\sigma = \gamma \circ \varphi$. Then $\sim$ is an equivalence relation.

Proof. ■

Definition 1.1.6. Using the terminology of Proposition ??, two paths γ and σ are equivalent if $\gamma \sim \sigma$. The function φ is called a change of parameter. An equivalence class of paths is called a curve.

Remark. The trace of a curve is the trace of any of its representatives. If f is continuous on the trace of a curve, then the integral over f over the curve is the integral of f over any representative of the curve. A curve is smooth if and only if one of its representatives is smooth.

Lemma 1.1.5. If G is an open set in $\mathbb{C}$, $\gamma : [a, b] \rightarrow G$ is a rectifiable path, and $f : G \rightarrow \mathbb{C}$ is continuous, then for every $\epsilon > 0$ there is a polygonal path Γ in G such that $\Gamma(a) = \gamma(a)$, $\Gamma(b) = \gamma(b)$, and $\left| \int_{\gamma} f - \int_{\Gamma} f \right| < \epsilon$.

Proof. ■

Theorem 1.1.6. Let $G \subseteq \mathbb{C}$ be open and let γ be a rectifiable path in G with initial and end points α and β respectively. If $f : G \rightarrow \mathbb{C}$ is a continuous function with primitive $F : G \rightarrow \mathbb{C}$, then:

$$\int_{\gamma} f = F(\beta) - F(\alpha),$$

(Recall that F is a primitive of f when $F' = f$.)

Proof. ■

Corollary 1.1.7. Let $G \subseteq \mathbb{C}$ be open and let γ be a closed rectifiable path in G . If $f : G \rightarrow \mathbb{C}$ is a continuous function with primitive $F : G \rightarrow \mathbb{C}$, then $\int_{\gamma} f = 0$.

§ 1.2. Winding Numbers and Cauchy's Integral Formula

Proposition 1.2.1. If $\gamma : [0, 1] \rightarrow \mathbb{C}$ is a closed rectifiable curve and $a \notin \{\gamma\}$, then:

$$\frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - a}$$

is an integer.

Proof. ■

Definition 1.2.1. If γ is a closed rectifiable curve in $\mathbf{C}$, then for $a \notin \{\gamma\}$, the index of γ with respect to the point a is:

$$n(\gamma; a) := \frac{1}{2\pi i} \int_{\gamma} (z - a)^{-1} dz.$$

Cauchy's Theorem Stuff...

§ 1.3. The Jordan Curve Theorem

Definition 1.3.1. Let $G \subseteq \mathbf{C}$ be open. A closed rectifiable curve γ is positively oriented if for every $a \in G \setminus \{\gamma\}$, $n(\gamma; a)$ is either 0 or 1. In this case the inside of γ is $\text{ins } \gamma := \{a \in \mathbf{C} \setminus \{\gamma\} : n(\gamma; a) = 1\}$. The outside of γ is $\text{out } \gamma := \{a \in \mathbf{C} \setminus \{\gamma\} : n(\gamma; a) = 0\}$.

Definition 1.3.2. A curve $\gamma : [0, 1] \rightarrow \mathbf{C}$ is simple if $\gamma(s) = \gamma(t)$ implies either:

- $s = t$,
- $s = 0$ and $t = 1$, or
- $s = 1$ and $t = 0$.

Theorem 1.3.1 (Jordan Curve Theorem). *If γ is a simple closed rectifiable curve, then $\mathbf{C} \setminus \{\gamma\}$ consists of exactly two connected components, each of which is bounded by $\{\gamma\}$.*

§ 1.4. Banach-Valued Integration

Definition 1.4.1. Let $G \subseteq \mathbf{C}$ be open and X a Banach space. If $f : G \rightarrow X$, define the derivative of f at z_0 to be:

$$f'(z_0) := \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h},$$

provided the limit exists. If the derivative of f exists for all $z \in G$, we say that f is differentiable. The function f is called analytic if f is continuously differentiable on G .

Remark. Let X be a Banach space. For any $x \in X$, recall that the evaluation map is $\hat{x} : X^* \rightarrow \mathbf{F}$ given by $\hat{x}(x^*) := x^*(x)$. We introduce the following notation for convenience:

- For $x^* \in X^*$, define $\langle \cdot, x^* \rangle : X \rightarrow \mathbf{F}$ by $x \mapsto x^*(x)$.
- For $x \in X$, define $\langle x, * \rangle : X^* \rightarrow \mathbf{F}$ by $x^* \mapsto \hat{x}(x^*) = x^*(x)$.

We call $\langle \cdot, \cdot \rangle$ the canonical evaluation map.

Proposition 1.4.1. *Let $G \subseteq \mathbf{C}$ be open and X a Banach space. If $f : G \rightarrow X$ is analytic and $x^* \in X^*$, then $z \mapsto \langle f(z), x^* \rangle$ is analytic on G and its derivative is $\langle f'(z), x^* \rangle$.*

Proof. As $z \mapsto \langle f(z), x^* \rangle$ is the composition of continuous maps, it is surely continuous. If such a function is denoted by g , then for any $z_0 \in G$ we have:

$$\begin{aligned} g'(z_0) &= \lim_{h \rightarrow 0} \frac{g(z_0 + h) - g(z_0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\langle f(z_0 + h), x^* \rangle - \langle f(z_0), x^* \rangle}{h} \\ &= \lim_{h \rightarrow 0} \left\langle \frac{f(z_0 + h) - f(z_0)}{h}, x^* \right\rangle \\ &= \left\langle \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}, x^* \right\rangle \\ &= \langle f'(z_0), x^* \rangle. \end{aligned}$$

Since $g'(z_0)$ exists for all $z_0 \in G$, g is differentiable. Since g' is the composition of continuous maps, it is continuous. Thus $z \mapsto \langle f(z), x^* \rangle$ is analytic. ■

Proposition 1.4.2. *Let $G \subseteq \mathbb{C}$ be open and X a Banach space. If $f : G \rightarrow X$ is a function such that $z \mapsto \langle f(z), x^* \rangle$ is analytic for each $x^* \in X^*$, then f is analytic.*

Proof. Since $G \subseteq \mathbb{C}$ is open, choose $r > 0$ such that $\bar{B}(z, r) \subseteq G$. Let $0 < \delta < \frac{r}{2}$, and let s, t satisfy $0 < |s|, |t| < \delta$. Fix $x^* \in X^*$ with $\|x^*\| \leq 1$ and let $g := \langle f(\cdot), x^* \rangle$. Since $z, z-s, z-t \in B^\circ(z, r)$, we can apply Cauchy's Integral Formula to $g(z)$, $g(z-s)$, and $g(z-t)$ (respectively). With this, observe that:

$$\begin{aligned} \left| \frac{g(z+s) - g(z)}{s} - \frac{g(z+t) - g(z)}{t} \right| &= \left| \frac{1}{2\pi i} \int_{\partial \bar{B}(z, r)} \frac{g(w)(s-t)}{(w-z)(w-z-s)(w-z-t)} dw \right| \\ &\leq \frac{1}{2\pi} \int_{\partial \bar{B}(z, r)} \frac{|g(w)||s-t|}{|w-z||w-z-s||w-z-t|} dw \end{aligned}$$

Note that $|s-t| \leq 2\delta$ and $|w-z||w-z-s||w-z-t| \geq \frac{r^3}{8}$. Hence:

$$\begin{aligned} \left| \frac{g(z+s) - g(z)}{s} - \frac{g(z+t) - g(z)}{t} \right| &\leq \frac{8\delta}{\pi r^3} \int_{\partial \bar{B}(z, r)} |g(w)| dw \\ &\leq c\delta \sup_{w \in \partial \bar{B}(z, r)} |g(w)| \\ &= c\delta \sup_{w \in \partial \bar{B}(z, r)} |\langle f(w), x^* \rangle| \end{aligned}$$

Note that $\langle f(w), \cdot \rangle : X^* \rightarrow \mathbb{C}$ is continuous for all $x^* \in X^*$. Hence $|\langle f(w), x^* \rangle| = |x^*(f(w))| \leq \|x^*\| \|f(w)\| \leq \|f(w)\|$. The above inequality can be written as:

$$\left| \left\langle \frac{f(z+s) - f(z)}{s} - \frac{f(z+t) - f(z)}{t}, x^* \right\rangle \right| \leq c\delta \sup_{w \in \partial \bar{B}(z, r)} \|f(w)\|.$$

Since the right-hand side is independent of $x^* \in X^*$, we have:

$$\sup_{\|x^*\| \leq 1} \left| \left\langle \frac{f(z+s) - f(z)}{s} - \frac{f(z+t) - f(z)}{t}, x^* \right\rangle \right| \leq c\delta \sup_{w \in \partial \bar{B}(z, r)} \|f(w)\|.$$

By a corollary of the Hahn-Banach Theorem, this gives:

$$\left\| \frac{f(z+s)-f(z)}{s} - \frac{f(z+t)-f(z)}{t} \right\| \leq c\delta \sup_{w \in \partial \overline{B}(z,r)} \|f(w)\|.$$

Given any $\epsilon > 0$, choose $0 < \delta < \frac{\epsilon}{2}$ such that $\delta \leq \frac{\epsilon}{\sup_{w \in \partial \overline{B}(z,r)} \|f(w)\|}$. If $0 < |s|, |t| < \delta$, then $\left\| \frac{f(z+s)-f(z)}{s} - \frac{f(z+t)-f(z)}{t} \right\| < \epsilon$. Since X is complete, by the Cauchy Criterion for limits $f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h)-f(z)}{h} = 0$. Since $z \in G$ was arbitrary, f is analytic. ■

Theorem 1.4.3. *Let $G \subseteq \mathbf{C}$ be open and X a Banach space. Let γ be a rectifiable curve in G and f a continuous function defined in a neighborhood of $\{\gamma\}$ with values in X . There exists a unique $I \in X$ such that, for every $\epsilon > 0$, there exists $\delta > 0$ with the following property: if $\{t_0 < t_1 < \dots < t_m\}$ is a partition of $[0, 1]$ with $\max\{t_k - t_{k-1} : 1 \leq k \leq m\} < \delta$, then for every choice of points $x_k \in [t_{k-1}, t_k]$ we have:*

$$\left\| I - \sum_{k=1}^m f(\gamma(x_k))(\gamma(t_k) - \gamma(t_{k-1})) \right\| < \epsilon.$$

Proof. ■

Definition 1.4.2. The element $I \in X$ established in Theorem ?? is called the line integral of f along γ and is denoted by:

$$\int_0^1 f(\gamma(t)) d\gamma(t).$$

This line integral is also denoted by $\int_\gamma f = \int_\gamma f(z) dz$.

Proposition 1.4.4. *Let $G \subseteq \mathbf{C}$ be open and γ a rectifiable curve in G . Suppose f is a continuous function defined in a neighborhood of $\{\gamma\}$ with values in X . For every $x^* \in X^*$, we have:*

$$\left\langle \int_\gamma f, x^* \right\rangle = \int_\gamma \langle f(\cdot), x^* \rangle.$$

Theorem 1.4.5 (Cauchy's Theorem). *If X is a Banach space, G is an open subset of $\mathbf{C}$, $f : G \rightarrow X$ is an analytic function, and $\gamma_1, \dots, \gamma_m$ are closed rectifiable curves in G such that $\sum_{j=1}^m n(\gamma_j; a) = 0$ for all $a \in \mathbf{C} \setminus G$, then $\sum_{j=1}^m \int_{\gamma_j} f = 0$.*

Theorem 1.4.6 (Cauchy's Integral Formula). *If X is a Banach space, G is an open subset of $\mathbf{C}$, $f : G \rightarrow X$ is analytic, γ is a closed rectifiable curve in G such that $n(\gamma; a) = 0$ for every $a \in \mathbf{C} \setminus G$, and $\lambda \in G \setminus \{\gamma\}$, then for every integer $k \geq 0$:*

$$n(\gamma; \lambda) f^{(k)}(\lambda) = \frac{k!}{2\pi i} \int_\gamma (z - \lambda)^{-(k+1)} f(z) dz.$$

Chapter 2

Banach Algebras

§ 2.1. Basic Definitions and Results

Definition 2.1.1. An algebra over $\mathbf{F}$ is a vector space $\mathcal{A}$ over $\mathbf{F}$ that has a multiplication operation making $\mathcal{A}$ into a ring such that, for any $\alpha \in \mathbf{F}$ and $a, b \in \mathcal{A}$, we have $\alpha(ab) = (\alpha a)b = a(\alpha b)$. If $\mathcal{A}$ admits an identity element, we say $\mathcal{A}$ is unital.

Definition 2.1.2. A normed algebra is an algebra $\mathcal{A}$ over $\mathbf{F}$ which has a norm $\|\cdot\|$ making $\mathcal{A}$ into a normed linear space and also admitting the property that, for any $a, b \in \mathcal{A}$, we have $\|ab\| \leq \|a\|\|b\|$. If $\mathcal{A}$ is a Banach space, then $\mathcal{A}$ is called a Banach algebra. If $\mathcal{A}$ has an identity e , then it is assumed that $\|e\| = 1$.

Proposition 2.1.1. If $\mathcal{A}$ is an algebra without identity, define algebraic operations on $\mathcal{A} \times \mathbf{F}$ by:

- (i) $(a, \alpha) + (b, \beta) := (a + b, \alpha + \beta)$,
- (ii) $\beta(a, \alpha) := (\beta a, \beta \alpha)$,
- (iii) $(a, \alpha)(b, \beta) := (ab + \alpha\beta + \beta a, \alpha\beta)$.

Then $\mathcal{A} \times \mathbf{F}$ is an algebra with identity $(0, 1)$.

Proof. ■

Proposition 2.1.2. Let $\mathcal{A}$ be an algebra over $\mathbf{F}$ in which $(\mathcal{A}, \|\cdot\|_A)$ is a Banach space. For every $a \in \mathcal{A}$, suppose that the maps $x \mapsto ax$ and $x \mapsto xa$ of $\mathcal{A} \rightarrow \mathcal{A}$ are continuous. Let $\mathcal{A} \times \mathbf{F}$ be the algebra with operations given from Proposition ???. For $a \in \mathcal{A}$, define $L_a : \mathcal{A} \times \mathbf{F} \rightarrow \mathcal{A} \times \mathbf{F}$ by $L_a(x, \xi) := (ax + \xi a, 0)$.

- (1) Define $\|(a, \alpha)\|_{\mathcal{A} \times \mathbf{F}} := \|a\|_A + |\alpha|$. Then $\mathcal{A} \times \mathbf{F}$ is a Banach space and an algebra with identity $(0, 1)$.
- (2) The map $a \rightarrow (a, 0)$ is an isometric isomorphism between $\mathcal{A}$ and $\mathcal{A} \times \{0\}$.
- (3) For each $a \in \mathcal{A}$, we have $L_a \in \mathcal{B}(\mathcal{A} \times \mathbf{F})$.
- (4) Define $\|a\|_* := \|L_a\|_{\mathcal{B}(\mathcal{A} \times \mathbf{F})}$. Then $\|\cdot\|_*$ is equivalent to the norm $\|\cdot\|_A$, and in particular $(\mathcal{A}, \|\cdot\|_*)$ is a Banach algebra.

Proof. ■

Parts (1) and (2) show that $\mathcal{A}$ may be identified isometrically with the ideal $\mathcal{A} \times \{0\}$ of the unital algebra $\mathcal{A} \times \mathbb{F}$, whose identity is $(0, 1)$. Thus, every algebra may be embedded isometrically into a unital algebra. When an algebra is already unital, it makes sense then to denote its identity by 1. We will use this convention throughout the rest of the paper.

Parts (3) and (4) show that the usual submultiplicativity condition on the norm need not be assumed. Indeed, if $\mathcal{A}$ is complete and multiplication is continuous, then the norm $\|a\|_*$ is equivalent to the original norm on $\mathcal{A}$ satisfying the submultiplicativity condition.

§ 2.2. Ideals and Quotients

Definition 2.2.1. Let $\mathcal{A}$ be an algebra. A left ideal (resp. right ideal) of $\mathcal{A}$ is a subalgebra $\mathcal{M}$ of $\mathcal{A}$ such that, for all $a \in \mathcal{A}$ and $x \in \mathcal{M}$, then $ax \in \mathcal{M}$ (resp. $xa \in \mathcal{M}$). If $\mathcal{M}$ is both a left and right ideal, then $\mathcal{M}$ is simply called an ideal (or bilateral ideal).

Definition 2.2.2. Let $\mathcal{A}$ be a unital algebra. An element $a \in \mathcal{A}$ is left invertible (resp. right invertible) if there exists an element $x \in \mathcal{A}$ such that $xa = 1$ (resp. $ax = 1$). If an element $a \in \mathcal{A}$ is both left and right invertible, then a is simply called invertible.

Proposition 2.2.1. Let $\mathcal{A}$ be a unital algebra.

- (1) If $a \in \mathcal{A}$ is invertible, there is a unique element a^{-1} such that $aa^{-1} = a^{-1}a = 1$.
- (2) If $a \in \mathcal{A}$ is left invertible, and $\mathcal{M}$ is a left ideal in $\mathcal{A}$ containing a , then $\mathcal{M} = \mathcal{A}$.

Proof. (1) Since a is invertible, there exists $x, y \in \mathcal{A}$ such that $xa = 1 = ay$. Hence $y = 1y = (xa)y = x(ay) = x1 = x$.

(2) Since $a \in \mathcal{A}$ is left invertible, there exists $x \in \mathcal{A}$ such that $xa = 1$. Since $\mathcal{M}$ is a left ideal and $a \in \mathcal{M}$, it must be the case that $1 = xa \in \mathcal{M}$. Since $1 \in \mathcal{M}$, we have $y = y1 \in \mathcal{M}$ for all $y \in \mathcal{A}$. Thus $\mathcal{A} \subseteq \mathcal{M}$. ■

Part (2) of Proposition ?? says that every element of a proper left ideal is necessarily not left invertible.

Lemma 2.2.2. If $\mathcal{A}$ is a unital Banach algebra and $x \in \mathcal{A}$ such that $\|x - 1\| < 1$, then x is invertible.

Proof. Define $y := 1 - x$. Note that:

$$\begin{aligned} \|y\| &= \|1 - x\| \\ &= \|x - 1\| \\ &< 1. \end{aligned}$$

Since $\mathcal{A}$ is a Banach algebra, it follows via induction that:

$$\|y^n\| \leq \|y\|^n.$$

Since $\|y\| < 1$, the series $\sum_{n=0}^{\infty} \|y\|^n$ converges, hence by the comparison test $\sum_{n=0}^{\infty} \|y^n\|$ converges. Because $\mathcal{A}$ is complete, $z := \sum_{n=0}^{\infty} y^n$ also converges. Taking $z_k := \sum_{n=0}^k y^n$, we can

see:

$$\begin{aligned} z_k(1-y) &= \sum_{n=0}^k y^n - \sum_{n=0}^k y^{n+1} \\ &= 1 - y^{k+1}. \end{aligned}$$

It is clear that $\sum_{n=0}^{\infty} y^{n+1}$ converges by the same reasoning from above. Hence $y^{n+1} \rightarrow 0$ as $n \rightarrow \infty$. This gives:

$$\begin{aligned} z(1-y) &= \lim_{n \rightarrow \infty} z_n(1-y) \\ &= \lim_{n \rightarrow \infty} 1 - y^{n+1} \\ &= 1. \end{aligned}$$

An identical argument will show that $(1-y)z = 1$. Thus $(1-y) = 1 - (1-x) = x$ is invertible. ■

The following fact surfaced in the preceding proof.

Corollary 2.2.3. *Let $\mathcal{A}$ be a unital Banach algebra. If $a \in \mathcal{A}$ and $\|a\| < 1$, then $(1-a)^{-1} = \sum_{k=0}^{\infty} a^k$.*

Lemma 2.2.4. *Let $\mathcal{A}$ be a unital Banach algebra. If $b_0 a_0 = 1$ and $\|a - a_0\| < \|b_0\|^{-1}$, then a is left invertible.*

Proof. Rearranging $\|a - a_0\| < \|b_0\|^{-1}$ yields $\|b_0 a - 1\| < 1$. By Lemma 2.2.2, the element $b_0 a$ is invertible. Hence there exists $x \in \mathcal{A}$ such that $x b_0 a = 1$, implying that a is left invertible. ■

Theorem 2.2.5. *If $\mathcal{A}$ is a unital Banach algebra, then:*

- (i) $G_l := \{a \in \mathcal{A} : a \text{ is left invertible}\},$
- (ii) $G_r := \{a \in \mathcal{A} : a \text{ is right invertible}\},$
- (iii) $G := \{a \in \mathcal{A} : a \text{ is invertible}\},$

are all open subsets of $\mathcal{A}$. Moreover, the map $a \mapsto a^{-1}$ of $G \rightarrow G$ is continuous.

Proof. Let $a_0 \in G_l$. If $a \in B(a_0, \|b_0\|^{-1})$, then $\|a - a_0\| < \|b_0\|^{-1}$, so by Lemma 2.2.4 a is left invertible. Thus $a_0 \in B(a_0, \|b_0\|^{-1}) \subseteq G_l$, establishing G_l as open. It follows similarly that G_r is open, and since $G = G_l \cap G_r$, it is open as well.

To prove that $a \mapsto a^{-1}$ is continuous, we first prove an intermediary result. Let $\{a_n\}$ be a sequence in G such that $a_n \rightarrow 1$. Let $0 < \delta < 1$ and suppose $\|a_n - 1\| < \delta$. By Corollary 2.2.3:

$$\begin{aligned} a_n^{-1} &= (1 - (1 - a_n))^{-1} \\ &= \sum_{k=0}^{\infty} (1 - a_n)^k \\ &= 1 + \sum_{k=1}^{\infty} (1 - a_n)^k. \end{aligned}$$

Hence $a_n^{-1} - 1 = \sum_{k=1}^{\infty} (1 - a_n)^k$. Thus:

$$\begin{aligned} \|a_n^{-1} - 1\| &= \left\| \sum_{k=1}^{\infty} (1 - a_n)^k \right\| \\ &\leq \sum_{k=1}^{\infty} \|1 - a_n\|^k \\ &= \sum_{k=1}^{\infty} \delta^k \\ &= \frac{\delta}{1 - \delta}. \end{aligned}$$

If $\epsilon > 0$ is given, then δ can be chosen such that $\frac{\delta}{1 - \delta} < \epsilon$. So $\|a_n - 1\| < \delta$ implies $\|a_n^{-1} - 1\| < \epsilon$. Hence $a_n^{-1} \rightarrow 1$. With this, let $a \in G$ be arbitrary and suppose $\{a_n\}$ is a sequence in G such that $a_n \rightarrow a$. Then $a^{-1}a_n \rightarrow 1$. The intermediary result above gives $a_n^{-1}a = (a^{-1}a_n)^{-1} \rightarrow 1$. Hence $a_n^{-1} = a_n^{-1}aa^{-1} \rightarrow a^{-1}$. ■

Definition 2.2.3. A maximal ideal is a proper ideal that is contained in no larger proper ideal.

Corollary 2.2.6. Let $\mathcal{A}$ be a unital Banach algebra.

- (1) The closure of a proper left (resp. right, bilateral) ideal is a proper left (resp. right, bilateral) ideal.
- (2) A maximal left (resp. right, bilateral) ideal is closed.

Proof. (1) Let $\mathcal{M}$ be a proper left ideal of $\mathcal{A}$. The proof showing $\overline{\mathcal{M}}$ is a subalgebra is omitted, but the idea can be replicated from the following. Let $a \in \mathcal{A}$ and $x \in \overline{\mathcal{M}}$. There exists a sequence $\{x_n\}$ in $\mathcal{M}$ such that $x_n \rightarrow x$. Then $\{ax_n\}$ is a sequence in $\mathcal{M}$ such that $ax_n \rightarrow ax$. Thus $ax \in \overline{\mathcal{M}}$, so $\overline{\mathcal{M}}$ is a left ideal. Let G_l denote the elements of $\mathcal{A}$ which are left invertible. By part (2) of Proposition 2.2.1, $\mathcal{M} \cap G_l = \emptyset$, which implies that $\mathcal{M} \subseteq \mathcal{A} \setminus G_l$. Since $\mathcal{A} \setminus G_l$ is closed, and since $\overline{\mathcal{M}}$ is the smallest closed set containing $\mathcal{M}$, we have $\overline{\mathcal{M}} \subseteq \mathcal{A} \setminus G_l$. Since G_l is necessarily nonempty, $\overline{\mathcal{M}} \neq \mathcal{A}$, hence it is proper.

(2) If $\mathcal{M}$ is a maximal left ideal, then $\overline{\mathcal{M}}$ is a proper ideal by (1). Hence $\mathcal{M} = \overline{\mathcal{M}}$ by maximality. ■

Theorem 2.2.7. *If $\mathcal{A}$ is a Banach algebra and $\mathcal{M}$ is a proper closed ideal in $\mathcal{A}$, then $\mathcal{A}/\mathcal{M}$ is a Banach algebra with norm given by:*

$$\|a + \mathcal{M}\| := \inf\{\|a + x\| : x \in \mathcal{M}\}.$$

If $\mathcal{A}$ has an identity, so does $\mathcal{A}/\mathcal{M}$.

Proof. The proof that $\mathcal{A}/\mathcal{M}$ is a normed linear space with well-defined operations given by $(a + \mathcal{M}) + (b + \mathcal{M}) = (a + b) + \mathcal{M}$ and $\alpha(a + \mathcal{M}) = (\alpha a) + \mathcal{M}$ is omitted. We will first show that $\mathcal{A}/\mathcal{M}$ is a Banach space. Let $\{a_n + \mathcal{M}\}_n$ be a Cauchy sequence in $\mathcal{A}/\mathcal{M}$. There is a subsequence $\{a_{n_k} + \mathcal{M}\}_k$ such that:¹

$$\|(a_{n_k} + \mathcal{M}) - (a_{n_{k+1}} + \mathcal{M})\| = \|(a_{n_k} - a_{n_{k+1}}) + \mathcal{M}\| < 2^{-k}.$$

Let $y_1 = 0$. By the definition of $\|\cdot\|$ on $\mathcal{A}/\mathcal{M}$, choose $y_2 \in \mathcal{M}$ such that:

$$\|a_{n_1} - a_{n_2} + y_2\| \leq \|a_{n_1} - a_{n_2} + \mathcal{M}\| + 2^{-1} < 2 \cdot 2^{-1}.$$

With y_2 fixed, choose $y_3 \in \mathcal{M}$ such that:

$$\begin{aligned} \|(a_{n_2} + y_2) - (a_{n_3} + y_2)\| &= \|a_{n_2} - a_{n_3} + (y_2 - y_3)\| \\ &\leq \|a_{n_2} - a_{n_3} + \mathcal{M}\| + 2^{-2} \\ &< 2 \cdot 2^{-2}. \end{aligned}$$

Inductively, we obtain a sequence $\{y_k\}_k$ in $\mathcal{M}$ such that:

$$\|(a_{n_k} + y_k) - (a_{n_{k+1}} + y_{k+1})\| < 2 \cdot 2^{-k}.$$

In particular, for $j < k$ we have:

$$\begin{aligned} \|(a_{n_k} + y_j) - (a_{n_k} + y_k)\| &\leq \|(a_{n_j} + y_j) - (a_{n_{j+1}} + y_{j+1})\| + \dots + \|(a_{n_{k-1}} + y_{k-1}) - (a_{n_k} + y_k)\| \\ &\leq \sum_{i=j}^k 2 \cdot 2^{-i} \\ &\leq \sum_{i=j}^{\infty} 2 \cdot 2^{-i} \\ &= 2^{2-j}. \end{aligned}$$

Thus $\{a_{n_k} + y_k\}_k$ is a Cauchy sequence in $\mathcal{A}$. Since $\mathcal{A}$ is complete, there exists $a_0 \in \mathcal{A}$ such that $a_{n_k} + y_k \rightarrow a_0$. Since the projection $\pi : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{M}$ is continuous, we have:

$$\begin{aligned} \lim_{k \rightarrow \infty} a_{n_k} + \mathcal{M} &= \lim_{k \rightarrow \infty} \pi(a_{n_k} + y_k) \\ &= \pi\left(\lim_{k \rightarrow \infty} a_{n_k} + y_k\right) \\ &= \pi(a_0) \\ &= a_0 + \mathcal{M}. \end{aligned}$$

¹Suppose $\{x_n\}$ is a Cauchy sequence. For $\epsilon = 2^{-1}$, choose N_1 such that $m, n \geq N_1$ implies $\|x_m - x_n\| < 2^{-1}$. Now choose $n_1 \geq N_1$. For $\epsilon = 2^{-2}$, choose N_2 such that $m, n \geq N_2$ implies $\|x_m - x_n\| < 2^{-2}$. Choose $n_2 \geq \max\{N_2, n_1 + 1\}$. Since $n_1, n_2 \geq N_1$, we have $\|x_{n_2} - x_{n_1}\| < 2^{-1}$. Inductively, we obtain a subsequence $\{x_{n_k}\}_k$ satisfying the above property.

Since $\{a_n + \mathcal{M}\}_n$ is a Cauchy sequence with a convergent subsequence, it is convergent. Thus $\mathcal{A}/\mathcal{M}$ is a Banach space.

The proof that $\mathcal{A}/\mathcal{M}$ is an algebra with well-defined multiplication given by $(a + \mathcal{M})(b + \mathcal{M}) = ab + \mathcal{M}$ is omitted. Let $a, b \in \mathcal{A}$ and $u, v \in \mathcal{M}$. Then $(a + u)(b + v) = ab + (av + ub + uv) \in ab + \mathcal{M}$. Hence:

$$\begin{aligned} \|(a + \mathcal{M})(b + \mathcal{M})\| &= \|ab + \mathcal{M}\| \\ &\leq \|(a + u)(b + v)\|. \end{aligned}$$

Taking the infimum over all $u, v \in \mathcal{M}$ gives $\|(a + \mathcal{M})(b + \mathcal{M})\| \leq \|a + \mathcal{M}\| \|b + \mathcal{M}\|$. Thus $\mathcal{A}/\mathcal{M}$ is a Banach algebra. Moreover, $1 + \mathcal{M}$ is indeed the identity of $\mathcal{A}/\mathcal{M}$ as $(1 + \mathcal{M})(a + \mathcal{M}) = (1a + \mathcal{M}) = a + \mathcal{M} = (a1 + \mathcal{M}) = (a + \mathcal{M})(1 + \mathcal{M})$ ■

§ 2.3. The Spectrum

Definition 2.3.1. If $\mathcal{A}$ is a unital Banach algebra and $a \in \mathcal{A}$, the spectrum of a is:

$$\sigma(a) := \{a \in \mathbf{F} : a - \alpha \text{ is not invertible}\}.$$

The left spectrum of a is:

$$\sigma_l(a) := \{a \in \mathbf{F} : a - \alpha \text{ is not left invertible}\}.$$

The right spectrum of a , denoted $\sigma_r(a)$, is defined similarly.

Definition 2.3.2. If $\mathcal{A}$ is a unital Banach algebra and $a \in \mathcal{A}$, the resolvent set of a is $\rho(a) := \mathbf{F} \setminus \sigma(a)$. The left and right resolvents of a are $\rho_l(a) := \mathbf{F} \setminus \sigma_l(a)$ and $\rho_r(a) := \mathbf{F} \setminus \sigma_r(a)$.

Example 2.3.1. Let X be compact. If $f \in C(X)$, then $\sigma(f) = f(X)$. Indeed, let $\alpha \in f(X)$. Then $\alpha = f(x_0)$ for some $x_0 \in X$. Hence $0 = f(x_0) - \alpha \in (f - \alpha)(X)$. It must be the case that $(f - \alpha)^{-1}$ does not exist, so $\alpha \in \sigma(f)$. Conversely, if $f - \alpha$ is not invertible, then $(f - \alpha)^{-1}$ does not exist. This means $0 = f(x_0) - \alpha$ for some $x_0 \in X$, hence $\alpha = f(x_0)$.

Example 2.3.2. Let X be a Banach space. If $A \in \mathcal{B}(X)$, then $\sigma(A) = \{\alpha \in \mathbf{F} : \text{either } \ker(A - \alpha) \neq 0 \text{ or } \operatorname{im}(A - \alpha) \neq X\}$. Indeed, if $A - \alpha$ is not invertible, then it is either not injective or not surjective. In fact, this means that $\rho(A) = \mathbf{F} \setminus \sigma(A) = \{\alpha \in \mathbf{F} : A - \alpha \text{ is bijective}\}$. If $\alpha \in \rho(A)$, then $\alpha \notin \sigma(A)$, hence $\ker(A - \alpha) = 0$ and $\operatorname{im}(A - \alpha) = X$. Hence $A - \alpha$ is bijective. Conversely, suppose $A - \alpha$ is bijective. By the Open Mapping Theorem, $A - \alpha$ is an open map. For any open subset $U \subseteq X$, the set $((A - \alpha)^{-1})^{-1}(U) = (A - \alpha)(U)$ is open. Thus $(A - \alpha)^{-1} \in \mathcal{B}(X)$, hence $\alpha \in \rho(A)$.

Proposition 2.3.1. Let $\mathcal{A}$ be a Banach algebra with identity

§ 2.4. The Riesz Functional Calculus

Definition 2.4.1. A closed rectifiable curve γ is positively oriented if...