

# 432 Class 08 Slides

[thomaseLove.github.io/432](https://thomaseLove.github.io/432)

2021-02-25

# Setup

```
library(magrittr); library(janitor)
library(here); library(knitr)
library(naniar); library(simputation)

library(rms)
library(tidyverse)
```

# Today's Data

## Heart and Estrogen/Progestin Study (HERS)

- Clinical trial of hormone therapy for the prevention of recurrent heart attacks and deaths among 2763 post-menopausal women with existing coronary heart disease (see Hulley et al 1998 and many subsequent references, including Vittinghoff, Chapter 4.)
- We're excluding the women in the trial with a diabetes diagnosis.

```
hers_raw <- read_csv(here("data/hersdata.csv")) %>%  
  clean_names()
```

```
hers1 <- hers_raw %>%  
  filter(diabetes == "no") %>%  
  select(subject, ldl, ht, age, smoking, drinkany, sbp,  
         physact, bmi, diabetes)
```

# The Codebook (n = 2032)

Variable	Description
subject	subject code
HT	factor: hormone therapy or placebo
diabetes	yes or no (all are no in our sample)
ldl	LDL cholesterol in mg/dl
age	age in years
smoking	yes or no
drinkany	yes or no
sbp	systolic BP in mm Hg
physact	5-level factor, details next slide
bmi	body-mass index in kg/m <sup>2</sup>

**Goal** Predict ldl using age, smoking, drinkany, sbp, physact and bmi, across both HT levels but restricted to women without diabetes.

# The physact variable

```
hers1 %>% count(physact)
```

```
# A tibble: 5 x 2
```

	physact	n
* <chr>		<int>
1	about as active	674
2	much less active	107
3	much more active	252
4	somewhat less active	322
5	somewhat more active	677

Comparison is to activity levels for these women just before menopause.

# Any missing data?

```
miss_var_summary(hers1)
```

```
# A tibble: 10 x 3
  variable n_miss pct_miss
  <chr>      <int>    <dbl>
1 ldl          7    0.344
2 drinkany     2    0.0984
3 bmi          2    0.0984
4 subject      0     0
5 ht           0     0
6 age          0     0
7 smoking      0     0
8 sbp          0     0
9 physact      0     0
10 diabetes    0     0
```

# Single Imputation for drinkany, bmi and ldl

Since drinkany is a factor, we have to do some extra work to impute.

```
set.seed(432092)
```

```
hers2 <- hers1 %>%  
  mutate(drinkany_n =  
    ifelse(drinkany == "yes", 1, 0)) %>%  
  impute_pmm(drinkany_n ~ age + smoking) %>%  
  mutate(drinkany =  
    ifelse(drinkany_n == 1, "yes", "no")) %>%  
  impute_rlm(bmi ~ age + smoking + sbp) %>%  
  impute_rlm(ldl ~ age + smoking + sbp + bmi)
```

# Now, check missingness...

```
miss_var_summary(hers2)
```

```
# A tibble: 11 x 3
```

	variable <chr>	n_miss <int>	pct_miss <dbl>
1	subject	0	0
2	ldl	0	0
3	ht	0	0
4	age	0	0
5	smoking	0	0
6	drinkany	0	0
7	sbp	0	0
8	physact	0	0
9	bmi	0	0
10	diabetes	0	0
11	drinkany_n	0	0



# Multiple Imputation using `aregImpute` from `Hmisc`

Model to predict all missing values of any variables, using additive regression bootstrapping and predictive mean matching.

Steps are:

- 1 `aregImpute` draws a sample with replacement from the observations where the target variable is observed, not missing.
- 2 It then fits a flexible additive model to predict this target variable while finding the optimum transformation of it.
- 3 It then uses this fitted flexible model to predict the target variable in all of the original observations.
- 4 Finally, it imputes each missing value of the target variable with the observed value whose predicted transformed value is closest to the predicted transformed value of the missing value.

# Fitting a Multiple Imputation Model

```
set.seed(4320132)
dd <- datadist(hers1)
options(datadist = "dd")
fit3 <- aregImpute(~ ldl + age + smoking + drinkany +
                  sbp + physact + bmi,
                  nk = c(0, 3:5), tlinear = FALSE,
                  data = hers1, B = 10, n.impute = 20)
```

Iteration 1 Iteration 2 Iteration 3 Iteration 4 Iteration 5 Iteration 6

# Multiple Imputation using `aregImpute` from `Hmisc`

`aregImpute` requires specifications of all variables, and several other details:

- `n.impute` = number of imputations, we'll run 20
- `nk` = number of knots to describe level of complexity, with our choice `nk = c(0, 3:5)` we'll fit both linear models and models with restricted cubic splines with 3, 4, and 5 knots
- `tlinear` = `FALSE` allows the target variable to have a non-linear transformation when `nk` is 3 or more
- `B` = 10 specifies 10 bootstrap samples will be used
- `data` specifies the source of the variables

# aregImpute Imputation Results (1 of 4)

```
fit3
```

Multiple Imputation using Bootstrap and PMM

```
aregImpute(formula = ~ldl + age + smoking + drinkany + sbp +  
  physact + bmi, data = hers1, n.impute = 20, nk = c(0, 3:5),  
  tlinear = FALSE, B = 10)
```

n: 2032      p: 7      Imputations: 20      nk: 0

Number of NAs:

ldl	age	smoking	drinkany	sbp	physact	bmi
7	0	0	2	0	0	2

## fit3 Imputation Results (2 of 4)

R-squares for Predicting Non-Missing Values for Each  
Variable Using Last Imputations of Predictors

ldl	drinkany	bmi
0.041	0.014	0.109

## fit3 Imputation Results (3 of 4)

Resampling results for determining the complexity of imputation models

Variable being imputed: ldl

Bootstrap bias-corrected summaries:

Statistic	nk = 0	nk = 3	nk = 4	nk = 5
R <sup>2</sup>	0.0139	0.0149	0.00776	0.0124
mean absolute error	28.3594	42.9139	44.09937	39.8266
median abs. error	22.8301	35.5441	38.85302	32.6386

10-fold cross-validated:

Statistic	nk = 0	nk = 3	nk = 4	nk = 5
R <sup>2</sup>	0.0214	0.0180	0.01517	0.0191
mean absolute error	145.7176	43.5007	45.02428	44.2456
median abs. error	141.4238	36.4102	38.88053	37.3141

## fit3 Imputation Results (4 of 4)




Variable being imputed: drinkany

	nk=0	nk=3	nk=4	nk=5
Bootstrap $R^2$	0.0163	0.0113	0.0102	0.00986
10-fold cv $R^2$	0.0205	0.0249	0.0163	0.01358
Bootstrap mean  error	0.4470	0.4568	0.4558	0.46624
10-fold cv mean  error	0.4450	0.4454	0.4476	0.44676
Bootstrap median  error	0.0000	0.0000	0.0000	0.00000
10-fold cv median  error	0.0000	0.0500	0.1000	0.00000

Variable being imputed: bmi

	nk=0	nk=3	nk=4	nk=5
Bootstrap $R^2$	0.0845	0.0932	0.0946	0.0847
10-fold cv $R^2$	0.0864	0.0903	0.0968	0.0899
Bootstrap mean  error	3.7829	4.8119	4.9226	5.1775
10-fold cv mean  error	27.6776	4.8359	4.9390	5.1136
Bootstrap median  error	2.9955	3.9704	3.9371	4.2634
10-fold cv median  error	27.0143	3.9894	3.9431	4.1876

# A plot of the imputed values... (results)





## A plot of the imputed values... (code)

```
par(mfrow = c(1,3))  
plot(fit3)  
par(mfrow = c(1,1))
```

- For `ldl`, we imputed most of the 7 missing subjects in most of the 20 imputation runs to values within a range of around 120 through 200, but occasionally, we imputed values that were substantially lower than 100.
- For `drinkany` we imputed about 70% no and 30% yes.
- For `bmi`, we imputed values ranging from about 23 to 27 in many cases, and up near 40 in other cases.
- This method never imputes a value for a variable that doesn't already exist in the data.

# Kitchen Sink Model (Main Effects only)

```
mod_ks <- ols(ldl ~ age + smoking + drinkany + sbp +  
              physact + bmi, data = hers2)  
anova(mod_ks)
```

## Analysis of Variance

Response: ldl

Factor	d.f.	Partial SS	MS	F	P
age	1	9330.911	9330.911	6.93	0.0085
smoking	1	8199.755	8199.755	6.09	0.0137
drinkany	1	6444.424	6444.424	4.79	0.0288
sbp	1	9274.287	9274.287	6.89	0.0087
physact	4	10874.528	2718.632	2.02	0.0891
bmi	1	15876.957	15876.957	11.80	0.0006
REGRESSION	9	60077.708	6675.301	4.96	<.0001
ERROR	2022	2721037.890	1345.716		

# Spearman $\rho^2$ Plot


How should we prioritize the degrees of freedom we spend on non-linearity?

```
plot(spearman2(ldl ~ age + smoking + drinkany + sbp +  
             physact + bmi, data = hers2))
```

Plot's on the next page.

- Note the use of the simple imputation `hers2` data here. Why?

# Spearman $\rho^2$ Plot Result



# Spending Degrees of Freedom

We're spending 9 degrees of freedom in our kitchen sink model. (We can verify this with `anova` or the plot.)

- Each quantitative main effect costs 1 df to estimate
- Each binary categorical variable also costs 1 df
- Multi-categorical variables with  $L$  levels cost  $L-1$  df to estimate

Suppose we're willing to spend up to a total of **14** degrees of freedom (i.e. a combined 5 more on interaction terms and other ways to capture non-linearity.)

What should we choose?

# What did we see in the Spearman $\rho^2$ Plot?

Group 1 (largest adjusted  $\rho^2$ )

- bmi, a quantitative predictor, is furthest to the right

Group 2 (next largest)

- smoking, a binary predictor, is next, followed closely by
- age, a quantitative predictor

Other predictors (rest of the group)

- sbp, quantitative
- drinkany, binary
- physact, multi-categorical (5 levels)

# Impact of Adding Non-Linear Terms on Spent DF

What happens when we add a non-linear term?

- Adding a polynomial of degree  $D$  costs  $D$  degrees of freedom.
  - So a polynomial of degree 2 (quadratic) costs 2 df, or 1 more than the main effect alone.
- Adding a restricted cubic spline with  $K$  knots costs  $K-1$  df.
  - So adding a rcs with 4 knots uses 3 df, or 2 more than the main effect.
  - We restrict ourselves to considering splines with 3, 4, or 5 knots.
- Adding an interaction (product term) depends on the main effects of the predictors we are interacting
  - If the product term's predictors have  $df1$  and  $df2$  degrees of freedom, product term adds  $df1 \times df2$  degrees of freedom.
  - An interaction of a binary and quantitative variable adds  $1 \times 1 = 1$  additional degree of freedom to the main effects model.
  - When we use a quantitative variable in a spline and interaction, we'll do the interaction on the main effect, not the spline.

# Model we'll fit with `ols`

Fitting a model to predict `ldl` using

- `bmi` with a restricted cubic spline, 5 knots
- `age` with a quadratic polynomial
- `sbp` as a linear term
- `drinkany` indicator
- `physact` factor
- `smoking` indicator and its interaction with the main effect of `bmi`

We can fit this to the data

- restricted to complete cases (`hers1`, effectively)
- after simple imputation (`hers2`)
- after our multiple imputation (`fit3`)



## Using only the Complete Cases

# Fitting the model to the complete cases

```
d <- datadist(hers1)
options(datadist = "d")

m1 <- ols(ldl ~ rcs(bmi, 5) + pol(age, 2) + sbp +
          drinkany + physact + smoking +
          smoking %ia% bmi, data = hers1,
          x = TRUE, y = TRUE)
```

where %ia% identifies the linear interaction alone.

# m1 results (screen 1/2)

m1

```
Frequencies of Missing Values Due to Each Variable
  1d1      bmi      age      sbp drinkany  physact  smoking
    7        2        0        0         2         0         0

Linear Regression Model

ols(formula = 1d1 ~ rcs(bmi, 5) + pol(age, 2) + sbp + drinkany +
      physact + smoking + smoking %ia% bmi, data = hers1, x = TRUE,
      y = TRUE)

              Model Likelihood      Discrimination
              Ratio Test              Indexes
Obs      2021    LR chi2      52.61    R2      0.026
sigma36.7430    d.f.      14    R2 adj  0.019
d.f.      2006    Pr(> chi2) 0.0000    g      6.629

Residuals

      Min      1Q    Median      3Q      Max
-113.440 -24.519  -3.778   20.940  197.087
```

## m1 results (screen 2/2)

m1

	Coef	S.E.	t	Pr(> t )
Intercept	121.6057	68.2000	1.78	0.0747
bmi	1.5687	1.0107	1.55	0.1208
bmi'	-8.6685	9.1577	-0.95	0.3440
bmi''	40.5712	37.4468	1.08	0.2787
bmi'''	-55.8872	44.5946	-1.25	0.2103
age	-0.5791	1.9657	-0.29	0.7683
age^2	0.0018	0.0149	0.12	0.9024
sbp	0.1221	0.0453	2.69	0.0072
drinkany=yes	-3.7427	1.6629	-2.25	0.0245
physact=much less active	-4.5660	3.8904	-1.17	0.2407
physact=much more active	-0.3291	2.7521	-0.12	0.9048
physact=somewhat less active	-0.0160	2.5270	-0.01	0.9950
physact=somewhat more active	3.7731	2.0293	1.86	0.0631
smoking=yes	-7.0832	12.0586	-0.59	0.5570
smoking=yes * bmi	0.4961	0.4391	1.13	0.2587

# Fit Model after Single Imputation

# Fitting the model after simple imputation

```
dd <- datadist(hers2)
options(datadist = "dd")

m2 <- ols(ldl ~ rcs(bmi, 5) + pol(age, 2) + sbp +
          drinkany + physact + smoking +
          smoking %ia% bmi, data = hers2,
          x = TRUE, y = TRUE)
```

where, again, %ia% identifies the linear interaction alone.

## m2 results (screen 1/2)

m2

## Linear Regression Model

```
ols(formula = ldl ~ rcs(bmi, 5) + pol(age, 2) + sbp + drinkany +  
      physact + smoking + smoking %ia% bmi, data = hers2, x = TRUE,  
      y = TRUE)
```

		Model Likelihood Ratio Test	Discrimination Indexes		
Obs	2032	LR chi2	53.14	R2	0.026
sigma36.6503		d.f.	14	R2 adj	0.019
d.f.	2017	Pr(> chi2)	0.0000	g	6.631

## Residuals

Min	1Q	Median	3Q	Max
-113.379	-24.326	-3.835	20.832	197.097

## m2 results (screen 2/2)

m2

	Coef	S.E.	t	Pr(> t )
Intercept	120.2662	67.6113	1.78	0.0754
bmi	1.5508	1.0071	1.54	0.1237
bmi'	-8.4486	9.0978	-0.93	0.3532
bmi''	39.6413	37.1378	1.07	0.2859
bmi'''	-54.8924	44.2677	-1.24	0.2151
age	-0.5249	1.9490	-0.27	0.7877
age^2	0.0014	0.0148	0.10	0.9233
sbp	0.1209	0.0451	2.68	0.0074
drinkany=yes	-3.7023	1.6544	-2.24	0.0253
physact=much less active	-4.7408	3.8621	-1.23	0.2198
physact=much more active	-0.2635	2.7391	-0.10	0.9234
physact=somewhat less active	0.0130	2.5101	0.01	0.9959
physact=somewhat more active	3.8031	2.0193	1.88	0.0598
smoking=yes	-6.8961	12.0196	-0.57	0.5662
smoking=yes * bmi	0.4892	0.4375	1.12	0.2636



# ANOVA results for m2 from ols

```
anova(m2)
```

Analysis of Variance	Response: ldl				
Factor	d.f.	Partial SS	MS	F	P
bmi (Factor+Higher Order Factors)	5	2.758824e+04	5517.64861	4.11	0.0010
All Interactions	1	1.679813e+03	1679.81344	1.25	0.2636
Nonlinear	3	9.735452e+03	3245.15068	2.42	0.0647
age	2	9.175762e+03	4587.88077	3.42	0.0330
Nonlinear	1	1.244351e+01	12.44351	0.01	0.9233
sbp	1	9.657476e+03	9657.47569	7.19	0.0074
drinkany	1	6.726918e+03	6726.91809	5.01	0.0253
physact	4	9.709992e+03	2427.49791	1.81	0.1247
smoking (Factor+Higher Order Factors)	2	1.085405e+04	5427.02463	4.04	0.0177
All Interactions	1	1.679813e+03	1679.81344	1.25	0.2636
smoking * bmi (Factor+Higher Order Factors)	1	1.679813e+03	1679.81344	1.25	0.2636
TOTAL NONLINEAR	4	9.738807e+03	2434.70175	1.81	0.1237
TOTAL NONLINEAR + INTERACTION	5	1.171134e+04	2342.26845	1.74	0.1214
REGRESSION	14	7.178905e+04	5127.78931	3.82	<.0001
ERROR	2017	2.709327e+06	1343.24569		

# Validation of summary statistics

```
set.seed(432); validate(m2)
```

	index.orig	training	test	optimism
R-square	0.0258	0.0307	0.0182	0.0125
MSE	1333.3300	1323.5182	1343.7711	-20.2529
g	6.6306	7.1676	5.8338	1.3338
Intercept	0.0000	0.0000	26.5316	-26.5316
Slope	1.0000	1.0000	0.8174	0.1826

	index.corrected	n
R-square	0.0133	40
MSE	1353.5829	40
g	5.2968	40
Intercept	26.5316	40
Slope	0.8174	40

	index.orig	training	test	optimism	index.corrected	n
R-square	0.0258	0.0307	0.0182	0.0125	0.0133	40
MSE	1333.3300	1323.5182	1343.7711	-20.2529	1353.5829	40
g	6.6306	7.1676	5.8338	1.3338	5.2968	40

# summary(m2) results

```
summary(m2)
```

```
Effects                Response : ldl


Factor               Low  High  Diff.  Effect  S.E.   Lower 0.95  Upper 0.95
bmi                  24.2  30.263  6.0625  5.1862  2.2217    0.82921  9.54330
age                   62.0  72.000 10.0000 -3.3412  1.3450   -5.97890 -0.70357
sbp                  120.0 145.000 25.0000  3.0218  1.1270    0.81165  5.23190
drinkany - yes:no      1.0   2.000    NA -3.7023  1.6544   -6.94690 -0.45779
physact - about as active:somewhat more active  5.0   1.000    NA -3.8031  2.0193   -7.76310  0.15695
physact - much less active:somewhat more active  5.0   2.000    NA -8.5439  3.9035  -16.19900 -0.88862
physact - much more active:somewhat more active  5.0   3.000    NA -4.0666  2.7125   -9.38630  1.25310
physact - somewhat less active:somewhat more active  5.0   4.000    NA -3.7901  2.5633   -8.81720  1.23690
smoking - yes:no       1.0   2.000    NA  6.2635  2.4009    1.55500 10.97200

Adjusted to: bmi=26.9 smoking=no
```

- Of course, these should really be plotted...


# Effect Size Plot for m2

```
plot(summary(m2))
```



Adjusted to: bmi=26.9 smoking=no

# Nomogram for $m_2$



# Making Predictions for an Individual

Suppose now that we want to use R to get a prediction for a new individual subject with `bmi = 30`, `age = 50`, `smoking = yes` and `physact = about as active`, `drinkany = yes` and `sbp` of 150.

```
predict(m2, expand.grid(bmi = 30, age = 50, smoking = "yes",  
                        physact = "about as active",  
                        drinkany = "yes", sbp = 150),  
        conf.int = 0.95, conf.type = "individual")
```

\$linear.predictors	\$lower	\$upper
160.9399	88.48615	233.3936

# Making Predictions for a Long-Run Mean

The other kind of prediction we might wish to make is for the mean of a series of subjects whose  $\text{bmi} = 30$ ,  $\text{age} = 50$ ,  $\text{smoking} = \text{yes}$  and  $\text{physact} = \text{about as active}$ ,  $\text{drinkany} = \text{yes}$  and  $\text{sbp}$  of 150.


```
predict(m2, expand.grid(bmi = 30, age = 50, smoking = "yes",  
                        physact = "about as active",  
                        drinkany = "yes", sbp = 150),  
        conf.int = 0.95, conf.type = "mean")
```

\$linear.predictors	\$lower	\$upper
160.9399	151.8119	170.0679

Of course, the confidence interval will always be narrower than the prediction interval given the same predictor values.

# Residuals vs. Fitted Values?

```
plot(resid(m2) ~ fitted(m2))
```





# Influential Points?

```
which.influence(m2, cutoff = 0.4)
```

```
$Intercept
```

```
[1] 1135
```

```
$age
```

```
[1] 1135
```

```
$smoking
```

```
[1] 132
```

```
$`smoking * bmi`
```

```
[1] 132
```

# Using Multiple Imputation

# Fitting the Model using Multiple Imputation

What do we have now?

- An imputation model `fit3`

```
fit3 <- aregImpute(~ ldl + age + smoking + drinkany + sbp +  
  physact + bmi, nk = c(0, 3:5), tlinear = FALSE,  
  data = hers1, B = 10, n.impute = 20, x = TRUE)
```

- A prediction model (from `m1` or `m2`)

```
ols(ldl ~ rcs(bmi, 5) + pol(age, 2) + sbp +  
  drinkany + physact + smoking + smoking %ia% bmi,  
  x = TRUE, y = TRUE)
```

Now we put them together with the `fit.mult.impute` function...

# Linear Regression & Imputation Model

```
m3imp <-  
  fit.mult.impute(ldl ~ rcs(bmi, 5) + pol(age, 2) + sbp +  
    drinkany + physact + smoking +  
    smoking %ia% bmi,  
    fitter = ols, xtrans = fit3,  
    data = hers1, pr = FALSE)
```

- When you run this without the `pr = FALSE` it generates considerable output related to the imputations, which we won't use today.
- Let's look at the rest of the output this yields...

# m3imp results (screen 1/2)

## m3imp

### Linear Regression Model

```
fit.mult.impute(formula = ldl ~ rcs(bmi, 5) + pol(age, 2) + sbp +  
  drinkany + physact + smoking + smoking %ia% bmi, fitter = ols,  
  xtrans = fit3, data = hers1, pr = FALSE)
```

		Model Likelihood		Discrimination	
		Ratio Test		Indexes	
Obs	2032	LR chi2	52.74	R2	0.026
sigma	36.7331	d.f.	14	R2 adj	0.019
d.f.	2017	Pr(> chi2)	0.0000	g	6.621

### Residuals

	Min	1Q	Median	3Q	Max
	-113.345	-24.510	-3.803	20.777	197.295

## m3imp results (screen 2/2)

m3imp

	Coef	S.E.	t	Pr(> t )
Intercept	119.8951	67.8409	1.77	0.0773
bmi	1.5436	1.0097	1.53	0.1265
bmi'	-8.3664	9.1409	-0.92	0.3602
bmi''	39.2149	37.3458	1.05	0.2938
bmi'''	-54.2873	44.5323	-1.22	0.2230
age	-0.5002	1.9555	-0.26	0.7981
age^2	0.0012	0.0148	0.08	0.9351
sbp	0.1198	0.0454	2.64	0.0083
drinkany=yes	-3.7196	1.6613	-2.24	0.0253
physact=much less active	-4.7109	3.8716	-1.22	0.2238
physact=much more active	-0.2328	2.7512	-0.08	0.9326
physact=somewhat less active	-0.0417	2.5246	-0.02	0.9868
physact=somewhat more active	3.8197	2.0286	1.88	0.0599
smoking=yes	-6.8967	12.0503	-0.57	0.5672
smoking=yes * bmi	0.4866	0.4389	1.11	0.2677

# ANOVA results for m3imp

```
anova(m3imp)
```

Analysis of Variance		Response: ldl			
Factor	d.f.	Partial SS	MS	F	P
bmi (Factor+Higher Order Factors)	5	2.728300e+04	5456.600791	4.04	0.0012
All Interactions	1	1.658459e+03	1658.458931	1.23	0.2677
Nonlinear	3	9.585703e+03	3195.234412	2.37	0.0690
age	2	9.320445e+03	4660.222299	3.45	0.0318
Nonlinear	1	8.950493e+00	8.950493	0.01	0.9351
sbp	1	9.407603e+03	9407.602954	6.97	0.0083
drinkany	1	6.763854e+03	6763.853503	5.01	0.0253
physact	4	9.698175e+03	2424.543639	1.80	0.1268
smoking (Factor+Higher Order Factors)	2	1.031090e+04	5155.452328	3.82	0.0221
All Interactions	1	1.658459e+03	1658.458931	1.23	0.2677
smoking * bmi (Factor+Higher Order Factors)	1	1.658459e+03	1658.458931	1.23	0.2677
TOTAL NONLINEAR	4	9.587178e+03	2396.794504	1.78	0.1309
TOTAL NONLINEAR + INTERACTION	5	1.152744e+04	2305.487432	1.71	0.1293
REGRESSION	14	7.030149e+04	5021.535034	3.72	<.0001
ERROR	2017	2.721574e+06	1349.317884		


# Summary of Effect Estimates for m3imp

```
summary(m3imp)
```

Effects		Response : ldl						
Factor		Low	High	Diff.	Effect	S.E.	Lower 0.95	Upper 0.95
bmi		24.2	30.263	6.0625	5.1643	2.2300	0.79099	9.53750
age		62.0	72.000	10.0000	-3.3824	1.3518	-6.03340	-0.73144
sbp		120.0	145.000	25.0000	2.9955	1.1345	0.77068	5.22040
drinkany - yes:no		1.0	2.000	NA	-3.7196	1.6613	-6.97780	-0.46150
physact - about as active:somewhat more active		5.0	1.000	NA	-3.8197	2.0286	-7.79800	0.15861
physact - much less active:somewhat more active		5.0	2.000	NA	-8.5306	3.9152	-16.20900	-0.85228
physact - much more active:somewhat more active		5.0	3.000	NA	-4.0525	2.7260	-9.39850	1.29350
physact - somewhat less active:somewhat more active		5.0	4.000	NA	-3.8614	2.5796	-8.92030	1.19760
smoking - yes:no		1.0	2.000	NA	6.1923	2.4427	1.40190	10.98300
Adjusted to: bmi=26.9 smoking=no								



```
plot(summary(m3imp))
```





Adjusted to:bmi=26.9 smoking=no


# Evaluation via Partial $R^2$ and AIC (code)

```
par(mfrow = c(1,2))  
plot(anova(m3imp), what="partial R2")  
plot(anova(m3imp), what="aic")  
par(mfrow = c(1,1))
```


# Evaluation via Partial R<sup>2</sup> and AIC (result)



# plot(nomogram(m3imp))



```
plot(resid(m3imp) ~ fitted(m3imp))
```



# Next Step

Can we do all of this for a logistic regression model?