

# Atlas

David Ledesma

3/22/2021

## Load in data

```
library(curl)
library(readxl)
Amb_linear_data <- curl("https://raw.githubusercontent.com/TIMAVID/Ambystoma/master/Linear_data/Data/Amb_linear_data.csv")
Amb_linear_data <- read.csv(Amb_linear_data)
```

## Tidy data

```
##
## Attaching package: 'dplyr'

## The following objects are masked from 'package:stats':
##
##   filter, lag

## The following objects are masked from 'package:base':
##
##   intersect, setdiff, setequal, union
```


## PCA's

```
Atlas.pca <- prcomp(Atlas_wofossil_noNA[c(1:7)], center = TRUE, scale = TRUE) # PCA
# Summary stats #
summary(Atlas.pca)
```


```
## Importance of components:
##              PC1      PC2      PC3      PC4      PC5      PC6      PC7
## Standard deviation    2.3031 1.0447 0.55567 0.36920 0.25577 0.23039 0.20233
## Proportion of Variance 0.7577 0.1559 0.04411 0.01947 0.00935 0.00758 0.00585
## Cumulative Proportion 0.7577 0.9136 0.95775 0.97722 0.98657 0.99415 1.00000
```

```
sd <- Atlas.pca$sdev
loadings <- Atlas.pca$rotation
rownames(loadings) <- colnames(Atlas[c(1:7)])
scores <- Atlas.pca$x
# Show variance explained by PC's #
var <- sd^2
varPercent <- var/sum(var) * 100
barplot(varPercent, xlab = "PC", ylab = "Percent Variance", names.arg = 1:length(varPercent),
```

```
las = 1, ylim = c(0, max(varPercent)), col = "gray")
abline(h = 1/ncol(Atlas[c(1:7)]) * 100, col = "red")
```



```
fviz_eig(Atlas.pca, xlab = "Principal Components")
```



```
# Show loadings #
loadings
```


```
##          PC1          PC2          PC3          PC4
## tub_interglen_extension 0.03870518 -0.94714699  0.1063967 -0.23874181
## X1                    0.41652039  0.14147859  0.2646479 -0.04799568
## X2                    0.42197173 -0.04128891  0.1441653  0.29389133
## X3                    0.40400635 -0.21003360 -0.0244320  0.71347417
## X4                    0.41821678  0.02044259  0.1837650 -0.49111174
## X5                    0.41053379  0.19060650  0.2312434 -0.26560742
## X6                    0.37455058 -0.01813215 -0.9000098 -0.18314807
##          PC5          PC6          PC7
## tub_interglen_extension 0.01387657 -0.16293082 -0.07973435
## X1                    0.48873145 -0.09764476 -0.69696451
## X2                    0.47031170 -0.27244438  0.64626642
## X3                   -0.39377589  0.30964846 -0.17911119
## X4                   -0.03666985  0.70273932  0.23351511
## X5                   -0.61565777 -0.53937106  0.03398159
## X6                    0.06570612 -0.09511705 -0.04957562
```

```
sqrt(1/ncol(Atlas[c(1:7)])) # cutoff for 'important' loadings
```


```
## [1] 0.3779645
```

```
# dev.new(height=7, width=7)
```

```
biplot(scores[, 1:2], loadings[, 1:2], cex = 0.9)
```




```
ggbiplot(Atlas.pca, ellipse = FALSE, groups = Atlas_wofossil_noNA$species)
```



## Plot


```
scores <- as.data.frame(scores)
scores$species <- Atlas_wofossil_noNA$species # reattach species
theme <- theme(panel.background = element_blank(), panel.border = element_rect(fill = NA),
  panel.grid.major = element_blank(), panel.grid.minor = element_blank(),
  strip.background = element_blank(), axis.text.x = element_text(colour = "black"),
  axis.text.y = element_text(colour = "black"), axis.ticks = element_line(colour = "black"),
  plot.margin = unit(c(1, 1, 1, 1), "line"))
percentage <- paste(colnames(scores), "(", paste(as.character(round(varPercent)),
  "%", " )", sep = ""))
n <- 14
palette <- distinctColorPalette(n)
p <- ggplot(scores, aes(x = PC1, y = PC2, color = species))
p <- p + geom_point(size = 5) + theme + xlab(percentage[1]) + ylab(percentage[2]) +
  scale_color_manual(values = palette)
# p + stat_ellipse()
p
```



```
# alternative plot fviz_pca_ind(Atlas.pca) fviz_pca_ind(Atlas.pca,
# label='none', habillage=Atlas_wofossil_noNA$species, addEllipses=TRUE,
# ellipse.level=0.95, palette = palette) alternative plot 2
autoplot(Atlas.pca, data = Atlas_wofossil_noNA, colour = "species", frame = TRUE,
label = TRUE)
```


```
## Warning: `select_()` was deprecated in dplyr 0.7.0.
## Please use `select()` instead.

## Warning: `group_by_()` was deprecated in dplyr 0.7.0.
## Please use `group_by()` instead.
## See vignette('programming') for more help
```



### No tuberculum interglenoideum ventral extent measurement

```
Atlas_wofossil_noTub <- Atlas_wofossil[, -1]
Atlas_wofossil_noTub <- na.omit(Atlas_wofossil_noTub) # remove rows with N/A's
Atlas_wofossil_noTub$species <- as.factor(Atlas_wofossil_noTub$species)
Atlas_wofossil_noTub$species <- factor(Atlas_wofossil_noTub$species, levels = c("A.gracile",
  "A.talpoideum", "A.maculatum", "A.macrodactylum", "A.opacum", "A.jeffersonianum",
  "A.laterale", "A.mabeei", "A.texanum", "A.annulatum", "A.tigrinum", "A.mavortium",
  "A.ordinarium", "A.subsalsum")) # Reorder species
Atlas.pca_2 <- prcomp(Atlas_wofossil_noTub[c(1:6)], center = TRUE, scale = FALSE) # PCA
autoplot(Atlas.pca_2, data = Atlas_wofossil_noTub, colour = "species", frame = TRUE,
  label = TRUE)
```



## Statistical Tests

```
# Removing 'A.laterale|A.talpoideum|A.subsalsum|A.ordinarium' due to low
# sample sizes#
Atlas_wofossil_noTub_sub <- dplyr::filter(Atlas_wofossil_noTub, !grepl("A.laterale|A.talpoideum|A.subsalsum|A.ordinarium",
species))
Atlas_wofossil_noTub_sub$species <- factor(Atlas_wofossil_noTub_sub$species,
levels = c("A.gracile", "A.maculatum", "A.macrodactylum", "A.opacum", "A.jeffersonianum",
"A.mabeei", "A.texanum", "A.annulatum", "A.tigrinum", "A.mavortium")) # Reorder species
```

## Various checks for MANOVA

```
# Atlas_wofossil_noTub_sub <- Atlas_wofossil_noTub_sub %>% add_column(id =
# rownames(Atlas_wofossil_noTub_sub), .after = 8)
# Check sample sizes:PASS
Atlas_wofossil_noTub_sub %>% dplyr::group_by(species) %>% dplyr::summarise(N = n())

## # A tibble: 10 x 2
##   species      N
##   <fct>      <int>
## 1 A.gracile      8
## 2 A.maculatum    14
## 3 A.macrodactylum 12
```

```
## 4 A.opacum 13
## 5 A.jeffersonianum 11
## 6 A.mabeei 9
## 7 A.texanum 9
## 8 A.annulatum 6
## 9 A.tigrinum 17
## 10 A.mavortium 12
```

```
# Identify univariate outliers for each variable:FAIL
```

```
Atlas_wofossil_noTub_sub %>% dplyr::group_by(species) %>% identify_outliers(5) #input variable column
```

```
## # A tibble: 15 x 10
```

```
##   species      M1      M2      M3      M4      M5      M6 specimen_num is.outlier
##   <fct>      <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <chr>          <lgl>
## 1 A.gracile  2.27  2.34  0.86  1.67  2.68  0.98 A.gracile_UTEP_16~ TRUE
## 2 A.maculatum 2.58  2.97  1.24  1.95  2.47  0.97 A.maculatum_UTEP_~ TRUE
## 3 A.macrodac~ 2.62  2.68  0.98  1.86  2.28  0.96 A.macrodactylum_U~ TRUE
## 4 A.opacum    1.93  2.18  0.85  1.47  2.13  0.83 A.opacum_UTEP_1276 TRUE
## 5 A.opacum    2.05  2.34  0.78  1.25  2.09  0.78 A.opacum_UTEP_1280 TRUE
## 6 A.opacum    2.54  2.53  0.87  1.74  2.49  0.84 A.opacum_UTEP_1285 TRUE
## 7 A.opacum    3.46  3.39  1.27  2.6   3.35  1.17 A.opacum_UTEP_1310 TRUE
## 8 A.jefferso~ 2.89  2.79  1.14  2.11  2.63  0.99 A.jeffersonianum_~ TRUE
## 9 A.mabeei    1.75  1.44  0.6   1.19  1.72  0.78 A.mabeei_UTEP_1576 TRUE
## 10 A.mabeei   2.51  2.71  1.01  1.94  2.6   1.15 A.mabeei_UTEP_1584 TRUE
## 11 A.texanum   2.8   3.19  1.28  2.31  2.62  1.26 A.texanum_UTEP_17~ TRUE
## 12 A.tigrinum  4.52  4.54  1.87  3.63  5.38  1.73 A.tigrinum_UTEP_1~ TRUE
## 13 A.tigrinum  3.69  3.67  1.47  3.35  4.49  1.51 A.tigrinum_UTEP_1~ TRUE
## 14 A.tigrinum  4.46  4.35  1.66  3.36  4.91  1.79 A.tigrinum_UTEP_1~ TRUE
## 15 A.mavortium 5.65  5.13  1.74  4.97  6.03  2.18 A.mavortium_UTEP_~ TRUE
## # ... with 1 more variable: is.extreme <lgl>
```

```
Atlas_wofossil_noTub_sub %>% dplyr::group_by(species) %>% identify_outliers(6) #input variable column
```

```
## # A tibble: 7 x 10
```

```
##   species      M1      M2      M3      M4      M5      M6 specimen_num is.outlier is.extreme
##   <fct>      <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <chr>          <lgl>      <lgl>
## 1 A.macu~    3.68  4.18  1.58  3.13  3.71  3.29 A.maculatum_~ TRUE      TRUE
## 2 A.mabe~    2.51  2.71  1.01  1.94  2.6   1.15 A.mabeei_UT~ TRUE      FALSE
## 3 A.annu~    3.38  3.38  1.17  2.32  2.73  1.08 A.annulatum_~ TRUE      FALSE
## 4 A.tigr~    4.52  4.54  1.87  3.63  5.38  1.73 A.tigrinum_~ TRUE      FALSE
## 5 A.tigr~    3.45  2.99  1.51  2.18  3.56  1.11 A.tigrinum_~ TRUE      FALSE
## 6 A.tigr~    4.46  4.35  1.66  3.36  4.91  1.79 A.tigrinum_~ TRUE      FALSE
## 7 A.mavo~    4.92  5.12  2.22  3.87  5.32  3.65 A.mavortium_~ TRUE      TRUE
```

```
# ...
```

```
# Detect multivariate outliers:MOSTLY PASS
```

```
head(mahalanobis_distance(Atlas_wofossil_noTub_sub[, 1:6]))
```

```
##   M1      M2      M3      M4      M5      M6 mahal.dist is.outlier
## 1 3.38 3.38 1.17 2.32 2.73 1.08      6.035      FALSE
## 2 3.36 3.21 1.31 2.49 2.91 1.22      6.431      FALSE
## 3 3.50 3.42 1.29 2.46 2.96 1.30      4.895      FALSE
## 4 3.42 3.61 1.30 2.57 2.87 1.28      4.941      FALSE
## 5 3.12 3.44 1.24 2.29 2.77 1.24      2.674      FALSE
## 6 3.05 3.18 1.24 2.27 2.61 1.38      4.128      FALSE
```



```
# Check univariate normality assumption:FAIL
Atlas_wofossil_noTub_sub %>% group_by(species) %>% shapiro_test(M1, M2, M3,
  M4, M5, M6) %>% arrange(variable)
```

```
## # A tibble: 60 x 4
##   species      variable statistic      p
##   <fct>      <chr>      <dbl>    <dbl>
## 1 A.gracile    M1          0.782 0.0183
## 2 A.maculatum  M1          0.787 0.00350
## 3 A.macrodactylum M1          0.968 0.894
## 4 A.opacum     M1          0.957 0.703
## 5 A.jeffersonianum M1          0.945 0.576
## 6 A.mabeei     M1          0.909 0.308
## 7 A.texanum    M1          0.972 0.910
## 8 A.annulatum  M1          0.885 0.293
## 9 A.tigrinum   M1          0.787 0.00136
## 10 A.mavortium M1          0.961 0.791
## # ... with 50 more rows
```

```
# Check Multivariate normality:FAIL
Atlas_wofossil_noTub_sub %>% dplyr::select(, 1:6) %>% mshapiro_test()
```

```
## # A tibble: 1 x 2
##   statistic p.value
##   <dbl>    <dbl>
## 1      0.482 4.95e-18
```

```
# Identify multicollinearity:FAIL
Atlas_wofossil_noTub_sub %>% cor_test(, 1:6)
```

```
## # A tibble: 36 x 8
##   var1 var2 cor statistic      p conf.low conf.high method
##   <chr> <chr> <dbl>    <dbl>    <dbl>    <dbl>    <dbl>    <chr>
## 1 M1    M1    1      Inf    0.      1      1      Pearson
## 2 M1    M2    0.94   30.2 1.02e-54 0.921    0.962 Pearson
## 3 M1    M3    0.85   17.1 1.29e-32 0.793    0.897 Pearson
## 4 M1    M4    0.93   25.9 2.59e-48 0.896    0.950 Pearson
## 5 M1    M5    0.93   27.0 4.50e-50 0.903    0.953 Pearson
## 6 M1    M6    0.76   12.1 7.33e-22 0.664    0.827 Pearson
## 7 M2    M1    0.94   30.2 1.02e-54 0.921    0.962 Pearson
## 8 M2    M2    1      Inf    0.      1      1      Pearson
## 9 M2    M3    0.92   23.8 5.21e-45 0.880    0.942 Pearson
## 10 M2   M4    0.92   24.8 1.22e-46 0.888    0.946 Pearson
## # ... with 26 more rows
```

```
# PROBLEM!! Absence of multicollinearity. The dependent (outcome) variables
# cannot be too correlated to each other. No correlation should be above r =
# 0.90 [Tabachnick and Fidell (2012)].
cor(Atlas_wofossil_noTub_sub[, 1:6])
```

```
##           M1           M2           M3           M4           M5           M6
## M1 1.0000000 0.9449857 0.8533793 0.9272517 0.9326445 0.7568932
## M2 0.9449857 1.0000000 0.9158753 0.9217077 0.8823482 0.7989239
## M3 0.8533793 0.9158753 1.0000000 0.8476626 0.8159639 0.7816608
## M4 0.9272517 0.9217077 0.8476626 1.0000000 0.9208258 0.7833759
## M5 0.9326445 0.8823482 0.8159639 0.9208258 1.0000000 0.7618434
```

```
## M6 0.7568932 0.7989239 0.7816608 0.7833759 0.7618434 1.0000000
```

MANOVA# : \*Failed multiple checks

```
Atlas.man <- manova(cbind(M1, M2, M3, M4, M5, M6) ~ species, data = Atlas_wofossil_noTub_sub)
summary(Atlas.man)
```

```
##              Df Pillai approx F num Df den Df      Pr(>F)
## species      9 2.1061      6.07    54    606 < 2.2e-16 ***
## Residuals 101
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
summary.aov(Atlas.man)
```

```
## Response M1 :
##              Df Sum Sq Mean Sq F value    Pr(>F)
## species      9 40.104  4.4560  27.517 < 2.2e-16 ***
## Residuals 101 16.355  0.1619
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

```
## Response M2 :
##              Df Sum Sq Mean Sq F value    Pr(>F)
## species      9 37.119  4.1244  26.124 < 2.2e-16 ***
## Residuals 101 15.945  0.1579
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

```
## Response M3 :
##              Df Sum Sq Mean Sq F value    Pr(>F)
## species      9 5.5115 0.61239  21.615 < 2.2e-16 ***
## Residuals 101 2.8615 0.02833
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

```
## Response M4 :
##              Df Sum Sq Mean Sq F value    Pr(>F)
## species      9 21.584 2.39828  16.607 < 2.2e-16 ***
## Residuals 101 14.586 0.14441
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

```
## Response M5 :
##              Df Sum Sq Mean Sq F value    Pr(>F)
## species      9 52.724  5.8582  35.407 < 2.2e-16 ***
## Residuals 101 16.711  0.1655
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

```
## Response M6 :
##              Df Sum Sq Mean Sq F value    Pr(>F)
## species      9  8.1785 0.90872   9.1615 4.933e-10 ***
## Residuals 101 10.0181 0.09919
```

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Permutation MANOVA ## \*Overcome failed checks

```
# Permutational Multivariate Analysis of Variance Using Distance Matrices
adonis(Atlas_wofossil_noTub_sub[, 1:6] ~ species, data = Atlas_wofossil_noTub_sub,
       method = "euclidean")
```

```
##
## Call:
## adonis(formula = Atlas_wofossil_noTub_sub[, 1:6] ~ species, data = Atlas_wofossil_noTub_sub,
##
## Permutation: free
## Number of permutations: 999
##
## Terms added sequentially (first to last)
##
##              Df SumsOfSqs MeanSqs F.Model      R2 Pr(>F)
## species        9   165.222  18.3580  24.245 0.68359 0.001 ***
## Residuals    101    76.477   0.7572      0.31641
## Total       110   241.698      1.00000
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
# pairwise comparisons between group levels with corrections for multiple
# testing
pairwise.perm.manova(Atlas_wofossil_noTub_sub[, 1:6], Atlas_wofossil_noTub_sub$species,
                    nperm = 50, progress = FALSE) #needs more permutation but takes a long time
```

```
##
## Pairwise comparisons using permutation MANOVAs (test: Pillai)
##
## data:  Atlas_wofossil_noTub_sub[, 1:6] by Atlas_wofossil_noTub_sub$species
## 50 permutations
##
##              A.gracile A.maculatum A.macrodactylum A.opacum
## A.maculatum      0.022      -            -            -
## A.macrodactylum 0.022      0.022          -            -
## A.opacum          0.040      0.022          0.022          -
## A.jeffersonianum 0.022      0.022          0.022          0.040
## A.mabeei         0.022      0.022          0.059          0.022
## A.texanum        0.022      0.022          0.022          0.022
## A.annulatum      0.022      0.022          0.022          0.022
## A.tigrinum       0.022      0.022          0.022          0.022
## A.mavortium      0.040      0.022          0.022          0.022
##
##              A.jeffersonianum A.mabeei A.texanum A.annulatum A.tigrinum
## A.maculatum      -            -            -            -
## A.macrodactylum -            -            -            -
## A.opacum         -            -            -            -
## A.jeffersonianum -            -            -            -
## A.mabeei         0.022      -            -            -
## A.texanum        0.022      0.022      -            -
## A.annulatum      0.022      0.022      0.040      -            -
```


```
## A.tigrinum      0.022      0.022      0.022      0.022      -
## A.mavortium     0.022      0.022      0.022      0.022      0.022
##
## P value adjustment method: fdr
# or using euclidean distances
AtlasPPM <- pairwise.perm.manova(dist(Atlas_wofossil_noTub_sub[, 1:6], "euclidean"),
  Atlas_wofossil_noTub_sub$species, nperm = 999)
AtlasPPM

##
## Pairwise comparisons using permutation MANOVAs on a distance matrix
##
## data: dist(Atlas_wofossil_noTub_sub[, 1:6], "euclidean") by Atlas_wofossil_noTub_sub$species
## 999 permutations
##
##      A.gracile A.maculatum A.macrodactylum A.opacum
## A.maculatum   0.0087      -                -                -
## A.macrodactylum 0.0020    0.0020          -                -
## A.opacum       0.0106    0.0020      0.0020          -
## A.jeffersonianum 0.2470    0.0037      0.0020      0.0047
## A.mabeei       0.0047    0.0020      0.1183      0.0095
## A.texanum      0.0020    0.0020      0.0020      0.0020
## A.annulatum    0.0836    0.0075      0.0020      0.0047
## A.tigrinum     0.0208    0.0252      0.0020      0.0020
## A.mavortium    0.0095    0.0201      0.0020      0.0020
##
##      A.jeffersonianum A.mabeei A.texanum A.annulatum A.tigrinum
## A.maculatum          -          -          -          -
## A.macrodactylum    -          -          -          -
## A.opacum             -          -          -          -
## A.jeffersonianum    -          -          -          -
## A.mabeei            0.0020      -          -          -
## A.texanum           0.0122      0.0020      -          -
## A.annulatum         0.1319      0.0020    0.0112      -
## A.tigrinum          0.0020      0.0020    0.0020    0.0047
## A.mavortium         0.0020      0.0037    0.0047    0.0112    0.0236
##
## P value adjustment method: fdr
```

## DFA

```
# PROBLEM:Check Multivariate normality:FAIL
mqqnorm(Atlas_wofossil_noTub_sub[, 1:6], main = "Multi-normal Q-Q Plot")
```

## Multi-normal Q-Q Plot



```
## [1] 50 61
```

```
# DFA# With MASS
```

```
library(MASS)
```

```
##
```

```
## Attaching package: 'MASS'
```

```
## The following object is masked from 'package:rstatix':
```

```
##
```

```
## select
```

```
## The following object is masked from 'package:dplyr':
```

```
##
```

```
## select
```

```
AtlasLDA <- lda(species ~ M1 + M2 + M3 + M4 + M5 + M6, data = Atlas_wofossil_noTub_sub,
  CV = FALSE) #DFA no jackknife
```

```
AtlasLDA
```

```
## Call:
```

```
## lda(species ~ M1 + M2 + M3 + M4 + M5 + M6, data = Atlas_wofossil_noTub_sub,
## CV = FALSE)
```

```
##
```

```
## Prior probabilities of groups:
```

## A.gracile	A.maculatum	A.macrodactylum	A.opacum
## 0.07207207	0.12612613	0.10810811	0.11711712
## A.jeffersonianum	A.mabeei	A.texanum	A.annulatum
## 0.09909910	0.08108108	0.08108108	0.05405405
## A.tigrinum	A.mavortium		
## 0.15315315	0.10810811		

```
##
```

```
## Group means:
##           M1           M2           M3           M4           M5           M6
## A.gracile      3.301250 3.227500 1.1475000 2.425000 3.305000 1.1750000
## A.maculatum    3.542143 3.787143 1.4935714 2.729286 3.442857 1.4521429
## A.macrodactylum 2.200833 2.234167 0.8316667 1.655000 2.088333 0.8066667
## A.opacum       2.709231 2.836154 1.0546154 1.982308 2.799231 1.0130769
## A.jeffersonianum 3.144545 3.211818 1.1481818 2.434545 2.997273 1.1618182
## A.mabeei       2.218889 2.440000 0.9411111 1.713333 2.268889 0.9200000
## A.texanum      3.011111 3.490000 1.3822222 2.382222 2.864444 1.3155556
## A.annulatum    3.305000 3.373333 1.2583333 2.400000 2.808333 1.2500000
## A.tigrinum     3.534118 3.487059 1.3176471 2.744706 3.968235 1.4082353
## A.mavortium    4.222500 4.206667 1.5216667 3.038333 4.310833 1.7458333
##
## Coefficients of linear discriminants:
##           LD1           LD2           LD3           LD4           LD5           LD6
## M1  0.55663683 -0.8606379 -5.5795956 -0.3483209  2.989576  1.097501598
## M2  0.86359598 -2.5415371  1.9312656 -3.1297756 -4.765508  0.001897505
## M3  4.33837399 -3.6814217  4.6634719  2.8925828  6.711399  2.898321950
## M4  3.68648363  0.3634334 -0.8034773  5.2104307 -1.277601 -1.351860806
## M5 -7.27345664  2.1171683  2.6638922 -0.7621234 -0.491493  0.461170573
## M6 -0.01259303  0.2267898 -0.7343062 -1.3772889  1.482124 -3.660023730
##
## Proportion of trace:
##           LD1           LD2           LD3           LD4           LD5           LD6
## 0.6619 0.2516 0.0620 0.0176 0.0047 0.0023

AtlasLDA_jack <- lda(species ~ M1 + M2 + M3 + M4 + M5 + M6, data = Atlas_wofossil_noTub_sub,
  CV = TRUE) #DFA with jackknife
# Assess the accuracy of jackknife #
accAtlasLDA <- table(Atlas_wofossil_noTub_sub$species, AtlasLDA_jack$class)
accAtlasLDA

##
##           A.gracile A.maculatum A.macrodactylum A.opacum
## A.gracile           3           0           0           2
## A.maculatum         0          12           0           0
## A.macrodactylum    0           0           9           0
## A.opacum             0           1           0           9
## A.jeffersonianum     1           2           0           2
## A.mabeei             0           0           3           1
## A.texanum            0           2           0           0
## A.annulatum          0           0           0           0
## A.tigrinum           2           0           0           0
## A.mavortium          1           2           0           1
##
##           A.jeffersonianum A.mabeei A.texanum A.annulatum A.tigrinum
## A.gracile                 2           0           0           0
## A.maculatum                0           0           2           0
## A.macrodactylum           0           3           0           0
## A.opacum                   2           1           0           0
## A.jeffersonianum           5           1           0           0
## A.mabeei                   0           5           0           0
## A.texanum                   0           0           7           0
## A.annulatum                 0           0           1           5
## A.tigrinum                  0           0           0           0
##                                     14
```

```
## A.mavortium          0          0          0          0          4
##
##           A.mavortium
## A.gracile          1
## A.maculatum        0
## A.macrodactylum   0
## A.opacum            0
## A.jeffersonianum    0
## A.mabeei            0
## A.texanum           0
## A.annulatum         0
## A.tigrinum          1
## A.mavortium         4
```

```
diag(prop.table(accAtlasLDA, 1))
```

```
##           A.gracile      A.maculatum  A.macrodactylum      A.opacum
##           0.3750000      0.8571429      0.7500000      0.6923077
## A.jeffersonianum      A.mabeei      A.texanum      A.annulatum
##           0.4545455      0.5555556      0.7777778      0.8333333
##           A.tigrinum      A.mavortium
##           0.8235294      0.3333333
```

```
sum(accAtlasLDA[row(accAtlasLDA) == col(accAtlasLDA)])/sum(accAtlasLDA)
```


```
## [1] 0.6576577
```

```
# DFA# With MORPHO
```

```
library(Morpho)
```

```
Atlascva = CVA(Atlas_wofossil_noTub_sub[, 1:6], groups = Atlas_wofossil_noTub_sub$species,
  rounds = 10000, cv = TRUE)
```

```
barplot(Atlascva$Var[, 2]) # Variance explained by the canonical roots
```



```
# get the typicality probabilities and resulting classifications all
# specimens with a probability of < 0.01 as outliers (assigned to no class)
typprobs <- typprobClass(Atlascva$CVscores, groups = Atlas_wofossil_noTub_sub$species,
  outlier = 0.01, cv = TRUE)
print(typprobs)
```

```
## cross-validated classification results in frequencies
```

```

##
##      A.annulatum A.gracile A.jeffersonianum A.mabeei
## A.gracile      0      3      2      0
## A.maculatum    0      0      0      0
## A.macrodactylum 0      0      0      3
## A.opacum       0      0      2      1
## A.jeffersonianum 0      2      6      1
## A.mabeei       0      0      0      7
## A.texanum      1      0      0      0
## A.annulatum    5      0      0      0
## A.tigrinum     0      2      0      0
## A.mavortium    0      1      0      0
##
##      A.macrodactylum A.maculatum A.mavortium A.opacum A.texanum
## A.gracile      0      0      1      2      0
## A.maculatum    0      11     0      0      2
## A.macrodactylum 9      0      0      0      0
## A.opacum       0      1      0      9      0
## A.jeffersonianum 0      1      0      0      1
## A.mabeei       2      0      0      0      0
## A.texanum      0      1      0      0      7
## A.annulatum    0      0      0      0      1
## A.tigrinum     0      0      1      0      0
## A.mavortium    0      1      4      1      0
##
##      A.tigrinum none
## A.gracile      0      0
## A.maculatum    0      1
## A.macrodactylum 0      0
## A.opacum       0      0
## A.jeffersonianum 0      0
## A.mabeei       0      0
## A.texanum      0      0
## A.annulatum    0      0
## A.tigrinum     12     2
## A.mavortium    3      2
##
## cross-validated classification result in %
##
##      A.annulatum A.gracile A.jeffersonianum A.mabeei
## A.gracile      0.0000 37.5000      25.0000  0.0000
## A.maculatum    0.0000  0.0000      0.0000  0.0000
## A.macrodactylum 0.0000  0.0000      0.0000 25.0000
## A.opacum       0.0000  0.0000     15.3846  7.6923
## A.jeffersonianum 0.0000 18.1818     54.5455  9.0909
## A.mabeei       0.0000  0.0000      0.0000 77.7778
## A.texanum      11.1111  0.0000      0.0000  0.0000
## A.annulatum    83.3333  0.0000      0.0000  0.0000
## A.tigrinum     0.0000 11.7647      0.0000  0.0000
## A.mavortium    0.0000  8.3333      0.0000  0.0000
##
##      A.macrodactylum A.maculatum A.mavortium A.opacum A.texanum
## A.gracile      0.0000  0.0000     12.5000 25.0000  0.0000

```



```
## A.maculatum 0.0000 78.5714 0.0000 0.0000 14.2857
## A.macrodactylum 75.0000 0.0000 0.0000 0.0000 0.0000
## A.opacum 0.0000 7.6923 0.0000 69.2308 0.0000
## A.jeffersonianum 0.0000 9.0909 0.0000 0.0000 9.0909
## A.mabeei 22.2222 0.0000 0.0000 0.0000 0.0000
## A.texanum 0.0000 11.1111 0.0000 0.0000 77.7778
## A.annulatum 0.0000 0.0000 0.0000 0.0000 16.6667
## A.tigrinum 0.0000 0.0000 5.8824 0.0000 0.0000
## A.mavortium 0.0000 8.3333 33.3333 8.3333 0.0000
##
## A.tigrinum none
## A.gracile 0.0000 0.0000
## A.maculatum 0.0000 7.1429
## A.macrodactylum 0.0000 0.0000
## A.opacum 0.0000 0.0000
## A.jeffersonianum 0.0000 0.0000
## A.mabeei 0.0000 0.0000
## A.texanum 0.0000 0.0000
## A.annulatum 0.0000 0.0000
## A.tigrinum 70.5882 11.7647
## A.mavortium 25.0000 16.6667
##
##
## overall classification accuracy: 65.76577 %
```

```
# Assess the accuracy of jackknife #
```

```
accJack <- table(Atlascva$groups, Atlascva$class)
accJack
```

```
##
## A.gracile A.maculatum A.macrodactylum A.opacum
## A.gracile 3 0 0 2
## A.maculatum 0 12 0 0
## A.macrodactylum 0 0 9 0
## A.opacum 0 1 0 9
## A.jeffersonianum 1 2 0 2
## A.mabeei 0 0 3 1
## A.texanum 0 2 0 0
## A.annulatum 0 0 0 0
## A.tigrinum 2 0 0 0
## A.mavortium 1 2 0 1
##
## A.jeffersonianum A.mabeei A.texanum A.annulatum A.tigrinum
## A.gracile 2 0 0 0 0
## A.maculatum 0 0 2 0 0
## A.macrodactylum 0 3 0 0 0
## A.opacum 2 1 0 0 0
## A.jeffersonianum 5 1 0 0 0
## A.mabeei 0 5 0 0 0
## A.texanum 0 0 7 0 0
## A.annulatum 0 0 1 5 0
## A.tigrinum 0 0 0 0 14
## A.mavortium 0 0 0 0 4
##
## A.mavortium
```

```
## A.gracile 1
## A.maculatum 0
## A.macrodactylum 0
## A.opacum 0
## A.jeffersonianum 0
## A.mabeei 0
## A.texanum 0
## A.annulatum 0
## A.tigrinum 1
## A.mavortium 4
```

```
diag(prop.table(accJack, 1))
```


```
## A.gracile A.maculatum A.macrodactylum A.opacum
## 0.3750000 0.8571429 0.7500000 0.6923077
## A.jeffersonianum A.mabeei A.texanum A.annulatum
## 0.4545455 0.5555556 0.7777778 0.8333333
## A.tigrinum A.mavortium
## 0.8235294 0.3333333
```

```
sum(accJack[row(accJack) == col(accJack)])/sum(accJack)
```


```
## [1] 0.6576577
```

#### Plot first two DF axes

```
AT_cva <- data.frame(AtlasCva$CVscores, species = AtlasCva$groups)
ggplot(AT_cva, aes(CV.1, CV.2)) + geom_point(size = 5, aes(color = species)) +
  theme_classic() + scale_color_brewer(palette = "Paired")
```




```
# alternative plot
plot(Atlascva$CVscores, col = Atlas_wofossil_noTub_sub$species, pch = as.numeric(Atlas_wofossil_noTub_sub$species),
     typ = "n", asp = 1, xlab = paste("1st canonical axis", paste(round(Atlascva$Var[1, 2], 1), "%")), ylab = paste("2nd canonical axis", paste(round(Atlascva$Var[2, 2], 1), "%")))
text(Atlascva$CVscores, as.character(Atlas_wofossil_noTub_sub$species), col = as.numeric(Atlas_wofossil_noTub_sub$species), cex = 0.7)
```



Plot Mahalanobis distances as dendrogram

```
dendroS = hclust(Atlascva$Dist$GroupdistMaha)
dendroS$labels = levels(Atlas_wofossil_noTub_sub$species)
par(mar = c(6.5, 4.5, 1, 1))
dendroS = as.dendrogram(dendroS)
plot(dendroS, main = "", sub = "", xlab = "", ylab = "Mahalanobis distance")
```



## Random Forest ###:Non-parametric

```
library(randomForest)

## randomForest 4.6-14
## Type rfNews() to see new features/changes/bug fixes.
##
## Attaching package: 'randomForest'
## The following object is masked from 'package:gridExtra':
##
##   combine
## The following object is masked from 'package:psych':
##
##   outlier
## The following object is masked from 'package:ggplot2':
##
##   margin
## The following object is masked from 'package:dplyr':
##
##   combine
Atlas.rf <- randomForest(species ~ M1 + M2 + M3 + M4 + M5 + M6, data = Atlas_wofossil_noTub_sub,
  importance = TRUE, proximity = TRUE)
```

```
print(Atlas.rf)
```

```
##
## Call:
## randomForest(formula = species ~ M1 + M2 + M3 + M4 + M5 + M6,      data = Atlas_wofossil_noTub_sub,
##               Type of random forest: classification
##               Number of trees: 500
## No. of variables tried at each split: 2
##
## OOB estimate of error rate: 34.23%
## Confusion matrix:
##               A.gracile A.maculatum A.macrodactylum A.opacum
## A.gracile           0           1           0           0
## A.maculatum          0          12           0           1
## A.macrodactylum     0           0          10           0
## A.opacum              1           0           2           7
## A.jeffersonianum      1           0           0           2
## A.mabeei              0           0           3           1
## A.texanum             0           0           0           0
## A.annulatum           0           0           0           0
## A.tigrinum            0           1           0           0
## A.mavortium           0           0           0           1
##
##               A.jeffersonianum A.mabeei A.texanum A.annulatum A.tigrinum
## A.gracile                     2           1           0           0           3
## A.maculatum                    0           0           0           0           0
## A.macrodactylum               0           2           0           0           0
## A.opacum                       1           2           0           0           0
## A.jeffersonianum               6           0           0           2           0
## A.mabeei                       0           5           0           0           0
## A.texanum                      0           0           8           1           0
## A.annulatum                    1           0           1           4           0
## A.tigrinum                     0           0           0           0          14
## A.mavortium                    1           0           1           0           2
##
##               A.mavortium class.error
## A.gracile           1  1.0000000
## A.maculatum          1  0.1428571
## A.macrodactylum     0  0.1666667
## A.opacum              0  0.4615385
## A.jeffersonianum      0  0.4545455
## A.mabeei              0  0.4444444
## A.texanum             0  0.1111111
## A.annulatum           0  0.3333333
## A.tigrinum            2  0.1764706
## A.mavortium           7  0.4166667
```

```
rf_acc <- Atlas.rf$confusion
rf_acc <- 1 - rf_acc[, 11] # percent correct classification
rf_acc
```

```
##           A.gracile      A.maculatum A.macrodactylum      A.opacum
##           0.0000000      0.8571429      0.8333333      0.5384615
## A.jeffersonianum      A.mabeei      A.texanum      A.annulatum
##           0.5454545      0.5555556      0.8888889      0.6666667
##           A.tigrinum      A.mavortium
```


```
##          0.8235294          0.5833333
```

```
# Look at variable importance
round(importance(Atlas.rf), 2)
```

```
##      A.gracile A.maculatum A.macrodactylum A.opacum A.jeffersonianum A.mabeei
## M1      1.97      11.13          5.12      8.44          4.47      16.07
## M2     -0.87      18.45          18.73      7.38          0.02      6.89
## M3     -2.16      23.27          11.84     10.25          11.06     -0.09
## M4     -0.73      9.98           7.36      7.72          4.82      6.69
## M5      5.73      15.79          22.77     13.50          12.47     10.31
## M6     -2.74      6.69           11.86     10.87          2.02      2.89
##      A.texanum A.annulatum A.tigrinum A.mavortium MeanDecreaseAccuracy
## M1      10.64      3.55          6.12      18.74          25.80
## M2      8.84      3.57          1.05      15.18          26.46
## M3      16.00      9.56         -0.89      3.30          27.70
## M4      5.06      5.19          8.20      1.26          16.37
## M5      16.91     10.89         31.19     16.91          44.81
## M6      3.63      0.55          8.24     10.03          18.71
##      MeanDecreaseGini
## M1              16.65
## M2              15.89
## M3              15.55
## M4              12.82
## M5              25.95
## M6              11.39
```

```
varImpPlot(Atlas.rf)
```

Atlas.rf




## K Nearest neighbor ###Non-parametric

```
library(tidyverse)
library(caret)

##
## Attaching package: 'caret'
## The following object is masked from 'package:vegan':
##
##   tolerance
## The following object is masked from 'package:purrr':
##
##   lift
Atlas_wofossil_noTub_sub <- column_to_rownames(Atlas_wofossil_noTub_sub, var = "specimen_num")

#make KNN model using LOOCV to find optimal k
KNNmodel <- train(
  species ~., data = Atlas_wofossil_noTub_sub, method = "knn",
  trControl = trainControl("LOOCV", number = 1),
  preProcess = c("center"), #center the data
  tuneLength = 15)

plot(KNNmodel) # plot accuracy vs k
```



```
KNNmodel$bestTune # optimal k
```

```
## k
## 2 7
```



```
predicted.classes <- KNNmodel %>% predict(Atlas_wofossil_noTub_sub[,1:6]) # predict class based on KNN
head(predicted.classes)
```

```
## [1] A.annulatum      A.jeffersonianum A.annulatum      A.annulatum
## [5] A.texanum          A.texanum
## 10 Levels: A.gracile A.maculatum A.macrodactylum A.opacum ... A.mavortium
```

```
mean(predicted.classes == Atlas_wofossil_noTub_sub$species) #overall accuracy
```

```
## [1] 0.7477477
```

```
# assess accuracy per species
```

```
accKNN <- table(Atlas_wofossil_noTub_sub$species,predicted.classes)
accKNN
```

```
##               predicted.classes
##               A.gracile A.maculatum A.macrodactylum A.opacum
## A.gracile           1           1           0           0
## A.maculatum          0          13           0           0
## A.macrodactylum     0           0          11           0
## A.opacum              0           1           2           8
## A.jeffersonianum      0           1           0           3
## A.mabeei              0           0           1           0
## A.texanum             0           0           0           0
## A.annulatum           0           0           0           0
## A.tigrinum            0           0           0           0
## A.mavortium           0           1           0           1
##               predicted.classes
##               A.jeffersonianum A.mabeei A.texanum A.annulatum A.tigrinum
## A.gracile                     2           1           0           0           3
## A.maculatum                   1           0           0           0           0
## A.macrodactylum              0           1           0           0           0
## A.opacum                      1           1           0           0           0
## A.jeffersonianum              7           0           0           0           0
## A.mabeei                     0           8           0           0           0
## A.texanum                    0           0           9           0           0
## A.annulatum                   1           0           2           3           0
## A.tigrinum                    0           0           0           0          15
## A.mavortium                   1           0           0           0           1
##               predicted.classes
##               A.mavortium
## A.gracile                0
## A.maculatum               0
## A.macrodactylum          0
## A.opacum                  0
## A.jeffersonianum          0
## A.mabeei                  0
## A.texanum                 0
## A.annulatum               0
## A.tigrinum                2
## A.mavortium              8
```

```
diag(prop.table(accKNN, 1))
```

```
##           A.gracile      A.maculatum A.macrodactylum      A.opacum
##           0.1250000      0.9285714      0.9166667      0.6153846
```

##	A.jeffersonianum	A.mabeei	A.texanum	A.annulatum
##	0.6363636	0.8888889	1.0000000	0.5000000
##	A.tigrinum	A.mavortium		
##	0.8823529	0.6666667		