

Fast method in evaluating density of multivariate Gaussian

- ▶ Given data $x \in \mathbb{R}^d$, the likelihood that it comes from a multivariate Gaussian density with mean vector $\mu \in \mathbb{R}^d$ and covariance matrix $\Sigma \in \mathbb{R}^{d \times d}$ is

$$\mathcal{N}(x; \mu, \Sigma) = \frac{1}{\sqrt{(2\pi)^d \det(\Sigma)}} e^{-\frac{1}{2}(x-\mu)^T \Sigma^{-1}(x-\mu)}$$

- ▶ The most expensive part to compute this is to evaluate Σ^{-1} , which has a complexity $\mathcal{O}(d^3)$.
- ▶ Moreover, when Σ is rank-deficient, i.e., there are close-to-zero eigenvalues, computing Σ^{-1} will return NAN (you cannot invert the matrix)

- ▶ Now let's make it faster and avoid the numerical issues by compute using “low-rank approximation”
- ▶ Compute eigendecomposition of

$$\Sigma = U\Lambda U^T$$

where $\Lambda = \text{diag}\{\lambda_1, \dots, \lambda_d\}$ and the eigenvalues are ordered

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d$$

- ▶ The rank- r approximation ($r < d$) of Σ is

$$\tilde{\Sigma} = \tilde{U}\tilde{\Lambda}\tilde{U}^T$$

where \tilde{U} is a d -by- r matrix formed by the first r columns of U , $\tilde{\Lambda} = \text{diag}\{\lambda_1, \dots, \lambda_r\}$.

- ▶ Typically we will choose r such that at least $\lambda_r > 0$

- Now compute transform of data and parameters

$$\tilde{x} = \tilde{U}^T x$$

$$\tilde{\mu} = \tilde{U}^T \mu$$

<https://www.cse.huji.ac.il/~csip/tirgul34.pdf>

- Compute $\tilde{\Lambda}^{-1} = \text{diag}\{\lambda_1^{-1}, \dots, \lambda_r^{-1}\}$
- Note that

$$\det(\Sigma) = \prod_{i=1}^d \lambda_i, \quad \det(\tilde{\Sigma}) = \prod_{i=1}^r \lambda_i$$

- Finally, the density calculated by replacing Σ with $\tilde{\Sigma}$ is:

$$\mathcal{N}(x; \mu, \Sigma) \approx \frac{1}{\sqrt{(2\pi)^d \prod_{i=1}^r \lambda_i}} \exp \left\{ -\frac{1}{2} \sum_{i=1}^r \frac{(\tilde{x}_i - \tilde{\mu}_i)^2}{\lambda_i} \right\}$$

where \tilde{x}_i and $\tilde{\mu}_i$ denote the i th entry of \tilde{x} and $\tilde{\mu}$, respectively.

- Note: you can play with different r to have a good tradeoff between accuracy and speed

- Note that above we have used the following basic identity from linear algebra

$$\tilde{\Sigma}^{-1} = \tilde{U} \tilde{\Lambda}^{-1} \tilde{U}^T$$

and

$$\begin{aligned} & (x - \mu)^T \Sigma^{-1} (x - \mu) \\ &= (x - \mu)^T \tilde{U} \tilde{\Lambda}^{-1} \tilde{U}^T (x - \mu) \\ &= [\tilde{U}^T (x - \mu)]^T \tilde{\Lambda}^{-1} [\tilde{U}^T (x - \mu)] \\ &= [\tilde{x} - \tilde{\mu}]^T \tilde{\Lambda}^{-1} [\tilde{x} - \tilde{\mu}] \\ &= \sum_{i=1}^r \frac{(\tilde{x}_i - \tilde{\mu}_i)^2}{\lambda_i} \end{aligned}$$