a Genetic correlation with adjusted ALM

Model	Construct	Genetic correlation	r_g (95% CI)	SE
Right	Regional FFM factor		0.779 (0.760–0.797)	0.0095
	Residual FFM		0.986 (0.972–1.001)	0.0073
Left	Regional FFM factor		0.745 (0.725–0.766)	0.0103
	Residual FFM		0.980 (0.965–0.996)	0.0080



b Unstandardized genetic covariance with adjusted ALM

Model	Construct	Genetic covariance	Covariance (95% CI)	SE
Right	Regional FFM factor		0.245 (0.224–0.267)	0.0108
	Residual FFM		0.223 (0.204–0.241)	0.0096
Left	Regional FFM factor		0.236 (0.216–0.257)	0.0106
	Residual FFM		0.211 (0.193–0.229)	0.0091



c Paired residual-minus-factor genetic correlation

Model	Paired Δr_g	Δr_g (95% CI)	P
Right primary		+0.208 (0.186–0.230)	<0.001
Left-side model		+0.235 (0.210–0.260)	<0.001



d Paired residual-minus-factor genetic covariance

Model	Paired ΔCov_g	ΔCov_g (95% CI)	P
Right primary		−0.0228 (−0.0308 to −0.0147)	<0.001
Left-side model		−0.0257 (−0.0342 to −0.0171)	<0.001