

Tolosat's Gravimetry Payload Report

November 11, 2021

Abstract

Tolosat's gravimetry payload aims to build a spherical harmonic model of the Earth's gravitational potential.

1 Introduction

Accurate knowledge of the gravitational potential of the Earth, on a global scale and at very high resolution, is a fundamental prerequisite for various geodetic, geophysical and oceanographic investigations and applications. Over the past 50 or so years, continuing improvements and refinements to the basic gravitational modeling theory have been paralleled by the availability of more accurate and complete data and by dramatic improvements in the computational resources available for numerical modeling studies.

These advances have brought the state-of-the-art from the early spherical harmonic models of degree 8 [?]

2 Method for unnoised model

The Earth's external gravitational potential, V , at a point P defined by its geocentric distance (r), geocentric co-latitude (θ) (defined as 90° -latitude), and longitude (λ), is given by:

$$V(r, \theta, \lambda) = \frac{GM}{r} \left[1 + \sum_{n=2}^{\infty} \left(\frac{R}{r} \right)^n \sum_{m=0}^n P_{nm}(\cos(\theta)) [C_{nm} \cos(m\lambda) + S_{nm} \sin(m\lambda)] \right] \quad (1)$$

where

$$\left\{ \begin{array}{l} R = \text{the Earth's equatorial radius} \\ P_{nm} = \text{the fully normalized associated Legendre functions of degree } n \text{ and order } m \\ C_{nm} S_{nm} \text{ the geopotential harmonic coefficients.} \end{array} \right.$$

We rewrite the geopotential harmonic series (1) :

$$V(r) = \sum_{n,m} [C_{nm} V_{nm}^c(r) + S_{nm} V_{nm}^s(r)] \quad (2)$$

This form shows explicitly the linearity of the harmonic coefficients C_{nm} et S_{nm}

The acceleration approach is based on the Newton's second law of motion, which links the acceleration vector to the gradient of the gravitational potential.

we obtain :

$$\frac{d^2 r}{dt^2} = a_{grav} + a_{lunisolar \text{ perturbations}} + a_{solid \text{ Earth and ocean tides}} + a_{\text{correction due to general relativity}} + a_{\text{non gravitational acceleration}}$$

2.1 A hypothesis on the accelerations

We studied the weight and importance of the accelerations. We conclude:

$$a_{grav} \gg \gg a_{others}$$

So, for the rest of the work we will consider that any accelerations other than a_{grav} is negligible.

2.2 A numerical approximation of the second derivative of the position

$$\frac{d^2 r}{dt^2} \approx \frac{d^2 Q(r_{gps})}{dt^2} = F * r_{gps} \quad (3)$$

Where r_{gps} is the GPS position. We do this approximation using a polynomial smoothing filter. We use as a filter the Savitzky-Golay filter

We obtain the observation equation

$$\frac{d^2 Q(r_{gps})}{dt^2} - a_{other} = \sum_{n,m} [C_{nm} \nabla V_{nm}^c(r) + S_{nm} \nabla V_{nm}^s(r)] \quad (4)$$

By solving this system, we obtain the harmonic coefficients.

We write the model:

$$y = Ax \quad (5)$$

the vector of observations $y = \frac{d^2 Q(r_{gps})}{dt^2}$

the design matrix $A = [\nabla V_{nm}^c; \nabla V_{nm}^s]$

the estimated parameters $x = [C_{nm}; S_{nm}]$

Intuitively, one can think about solving the problem with:

$$x = (A^T A)^{-1} A^T y$$

However, the dimensions of the matrix A are big. Numerical calculus would be very sensible to numerical errors.

We use a classic least square method to solve this problem.

2.3 Resultats

Our first experiences showed that this model is very sensible towards small variations of θ, ϕ .

We suggest to use a projection on a space.

Let this projection be:

$$P_E : \left| \begin{array}{ccc} \mathbb{R}^3 & \longrightarrow & \mathbb{R}^3 \\ \left(\begin{array}{c} r \\ \theta \\ \phi \end{array} \right) & \longmapsto & \left(\begin{array}{c} r \\ 0 \\ 0 \end{array} \right) \end{array} \right. \quad (6)$$

In other words, we focus on the radial acceleration.

Let

$$Ax = y \Leftrightarrow P_E Ax = P_E y = \tilde{y} \implies x \approx \tilde{x} = Px$$

3 Method for noised model

Let us suppose, for instance, that the measurements of the positions are noised.
i.e

$$\begin{aligned} p &= \tilde{p} + \epsilon \\ Sp &= S\tilde{p} + S\epsilon \quad \text{where S is the Savitzky Golay Filter} \\ x &= \tilde{x} + S\epsilon \\ x &= (A^T A)^{-1} A^T S\tilde{p} + (A^T A)^{-1} A^T S\epsilon \end{aligned}$$

If we suppose that $\epsilon \sim \mathcal{N}(0, \sigma I)$ we can obtain ,

$$\epsilon_2 = Q\epsilon \quad \mathbb{E}[\epsilon\epsilon^T] = \sigma^2 Q Q^T$$

Where $Q = (A^T A)^{-1} S$

Note that we should add in the equation a matrix B of noise decorrelation