

Combinatorial Analysis

A Complete Reference Manual on Permutations, Combinations,
Decision Logic, and Pitfalls

Department of Mathematics
Advanced Combinatorics & Problem Solving Division

September 19, 2026

Abstract

This reference document presents a rigorous, exhaustive synthesis of Permutations and Combinations (Combinatorial Analysis). It incorporates fundamental counting principles, factorial algebra, distinct and non-distinct linear/circular arrangements, combinatorial identities, partition and distribution theory (including Stars and Bars), derangements, geometric applications, operational decision frameworks, and an extensive catalog of mathematical pitfalls and competitive traps.

Contents

1	Fundamental Principles of Counting (FPC)	3
1.1	The Rule of Product (Multiplication Principle)	3
1.1.1	Generalization to k Stages	3
1.1.2	Operational Conditions	3
1.1.3	Application Methodology	3
1.2	The Rule of Sum (Addition Principle)	3
1.2.1	Generalization and Inclusion-Exclusion Principle	4
2	Factorial Algebra and Divisibility Theory	4
2.1	Factorial Function Definition and Properties	4
2.1.1	Key Algebraic Identities	4
2.2	Legendre's Formula (Exponent of Prime p in $n!$)	4
2.2.1	Application: Number of Trailing Zeros in $n!$	4
3	Permutations (Ordered Arrangements)	4
3.1	Permutations of Distinct Objects	5
3.1.1	Without Repetition	5
3.1.2	With Unlimited Repetition	5
3.2	Permutations of Multisets (Objects Not All Distinct)	5
3.3	Circular Permutations	5

3.3.1	Case 1: Clockwise and Counter-Clockwise Orders Are Distinct	5
3.3.2	Case 2: Clockwise and Counter-Clockwise Orders Are Indistinguishable	5
3.4	Special Structural Techniques for Constrained Permutations	6
3.4.1	1. The Grouping Method (String / Tie Method)	6
3.4.2	2. The Gap Method	6
3.4.3	3. Lexicographical Ordering (Dictionary Rank Algorithm)	6
4	Combinations (Unordered Selections)	6
4.1	Fundamental Formula	7
4.2	Essential Combinatorial Identities	7
4.3	Summation Properties of Binomial Coefficients	7
4.4	Selections from Groups (At Least One Object)	7
4.4.1	Selecting from n Distinct Objects	7
4.4.2	Selecting from Identical Objects	8
4.4.3	Combined Distinct and Identical Objects	8
5	Partitioning, Group Division, and Distribution	8
5.1	Division of Distinct Objects into Groups	8
5.1.1	Unequal Groups	8
5.1.2	Equal Groups (The Symmetry Factor Trap)	8
5.2	Distribution of Identical Objects (Stars and Bars Theorem)	8
6	Derangements and Inclusion-Exclusion	9
6.1	Derangement Formula (D_n)	9
6.2	Recurrence Relations and Asymptotic Value	9
6.2.1	Small Values Lookup Table	9
6.3	Partial Derangements	9
7	Geometric Applications of Combinatorics	10
8	Master Decision Logic Matrix	10
9	Exhaustive Guide to Tricks, Traps, and Pitfalls	10
10	Summary Checklist for Problem Solving	12

1 Fundamental Principles of Counting (FPC)

Counting without explicit enumeration forms the cornerstone of discrete mathematics. The fundamental counting framework relies on two complementary rules: the Rule of Product and the Rule of Sum.

1.1 The Rule of Product (Multiplication Principle)

Definition 1.1 (Sequential Multi-Stage Counting). If an event E_1 can occur in m distinct ways, and following its occurrence, a second independent event E_2 can occur in n distinct ways, then the total number of ordered pairs of occurrences (E_1, E_2) in the specified order is:

$$N = m \times n \quad (1)$$

1.1.1 Generalization to k Stages

For a sequence of k events $E_1, E_2, \dots, E_k$, where event E_i can occur in n_i ways regardless of the outcomes of prior stages:

$$N = \prod_{i=1}^k n_i = n_1 \times n_2 \times \dots \times n_k \quad (2)$$

1.1.2 Operational Conditions

- **Sequential Dependence of Process:** Events occur in a multi-step sequence or simultaneous combination.
- **Choice Invariance:** The *number* of choices n_i at stage i must remain constant regardless of which specific choice was made in preceding stages $1, 2, \dots, i - 1$.

1.1.3 Application Methodology

1. Identify the total number of decision positions or slots $(S_1, S_2, \dots, S_k)$.
2. Analyze constraints on individual slots (e.g., non-zero leading digits, even unit digits).
3. **Fill the most restricted slots first.**
4. Compute choices for remaining slots and apply $\prod n_i$.

1.2 The Rule of Sum (Addition Principle)

Definition 1.2 (Disjoint Mutually Exclusive Alternatives). If an event E can occur either via Case A (in m distinct ways) or via Case B (in n distinct ways), where Case A and Case B are mutually exclusive ($A \cap B = \emptyset$), then the total number of ways event E can occur is:

$$N = m + n \quad (3)$$

1.2.1 Generalization and Inclusion-Exclusion Principle

For k mutually exclusive cases $C_1, C_2, \dots, C_k$:

$$N = \sum_{i=1}^k n_i = n_1 + n_2 + \dots + n_k \quad (4)$$

When cases overlap ($A \cap B \neq \emptyset$), the Principle of Inclusion-Exclusion (PIE) must be applied:

$$|A \cup B| = |A| + |B| - |A \cap B| \quad (5)$$

$$|A \cup B \cup C| = |A| + |B| + |C| - (|A \cap B| + |B \cap C| + |C \cap A|) + |A \cap B \cap C| \quad (6)$$

2 Factorial Algebra and Divisibility Theory

2.1 Factorial Function Definition and Properties

For any non-negative integer $n \in \mathbb{Z}_{\geq 0}$, the factorial function $n!$ is defined as:

$$n! = \begin{cases} 1 & \text{if } n = 0 \\ \prod_{k=1}^n k = 1 \times 2 \times 3 \times \dots \times n & \text{if } n \geq 1 \end{cases} \quad (7)$$

2.1.1 Key Algebraic Identities

- **Recurrence Relation:** $n! = n \cdot (n-1)!$ for $n \geq 1$.
- **Double Factorial:** For odd n : $n!! = 1 \cdot 3 \cdot 5 \dots n$. For even n : $n!! = 2 \cdot 4 \cdot 6 \dots n = 2^{n/2} (n/2)!$.
- **Gamma Function Extension:** $\Gamma(n) = (n-1)!$, with $\Gamma(1) = 0! = 1$.

2.2 Legendre's Formula (Exponent of Prime p in $n!$)

To determine the highest power of a prime p that divides $n!$, denoted $E_p(n!)$ or $v_p(n!)$:

Formula 2.1 (Legendre's Theorem).

$$E_p(n!) = \sum_{k=1}^{\lfloor \log_p n \rfloor} \left\lfloor \frac{n}{p^k} \right\rfloor = \left\lfloor \frac{n}{p} \right\rfloor + \left\lfloor \frac{n}{p^2} \right\rfloor + \left\lfloor \frac{n}{p^3} \right\rfloor + \dots \quad (8)$$

where $\lfloor x \rfloor$ denotes the floor function (greatest integer $\leq x$).

2.2.1 Application: Number of Trailing Zeros in $n!$

The number of trailing zeros in $n!$ in base 10 equals $\min(E_2(n!), E_5(n!)) = E_5(n!)$:

$$Z(n!) = \left\lfloor \frac{n}{5} \right\rfloor + \left\lfloor \frac{n}{25} \right\rfloor + \left\lfloor \frac{n}{125} \right\rfloor + \dots \quad (9)$$

3 Permutations (Ordered Arrangements)

A permutation is an arrangement of a set of objects into a specified order.

3.1 Permutations of Distinct Objects

3.1.1 Without Repetition

The number of ways to arrange r objects selected from n distinct objects ($0 \leq r \leq n$):

$${}^nP_r = P(n, r) = \frac{n!}{(n-r)!} = n(n-1)(n-2) \dots (n-r+1) \quad (10)$$

Operational Conditions:

1. All n source objects are completely distinct.
2. No object can be used more than once ($r \leq n$).
3. The ordering of elements within chosen positions creates distinct outcomes.

3.1.2 With Unlimited Repetition

The number of arrangements of r positions using n distinct objects where each object can be reused indefinitely:

$$P_{\text{rep}}(n, r) = n^r \quad (11)$$

3.2 Permutations of Multisets (Objects Not All Distinct)

Formula 3.1 (Multiset Permutations). The number of distinct linear permutations of n objects, where p_1 objects are of type 1, p_2 objects are of type 2, ..., p_k objects are of type k (with $\sum_{i=1}^k p_i = n$), is given by the multinomial coefficient:

$$N = \binom{n}{p_1, p_2, \dots, p_k} = \frac{n!}{p_1! p_2! p_3! \dots p_k!} \quad (12)$$

3.3 Circular Permutations

When objects are arranged in a continuous loop, shifting all objects by one position does not yield a new arrangement.

3.3.1 Case 1: Clockwise and Counter-Clockwise Orders Are Distinct

For seating people around a circular table:

$$P_{\text{circ}}(n) = (n-1)! \quad (13)$$

If selecting and arranging r objects out of n distinct objects in a circle:

$$P_{\text{circ}}(n, r) = \frac{{}^nP_r}{r} = \frac{n!}{r(n-r)!} \quad (14)$$

3.3.2 Case 2: Clockwise and Counter-Clockwise Orders Are Indistinguishable

For arrangements like beads on a necklace, keys on a ring, or flowers in a garland (where flipping the ring over reverses clockwise to counter-clockwise):

$$P_{\text{circ, flip}}(n) = \frac{(n-1)!}{2} \quad (15)$$

3.4 Special Structural Techniques for Constrained Permutations

3.4.1 1. The Grouping Method (String / Tie Method)

Condition: When k specific objects MUST ALWAYS occur together in a block.

Procedure:

1. Tie the k specific objects together to form 1 single "super object".
2. Total objects to arrange = $(n - k + 1)$.
3. Arrange these $(n - k + 1)$ objects in $(n - k + 1)!$ ways.
4. Arrange the k objects *internally* inside the block in $k!$ ways.
5. Total Permutations:

$$N_{\text{together}} = (n - k + 1)! \times k! \quad (16)$$

3.4.2 2. The Gap Method

Condition: When k specific objects MUST NEVER occur adjacent to each other.

Procedure:

1. First arrange the remaining $(n - k)$ unconstrained objects in $(n - k)!$ ways.
2. This creates $(n - k + 1)$ available gaps (including ends):

$$_ O_1 _ O_2 _ O_3 _ \dots _ O_{n-k} _$$

3. Select k gaps out of $(n - k + 1)$ and place the k constrained objects:

$$N_{\text{separated}} = (n - k)! \times {}^{n-k+1}P_k = (n - k)! \times \frac{(n - k + 1)!}{(n - 2k + 1)!} \quad (17)$$

4. **Constraint Check:** This method requires $k \leq n - k + 1 \implies 2k \leq n + 1$.

3.4.3 3. Lexicographical Ordering (Dictionary Rank Algorithm)

To compute the rank of a word formed by distinct letters in alphabetical order:

1. List all letters of the word in alphabetical order.
2. Fix letters prior to the target letter at the current position, and count permutations of remaining letters.
3. Sum the counts for all preceding possibilities and add 1 for the target word.

4 Combinations (Unordered Selections)

A combination is a selection of items where order does not matter.

4.1 Fundamental Formula

The number of ways to select r unordered objects from n distinct objects ($0 \leq r \leq n$):

$${}_nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{{}^nP_r}{r!} \quad (18)$$

4.2 Essential Combinatorial Identities

Theorem 4.1 (Standard Identities). For non-negative integers n, r :

$$\text{Symmetric Property: } \binom{n}{r} = \binom{n}{n-r} \quad (19)$$

$$\text{Pascal's Identity: } \binom{n}{r} + \binom{n}{r-1} = \binom{n+1}{r} \quad (20)$$

$$\text{Equality Condition: } \binom{n}{x} = \binom{n}{y} \iff x = y \text{ or } x + y = n \quad (21)$$

$$\text{Ratio Rule: } \frac{\binom{n}{r}}{\binom{n}{r-1}} = \frac{n-r+1}{r} \quad (22)$$

$$\text{Absorption Identity: } \binom{n}{r} = \frac{n}{r} \binom{n-1}{r-1} \quad (23)$$

$$\text{Hockey-Stick Identity: } \sum_{i=r}^n \binom{i}{r} = \binom{r}{r} + \binom{r+1}{r} + \cdots + \binom{n}{r} = \binom{n+1}{r+1} \quad (24)$$

$$\text{Vandermonde's Identity: } \sum_{k=0}^r \binom{m}{k} \binom{n}{r-k} = \binom{m+n}{r} \quad (25)$$

4.3 Summation Properties of Binomial Coefficients

$$\sum_{r=0}^n \binom{n}{r} = \binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n \quad (26)$$

$$\sum_{r=0}^n (-1)^r \binom{n}{r} = \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \cdots = 0 \quad (27)$$

$$\sum_{r \text{ even}} \binom{n}{r} = \sum_{r \text{ odd}} \binom{n}{r} = 2^{n-1} \quad (28)$$

4.4 Selections from Groups (At Least One Object)

4.4.1 Selecting from n Distinct Objects

The total number of ways to select one or more objects from n distinct objects:

$$\sum_{r=1}^n \binom{n}{r} = 2^n - 1 \quad (29)$$

4.4.2 Selecting from Identical Objects

If there are p identical objects of type 1, q identical of type 2, r identical of type 3:

- Number of ways to select any number of objects (including zero): $(p + 1)(q + 1)(r + 1)$.
- Number of ways to select **at least one object**:

$$N = (p + 1)(q + 1)(r + 1) - 1 \quad (30)$$

4.4.3 Combined Distinct and Identical Objects

For p identical of type 1, q identical of type 2, and k distinct objects:

$$N_{\text{at least 1}} = (p + 1)(q + 1)2^k - 1 \quad (31)$$

5 Partitioning, Group Division, and Distribution

5.1 Division of Distinct Objects into Groups

5.1.1 Unequal Groups

The number of ways to divide $(m + n + p)$ distinct objects into three specified unequal groups of sizes m, n, p :

$$N = \binom{m + n + p}{m} \binom{n + p}{n} \binom{p}{p} = \frac{(m + n + p)!}{m! n! p!} \quad (32)$$

5.1.2 Equal Groups (The Symmetry Factor Trap)

When dividing objects into groups of equal sizes, we must distinguish between **unnamed (unlabelled)** groups and **named (labelled / distributed)** groups.

Formula 5.1 (Group Division Rules). For dividing $k \cdot m$ distinct objects into k groups of size m each:

1. Unnamed / Unlabelled Groups (Partitioning only):

$$N_{\text{unnamed}} = \frac{(km)!}{k! (m!)^k} \quad (33)$$

Reason: The $k!$ in the denominator corrects for overcounting due to identical group sizes.

2. Named Groups / Distributed to k Distinct Recipients:

$$N_{\text{distributed}} = N_{\text{unnamed}} \times k! = \frac{(km)!}{(m!)^k} \quad (34)$$

5.2 Distribution of Identical Objects (Stars and Bars Theorem)

Theorem 5.1 (Stars and Bars / Beggar's Method). Let n be the number of identical objects to be distributed among r distinct recipients (or bins).

1. **Non-Negative Integer Solutions (Empty bins allowed, $x_i \geq 0$):** The number of non-negative integer solutions to $x_1 + x_2 + \cdots + x_r = n$ is:

$$N(x_i \geq 0) = \binom{n+r-1}{r-1} = \binom{n+r-1}{n} \quad (35)$$

2. **Positive Integer Solutions (No empty bins, $x_i \geq 1$):** The number of positive integer solutions to $x_1 + x_2 + \cdots + x_r = n$ (where $n \geq r$) is:

$$N(x_i \geq 1) = \binom{n-1}{r-1} \quad (36)$$

3. **Lower Bound Constrained Solutions ($x_i \geq k_i$):** Substitute $y_i = x_i - k_i \geq 0$. The equation becomes $\sum y_i = n - \sum k_i = n'$.

$$N = \binom{n' + r - 1}{r - 1} \quad \text{where } n' = n - \sum_{i=1}^r k_i \quad (37)$$

6 Derangements and Inclusion-Exclusion

A derangement is a permutation of a set of objects such that no object appears in its original or natural position.

6.1 Derangement Formula (D_n)

Formula 6.1 (De Montmort's Derangement Formula). The number of derangements of n distinct objects, denoted D_n or $!n$, is:

$$D_n = n! \sum_{k=0}^n \frac{(-1)^k}{k!} = n! \left(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots + (-1)^n \frac{1}{n!} \right) \quad (38)$$

6.2 Recurrence Relations and Asymptotic Value

$$\text{First Recurrence: } D_n = (n-1)(D_{n-1} + D_{n-2}) \quad \text{for } n \geq 2 \quad (39)$$

$$\text{Second Recurrence: } D_n = nD_{n-1} + (-1)^n \quad \text{for } n \geq 1 \quad (40)$$

$$\text{Asymptotic Value: } \lim_{n \rightarrow \infty} \frac{D_n}{n!} = \frac{1}{e} \approx 0.367879 \quad (41)$$

6.2.1 Small Values Lookup Table

n	1	2	3	4	5	6
D_n	0	1	2	9	44	265

6.3 Partial Derangements

The number of ways to arrange n objects such that **exactly** r objects stay in their original positions and the remaining $(n-r)$ objects are deranged:

$$N_{\text{partial}} = \binom{n}{r} D_{n-r} \quad (42)$$

7 Geometric Applications of Combinatorics

For n points located in a 2D plane:

Formula 7.1 (Geometric Formulations). 1. **Straight Lines:** If no 3 points are collinear, total lines = $\binom{n}{2}$. If m points are collinear:

$$L = \binom{n}{2} - \binom{m}{2} + 1 \quad (43)$$

2. **Triangles:** If no 3 points are collinear, total triangles = $\binom{n}{3}$. If m points are collinear:

$$T = \binom{n}{3} - \binom{m}{3} \quad (44)$$

3. **Diagonals of an n -sided Polygon:**

$$D = \binom{n}{2} - n = \frac{n(n-3)}{2} \quad (45)$$

4. **Interior Intersection Points of Diagonals:** For a convex n -sided polygon where no three diagonals intersect at a single interior point:

$$I = \binom{n}{4} \quad (46)$$

5. **Parallelograms Formed by Intersecting Parallel Lines:** Given m parallel lines intersected by n parallel lines:

$$P = \binom{m}{2} \times \binom{n}{2} \quad (47)$$

8 Master Decision Logic Matrix

When approaching a combinatorial problem, use the following matrix to identify the appropriate formula:

9 Exhaustive Guide to Tricks, Traps, and Pitfalls

Pitfall & Trap 9.1 (The Leading Zero Trap in Digit Problems). When forming n -digit numbers using digits including 0, the leading digit position cannot be 0.

- **Incorrect Method:** Treating the leading digit as having n choices.
- **Correct Method:** Fill the highest place value first with $(n-1)$ available non-zero choices, then fill the remaining $(n-1)$ places.

Pitfall & Trap 9.2 (Overcounting Group Divisions (Symmetry Factor Error)). Dividing $2n$ distinct items into 2 equal groups of n items each.

- **Mistake:** Writing $\frac{(2n)!}{n!n!} = \binom{2n}{n}$. This assumes the two groups are distinct (e.g., Group A and Group B).

Problem Type	Objects	Bins/Positions	Order?	Master Formula
Linear Selection	Distinct	–	No	$\binom{n}{r} = \frac{n!}{r!(n-r)!}$
Linear Arrangement	Distinct	–	Yes	${}^nP_r = \frac{n!}{(n-r)!}$
Arrangement w/ Rep.	Distinct	–	Yes	n^r
Multiset Permutation	p_1, p_2 Identical	–	Yes	$\frac{n!}{p_1! p_2! \dots}$
Circular Arrangement	Distinct	Circle	Yes	$(n-1)!$
Necklace / Garland	Distinct	Circle (Flip)	Yes	$\frac{(n-1)!}{2}$
Identical Distribution	Identical (n)	Distinct (r)	No	$\binom{n+r-1}{r-1}$ (empty allowed)
Identical Dist. (No empty)	Identical (n)	Distinct (r)	No	$\binom{n-1}{r-1} (x_i \geq 1)$
Equal Group Division	Distinct (km)	Unnamed (k)	No	$\frac{(km)!}{k! (m!)^k}$
Equal Group Dist.	Distinct (km)	Named (k)	No	$\frac{(km)!}{(m!)^k}$
Derangement	Distinct	Fixed positions	Yes	$n! \sum_{k=0}^n \frac{(-1)^k}{k!}$

Table 1: Decision Matrix for Problem Classification

- **Correction:** If groups are unlabelled, divide by $2!$ to yield $\frac{(2n)!}{2! (n!)^2}$.

Pitfall & Trap 9.3 (Misapplication of the Gap Method). Placing constrained items into gaps before arranging the unconstrained base items.

- **Mistake:** Calculating gaps based on total elements n instead of unconstrained elements $(n - k)$.
- **Correction:** Always place the $(n - k)$ unconstrained elements first, count the resulting $(n - k + 1)$ gaps, and then choose k gaps.

Pitfall & Trap 9.4 (Forgetting Internal Permutations in Grouping). Grouping k items together and forgetting that they can swap positions among themselves.

- **Mistake:** Writing $(n - k + 1)!$.
- **Correction:** Multiply by internal arrangements $k! \implies (n - k + 1)! \times k!$.

Pitfall & Trap 9.5 (Base vs. Exponent Confusion in n^r). Distributing r distinct items into n distinct boxes with repetition allowed.

- **Mistake:** Writing r^n .
- **Rule:** Formula = (Choices per item)^(Number of items) = n^r .

Pitfall & Trap 9.6 (Misapplying Stars and Bars to Distinct Objects). Using $\binom{n+r-1}{r-1}$ when the objects being distributed are distinct.

- **Correction:** Stars and Bars applies **ONLY** to identical objects. For distinct objects into distinct boxes, use exponential distribution (r^n) or Stirling numbers of the second kind.

10 Summary Checklist for Problem Solving

1. **Identify Order Relevance:** Does changing the order of selected elements create a new state? If yes → Permutations. If no → Combinations.
2. **Identify Object Nature:** Are the objects distinct or identical?
3. **Identify Repetition Limits:** Is repetition permitted, forbidden, or multi-set constrained?
4. **Check Boundary Cases:** Check for $0!$, empty bins ($x_i = 0$), leading zeros, and symmetry factors in equal group divisions.