

Disclaimer

Please ensure you read and understand the following information prior to completing the medical assessment. The consultation should not take longer than 3-5 minutes to complete.

It is important that you answer all the questions correctly and honestly to allow the Pharma Health clinical team to assess your needs and ensure the treatment is safe and effective for you.

The information you provide us is treated with the utmost confidentiality & will be reviewed by a UK registered pharmacist prescriber. The questions listed in this medical assessment will help provide the pharmacist prescriber with an appropriate level of information to make an informed decision on whether the treatment is suitable for your condition.

Confirmation

By clicking “Continue”, you are confirming that you:

General

- Are 18 years old or above.
 - Are completing this questionnaire for yourself and to the best of your knowledge.
 - Understand that any treatment prescribed for you is for your personal use only.
 - Will disclose any serious illnesses or operations you have had.
 - Will disclose any medications (including prescription and non-prescription medications) you currently take.
 - Are happy for us to contact you if we have any queries about your condition.
 - Agree to follow any additional advice given by the Pharma Health clinical team.
 - Have read, understood and agreed to our Terms & Conditions, Privacy Notice and Terms of Sale.
-
- Was assigned male at birth.
 - Are experiencing some form of Erectile Dysfunction (ED) i.e. cannot regularly get or maintain an erection that is hard enough for sexual intercourse.
 - Understand that you should not take any combination of Viagra/Sildenafil, Cialis/Tadalafil, Levitra/Vardenafil or use any other ED treatment at the same time.
 - Understand that a higher dose of Sildenafil, Tadalafil (non
 - daily), Vardenafil and its equivalent brands (Viagra, Cialis and Levitra) should only be tried if a lower dose has been tried several times in the past year and found to be ineffective.

- Understand that you should always start Tadalafil daily at a higher dose and taper down to the lower dose once you have achieved a good effect.
- Understand that although unlikely, some side effects of these ED medications include: headaches, facial flushing, indigestion, nausea, lightheadedness, and changes in vision.

Generated on: 06-29-2024 10:00:02 am