

Disclaimer

Please ensure you read and understand the following information prior to completing the medical assessment. The consultation should not take longer than 3-5 minutes to complete.

It is important that you answer all the questions correctly and honestly to allow the Pharma Health clinical team to assess your needs and ensure the treatment is safe and effective for you.

The information you provide us is treated with the utmost confidentiality & will be reviewed by a UK registered pharmacist prescriber. The questions listed in this medical assessment will help provide the pharmacist prescriber with an appropriate level of information to make an informed decision on whether the treatment is suitable for your condition.

Confirmation

By clicking “Continue”, you are confirming that you:

General

- Are 18 years old or above.
- Are completing this questionnaire for yourself and to the best of your knowledge.
- Understand that any treatment prescribed for you is for your personal use only.
- Will disclose any serious illnesses or operations you have had.
- Will disclose any medications (including prescription and non-prescription medications) you currently take.
- Are happy for us to contact you if we have any queries about your condition.
- Agree to follow any additional advice given by the Pharma Health clinical team.
- Have read, understood and agreed to our Terms & Conditions, Privacy Notice and Terms of Sale.
- Understand that this service should not be used if you think you've had a sudden allergic reaction to something, or if you're systemically unwell (fever, nausea, dizziness etc.). You should contact your GP as soon as possible instead.
- Understand that this is not a mole
- checking service as we are unable to provide accurate and safe mole advice using this service. If you have any concerns about a mole or skin cancer, please book contact your GP.
- Are aware that oral Lymecline antibiotics should not be taken whilst using Zineryt solution or Duac gel. Oral Lymecline antibiotics are best used with topical Differin, Epiduo or Benzoyl Peroxide Gels. This reduces the

chance of the bacteria developing antibiotic resistance.

- Are you aware that antibiotic treatments (Lymecycline, Zineryt and Duac) should not normally be used continuously for more than 3 months.
- Will stop using Duac if severe erythema (severe redness), dryness, itching or burning of the affected area occurs.
- Understand that Duac treatment can cause irritation on the skin. If this occurs, either reduce frequency of use to alternate days until symptoms improve, or stop altogether and try applying on alternate days once skin recovered. You can continue to use for up to 12 weeks at the tolerated dose.
- Understand that this medication can make your skin more sensitive to sunlight (photosensitivity). You should avoid sunlight and sunlamps while using this gel and take care to use sunscreen and cover in protective clothing if sunlight exposure is necessary.
- Are aware it usually takes 8 weeks of continuous usage for acne treatments to work. Acne treatment should be stopped or reduced after 4
- 6 months, if acne has cleared up.

Generated on: 10-17-2024 12:51:40 pm