

Math Notes

Tyler Wilson

Contents

1	Complex Analysis	1
1.1	Complex Algebra	1
1.1.1	Complex Arithmetic	1
1.1.2	De Moirre's Formula	6
1.1.3	Geometry in the Complex Plane	10
1.1.4	Roots of a Complex Number	11
1.2	Complex Functions	13
1.2.1	Mapping Properties of Simple Functions	13
1.2.2	Calculus of Complex Functions	19
1.2.3	Conformal Mappings	21
1.2.4	Conformal Mapping to Solve Laplace's Equation	24
1.2.5	Sinusoidal Functions	26

1 Complex Analysis

1.1 Complex Algebra

Complex numbers arise from the roots of polynomials.

Ex: $x^2 + 1 = 0 \Rightarrow x^2 = -1 \Rightarrow x = \pm\sqrt{-1}$. This polynomial has no real roots, however, we can introduce an imaginary number i such that $i^2 = -1$. Then we will have the solution $x = \pm i$

We can introduce *complex numbers* which are numbers in the form $z = x + iy$, where x is the real part of z , $\Re(z)$, and y is the imaginary part of z , $\Im(z)$. These numbers can also be expressed in vector notation along the complex plane.

1.1.1 Complex Arithmetic

Addition, subtraction, and multiplication work the same, just with the addition of the fact $i^2 = -1$. For division, we require what is called the conjugate.

The conjugate of a complex number is the same number, just with the sign of the imaginary component flipped.

$$\bar{z} = x - yi$$

where \bar{z} is the conjugate of z .

Similarly to vectors, we can also define the modulus (length) of a complex number

$$|z|^2 = x^2 + y^2 = z \cdot \bar{z}$$

Using this, we can define the division of a complex number and also define the real and imaginary components of a complex number.

The general expression for division is:

$$\frac{u}{z} = \frac{s + it}{x + iy} = \frac{(s + it)(x - iy)}{(x + iy)(x - iy)} = \frac{u\bar{z}}{x^2 + y^2} = \frac{u\bar{z}}{|z|^2}$$

The real and imaginary components can be computed as

$$z = x + iy$$

$$\bar{z} = x - iy$$

$$\Re(z) = \frac{z + \bar{z}}{2}$$

$$\Im(z) = \frac{z - \bar{z}}{2i}$$

Ex1: Simplify $(1 + 2i)(3 + i)(2 - 3i)$

$$\begin{aligned}(1 + 2i)(3 + i)(2 - 3i) &= (3 + i + 6i - 2)(2 - 3i) = (1 + 7i)(2 - 3i) \\ &= 2 - 3i + 14i + 21 \\ &= 23 + 11i\end{aligned}$$

Ex2: Simplify $\left(\frac{2+i}{1+i}\right)^2$

$$\begin{aligned}\left(\frac{2+i}{1+i}\right)^2 &= \left(\frac{(2+i)(1-i)}{(1+i)(1-i)}\right)^2 = \left(\frac{2+i-2i+1}{2}\right)^2 = \left(\frac{3-i}{2}\right)^2 = \frac{9-6i-1}{4} \\ &= 2 - \frac{3}{2}i\end{aligned}$$

Ex3: Simplify $(1 + 2i)^5$

$$\begin{aligned}(1 + 2i)^5 &= 1^5 + 5(1)^4(2i) + 10(1)^3(2i)^2 + 10(1)^2(2i)^3 + 5(1)(2i)^4 + (2i)^5 \\ &= 1 + 10i + 10(-4) + 10(-8i) + 5(16) + 32i \\ &= 1 + 10i - 40 - 80i + 80 + 32i \\ &= 41 - 38i\end{aligned}$$

Ex4: Prove

$$\text{if } |z| = 1 \text{ then } \Re\left(\frac{1}{1+z}\right) = \frac{1}{2}$$

Proof.

$$\begin{aligned}\Re(w) &= \frac{w + \bar{w}}{2} \\ \frac{1}{1+z} &= \frac{1}{1+\bar{z}}\end{aligned}$$

$$\Re\left(\frac{1}{1+z}\right) = \frac{\frac{1}{1+z} + \frac{1}{1+\bar{z}}}{2} = \frac{1+\bar{z}+1+z}{2(1+z)(1+\bar{z})} = \frac{2+z+\bar{z}}{2(1+z+\bar{z}+|z|^2)}$$

$$|z|=1 \Rightarrow \Re\left(\frac{1}{1+z}\right) = \frac{2+z+\bar{z}}{2(2+z+\bar{z})} = \frac{1}{2}$$

□

Some properties of \bar{z} :

- $\overline{\bar{z}} = z$
- $\overline{\bar{z}_1 z_2} = \bar{z}_1 \bar{z}_2$
- $|z_1 z_2| = |z_1| |z_2|$

Some common inequalities come from the triangle inequality:

$$|z_1 + z_2| \leq |z_1| + |z_2|$$

Proof.

$$\begin{aligned} |z_1 + z_2| &\leq |z_1| + |z_2| \\ |z_1 + z_2| &= \sqrt{(x_1 + x_2)^2 + (y_1 + y_2)^2} \\ \sqrt{(x_1 + x_2)^2 + (y_1 + y_2)^2} &\leq \sqrt{x_1^2 + y_1^2} + \sqrt{x_2^2 + y_2^2} \\ |z_1 + z_2| &\leq |z_1| + |z_2| \end{aligned}$$

□

$$\Rightarrow |z_1 \pm z_2| \geq ||z_1| - |z_2||$$

This leads to a general upper and lower bound that can be derived from these inequalities:

$$||z_1| - |z_2|| \leq |z_1 \pm z_2| \leq |z_1| + |z_2|$$

Another way to represent complex numbers is through polar form. To use this we must first introduce Euler's identity:

$$e^{i\varphi} = \cos \varphi + i \sin \varphi$$

This then helps us with the equation for the polar form

$$z = a + ib = |z|(\cos \varphi + i \sin \varphi) = |z|e^{i\varphi}$$

For $z = re^{i\varphi}$ We call r the magnitude of the complex number and φ is the argument. This can be especially useful for simplifying some complex numbers.

Ex:

$$\left| \frac{(1 + \sqrt{3}i)^{100}}{(\sqrt{3} - i)^{100}} \right| = \frac{|1 + \sqrt{3}i|^{100}}{|\sqrt{3} - i|^{100}} = \frac{2^{100}}{2^{100}} = 1$$

Note that because sinusoidal functions are periodic every 2π this means that there infinite ways to express a function in polar coordinates.

$$e^{i2\pi k} = 1, \quad k \in \mathbb{Z} \Rightarrow z = re^{i(\varphi+2\pi k)}$$

To get around the issue of having infinite possible polar forms for every complex number we define what's called the principal argument to be

$$\text{Arg}(z) = \varphi \in (-\pi, \pi]$$

We define the regular argument to be

$$\arg(z) = \text{Arg}(z) + 2k\pi, \quad k \in \mathbb{Z}$$

Note that $\text{Arg}(z)$ is singular valued while $\arg(z)$ is multi-valued.

We can define $\text{Arg}(z)$ in terms of the real and imaginary parts (x and y) as

$$\text{Arg}(z) = \arctan\left(\frac{y}{x}\right) \pm k\pi$$

What it is specifically depends on what quadrant of the complex plane the point lies in.

- QI: $\text{Arg}(z) = \arctan\left(\frac{y}{x}\right)$
- QII: $\text{Arg}(z) = \arctan\left(\frac{y}{x}\right) + \pi$
- QIII: $\text{Arg}(z) = \arctan\left(\frac{y}{x}\right) - \pi$
- QIV: $\text{Arg}(z) = \arctan\left(\frac{y}{x}\right)$

Ex:

$$\text{Arg}(-1 - \sqrt{3}i) = \arctan(\sqrt{3}) - \pi = -\frac{2\pi}{3}$$

Ex2:

$$\begin{aligned} \arg(1 - \sqrt{3}i) \\ \text{Arg}(1 - \sqrt{3}i) &= -\frac{\pi}{3} \\ \arg(1 - \sqrt{3}i) &= -\frac{\pi}{3} + 2k\pi, \quad k \in \mathbb{Z} \end{aligned}$$

Ex3:

$$\begin{aligned} \arg(-1 + 2i) \\ \text{Arg}(-1 + 2i) &= \arctan(-2) + \pi \\ \arg(-1 + 2i) &= \pi - \arctan(2) + 2k\pi \end{aligned}$$

Ex4: Simplify

$$z = -3 + 3i$$

$$\text{Arg}(z) = \arctan\left(\frac{3}{-3}\right) + \pi = \frac{3\pi}{4}$$

$$|z| = 3\sqrt{2}$$

$$z = 3\sqrt{2}e^{\frac{3\pi}{4}i}$$

Ex5: Simplify

$$z = -3 - 3i$$

$$\text{Arg}(z) = \arctan\left(\frac{-3}{-3}\right) - \pi = -\frac{3\pi}{4}$$

$$|z| = 3\sqrt{2}$$

$$z = 3\sqrt{2}e^{-\frac{3\pi}{4}i}$$

Ex6: Simplify

$$z = \frac{1-i}{-\sqrt{3}+i} = \frac{u}{v}$$

$$\arg(z) = \arg(u) - \arg(v) = \arctan\left(\frac{-1}{1}\right) - \left(\arctan\left(\frac{1}{-\sqrt{3}}\right) + \pi\right) + 2k\pi$$

$$= -\frac{\pi}{4} - \frac{5\pi}{6} + 2k\pi = -\frac{13\pi}{12} + 2k\pi$$

$$\text{Arg}(z) = \frac{11\pi}{12}$$

$$|z| = \frac{|u|}{|v|} = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$$

$$z = \frac{e^{\frac{11\pi}{12}i}}{\sqrt{2}}$$

Ex7: Simplify

$$z = (\sqrt{3} - i)^2 = w^2$$

$$\arg(w) = \arctan\left(\frac{-1}{\sqrt{3}}\right) + 2k\pi = -\frac{\pi}{6}$$

$$\Rightarrow \arg(z) = 2\arg(w) = -\frac{\pi}{3} + 4k\pi$$

$$\text{Arg}(z) = -\frac{\pi}{3}$$

$$|w| = 2 \Rightarrow |z| = |w|^2 = 4$$

$$z = 4e^{-\frac{\pi}{3}i}$$

Properties of $\text{Arg}(z)$ and $\arg(z)$

- $\text{Arg}(z_1 z_2) \neq \text{Arg}(z_1) + \text{Arg}(z_2)$

Proof. Proof by contradiction: assume that $\text{Arg}(z_1 z_2) = \text{Arg}(z_1) + \text{Arg}(z_2)$ is true.

Take $z_1 = z_2 = -1$.

$$\text{Arg}(z_1) = \text{Arg}(z_2) = \pi$$

$$\begin{aligned}\Rightarrow \text{Arg}(z_1) + \text{Arg}(z_2) &= 2\pi \\ \text{Arg}(z_1 z_2) &= \text{Arg}(1) = 0 \\ \text{Arg}(z_1 z_2) &\neq \text{Arg}(z_1) + \text{Arg}(z_2) \quad \forall z_1, z_2 \neq 0 \in \mathbb{C}\end{aligned}$$

□

- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$
- $\text{Arg}(\bar{z}) \neq -\text{Arg}(z)$

Proof. Proof by contradiction: assume that $\text{Arg}(\bar{z}) = -\text{Arg}(z)$ is true.
Take $z = -1$

$$\begin{aligned}\bar{z} &= z = -1 \\ \text{Arg}(z) &= \pi \\ \text{Arg}(\bar{z}) &= \pi \\ -\text{Arg}(z) &= -\pi \\ \Rightarrow \text{Arg}(\bar{z}) &\neq -\text{Arg}(z) \quad \forall z \in \mathbb{C}\end{aligned}$$

□

- $\arg(z) = -\arg(\bar{z})$

Proof.

$$\begin{aligned}z &= |z|e^{i\arg(z)} \quad \forall z \in \mathbb{C} \\ \bar{z} &= |z|e^{-i\arg(z)} \\ \Rightarrow \arg(\bar{z}) &= -\arg(z)\end{aligned}$$

□

1.1.2 De Moirre's Formula

Using Euler's identity we can derive a powerful formula called De Moirre's Formula as follows:

$$\begin{aligned}e^{iN\varphi} &= \cos(N\varphi) + i\sin(N\varphi) \\ e^{iN\varphi} &= (e^{i\varphi})^N = (\cos \varphi + i\sin \varphi)^N \\ (\cos \varphi + i\sin \varphi)^N &= \cos(N\varphi) + i\sin(N\varphi)\end{aligned}$$

Applications of De Moirre's Formula:

Binomial expansion:

$$\begin{aligned}N &= 2 : \\ \cos(2\theta) &= \cos^2 \theta - \sin^2 \theta \\ \sin(2\theta) &= 2\cos \theta \sin \theta\end{aligned}$$

$N = 3 :$

$$(\cos \theta + i \sin \theta)^3 = \cos^3 \theta + 3 \cos^2 \theta (i \sin \theta) + 3 \cos \theta (i \sin \theta)^2 + (i \sin \theta)^3$$

$$\cos(3\theta) = \cos^3 \theta - 3 \cos \theta \sin^2 \theta$$

$$\sin(3\theta) = 3 \cos^2 \theta \sin \theta - \sin^3 \theta$$

Ex: Prove

$$\sin(3\theta) = 3 \sin \theta - 4 \sin^3 \theta$$

Proof. Using De Moivre's formula with $N = 3$

$$(\cos \theta + i \sin \theta)^3 = \cos(3\theta) + i \sin(3\theta)$$

$$(\cos \theta)^3 + 3(\cos \theta)^2(i \sin \theta) + 3 \cos \theta (i \sin \theta)^2 + (i \sin \theta)^3 = \cos(3\theta) + i \sin(3\theta)$$

$$\cos^3 \theta - 3 \cos \theta \sin^2 \theta + i(3 \cos^2 \theta \sin \theta - \sin^3 \theta) = \cos(3\theta) + i \sin(3\theta)$$

$$\Im \{ \cos^3 \theta - 3 \cos \theta \sin^2 \theta + i(3 \cos^2 \theta \sin \theta - \sin^3 \theta) \} = \Im \{ \cos(3\theta) + i \sin(3\theta) \}$$

$$3 \cos^2 \theta \sin \theta - \sin^3 \theta = \sin(3\theta)$$

$$\sin^2 \theta + \cos^2 \theta = 1 \Rightarrow \cos^2 \theta = 1 - \sin^2 \theta$$

$$3(1 - \sin^2 \theta) \sin \theta - \sin^3 \theta = \sin(3\theta)$$

$$3 \sin \theta - 3 \sin^3 \theta - \sin^3 \theta = \sin(3\theta)$$

$$3 \sin \theta - 4 \sin^3 \theta = \sin(3\theta)$$

□

Computing trigonometric integrals:

Ex:

$$\int_0^{2\pi} \cos^8 \varphi d\varphi$$

$$e^{i\varphi} = \cos \varphi + i \sin \varphi$$

$$e^{-i\varphi} = \cos \varphi - i \sin \varphi$$

$$\cos \varphi = \frac{e^{i\varphi} + e^{-i\varphi}}{2}$$

$$\sin \varphi = \frac{e^{i\varphi} - e^{-i\varphi}}{2i}$$

$$\int_0^{2\pi} \left(\frac{e^{i\varphi} + e^{-i\varphi}}{2} \right)^8 d\varphi = \frac{1}{2^8} \int_0^{2\pi} (e^{i\varphi} + e^{-i\varphi})^8 d\varphi$$

$$= \frac{1}{2^8} \int_0^{2\pi} (e^{i8\varphi} + {}_1C_8 e^{i7\varphi} e^{-i\varphi} + \dots + {}_7C_8 e^{i\varphi} e^{-i7\varphi} + e^{-i8\varphi}) d\varphi$$

$$= \frac{1}{2^8} (0 + \dots + {}_4C_8 2\pi + \dots + 0) = \frac{{}_4C_8}{2^7} \pi$$

Ex2:

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}$$

$$\begin{aligned}
\int_0^{2\pi} \cos^6 \theta d\theta &= \int_0^{2\pi} \left(\frac{e^{i\theta} + e^{-i\theta}}{2} \right) \\
&= \frac{1}{2^6} \int_0^{2\pi} \sum_{k=0}^6 \binom{6}{k} e^{ik\theta} e^{-i\theta(6-k)} d\theta \\
&= \frac{1}{2^6} \sum_{k=0}^6 \binom{6}{k} \int_0^{2\pi} e^{i\theta(2k-6)} d\theta \\
\int_0^{2\pi} e^{ik\theta} d\theta &= \frac{e^{ik\theta}}{ik} \Big|_0^{2\pi} = \frac{e^{2\pi ik} - 1}{ik} \\
e^{2\pi ik} &= 1, \quad k \in \mathbb{Z} \Rightarrow \int_0^{2\pi} e^{ik\theta} d\theta = 0, \quad k \neq 0 \in \mathbb{Z} \\
\Rightarrow \int_0^{2\pi} \cos^6 \theta d\theta &= \frac{1}{2^6} \binom{6}{3} \int_0^{2\pi} d\theta = \frac{(20)(2\pi)}{2^6} = \frac{5\pi}{8}
\end{aligned}$$

Ex3:

$$\begin{aligned}
&\int_0^{2\pi} \sin^6(2\theta) d\theta \\
\sin(2\theta) &= \frac{e^{2i\theta} - e^{-2i\theta}}{2i} \\
\int_0^{2\pi} \sin^6(2\theta) d\theta &= \int_0^{2\pi} \left(\frac{e^{2i\theta} - e^{-2i\theta}}{2i} \right)^6 d\theta \\
&= \frac{1}{(2i)^6} \int_0^{2\pi} \sum_{k=0}^6 \binom{6}{k} (-1)^{6-k} e^{2ik\theta} e^{-2i\theta(6-k)} d\theta \\
&= -\frac{1}{2^6} \sum_{k=0}^6 \binom{6}{k} (-1)^{6-k} \int_0^{2\pi} e^{i\theta(4k-12)} d\theta \\
\int_0^{2\pi} e^{ik\theta} d\theta &= \frac{e^{ik\theta}}{ik} \Big|_0^{2\pi} = \frac{e^{2\pi ik} - 1}{ik} \\
e^{2\pi ik} &= 1, \quad k \in \mathbb{Z} \Rightarrow \int_0^{2\pi} e^{ik\theta} d\theta = 0, \quad k \neq 0 \in \mathbb{Z} \\
\Rightarrow \int_0^{2\pi} \sin^6(2\theta) d\theta &= -\frac{1}{2^6} \binom{6}{3} (-1)^3 \int_0^{2\pi} d\theta \\
&= \frac{5\pi}{8}
\end{aligned}$$

Ex4: Prove

$$\sum_{k=0}^n \cos(k\theta) = \frac{1}{2} + \frac{\sin\left(\left(n + \frac{1}{2}\right)\theta\right)}{2\sin\left(\frac{\theta}{2}\right)}$$

Proof. De Moivre's formula states

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta)$$

If we take the conjugate of both sides we get

$$(\cos \theta - i \sin \theta)^n = \cos(n\theta) - i \sin(n\theta)$$

Summing these two equations gives

$$\begin{aligned} (\cos \theta + i \sin \theta)^k + (\cos \theta - i \sin \theta)^k &= 2 \cos(k\theta) \\ \cos \theta \pm i \sin \theta &= e^{\pm i\theta} \\ 2 \cos(k\theta) &= (e^{i\theta})^k + (e^{-i\theta})^k \end{aligned}$$

We can sum both sides of this to get

$$2 \sum_{k=0}^n \cos(k\theta) = \sum_{k=0}^n (e^{i\theta})^k + \sum_{k=0}^n (e^{-i\theta})^k$$

The formula for the geometric sum is

$$\sum_{k=0}^n z^k = \frac{1 - z^{n+1}}{1 - z}$$

Applying this we get

$$\begin{aligned} 2 \sum_{k=0}^n \cos(k\theta) &= \frac{1 - (e^{i\theta})^{n+1}}{1 - e^{i\theta}} + \frac{1 - (e^{-i\theta})^{n+1}}{1 - e^{-i\theta}} \\ 2 \sum_{k=0}^n \cos(k\theta) &= \frac{(1 - e^{i\theta(n+1)})(1 - e^{-i\theta}) + (1 - e^{-i\theta(n+1)})(1 - e^{i\theta})}{(1 - e^{i\theta})(1 - e^{-i\theta})} \\ 2 \sum_{k=0}^n \cos(k\theta) &= \frac{1 - e^{i\theta(n+1)} - e^{i\theta} + e^{i\theta n} + 1 - e^{-i\theta(n+1)} - e^{-i\theta} + e^{-i\theta n}}{1 - e^{i\theta} - e^{-i\theta} + 1} \\ 2 \sum_{k=0}^n \cos(k\theta) &= 1 + \frac{e^{i\theta n} + e^{-i\theta n} - e^{i\theta(n+1)} - e^{-i\theta(n+1)}}{2 - e^{i\theta} - e^{-i\theta}} \\ 2 \sum_{k=0}^n \cos(k\theta) &= 1 + \frac{e^{i\frac{\theta}{2}} \left(e^{-i\theta(n+\frac{1}{2})} - e^{i\theta(n+\frac{1}{2})} \right) + e^{-i\frac{\theta}{2}} \left(e^{i\theta(n+\frac{1}{2})} - e^{-i\theta(n+\frac{1}{2})} \right)}{e^{i\frac{\theta}{2}} \left(e^{-i\frac{\theta}{2}} - e^{i\frac{\theta}{2}} \right) + e^{-i\frac{\theta}{2}} \left(e^{i\frac{\theta}{2}} - e^{-i\frac{\theta}{2}} \right)} \\ 2 \sum_{k=0}^n \cos(k\theta) &= 1 + \frac{\left(e^{-i\frac{\theta}{2}} - e^{i\frac{\theta}{2}} \right) \left(e^{i\theta(n+\frac{1}{2})} - e^{-i\theta(n+\frac{1}{2})} \right)}{- \left(e^{-i\frac{\theta}{2}} - e^{i\frac{\theta}{2}} \right)^2} \\ 2 \sum_{k=0}^n \cos(k\theta) &= 1 + \frac{e^{i\theta(n+\frac{1}{2})} - e^{-i\theta(n+\frac{1}{2})}}{e^{i\frac{\theta}{2}} - e^{-i\frac{\theta}{2}}} \\ 2 \sum_{k=0}^n \cos(k\theta) &= 1 + \frac{\frac{1}{2i} \left(e^{i\theta(n+\frac{1}{2})} - e^{-i\theta(n+\frac{1}{2})} \right)}{\frac{1}{2i} \left(e^{i\frac{\theta}{2}} - e^{-i\frac{\theta}{2}} \right)} \\ \sin(x) &= \frac{e^{ix} - e^{-ix}}{2i} \end{aligned}$$

$$2 \sum_{k=0}^n \cos(k\theta) = 1 + \frac{\sin(\theta(n + \frac{1}{2}))}{\sin(\frac{\theta}{2})}$$

$$\sum_{k=0}^n \cos(k\theta) = \frac{1}{2} + \frac{\sin((n + \frac{1}{2})\theta)}{2 \sin(\frac{\theta}{2})}$$

□

1.1.3 Geometry in the Complex Plane

Using the notation such that $z = x + iy$ where $\Re(z) = x$ and $\Im(z) = y$ we can define a circle in the complex plane as

$$(x - x_0)^2 + (y - y_0)^2 = r_0^2$$

This is analogous to writing

$$|z - z_0| = r_0$$

The two can be related as follows:

$$|z - z_0| = r_0$$

$$|x + iy - x_0 - iy_0| = r_0$$

$$\sqrt{(x - x_0)^2 + (y - y_0)^2} = r_0$$

$$(x - x_0)^2 + (y - y_0)^2 = r_0^2$$

Ex: describe the circle formed by $2|z| = |z + 1|$

$$2|z| = |z + 1|$$

$$4|z|^2 = |z + 1|^2$$

$$4x^2 + 4y^2 = (x + 1)^2 + y^2$$

$$4x^2 + 4y^2 = x^2 + 2x + 1 + y^2$$

$$3x^2 + 3y^2 - 2x - 1 = 0$$

$$3x^2 - 2x + \frac{1}{3} + 3y^2 - \frac{4}{3} = 0$$

$$3\left(x - \frac{1}{3}\right)^2 + 3y^2 = \frac{4}{3}$$

$$\left(x - \frac{1}{3}\right)^2 + y^2 = \frac{4}{9}$$

A line in the complex plane can be written as

$$ax + by = c \longleftrightarrow a \frac{z + \bar{z}}{2} + b \frac{z - \bar{z}}{2i} = c$$

Ex: describe the line formed by $|z - 1 + i| = |z - 2i|$

$$|z - 1 + i| = |z - 2i|$$

$$\begin{aligned}
|z - 1 + i|^2 &= |z - 2i|^2 \\
(x - 1)^2 + (y + 1)^2 &= x^2 + (y - 2)^2 \\
x^2 - 2x + 1 + y^2 + 2y + 1 &= x^2 + y^2 - 4y + 4 \\
-2x + 2 + 2y &= -4y + 4 \\
6y &= 2x + 2 \\
y &= \frac{1}{3}(x + 1)
\end{aligned}$$

We can define an ellipse in the complex plane as

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

If we have $a > b$ then we will have a horizontal ellipse and if $b > a$ then we will have a vertical ellipse.

Assuming that $a > b$ then we can define the foci points to be at

$$\begin{aligned}
+F &= (\sqrt{a^2 - b^2}, 0) \\
-F &= (-\sqrt{a^2 - b^2}, 0)
\end{aligned}$$

The equation of an ellipse can also be described by

$$|z - F| + |z + F| = 2a$$

Ex: describe the ellipse formed by $|z - 1| + |z + 1| = 4$

$$\begin{aligned}
|z - 1| + |z + 1| &= 4 \\
|z - F| + |z + F| &= 2a \Rightarrow 2a = 4 \Rightarrow a = 2 \\
F = \sqrt{a^2 - b^2} &= 1 \Rightarrow 1 = 4 - b^2 \Rightarrow b^2 = 3 \\
\frac{x^2}{4} + \frac{y^2}{3} &= 1
\end{aligned}$$

Ex2: describe the ellipse formed by $|z - 1| + |z + 3| = 6$

$$\begin{aligned}
|z - 1| + |z + 3| &= 6 \\
\text{note that this ellipse is not centered at the origin so we need to shift it} \\
|z + 1 - 2| + |z + 1 + 2| &= 6 \\
2a = 6 &\Rightarrow a = 3 \\
F = 2 = \sqrt{a^2 - b^2} &\Rightarrow 4 = 9 - b^2 \Rightarrow b^2 = 5 \\
\frac{(x + 1)^2}{9} + \frac{y^2}{5} &= 1
\end{aligned}$$

1.1.4 Roots of a Complex Number

Given $z_0 = r_0 e^{i\varphi_0}$, what is $z_0^{\frac{1}{n}}$?

If we let $w = z_0^{\frac{1}{n}}$ then $w^n = z_0$

$$w = r e^{i\varphi}, \quad w^n = r^n e^{in\varphi}$$

$$\begin{aligned}
w^n &= z_0 \Rightarrow r^n e^{in\varphi} = r_0 e^{i\varphi_0} \\
&\Rightarrow r^n = r_0 \Rightarrow r_0^{\frac{1}{n}} \\
e^{in\varphi} &= e^{i\varphi_0} \Rightarrow n\varphi = \varphi_0 + 2k\pi \\
\varphi &= \frac{\varphi_0}{n} + \frac{2k\pi}{n}
\end{aligned}$$

So all solutions to $w^n = z_0$ are given by

$$w = r_0^{\frac{1}{n}} e^{i(\frac{\varphi_0}{n} + \frac{2k\pi}{n})}, \quad k \in \mathbb{Z}$$

If we normalize $\varphi_0 = \text{Arg}(z_0)$ then k will be in the range $k \in \{0, 1, \dots, n-1\}$.
Note that the expression

$$w = r_0^{\frac{1}{n}} e^{i(\frac{\varphi_0}{n} + \frac{2k\pi}{n})}, \quad k \in \mathbb{Z}$$

is multi-valued. If we want to avoid this, we can take what's called the principal value which is the value when $k = 0$:

$$z_0^{\frac{1}{n}} = r_0^{\frac{1}{n}} e^{i\frac{\varphi_0}{n}}$$

Ex: Compute $(-1)^{\frac{1}{2}}$

$$z_0 = -1 = 1^{\frac{1}{2}} e^{i(\frac{\pi}{2} + \frac{2k\pi}{2})} = e^{i(\frac{\pi}{2} + k\pi)} = \left\{ \dots, e^{-i\frac{3\pi}{2}}, e^{-i\frac{\pi}{2}}, e^{i\frac{\pi}{2}}, e^{i\frac{3\pi}{2}}, e^{i\frac{5\pi}{2}}, \dots \right\}$$

note that there are only 2 unique values

$$(-1)^{\frac{1}{2}} = e^{i(\frac{\pi}{2} + k\pi)}, \quad k \in \{0, 1\}$$

The principal value (when $k = 0$) of this equation works out to be i .

Ex2: Find all solutions to

$$\begin{aligned}
z^7 &= i - 1 \\
z^7 &= \sqrt{2} e^{i(\frac{3\pi}{4} + 2\pi k)} \\
z &= 2^{1/14} e^{i(\frac{3\pi}{28} + \frac{2\pi}{7}k)}, \quad k \in \{0, 1, 2, 3, 4, 5, 6\}
\end{aligned}$$

Ex3: Find all solutions to

$$\begin{aligned}
z^5 &= \frac{2i}{-1 - \sqrt{3}i} \\
z^5 &= \frac{2e^{i\frac{\pi}{2}}}{2e^{-i\frac{2\pi}{3}}} = e^{i\frac{7\pi}{6}} = e^{-i(\frac{5\pi}{6} + 2\pi k)} \\
z &= e^{-i(\frac{\pi}{6} + \frac{2\pi}{5}k)}, \quad k \in \{0, 1, 2, 3, 4\}
\end{aligned}$$

Ex4: Find all solutions to

$$\left(\frac{z}{z+1} \right)^2 = i$$

$$\begin{aligned} \left(\frac{z}{z+1}\right)^2 &= e^{i(\frac{\pi}{2}+2\pi k)} \\ \frac{z}{z+1} &= e^{i(\frac{\pi}{4}+\pi k)}, \quad k \in \{0, 1\} \\ e^{i(\frac{\pi}{4}+\pi k)} &= \left\{ \frac{1}{\sqrt{2}}(1+i), \frac{1}{\sqrt{2}}(-1-i) \right\} \\ z &= e^{i(\frac{\pi}{4}+k\pi)}(z+1) \\ z &= \frac{e^{i(\frac{\pi}{4}+k\pi)}}{1 - e^{i(\frac{\pi}{4}+k\pi)}} = \left\{ \frac{1+i}{\sqrt{2}-1-i}, \frac{-1-i}{\sqrt{2}+1+i} \right\} \end{aligned}$$

Ex5: Find all solutions to

$$\begin{aligned} z^2 + 4iz + 1 &= 0 \\ (z^2 + 4iz - 4) + 4 + 1 &= 0 \\ (z + 2i)^2 + 5 &= 0 \\ (z + 2i)^2 &= -5 = 5e^{i(\pi+2\pi k)} \\ z + 2i &= \sqrt{5}e^{i(\frac{\pi}{2}+\pi k)} = \left\{ \sqrt{5}i, -\sqrt{5}i \right\} \\ z &= \left\{ (\sqrt{5}-2)i, -(\sqrt{5}+2)i \right\} \end{aligned}$$

1.2 Complex Functions

1.2.1 Mapping Properties of Simple Functions

Similar to how functions with real variables map values to a different set of values, complex functions do the same. The main difference is that with complex functions we're mapping a 2 dimensional set of inputs to a 2 dimensional set of outputs.

$$w = f(z) = u + iv$$

We define $z \in \mathcal{S}$ the image of \mathcal{S} under w .

Some common mappings:

- The identity map

$$\begin{aligned} w = f(z) &= z \\ \begin{cases} u = x \\ v = y \end{cases} \end{aligned}$$

- Translation by z_0

$$\begin{aligned} w = f(z) &= z + z_0 \\ \begin{cases} u = x + x_0 \\ v = y + y_0 \end{cases} \end{aligned}$$

- Stretching ($a > 1$) or contraction ($a < 1$)

$$w = f(z) = az = are^{i\varphi}, \quad a \in \mathbb{R}$$


$$\begin{cases} u = ax \\ v = ay \end{cases}$$

- Rotation by φ_0

$$w = f(z) = e^{i\varphi_0} z = e^{i(\varphi+\varphi_0)}$$

Using these basic mapping principles we are able to lay the foundation for some more complicated mappings.

Ex: Find the image of $S = \{|z - 1| \geq 1\}$ under the mapping $f(z) = \frac{1}{z}$



$$z = \frac{1}{w} = \frac{1}{u + iv} = \frac{u - iv}{u^2 + v^2}$$

$$x = \frac{u}{u^2 + v^2}$$

$$y = -\frac{v}{u^2 + v^2}$$

$$|z - 1| \geq 1 \Rightarrow \left| \frac{1}{w} - 1 \right| \geq 1$$

$$\left| \frac{1 - w}{w} \right| \geq 1$$

$$|1 - w| \geq |w|$$


$$|1 - w|^2 \geq |w|^2$$

$$(1 - u)^2 + v^2 \geq u^2 + v^2$$


$$-2u + 1 \geq 0$$

$$u \leq \frac{1}{2}$$

$$S' = \left\{ u \leq \frac{1}{2} \right\}$$



Ex2: Find the image of $S = \{x \leq 1\}$ under the mapping $f(z) = \frac{1}{z}$




$$z = \frac{1}{w} = \frac{1}{u + iv} = \frac{u - iv}{u^2 + v^2}$$

$$x = \frac{u}{u^2 + v^2}$$

$$x \leq 1 \Rightarrow \frac{u}{u^2 + v^2} \leq 1 \Rightarrow u^2 + v^2 \geq u$$

$$(u - \frac{1}{2})^2 + v^2 \geq \frac{1}{4}$$



We see from the previous two examples that circles map to lines and lines map to circles. Let's see why this is the case.

$$a(x^2 + y^2) + bx + cy + d = 0$$

$$a|z|^2 + b\frac{z + \bar{z}}{2} + c\frac{z - \bar{z}}{2i} + d = 0$$

In the case where $a = 0$, we have a line. In the case where $a \neq 0$, we have a circle.

$$w = \frac{1}{z} \Rightarrow z = \frac{1}{w}$$

$$\begin{aligned}
z\bar{z} &= |z|^2 = \frac{1}{|w|^2} = \frac{1}{w\bar{w}} \\
a\frac{1}{w\bar{w}} + b\frac{z+\bar{z}}{2} + c\frac{z-\bar{z}}{2i} + d &= 0 \\
a\frac{1}{w\bar{w}} + \frac{b}{2}\left(\frac{1}{w} + \frac{1}{\bar{w}}\right) + \frac{c}{2i}\left(\frac{1}{w} - \frac{1}{\bar{w}}\right) + d &= 0 \\
a + \frac{b}{2}(w + \bar{w}) + \frac{c}{2i}(w - \bar{w}) + d(w\bar{w}) &= 0
\end{aligned}$$

If we have a linear transformation of the form $az + b$ it corresponds to the scaling and translation of the set only. A line will map to a line and a circle will map to a circle.

We can combine this with the $w = \frac{1}{z}$ transformation property to get a more general transformation. We call this the *Mobius transformation*:

$$f(z) = \frac{az + b}{cz + d} = \frac{a}{c} + \frac{b - \frac{ad}{c}}{cz + d}$$

where $a, b, c, d \in \mathbb{C}$

Ex: Find the mapping of $f(z) = \frac{1}{z+1}$ on the set $S = \{\Re(z) > 0\}$

$$\begin{aligned}
u + iv &= \frac{1}{x + 1 + iy} \Rightarrow x + 1 + iy = \frac{1}{u + iv} \\
x + 1 &= \frac{u}{u^2 + v^2} \\
x > 0 &\Rightarrow x + 1 > 1 \\
\frac{u}{u^2 + v^2} &> 1 \Rightarrow u > u^2 + v^2 \\
u^2 + v^2 - u + \frac{1}{4} &< \frac{1}{4} \\
\left(u - \frac{1}{2}\right)^2 + v^2 &< \frac{1}{4} \\
S' &= \left\{ w = u + iv \mid \left(u - \frac{1}{2}\right)^2 + v^2 < \left(\frac{1}{2}\right)^2 \right\}
\end{aligned}$$

Ex2: Find the mapping of $f(z) = \frac{z-i}{z+i}$ on $S = \{|z| < 3\}$

$$\begin{aligned}
wz + iw &= z - i \Rightarrow z(w - 1) = -i - iw \Rightarrow z = \frac{i(w + 1)}{1 - w} \\
|z| &= \frac{|w + 1|}{|w - 1|} < 3 \\
|w + 1| &< 3|w - 1| \Rightarrow |w + 1|^2 < 9|w - 1|^2 \\
(u + 1)^2 + v^2 &< 9(u - 1)^2 + 9v^2 \\
u^2 + 2u + 1 + v^2 &< 9u^2 - 18u + 9 + 9v^2 \\
0 &< 8u^2 - 20u + 8 + 8v^2 \Rightarrow 0 < u^2 - \frac{5}{2}u + 1 + v^2
\end{aligned}$$

$$\frac{9}{16} < u^2 - \frac{5}{2}u + \frac{25}{16} + v^2$$

$$\frac{9}{16} < \left(u - \frac{5}{4}\right)^2 + v^2$$

$$S' = \left\{ w = u + iv \mid \left(u - \frac{5}{4}\right)^2 + v^2 > \left(\frac{3}{4}\right)^2 \right\}$$

Another common mapping is the $f(z) = z^2$ or more generally $f(z) = z^n$ mapping. For $w = z^2$,

$$w = z^2 = r^2 e^{2i\varphi} \Rightarrow \begin{cases} |w| = |z|^2 \\ \arg(w) = 2 \arg(z) \end{cases}$$

This mapping scales the magnitude but more notably, it doubles the argument. This means that the mapping of a half circle will now be a full circle.

Ex: Find the mapping of $f(z) = z^2$ on $S = \{0 \leq \Re(z) \leq 1, \Im(z) = 1\}$

$$w = x^2 + i2xy - y^2$$

$$u = x^2 - y^2 = x^2 - 1 \Rightarrow -1 \leq u \leq 0$$

$$v = 2xy = 2x \Rightarrow 0 \leq v \leq 2$$

$$S' = \{w = u + iv \mid -1 \leq u \leq 0, 0 \leq v \leq 2\}$$

Ex2: Find the mapping of $f(z) = -2z^5$ on $S = \{|z| < 1, 0 < \text{Arg}(z) < \frac{\pi}{2}\}$

$$z^5 = -\frac{w}{2} \Rightarrow |z|^5 = \frac{|w|}{2} < 1 \Rightarrow |w| < 2$$

$$5 \arg(z) = \arg(w) \pm \pi$$

$$0 < \arg(w) \pm \pi < \frac{5\pi}{2}$$


$$-\pi < \arg(w) < \frac{3\pi}{2}$$

$$S' = \{|w| < 2\}$$

Another common mapping is the $f(z) = e^z$ mapping.

$$w = e^z = e^{x+iy} = e^x e^{iy}$$

$$\begin{cases} |w| = e^x \\ \arg(w) = y \end{cases}$$



This mapping has the property that the magnitude is only dependent on x and the argument is exactly y .

Ex: Find the mapping of $f(z) = e^z$ on $S = \{\Re(z) = 1\}$

$$w = e^x e^{iy}$$

$$|w| = e, \arg(w) = y$$

$$S' = \{|w| = e\}$$

Ex2: Find the mapping of $f(z) = e^z$ on $S = \{0 \leq \Im(z) \leq \frac{\pi}{4}\}$

$$|w| = x$$

$$\arg(w) = y \Rightarrow 0 \leq \arg(w) \leq \frac{\pi}{4}$$

$$S' = \left\{0 \leq \text{Arg}(w) \leq \frac{\pi}{4}\right\}$$

Ex3: Find the mapping of $f(z) = e^{iz}$ on $S = \{z : -\frac{\pi}{2} \leq \Re(z) \leq \pi, -1 \leq \Im(z) \leq 1\}$
(Note that multiplying z by i rotates it by 90°)

$$w = e^{iz} = e^{ix} e^{-y}$$

$$|w| = e^{-y} \Rightarrow e^{-1} \leq |w| \leq e$$

$$\arg(w) = x \Rightarrow -\frac{\pi}{2} \leq \arg(w) \leq \pi$$

$$S' = \left\{w \mid e^{-1} \leq |w| \leq e, -\frac{\pi}{2} \leq \arg(w) \leq \pi\right\}$$

Ex4: Prove

$$|e^{-z^3}| \leq 1 \quad \forall \left\{z \mid -\frac{\pi}{6} \leq \text{Arg}(z) \leq \frac{\pi}{6}\right\}$$

Proof. We can express $-z^3$ as some complex number $a + ib$ where $a = \Re(-z^3)$ and $b = \Im(-z^3)$. Taking the magnitude gives

$$|e^{-z^3}| = |e^{a+ib}| = |e^a e^{ib}| = |e^a| |e^{ib}| = |e^a| = |e^{\Re(-z^3)}|$$

z can be written as

$$z = |z| e^{i \text{Arg}(z)}$$

$$z^3 = |z|^3 e^{i3 \text{Arg}(z)} = |z|^3 (\cos(3 \text{Arg}(z)) + i \sin(3 \text{Arg}(z)))$$

$$-z^3 = -|z|^3 (\cos(3 \text{Arg}(z)) + i \sin(3 \text{Arg}(z)))$$

$$\Re(-z^3) = -|z|^3 \cos(3 \text{Arg}(z))$$

$$\text{Arg}(z) \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right]$$

$$\Rightarrow 3 \text{Arg}(z) \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

$$\cos(3 \text{Arg}(z)) \in [0, 1]$$

$$\begin{aligned}
|z|^3 &\in \{x \in \mathbb{R} | x \geq 0\} \\
\Re(-z^3) &= -|z|^3 \cos(3 \operatorname{Arg}(z)) \in \{x \in \mathbb{R} | x \leq 0\} \\
e^{\Re(-z^3)} &\in [0, 1] \\
\Rightarrow |e^{-z^3}| &\leq 1
\end{aligned}$$

□

1.2.2 Calculus of Complex Functions

We define the limit of a complex function to be

$$\begin{aligned}
w &= f(z) = u + iv \\
\lim_{z \rightarrow z_0} f(z) &= \lim_{(x,y) \rightarrow (x_0,y_0)} u(x,y) + i \lim_{(x,y) \rightarrow (x_0,y_0)} v(x,y)
\end{aligned}$$

Note that the notation $(x, y) \rightarrow (x_0, y_0)$ means that the limit is taken as (x, y) approaches (x_0, y_0) along *any* path.

The usual limit arithmetic rules are able to be applied as with real numbers.

In order for $\lim_{z \rightarrow z_0} f(z)$ to exist, we require that both $\lim_{(x,y) \rightarrow (x_0,y_0)} u(x,y)$ and $\lim_{(x,y) \rightarrow (x_0,y_0)} v(x,y)$ exist.

If we define $z_0 = x_0 + iy_0$ then we can define the derivative of a complex function as

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

If this limit exists then the function is said to be differentiable at z_0 .

Ex: $f(z) = z$

$$\begin{aligned}
f'(z_0) &= \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} = \frac{z_0 + \Delta z - z_0}{\Delta z} = 1 \\
\Rightarrow f'(z_0) &= 1
\end{aligned}$$

Ex2: $f(z) = \bar{z}$

$$\begin{aligned}
f'(z) &= \lim_{\Delta z \rightarrow 0} \frac{f(z + \Delta z) - f(z)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{\overline{\Delta z}}{\Delta z} \\
\Delta z &= h_1 + ih_2 \Rightarrow \overline{\Delta z} = h_1 - ih_2 \\
h_2 = 0 : \lim_{\Delta z \rightarrow 0} \frac{\overline{\Delta z}}{\Delta z} &= \lim_{h_1 \rightarrow 0} \frac{h_1}{h_1} = 1 \\
h_1 = 0 : \lim_{\Delta z \rightarrow 0} \frac{\overline{\Delta z}}{\Delta z} &= \lim_{h_2 \rightarrow 0} \frac{-ih_2}{ih_2} = -1 \\
\lim_{h_1 \rightarrow 0} \frac{h_1}{h_1} &\neq \lim_{h_2 \rightarrow 0} \frac{-ih_2}{ih_2} \therefore \text{the derivative does not exist}
\end{aligned}$$

An easy way to determine if a function is differentiable is to use the Cauchy-Riemann equations.

Any path that can be taken to approach z_0 can be written as a linear combination of the paths $\Delta z = \Delta x$ and $\Delta z = i\Delta y$ so the derivative must satisfy both of these paths.

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

$$f(z_0) = u(x_0, y_0) + iv(x_0, y_0)$$

Define $\Delta z = \Delta x$

$$f'(z_0) = \lim_{\Delta x \rightarrow 0} \frac{u(x_0 + \Delta x, y_0) + iv(x_0 + \Delta x, y_0) - u(x_0, y_0) - iv(x_0, y_0)}{\Delta x}$$

$$f'(z_0) = \lim_{\Delta x \rightarrow 0} \frac{u(x_0 + \Delta x, y_0) - u(x_0, y_0)}{\Delta x} + i \lim_{\Delta x \rightarrow 0} \frac{v(x_0 + \Delta x, y_0) - v(x_0, y_0)}{\Delta x}$$

$$f'(z_0) = \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0)$$

Define $\Delta z = i\Delta y$

$$f'(z_0) = \lim_{\Delta y \rightarrow 0} \frac{u(x_0, y_0 + \Delta y) + iv(x_0, y_0 + \Delta y) - u(x_0, y_0) - iv(x_0, y_0)}{i\Delta y}$$

$$f'(z_0) = \lim_{\Delta y \rightarrow 0} \frac{u(x_0, y_0 + \Delta y) - u(x_0, y_0)}{i\Delta y} + i \lim_{\Delta y \rightarrow 0} \frac{v(x_0, y_0 + \Delta y) - v(x_0, y_0)}{i\Delta y}$$

$$f'(z_0) = -i \frac{\partial u}{\partial y}(x_0, y_0) + \frac{\partial v}{\partial y}(x_0, y_0)$$

$$\Rightarrow \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0) = -i \frac{\partial u}{\partial y}(x_0, y_0) + \frac{\partial v}{\partial y}(x_0, y_0)$$

Splitting the real and imaginary parts we get that the Cauchy-Riemann equations are

$$\begin{cases} u_x = v_y \\ u_y = -v_x \end{cases}$$

If the Cauchy-Riemann equations are satisfied and the partial derivatives are continuous then the function is differentiable.

Some functions are not differentiable everywhere, but are differentiable at a point or a set of points.

- If $f(z)$ is differentiable everywhere in the complex plane then it is said to be **entire**.
- If $f(z)$ is differentiable in some region R then it is said to be **analytic** in R .
(note that this region cannot be a single point, as the Cauchy-Riemann equations require the partial derivatives to be continuous)

Ex: Show using the Cauchy-Riemann equations that $f(z) = \bar{z}$ is not differentiable anywhere.

$$\bar{z} = x - iy$$

$$u_x = 1 \neq v_y = -1$$

Ex2: Show that $f(z) = z^2$ is entire

$$f(z) = z^2 = (x + iy)^2 = x^2 - y^2 + 2ixy$$

$$u(x, y) = x^2 - y^2$$

$$v(x, y) = 2xy$$

$$u_x = 2x = v_y$$

$$u_y = -2y = -v_x$$

Ex3: Show that $f(z) = \bar{z}$ is differentiable but not analytic at $z_0 = 0$

$$\begin{aligned} |z|^2 + 2z &= x^2 + 2x + y^2 + i2y \\ u_x &= 2x + 2 = v_y = 2 \Rightarrow x = 0 \\ u_y &= 2y = -v_x = 0 \Rightarrow y = 0 \\ &\text{differentiable but not analytic on } z = \{0\} \end{aligned}$$

1.2.3 Conformal Mappings

Using the Cauchy-Riemann equations,

$$\begin{cases} u_x = v_y \\ u_y = -v_x \end{cases}$$

we can get the Laplacian of u and v ,

$$\begin{aligned} u_{xx} + u_{yy} &= v_{yx} - v_{xy} = 0 \\ v_{xx} + v_{yy} &= -(u_{yx} + u_{xy}) = 0 \end{aligned}$$

If the Laplacian of u and v are both zero then the function is said to be **harmonic**.

If $\nabla^2 u = 0$ then we can use the Cauchy-Riemann equations to find its harmonic conjugate v . Ex:
Find the harmonic conjugate of $u = xy - x + y$

$$\begin{aligned} u_x &= y - 1 = v_y \\ v &= \int y - 1 dy = \frac{y^2}{2} - y + h(x) \\ u_y &= x + 1 = -v_x = -h'(x) \\ h(x) &= \int -x - 1 dx = -\frac{x^2}{2} - x + C \\ v &= \frac{y^2}{2} - \frac{x^2}{2} - y - x + C \end{aligned}$$

Ex2: Find the harmonic conjugate of $u = \ln \sqrt{x^2 + y^2}$

$$\begin{aligned} u_x &= \frac{x}{x^2 + y^2} = v_y \\ v &= \int \frac{x}{x^2 + y^2} dy = \int \frac{1/x}{1 + \frac{y^2}{x^2}} dy = \arctan\left(\frac{y}{x}\right) + h(x) \\ u_y &= \frac{y}{x^2 + y^2} = -v_x = -\frac{y}{1 + \frac{y^2}{x^2}} \left(\frac{-1}{x^2}\right) - h'(x) = \frac{y}{x^2 + y^2} - h'(x) \Rightarrow h'(x) = 0 \\ h(x) &= C \\ v &= \arctan\left(\frac{y}{x}\right) + C \\ v &= \arg(z) + C \end{aligned}$$

Ex3: Find the harmonic conjugate of $u = \sin x \cosh y$

$$u = \sin x \cosh y$$

$$u_x = \cos x \cosh y = v_y$$

$$v = \int \cos x \cosh y dy = \cos x \sinh y + h(x)$$

$$u_y = \sin x \sinh y = -v_x = -(-\sin x \sinh y) - h'(x) \Rightarrow h'(x) = 0$$

$$h(x) = C$$

$$v = \cos x \sinh y + C$$

Another property is that

$$|f'(z)|^2 = |\nabla u|^2 = |\nabla v|^2$$

This also implies that

$$\nabla u \cdot \nabla v = 0$$

$$f'(z) = u_x + iv_x = \frac{1}{i}(u_y + iv_y)$$

$$|f'(z)|^2 = u_x^2 + v_x^2 = u_y^2 + v_y^2 = |\nabla u|^2 = |\nabla v|^2$$

$$\nabla u \cdot \nabla v = u_x u_y + v_x v_y = u_x v_x + (-v_x)(u_x) = 0$$

A conformal mapping is a mapping between two regions that preserves angles. If we have some function $f(z) = u + iv$ that is analytic then we can create a function of a function as

$$\Phi(u(x, y), v(x, y)) = \phi(x, y)$$

where $\phi(x, y)$ is a conformal mapping.

A conformal mapping will have the property that

$$\phi_{xx} + \phi_{yy} = |f'(z)|^2(\Phi_{uu} + \Phi_{vv})$$

where $|f'(z)|^2$ is known as the *conformal factor*.

This can be shown as follows:

$$\phi_x = \Phi_u u_x + \Phi_v v_x$$

$$\phi_{xx} = u_{xx} \Phi_u^2 + 2u_x v_x \Phi_u \Phi_v + v_{xx} \Phi_v^2 + \Phi_u u_{xx} + \Phi_v v_{xx}$$

$$\phi_y = \Phi_u u_y + \Phi_v v_y$$

$$\phi_{yy} = u_{yy} \Phi_u^2 + 2u_y v_y \Phi_u \Phi_v + v_{yy} \Phi_v^2 + \Phi_u u_{yy} + \Phi_v v_{yy}$$

$$\phi_{xx} + \phi_{yy} = \Phi_u \nabla^2 u + \Phi_v \nabla^2 v + \Phi_{uu} |\nabla u|^2 + 2\Phi_u \Phi_v \nabla u \cdot \nabla v + \Phi_{vv} |\nabla v|^2$$

$$\phi_{xx} + \phi_{yy} = |f'(z)|^2(\Phi_{uu} + \Phi_{vv})$$

Ex: $f(z) = z^2$ under $\Phi(u, v) = e^u + v^2$

$$f(z) = x^2 - y^2 + i(2xy)$$

$$u = x^2 - y^2$$

$$v = 2xy$$

$$\phi(x, y) = e^{x^2 - y^2} + (2xy)^2$$

$$\phi_{xx} + \phi_{yy} = 4|z|^2(e^u + 2)$$

$f(z)$ is considered a conformal mapping if f is analytic and $f'(z) \neq 0$. As a consequence, if $\phi_{xx} + \phi_{yy} = 0$ then $\Phi_{uu} + \Phi_{vv} = 0$.

If f is a conformal mapping then it will preserve angles.

If we have two curves C_1 and C_2 described by the parametrized functions $z_1(t)$ and $z_2(t)$ that intersect at a point z_0 then the angle between the two curves is given by $\theta = \arg(z_2'(t)) - \arg(z_1'(t))$.

If we then apply a conformal mapping $w = f(z)$ to the curves then we get $w_1(t) = f(z_1(t))$ and $w_2(t) = f(z_2(t))$ and the angle between the two curves is given by $\theta_w = \arg(w_2'(t)) - \arg(w_1'(t))$.

$$\theta_w = \arg(f'(z_2)z_2'(t)) - \arg(f'(z_1)z_1'(t))$$

$$\theta_w = \arg(f'(z_2)) - \arg(f'(z_1)) + \arg(z_2'(t)) - \arg(z_1'(t))$$

$$\arg(f'(z_1)) = \arg(f'(z_2))$$

$$\Rightarrow \theta_w = \arg(z_2'(t)) - \arg(z_1'(t)) = \theta$$

So the angle between the two curves is preserved under a conformal mapping.

Note that $f'(z) \neq 0$ is a necessary condition in this proof.

If we have a nonconformal mapping then the angle between the two curves will not be preserved. One such case is that of $f(z) = z^2$ at $z_0 = 0$. At this point, $f'(z_0) = 0$ and the angle between the two curves is doubled.

Conformal mappings also have the property that they map Neumann boundary conditions to Neumann boundary conditions.

If we let \hat{n} represent the normal vector to the curve $\phi(x, y)$ then they are related by

$$\frac{\partial \phi}{\partial n} = \nabla \phi \cdot \hat{n}$$

and the mapping is similarly related brcurly

$$\begin{aligned} \frac{\partial \Phi}{\partial n'} &= \nabla \Phi \cdot \hat{n}' \\ \frac{\partial \phi}{\partial n} &= |f'(z)| \frac{\partial \Phi}{\partial n'} \end{aligned}$$

So for Neumann boundary conditions, we will have

$$\frac{\partial \phi}{\partial n} = 0 \Rightarrow \frac{\partial \Phi}{\partial n'} = 0$$

Some examples of conformal mappings come from harmonic functions (having the property that $\nabla^2 \phi = 0$).

Some common harmonic functions are:

- $\phi = C$
- $\phi = ax + by + c$

- $\ln \sqrt{x^2 + y^2}$, $\mathbb{C} \setminus 0$
- $\phi = \text{Arg}(z)$, $\mathbb{C} \setminus (-\infty]$
- $\phi = x^2 - y^2$

1.2.4 Conformal Mapping to Solve Laplace's Equation

Given the useful properties of matching boundary conditions, we can use conformal mappings to help us solve Laplace's equation for a given region.

Ex: Given $\nabla^2 \phi = 0$ for $1 < x^2 - y^2 < 4$ and $\phi = 1$ on $x^2 - y^2 = 1$ and $\phi = 3$ on $x^2 - y^2 = 4$, find $\phi(x, y)$.

$$\text{choose } \phi = x^2 - y^2$$

$$\text{choose } \Phi(u, v) = Au + B$$

$$\begin{cases} A(1) + B = 1 \\ A(4) + B = 3 \end{cases} \Rightarrow A = \frac{2}{3}, B = \frac{1}{3}$$

$$\Phi(u, v) = \frac{2}{3}u + \frac{1}{3}$$

$$u = x^2 - y^2$$

$$\phi(x, y) = \frac{2}{3}(x^2 - y^2) + \frac{1}{3}$$

Ex2: Given $\nabla^2 \phi = 0$ within the circular region $\mathcal{D} = \{1 < x^2 + y^2 < 4\}$ with $\phi = 1$ on $x^2 + y^2 = 1$ and $\phi = -2$ on $x^2 + y^2 = 4$, find $\phi(x, y)$.

$$\phi(x, y) = A_1 \ln r + A_2$$

$$r = \sqrt{x^2 + y^2}$$

$$r = 1 : \phi = 1 \Rightarrow A_1 \ln(1) + A_2 = 1$$

$$r = 2 : \phi = -2 \Rightarrow A_1 \ln(2) + A_2 = -2$$

$$\Rightarrow A_2 = 1, A_1 = -\frac{3}{\ln 2}$$

$$\phi(x, y) = -\frac{3}{\ln 2} \ln \sqrt{x^2 + y^2} + 1$$

Ex3: Given $\nabla^2 = 0$ within the strip described by $\{z : -3 \leq 3\Re(z) - 4\Im(z) \leq 2\}$ with $\phi = 0$ for $\{z : -3 = 3\Re(z) - 4\Im(z)\}$ and $\phi = 4$ for $\{z : 3\Re(z) - 4\Im(z) = 2\}$ find $\phi(x, y)$

$$u = 3x - 4y$$

$$u(-3) = 0, u(2) = 4$$

$$\Phi = Au + B$$

$$\Phi(-3) = -3A + B = 0 \Rightarrow B = 3A$$


$$\Phi(2) = 2A + 3A = 5A = 4 \Rightarrow A = \frac{4}{5} \Rightarrow B = \frac{12}{5}$$

$$\phi(x, y) = \frac{4}{5}(3x - 4y) + \frac{12}{5}$$

Ex4: Given $\nabla^2 \phi = 0$ for the upper half-plane described by $\{y > 0 \wedge x \in \mathbb{R}\}$ with the boundary conditions along the x-axis given by

$$\phi(x, 0) = \begin{cases} -1 & x < -5 \\ 0 & -5 < x < -1 \\ 2 & -1 < x < 2 \\ 1 & x > 2 \end{cases}$$

find $\phi(x, y)$.



One trick to solve a problem like this is to choose a linear combination of functions of the form $\text{Arg}(z - z_0)$ with a different point z_0 for every place where the boundary condition changes along the x-axis.

$$\phi = A_1 \text{Arg}(z + 5) + A_2 \text{Arg}(z + 1) + A_3 \text{Arg}(z - 2) + A_4$$

$$\phi(x > 2, 0) = A_4 = 1$$

$$\phi(-1 < x < 2, 0) = \pi A_3 + 1 = 2 \Rightarrow A_3 = \frac{1}{\pi}$$

$$\phi(-5 < x < -1, 0) = \pi A_2 + 1 + 1 = 0 \Rightarrow A_2 = -\frac{2}{\pi}$$

$$\phi(x < -5, 0) = \pi A_1 - 2 + 2 = -1 \Rightarrow A_1 = -\frac{1}{\pi}$$

$$\phi(x, y) = -\frac{1}{\pi} \text{Arg}(z + 5) - \frac{2}{\pi} \text{Arg}(z + 1) + \frac{1}{\pi} \text{Arg}(z - 2) + 1$$

We can also apply these techniques to other types of boundary conditions in some cases.

Ex5: Given $\nabla^2 \phi = 0$ in the circular region $\{z : 1 \leq |z| \leq 2\}$ with the boundary conditions $\phi = 1$ for $|z| = 1$ and $\frac{\partial \phi}{\partial r} = 2$ for $|z| = 2$, find $\phi(x, y)$

$$\phi = A \ln r + B$$


$$\phi(1) = B = 1$$

$$\frac{\partial \phi(2)}{\partial r} = \frac{A}{2} = 2 \Rightarrow A = 4$$

$$\phi = 4 \ln r + 1$$

$$\phi(x, y) = 4 \ln \sqrt{x^2 + y^2} + 1$$

If we have a region that has a semicircle in it we can use the Joukowski mapping to transform it into a region that is easier to work with.



Ex6: Given $\nabla^2 \phi = 0$ in the upper region of the plane described by $\{y > 0 \wedge x^2 + y^2 > 9\}$ with the boundary conditions $\phi = -1$ for $x < -3$, $\phi = 0$ for $x^2 + y^2 = 9$, and $\phi = 2$ for $x > 3$, find $\phi(x, y)$.

$$\begin{cases} \phi(x, 0) = -1 & x < -3 \\ \phi(x, y) = 0 & x^2 + y^2 = 9 \\ \phi(x, 0) = 2 & x > 3 \end{cases}$$

$$w = \frac{1}{2} \left(\frac{z}{3} + \frac{3}{z} \right) = u + iv$$

$$\Phi = A_1 \operatorname{Arg}(w + 1) + A_2 \operatorname{Arg}(w - 1) + A_3$$

$$\Phi(u > 1, v) = A_3 = 2$$


$$\Phi(-1 < u < 1, v) = \pi A_2 + 2 = 0 \Rightarrow A_2 = -\frac{2}{\pi}$$

$$\Phi(u < -1, v) = \pi A_1 - 2 + 2 = -1 \Rightarrow A_1 = -\frac{1}{\pi}$$

$$\Phi = -\frac{1}{\pi} \operatorname{Arg}(w + 1) - \frac{2}{\pi} \operatorname{Arg}(w - 1) + 2$$

$$\phi(z) = -\frac{1}{\pi} \operatorname{Arg} \left(\frac{1}{2} \left(\frac{z}{3} + \frac{3}{z} \right) + 1 \right) - \frac{2}{\pi} \operatorname{Arg} \left(\frac{1}{2} \left(\frac{z}{3} + \frac{3}{z} \right) - 1 \right) + 2$$

In the case that we have a semicircle not of radius 1 we can apply a scaling before the Joukowski mapping to get the correct radius.



$$w = \frac{1}{2} \left(\frac{z}{a} + \frac{a}{z} \right)$$

1.2.5 Sinusoidal Functions

If we recall Euler's formula $e^{ix} = \cos x + i \sin x$ we can use with complex numbers to get the following identities for complex sinusoids:

- $\sin z = \frac{e^{iz} - e^{-iz}}{2i}$
- $\cos z = \frac{e^{iz} + e^{-iz}}{2}$
- $\sinh z = \frac{e^z - e^{-z}}{2}$
- $\cosh z = \frac{e^z + e^{-z}}{2}$
- $\cos z = \sin\left(\frac{\pi}{2} - z\right)$
- $\sinh z = -i \sin(iz)$
- $\cosh z = \cos(iz) = \sin\left(\frac{\pi}{2} - iz\right)$
- $\frac{d}{dz} \sin z = \cos z$
- $\frac{d}{dz} \cos z = -\sin z$
- $\cos^2 z + \sin^2 z = 1$
- $\frac{d}{dz} \sinh z = \cosh z$
- $\frac{d}{dz} \cosh z = \sinh z$
- $\cosh^2 z - \sinh^2 z = 1$

The most notable difference between the real and complex versions of these functions is that $|\sin z| \not\leq 1$. We will see cases of this soon but it is easy to see once we write out the real and imaginary components of $\sin z$.

$$\begin{aligned}\sin z &= \sin(x + iy) = \frac{e^{i(x+iy)} - e^{-i(x+iy)}}{2i} = \frac{e^{-y}e^{ix} - e^ye^{-ix}}{2i} \\ \sin z &= \frac{e^{-y}(\cos x + i \sin x) - e^y(\cos x - i \sin x)}{2i} = \frac{e^{-y} - e^y}{2} \cos x + \frac{e^{-y} + e^y}{2i} \sin x \\ \sin z &= \sin x \cosh y + i \cos x \sinh y\end{aligned}$$