

# MONTE CARLO SIMULATION

## 3D Simulation.

### Final Results

#### QUESTION:

Point source is anisotropic: all electrons are emitted along one direction; electron-atom scattering is anisotropic: Theta,  $\theta$  is determined from  $\sin(\theta/2) = a_3$ , (where  $a_1$ ,  $a_2$  and  $a_3$  are random numbers selected between 0 and 1)

Determination of  $\lambda$  and  $\phi$  is given by:

Lambda,  $\lambda = -\ln a_1$ , phi,  $\phi = 2\pi a_2$ .

It is necessary to make simulation for the anisotropic case and to compare results with [previous results for isotropic case](#).

Lambda is the length between the collisions, phi and theta are the angular displacement in x-y, x-z coordinates respectively.

From the sine identity formula, we have that:


$$\sin(\theta/2) = a_3$$

$$(\theta/2) = \arcsin(a_3)$$


$$\theta = 2 * \arcsin(a_3)$$

## Anisotropic Electron-Atom Scattering in 3D for 100, 1000, and 10000 collisions.


For 100 collisions:



The Distance from the Source against Number of Collisions: Absolute Values




The graph of  $\log(L)$  versus  $\log(N)$




**The Slope ( $\alpha$ ): 0.5681**


**For 1000 collisions:**



The Distance from the Source against Number of Collisions: Absolute Values




The graph of  $\log(L)$  versus  $\log(N)$




**The Slope ( $\alpha$ ): 0.5414**


**For 10000 collisions:**



The Distance from the Source against Number of Collisions: Absolute Values



The graph of  $\log(L)$  versus  $\log(N)$



**The Slope ( $\alpha$ ): 0.5302**

The codes for these Results can be found in [My GitHub](#)

*By Mfeuter Joseph, Tachia*

*Moscow Institute of Physics and Technology (MIPT)*