

Problem 1 (8 points, 1 point for each part)

This problem consists of some true/false questions regarding concepts of statistical inference. Indicate if a statement is true or false. You do not need to explain why it is true or false.

- 1.1 In a hypothesis test, the p value is 0.043. This means that the null hypothesis would be rejected at $\alpha = 0.05$.
- 1.2 If the null hypothesis is rejected by a one-tailed hypothesis test at α , then it will also be rejected by a two-tailed test at α .
- 1.3 If a null hypothesis is rejected at the 0.01 level of significance, it will also be rejected at the 0.05 level of significance.
- 1.4 The probability of a type II error is controlled in a hypothesis test by establishing a specific significance level.
- 1.5 If we decrease the confidence coefficient for a fixed n , we decrease the width of the confidence interval.
- 1.6 If a 95% confidence interval on μ was from 50.5 to 60.6, we would reject the null hypothesis that $\mu = 60$ at the 0.05 level of significance.
- 1.7 If the sample size is increased and the level of confidence is decreased, the width of the confidence interval will increase.
- 1.8 A research article reports that a 95% confidence interval for mean reaction time is from 0.25 to 0.29 seconds. Therefore, about 95% of individuals will have reaction times in this interval.

Problem 2 (9 points, 1 point for each part)

Multiple choice questions. Only one of the statements is correct.

- 2.1 In a hypothesis test the p value is 0.043. This means that we can find statistical significance at:
 - a) both the 0.05 and 0.01 levels
 - b) the 0.05 but not at the 0.01 level
 - c) the 0.01 but not at the 0.05 level
 - d) neither the 0.05 or 0.01 levels
 - e) none of the above

2.2 A research report states: The differences between public and private school seventh graders' attitudes toward minority groups were statistically significant at the $\alpha = 0.05$ level. This means that:

- a) It has been proven that the two groups are different.
- b) There is a probability of 0.05 that the attitudes of the two groups are different.
- c) There is a probability of 0.95 that the attitudes of the two groups are different.
- d) If there is no difference between the groups, the difference observed in the sample would occur by chance with a probability of no more than 0.05.
- e) None of the above is correct.

2.3 If the null hypothesis is really false, which of these statements characterizes a situation where the value of the test statistic falls in the rejection region?

- a) The decision is correct.
- b) A type I error has been committed.
- c) A type II error has been committed.
- d) Insufficient information has been given to make a decision.
- e) None of the above is correct.

2.4 If the null hypothesis is really false, which of these statements characterizes a situation where the value of the test statistic does not fall in the rejection region?

- a) The decision is correct.
- b) A type I error has been committed.
- c) A type II error has been committed.
- d) Insufficient information has been given to make a decision.
- e) None of the above is correct.

2.5 If the value of any test statistic does not fall in the rejection region, the decision is:

- a) Reject the null hypothesis.
- b) Reject the alternative hypothesis.
- c) Fail to reject the null hypothesis.
- d) Fail to reject the alternative hypothesis.
- e) There is insufficient information to make a decision.

2.6 For a particular sample, the 95% two-sided confidence interval for the population mean is from 11 to 17. You are asked to test the hypothesis that the population mean is 18 against a two-sided alternative. Your decision is:

- a) Fail to reject the null hypothesis, $\alpha = 0.05$.
- b) Reject the null hypothesis, $\alpha = 0.05$.
- c) There is insufficient information to decide.

2.7 If we decrease the confidence level, the width of the confidence interval will:

- a) Increase
- b) remain unchanged
- c) decrease
- d) double
- e) none of the above

2.8 If the value of the test statistic falls in the rejection region, then:

- a) We cannot commit a type I error.
- b) We cannot commit a type II error.
- c) We have proven that the null hypothesis is true.
- d) We have proven that the null hypothesis is false.
- e) None of the above is correct.

2.9 You are reading a research article that states that there is no significant evidence that the median income in the two groups differs, at $\alpha = 0.05$. You are interested in this conclusion, but prefer to use $\alpha = 0.01$.

- a) You would also say there is no significant evidence that the medians differ.
- b) You would say there is significant evidence that the medians differ.
- c) You do not know whether there is significant evidence or not, until you know the p value.

Problem 3 (10 points, 5 points for each part)

Suppose that for a given population with $\sigma = 7.2$ we want to test $H_0: \mu = 80$ against $H_1: \mu < 80$ based on a sample of $n = 100$. For this problem, you can use R to find the probabilities associated with the standard normal and z_α but you need to present sufficient details about your calculations.

- (1) If the null hypothesis is rejected when $\bar{y} < 76$, what is the probability of a type I error?
- (2) What would be the rejection region if we wanted to have a level of significance of exactly 0.05?

Problem 4 (25 points)

The family incomes in a certain city in 1970 had a mean of \$14,200 with a standard deviation of \$2600. A random sample of 75 families taken in 1975 produced $\bar{y} = \$15,300$ (adjusted for inflation). For this problem, you can use R to find the probabilities associated with the standard normal and z_α but you still need to present sufficient details about your calculations.

- (1) **(11 points)** Assume the standard deviation has remained unchanged. Use the rejection region method to test if the mean family income has changed at a 0.05 level of significance. Please clearly specify 5 steps used in the test.
- (2) **(6 points)** Calculate the power of the test from (1) for the mean income of \$15300 and \$15600.
- (3) **(3 points)** Find the p -value associated with this test.
- (4) **(5 points)** Construct a 99% two-sided confidence interval on the mean family income in 1975. Based on this confidence interval, state if you would like to reject the null hypothesis from part (1) with a significance level of 0.01.