

# Homework 2

Your Name

February 27, 2025

We will now start reflecting on the coding questions of Homework 2. The code base that can be used to replicate the result can be found in the following link: <https://github.com/TagoreZhao/STAT260/tree/main/HW2>

## 1 Problem 6, 7, and 8

The code needed for generating these three matrices is given below:

```
1 import numpy as np
2
3 def generate_covariance_matrix(d):
4     indices = np.arange(d)
5     Sigma = 2 * 0.5 ** np.abs(indices[:, None] - indices[None, :])
6     return Sigma
7
8 def generate_gaussian_A(n, d, seed=1234):
9     rng = np.random.default_rng(seed)
10    Sigma = generate_covariance_matrix(d)
11    mean = np.ones(d)
12    A = rng.multivariate_normal(mean, Sigma, size=n)
13    return A
14
15 def generate_t_distribution_A(n, d, df, seed=1234):
16     rng = np.random.default_rng(seed)
17     Sigma = generate_covariance_matrix(d)
18     mean = np.ones(d)
19     z = rng.multivariate_normal(mean, Sigma, size=n)
20     chi2_samples = rng.chisquare(df, size=(n, 1))
21     A = z / np.sqrt(chi2_samples / df)
22     return A
```

Listing 1: Python code for generating matrices

Since numpy does not provide built in functions for generating t-distributed random variables, we have to generate the random variables ourselves. The Gaussian random variables are generated using the `multivariate_normal` function, while the t-distributed random variables are generated using the formula  $A = Z / \sqrt{\chi^2 / df}$ , where  $Z$  is the Gaussian random variable,  $\chi^2$  is the chi-squared random variable, and  $df$  is the degrees of freedom.

## Problem 9

We